Effects of phenotypic variation on consumer coexistence and prey community structure

Shane L. Hogle[1,6,*], Iina Hepolehto[2,3,6], Lasse Ruokolainen[3], Johannes Cairns[4,5], Teppo Hiltunen[1,*]

[1]Department of Biology, University of Turku, Turku, Finland
[2]Department of Microbiology, University of Helsinki, Helsinki, Finland
[3]Faculty of Biological and Environmental Sciences, University of Helsinki, Helsinki, Finland
[4]Organismal & Evolutionary Biology Research Programme, University of Helsinki, Finland
[5]Department of Computer Science, University of Helsinki, Finland
[6]These authors contributed equally

[*]**Corresponding Authors:** Shane Hogle and Teppo Hiltunen

Author Contributions: SLH: conceptualization, supervision, software, formal analysis, writing - original draft. IH: conceptualization, investigation - lead, visualization, writing - review & editing. LR: supervision, formal analysis, funding acquisition, writing - review & editing. JC: conceptualization, writing - review & editing. TH: conceptualization, supervision - lead, project administration, funding acquisition, writing - review & editing. All authors provide approval for publication.

Competing Interest Statement: The authors declare no conflict of interest.

Preprint Servers: The article is available on the bioRxiv preprint server: https://www.biorxiv.org/content/10.1101/2021.06.09.447767v1.

Data Availability: Raw sequencing data is available from the NCBI Sequence Read Archive (SRA) under the BioProject accession number PRJNA725120. Preprocessed count tables are available from https://github.com/slhogle/consumer-competition. Scripts reproducing all figures and steps in the data analysis are available from https://github.com/slhogle/consumer-competition.

This PDF file includes:
Main Text
Figures 1 to 6

Abstract

A common expectation in ecology is that trait variation among individuals from the same species promotes the coexistence of competing species. However, theoretical and empirical studies testing this expectation have been equivocal. One complicating factor is that species exist within multitrophic networks characterized by direct and indirect linkages across trophic levels. It is currently unknown how intraspecific trait diversity at the consumer trophic level affects interspecific consumer competition and population dynamics at lower trophic levels. Here we use an experimental microbial ecosystem consisting of ciliated protozoa, nematode worms, and 24 species of bacterial prey to mechanistically show how intraspecific consumer variation can 1) modulate competitive and niche differences between competing consumers and 2) produce cascading effects on ecological dynamics at lower trophic levels. Although the nematode could competitively exclude ciliates with low trait variation, we found that ciliate trait variation inverted the consumer hierarchy, ultimately excluding the nematode. Competition between the high trait variation ciliate and the nematode altered the temporal trajectories of individual prey species in non-additive ways, mediated by prey traits related to growth rate and defense. We connected the competitive outcome to consumer prey preferences by performing pairwise co-cultures with each consumer and prey species and measuring the clearance of prey biomass. High trait variation in the ciliate increased the mean and variance of the species-specific prey clearance rate over the low trait variation ciliate, which resulted in increased selective foraging behavior. These findings provide a new multitrophic perspective on trait diversity and species coexistence.

Significance

Ecological theory posits that a balance of trait differences influencing competition (competitive differences) and the use of the environment (niche differences) allow different species to coexist in time and space. However, a substantial portion of trait variation exists within species. We manipulated intraspecific trait diversity in a microscopic ciliate competing with a nematode for bacterial prey and found that consumer trait variation inverted the competitive hierarchy that existed under low trait variation. These altered consumer competitive dynamics produced non-additive ecological effects on prey abundance, resulting in prey communities distinct from those forming under either predator individually. Our results suggest that predicting consumer competitive population dynamics and the assembly of prey communities will require understanding the extent of trait variation within consumer species.

Introduction

Comparative studies of species traits have historically focused on trait differences between species. However, ecologists increasingly recognize that a large amount of the trait variation in ecological communities exists between individuals from the same species. For example, trait variation within species accounts for up to one-fourth of total trait variation in plant communities[1] and can have ecological consequences comparable in magnitude to those from separate species[2]. Intraspecific trait variation may emerge from phenotypic plasticity (e.g., maternal effects or learned behavior) and heritable evolutionary change (e.g., intergenerational changes in allele frequencies).

While intraspecific trait variation is ubiquitous in natural ecological communities, its relevance for competitive interactions and species coexistence is unclear. Theory has considered intraspecific variation and species coexistence through the evolutionary lens of character displacement[3], but the purely ecological effects of intraspecific trait variation have historically been ignored. Work built from coexistence theory[4] suggests that the ecological effects of intraspecific variation should hinder species

coexistence and mostly reinforce existing competitive hierarchies[5, 6]. However, other studies have argued that intraspecific variation should promote coexistence - for example, by blending average competitive differences between species, reducing the fraction of individuals that strongly compete, or processes akin to bet-hedging [7–9]. From the perspective of coexistence theory, intraspecific trait variation can have both positive and negative effects on average trait values that drive both niche and competitive differences between species[10]. Therefore, intraspecific trait variation may have complex effects that can both promote and hinder species coexistence[11].

Studies on the effects of intraspecific trait variation have primarily focused on ecological effects at a single trophic level, predominantly with plants[10]. The handful of studies investigating the consequences of intraspecific variation for trophic interactions have concentrated on variable prey defense traits that reduce consumption risk[12]. Very few studies have examined trophic interaction from the perspective of consumers and offensive trait variation (but see [13, 14]). Furthermore, the simultaneous effects of consumer trait variation on trophic interactions and consumer competition are entirely unknown. This is important because most natural ecosystems contain multiple consumer species competing for prey[15]. Furthermore, a fundamental understanding of natural community assembly processes requires adopting multitrophic perspectives that account for indirect and direct linkages spanning trophic levels[16].

Motivated by these knowledge gaps, we conducted an experiment to determine 1) how intraspecific consumer trait variation influences the competitive dynamics of two consumer species and 2) how intraspecific trait variation and consumer species composition impacts bacterial species diversity at the prey trophic level (Fig. 1). We combined a 24-species isogenic bacterial prey community (Table S1) with an isogenic nematode worm, *Caenorhabditis elegans*, and an isogenic ciliated protozoan *Tetrahymena thermophila* - hereafter Low Trait Variation (LTV) ciliate. In some treatments, we replaced the LTV ciliate consumer with an even mixture of 20 different ciliate populations, each displaying high intraspecific phenotype variation - hereafter High Trait Variation (HTV) ciliate[17]. In this way, we could manipulate the standing stock of trait diversity in the ciliate consumer in the presence or absence of competition from the nematode. We then combined the LTV ciliate, the HTV ciliate, the nematode, and the prey bacterial community in experimental microcosms with a full-factorial design (six conditions) and followed consumer densities, prey density, and bacterial community composition over 60 days. We also measured key functional traits from both prey and consumer species to determine how community assembly and consumer composition were related to species phenotypes.

Results

Consumer trait variation flips the consumer competitive hierarchy   The magnitude of trait variation in the ciliate consumer was sufficient to alter ecological dynamics of the competing consumer. Each consumer by itself (nematode, LTV ciliate, HTV ciliate) could feed and grow successfully on the bacterial prey community (Fig. 2A), and ciliate trait variation did not significantly alter the carrying capacity for the ciliate in the absence of competition (Table S3). Prey biomass differed by less than twofold across all consumer treatments (mean fold difference of 1.08, Table S2), indicating that each consumer or consumer pair drew down total prey resources to similar levels. The ciliate and nematode grew from starting densities of 10^4 and 10 individuals ml^{-1}, respectively. In the absence of competition, the LTV ciliate, the HTV ciliate, and the nematode stabilized at $4.2 \pm 2.3 \times 10^4$, $5.0 \pm 2.3 \times 10^4$, and $6.0 \pm 2.5 \times 10^3$ individuals ml^{-1}, respectively, after an exponential growth period of about 17 days (Fig. 2A). However, the density of each consumer was strongly dependent upon the competitive arrangement between species (Tables S3-S4). Competition between the LTV ciliate and the nematode excluded the ciliate from the microcosms by day 11, and by day 24, the nematode density had reached competition-free densities. However, competition between the HTV ciliate and the nematode flipped the consumer hierarchy, ultimately excluding the nematode but promoting longer species coexistence.

Figure 1. textbfStudy setup
A) The experimental bacterial community with different consumer combinations. Ctrl - bacteria only, N - bacteria with nematode, C_{LTV} - bacteria with low trait diversity isogenic ciliate, C_{HTV} - bacteria high trait diversity mixed ciliate populations, NC_{LTV} - bacteria + nematode + low trait diversity ciliate, NC_{HTV} - bacteria + nematode + high trait diversity ciliate. **B)** Schematic representation of the serial transfer experiment. Each experimental condition from A) was performed in four biological replicates and transferred/sampled every four days. **C)** Traits of individual bacterial species were measured in monoculture and inferred from genome sequences (see methods). **D)** Consumers were grown on each bacterial species to assess consumer grazing efficiency (see methods).

We next attributed consumer dynamics to changes in niche differences and competitive differences following coexistence theory. Coexistence theory describes the coexistence of two species as a balance between stabilizing and equalizing mechanisms[4]. Equalizing mechanisms relate to inherent competitive differences between species that describe dominance hierarchies in a community. Stabilizing mechanisms (i.e., niche differences) relate to the ability of each species to recover from low density after perturbation - e.g., when species can use the environment in different ways to avoid competition. Niche and competitive differences can be quantitatively estimated by fitting population density data to a model of competitive population dynamics, the parameters of which represent within-species and between-species interactions[18]. Using consumer density data and coefficients derived from the model we estimated niche differences $(1 - \rho)$ and the competitive differences $(k_{ciliate}/k_{nematode})$ between the HTV or LTV ciliate and the nematode (see methods). Results indicate that relatively strong niche differences separated the LTV ciliate and nematode, but these differences were more than compensated by a strong competitive advantage for the nematode (Fig. 2B). Surprisingly, the model predicted the LTV ciliate and nematode were at or near the boundary of coexistence. In both HTV and LTV competitive arrangements we observed very low abundances of the inferior competitor (< 100 ciliates ml^{-1}, < 5 nematodes ml^{-1}) in some of the replicates near day 40. Accurate manual counting is challenging at these low densities, and it may be that the inferior competitor never truly became extinct. Finally, high trait variation in the ciliate strongly increased competitive differences between the ciliate and nematode while simultaneously *decreasing* niche differences. The magnitude of the competitive change was much larger than the niche change, which suggests that high trait variation in the ciliate predominantly drove equalizing competitive

Figure 2. Consumer competitive hierarchy depends upon ciliate trait variation
A) Consumer and bacterial biomass during the course of the experiment. Points are observations, lines are estimated marginal means (mean response and 95% confidence levels adjusted for all other variables) from generalized additive models. Model summaries and contrasts of marginal means are available in Tables S2-S4). **B)** Niche and competitive differences between the ciliate and the nematode depending on whether the ciliate had low trait variation (NC_{LTV}) or high trait variation (NC_{HTV}). Coexistence (grey region) occurs when when stabilizing niche differences $(1 - \rho)$ are greater than competitive differences $(k_{ciliate}/k_{nematode})$. Coexistence is defined by the inequality $\rho < k_{ciliate}/k_{nematode} < 1/\rho$. Points and line ranges shows the mean and standard deviation of niche and competitive differences from four biological replicates.

differences between competitors rather than stabilizing niche differences (Fig. 2B).

Prey community assembly follows consumer dynamics: Our experimental setup allowed us to investigate the prey community response to consumer trait variation. We first asked how consumer trait variation influenced the repeatability of prey responses across biological replicates. Overall, the prey species abundance distribution across all populations/treatments followed the canonical form[19] with a handful of abundant species and many rare species (Fig. 3A). We quantified the repeatability of prey

communities by calculating the scaled Shannon diversity dissimilarity (D') across treatments at each   109
sampling point. Replicate prey communities were highly consistent in composition across all consumer   110
combinations and time (Fig. S1, Table S5). These results show that 1) prey communities generally   111
assembled under each treatment condition so that the majority of prey species were represented by the   112
minority of sequencing reads, 2) technical variation in laboratory conditions (e.g., temperature or   113
transfer volumes) were minimal for these experiments, and 3) the process of prey community assembly   114
alongside each consumer or consumer pair was strongly deterministic.   115

Next, we asked how consumers affected both the local abundance and evenness of prey species present in   116
each treatment (i.e., alpha diversity). Generally, we observed a rapid peak and then monotonic decline of   117
the rarest species during the first 15 days prior to the stabilization of the consumer (Fig. 3B). In the last   118
30 days, these rare species would briefly increase in abundance above 0.01%, then rapidly decline. The   119
more common species generally established themselves at high abundances ($> 10\%$) after 14 days when   120
both total consumer and prey densities had stabilized. Afterwards, these common species displayed   121
dynamic peaks and declines for the remainder of the experiment. We next noticed that prey Shannon   122
diversity varied in two distinct phases across treatments, which we then operationally defined using   123
multiple change point regression (Fig. S2). Prey Shannon diversity declined linearly with time in the   124
first phase - hereafter sorting phase - which was followed by oscillation around a temporally zero-trend   125
mean in the second phase - hereafter equilibrium phase (Fig. 3B, Supplementary methods). We found   126
that the presence of a a consumer or co-consumer pair slowed the rate of Shannon diversity decline in   127
the sorting phase relative to the control, with the nematode slowing decline by the largest magnitude on   128
average (Fig. S3, Table S6). In the equilibrium phase, predation generally supported higher mean   129
Shannon diversity, with the exception of the LTV ciliate, which decreased prey diversity to the lowest   130
mean value of all treatments (Tables S7). Consumer competition did not change either the rate of   131
diversity decline in the sorting phase or the mean diversity in the equilibrium phase relative to consumer   132
monocultures. Thus, the effect of two competing consumers on the Shannon diversity of the prey   133
community was generally equivalent to that of a single predator.   134

We then examined how the differentiation of prey community composition (i.e., beta diversity) changed   135
between consumer treatments and through time. We used ordination plots to visualize the temporal   136
trajectory of each community through system phase space (i.e., community space) and to quantify   137
relative state changes under each consumer treatment[20]. The LTV ciliate prey community traversed   138
the longest distance in community space followed by the no-consumer control and then the combination   139
of the LTV ciliate and nematode (Fig. 3C). The prey communities assembling with only the HTV ciliate   140
or the nematode both traversed the smallest distance and had significantly smaller jumps through   141
community space than the no-consumer control and the LTV ciliate (Table S8). Thus, the HTV ciliate   142
or the nematode rapidly stabilized the prey community in a relatively small region of community state   143
space. Alternatively, the LTV ciliate appeared to drive the prey community across a wider range of   144
community states similar to the no-consumer treatment. The effect of two consumers was generally to   145
reduce the distance traversed in prey community from the no-consumer control, but not to the extent of   146
the HTV ciliate or nematode alone.   147

Consumer trait variation results in non-additive effects on prey abundances: We next   158
asked how ciliate trait variation affected the response of individual prey species to the presence of   159
multiple predators. In animal communities[21, 22], the effects of multiple predators on prey often do not   160
combine additively but result in higher (risk reduction) or lower (risk enhancement) prey abundances   161
than would be predicted by a linear combination of individual predator effects. These emergent multiple   162
predator effects[15] cannot be predicted based on the knowledge of the effects of a single predator on   163
prey but are dependent upon the context of the consumer arrangement. We used additive longitudinal   164
Gaussian process regression[23] to decompose prey abundance trajectories into components   165
corresponding to the effect of each consumer and a statistical interaction (i.e., an emergent multiple   166
predator effect) between consumer types.   167

Figure 3. Prey community response to consumer trait variation

A) Prey species abundance distributions from each experimental treatment. X-axis ranks prey species (1977 most abundant) by average across all conditions. Y-axis shows relative abundance. **B)** Replicate mean abundance trajectories over time (normalized to each species maximum) for each of the 24 bacterial prey species. **C)** Non-metric multidimensional scaling ordination of trajectories through community space. Larger, outlined circles show means across replicates, while small circles are replicates. Colors represent days from the experiment start. Means are connected in chronological order.

We found that different consumer combinations resulted in significant non-additive effects - i.e., emergent multiple predator effects - for both abundant and rare prey species (Fig. S4). The effects were generally strongest when the HTV ciliate was paired with the nematode. For example, the ciliate had the largest predicted effect on *Aeromonas caviae* 1972 during exponential growth of the consumer (first 20 days) and before the ciliate density had stabilized (Fig. 4). The nematode had the largest predicted effect on *A. caviae* 1972 after its exponential growth phase (30 days). The model did not support the existence of a significant statistical interaction between the LTV ciliate and nematode - i.e., the effect of the combined consumer arrangement was effectively the sum of the two individual consumers. However, competition between the HTV ciliate and nematode resulted in a substantial risk enhancement for *A. caviae* 1972 (Fig. 4), which explained the greatest proportion of variance in the model. In the case of *C. testosterone* 0403, there was a significant but small non-additive effect between the LTV ciliate and nematode, but the greatest amount of variance was explained by the statistical interaction between the HTV ciliate and the nematode model terms. Surprisingly, this effect was persistent after the nematode was effectively excluded from the microcosm (Fig. S5), implying that prior dynamics from the co-consumer arrangement influenced the state of the microcosms long after one consumer had been competitively excluded. Finally, we also observed instances where a prey species could only exist in the presence of two consumers but not with either consumer alone. For example, the rare species *Niabella yanshanensis* 3031 only reached appreciable abundance in the presence of both the HTV ciliate and nematode, making it

impossible to predict the longitudinal trajectory of this strain from individual consumer effects or the effect of trait variance alone (Fig. S6). Thus, the dynamics of *Niabella yanshanensis* 3031 were an emergent property of the HTV ciliate/nematode system and provide a concrete example of how trait variation at one trophic level can have nonlinear ecological effects on a lower trophic level.

Figure 4. Nonadditive effects of consumer trait variation and consumer competition on prey trajectories

A) f(1) - f(7) are decomposed function components inferred from longitudinal Gaussian process regression for the abundance of *A. caviae* 1972. Components include a microcosm-specific deviation from the shared time effect f(1) and the shared time effect f(2). The remaining components are deviations from the shared time effect and include the low trait diversity ciliate specific effect f(3), the high trait diversity ciliate effect f(4), the nematode effect f(5), an interaction effect for low trait diversity ciliate and nematode f(6), and an interaction effect for high trait diversity ciliate and nematode. Red and dashed lines show when a consumer was excluded or stabilized as in Fig. 3. N.S. indicates that the term explained insufficient variance to be selected in the final model. The sum f(Sum) of the decomposed components is shown in the bottom row. Shading is 2x std. deviations from the posterior mean. Solid lines are posterior means, and points are experimental observations. Term relevances are shaded if combined with noise they explain over 95% of variance of the data.

Overall, the consumer arrangement had an important effect on the abundance trajectories for individual prey species and frequently manifested as non-additive emergent multiple predator effects. Different consumer combinations produced a distinct compositional succession of both rare and abundant prey species relative to the control (Fig. 3), even though bacteria and consumer densities were similar across treatments (Fig. 2). Most prey species displayed multiple predator effects due to the presence of both the nematode and ciliate, but usually, the relevance of one of these interaction terms to the model was significantly larger than the other (Fig. S4). Thus, risk modifying emergent multiple predator effects were universal to this ecological system, but the magnitude and direction of these effects could be modulated at the prey level via consumer trait diversity and competition.

Prey community response to consumer trait variation is mediated through defense traits: We next asked how consumer trait variation and competition influenced community-scale prey responses in the microcosms. We used a joint species distribution model (JSDM) framework[24] to simultaneously

evaluate prey species and community responses to different consumer competitive arrangements. We   217
modeled the sorting and equilibrium phases separately to account for the strong temporal dependence in   218
the sorting phase (supplementary text). The average explanatory power was excellent for the sorting   219
phase ($R^2 = 0.90 \pm 0.09$) due to the strong species changes with time (Fig. 5A). In the equilibrium   220
phase, the average model power was lower ($R^2 = 0.58 \pm 0.19$) but still appreciable. Models of both   221
phases showed the most predictive power through their fixed effects, with random effects contributing   222
largely in the equilibrium phase due to temporal oscillations around the mean (Table S9). Variance   223
partitioning over the explanatory variables included in the models showed that the fixed effects explained   224
a substantial amount of variation in the abundance model. Time and treatment-specific differences   225
accounted for most variation in the sorting phase, which reflected 1) the rapid decline in abundance of   226
the rare species under all treatments and 2) the condition-specific responses of the more abundant   227
species (Fig. 5A). The direction of these effects generally reflected the same patterns we observed from   228
the individual species models. Specifically, ciliate evolution generally strengthened the non-additivity of   229
multiple consumers on prey abundances (supplementary text).   230

Our primary goal with the JSDM was to obtain a mechanistic understanding of the prey community   231
assembly process by identifying specific prey traits associated with the different consumer arrangements.   232
We used the JSDM framework to ask whether phylogenetic relationships and prey traits were associated   233
with specific community responses to the different consumer arrangements. This is important as   234
bacterial taxonomy is an important determinant for colonization success of the *C. elegans* intestine[25],   235
certain bacterial taxonomic classes are associated with different stages of *C. elegans* growth in the   236
wild[26], and prey trait variation affects prey coexistence under *T. thermophila* predation[27]. In the   237
final JSDMs, we included traits for defense, growth rate, diversity of carbon compounds used, and   238
biofilm formation. These particular traits represented approximately 80% of the variance from the total   239
collection of traits acquired for the synthetic community (Fig. S7). The estimated posterior of   240
phylogenetic strength was 0.05 ± 0.11 in sorting and 0.32 ± 0.34 in equilibrium with the 95% credibility   241
interval including zero for both phases. Thus, closely related bacteria were no more likely to share a   242
common response to the experimental consumer arrangements than distant relatives. In contrast to   243
phylogeny, prey traits explained a substantial amount of variation in both experimental phases (Table   244
S10, Fig. 5C,F), indicating a shared phenotypic response of the prey community under each treatment.   245
In the sorting phase, the diversity of carbon compounds used was positively associated with the average   246
community response to the LTV ciliate, which reflected the dominance of *Morganella morganii* 1292 and   247
Pseudomonas chlororaphis 1977 - the two strains that could use the most carbon compounds. However,   248
carbon use versatility was negatively associated with changes in prey abundance with the nematode,   249
which mirrored the higher species diversity supported by the nematode Fig. 5C). Prey defense and   250
biofilm traits were consistently associated with community responses in both the sorting and equilibrium   251
phases. These traits were mostly associated with the community response to the HTV ciliate either alone   252
or through an interaction with the nematode (Fig. 5C,F). Overall traits explained 22% and 44% of the   253
variation in species abundances in the sorting and equilibrium phases, respectively. Traits explained the   254
greatest proportion of the community response to the HTV ciliate and the combination of HTV ciliate   255
and nematode (Table S10).   256

Figure 5. Determinants of prey community assembly

JSDM results from **A-C)** the sorting phase (days 0-13) and **D-F)** the equilibrium phase (days 17-61). **A** and **D** show the explanatory power for each bacterial prey species in the JSDM. For each species, the bars are colored by the proportion of variance explained by fixed (i.e. consumer evolution and/or co-consumer competition, etc) and random effects (time, sequencing effort, etc) and their interactions. **B)** and **E)** show the influence of fixed effect covariates on bacterial prey species. **C)** and **D)** show the influence of prey traits on bacterial prey species association with fixed effects from the JSDM. The sign of the coefficient (i.e the direction of the effect) is brown if positive and green if negative. Coefficients are included only if the 95% credibility interval excludes zero (i.e the probability the effect is not zero is 95%). **D)**, defense against C_{LTV}; N carbon, number of carbon sources the strain can use; r, maximum specific growth rate; biofilm, biofilm production.

The JSDM results were generally consistent with a growth/defense trade-off scenario in the prey community. Rapidly growing, generalist prey species (1977 and 1292) quickly dominated the community in the absence of a predator and with the LTV ciliate even though these species were relatively undefended against the LTV ciliate (Fig. S6). The most defended prey species were less abundant with the HTV ciliate than with the LTV ciliate, presumably because the HTV ciliate was better adapted to consume these prey species[17, 27]. However, biofilm-forming prey species were more abundant with the HTV ciliate, potentially because it was poorly adapted to consuming prey biofilms. Although biofilm formation is one specific defense mechanism against protozoan grazing, bacterial defense against consumers ranges from motility to toxin production to cell surface masking[28]. Surprisingly, defended prey were more abundant than would be predicted from the summed effects of individual consumers in the presence of the HTV ciliate and nematode, while the opposite was true for biofilm-forming prey. In the sorting phase, this pattern also held for the interaction between the HTV ciliate and nematode. This may reflect prey risk enhancement and reduction from predation by multiple consumers[15]. For example, the nematode may have been able to mechanically disrupt prey aggregates while foraging, subsequently rendering these species more vulnerable to the HTV ciliate. Alternatively, nematode foraging behavior may have induced new traits or modulated existing traits in well-defended prey species that also increased resistance to the HTV ciliate. Overall, the model revealed that prey traits were key components mediating the prey community response to both consumer evolution and interspecific competition.

Ciliate trait diversity alters prey preference: We next asked whether the competitive outcome between consumer pairs was correlated with differences in specific consumer traits. We experimentally assayed the ability of each consumer to clear prey biomass in pairwise grazing trials with each of the 24 bacterial strains (Fig. 6A). We defined prey clearance as the ratio of removed bacteria to consumer individuals at the end of a 144-hour incubation for each consumer/prey species combination (see methods). The nematode had the highest prey clearance ($7.6 \pm 5.3 \times 10^4$, observed median and std. dev.), followed by the HTV ciliate ($7.1 \pm 9.0 \times 10^3$), and the LTV ciliate ($2.7 \pm 1.6 \times 10^3$). Indeed, trait variation significantly increased prey clearance by the ciliate, while the nematode was the most efficient consumer overall (Table S11). The coefficient of variation of prey clearance was smallest for the LTV ciliate, then the nematode, and then the HTV ciliate ($CV_{LTV} = 0.59$, $CV_N = 0.70$, $CV_{HTV} = 1.25$), reflecting a range of per species prey clearances for each consumer. Overall, the consumers displayed a hierarchy of prey clearance rates: 1) the LTV ciliate removed the fewest prey cells per capita with the most uniform clearance rates across prey species, 2) trait variation significantly increased ciliate feeding efficiency on some prey species, which increased prey clearance variation overall, and 3) averaged across all prey species the nematode removed the most prey per capita but also grazed on some species more efficiently than others.

We might expect greater relative variation in consumer prey clearance to manifest as more uneven foraging across prey species in a community context. In the short term, a consumer with the most uniform clearance rates should remove each prey species in proportion to their abundance in the community. The consequence would be lower total prey biomass but a prey composition with greater resemblance to the prey community without consumers. Alternatively, a consumer displaying nonuniform prey preferences should remove prey in a way that deviates from this proportionality[29, 30]. This approach assumes limited indirect consumer effects, but this assumption should hold during exponential growth (sorting phase) when resources should not limiting for prey or consumers[31]. We estimated consumer foraging preferences from sequencing data by regressing the relative abundance of bacterial species in the presence and absence of each consumer across all time points in the sorting phase of the experiment. We found that the slope of this relationship was largest for the LTV ciliate(Fig. 6B), consistent with it having the smallest relative variation in clearance across all prey species. Both the slopes for the HTV ciliate and nematode were significantly smaller than the LTV ciliate slope suggesting that these consumers foraged on some higher abundance prey species preferentially (Table S12).

Our results suggest that trait variation in the HTV ciliate increased both per capita prey removal and

Figure 6. Consumer trait variation alters consumer prey preference
A) Distribution of prey clearances on the 24 different bacterial prey species. Prey clearance is the number of
bacterial cells removed per consumer individual after 144 hours. Species (connected by lines) in black/labeled are
those with > 1% mean relative abundance across consumer arrangements. Asterisks show significant differences
in mean prey clearance between consumers (Table S11). B) Relative abundance of prey species in the presence of
ciliate or nematode consumers (vertical axis) versus the corresponding relative abundance in the consumer-free
controls. The regression includes prey species > 1% relative abundance and samples from first 21 days. The
axes are arcsin square-root transformed. β, R^2 is the median for the posterior of the slope and Bayesian R^2,
respectively from linear regression (Table S12).

selective feeding behavior over the LTV ciliate. What physiological differences might account for this
outcome? Ciliate individuals from the HTV populations were, on average, 40 μm larger, 1.25$\times$ faster,
and turned less often than isogenic, LTV ciliate individuals[17]. These traits primarily influence the
"search phase" of prey capture[32] because they influence the search volume covered by each consumer.
We speculate that altered body size and swimming speed account for the greater average prey clearance
of the HTV ciliate relative to the LTV ciliate (Fig. 6A). The average *C. elegans* individual is over 100
times the volume of a *Tetrahymena* cell (BNIDs: 114967, 101424) [33], which likely explains why it
removed about an order of magnitude more prey cells per capita than the ciliate (Fig. 6A). The
physiological mechanisms driving the apparent shift in prey preference of the HTV ciliate are more
difficult to explain. Prior work has demonstrated that both *C. elegans* and *Tetrahymena* ciliates can
selectively forage for optimal prey[34–36]. Ciliate selective foraging occurs through multiple mechanistic
steps ranging from searching to prey handling to prey digestion[32], any of which may have been affected
by trait variation in the HTV ciliate.

Discussion

Many studies (e.g., discussed in [8, 10]) have focused on how the ecological effects of intraspecific trait
variation should promote species coexistence. Our results suggest that trait variation in a single
competitor (here, the LTV ciliate) can result in competitive exclusion. Following coexistence theory, we

estimated that the inversion of the competitive consumer hierarchy was due to the development of ₃₄₈
greater niche overlap between competitors and substantial competitive differences that favored the HTV ₃₄₉
ciliate over the nematode. We confirmed this prediction by measuring prey clearance rates for each ₃₅₀
consumer and morphological trait variation in the HTV and LTV ciliate. We found that 1) the ₃₅₁
nematode consumed the most prey per capita, 2) the HTV ciliate displayed both a broader range of ₃₅₂
clearance rates across prey species and a higher mean prey clearance than the isogenic LTV ciliate, 3) on ₃₅₃
average, the HTV ciliate was larger, faster, and turned less than the LTV ciliate[17] which likely allowed ₃₅₄
HTV individuals to forage across a larger habitat volume, and 4) the HTV ciliate and nematode ₃₅₅
suppressed the growth of the most abundant prey species from the no-consumer control while the LTV ₃₅₆
ciliate could not. We speculate that changes in ciliate morphological traits and prey preference increased ₃₅₇
the relative strength of competition with the nematode, while the convergence of HTV prey preferences ₃₅₈
towards the most abundant prey species also reduced the niche difference. ₃₅₉

Consumer trait variation also strongly influenced the community composition of the prey trophic level. ₃₆₀
The HTV ciliate or the nematode rapidly stabilized the composition of the prey community, while the ₃₆₁
LTV ciliate and no-consumer treatment caused the prey community to make the largest oscillations ₃₆₂
through community state space (Fig. 3C). The combination of two consumers caused the prey ₃₆₃
community to traverse state space in a way that was distinct from either consumer alone. Indeed, we ₃₆₄
found that the combined effects of the ciliate and nematode on individual prey species were often ₃₆₅
non-additive - i.e., the combined effects of the consumers were not simple linear extrapolations of their ₃₆₆
individual effects. Emergent multiple predator effects are well-known for macroscopic predators and their ₃₆₇
prey[15] but have not been documented at the microbial scale. Importantly, we also found that trait ₃₆₈
variation in the ciliate significantly altered the magnitude of non-additive consumer effects across prey ₃₆₉
species. We also found that prey community assembly in the presence of consumers was associated with ₃₇₀
distinct prey traits related to consumer defense and resource use. Thus, prey communities may assemble ₃₇₁
partly due to how consumer trait heterogeneity and consumer competition operate on the distribution of ₃₇₂
defense traits present in a prey community. Consumer trait diversity generally strengthened the ₃₇₃
non-additivity of how prey defense traits influenced prey community assembly in the presence of the ₃₇₄
competing consumer. This suggests that the role of trait variation at one trophic level may have complex ₃₇₅
downstream effects on other trophic levels in the wild. The number and magnitude of these nonlinear ₃₇₆
relationships may be critical feedbacks stabilizing the coexistence of many bacterial prey species in ₃₇₇
natural microbial communities. Our findings also challenge the simple pairwise rules often invoked to ₃₇₈
explain microbial community assembly[37, 38] and suggest that nonlinear interactions across trophic ₃₇₉
levels - for example, driven by consumer trait variation - are important determinants of structure in ₃₈₀
natural, more complex microbial communities[39]. ₃₈₁

Our study leaves many avenues open for future research. One major question is what role prey trait ₃₈₂
variation plays in community dynamics across trophic levels. Future studies should investigate ₃₈₃
community assembly and species coexistence while systematically manipulating trait variation in both ₃₈₄
consumers and prey. Furthermore, we could not examine the role of rapid contemporary evolution during ₃₈₅
the time course of our experiments. This is important because recent theory suggests that the rapid ₃₈₆
evolution of a single prey species can facilitate the coexistence of multiple consumers both by broadening ₃₈₇
niche differences [40] and equalizing consumer competitive differences[41]. Longer experiments coupled ₃₈₈
with genome sequencing will better allow us to characterize the role of rapid evolution during the ₃₈₉
experiment time course for species coexistence. Recent studies have shown consumer co-evolution ₃₉₀
contributes significantly to eco-evolutionary dynamics in one and two species systems[27, 42]. Here we ₃₉₁
show that feedback between consumer competition and consumer trait variation can also add a ₃₉₂
significant layer of complexity to our understanding of species coexistence and the maintenance of ₃₉₃
diversity across trophic levels in more complex communities. ₃₉₄

Materials and Methods

Study species: The prey bacterial community consisted of 24 species (Table S1) from the University of Helsinki Culture Collection. These species are derived from soil, aquatic, plant, animal, and human sources and have been described in detail elsewhere [43]. The consumer species consisted of the ciliate *Tetrahymena thermophila* strain 1630/1U (CCAP), obtained from the Culture Collection of Algae and Protozoa at the Scottish Marine Institute (Oban, Scotland, United Kingdom) and the clonal hermaphrodite nematode *Caenorhabditis elegans* strain N2, obtained from the *Caenorhabditis* Genetics Center (University of Minnesota, Minneapolis, MN, USA). The mixture of HTV genetically diverse ciliates is derived from 20 continuously transferred, clonally reproducing, long-term selection lines. Each of the ciliate lines is from a 600 generation co-evolution experiment initiated with the isogenic ciliate and one of seven different bacterial species (three replicates for each species, one ciliate line lost) with common anti-predatory defense mechanisms[17]. Only one of the bacterial species from the co-evolution experiment (*Comamonas testosterone* 0403) was included in the experimental prey community. These HTV ciliate populations each displayed altered phenotypic traits related to predation, including increased swimming speed, straighter swimming trajectories, and increased body size [17].

Culture conditions and experimental preparations: Bacteria were revived from cryopreserved glycerol stocks (-80°C) in protease peptone yeast (PPY) broth at 28°C with shaking for 24 hours. Individual colonies were picked from cultures plated on 50% PPY agar. Clones were then grown in 5% King's Broth (KB) until mid-log phase and cell densities (colony forming units) determined via plating on 50% PPY agar. Ciliates were revived from liquid nitrogen and cultured axenically at room temperature in PPY as described before[17, 44]. 50 ml of each revived ciliate culture was harvested by centrifugation (8 minutes, 3300 rpm, 4°C), then the cell pellet was resuspended in 25 ml M9 buffer. Ciliates were then counted by microscope using $40\times$ magnification. Cryopreserved (-80°C) nematode stocks were revived following the standard protocol [45]. Briefly, frozen nematodes were spread on NGM-plates with *E. coli* OP50, then individuals transferred to fresh plates by "chunking" after eight days. Eggs were collected from 7-8 plates after six days by bleaching live worms. Developmentally synchronized nematodes were hatched in 125 ml M9 buffer with gentle rotation in tissue flasks overnight. Flasks were chilled on ice, 75 ml of buffer was manually decanted, and the remaining buffer and nematodes were centrifuged (3 mins, 1300 rpm, 8°C). Afterward, half the remaining buffer was decanted, and nematodes were counted directly by microscope.

Experimental setup and sampling: The general experimental design with different consumer treatment combinations is outlined in Fig. 1. Each microcosm was performed in four biological replicates in 150 ml plastic bottles using 20 ml of 5% King's Broth (KB) liquid medium containing M9 salts and supplemented with cholesterol in ethanol (5 μg ml^{-1}) (5% King's B: 1 g L^{-1} peptone number 3 and 0.5 ml L^{-1} 85% glycerol). For each treatment/replicate we inoculated 1×104 cfu ml^{-1} of each 24 bacterial species (total $\approx 2.4 \times 10^5$ cfu ml^{-1}). Consumer densities were inoculated at 1×10^4 ciliate cells ml^{-1} and 10 nematode cells ml^{-1}. Microcosms were maintained at 22°C with gentle shaking. Two ml of each microcosm was transferred to fresh growth medium every four days for a total of 21 transfers over 61 days. One ml was sampled at each transfer, combined with 0.5 ml 85% glycerol, and frozen at -80 °C for later DNA extraction. Another 0.5 ml was sampled at each transfer for quantifying consumer density (ciliate: 1:3 vol/vol with 10% Lugol, nematode: 1:1 vol/vol with 20% Lugol). Nematode cell densities were measured directly under a microscope. Ciliate densities were measured using Fiji/ImageJ with the cell counter plugin[46]. Total bacterial biomass was estimated using optical density (OD$_{600}$) as a proxy.

Bacterial trait measurements and processing: We measured innate predator defense, carrying capacity and growth rate for each bacterial strain as described in detail earlier[27]. Briefly, we revived

each bacterial species in fresh medium (1% Kings Broth), acclimated them 24 hours, then pin replicated ⁴⁴⁰
strains to fresh medium in 100 well plates for use with a Bioscreen C spectrophotometer (Growth Curves ⁴⁴¹
ABLtd, Helsinki, Finland). Plates were incubated at 28°C at constant shaking in the Bioscreen C and ⁴⁴²
measurements were taken at 5 min intervals for 24 hours. After 48 hours, LTV ciliates were added to ⁴⁴³
Bioscreen C wells to estimate biomass loss due to predation. For each bacterial species k we quantified ⁴⁴⁴
growth potential using area under the logistic growth curve (AUC), and estimated innate defense of ⁴⁴⁵
species k as $AUC_{k+consumer}/AUC_k$. We inferred maximum growth rate, the doubling time, the carrying ⁴⁴⁶
capacity using growthcurveR[47]. We determined biofilm formation capacity for each using the ⁴⁴⁷
crystal-violet method[48]. We determined bacterial growth on 31 different carbon substrates using ⁴⁴⁸
BioLog EcoPlates (https://www.biolog.com). We predicted additional traits from bacterial genomes[43] ⁴⁴⁹
using Traitar[49]. ⁴⁵⁰

Consumer feeding efficiency experiments: The 24 bacteria species, LTV ciliate, HTV ciliate, and ⁴⁵¹
nematode stocks were revived from cryopreservation as described above. Each of the bacterial species ⁴⁵²
was grown as monocultures in 100 ml of Reasoner's 2A liquid medium[50] at room temperature for 96 ⁴⁵³
hours with shaking at 50 RPM. Afterward, bacteria were harvested by centrifugation (13000 RPM, 10 ⁴⁵⁴
minutes, 21°C), the supernatant was discarded, and cells were resuspended in 100 ml of M9 saline ⁴⁵⁵
solution. The optical density was then adjusted to $OD_{600} = 0.5$ with additional M9. We added 5 ml of ⁴⁵⁶
suspension from each bacterial species to 6 well culture plates and then added either 2.5×10^3 nematode ⁴⁵⁷
individuals or 2.5×10^4 of HTV or LTV ciliate individuals. One well for each bacterial species was ⁴⁵⁸
reserved as a no-consumer control. All conditions were assayed in biological triplicates. Plates were ⁴⁵⁹
incubated at room temperature with shaking at 50 RPM. After 144 hours of incubation, we measured ⁴⁶⁰
bacterial optical density (OD_{600}) and counted consumers as described above. We defined prey clearance ⁴⁶¹
as the difference between the OD_{600} in each consumer treatment and the no consumer control. We ⁴⁶²
converted optical density to cells ml⁻¹ using a conversion of 8×10^6 bacterial cells per 0.01 OD_{600} unit. ⁴⁶³
We defined prey clearance as the number of prey cells consumed per consumer individual after 144 hours. ⁴⁶⁴

DNA extraction and sequencing: Total DNA was extracted using the DNeasy Blood & Tissue ⁴⁶⁵
96-well extraction kit (Qiagen) according to the manufacturer's instructions. The V3-V4 region of the ⁴⁶⁶
bacterial 16S rRNA was amplified using a primer set with Illumina sequencing adapters and multiplexing ⁴⁶⁷
barcodes. Samples were sequenced on an Illumina MiSeq using v3 600 cycles reagent kit. The procedure ⁴⁶⁸
is outlined in detail in earlier studies[43, 51]. ⁴⁶⁹

DNA sequence analysis: Quality control and processing were performed using BBTools (version ⁴⁷⁰
38.61b (https://sourceforge.net/projects/bbmap/). 16S amplicon read pairs were processed using BBDuk ⁴⁷¹
to remove contaminants, trim reads that contained adapter sequence, and right quality trim reads where ⁴⁷²
quality drops below Q10. BBMerge [52] was used to merge surviving read pairs into single amplicon ⁴⁷³
sequences, and `msa.sh` and `cutprimers.sh` were used to remove any forward and reverse primer ⁴⁷⁴
sequences. VSEARCH [53] was used to filter out amplicons with more than two expected errors ⁴⁷⁵
(`-fastq_maxee 2`)[54] and excluded sequences outside of the 360-480 bp range. Quality controlled ⁴⁷⁶
amplicon sequences were assigned to bacterial species using BBMap by mapping against a database of 30 ⁴⁷⁷
full-length 16S rRNA sequences using the best possible mapping position. Only amplicons mapping ⁴⁷⁸
unambiguously between position 341 to 805 in the aligned 16S rRNA region with at least a 95% mapping ⁴⁷⁹
identity were retained and assigned to a species for subsequent analysis. Nearly all reads from each ⁴⁸⁰
sample were perfectly mapped to a single species 16S rRNA sequence using these criteria, and the ⁴⁸¹
greatest number of unmapped reads was 0.1% of the total. ⁴⁸²

Statistics and data analysis: Detailed descriptions of the statistics and data analysis procedures are ⁴⁸³
available in the supplementary text and `https://github.com/slhogle/consumer-competition`. All ⁴⁸⁴

statistical analyses were done in R version 3.6.1 (2019-07-05)[55]. Probabilistic modeling was performed ₄₈₅ using the Stan programming language v2.24[56] and the RStan v2.21.1 R interface to Stan. We assessed ₄₈₆ MCMC chain mixing and convergence using effective sample sizes [57] and potential scale reduction ₄₈₇ factors ($\hat{R} < 1.05$ for all covariates)[58]. For all Bayesian analyses, we used default priors. ₄₈₈

Ciliate, nematode and bacterial densities were modeled with hierarchical generalized additive models ₄₈₉ using mgcv v1.8-33[59]. All other regression was performed in the Bayesian framework using rstanarm ₄₉₀ v2.21-1 [60] or MCPv0.3-0[61]. Posterior distributions were summarized using BayestestR v0.8-2[62]. We ₄₉₁ estimated differences in consumer competitive ability and niche overlap by parameterizing ₄₉₂ phenomenological Lotka-Volterra competition models using gauseR v1.0[63]. We then used the inferred ₄₉₃ per capita growth rates, intraspecific, interspecific coefficients to calculate competitive and niche ₄₉₄ differences[18]. We estimated the compositional bias of amplicon-derived species abundances using ₄₉₅ metacal v0.1.0[64] using abundance information from control samples (day 0) where all prey species were ₄₉₆ inoculated at equal cell densities based on colony counts. We calculated the Shannon diversity ₄₉₇ dissimilarity index following Jasinska[65]. We estimated population level alpha and beta diversities ₄₉₈ (Shannon diversity and Bray Curtis dissimilarity) over the four biological replicates per treatment using ₄₉₉ DivNet v0.3.6 [66]. To identity sorting and equilibrium phases of the experiment, we performed ₅₀₀ segmented regressions of Shannon diversity using MCP v0.3-0[61]. We also calculated Beta diversity for ₅₀₁ individual replicates using Bray and Curtis dissimilarity on species relative abundances. We used ₅₀₂ non-metric multidimensional scaling for ordination. ₅₀₃

For modeling absolute species abundances with Gaussian Process Regression, we transformed calibrated ₅₀₄ species count matrices to the log scale using the regularized logarithm transform from DESeq2 v1.28.1 ₅₀₅ [67]. This transformation moderates variance across the mean abundance for each species and normalizes ₅₀₆ with respect to library size. We modeled the transformed abundance of each species using a longitudinal ₅₀₇ Gaussian process regression model implemented in lpgr v0.33.3 [68]. To infer whether covariates were ₅₀₈ informative for the models, we used a probabilistic covariate selection method which estimates the ₅₀₉ proportion of variance explained for each covariate and noise and selects those covariates that alongside ₅₁₀ noise explain 95% of the variance. ₅₁₁

For joint species distribution modeling[69], we used the Hierarchical Modelling of Species Communities ₅₁₂ (HMSC) package[70, 71]. We used a two-step hurdle model approach to account for zero inflation in the ₅₁₃ log transformed count data where we first fit a probit model for presence/absence and then a Gaussian ₅₁₄ model for regularized log-transformed species abundances conditional upon presence. We accounted for ₅₁₅ differences in sequencing depth by including log-transformed total reads as a fixed effect in the model. ₅₁₆ We followed the approach from earlier studies using sequencing count data with HMSC[70–72]. ₅₁₇

Data availability: ₅₁₈

Raw sequencing data is available from the NCBI Sequence Read Archive (SRA) under the BioProject ₅₁₉ accession number PRJNA725120. Preprocessed count tables are available from ₅₂₀ https://github.com/slhogle/consumer-competition. ₅₂₁

Code availability: ₅₂₂

Scripts reproducing all figures and steps in the data analysis are available from ₅₂₃ https://github.com/slhogle/consumer-competition. ₅₂₄

Acknowledgments

Elizaveta Zakharova: investigation - supporting, data curation. Olli Pitkänen: investigation - supporting, data curation. Ida-Marija Hyvönen investigation - supporting, data curation. Funding: Academy of Finland (grants #327741 to TH and #286405 to LR)

Author contributions:

SLH: conceptualization, supervision, software, formal analysis, writing - original draft. IH: conceptualization, investigation - lead, visualization, writing - review & editing. LR: supervision, formal analysis, funding acquisition, writing - review & editing. JC: conceptualization, writing - review & editing. TH: conceptualization, supervision - lead, project administration, funding acquisition, writing - review & editing. All authors provide approval for publication.

References

1. Siefert, A. *et al.* A global meta-analysis of the relative extent of intraspecific trait variation in plant communities. en. *Ecol. Lett.* **18,** 1406–1419 (Dec. 2015).

2. Des Roches, S. *et al.* The ecological importance of intraspecific variation. en. *Nat Ecol Evol* **2,** 57–64 (Jan. 2018).

3. Slatkin, M. Ecological character displacement. *Ecology* **61,** 163–177 (Feb. 1980).

4. Chesson, P. Mechanisms of Maintenance of Species Diversity. en. *Annu. Rev. Ecol. Syst.* **31,** 343–366 (Nov. 2000).

5. Lichstein, J. W., Dushoff, J., Levin, S. A. & Pacala, S. W. Intraspecific variation and species coexistence. en. *Am. Nat.* **170,** 807–818 (Dec. 2007).

6. Hart, S. P., Schreiber, S. J. & Levine, J. M. How variation between individuals affects species coexistence. en. *Ecol. Lett.* **19,** 825–838 (Aug. 2016).

7. Clark, J. S. Individuals and the variation needed for high species diversity in forest trees. en. *Science* **327,** 1129–1132 (Feb. 2010).

8. Bolnick, D. I. *et al.* Why intraspecific trait variation matters in community ecology. en. *Trends Ecol. Evol.* **26,** 183–192 (Apr. 2011).

9. Maynard, D. S., Serván, C. A., Capitán, J. A. & Allesina, S. Phenotypic variability promotes diversity and stability in competitive communities. en. *Ecol. Lett.* **22,** 1776–1786 (Nov. 2019).

10. Turcotte, M. M. & Levine, J. M. Phenotypic Plasticity and Species Coexistence. en. *Trends Ecol. Evol.* **31,** 803–813 (Oct. 2016).

11. Hausch, S., Vamosi, S. M. & Fox, J. W. Effects of intraspecific phenotypic variation on species coexistence. en. *Ecology* **99,** 1453–1462 (June 2018).

12. Fischer, B. B. *et al.* Phenotypic plasticity influences the eco-evolutionary dynamics of a predator-prey system. en. *Ecology* **95,** 3080–3092 (Nov. 2014).

13. Palkovacs, E. P. & Post, D. M. Experimental evidence that phenotypic divergence in predators drives community divergence in prey. en. *Ecology* **90,** 300–305 (Feb. 2009).

14. Chislock, M. F., Sarnelle, O., Olsen, B. K., Doster, E. & Wilson, A. E. Large effects of consumer offense on ecosystem structure and function. en. *Ecology* **94,** 2375–2380 (Nov. 2013).

15. Sih, A., Englund, G. & Wooster, D. Emergent impacts of multiple predators on prey. en. *Trends Ecol. Evol.* **13,** 350–355 (Sept. 1998).

16. Seibold, S., Cadotte, M. W., MacIvor, J. S., Thorn, S. & Müller, J. The Necessity of Multitrophic Approaches in Community Ecology. en. *Trends Ecol. Evol.* **33,** 754–764 (Oct. 2018).

17. Cairns, J., Moerman, F., Fronhofer, E. A., Altermatt, F. & Hiltunen, T. Evolution in interacting species alters predator life-history traits, behaviour and morphology in experimental microbial communities. en. *Proc. Biol. Sci.* **287,** 20200652 (June 2020).

18. Letten, A. D., Hall, A. R. & Levine, J. M. Using ecological coexistence theory to understand antibiotic resistance and microbial competition. en. *Nat Ecol Evol* (Feb. 2021).

19. McGill, B. J. *et al.* Species abundance distributions: moving beyond single prediction theories to integration within an ecological framework. en. *Ecol. Lett.* **10,** 995–1015 (Oct. 2007).

20. Gonze, D., Coyte, K. Z., Lahti, L. & Faust, K. Microbial communities as dynamical systems. en. *Curr. Opin. Microbiol.* **44,** 41–49 (Aug. 2018).

21. Vonesh, J. R. & Osenberg, C. W. Multi-predator effects across life-history stages: non-additivity of egg- and larval-stage predation in an African treefrog. en. *Ecol. Lett.* **6,** 503–508 (June 2003).

22. Anderson, T. L. & Whiteman, H. H. Non-additive effects of intra- and interspecific competition between two larval salamanders. en. *J. Anim. Ecol.* **84,** 765–772 (May 2015).

23. Timonen, J., Mannerström, H., Vehtari, A. & Lähdesmäki, H. lgpr: An interpretable nonparametric method for inferring covariate effects from longitudinal Data. *Bioinformatics* (Jan. 2021).

24. Ovaskainen, O. *et al.* How to make more out of community data? A conceptual framework and its implementation as models and software. en. *Ecol. Lett.* **20,** 561–576 (May 2017).

25. Ortiz, A., Vega, N. M., Ratzke, C. & Gore, J. Interspecies bacterial competition regulates community assembly in the C. elegans intestine. *ISME J.* (Feb. 2021).

26. Samuel, B. S., Rowedder, H., Braendle, C., Félix, M.-A. & Ruvkun, G. Caenorhabditis elegans responses to bacteria from its natural habitats. en. *Proc. Natl. Acad. Sci. U. S. A.* **113,** E3941–9 (July 2016).

27. Scheuerl, T., Cairns, J., Becks, L. & Hiltunen, T. Predator coevolution and prey trait variability determine species coexistence. en. *Proc. Biol. Sci.* **286,** 20190245 (May 2019).

28. Matz, C. & Kjelleberg, S. Off the hook–how bacteria survive protozoan grazing. en. *Trends Microbiol.* **13,** 302–307 (July 2005).

29. Bell, T. *et al.* Protists have divergent effects on bacterial diversity along a productivity gradient. en. *Biol. Lett.* **6,** 639–642 (Oct. 2010).

30. Saleem, M., Fetzer, I., Harms, H. & Chatzinotas, A. Diversity of protists and bacteria determines predation performance and stability. en. *ISME J.* **7,** 1912–1921 (Oct. 2013).

31. Wootton, J. T. The nature and consequences of indirect effects in ecological communities. *Annu. Rev. Ecol. Syst.* **25,** 443–466 (Nov. 1994).

32. Montagnes, D. J. S. *et al.* Selective feeding behaviour of key free-living protists: avenues for continued study. *Aquat. Microb. Ecol.* **53,** 83–98 (Sept. 2008).

33. Milo, R., Jorgensen, P., Moran, U., Weber, G. & Springer, M. BioNumbers - the database of key numbers in molecular and cell biology. en. *Nucleic Acids Res.* **38,** D750–3 (Jan. 2010).

34. Shtonda, B. B. & Avery, L. Dietary choice behavior in Caenorhabditis elegans. en. *J. Exp. Biol.* **209,** 89–102 (Jan. 2006).

35. Thurman, J., Parry, J. D., Hill, P. J. & Laybourn-Parry, J. The filter-feeding ciliates Colpidium striatum and Tetrahymena pyriformis display selective feeding behaviours in the presence of mixed, equally-sized, bacterial prey. en. *Protist* **161,** 577–588 (Oct. 2010).

36. Dopheide, A., Lear, G., Stott, R. & Lewis, G. Preferential feeding by the ciliates Chilodonella and Tetrahymena spp. and effects of these protozoa on bacterial biofilm structure and composition. en. *Appl. Environ. Microbiol.* **77,** 4564–4572 (July 2011).

37. Friedman, J., Higgins, L. M. & Gore, J. Community structure follows simple assembly rules in microbial microcosms. en. *Nat Ecol Evol* **1,** 109 (Mar. 2017).

38. Kehe, J. *et al. Positive interactions are common among culturable bacteria* en. June 2020.

39. Gralka, M., Szabo, R., Stocker, R. & Cordero, O. X. Trophic Interactions and the Drivers of Microbial Community Assembly. en. *Curr. Biol.* **30,** R1176–R1188 (Oct. 2020).

40. Klauschies, T., Vasseur, D. A. & Gaedke, U. Trait adaptation promotes species coexistence in diverse predator and prey communities. en. *Ecol. Evol.* **6,** 4141–4159 (June 2016).

41. Van Velzen, E. Predator coexistence through emergent fitness equalization. en. *Ecology* **101,** e02995 (May 2020).

42. Hiltunen, T. & Becks, L. Consumer co-evolution as an important component of the eco-evolutionary feedback. en. *Nat. Commun.* **5,** 5226 (Oct. 2014).

43. Cairns, J. *et al.* Construction and Characterization of Synthetic Bacterial Community for Experimental Ecology and Evolution. en. *Front. Genet.* **9,** 312 (Aug. 2018).

44. Cassidy-Hanley, D. M. in *Methods in Cell Biology* (ed Collins, K.) 237–276 (Academic Press, Jan. 2012).

45. Stiernagle, T. in *WormBook* (ed The C. elegans Research Community) (WormBook, Feb. 2006).

46. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. en. *Nat. Methods* **9,** 676–682 (June 2012).

47. Sprouffske, K. & Wagner, A. Growthcurver: an R package for obtaining interpretable metrics from microbial growth curves. en. *BMC Bioinformatics* **17,** 172 (Apr. 2016).

48. O'Toole, G. A. & Kolter, R. Initiation of biofilm formation in Pseudomonas fluorescens WCS365 proceeds via multiple, convergent signalling pathways: a genetic analysis. en. *Mol. Microbiol.* **28,** 449–461 (May 1998).

49. Weimann, A. *et al.* From Genomes to Phenotypes: Traitar, the Microbial Trait Analyzer. en. *mSystems* **1** (Nov. 2016).

50. Reasoner, D. J. & Geldreich, E. E. A new medium for the enumeration and subculture of bacteria from potable water. en. *Appl. Environ. Microbiol.* **49,** 1–7 (Jan. 1985).

51. Herlemann, D. P. *et al.* Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea. *ISME J.* **5,** 1571–1579 (Oct. 2011).

52. Bushnell, B., Rood, J. & Singer, E. BBMerge - Accurate paired shotgun read merging via overlap. en. *PLoS One* **12,** e0185056 (Oct. 2017).

53. Rognes, T., Flouri, T., Nichols, B., Quince, C. & Mahé, F. VSEARCH: a versatile open source tool for metagenomics. en. *PeerJ* **4,** e2584 (Oct. 2016).

54. Edgar, R. C. & Flyvbjerg, H. Error filtering, pair assembly and error correction for next-generation sequencing reads. *Bioinformatics* **31,** 3476–3482 (Nov. 2015).

55. R Core Team. *R: A Language and Environment for Statistical Computing* Vienna, Austria, 2019.

56. Carpenter, B. *et al.* Stan: A Probabilistic Programming Language. *Journal of Statistical Software, Articles* **76,** 1–32 (2017).

57. Brooks, S. P. & Gelman, A. General Methods for Monitoring Convergence of Iterative Simulations. *J. Comput. Graph. Stat.* **7,** 434–455 (Dec. 1998).

58. Gelman, A. & Rubin, D. B. Inference from Iterative Simulation Using Multiple Sequences. en. *Stat. Sci.* **7,** 457–472 (Nov. 1992).

59. Wood, S. N. *Generalized Additive Models: An Introduction with R* 2017.

60. Goodrich, B., Gabry, J., Ali, I. & Brilleman, S. *rstanarm: Bayesian applied regression modeling via Stan* 2020.

61. Lindeløv, J. K. *Mcp: An R package for regression with multiple change points* Jan. 2020.

62. Makowski, D., Ben-Shachar, M. & Lüdecke, D. BayestestR: Describing effects and their uncertainty, existence and significance within the Bayesian framework. *J. Open Source Softw.* **4,** 1541 (Aug. 2019).

63. Mühlbauer, L. K., Schulze, M., Harpole, W. S. & Clark, A. T. gauseR: Simple methods for fitting Lotka-Volterra models describing Gause's "Struggle for Existence". en. *Ecol. Evol.* **10,** 13275–13283 (Dec. 2020).

64. McLaren, M. R., Willis, A. D. & Callahan, B. J. Consistent and correctable bias in metagenomic sequencing experiments. en. *Elife* **8** (Sept. 2019).

65. Jasinska, W. *et al.* Chromosomal barcoding of E. coli populations reveals lineage diversity dynamics at high resolution. en. *Nat Ecol Evol* **4,** 437–452 (Mar. 2020).

66. Willis, A. D. & Martin, B. D. Estimating diversity in networked ecological communities. en. *Biostatistics* (May 2020).

67. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15,** 550 (2014).

68. Timonen, J., Mannerström, H., Vehtari, A. & Lähdesmäki, H. An interpretable probabilistic machine learning method for heterogeneous longitudinal studies. arXiv: `1912.03549 [stat.ML]` (Dec. 2019).

69. Warton, D. I. *et al.* So Many Variables: Joint Modeling in Community Ecology. en. *Trends Ecol. Evol.* **30,** 766–779 (Dec. 2015).

70. Ovaskainen, O. & Abrego, N. *Joint Species Distribution Modelling: With Applications in R* (Cambridge University Press, Cambridge, 2020).

71. Minard, G., Tikhonov, G., Ovaskainen, O. & Saastamoinen, M. The microbiome of the Melitaea cinxia butterfly shows marked variation but is only little explained by the traits of the butterfly or its host plant. en. *Environ. Microbiol.* **21,** 4253–4269 (Nov. 2019).

72. Abrego, N. *et al.* Fungal communities decline with urbanization-more in air than in soil. en. *ISME J.* **14,** 2806–2815 (Nov. 2020).