The genome sequence of the avian vampire fly (*Philornis downsi*), an invasive nest

parasite of Darwin's finches in Galápagos

Melia Romine[1], Sarah A. Knutie[2,3], Carly M. Crow[4], Grace J. Vaziri[2], Jaime Chaves[5,6],

Jennifer A.H. Koop[4*] & Sangeet Lamichhaney[1,7*#]

[1]School of Biomedical Sciences, Kent State University, Kent, OH, USA

[2]Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs, CT USA

[3]Institute for Systems Genomics, University of Connecticut, Storrs, CT USA

[4]Department of Biological Sciences, Northern Illinois University, DeKalb, IL, USA

[5]Department of Biology, San Francisco State University, San Francisco, CA, USA

[6]Colegio de Ciencias Biológicas y Ambientales, Universidad San Francisco de Quito, Quito,

Ecuador

[7]Department of Biological Sciences, Kent State University, Kent, OH, USA

[*]co-senior authors

[#]corresponding author (slamichh@kent.edu)

Abstract

The invasive avian vampire fly (*Philornis downsi*) is considered one of the greatest threats to the unique and endemic avifauna of the Galápagos Islands, Ecuador. The fly parasitizes nearly every passerine species, including Darwin's finches, in the Galápagos. The fly is thought to have been introduced from mainland Ecuador, although the full pathway of invasion is not yet known. The majority of research to date has focused on the effects of the fly on the fitness of avian hosts and explorations of mitigation methods. A lag in research related to the genetics of this invasion demonstrates, in part, a need to develop full-scale genomic resources with which to address further questions within this system. In this study, an adult *P. downsi* collected from San Cristobal Island within the Galápagos archipelago was sequenced to generate a high-quality genome assembly. We examined various features of the genome (e.g., coding regions, non-coding transposable elements) and carried out comparative genomics analysis against other dipteran genomes. We identified lists of gene families that are significantly expanding/contracting in *P. downsi* that are related to insecticide resistance, detoxification, and potential feeding ecology and counter defense against host immune responses. The *P. downsi* genome assembly provides an important foundational resource for studying the molecular basis of its successful invasion in the Galápagos and the dynamics of its population across multiple islands. The findings of significantly changing gene families associated with insecticide resistance and immune responses highlight the need for further investigations into the role of different gene families in aiding the fly's successful invasion. Furthermore, this genomic resource will also better help inform future research studies and mitigation strategies aimed at minimizing the fly's impact on the birds of the Galápagos.

Keywords: Avian vampire fly genome, parasitic invasion, Galápagos, insecticide resistance, Darwin's finches

Introduction

The invasive avian vampire fly (*Philornis downsi*) (Figure 1a) is considered among the greatest threats to the unique and endemic avifauna of the Galápagos Islands, Ecuador (Causton *et al.*, 2013). The fly parasitizes eleven species of Darwin's finches (Figure 1b) as well as nearly every other Galápagos passerine species (Causton *et al.*, 2013; Fessl *et al.*, 2018). The earliest record of adult flies from the Galápagos occurred in 1964, but the fly was not observed in the nests of birds until 1997 (Fessl and Tebbich, 2002; Causton *et al.*, 2006). The fly has now been found on all major islands within the archipelago except Española, Genovesa, Darwin, and Wolf (Causton *et al.*, 2013). The impact of parasitism by the fly has been severe in some populations of birds in Galápagos and while the effects are variable, some studies have reported near-total nest failure rates due to parasitism (Dudaniec *et al.*, 2007; Koop *et al.*, 2011; Koop, Le Bohec, *et al.*, 2013; O'Connor *et al.*, 2014; Knutie *et al.*, 2016a; Heimpel *et al.*, 2017; Addesso *et al.*, 2020). The fly has also been implicated in the decline of the medium tree finch (*Camarhyncus pauper*), the warbler finch (*Certhidia olivacea*), and the mangrove finch (*Camarhyncus heliobates*) (Dvorak *et al.*, 2004; Grant *et al.*, 2005; Cunninghame *et al.*, 2017; Peters and Kleindorfer, 2018; Bulgarella *et al.*, 2019). Furthermore, the potential for population-level declines of even relatively prominent bird species, e.g., the medium ground finch, have also been demonstrated using predictive models (Koop *et al.*, 2015).

The genus *Philornis* includes approximately 50 species found primarily in the Neotropics and into North America (Dodge, 1955, 1963; Spalding *et al.*, 2002; Dudaniec and Kleindorfer, 2006; Couri *et al.*, 2007). Of the described species within *Philornis*, only two species (including *P. downsi*) are free-living ectoparasites within the nests of their hosts (Fessl *et al.*, 2006), while most others are subcutaneous on their hosts. Several *Philornis* species are found in mainland

Ecuador (Bulgarella *et al.*, 2015, 2017), but *P. downsi* remains the only recorded species of the

genus present, to date, in the Galápagos. *P. downsi* is thought to have been introduced from

mainland Ecuador, though the full pathway of invasion is not yet known (Fessl *et al.*, 2018).

Preliminary population genetics studies show that flies within the archipelago have a high degree

of relatedness relative to those on the mainland, which supports the hypothesis of a relatively

recent invasion and also the possibility of continued gene flow between populations in the

Galápagos (Dudaniec *et al.*, 2008; Koop *et al.*, 2021).

A collaborative research effort has been made to continue monitoring the effects of the avian

vampire fly on bird populations in the Galápagos. These efforts aim to identify source

population(s), the pathway and mechanism of invasion, and possible long-term mitigation

methods, including the Sterile Insect Technique and biocontrol (Causton *et al.*, 2013, 2019).

Current stopgap mitigation efforts in the Galápagos include the direct application of insecticide

to bird nests. Newly hatched nestlings are removed, and a permethrin solution is sprayed inside

the nest and allowed to dry, at which point the nestlings are placed back in the nest. This method

has been shown to effectively reduce parasites in the nest and increase fledging success (Koop *et*

al., 2011). A continuation of this method relies on birds incorporating permethrin-treated cotton

into their nests, achieving similar increases in fledging success (Knutie *et al.*, 2014). Both

methods, with subtle variations, are being used in attempts to increase the nesting success of the

critically endangered mangrove finch, currently one of the most endangered birds in the world

(Fessl *et al.*, 2018). The same method is also being used successfully to protect endangered bird

species against their respective parasites in Australia (Alves *et al.*, 2020).

Despite the robust number of studies that explored the effects of these flies on hosts in the

Galápagos, questions remain about the underlying ecological and evolutionary mechanisms of

their successful invasion in the Galápagos. This knowledge gap demonstrates the need to develop

a full-scale genomic resource of the fly as it provides a critical knowledge base from which to

explore these questions, similar to other parasites of concern (Scott *et al.*, 2020). In this study,

we generated a high-quality draft genome of the avian vampire fly, which is expected to become

an important resource of future molecular studies in this system. We further carried out

comparative genomics analysis with additional published dipteran genomes that identified

evidence of significantly changing gene families associated with insecticide resistance,

detoxification, and possible counter defense against host immune responses. These results serve

as the first step toward investigations of this fly's ability to rapidly evolve traits associated with

its successful invasion in the Galápagos. From an applied aspect, these genomic resources will

also help inform future research studies and mitigation strategies aimed at minimizing the fly's

impact on the birds of the Galápagos.

Materials and Methods

Sampling and DNA extraction for genome sequencing

Flies were collected in Jardín de las Opuntias on San Cristobal Island, Galápagos,

Ecuador (-0.9491651°, -89.5528454°) in March-April of 2019. Adult flies were reared from

pupae collected in the nests of small ground finches (*Geospiza fuliginosa*). When the nests were

empty because nestlings died or fledged, all larvae and pupae were extracted from the nest and

placed in a falcon tube with a modified lid that allowed airflow. After flies eclosed from their

pupal case, they were placed in the freezer, then preserved in 95% ethanol. Preserved flies were

transported to the University of Connecticut, then shipped to Northern Illinois University for

further processing. DNA was extracted using DNeasy Blood and Tissue Kit (Qiagen, Valencia,

California, USA) on whole fly samples after wings and legs were removed using forceps. All

samples were treated with Monarch® RNase A (New England Biolabs) to remove RNA from

genomic DNA samples. Multiple samples were extracted and those with low yields of DNA or

contamination were discarded from further processing. Quantification of double-stranded DNA

was done using QuBit® and samples were run on a gel to assess fragmentation.

Estimate of genome size and ploidy

We prepared a 10X Chromium GEM library from the extracted DNA above according to

the manufacturer's recommended protocols. The library was sequenced using the Illumina 10X

platform to generate paired-end 150 bp reads. We used a k-mer based approach to estimate

genome size, heterozygosity, and repeat content from unprocessed short sequencing reads

(Vurture *et al.*, 2017). We also used the k-mer distribution to extract heterozygous k-mer pairs to

assess the ploidy level in *P. downsi* (Ranallo-Benavidez *et al.*, 2020). These estimates of genome

size and ploidy were then used to choose parameters for the downstream genome assembly

pipeline.

Genome assembly and annotation

The 10x Chromium linked reads generated above were used to generate a reference

genome using Supernova.v.2.1.1 (Weisenfeld *et al.*, 2017) with default parameters. The genome

contiguity statistics such as scaffold N_{50} and the total number of scaffolds etc. were calculated

using custom Perl scripts. We further compared the draft genome assembly against a set of

conserved genes in Diptera using BUSCO (Waterhouse *et al.*, 2017) to assess gene-space

completeness. We also mapped the short sequencing reads back to the draft genome assembly

using BWA (Li and Durbin, 2009) to estimate how many short-sequence reads were used to

build the reference genome and infer genome completeness. The annotation of the *P. downsi*

genome was done using the MAKER annotation pipeline (Cantarel *et al.*, 2008). We combined

protein homology evidence (using publicly available protein datasets from major dipteran

lineages, downloaded from the Ensemble database (Cunningham *et al.*, 2019) and *ab-initio* gene

predictions models to annotate the genome.

Transposable elements (TE) in *P. downsi*

We combined homology-based and de novo approaches using Repeatmasker (Smit

Hubley P., 2013; Flynn *et al.*, 2019) to characterize TEs in *P. downsi*. We used two different sets

of repeat libraries, (a) reference repeat library downloaded from the Repbase database (Bao *et*

al., 2015) and (b) *de novo* custom-built species-specific repeat library generated for *P. downsi*

using RepeatModeler (Flynn *et al.*, 2019). The usage of the *de novo* species-specific repeat

library increased the accuracy of detection and annotation of transposable elements. For an

unbiased comparison of repeat landscapes among all species, we used similar approaches to

detect TEs in the Housefly *(M. domestica)*, Stable fly *(S. calcitrans)*, Tsetse fly *(G. morsitans)*,

Mediterranean fruit fly *(C. capitata)*, Fruit fly *(D. melanogaster)* and Yellow fever mosquito *(A.*

aegypti), together with an outgroup, Postman butterfly *(H. melpomene)*.

Orthologs to other dipteran genomes

We inferred orthogroups and orthologs by comparative analysis of proteomes in *P.*

downsi and six additional dipteran species (*M. domestica, S. calcitrans, G. morsitans, C.*

capitata, D. melanogaster, and A. aegypti), together with an outgroup, *H melpomene.* This

analysis used the draft proteome of *P. downsi* generated from our genome annotation pipeline

described above. Complete proteomes of each of the remaining six species were downloaded

from the Ensemble database (Cunningham *et al.*, 2019). The orthologs inference was carried out

using Orthofinder (Emms and Kelly, 2015).

Gene family evolution

To explore the evolution of gene families in *P. downsi* to other Dipterans, we first

constructed a maximum likelihood phylogeny using 3,070 single-copy orthologs among these

eight species used for inferring orthologues (Figure 3a). We then used this phylogeny to analyze

the changes in gene family size across lineages leading to each of these eight species using a

maximum likelihood approach (CAFÉ) (De Bie *et al.*, 2006) that uses a birth and death process

to model gene gain and loss across a phylogenetic tree.

Results

Genome sequencing and assembly

We sampled an adult *P. downsi* fly in Jardín de las Opuntias on San Cristobal Island,

Galápagos. DNA library was prepared using 10X Chromium linked-read approach (Zheng *et al.*,

2016) and sequenced using Illumina 10X platform to generate paired-end 150 bp reads which

resulted in ~479 million read pairs (~72 Gb of raw sequence data). We first generated a k-mer

distribution based on these short-sequencing reads for preliminary characterization of genome

structure in *P. downsi* (e.g., genome size, an abundance of repetitive elements, rate of

heterozygosity) (Supplementary Table 1, Supplementary Figure 1) that further allowed us to

make informed decisions on parameters needed for building a reference genome. We also

utilized the k-mer distribution to estimate the ploidy level in *P. downsi,* which indicated it to be a

diploid species (Supplementary Figure 2).

Based on genome size estimates using k-mer distributions from short sequence reads

(Supplementary Table 1), we had ~ 91X sequence coverage for generating the draft *de-novo*

genome assembly. The total estimated length of the draft assembly was 971.6 Mb. The assembly

contained 41,176 total scaffolds (minimum 1000 bp to maximum 8.6 Mb) with scaffold N_{50} of

1.3 Mb. We further assessed genome contiguity and gene-space completeness using

benchmarking universal single-copy orthologs (BUSCO) (Waterhouse *et al.*, 2017). Among

3,258 genes highly conserved across Diptera, 3,147 (95.8%) full-length genes were detected in

the *P. downsi* draft genome assembly. Partial sequences of 71 genes were identified (2.2%) and

only 67 (2.0%) were missing, indicating a high degree of contiguity in the *P. downsi* genome. In

addition, 99.16% of total short sequencing reads aligned back to the draft genome indicating a

high degree of genome completeness as well.

Genome annotation

We combined protein homology-based evidence and *ab-initio* gene prediction models

using the MAKER genome annotation pipeline (Cantarel *et al.*, 2008) which identified 15,774

protein-coding genes in the *P. downsi* genome. We compared the genome annotation of *P.*

downsi with two other published and highly curated dipteran genomes, *Musca domestica* (Scott

et al., 2014) and *D. melanogaster* (Adams *et al.*, 2000). We annotated fewer genes in the *P.*

downsi genome (15,774) compared to that of *M. domestica* (17,283) or *D. melanogaster* (17,468)

(Table 1). The average length of genes is similar among *P. downsi* and *D. melanogaster,* whereas

M. domestica typically has longer genes. The mean number of exons per gene is fewer in *P.*

downsi compared to the other two species.

Table 1: Statistics of genomic features among three fly genomes

Genome statistics	P. downsi	M. domestica (Scott et al. 2014)	D. melanogaster (Adams et al. 2000)
Genome size (Gb)	0.97	0.69	0.18
Total genes	15,774	17,283	17,468
Avg. gene length (bp)	4,789	13,747	5,830
Total exons	56,595	123,831	188,405
Avg. exon length (bp)	367	453	485
Number of Exons per gene	3.59	7.16	10.79

Transposable elements in *P. downsi*

Mobile transposable elements (TE) are key features of eukaryotic genomes, being major

determinants of genome size variation (Kapusta *et al.*, 2017; Lamichhaney *et al.*, 2021), and

important contributors to the evolutionary potential of an organism (Pourrajab and

Hekmatimoghaddam, 2021). We characterized the transposable elements in the *P. downsi*

genome using homology-based (Smit Hubley P., 2013) and *de-novo* approaches (Flynn *et al.*,

2019). More than half of the genome (51.7%) of *P. downsi* consists of transposable elements

(Supplementary Table 2). Among these sequences, 9.3% of the genome are retroelements (7.7%

LINEs and 1.6% LTRs) and 23.4% DNA transposons. Short interspersed nuclear elements

(SINEs), a major category of retroelements, were not detected in the *P. downsi* genome.

We also used similar methods to detect, annotate, and compare the repeat content across

several other species including the house fly *(Musca domestica)*, stable fly *(Stomoxys calcitrans)*,

tsetse fly *(Glossina morsitans)*, Mediterranean fruit fly *(Ceratitis capitata)*, fruit fly *(Drosophila*

melanogaster) and yellow fever mosquito *(Aedes aegypti)*. The postman butterfly *(Heliconius*

melpomene) was used as an outgroup. Transposable elements are known to be highly correlated

with genome size across the tree of life (Kidwell, 2002; Lynch, 2007), and our results across

various dipteran genomes are consistent with this pattern (Figure 2a).

We also compared different categories of transposable elements across these seven

genomes (Figure 2b). *P. downsi* had the highest proportion of DNA transposons (23.4%) among

all taxa analyzed. In comparison, only 6.36% of the *S. calcitrans* genome and 15.96% of the *M.*

domestica genome consisted of DNA transposons. Although *P. downsi* and *S. calcitrans* have

similar genome sizes, *S. calcitrans* had slightly higher amounts of repeat content in its genome

(58.3%), compared to *P. downsi* (51.7%) (Figure 2b). *S. calcitrans* had higher amounts of long

interspersed nuclear elements (LINEs) (23.8%), compared to *P. downsi* (7.7%). We observed

that most Diptera (except *Aedes*) have only a few SINE elements, one of the major classes of

transposable elements (Figure 2b). LTR elements were relatively common in *D. melanogaster*,

whereas other dipterans had a low amount of LTR elements (including *P. downsi*).

Orthologs to other dipteran genomes and outgroup *Heliconius*

Comparative genomics analysis of 113,047 protein sequences from *P. downsi* and seven

other species (listed above) identified 11,112 orthogroups. A total of 95,567 proteins (out of

113,047, 86.3%) were assigned to these orthogroups. The mean size of an orthogroup was 8.8

genes/species, and 3,069 orthogroups had singly copy genes in each species. A total of 5,754

orthogroups were shared among all eight species. Only a few orthogroups (minimum 496 in *A.*

aegypti and a maximum of 1,445 in *M. domestica*) were present in fewer than four species

(Figure 3a). The number of unique orthologs in each species was consistent with their

phylogenetic relationships (e.g., *A. aegypti* and the outgroup *H. melpomene* had the highest

number of unique orthologs). The number of shared orthologs among each pair of species is

presented in Supplementary Table 3. The number of orthologs identified in *P. downsi* is

consistent with other published dipteran genomes (Figure 3a, Supplementary Table 3), indicating

no major bias in the genome assembly and annotation pipeline used in this study.

Within *P. downsi*, 13,706 out of 15,774 (86.9%) annotated genes were assigned to

orthogroups. We expect the remaining missing genes to either be the most recently evolved

orphan genes in the branch leading to the *P. downsi* lineage or the consequence of a lack of

inclusion of enough closely related species of *P. downsi* in the analysis. The distribution of these

unique genes in the *P. downsi* genome is random and does not show specific clustering patterns

across various locations of the genome.

We also compared the number of pairwise orthogroups that are uniquely shared among

all eight species (Figure 3b). A total of 993 orthogroups were unique to Diptera (after excluding

the outlier *H. melpomene),* 79 orthogroups were shared only between *M. domestica* and *P.*

downsi and 12 putative gene families were unique to only *P. downsi.* These 12 gene families

consisted of 27 genes and the great majority had the best BLAST hits against "uncharacterized"

or "hypothetical" proteins in other related species (Supplementary Table 4). This result indicates

that these gene families that appeared "unique" in *P. downsi* are likely due to the lack of proper

gene annotation in other species.

We further examined the gene ontology terms of the 993 orthogroups unique to Diptera

using the PANTHER gene ontology database (Mi *et al.*, 2021). The common biological

processes of these genes included localization, locomotion, immune system processes, response

to stimulus, and reproduction (Supplementary Figure 3), many of which are likely key genes for

the overall development and function of dipterans.

Gene family evolution

We used a maximum likelihood approach to analyze changes in gene family size among the same eight species used for inferring orthologs (De Bie *et al.*, 2006). This method uses a statistical approach to model gene gain and loss, accounting for phylogenetic history and assess the significance of the observed gene family size differences among taxa. Across 11,112 orthogroups we identified before among eight taxa; 101 gene families were significantly expanding/contracting ($p < 0.01$) across the phylogenetic tree. 25 out of these 101 gene families were identified in the branch leading to *P. downsi*. The list of these gene families in *P. downsi* included those associated with insecticide resistance or detoxification and host defense or immunity proteins (Table 2).

Table 2: *List of* significantly expanding/contracting ($p < 0.01$) *gene families in P. downsi*

Functional Categories	Gene Family
Insecticide Resistance and Detoxification	Cytochrome P450
	Glutathione S Transferase
	Cuticular Protein
Defense against host Immunity	Fibrinogen C-terminal Domain-Containing Protein
	Scp Domain-Containing Protein

Gene families associated with insecticide resistance and detoxification

Cytochrome P450 gene family

We examined the number of genes in the cytochrome P450 (CYP450) family across the seven Diptera and the *H. melpomene* outgroup. *P. downsi and M. domestica* have an expanded CYP450 gene family in comparison to their most recent common ancestor with *G. morsitans* (family Glossinidae) (Figure 4a). For example, The *P. downsi* CYP450 family is composed of 102 genes in comparison to 66 in *G. morsitans.* An even greater level of expansion was observed in two other members of Muscidae *(M. domestica:* 143 genes and *S. calcitrans:* 193 genes).

Compared to *G. morsitans*, the expansion of CYP450 genes is mainly found in CYP4, CYP6,

and CYP28 genes (Figure 4b). The CYP6 subfamily in *P. downsi* is composed of 25 genes,

almost doubling the number from the 14 genes present in *G. morsitans.* We also found an

expansion in the CYP4 subfamily from 15 genes in *G. morsitans* to 31 genes in *P. downsi.* A

similar expansion was seen in the CYP28 subfamily (4 genes in *G. morsitans* to 8 genes in *P.*

downsi).

Glutathione S-transferase gene family

The *P. downsi* Glutathione S-transferase family consists of 25 genes and the pattern

across the dipterans is similar to observations of the Cytochrome P450 gene family expansion

(Figure 4c). For example, in comparison to *G. morsitans* (15 genes), *P. downsi and M. domestica*

have an expanded number of genes (25 genes in *P. downsi,* 31 genes in *M. domestica*). The

GSTs are grouped into six subclasses (Delta, Epsilon, Omega, Sigma, Theta, and Zeta). The

expansion of GSTs in *P. downsi* occurred mainly in the Delta, Epsilon, and Zeta subclasses

(Figure 4d).

Cuticular Protein

The *P. downsi* cuticular gene family consists of 214 genes and the pattern across the

dipterans is similar to observations of the Cytochrome P450 and GST gene family expansions

(Supplementary Figure 4). For example, the *Muscidae* family (represented here by *P. downsi, M.*

domestica, and S. calcitrans) has an expanded gene family in comparison to their most recent

common ancestor with *G. morsitans* (115 genes). An even greater level of expansion was

observed by two other members of *Muscidae (M. domestica* – 357 genes and *S. calcitrans* – 270

genes).

Gene families associated with immunity

Two gene families (Fibrinogen C-terminal Domain-Containing Protein and Scp Domain-

Containing Protein) associated with immunity were also identified as significantly changing in P.

downsi (Table 2). Opposite of genes associated with insecticide resistance and detoxification

(Cytochrome P450 and Glutathione S-transferase), these gene families showed a reduction in the

number of genes in *P. downsi* compared to the most recent common ancestor with *G. morsitans*

(Supplementary Table 5). They also show contraction in comparison with two included members

of the Family Muscidae *(M. domestica and S. calcitrans)*.

Discussion

In this study, we report the first genome sequence of the avian vampire fly, a highly

invasive parasitic nest fly that threatens endemic avifauna of the Galápagos Islands (Fessl *et al.*,

2018; Causton *et al.*, 2019). This genome is meant to serve as an important resource to research

efforts aimed at characterizing the molecular mechanisms of the fly's successful invasion in the

Galápagos. The genome size of *P. downsi* is 971.6 Mb and its high quality is reflected both in

terms of genome contiguity (scaffold N_{50} of 1.3 Mb) and completeness (98% BUSCO gene-space

score). Interestingly, the total number of annotated genes in *P. downsi* is slightly lower compared

to other published fly genomes (*M. domestica* and *D. melanogaster*) (Table 1). The low gene

count likely reflects the lack of *P. downsi* specific transcriptome data in this study, which

perhaps led to reduced gene predictions. We aim to improve the *P. downsi* genome annotations

in the future using additional transcriptomic resources using additional flies collected from the

Galápagos.

Transposable elements (TE) are typically non-coding sequences that can insert themselves in

various places of the genome, often with neutral or deleterious phenotypic consequences

(Bourque *et al.*, 2018). The role of TEs, as well as their evolution across insect genomes, is still

an area of major research, but they are thought to be important drivers of genomic architecture

depending on the location of the genome to which they insert themselves (i.e., coding versus

non-coding regions). Furthermore, TE may also be a critical mechanism of adaptive evolution, as

has been shown in an invasive ant species (Schrader *et al.*, 2014). Analysis of transposable

elements in *P. downsi* and across other dipteran genomes showed a strong positive correlation

between genome size and repeat content (Figure 2a), consistent with similar findings across other

taxa (Lynch, 2007; Lamichhaney *et al.*, 2021). Interestingly, *P. downsi* had a higher number of

DNA transposons (Class II TEs) than any other compared genome, including *M. domestica, S.*

calcitrans, and *G. morsitans*. While long terminal repeat (LTR) transposons, LINEs, and SINEs)

were present in the species studied, *P. downsi* had no SINEs, a finding consistent with a study by

Petersen and colleagues (Petersen *et al.*, 2019) showing that SINEs contribute less than 1% to the

TE content of dipterans. However, it is important to note that some SINEs may be present in *P.*

downsi but are currently masked as unclassified. Future research should explore the role of TEs,

especially DNA transposons, in aiding the invasion of *P. downsi* to the Galápagos.

Comparative genomics analysis of *P. downsi* and other additional dipteran genomes

allowed us to identify gene families that were significantly expanded/contracted (p < 0.01) in *P.*

downsi. The list of these gene families in *P. downsi* included those associated with insecticide

resistance or detoxification and host defense or immunity proteins, and we predict that these

gene families are associated with a successful invasion of *P. downsi* in the Galápagos (Table 1)*.*

E.g. CYP450 mono-oxygenases are a diverse superfamily of proteins, including enzymes,

associated with catabolism and anabolism of xenobiotics and endogenous compounds. These

monooxygenase-mediated metabolisms have allowed numerous insect species to develop

insecticide resistance and detoxification (Scott, 1999; Wen *et al.*, 2011). The expansion of

CYP450 genes in *P. downsi* is mainly found in CYP4 and CYP6 gene subfamilies (Figure 4b),

which has also been shown in other species from the family Muscidae (Scott *et al.*, 2014;

Olafson *et al.*, 2021). Interestingly, in higher Diptera, many of the genes within the CYP6

subfamily (e.g., CYP6A, CYP6G, CYP6D) are associated with insecticide resistance

(Feyereisen, 2012). For example, the Cyp6g1 gene is involved in resistance to the insecticide

Dichlorodiphenyltrichloroethane (DDT) in *D. melanogaster* (Festucci-Buselli *et al.*, 2005). The

CYP4 subfamily can also influence the breakdown of synthetic insecticides (Iga and Kataoka,

2012). Cyp4d4v2, Cyp4g2, and Cyp6a38 can be co-up-regulated in-house flies that are resistant

to the insecticide permethrin, which is used to control *P. downsi* in the Galapagos (Zhu *et al.*,

2008). Overall, the expansion of CYP4 and CYP6 subfamilies may indicate the evolution of

insecticide resistance in *P. downsi* over macroevolutionary time (i.e., before arriving in the

Galápagos), which could have facilitated its invasion to the Galápagos and might affect its

management on the islands.

Glutathione S-transferase (GSTs) was another major expanding gene family in *P. downsi.*

GSTs are a highly conserved, large family of dimeric enzymes associated with detoxification of

endogenous and/or xenobiotic compounds, such as insecticides (Ketterman *et al.*, 2011). The

GST family is further grouped into six subclasses (Delta, Epsilon, Omega, Sigma, Theta, and

Zeta), with Delta and Epsilon being specific subclasses found in the class Insecta. We observed

an expansion of GST genes in *P. downsi* relative to *G. morsitans*, but fewer than were found in

M. domestica and S. calcitrans. The major expansions of the GST family in *P. downsi* were

observed in Delta and Epsilon subclasses (Figure 4d). Insecticides, such as permethrin, are used

to experimentally manipulate *P. downsi* abundance in bird nests to study the effects of the

parasite on the health of the birds (Fessl *et al.*, 2010; Koop *et al.*, 2011; Koop, Le Bohec, *et al.*,

2013; Koop, Owen, *et al.*, 2013; Knutie *et al.*, 2014, 2016b; O'Connor *et al.*, 2014; McNew *et*

al., 2020; Addesso *et al.*, 2020). Increased expression of GSTs following permethrin exposure

has been documented in several insect species including oriental fruit flies (*Bactrocera dorsalis*)

(Hu *et al.*, 2008). Expansion of the GST gene family in *P. downsi* may be a result of such

permethrin exposure. However, gene expression studies that further explore the role of the GST

gene family in insecticide resistance and detoxification are needed across insect taxa including *P.*

downsi. It is important to note that the expansion of CYP450 and GST families observed in *P.*

downsi may be an artifact of phylogenetic relationships rather than ecological adaptations. Still,

given the observed and predicted impacts of *P. downsi* on native endemic host populations, it is

important to consider the implications of expanded gene families related to detoxification and

insecticide resistance.

Previous studies have shown that Darwin's finch species can increase specific antibody

responses to parasitism (Koop, Owen, *et al.*, 2013). This response is most prominent in adult

females that are likely parasitized while brooding nestlings or eggs on the nest. However, little is

known about the ability of host bird immunological responses to effectively reduce fly fitness, in

part, because so little is known about the fly itself. One of the most prominent questions is

whether the *P. downsi* possesses the ability to counter defend against host immune responses.

We identified a reduction in the size of two additional gene families (Fibrinogen C-terminal

Domain-Containing and SCP domain-containing gene family), both with immune function

properties in *P.downsi* (Table 1, Supplementary Table 5). Fibrinogen plays a key role in blood

clot formation through the conversion of fibrinogen to insoluble fibrin (Weisel and Litvinov,

2017) and the C-terminal domain of fibrinogen is the primary binding site of platelets

(Hanington and Zhang, 2011). The Sperm-coating glycoprotein (Scp) family contains, among

other proteins, antigen 5 (Ag5), which is associated with the venom secretory ducts of stinging

insects (Gibbs *et al.*, 2008). However, the interpretation of these results is difficult without

further investigation into whether these genes are associated with innate immune responses of *P.*

downsi toward their hosts or whether they might be important components of their feeding

ecology. The goal of our study was not to make such inferences, but rather to highlight

promising avenues of future research.

The invasion of *P. downsi* has had dramatic negative effects on the endemic avifauna of

the Galápagos, including Darwin's finches. As researchers work to better understand the

pathway of invasion and the ecological and evolutionary processes that may have facilitated its

invasion to the Galápagos, the need for a high-quality whole genome sequence has grown. The

addition of this resource is therefore meant to provide the foundation for further investigations

using genomics tools in this system. These genomic resources will further allow us to understand

the evolution of the *P. downsi* defense in response to host defenses or disease resistance itself.

Gene expression studies could shed light on the development of larvae and at what stage they are

most vulnerable to host defenses. Further population-scale resequencing of various populations

of *P. downsi* will also allow us to explore mechanisms of local adaptation of the parasite to the

environment across islands.

Data availability

All raw data generated in this study (raw short sequence reads and draft genome assembly) has been deposited to NCBI, under accession number PRJNAXXX. The final genome assembly and annotation can be found under the accession number GCA_XXX. Supplementary material is available on figshare.

Acknowledgment

We would like to thank Lauren Albert, Taylor Verrett, and Corrine Arthur for field assistance and the Galápagos Science Center and the Galápagos National Park for logistical support. We also thank Noah K. Whiteman for his helpful comments on the manuscript. The sampling was done under Galápagos National Parks permits PC 28-19 and Genetic Access permit MAE-DNB-CM-2016-0041. This project was supported by the Department of Biological Sciences, Kent State University to SL, and the University of Connecticut to SAK.

Competing interest statement

The authors declare no competing interests.

Author contributions

SL and JAHK conceived the idea. SAK, GJV, and JC collected the specimens. CMC and JAHK carried out the laboratory work. MR and SL designed and performed the bioinformatic analyses with the support of CMC and JAHK. SL and JAHK prepared the manuscript, and all authors edited and approved the final version.

List of Figures

Figure 1: The avian vampire fly, Philornis downsi **(a)**, parasitizes many endemic bird species of

the Galapagos Islands, including the medium ground finch, Geospiza fortis **(b)**. The fly is

parasitic in its larval forms **(c**, bottom three) when it feeds on the blood and other fluids of its

avian hosts. The larva then pupates **(c**, second from top) and eclose **(c**, top) as adult flies.

Figure 2: Landscape of Transposable elements in *P. downsi* **(a)** Comparison of repeat content

and genome size across Diptera and its outgroup **(b)** Repeat statistics on various classes of

transposable elements across dipteran genomes.

Figure 3: Orthogroups in *P. downsi* **(a)** Phylogenetic relationship between *P. downsi* and other

seven published Diptera genomes, estimated using alignments from 3,069 orthogroups had

singly copy orthogroups in each species. Horizontal bars for each species show number of

orthogroups that are single-copy orthologs in all species, present in all species, present in the

majority of species, present in few species, and unique to the species **(b)** Number of shared

orthologs among all species.

Figure 4: Gene family evolution **(a)** Number of genes in Cytochrome P450 gene family across

Diptera **(b)** Number of genes in various subfamilies of Cytochromes P450 gene family in *G.*

morsitans and *P. downsi, M. domestica* and *D. melanogaster* **(c)** Number of genes in Glutathione

S-transferase gene family across Diptera **(d)** Number of genes in various subfamilies of

Glutathione S-transferase gene family in *G. morsitans* and *P. downsi, M. domestica* and *D.*

melanogaster.

References

Adams MD, Celniker SE, Holt RA, Evans CA, Gocayne JD, Amanatides PG, *et al.* (2000). The
genome sequence of Drosophila melanogaster. *Science* **287**: 2185–95.

Addesso AM, Harvey JA, Vaziri GJ, Verrett TB, Albert L, Arthur C, *et al.* (2020). Effect of
introduced parasites on the survival and microbiota of nestling cactus finches (Geospiza
scandens) in the Galápagos Islands. *Journal of Ornithology* **161**: 1011–1019.

Alves F, Langmore N, Heinsohn R, Stojanovic D (2020). 'Self-fumigation' of nests by an
endangered avian host using insecticide-treated feathers increases reproductive success
more than tenfold. *Animal Conservation*.

Bao W, Kojima KK, Kohany O (2015). Repbase Update, a database of repetitive elements in
eukaryotic genomes. *Mobile DNA* **6**: 11.

Bourque G, Burns KH, Gehring M, Gorbunova V, Seluanov A, Hammell M, *et al.* (2018). Ten
things you should know about transposable elements. *Genome Biol* **19**: 199–199.

Bulgarella M, Quiroga MA, Boulton RA, Ramirez IE, Moon RD, Causton CE, *et al.* (2017). Life
Cycle and Host Specificity of the Parasitoid Conura annulifera (Hymenoptera:
Chalcididae), a Potential Biological Control Agent of Philornis downsi (Diptera:
Muscidae) in the Galapagos Islands. *Ann Entomol Soc Am* **110**: 317–328.

Bulgarella M, Quiroga MA, Brito vera GA, Dregni JS, Cunningham F, Mosquera Munoz DA, *et*
al. (2015). Philornis downsi (Diptera: Muscidae), an avian nest parasite invasive to the
Galapagos islands, in mainland Ecuador. *Annals of the Entomological Society of*
America.

Bulgarella M, Quiroga MA, Heimpel GE (2019). Additive negative effects of Philornis nest
parasitism on small and declining Neotropical bird populations. *Bird Conservation*
International **29**: 339–360.

Cantarel BL, Korf I, Robb SMC, Parra G, Ross E, Moore B, *et al.* (2008). MAKER: an easy-to-
use annotation pipeline designed for emerging model organism genomes. *Genome Res*
18: 188–196.

Causton CE, Cunninghame F, Tapia W (2013). *Management of the avian parasite Philornis*
downsi in the Galapagos Islands: a collaborative and strategic action plan. GNPS,
GCREG, CDF, and GC: Puerto Ayora, Galapagos, Ecuador.

Causton CE, Moon RD, Cimadom A, Boulton RA, Cedeño D, Lincango MP, *et al.* (2019).
Population dynamics of an invasive bird parasite, Philornis downsi (Diptera: Muscidae),
in the Galapagos Islands. *PLoS One* **14**: e0224125.

Causton CE, Peck SB, Sinclair BJ, Roque-Albelo L, Hodgson CJ, Landry B (2006). Alien
insects: Threats and implications for conservation of Galápagos Islands. *Annals of the*
Entomological Society of America **99**: 121–143.

Couri MS, de Carvalho CJB, Lowenberg-Neto P (2007). Phylogeny of Philornis Meinert species
(Diptera: Muscidae). *Zootaxa*: 19–26.

Cunningham F, Achuthan P, Akanni W, Allen J, Amode MR, Armean IM, *et al.* (2019).
Ensembl 2019. *Nucleic Acids Research* **47**: D745–D751.

Cunninghame F, Fessl B, Sevilla CR, Young GR, La Greco N (2017). *Manejo de la*
conservacion a largo plazo para salvar al pinzon de manglar (Camarhynchus heliobates)
en peligro critico de extincion. DPNG, CGREG, FCD and GC: Puerto Ayora, Galapagos,
Ecuador.

De Bie T, Cristianini N, Demuth JP, Hahn MW (2006). CAFE: a computational tool for the
study of gene family evolution. *Bioinformatics (Oxford, England)* **22**: 1269–1271.

Dodge HR (1955). New Muscid flies from Florida and the West Indies (Diptera: Muscidae). *The*
Florida Entomologist **38**: 147–151.

Dodge HR (1963). A new Philornis with coprophagous larva, and some related species (Diptera:
Muscidae). *Journal of the Kansas Entomological Society* **36**: 239–247.

Dudaniec RY, Fessl B, Kleindorfer S (2007). Interannual and interspecific variation in intensity
of the parasitic fly, Philornis downsi, in Darwin's finches. *Biological Conservation* **139**:
325–332.

Dudaniec RY, Gardner MG, Donnellan S, Kleindorfer S (2008). Genetic variation in the invasive
avian parasite, Philornis downsi (Diptera, Muscidae) on the Galapagos archipelago. *BMC*
Ecology **8**: 13.

Dudaniec RY, Kleindorfer S (2006). Effects of the parasitic flies of the genus Philornis (Diptera :
Muscidae) on birds. *Emu* **106**: 13–20.

Dvorak M, Vargas H, Fessl B, Tebbich S (2004). On the verge of extinction: a survey of the
mangrove finch Cactospiza heliobates and its habitat on the Galápagos Islands. *Oryx* **38**:
1–9.

Emms DM, Kelly S (2015). OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. *Genome Biology* **16**:
157.

Fessl B, Heimpel G, Causton C (2018). Invasion of an avian nest parasite, Philornis downsi, to
the Galapagos Islands: colonization history, adaptations to novel ecosystems, and
conservation challenges. In: Parker P (ed) *Disease Ecology, Social and Ecological*
Interactions in the Galapagos Islands, Springer International Publishing.

Fessl B, Sinclair BJ, Kleindorfer S (2006). The life-cycle of Philornis downsi (Diptera :
Muscidae) parasitizing Darwin's finches and its impacts on nestling survival.
Parasitology **133**: 739–747.

Fessl B, Tebbich S (2002). Philornis downsi - a recently discovered parasite on the Galápagos
archipelago - a threat for Darwin's finches? *Ibis* **144**: 445–451.

Fessl B, Young GH, Young RP, Rodríguez-Matamoros J, Dvorak M, Tebbich S, *et al.* (2010).
How to save the rarest Darwin's finch from extinction: the mangrove finch on Isabela
Island. *Philos Trans R Soc Lond B Biol Sci* **365**: 1019–1030.

Festucci-Buselli RA, Carvalho-Dias AS, de Oliveira-Andrade M, Caixeta-Nunes C, Li HM,
Stuart JJ, *et al.* (2005). Expression of Cyp6g1 and Cyp12d1 in DDT resistant and
susceptible strains of Drosophila melanogaster. *Insect Mol Biol* **14**: 69–77.

Feyereisen R (2012). Insect CYP genes and P450 enzymes. In: Gilbert LI, editor. *Insect*
Molecular Biology and Biochemistry: 236–316.

Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, *et al.* (2019).
RepeatModeler2: automated genomic discovery of transposable element families.
bioRxiv: 856591.

Gibbs GM, Roelants K, O'Bryan MK (2008). The CAP Superfamily: Cysteine-Rich Secretory
Proteins, Antigen 5, and Pathogenesis-Related 1 Proteins—Roles in Reproduction,
Cancer, and Immune Defense. *Endocrine Reviews* **29**: 865–897.

Grant PR, Grant BR, Petren K, Keller LF (2005). Extinction behind our backs: the possible fate
of one of the Darwin's finch species on Isla Floreana, Galápagos. *Biological*
Conservation **122**: 499–503.

Hanington PC, Zhang S-M (2011). The primary role of fibrinogen-related proteins in
invertebrates is defense, not coagulation. *J Innate Immun* **3**: 17–27.

Heimpel GE, Hillstrom A, Freund D, Knutie SA, Clayton DH (2017). Invasive Parasites and the
Fate of Darwin's Finches in the Galapagos Islands: The Case of the Vegetarian Finch
(Platyspiza crassirostris). *Wilson J Ornithol* **129**: 345–349.

Hu J, Zhang JL, Nardi F, Zhang RJ (2008). Population genetic structure of the melon fly,
Bactrocera cucurbitae (Diptera: Tephritidae), from China and Southeast Asia. *Genetica*
134: 319–24.

Iga M, Kataoka H (2012). Recent Studies on Insect Hormone Metabolic Pathways Mediated by
Cytochrome P450 Enzymes. *Biological and Pharmaceutical Bulletin* **35**: 838–843.

Kapusta A, Suh A, Feschotte C (2017). Dynamics of genome size evolution in birds and
mammals. *Proceedings of the National Academy of Sciences* **114**: E1460 LP-E1469.

Ketterman AJ, Saisawang C, Wongsantichon J (2011). Insect glutathione transferases. *Drug*
Metabolism Reviews **43**: 253–265.

Kidwell MG (2002). Transposable elements and the evolution of genome size in eukaryotes.
Genetica **115**: 49–63.

Knutie SA, McNew SM, Bartlow AW, Vargas DA, Clayton DH (2014). Darwin's finches
combat introduced nest parasites with fumigated cotton. *Current Biology* **24**: R355–
R356.

Knutie SA, Owen JP, McNew SM, Bartlow AW, Arriero E, Herman JM, *et al.* (2016a).
Galapagos mockingbirds tolerate introduced parasites that affect Darwin's finches.
Ecology **97**: 940–950.

Koop JAH, Causton CE, Bulgarella M, Cooper E, Heimpel GE (2021). Population structure of a
nest parasite of Darwin's finches within its native and invasive ranges. *Conservation*
Genetics **22**: 11–22.

Koop JAH, Huber SK, Laverty SM, Clayton DH (2011). Experimental demonstration of the
fitness consequences of an introduced parasite of Darwin's finches. *PLoS ONE* **6**: e19706
doi:10.1371/journal.pone.0019706.

Koop JAH, Kim PS, Knutie SA, Adler F, Clayton DH (2015). Introduced parasitic fly may lead
to local extinction of Darwin's finch populations. *Journal of Applied Ecology*.

Koop JAH, Le Bohec C, Clayton DH (2013). Dry year does not reduce invasive parasitic fly
prevalence or abundance in Darwin's finch nests. *Reports in Parasitology* **3**: 11–17.

Koop JAH, Owen JP, Knutie SA, Aguilar MA, Clayton DH (2013). Experimental demonstration
of a parasite-induced response in wild birds: Darwin's finches and introduced nest flies.
Ecology and Evolution **3**: 2514–2523.

Lamichhaney S, Catullo R, Keogh JS, Clulow S, Edwards SV, Ezaz T (2021). A bird-like
genome from a frog: Mechanisms of genome size reduction in the ornate burrowing frog,
Platyplectrum ornatum. *Proceedings of the National Academy of Sciences* **118**:
e2011649118.

Li H, Durbin R (2009). Fast and accurate short read alignment with Burrows–Wheeler transform.
Bioinformatics **25**: 1754–1760.

Lynch M (2007). *The Origins of Genome Architecture*, 1 edition. Sinauer Associates Inc:
Sunderland.

McNew SM, Knutie SA, Clayton DH (2020). No evidence of sex ratio manipulation by
Galápagos mockingbirds in response to environment. *Journal of Avian Biology* **51**.

Mi H, Ebert D, Muruganujan A, Mills C, Albou L-P, Mushayamaha T, *et al.* (2021). PANTHER
version 16: a revised family classification, tree-based classification tool, enhancer regions
and extensive API. *Nucleic Acids Research* **49**: 394–403.

O'Connor JA, Robertson J, Kleindorfer S (2014). Darwin's finch begging intensity does not
honestly signal need in parasitised nests. *Ethology* **120**: 228–237.

Olafson PU, Aksoy S, Attardo GM, Buckmeier G, Chen X, Coates CJ, *et al.* (2021). The genome
of the stable fly, Stomoxys calcitrans, reveals potential mechanisms underlying
reproduction, host interactions, and novel targets for pest control. *BMC Biology* **19**: 41.

Peters KJ, Kleindorfer S (2018). Avian population trends in Scalesia forest on Floreana Island
(2004-2013): Acoustical surveys cannot detect hybrids of Darwin's tree finches
Camarhynchus spp. *Bird Conservation International* **28**: 319–335.

Petersen M, Armisén D, Gibbs RA, Hering L, Khila A, Mayer G, *et al.* (2019). Diversity and
evolution of the transposable element repertoire in arthropods with particular reference to
insects. *BMC Evol Biol* **19**: 11–11.

Pourrajab F, Hekmatimoghaddam S (2021). Transposable elements, contributors in the evolution
of organisms (from an arms race to a source of raw materials). *Heliyon* **7**: e06029–
e06029.

Ranallo-Benavidez TR, Jaron KS, Schatz MC (2020). GenomeScope 2.0 and Smudgeplot for
reference-free profiling of polyploid genomes. *Nature Communications* **11**: 1432.

Schrader L, Kim JW, Ence D, Zimin A, Klein A, Wyschetzki K, *et al.* (2014). Transposable
element islands facilitate adaptation to novel environments in an invasive species. *Nature
Communications* **5**: 5495.

Scott JG (1999). Cytochromes P450 and insecticide resistance. *Insect Biochem Mol Biol* **29**:
757–77.

Scott MJ, Benoit JB, Davis RJ, Bailey ST, Varga V, Martinson EO, *et al.* (2020). Genomic
analyses of a livestock pest, the New World screwworm, find potential targets for genetic
control programs. *Communications Biology* **3**: 424–424.

Scott JG, Warren Wc Fau - Beukeboom LW, Beukeboom Lw Fau - Bopp D, Bopp D Fau - Clark
AG, Clark Ag Fau - Giers SD, Giers Sd Fau - Hediger M, *et al.* (2014). Genome of the
house fly, Musca domestica L., a global vector of diseases with adaptations to a septic
environment.

Smit Hubley P. AFA R& Green (2013). RepeatMasker Open-4.0,.

Spalding MG, Mertins JW, Walsh PB, Morin KC, Dunmore DE, Forrester DJ (2002). Burrowing
fly larvae (Philornis porteri) associated with mortality of eastern bluebirds in Florida.
Journal of Wildlife Diseases **38**: 776–783.

Vurture GW, Sedlazeck FJ, Nattestad M, Underwood CJ, Fang H, Gurtowski J, *et al.* (2017).
651   GenomeScope: fast reference-free genome profiling from short reads. *Bioinformatics* **33**:
652   2202–2204.

653   Waterhouse RM, Seppey M, Simao FA, Manni M, Ioannidis P, Klioutchnikov G, *et al.* (2017).
654   BUSCO applications from quality assessments to gene prediction and phylogenomics.
655   *Molecular biology and evolution.*

656   Weisel JW, Litvinov RI (2017). Fibrin Formation, Structure and Properties. *Subcell Biochem* **82**:
657   405–456.

658   Weisenfeld NI, Kumar V, Shah P, Church DM, Jaffe DB (2017). Direct determination of diploid
659   genome sequences. *Genome Research* **27**: 757–767.

660   Wen Z, Zhang X, Zhang Y (2011). P450—mediated Insecticide Detoxification and Its
661   Implication in Insecticide Efficacy. In: Liu T, Kang L (eds) *Recent Advances in*
662   *Entomological Research: From Molecular Biology to Pest Management*, Springer Berlin
663   Heidelberg: Berlin, Heidelberg, pp 229–245.

664   Zheng GXY, Lau BT, Schnall-Levin M, Jarosz M, Bell JM, Hindson CM, *et al.* (2016).
665   Haplotyping germline and cancer genomes with high-throughput linked-read sequencing.
666   *Nature Biotechnology* **34**: 303–311.

667   Zhu F, Li T, Zhang L, Liu N (2008). Co-up-regulation of three P450 genes in response to
668   permethrin exposure in permethrin resistant house flies, Musca domestica. *BMC Physiol*
669   **8**: 18.

Figure 1

a

b

	Heliconius	*Aedes*	*Drosophila*	*Ceratitis*	*Glossina*	*Philornis*	*Stomoxyx*	*Musca*
■ *SINE*	13.67	2.36	0	0.41	0.01	0	0.02	0.02
■ *LINE*	4.22	16.92	4.95	11.91	3.22	7.67	23.81	3.23
■ *LTR elements*	0.81	9.97	12.05	2.89	0.43	1.62	0.82	2.82
■ *DNA transposons*	3.59	23.72	2.08	13.47	8.15	23.39	6.36	15.96
■ *Unclassified*	9	12.18	1.2	6.34	9.53	17.15	27.6	32.15

Figure 2

Figure 3

Figure 4

Supplementary Materials

The genome sequence of the avian vampire fly (Philornis downsi), an invasive nest parasite of Darwin's finches

5
6
7 **Supplementary Table 1:** *Genome Statistics of P. downsi estimated from short-sequencing reads*

Property	min	max
Heterozygosity	1.40879%	1.42194%
Genome Haploid Length	796,797,589 bp	799,050,413 bp
Genome Repeat Length	232,405,721 bp	233,062,812 bp
Genome Unique Length	564,391,869 bp	565,987,600 bp
Model Fit	95.7713%	98.7012%
Read Error Rate	1.228%	1.228%

16
17

Supplementary Table 2: *Repeat statistics in P. downsi genome*

sequences: 41,176

total length: 971,6346,46 bp

GC level: 35.10 %

bases masked: 502,200,627 bp (51.69 %)

==

	number of elements*	length occupied	percentage of sequence
Retroelements	224657	90253146 bp	9.29 %
SINEs:	**58**	**6761 bp**	**0.00 %**
Penelope	3187	661022 bp	0.07 %
LINEs:	**196268**	**74550945 bp**	**7.67 %**
CRE/SLACS	0	0 bp	0.00 %
L2/CR1/Rex	43733	18550153 bp	1.91 %
R1/LOA/Jockey	15852	7502835 bp	0.77 %
R2/R4/NeSL	834	673762 bp	0.07 %
RTE/Bov-B	102395	29493616 bp	3.04 %

36	L1/CIN4	1415	94740 bp	0.01 %
37	**LTR elements:**	**28331**	**15695440 bp**	**1.62 %**
38	BEL/Pao	12565	7084057 bp	0.73 %
39	Ty1/Copia	1561	775084 bp	0.08 %
40	Gypsy/DIRS1	9951	7481940 bp	0.77 %
41	Retroviral	2481	143387 bp	0.01 %
42				
43	**DNA transposons**	621139	227262774 bp	23.39 %
44	hobo-Activator	8749	1542960 bp	0.16 %
45	Tc1-IS630-Pogo	576750	212960682 bp	21.92 %
46	En-Spm	0	0 bp	0.00 %
47	MuDR-IS905	0	0 bp	0.00 %
48	PiggyBac	1280	395680 bp	0.04 %
49	Tourist/Harbinger	4060	1924044 bp	0.20 %
50	Other	275	11451 bp	0.00 %
51				
52	Rolling-circles	19653	4597029 bp	0.47 %
53				
54	**Unclassified:**	**828801**	**166508339 bp**	**17.14 %**
55				
56	Total interspersed repeats:		484024259 bp	49.82 %
57				
58	Small RNA:	7148	2907047 bp	0.30 %
59				
60	Satellites:	796	124775 bp	0.01 %
61	Simple repeats:	213281	8909635 bp	0.92 %
62	Low complexity:	34278	1638267 bp	0.17 %

65 **Supplementary Table 3:** *Number of shared orthogroups among various dipteran species and their*

66 *outgroup*

	H.melpomene	A.aegypti	C.capitata	S.calcitrans	M.domestica	P.downsi	G.morsitans	D.melanogaster
H.melpomene		6,978	6,874	6,883	6,910	6,800	6,337	6,859
A.aegypti	6,978		8,015	8,008	8,018	7,796	7,228	7,995
C.capitata	6,874	8,015		9,123	9,186	8,855	8,014	9,096
S.calcitrans	6,883	8,008	9,123		9,650	9,197	8,087	9,041
M.domestica	6,910	8,018	9,186	9,650		9,295	8,144	9,117
P.downsi	6,800	7,796	8,855	9,197	9,295		8,103	8,770
G.morsitans	6,337	7,228	8,014	8,087	8,144	8,103		7,940
D.melanogaster	6,859	7,995	9,096	9,041	9,117	8,770	7,940	

71 **Supplementary Table 4:** *Gene families unique only to P. downsi*

EGT47331	hypothetical protein CAEBREN_08836 [Caenorhabditis brenneri]
GBO13109	hypothetical protein AVEN_233885-1 [Araneus ventricosus]
GBP44057	PiggyBac transposable element-derived protein 4 [Eumeta japonica]
KMQ83311	transposable element tc3 transposase [Lasius niger]
KNC20639	hypothetical protein FF38_06891, partial [Lucilia cuprina]
KNC20966	hypothetical protein FF38_09018 [Lucilia cuprina]
KNC26906	hypothetical protein FF38_10930 [Lucilia cuprina]
KXJ68891	hypothetical protein RP20_CCG001210 [Aedes albopictus]
OAF68034	hypothetical protein A3Q56_04227 [Intoshia linei]
PCG68510	hypothetical protein B5V51_5157, partial [Heliothis virescens]
XP_011295145	PREDICTED: tigger transposable element-derived protein 6-like isoform X2 [Musca domestica]
XP_011295374	PREDICTED: tigger transposable element-derived protein 6 [Musca domestica]
XP_013109704	PREDICTED: uncharacterized protein LOC106088638 [Stomoxys calcitrans]
XP_017475229	PREDICTED: uncharacterized protein LOC108365650 [Rhagoletis zephyria]
XP_017478109	PREDICTED: uncharacterized protein LOC108367917 [Rhagoletis zephyria]
XP_017478991	PREDICTED: uncharacterized protein LOC108368617 [Rhagoletis zephyria]
XP_017479715	PREDICTED: uncharacterized protein LOC108369194 [Rhagoletis zephyria]
XP_017481195	PREDICTED: RNA-directed DNA polymerase from mobile element jockey-like [Rhagoletis zephyria]
XP_019891578	PREDICTED: ATP-binding cassette sub-family A member 3-like [Musca domestica]
XP_019894716	PREDICTED: uncharacterized protein LOC105262305 isoform X1 [Musca domestica]
XP_021704105	protein ALP1-like [Aedes aegypti]
XP_022823959	piggyBac transposable element-derived protein 4-like [Spodoptera litura]
XP_022834134	uncharacterized protein LOC111361914 [Spodoptera litura]
XP_033325321	uncharacterized protein LOC117219890 [Megalopta genalis]
XP_036214104	trypsin zeta-like [Bactrocera oleae]
XP_036337749	uncharacterized protein LOC118747737 isoform X3 [Rhagoletis pomonella]
XP_036342696	uncharacterized protein LOC118751975 [Rhagoletis pomonella]

73
74

75 **Supplementary Table 5:** Number of genes in *Fibrinogen C-Terminal Domain-Containing* gene family
76 and *SCP domain-containing* gene family

Species	Fibrinogen C-Terminal Domain-Containing gene family	SCP domain-containing gene family
H.melpomene	1	2
A.aegypti	33	20
C.capitata	10	6
S.calcitrans	45	17
M.domestica	32	25
P.downsi	5	7
G.morsitans	7	7
D.melanogaster	11	15

80 **Supplementary Figure 1**: *k-mer spectrum and fitted modelling used for estimating genome parameters*

81 *of P. downsi from short sequencing reads*

84 **Supplementary Figure 2:** *Distribution of the total coverage of the k-mer pair (y-axis) against relative*

85 *minor k-mer coverage (x-axis) providing evidence of diploidy in P. downsi.*

89   **Supplementary Figure 3:** *Gene ontology terms associated with 993 gene families unique to diptera*

90
91

94   **Supplementary Figure 4:** *Number of genes in Cuticular gene family across dipterans*