A robust field-based method to screen heat tolerance in wheat

Najeeb Ullah[1], Jack Christopher[1], Troy Frederiks[2], Shangyu Ma[1,3], Daniel KY Tan[4], Karine

Chenu[1,*]

[1] The University of Queensland, Queensland Alliance for Agriculture and Food Innovation (QAAFI), Leslie

Research Facility, 13 Holberton street, Toowoomba, QLD 4350, Australia

[2] Department of Agriculture and Fisheries, 203 Tor street, Toowoomba, QLD 4350, Australia

[3] School of Agronomy, Anhui Agricultural University, Hefei, Anhui, China

[4] The University of Sydney, Plant Breeding Institute, Sydney Institute of Agriculture, School of Life and

Environmental Sciences, Faculty of Science, Sydney, NSW 2006, Australia

[*] Corresponding author: karine.chenu@uq.edu.au

Abstract

Wheat is highly sensitive to elevated temperatures, particularly during pollen meiosis and

early-to-mid grain filling. The impact of heat stress greatly depends on the plant developmental

stage. Thus, germplasm ranking for heat tolerance in field trials may be confounded by

variations in developmental phase between genotypes at the time of heat events. A

photoperiod-extension method (PEM) was developed allowing screening of 35 diverse

genotypes at matched developmental phase despite phenological variations. Paired trials were

conducted to compare the new PEM against conventional field screening in plots. In the PEM,

plants were sown in single rows or small plots. Artificial lighting was installed at one end of

each row or plot to extend day length, inducing a gradient of flowering times with distance

from the lights. Individual stems or plot quadrats of each genotype were tagged at flowering.

Late-sown plants received more heat shocks during early to mid grain filling than earlier

sowings, suffering reductions in both individual grain weight (IGW) and yield. IGW was

reduced by 1.5 mg for each additional post-flowering day with temperature $> 30^{\circ}$C. Significant

genotypic differences in heat tolerance ranking were observed between PEM versus

conventional plot screening. Strong correlations between trials experiencing similar degree of

heat were found both for IGW and for total grain weight with the PEM either with individual-

stem tagging (e.g. average r of 0.59 and 0.54, respectively for environments with moderate

postflowering heat) or quadrat tagging (r of 0.53 and 0.47). However, correlations for IGW

and yield in these environments were either poor or negative for conventional trials (e.g.

average r of 0.11 and 0.12, respectively for environments with moderate postflowering heat).

Accordingly, a PCA grouped genotypes consistently for heir performance across environments
with similar heat stress in PEM trials but not in conventional trials. In this study, most
consistent genotype ranking for heat tolerance was achieved with the PEM with tagging and
harvesting individual spikes at matched developmental phase. The PEM with quadrat sampling
provided slightly less consistent rankings but appears overall more suitable for high-throughput
phenotyping. The method promises to improve the efficiency of heat tolerance field screening,
particularly when comparing genotypes of different maturity types.

Keyword: Phenotyping, heat stress, genotype x environment interaction, crop improvement,
breeding, photoperiod extension.

Introduction

Climate variability is among the major determinants of global crop yields, including wheat
(Najeeb et al., 2019; Ray et al., 2015) and strongly impedes plant breeding (Chapman et al.,
2012). A significant increase in the frequency of heat stress events, particularly during grain
filling has been recorded in the major wheat production regions such as Australia during the
past 30 years (Ababaei and Chenu, 2020). These heat events have substantially affected the
growth, development and ultimately yield of wheat crops (e.g. Ababaei and Chenu, 2020;
Zheng et al., 2016), and are impacting optimal management practices (e.g. Collins and Chenu,
2021; Lobell et al., 2015; Zheng et al., 2012). With the recent rate of climate change, a further
increase in the frequency of these heat events is projected in the near future both in Australia
(Collins and Chenu, 2021) and globally (Field et al., 2012). Thus, developing wheat genotypes
with superior heat tolerance during grain filling is critical for sustaining wheat grain yields and
maintaining food security in future hot climates (Ullah et al., 2020).

Wheat is highly sensitive to elevated temperatures and the impact of heat stress is strongly
dependent on the crop developmental stage (e.g. Chenu and Oudin, 2019; Farooq et al., 2011;
Prasad and Djanaguiraman, 2014). Reproductive and grain filling phases of wheat are
extremely sensitive to heat and even a mild increase in the atmospheric temperature during
these stages can significantly reduce grain yield. For example, a single hot day (>30°C) during
early reproductive stage or onset of meiosis in pollen or micro / megaspore development, can
completely sterilise the developing wheat pollen (Saini and Aspinall, 1982). The effect of high
temperature on pollen typically translates into a poor grain set and grain yield loss (Guo et al.,
2016). Each degree increase (from 15–22°C) in mean temperature during pollen developmental
can reduce grain number per unit area by 4%, while a degree increase in maximum temperature

at mid anthesis can also result in a 4% reduction in grain number in wheat (Wheeler et al.,
1996).

Post-anthesis heat reduces grain yield primarily by limiting assimilate synthesis, translocation,
and starch deposition to developing grains (Sofield et al., 1977). Grain weight is most sensitive
to heat during early grain filling and becomes progressively less sensitive as grain filling
proceeds (Stone and Nicolas, 1998). A single hot day (maximum temperature of 40°C)
occurring 10–13 days after anthesis can reduce individual grain weight (IGW) by 14% (Stone
and Nicolas, 1998). In addition, IGW may be reduced by 0.5% for each day delay in exposure
to heat stress between 15 and 35 days after anthesis (Stone and Nicolas, 1998). Reduction in
IGW of heat stressed plants is strongly linked with a shortened grain filling duration (Girousse
et al., 2021; Stone and Nicolas, 1998). For each °C rise in mean daily temperature above
optimum (15–20°C), a two to eight day reduction in grain filling duration has been reported in
wheat crop (reviewed by Streck, 2005). In the Australian wheatbelt, a steady increase in the
frequency of hot days ($T_{max} > 26$°C) during the grain filling period of wheat crops has been
recorded over the past 30 years (Ababaei and Chenu, 2020). These authors also pointed out
that heat-induced yield losses due to reduced grain weight (18.1%) were greater than those due
to grain number (3.6%). This highlights the importance of developing wheat germplasm more
tolerant to heat post-flowering.

Conventionally, wheat genotypes are screened for heat tolerance by serial sowings, using heat
chambers in the field, or in controlled environments (e.g. Telfer et al., 2021, 2018;
Thistlethwaite et al., 2020). Ranking for heat tolerance is typically based on physiological or
morphological traits associated with plant function and performance (Bennett et al., 2012).
However, changes in these traits are strongly influenced by the environment and the
methodology used (Limpens et al., 2012; Poorter et al., 2016). Field-based screening methods
are generally considered more representative of the plant response to natural environments
(Passioura, 2006). However, screening of wheat genotypes with varying maturity types may be
complicated by the unpredictability of heat events under field conditions. The impact of heat
events is highly dependent on the developmental phase (e.g. Chenu and Oudin, 2019;
Djanaguiraman et al., 2014; Tashiro and Wardlaw, 1990), which is specific to each particular
wheat line being tested. Thus, ranking of wheat genotypes for heat tolerance may be
confounded by variation in developmental stage during a natural heat event. An improved

technique to screen for high temperature stress at matched developmental stages of wheat
genotypes in the field could accelerate the selection of heat tolerant genotypes.

Here, we developed and tested a photoperiod-extension method (PEM) that allows comparison
of the performance of wheat genotypes with varying maturity types at common developmental
stages during natural heat events. The method was tested across three different locations over
three consecutive years, harvesting either specific spikes that flowered at the same time or
small areas of plants (i.e. quadrats) flowering the same day (i.e. 50% of the spikes at anthesis).
Rankings of wheat genotypes with varying maturity type and heat tolerance were compared
using PEM and conventional plots.

Materials and Methods

Growth conditions and experimental design

Field trials were conducted over three consecutive years (2018 to 2020) at three locations
across southern Queensland, Australia. Conventional plots were estabilished adjacent to trials
using the newly developed photoperiod-extension method (PEM). Randomised block design
trials with two times of sowing blocks and four replicates were established each year. To
maximise the likelihood of heat during grain filling, trials were planted typically later in the
cropping season than industry practice (Table 1). At The University of Queensland Research
Farm, Gatton (27°34′50″S, 152°19′28″E), the first sowings (s1) were established between late
May or early July and the second sowings (s2) between late August or early September. At the
Hermitage Research Station, Warwick (28°12'40"S, 152°06'06"E), trials were sown early June
or mid-July (s1) or mid-August and mid-September (s2). At the Tosari Crop Research Centre,
Tummaville (27°49'09.1"S 151°26'14.9"E), plants were sown mid-July (s1) and early
September (s2). All trials were fully irrigated (except at Tosari) and cultivated under non-
limiting fertiliser conditions (Table 1). A boom irrigator (centre pivot sprinkler) was used for
irrigating plots at Gatton (2018-2020) as well as both plot and PEM trials at Tosari. PEM trials
at Gatton (2018-2020) and Warwick (2018) were irrigated with wobbler sprinklers. In 2020,
both PEM and conventional plots trials at Warwick were irrigated using a drip irrigation
system. At Tosari, trials were irrigated at sowing and pre-flowering, but no post-flowering
irrigation was applied, as water was then unavailable. For all trials, standard crop management
practices including weed, disease and pest control were adopted during the season.

With the PEM, each wheat genotypes was either hand sown in a 5 m single row (33 cm row

spacing) in 2018 and 2019, or machine planted in a four row plot (1×5 m, 2020 with 25 cm

row spacing).

Conventional field plots were planted at the same time and with similar management to the

PEM trials. In 2018 and 2020, the conventional field plots (2×6 m) were set up in Gatton and

Warwick, while in 2019, genotypes were tested in smaller plots (1×6 m) at Tosari and Gatton.

All conventional plots were planted at a 25 cm row spacing with a population density of 130

plants m^{-2} and in four independent replications. Due to unavailability of irrigation water, only

sowing 2 (s2) plots were established at Gatton during 2018.

Trials are given identifiers denoting the site (Gatton (GAT), Tosari (TOS) or Warwick (WAR),

and the year of the trial. Data may also be identified by time of sowing (s1 or s2) and the

tagging event (T1, T2, T3) when applicable.

Genotypes

Thirty-five wheat (*Triticum aestivum* L.) genotypes with contrasting phenology and adaptation

were used in the study (Table S1, Supplementary). These included high-performing spring

cultivars Suntop, Spitfire, Gregory, Janz, Hartog, EGA Wylie, Corack, Yitpi, Mace and Scout

widely cultivated in major cropping regions of Australia. A set of eight CIMMYT genotypes

described as heat tolerant under Australian environments was obtained from the University of

Sydney (Thistlethwaite et al., 2020). Other genotypes used in these trials included donors of a

multi-reference parent nested association mapping (MR-NAM) population developed for

screening for heat and drought tolerance in wheat (Christopher et al., 2015, 2021; Richard,

2017). In total, 35 wheat genotypes were tested in PEM and plot trials, with 32 genotypes at

each trial, except for the PEM trial with quadrat harvest in 2020 when only 20 selected

genotypes were used (Table S1, Supplementary). In conventional plot trials, 32 genotypes were

used in each trial.

The photoperiod-extension method

The novel photoperiod-extension method (PEM) was based on a method described by

Frederiks et al. (2012) originally developed to test for frost damage. At one end of each row or

plot, LED lamps (CLA LT401, 9W T40 LED LAMP, 3000K 760LM) with a lumen efficiency

≥ 80, were set up approximately 1 m above the ground level and at spacing of 0.8 m. These

lamps supplement light by extending the day length to 20 h (Fig. 1). The intensity of light

diminishes with the square of the distance from the lights along the test row, with maximum

effect closest to the lights and minimum or no impact at the other end of the row. This variation

in light intensity across the rows induced a gradient of flowering times within each row with

the plants closest to the light developing more rapidly.

Plant measurements

For each PEM trial and sowing date in 2018 and 2019, approximately 20 stems of each

genotypes were tagged at flowering (Zadoks decimal growth stage 65; Zadoks et al., 1974).

The induced gradient in phenology along the rows allowed tagging of genotypes multiple times

for plants in rows or plots from each sowing time. One-to-three cohorts of stems were tagged

at precisely matched flowering in rows or plots from each sowing time and trial. These

sequentially tagged coherts were termed as 'tagging 1' (T1), 'tagging 2' (T2) and 'tagging 3'

(T3). The spikes from tagged stems were manually harvested at maturity and processed for

grain yield components.

In addition, in each test row of the all PEM trials in 2018 and s1 trials at Tosari in 2019, a

quadrat consisting of a 0.5 m section of row that originally contained heads from tagging 1 was

manually harvested to estimate yield and its components. In 2020, ~ 0.5 m section of each plot

(four rows) was tagged at flowering (Zadoks 65), and a quadrat of two central rows (0.5 m

each) within the tagged region was manually harvested at crop maturity.

All plots from the conventional method were harvested using a small plot machine harvester at

maturity when grain moisture was approximately 11%. Grain samples were maunally counted

to calculate individual grain weight (IGW).

Environment variables

Local weather stations (Campbell Scientific) were set up at each site to record weather data for

each 10 min period. Light interception was measured with light sensors (Apogee SP-110

pyranometers, and Apogee SQ-110 for radiation and PAR measurements, respectively)

installed at 1.5 m height. HMP60 (Vaisala INTERCAP®) probes were used to measure air

temperature (Tair) and relative humidty (RH) at 1.5 m above the ground. Environmental

characteristics of each trial are presented in Table 1.

Thermal time was calculated in degree days (°Cd) using the following equation: (Jamieson et

al., 1995)

$Thermal\ time\ = \sum_{h=1}^{24} -0.0032 * Tair_h^3 + 0.1369 * Tair_h^2 + 0.3968 * Tair + 0.993$

190   (Eq 1)

Where $Tair_h$ is the hourly air temperature data.

Vapour pressure deficit (VPD) was calculated hourly during the day time as in Alduchov and

Eskridge (1996) by the following equation:

$VPD = 0.61094 \left(\frac{1-RH}{100} \right)^{17.625*Tair/(Tair+243.04)}$ (Eq 2)

Where Tair and RH were the hourly air temperature and hourly air relative humidity,

respectively, during the daytime.

 Statistical analysis

Data were analysed using R (R Core Team, 2018). Individual and interaction effects of

genotype and environments (sowing, location and tagging) were determined by analysis of

variance (ANOVA). Statistical differences were tested with student's t-tests at a 5% level.

Principle component analysis was computed to rank genotypes in different environments.

Results

Wheat crops experienced a wide range of heat events across locations and sowing times

Wheat genotypes at each location, season, sowing and tagging experienced varying air

temperatures and vapour pressure deficit (VPD) in the pre- and post-flowering periods (Table

1, Fig. 2 & Table S2). In all trials, plants from sowing 2 ('s2') experienced significantly higher

temperatures and VPD than those from the first sowing ('s1'). Post-flowering mean air

temperature was ~20% higher in s2 than s1 across all trials (Table 1). Similar differences in

post-flowering daily-maximum air temperature were also observed between s1 and s2, with the

exception of in the 2020 Warwick trial (WAR20), where temperatures were only 10% higher.

The number of hot days (with maximum temperature $>30°C$) across these trials also varied

greatly, with TOS19s2 in 2019 receiving the maximum number (22) of post-flowering hot days.

Higher temperature during s2 shortened the duration of both pre- and post-flowering periods,

although the reduction in time to flowering was relatively greater than the reduction in post-

flowering duration across all the tested locations (Table 1).

Environments with similar post-flowering stress were defined as (i) heat environment type 1

(HET1) that corresponded to environments with no or only late grain-filling heat stress (i.e.

less of 4 cumulated hours of temperature $>30°C$ between 0 and $450°Cd$ after flowering), (ii)

HET2 that included all environments with moderate heat stress during grain fill (5 to 15 days

with a maximum temperature $>30°C$ in our set of trials), and (iii) HET3 that corresponded to

environments severely stressed during grain fill (20 and 22 days with a maximum temperature

$>30°C$ in our set of trials). In addition to post-flowering heat, HET3 trials experienced high

pre-flowering temperatures (Table S2, Supplementary) including with hot days around stem

elongation and meiosis (Fig. 2) that significantly reduced grain set (Table S3 Supplementary).

Late-sown crops had lower individual grain weight and grain yield

Individual grain weight and grain yield varied significantly across trials, but plants typically

produced significantly smaller grains and lower grain yield in s2 compared with s1 (Fig. 3).

Trial-mean grain number in s2 was also significantly lower or not significantly different than
s1 in almost all trials (Table S3 Supplementary), probably due to shorter pre-flowering periods
associated to warmer temperatures (Table S2 Supplementary).

The maximum trial-means for IGW and grain yield were measured in HET1 environments. In
the PEM with quadrat tagging and harvests, a maximum of 45.5 mg for IGW and 571 g m^{-2} for
grain yield was recorded in GAT18s1 (Fig. 3). With the conventional plot method, genotypes
produced a maximum trial-mean IGW of 37.7 mg and a maximum grain yield of 484 g m^{-2} in
WAR20s1. Trial-mean reductions in IGW between s2 and s1 for PEM and conventional plot
trials were maximum at WAR18 (64%) and GAT19 (47%), respectively (Fig 3a & b). Grain
yield reduction between s2 and s1 was maximum in GAT18 for the PEM and in TOS19 for
conventional plots (Fig 3c & d).

All trials were fully irrigated, except TOS19, which only received pre-flowering supplementary
irrigation and was subjected to a post-flowering drought. Given the pre-flowering conditions,
s1 plants at TOS19 produced a high number of grains but significant reduction in grain size
was confounded by post-flowering heat and drought.

Extending the photoperiod increased allowed tagging of plants at a matched development
stage from genotypes contrasting maturity

Phenology data were collected from plants 0.5 m from each end of experimental rows in the
PEM, from next to the supplemented light and at the far end away from the supplemental lights
(Fig. 4-5). The phenology of the different genotypes significantly varied both under natural and
supplemented light. Across all PEM trials, the earliest maturing genotypes flowered
approximately 10.1 days (172^{o}Cd) and 7.8 days (145^{o}Cd) earlier than the latest maturing
genotypes under natural and supplemented lights (Fig. 4). The supplemented light accelerated
flowering by 7.9 days (152^{o}Cd) and 5.4 days (114^{o}Cd) on average in s1 and s2, respectively
(averaged across genotypes and trials). This gap between flowering times of plants under
natural and supplemented light allowed multiple tagging of the genotypes at a matched
developmental phase (Zadok 65) from a single time of sowing.

Supplemented light had a weaker effect in late sowings when the natural photoperiod was
already longer (Fig. 5). Under the shortest studied photoperiods (<10.5 h at sowing), plants
closer to the lights flowered approximately 8.8 days earlier than plants away from the lights
(averaged across genotypes and trials). The phenology effect of supplemented light was

reduced by approximately 49% when plants were planted under longer photoperiod (11.5 h or
more) at sowing.

A reduction in individual grain weight by 1.5 mg for every post-flowering heat day

A strong correlation ($r^2 = 0.90$) between the number of post-flowering days with maximum
temperature >30°C and individual grain weight was observed across PEM trials with
individual-spike harvests (Fig. 6a). Across all trials and tagging events, IGW was reduced by
1.5 mg for every additional post-flowering hot day (with maximum temperature >30°C).
Stems receiving 0-4 post-flowering hot days (HET1), produced the largest grains (40-45 mg;
Fig. 6a). Given that any heat event affecting these plants were very brief (Fig. 2), IGW
produced during this period is likely to be close to the potential grain weight. Stems with a
moderate numbers of heat events between 5 and 15 post-flowering hot days (HET2) had
significantly lower mean IGW and total grain weight per spike than those in HET1 (Fig. 6).
The most stressed stems experienced greater than 15 hot days (HET3, Fig. 3). In GAT19s2 and
TOS19s2, most plants experienced 20 and 22 post-flowering hot days, respectively and
suffered maximum reduction in IGW. On average, stems under these extremely hot conditions
produced 2.7 times smaller grains than the potential IGW (Fig. 6a).

While greatest environmental variations occurred between sites and sowings, variations in the
number of hot days and mean IGW were also observed among stems tagged at different times
(i.e. group of stems flowering at different dates) within a single sowing time. For instance, T2
stems of GAT19s1 produced 14% smaller grains compared with T1 stems (Fig. 6a, Table S3).

The photoperiod-extension method provided a stable genotype ranking within heat environment types

Genotypic rankings for IGW assessed using the PEM were relatively consistent between environments within heat environment types, i.e. environments experiencing similar heat stress (Fig. 7). For single spike harvests with the PEM, IGW correlations for stems experiencing mild or no heat stress during grain-filling (HET1), ranged from r of 0.64 to 0.89 (Fig. 7a). In HET2, r correlations ranged from 0.2 to 0.90 for fully-irrigated trials (i.e. all trials except TOS19 which experienced a post-flowering drought). These correlations were stronger (0.57-0.9) among the irrigated trials with a similar number of hot days (i.e. 9-13) and became more variable with the environment (GAT19s1T1) with five hot days only (r of 0.2-0.57). IGW of the two severely-stressed environments (HET3) were strongly correlated ($r = 0.46$, Fig 7a). Correlations between HET1, HET2 and HET3 varied, with moderate to high correlations between HET1 and HET2 environments and weaker correlations with HET3 environments.

IGW from PEM quadrat harvests were also positively correlated between the different tested environments, and these correlations were particularly strong among environments experiencing a similar number of hot days (Fig. 7b). For example, in HET1, these correlations were moderate to strong (r of 0.24-0.88) mainly because of a weaker correlation between GAT20s1 and WAR20s1 ($r = 0.24$). Similarly, all fully-irrigated HET2 environments were positively correlated with correlations ranging from 0.19 to 0.68.

In the conventional plot trials, genotype ranking for IGW varied widely across environments, both between sowings and sites (Fig. 7c). Correlations between non-stressed environments (HET1) were relatively high (r of 0.4-0.78). However, correlations between HET2 environments were highly variable or even negative (r of -0.15 to 0.42). Similarly, HET3 trials GAT19s2 and TOS19s2 had a correlation of 0.10.

Grain yield from PEM quadrat harvests for irrigated trials were also positively correlated. The correlations ranged from 0.18 to 0.72 within HET1, from 0.33 to 0.63 within HET2 (Fig. S2b, supplementary data). In contrast, for conventional plots, correlations for grain yield were either poor or negative for most trials, irrespective to their heat environment types (Fig. 7c). For instance, maximum correlations within HET1, HET2 and HET3 were 0.3, 0.21 and -0.36, respectively.

Genotype rankings across methods were also compared. Strong positive relationships (r^2 from 0.65 to 0.97 depending on the environment considered) were observed between the IGW from PEM trials with (i) individual spike harvest and (ii) quadrat harvest under the fully irrigated

313   conditions (Fig. 8a). In contrast, correlations for IGW between PEM quadrat harvest and

314   conventional plot trials under irrigated conditions were poorer in most trials (r^2 from 0.19 to

315   0.57) (Fig. 8b.

The photoperiod-extension method allowed genotypes to be discriminate according to their performance in different type of heat environments

Performance of wheat genotypes in terms of IGW and total grain weight were analysed using a principal component analysis (PCA, Fig. 9). The first two principle components (PC) combined explained more than 50% variance across environments in both IGW and grain yield for all the tested methods, except for grain yield in conventional field plots (Fig. 9).

For PEM spike harvests, strong positive correlations were observed between fully-irrigated HET1 and HET2 environments as indicated by the small angles between the vectors in the biplots (Fig. 9a & d). These correlations were stronger within HETs for IGW (Fig. 9a) than for total grain weight per spike (Fig. 9d). In contrast, a wider angle between vectors for HET3 and HET1-HET2 suggested weaker correlations between these environments (Fig. 9 a & d). Drought affected TOS19s1, which also experienced moderate post-flowering heat stress (HET2) better correlated with HET3 environments severely stressed by heat both for IGW and total grain weight per spike.

The projection of a genotype onto an environmental axis reflects the performance of that genotype in that environment. Genotypes were grouped based on their performance in environments from the PEM spike harvests. Biplots separated genotypes into three distinct groups for IGW and three partly differentiated groups for grain weight per spike (Fig. 9a & d). Top performing genotypes in HET1 and HET2 such as ZWB10-37 (in green in Fig. 9) project above the origin on axes corresponding to HET1 and HET2. In contrast, poorly performing genotypes such as Yitpi and EGA Wylie (in red in Fig. 9) project below the origin on the same axes. The top and poor performing genotypes identified with the PEM spike harvests were consistent with those identified from the PEM quadrat harvests data (Fig. 9b & e).

For the conventional plots, heat environment types (HET) were less clearly differentiated, particularly for grain yield, indicating less power to distinguish between HETs (Fig. 9c & f). Genotype rankings were also changed compared to ranking with the PEM. For instance, Suntop_1 ranked among the top performing genotypes for IGW in HET1 and HET2 in the conventional plot trials (Fig. 9c) but it ranked with the intermediate to poor group with the PEM (Fig. 9a). For grain yield in conventional plot trials, genotype performance was even more difficult to differentiate. For example, top performing genotypes in the PEM (in green) were

347 not distinct from others in conventional plots being scattered almost randomly across the biplot

348 (Fig. 9f).

Discussion

Impact of post-flowering heat events on individual grain weight and yield

Crop simulation studies identified post-flowering heat as the major determinant of wheat productivity in the Australian wheatbelt for current and future climates (Ababaei and Chenu, 2020; Collins and Chenu, 2021). Data from the current study demonstrates experimentally how heat intensity during the grain filling period is a major determinant for IGW and grain yield in irrigated field conditions (Fig. 6), agreeing with previous reports for crops grown under rainfed field conditions (Gebeyehou et al., 1982; Gerard et al., 2020; Telfer et al., 2018; Thistlethwaite et al., 2015).

In wheat, temperatures above 30°C negatively impact different processes during the grain filling period (Porter and Gawith, 1999; Girousse et al., 2021). In the tested conditions, under full irrigation, wheat genotypes did not experience any significant loss in IGW or grain yield in response to 0-4 hot days (maximum daily temperature $> 30°C$) during grain filling (HET1, Fig. 6a). These heat events were either very brief (1-2 h only in total from 0 to 450°Cd after flowering) or the heat event occurred late in development when grain filling was already well advanced i.e. $> 450°Cd$ after flowering (Fig. 2). By contrast, in environments with 5 or more days with maximum temperatures $>30°C$ between flowering and 500°C after flowering (≥ 4 cumulated hours of temperature $>30°C$; i.e. HET2 & HET3), wheat genotypes were substantially impacted. A significant and strong effect of the timing of heat stress has also been recorded for wheat IGW during early to mid grain filling in both controlled (Stone and Nicolas, 1996) and field conditions (Thistlethwaite et al., 2015). In the field conditions tested in the current study, each additional hot day during grain filling reduced IGW by 1.5 mg (Fig. 6a). This significant ($p<0.001$) reduction in IGW was also found in plants sown at single date but exposed to a range of natural heat events. For example, in GAT19s1, stems tagged one week apart (taggings 1 & 2) experienced different levels of heat stress, with stems from tagging 2 being subjected to four additional days of post-flowering heat and producing grain 14% smaller than stems from tagging 1 (Fig. 6a, Table S3). Developmental-phase-specific effects of heat shocks on IGW and grain yield were observed in this study, particularly for moderate stress in HET2 (data not shown). This highlights the importance of screening for heat tolerance at matched developmental phases. High heat sensitivity during early grain filling has also been reported by Talukder et al. (2013), who observed a single hot day (maximum temperature $> 35°C$) during this period can significantly reduce grain weight of wheat crops.

A new method to screen for heat tolerance at matched development stages

During both the reproductive and the grain filling phases, sensitivity of developing grains to heat events can change over a period as short as a few days (Stone and Nicolas, 1995; Chenu and Oudin, 2019). Since field-based techniques for screening heat tolerant germplasm generally rely serial sowings, it is hard to optimise sowing time given the unpredictable timing of heat events. Field-based screening for heat tolerance using this conventional method is further complicated when genotypes with varying maturity types are tested together. These genotypes are likely to be at different developmental stages when natural heat events occur. Thus, heat escape by being at a less sensitive stage can confound comparisons of heat tolerance, *per se*.

We developed a new method using supplemental lights that allows screening of wheat genotypes at matched developmental phases. Photoperiod was extended to 20 h at one end of test rows of plots. The light intensity diminishes with the square of distance (Niinemets and Keenan, 2012), generating a gradient of flowering times along the length of the test rows (Fig. 1 & 4). The range of flowering times within a single row of individual genotype, allowed comparison of the performance of genotypes with varying maturity types at matched developmental stages. The method was tested for sowing dates from late May to mid-September. The impact of extended photoperiod on flowering was associated with the natural photoperiod during vegetative crop growth ($r^2 = 0.32$; Fig. 5). A wide gap in flowering along the rows, of up to 8.8 days when averaged across all tested genotypes, was recorded when planting under short photoperiod (<10.50 h). This gap narrowed by 3.5 days with each increasing hour in photoperiod at later sowing dates (Fig. 5), so that all genotypes flowered within only 4.5 days of each other in the latest sowing dates tested. A wider flowering-time gradient is interesting as it allows; (i) multiple taggings within a single time of sowing, which may increase the probability of being able to screen for heat at a particular stage, (ii) more flexibility for operators to visit the trial during the appropriate window and tag all genotypes at a matched development stage, and/or (iii) a wider range of maturity types to be considered. Despite this additional flexibility, the robust genotype rankings generated with the PEM suggest that a single tagging is likely sufficient for reliable screening.

In the conditions tested, sowing dates for screening post-flowering heat tolerance between early July and late August provided the best discrimination. Sowings during this period were typically associated with moderate post-flowering heat stress (HET2) and a relatively wide flowering gradient for tagging stems or quadrats at a matched development stage. With earlier

sowings, very few heat events occurred. In contrast, later sowings had a narrower flowering gradient and a greater risk of exposure to a high number of severe heat events which can increase heat damage to a level where variation between genotypes is reduced. Late sowings are also usually more prone to pre-flowering heat (e.g. TOS19s2 and GAT19s2), thus resulting in confounding effects, with IGW varying due to both a decrease in grain number and direct effects of post-flowering heat stress. Optimum sowing dates to screen heat stress obviously depend on the targeted stress (e.g. with or without pre-flowering heat). They also depend on the test location that impacts both wheat phenology and the frequency of heat events. Crop models can help to identify locations and sowing windows for efficient screening of heat tolerance in particular target environments (e.g. Chauhan et al., 2017; Chenu et al., 2017; Collins and Chenu, 2021).

The PEM method allows reliable ranking of wheat genotypes under varying environments

With the PEM, strong correlations among trials for either IGW or grain yield (Fig. 7a & b, S2a & b) indicated a stable ranking of tested genotypes particularly within the environments receiving a similar degree of heat (Supplementary tables S4, S5 & S7). For instance, for the PEM with individual-spike harvests, the respective mean r correlations within fully-irrigated HET1, HET2 and HET3 environments were 0.80, 0.59 and 0.46 for IGW; and 0.75, 0.54 and 0.57 for total grain weight per spike. In contrast, these correlations were typically much weaker in conventional plots with r averages of 0.53, 0.11 and 0.1 for IGW, and 0.05, 0.12 and -0.36 for grain yield within HET1, HET2 and HET3, respectively (Fig. 7c, S2c, Supplementary tables S6 & S8). This indicates that in conventional plots, ranking of wheat genotypes was highly variable depending on environmental conditions, including for the targeted moderate grain-filling heat stress (HET2).

The PEM allowed the tested genotypes to be clustered into distinct groups based on their performance in these environments (Fig. 9a, b & d, e). In contrast, this genotype clustering was inconsistent with that found in conventional plots. The clustering in the conventional plots was also much less powerful at separating groups of genotypes as seen by the greater overlap of genotypes within each group (Fig. 9c & f). This indicates that the PEM ranked wheat genotypes more consistently across a wide range of heat-stressed environments compared to conventional plot trials.

Implications for breeding

A high-throughput and accurate method for screening heat tolerance is necessary for sustaining food security under changing environments, particularly for projected hot and dry environments (Collins and Chenu, 2021). The PEM was tested using single rows (hand planted) and small plots (machine planted) with tagging of either individual stems or small quadrats at distances from the lights where development was matched at the flowering stage. The most reliable genotype ranking was achieved by tagging and harvesting individual spikes (Fig. 7a, 9a & d, S2a). However, strong positive relationships for either IGW (r^2 from 0.65 to 0.97) or total grain weight (r^2 from 0.34 to 0.97 across environments) between individual-spike and quadrat harvests were observed for the PEM (Fig. 8a & S1), suggesting that reliable screening may be done with the PEM for small plot with quadrat harvest. This was supported by the results, as genotype ranking with quadrat harvest was also robust (Fig. 7b, 9b & e, S2b) with strong genetic correlations between tested environments (average r of 0.51 in HET1 and 0.53 in HET2 for IGW, and 0.47 and 0.36 for grain yield). Using the PEM, with genotypes sown in small plots with machinery, and harvested from plot segments tagged at a matched developmental stage could have great potential to be scaled-up for large number of genotypes.

In the tested conditions, the PEM allowed effective screening for post-flowering heat stress (HET2), which is the main type of heat stress targeted for Australian production regions in current and projected climates (Collins and Chenu, 2021). Robust screening was also performed for non-stressed environments (HET1) and severely stressed environments (HET3) that happened to be affected by both pre- and post-flowering heat in the tested conditions. Different genotype rankings were observed between heat environment types HET2 and HET3, highlighting the importance of screening in the relevant target HET. The occurrence of water stress also strongly impacted genotype ranking (e.g. Fig. 2 and S2), suggesting that heat and drought adaptations are at least partly regulated by different processes. This highlights the importance of understanding the physiology and genetics associated to heat tolerance, drought tolerance and their interaction.

The PEM described here offers an opportunity to select for heat tolerant wheat genotypes more reliably than can be done conventionally. This method could be adjusted and deployed in other regions and other crops, with sowing dates adapted to the targeted heat environment type.

Conclusions

A new field-based method was developed and tested to screen wheat genotypes for post-

flowering heat tolerance under natural heat events. In this method, supplemental light was used to manipulate crop phenology in a way that allowed genotypes with varying phenology to be tested at closely matched developmental stage when a heat event occurred. IGW of wheat genotypes tested in this study was highly sensitive to the number of hot days during early-to-mid grain filling, particularly in the most relevant heat environment for Australian wheat crops (HET2). With the PEM, strong genetic correlations were found between irrigated environments of similar heat stress, with respective r correlations within HET1, HET2 and HET3 averaging 0.8, 0.59 and 0.46 for IGW; and 0.75, 0.54 and 0.57 for total grain weight. By contrast these correlations were substantially weaker for conventional yield plots (average r of 53, 0.11 and 0.1 for IGW; and 0.05, 0.12 and -0.36 for grain yield). Clustering of genotypes across tested environments with a PCA further highlighted the ability of the PEM to differentiate genotypes based on their performance in similar heat-stress environments. In contrast, ranking substantially changed between environments with moderate (HET2) and severe (HET3) heat stress. Similarly, genotype rankings greatly varied between fully-irrigated HET2 environments and HET2 environments subjected to a post-flowering water stress.

With increasing frequency of post-flowering heat stress affecting wheat growing regions, the photoperiod-extension method promises to improve the efficiency of heat tolerance field screening, particularly when comparing genotypes of different maturity types.

Acknowledgments

The research was made possible thanks to the support of The University of Queensland and the Queensland Government, Department of Agriculture and Fisheries. Najeeb Ullah was supported by a Queensland Government, Advance Queensland Fellowship. We acknowledge the assistance of Ian Broad (Department of Agriculture and Fisheries Queensland) with operating of weather stations across the studied sites, Brian Collins (James Cook University) for data analysis and Thaís Helena Godoy Sanches for sample processing and data collection.

References

Ababaei, B., Chenu, K., 2020. Heat shocks increasingly impede grain filling but have little effect on grain setting across the Australian wheatbelt. Agric. For. Meteorol. 284, 107889. https://doi.org/10.1016/j.agrformet.2019.107889

Alduchov, O.A., Eskridge, R.E., 1996. Improved Magnus form approximation of saturation vapor pressure. J. Appl. Meteorol. https://doi.org/10.1175/1520-0450(1996)035<0601:IMFAOS>2.0.CO;2

Bennett, D., Reynolds, M., Mullan, D., Izanloo, A., Kuchel, H., Langridge, P., Schnurbusch, T., 2012.
Detection of two major grain yield QTL in bread wheat (Triticum aestivum L.) under heat,
drought and high yield potential environments. Theor. Appl. Genet.
https://doi.org/10.1007/s00122-012-1927-2

Chapman, S.C., Chakraborty, S., Dreccer, M.F., Howden, S.M., 2012. Plant adaptation to climate
changeopportunities and priorities in breeding, in: Crop and Pasture Science.
https://doi.org/10.1071/CP11303

Chauhan, Y., Allard, S., Williams, R., Williams, B., Mundree, S., Chenu, K., Rachaputi, N.C., 2017.
Characterisation of chickpea cropping systems in Australia for major abiotic production
constraints. F. Crop. Res. https://doi.org/10.1016/j.fcr.2017.01.008

Chenu, K., Oudin, F., 2019. Heat impact on yield components of fertile primary tillers in wheat can
inform crop modelling for future climates 23–26.

Chenu, K., Porter, J.R., Martre, P., Basso, B., Chapman, S.C., Ewert, F., Bindi, M., Asseng, S., 2017.
Contribution of Crop Models to Adaptation in Wheat. Trends Plant Sci.
https://doi.org/10.1016/j.tplants.2017.02.003

Christopher, J., Richard, C., Chenu, K., Christopher, M., Borrell, A., Hickey, L., 2015. Integrating
Rapid Phenotyping and Speed Breeding to Improve Stay-Green and Root Adaptation of Wheat
in Changing, Water-Limited, Australian Environments. Procedia Environ. Sci.
https://doi.org/10.1016/j.proenv.2015.07.246

Christopher, M., Paccapelo, V., Kelly, A., Macdonald, B., Hickey, L., Richard, C., Verbyla, A.,
Chenu, K., Borrell, A., Amin, A., Christopher, J., 2021. QTL identified for stay-green in a multi-
reference nested association mapping population of wheat exhibit context dependent expression
and parent-specific alleles. F. Crop. Res. 270, 108181. https://doi.org/10.1016/j.fcr.2021.108181

Collins, B., Chapman, S., Hammer, G., Chenu, K., 2021. Limiting transpiration rate in high
evaporative demand conditions to improve Australian wheat productivity. in silico Plants.
https://doi.org/10.1093/insilicoplants/diab006

Collins, B., Chenu, K., 2021. Improving productivity of Australian wheat by adapting sowing date
and genotype phenology to future climate. Clim. Risk Manag.
https://doi.org/10.1016/j.crm.2021.100300

Djanaguiraman, M., Vara Prasad, P. V., Murugan, M., Perumal, R., Reddy, U.K., 2014. Physiological
differences among sorghum (Sorghum bicolor L. Moench) genotypes under high temperature
stress. Environ. Exp. Bot. https://doi.org/10.1016/j.envexpbot.2013.11.013

Farooq, M., Bramley, H., Palta, J.A., Siddique, K.H.M., 2011. Heat stress in wheat during

reproductive and grain-filling phases. CRC. Crit. Rev. Plant Sci. 30, 491–507.

https://doi.org/10.1080/07352689.2011.615687

Field, C.B., Barros, V., Stocker, T.F., Dahe, Q., Jon Dokken, D., Ebi, K.L., Mastrandrea, M.D., Mach,

K.J., Plattner, G.K., Allen, S.K., Tignor, M., Midgley, P.M., 2012. Managing the risks of

extreme events and disasters to advance climate change adaptation: Special report of the

intergovernmental panel on climate change, Managing the Risks of Extreme Events and

Disasters to Advance Climate Change Adaptation: Special Report of the Intergovernmental

Panel on Climate Change. https://doi.org/10.1017/CBO9781139177245

Gebeyehou, G., Knott, D.R., Baker, R.J., 1982. Relationships among Durations of Vegetative and

Grain Filling Phases, Yield Components, and Grain Yield in Durum Wheat Cultivars 1 . Crop

Sci. https://doi.org/10.2135/cropsci1982.0011183x002200020021x

Gerard, G.S., Crespo-Herrera, L.A., Crossa, J., Mondal, S., Velu, G., Juliana, P., Huerta-Espino, J.,

Vargas, M., Rhandawa, M.S., Bhavani, S., Braun, H., Singh, R.P., 2020. Grain yield genetic

gains and changes in physiological related traits for CIMMYT's High Rainfall Wheat Screening

Nursery tested across international environments. F. Crop. Res.

https://doi.org/10.1016/j.fcr.2020.107742

Girousse, C., Inchboard, L., Deswarte, J.-C., Chenu, K., 2021. How does post-flowering heat impact

grain growth and its determining processes in wheat? J. Exp. Bot.

https://doi.org/10.1093/jxb/erab282

Guo, Z., Slafer, G.A., Schnurbusch, T., 2016. Genotypic variation in spike fertility traits and ovary

size as determinants of floret and grain survival rate in wheat. J. Exp. Bot. 67, 4221–4230.

https://doi.org/10.1093/jxb/erw200

Jamieson, P.D., Brooking, I.R., Porter, J.R., Wilson, D.R., 1995. Prediction of leaf appearance in

wheat: a question of temperature. F. Crop. Res. https://doi.org/10.1016/0378-4290(94)00102-I

Limpens, J., Granath, G., Aerts, R., Heijmans, M.M.P.D., Sheppard, L.J., Bragazza, L., Williams,

B.L., Rydin, H., Bubier, J., Moore, T., Rochefort, L., Mitchell, E.A.D., Buttler, A., van den

Berg, L.J.L., Gunnarsson, U., Francez, A.J., Gerdol, R., Thormann, M., Grosvernier, P.,

Wiedermann, M.M., Nilsson, M.B., Hoosbeek, M.R., Bayley, S., Nordbakken, J.F., Paulissen,

M.P.C.P., Hotes, S., Breeuwer, A., Ilomets, M., Tomassen, H.B.M., Leith, I., Xu, B., 2012.

Glasshouse vs field experiments: Do they yield ecologically similar results for assessing N

impacts on peat mosses? New Phytol. https://doi.org/10.1111/j.1469-8137.2012.04157.x

Lobell, D.B., Hammer, G.L., Chenu, K., Zheng, B., McLean, G., Chapman, S.C., 2015. The shifting

influence of drought and heat stress for crops in northeast Australia. Glob. Chang. Biol. 21,

4115–4127. https://doi.org/10.1111/gcb.13022

Najeeb, U., Tan, D.K.Y., Sarwar, M., Ali, S., 2019. Adaptation of crops to warmer climates:

Morphological and physiological mechanisms, in: Sustainable Solutions for Food Security:

Combating Climate Change by Adaptation. https://doi.org/10.1007/978-3-319-77878-5_2

Niinemets, Ü., Keenan, T.F., 2012. Measures of light in studies on light-driven plant plasticity in

artificial environments. Front. Plant Sci. https://doi.org/10.3389/fpls.2012.00156

Passioura, J.B., 2006. The perils of pot experiments. Funct. Plant Biol.

Poorter, H., Fiorani, F., Pieruschka, R., Wojciechowski, T., van der Putten, W.H., Kleyer, M., Schurr,

U., Postma, J., 2016. Pampered inside, pestered outside? Differences and similarities between

plants growing in controlled conditions and in the field. New Phytol.

https://doi.org/10.1111/nph.14243

Porter, J.R., Gawith, M., 1999. Temperatures and the growth and development of wheat: A review.

Eur. J. Agron. https://doi.org/10.1016/S1161-0301(98)00047-1

Prasad, P.V.V., Djanaguiraman, M., 2014. Response of floret fertility and individual grain weight of

wheat to high temperature stress: Sensitive stages and thresholds for temperature and duration.

Funct. Plant Biol. https://doi.org/10.1071/FP14061

Ray, D.K., Gerber, J.S., Macdonald, G.K., West, P.C., 2015. Climate variation explains a third of

global crop yield variability. Nat. Commun. https://doi.org/10.1038/ncomms6989

Richard, C., 2017. Breeding wheat for drought adaptation: Development of selection tools for root

architectural traits. https://doi.org/10.14264/uql.2017.1055

Saini, H., Aspinall, D., 1982. Sterility in Wheat (Triticum aestivum L.) Induced by Water Deficit or

High Temperature: Possible Mediation by Abscisic Acid. Funct. Plant Biol.

https://doi.org/10.1071/pp9820529

Sofield, I., Evans, L., Cook, M., Wardlaw, I., 1977. Factors Influencing the Rate and Duration of

Grain Filling in Wheat. Funct. Plant Biol. https://doi.org/10.1071/pp9770785

Stone, P.J., Nicolas, M.E., 1998. The effect of duration of heat stress during grain filling on two wheat

varieties differing in heat tolerance: Grain growth and fractional protein accumulation. Aust. J.

Plant Physiol. https://doi.org/10.1071/PP96114

Stone, P.J., Nicolas, M.E., 1996. Effect of timing of heat stress during grain filling on two wheat

varieties differing in heat tolerance. II. Fractional protein accumulation. Aust. J. Plant Physiol.

https://doi.org/10.1071/PP9960739

Stone, P.J., Nicolas, M.E., 1995. Effect of timing of heat stress during grain filling on two wheat
varieties differing in heat tolerance. I. Grain growth. Aust. J. Plant Physiol.
https://doi.org/10.1071/PP9950927

Streck, N.A., 2005. Climate change and agroecosystems: the effect of elevated atmospheric CO2 and
temperature on crop growth, development, and yield. Ciência Rural.
https://doi.org/10.1590/s0103-84782005000300041

Talukder, A.S.M.H.M., McDonald, G.K., Gill, G.S., 2013. Effect of short-term heat stress prior to
flowering and at early grain set on the utilization of water-soluble carbohydrate by wheat
genotypes. F. Crop. Res. 147, 1–11. https://doi.org/10.1016/j.fcr.2013.03.013

Tashiro, T., Wardlaw, I., 1990. The Response to High Temperature Shock and Humidity Changes
Prior to and During the Early Stages of Grain Development in Wheat. Funct. Plant Biol.
https://doi.org/10.1071/pp9900551

Team, R.C., 2018. R Development Core Team. R: A language and environment for statistical
computing. R Foundation for Statistical Computing, Vienna, Austria; 2014. Google Sch.

Telfer, P., Edwards, J., Bennett, D., Ganesalingam, D., Able, J., Kuchel, H., 2018. A field and
controlled environment evaluation of wheat (Triticum aestivum) adaptation to heat stress. F.
Crop. Res. https://doi.org/10.1016/j.fcr.2018.09.013

Telfer, P., Edwards, J., Norman, A., Bennett, D., Smith, A., Able, J.A., Kuchel, H., 2021. Genetic
analysis of wheat (Triticum aestivum) adaptation to heat stress. Theor. Appl. Genet.
https://doi.org/10.1007/s00122-021-03778-2

Thistlethwaite, R.J., Tan, D.K.Y., Bokshi, A.I., Ullah, S., Trethowan, R.M., 2020. A phenotyping
strategy for evaluating the high-temperature tolerance of wheat. F. Crop. Res.
https://doi.org/10.1016/j.fcr.2020.107905

Thistlethwaite, R.J., Tan, D.K.Y., Buckley, T.N., Trethowan, R.M., 2015. Identification of Genetic
Variation in Heat Stress and Mechanisms of Tolerance in Wheat. Procedia Environ. Sci. 29, 30.
https://doi.org/10.1016/j.proenv.2015.07.139

Ullah, N., Ababaei, B., Chenu, K., 2020. Increasing Heat Tolerance in Wheat to Counteract Recent
and Projected Increases in Heat Stress † 3390. https://doi.org/10.3390/proceedings2019036132

Wheeler, T.R., Batts, G.R., Ellis, R.H., Hadley, P., Morison, J.I.L., 1996. Growth and yield of winter
wheat (Triticum aestivum) crops in response to CO2 and temperature. J. Agric. Sci.
https://doi.org/10.1017/s0021859600077352

Zadoks, J.C., Chang, T.T., Konzak, C.F., 1974. A decimal code for the growth stages of cereals.

Weed Res. https://doi.org/10.1111/j.1365-3180.1974.tb01084.x

Zheng, B., Chenu, K., Chapman, S.C., 2016. Velocity of temperature and flowering time in wheat -
assisting breeders to keep pace with climate change. Glob. Chang. Biol.
https://doi.org/10.1111/gcb.13118

Zheng, B., Chenu, K., Fernanda Dreccer, M., Chapman, S.C., 2012. Breeding for the future: What are
the potential impacts of future frost and heat events on sowing and flowering time requirements
for Australian bread wheat (Triticum aestivium) varieties? Glob. Chang. Biol.
https://doi.org/10.1111/j.1365-2486.2012.02724.x

1
2 Figure 1: Layout of the photoperiod-extension method (PEM). Wheat genotypes were sown either in
3 a single row (as in this picutre) or in narrow plots of four rows. In the centre of the trial at the end of
4 each test row, at the central axis of the trial, LED lights were setup. These supplemental lights extend
5 the photoperiod to 20 h. The intensity of light diminishes along the row and induces a gradient of
6 flowering times within each test row, with plants closest to the light developing faster.
7

8
9 Figure 2: The number of hours each day when air temperatures exceeded 26°C (blue lines), 30°C (orange
10 lines) or 35°C (red lines) is plotted against thermal time (°Cd) from sowing. These are presented for
11 each location, season and sowing time. The vertical lines represent tagging of wheat genotypes at
12 flowering. T1, 1st cohort of stems tagged at flowering, T2: 2nd cohort of stems tagged at flowering, T3:
13 3rd cohort of stems tagged at flowering. Trial identifiers are as described in Table 1.
14

Figure 3: Effect of sowing time on (a, c) individual grain weight and (b, d) grain yield of tested wheat genotypes for trials with (b, d) conventional pots or (a, b) the photoperiod-extension method with quadrat tagging and harvests. Each boxplot displays data from 32 genotypes and four independent replicates. TOS19 crops only had supplementary irrigation and experienced a mild post-flowering water stress. In the boxplot, horizontal black lines insides each box denote median values; boxes extend from the 25th to the 75th percentile of each group's distribution of values; the whiskers the 10th and 90th percentiles; and the dots outside the whiskers represent individual values outside this range. ** corresponds to significant differences at *P*< 0.001 between the two sowing times within each trial. In 2018, s2 plots were established only at Gatton (GAT18) and in 2019, quadrat harvests were taken from the s1 PEM trials at Tosari (TOS19) only. Trial identifiers are as described in Table 1.

Figure 4: Flowering time of wheat genotypes with and without supplemented light. Horizontal dashed lines correspond to the day of tagging of all wheat genotypes at matched flowering. Data on flowering were collected for plants 0.5 m away from the light source (extended photoperiod, red) and plants 0.5 m from the end of the row (natural light, blue). Values correspond to the mean of four independent replicates ± confidence interval (95%). s1 and s2 represent sowing 1 and sowing 2, respectively.

Figure 5: Delay in flowering time due to supplemented light in response to the photoperiod measured at sowing in all trials and sowings. The delay in flowering was calculated as the difference in flowering dates for plants located 0.5 m (i.e. extended light, 20 h) and 4.5 m (i.e. natural light) away from the lights. Data correspond to the mean of 32 (2018 and 2019) or 20 (2020) genotypes and four independent replicates for all PEM plots. TOS19 crops only had supplementary irrigation and experienced a mild post-flowering water stress.

1
2   Figure 6: Changes in (a) individual grain weight and (b) spike grain weight, in response to the number of post-flowering hot days (0-500°Cd after flowering)
3   across locations and sowing times. Each point represents the mean of all genotypes for individual grain weight of spikes that flowered the same day (four
4   independent replicates of 20 stems each). Heat environment types are indicated by the horizontal bar at the top of each panel (HET1, green; HET2, orange;
5   HET3, red). TOS19 only had supplementary irrigation and experienced a mild post-flowering water stress.

Figure 7: Genetic Pearson's correlation coefficients (r) for individual grain weight between environments. Correlations for mean individual grain weight of genotypes between each pair of tested environments (i.e. site x year x sowing x tagging-event combinations). Individual grain weight was estimated from measurements at (a) spike level, and at (b) crop level with quadrat harvests in the photoperiod-extension method (PEM), as well as at (c) the plot level in the conventional plot trials. Below the heat maps, are indicated the number post-flowering hot days with maxima exceeding 30°C from 0 to 500°Cd after flowering for each environment, grouped by heat environment types are indicated by dashed, coloured boxes (HET1, green; HET2, orange; HET3, red). All environments were fully irrigated except TOS19 (framed in brown), which experienced a mild post-flowering water stress. Genetic correlations for total grain weight are presented in Figure S2.

15
16 Figure 8: Correlations for individual grain weight of all studied genotypes between either (a) data collected from individually tagged stems and tagged quadrats
17 at the matched development stage with the photoperiod-extension method (PEM), or between (b) the PEM with quadrat tagging and harvests and conventional
18 plots. Each data point represents the genotypic mean value of four independent replicates. Correlations for total grain weight are presented in Figure S1.

Figure 9: Principal component analysis biplots of (a-c) individual grain weight (IGW) and (d) grain weight per spike or (e-f) grain yield of studied wheat genotypes for the PEM spike harvests (a, d), PEM quadrat harvests (b, e) and conventional machine harvested plot trials (c, f). Environments corresponded to

combinations of sowing dates, sites and years (b, c, e, f) together with tagging events for single-spike harvests (a, d). Principal component loadings (arrows) were coloured and grouped based on heat environment type; HET1, green; HET2, orange; HET3, red. Genotypes were grouped for similarity in their performance for either IGW (a-c) or total grain weight (d-f) in PEM spike harvests, i.e. green, top performing genotypes in HET1 and HET2, blue, genotypes with inconsistent / poor performance, particularly across HET1 and HET2; red, genotypes with poor performance across most tested environments. Eclipse colours correspond to the performance groups for genotypes.

Table 1. Trial characteristics, including the trial identifier (Trial), site, irrigation treatment, sowing date and the types of trial conducted (i.e. PEM with tagging and harvesting of either single spikes or quadrates; and conventional plots). Also presented are mean and max daily temperature, day-time vapour pressure deficit (VPD) and radiation from sowing to maturity, as well as the mean duration of the pre- and post-flowering periods, and the heat environment type (HET) for the first tagging of trials with the photoperiod-extension method (PEM). Days to flowering and post-flowering duration were calculated from sowing to flowering and flowering to maturity, respectively. TOS19 trial only had supplementary pre-flowering irrigation and experienced a mild post-flowering water stress.

Trial	Site	Sowing	Trial type	Sowing date	Irrigation	Mean temp. (°C)	Mean daily max temp. (°C)	Mean VPD (kPa)	Mean daily radiation (MJ m^{-2})	Days to flowering (days)	Post-flowering duration (days)	HET
GAT18-s1*	Gatton	s1	Spike & quadrate PEM	03/07/2018	Full	19.3	26.3	0.74	23.8	73	42	HET1
GAT18-s2	Gatton	s2	Spike & quadrate PEM & plots	31/08/2018	Full	23.1	31.4	1.38	23.2	53	39	HET2
WAR18-s1	Warwick	s1	Spike & quadrate PEM	16/07/2018	Full	18.6	25.6	0.84	17.7	73	39	HET1
WAR18-s2	Warwick	s2	Spike & quadrate PEM	12/09/2018	Full	22.0	30.3	1.52	21.2	52	37	HET2
GAT19-s1	Gatton	s1	Spike PEM & plots	09/07/2019	Full	19.5	28.7	1.40	16.8	71	39	HET2
GAT19-s2	Gatton	s2	Spike PEM & plots	03/09/2019	Full	23.7	33.8	2.23	20.8	59	35	HET3
TOS19-s1	Tosari	s1	Spike & quadrate PEM & plots	16/07/2019	Supplementary	21.1	30.2	1.26	17.7	79	35	HET2
TOS19-s2	Tosari	s2	Spike PEM & plots	06/09/2019	Supplementary	26.5	36.7	2.27	16.2	62	35	HET3
GAT20-s1	Gatton	s1	Quadrate PEM & plots	26/05/2020	Full	17.9	26.8	1.08	17.3	79	49	HET1
GAT20-s2	Gatton	s2	Quadrate PEM & plots	04/08/ 2020	Full	22.2	31.2	1.35	19.7	65	37	HET2
WAR20-s1	Warwick	s1	Quadrate PEM & plots	08/06/ 2020	Full	19.3	26.3	0.74	13.6	73	42	HET1
WAR20-s2	Warwick	s2	Quadrate PEM & plots	12/08/2020	Full	23.1	31.4	1.38	11.6	53	39	HET1

Trial identifiers indicate site (Gatton (GAT), Tosari (TOS) or Warwick (WAR)), year conducted and sowing time (s1 or s2).