Functional redundancy of WDR31 with GTPase-activating proteins ELMOD and RP2 in

regulating IFT trains via BBSome

Sebiha Cevik[1], Xiaoyu Peng[2], Tina Beyer[3], Mustafa S. Pir[1], Ferhan Yenisert[1], Franziska Woerz[4],

Felix Hoffmann[3], Betul Altunkaynak[1], Betul Pir[1], Karsten Boldt[3], Asli Karaman[5], Miray

Cakiroglu[5], S. Sadik Oner[4,5], Ying Cao[2], Marius Ueffing[3], Oktay I. Kaplan[1]†

1- Rare Disease Laboratory, School of Life and Natural Sciences, Abdullah Gul University,

Kayseri, Turkey

2- School of Life Sciences and Technology, Tongji University, Shanghai 200092, China

3- Institute for Ophthalmic Research, Centre for Ophthalmology, University of Tuebingen,

Elfriede-Aulhorn-Strasse 7, D-72076 Tuebingen, Germany.

4- Goztepe Prof. Dr. Suleyman Yalcin City Hospital, Istanbul, Turkey

5- Science and Advanced Technology Application and Research Center, Istanbul Medeniyet

University, Istanbul

† Correspondence to oktay.kaplan@agu.edu.tr

Abstract

The correct intraflagellar transport (IFT) assembly at the ciliary base and the IFT turnaround at the ciliary tip are key for the IFT to perform its function, but we still have poor understanding about how these processes are regulated. Here, we identify WDR31 as a new ciliary protein, and analysis from zebrafish and *Caenorhabditis elegans* reveals the role of *WDR31* in regulating the cilia morphology. We find that loss of WDR-31 together with RP-2 and ELMD-1 (the sole ortholog ELMOD1-3) results in ciliary accumulations of IFT Complex B components and KIF17 kinesin, with fewer IFT/BBSome particles traveling along cilia in both anterograde and retrograde directions, suggesting that the IFT/BBSome entry into cilia and exit from cilia are impacted. Furthermore, anterograde IFT in the middle segment travel at increased speed in *wdr-31;rpi-2;elmd-1*. Remarkably, a non-ciliary protein leaks into cilia of *wdr-31;rpi-2;elmd-1* possible due to IFT defects. This work reveals WDR31-RP-2-ELMD-1 as IFT and BBSome trafficking regulators.

Introduction

Cilia are structurally and functionally distinct cellular projections, consisting of a microtubule-based axoneme extending from a centriole-derived basal body anchored at the plasma membrane. Cilia consist of multiple sub-compartments (basal body, transition zone, and ciliary tip) that display different protein compositions and structure (Blacque and Sanders, 2014; Rosenbaum and Witman, 2002; Satir and Christensen, 2007). Motile cilia mediate the movement of unicellular organisms such as *Chlamydomonas reinhardtii* or are involved in fluid movement across a tissue surface (Silflow and Lefebvre, 2001; Sleigh, 1989). Non-motile cilia, also known as primary cilia, possess mechanosensory, chemosensory, and osmosensory functions, and coordinate a range of extrinsic signaling pathways involved in cellular behavior, tissue development, and homeostasis such as those mediated by Hedgehog (Hh), Wnt, and receptor tyrosine kinase ligands (e.g. PDGFα) (Anvarian et al., 2019; Bloodgood, 2009; Nachury, 2014; Scholey, 2007).

The relation between cilia and human disorders has led to a greater understanding of their importance for human health. Both motile and primary cilia have been linked to the heterogeneous class of diseases known as ciliopathies, including Joubert Syndrome, Meckel Syndrome (MKS), and Nephronophthisis (NPHP). Owing to the presence of cilia on most cell types, ciliary defects result in varying multiorgan phenotypes such as kidney defects, retinitis pigmentosa, pancreatic cysts, hearing loss, congenital heart disease, and polydactyly (Reiter and Leroux, 2017; Wheway et al., 2019).

Over the last twenty years, there has been a large effort to reveal the molecular composition of cilia and its sub-compartments using several independent approaches, including clinical genomics, proteomics, functional genomics, and bioinformatics (Arnaiz et al., 2009; Avidor-Reiss et al.,

2004; Blacque et al., 2005; Breslow et al., 2018; Choksi et al., 2014; Jensen et al., 2016; Lambacher

et al., 2016; Li et al., 2004; Mick et al., 2015; Piasecki et al., 2010; Ruiz García et al., 2019;

Shaheen et al., 2016; Shamseldin et al., 2020; Sigg et al., 2017; UK10K Rare Diseases Group et

al., 2016; van Dam et al., 2019). Many proteins that make up the cilium, as well as many proteins

that regulate cilia biology, have been identified. These collective efforts have resulted in the

identification of 302 genes that are certain to be involved in cilia biogenesis, as well as over 180

ciliopathy genes. CiliaCarta estimates the total number of the ciliary genes to be about 1200 genes,

implying that many more ciliary proteins and ciliopathy genes are yet to be discovered. Indeed,

the genetic diagnosis of many ciliopathy disorders is still unknown (Shamseldin et al., 2020, 2020;

SYSCILIA Study Group et al., 2013; van Dam et al., 2019; Wheway et al., 2019).

Structural components of cilia and cilia cargos must be transported to cilia in order to construct

and sustain cilia, and mutations in genes involved in ciliary trafficking are commonly seen in

ciliopathies. Cilia has a one-of-a-kind protein delivery system called intraflagellar transport (IFT).

IFT is made up of multisubunit protein complexes that travel bidirectionally along the cilia. The

IFT complex contains two sub-complexes, IFT-A and IFT-B, consisting of 6 and 16 protein

subunits, respectively (Prevo et al., 2017). The IFT-B sub-complex and Kinesin-2 motors mediate

the motility of IFT and IFT cargos from the base of the cilia to the tip of the cilia (anterograde

IFT), while the IFT-A sub-complex and the cytoplasmic dynein-2 motor facilitate the retrograde

IFT transport (from the cilia tip to the base of the cilia) (Blacque, 2008; Rosenbaum and Witman,

2002). The IFT-A is involved in the transport of certain membrane proteins into cilia (Lee et al.,

2008; Liem et al., 2012; Mukhopadhyay et al., 2010). Mutations in genes encoding IFT

components lead to defects in cilia formation in all examined organisms, indicating the importance

of IFT for cilia assembly (Pazour et al., 2000; Prevo et al., 2017).

Bardet–Biedl syndrome (BBS) was classified as a ciliopathy in 2003, and eight of the highly

conserved Bardet-Biedl syndrome proteins (BBS1, BBS2, BBS4, BBS5, BBS7, BBS8, BBS9, and

BBIP10) establish a stable protein complex called the BBSome that undergoes IFT. Work from a

range of organisms implicates the BBSome in a variety of cilia related process including acting as

a cargo adaptor for removing proteins from cilia and as a regulator of the assembly and stability

of IFT trains (Ansley et al., 2003; Lechtreck et al., 2009; Loktev et al., 2008; Nachury et al., 2007;

Nozaki et al., 2019; Ou et al., 2007, 2005; Wei et al., 2012; Williams et al., 2014; Xu et al., 2015;

Ye et al., 2018). For example, in the nematode *C. elegans* mutants lacking *bbs-7* or *bbs-8*,

detachment of IFT-A and IFT-B in the amphid cilia was reported in the anterograde direction (Ou

et al., 2005). Analysis with a hypomorphic mutant (*bbs-1*) revealed that BBSome is involved in

attaching IFT-B components to the retrograde IFT machinery at the ciliary tips in the channel cilia

of *C. elegans* (Wei et al., 2012). Though it is known that the BBSome governs the IFT assembly

and returns of IFT from the cilia, we do not yet know which additional regulators control IFT

assembly at the ciliary base and IFT turnaround at the ciliary tip.

To identify new IFT regulators, we focused on our single-cell RNA-seq data that compared

expression profiles of ciliated cells with those of non-ciliated cells in the nematode *C. elegans,* and

this work revealed novel cilia genes and potential IFT regulators, including *WDR31* (manuscript

in preparation). Our gene discovery approach in combination with the use of *in vivo* IFT

microscopy analysis identified *WDR31* (WD repeat domain 31) and two GTPase activating

proteins (GAPs) as IFT regulators that facilitate the ciliary entry of IFT complex. Our work from

Zebrafish and *C. elegans* provides significant insight into the function of WDR31 in cilia

biogenesis. First, our work from *C. elegans* revealed that WDR-31 functions redundantly with two

GTPase activating proteins (GAPs) ELMD-1 (the sole ortholog of the human ELMOD proteins)

and RPI-2 (human retinitis pigmentosa 2 orthologue) to control the cilia morphology. Second,

knocking out *wdr-31* along with *elmd-1* or *elmd-1;rpi-2* causes IFT trafficking to be disrupted,

resulting in ciliary tip accumulations of IFT-B components and the OSM-3/KIF17 motor, as well

as significantly reduced BBSome recruitment to cilia. Third, fewer IFT/BBSome particles travel

along cilia in both anterograde and retrograde directions in *wdr-31;rpi-2;elmd-1* triple mutants,

indicating that the IFT/BBSome entry into and exit from cilia is affected. Third, anterograde IFT

in the middle segment moves faster in *wdr-31;rpi-2;elmd-1*. Finally, TRAM-1, a non-ciliary

membrane protein, penetrates *wdr-31;rpi-2;elmd-1* cilia. Taken together, this work identifies

WDR31 and two GAP proteins (ELMOD and RP2) as regulators for IFT and BBSome trafficking.

Material and Methods

C. elegans **strains, maintenance and genetic crossing**

For strain maintenance and genetic crosses, a standard procedure was followed, as described by

Sidney Brenner in 1974 (Brenner, 1974). After genetic cross with a marker to generate single,

double and triple mutants, we used the PCR strategy to trace the mutations in following mutants:

wdr-31(T05A8.5)(tm10423)II.; wdr-31(tur003)II.; wdr-31(syb1568)II.; nphp-4(tm925)V.; mks-

6(gk674)I.; elmd-1(syb630) III.; RB1550 rpi-2(K08D12.2)(ok1863) IV.;   and   *bbs-8(nx77) V.*

Primers can be found in Table S1.

Lipophilic fluorescent dye-uptake assay and rescue analysis

Healthy mixed-staged animals were collected using M9 buffer (3 g/L KH2PO4, 6 g/L Na2HPO4,

5 g/L NaCl, 1 mM MgSO4), centrifuged for 1 minute at 2000 rpm, and washed twice with M9

buffer to remove any bacterial contaminations. Worms were incubated for 45-60 minutes at room

temperature in an M9 buffer containing lipophilic dye (1:200 dilution in M9, Invitrogen™

Vybrant™ DiI Cell-Labeling Solution) (Herman and Hedgecock, 1990). The worms were then

washed twice with M9 before being moved to a new NGM plate. Wild type was always included

in Dye filling assay, and the dye uptake control for the wild type was performed under a stereotype

fluorescence microscope, followed by imaging with the fluorescence upright microscope. For the

rescue experiment, *N2;turEx24[arl-13p::GFP::elmd-1 (C56G7.3)::unc-54 3'UTR +rol-6}*

(OIK1045) were crossed into *wdr-31(T05A8.5)(tm10423)II., elmd-1(syb630) II,* double and *wdr-*

31(T05A8.5)(tm10423)II., elmd-1(syb630) II, rpi-2(K08D12.2)(ok1863) IV. triple mutants.

N2;turEx21[arl-13p::wdr-31 (T05A8.5)::GFP::unc-54 3'UTR +rol-6} (1 ng) (OIK1042) were

mated with *T05A8.5(syb1568)II., elmd-1(syb630) II, rpi-2(K08D12.2)(ok1863) IV.* triple mutants.

Plasmid (*arl-13p::wdr-31 (T05A8.5)::GFP::unc-54 3'UTR*) was directly microinjected into triple

mutant *T05A8.5(syb1568)II., elmd-1(syb630) II, rpi-2(K08D12.2)(ok1863) IV* (1 ng). *dpy-*

5(e907);nxEx386[rpi-2::gfp + dpy-5(+)] (MX352) were crossed into *wdr-*

31(T05A8.5)(tm10423)II., elmd-1(syb630) II, rpi-2(K08D12.2)(ok1863) IV. triple mutants. Three

independent Dye uptake assays were performed, and fluorescence filters were set for GFP and

Texas Red, followed by fluorescence imaging (20-150 heads and tails were counted). Dye uptake

of mutants with corresponding transgenics strains was compared to that of non-transgenic strains

in the same rescue plates **(Fig. S2C, D)**.

Generation of mutants using CRISPR/Cas9 in the nematode *Caenorhabditis elegans*

To generate *wdr-31(T05A8.5)(tur003)* allele, three sgRNAs targeting *C. elegans* T05A8.5 (human

WDR-31) were chosen using an online tool, Benchling [Biology Software] (2019), followed by

ordering complementary oligonucleotides (Macrogen, South Korea), and cloning of sgRNAs into

an empty sgRNA vector pRB1017. The successful sgRNA insert was confirmed with colony PCR,

followed by plasmid isolation. Three sgRNAs (each 50 ng/µl) were injected into the gonads of

wild type together with pDD162 (*Peft-3::Cas9;* 15 ng/ µl plasmid pRF4) and pRF4 (50 ng/ µl

plasmid pRF4) (Dickinson et al., 2013). F1s with the roller phenotype were identified, and after

they generated enough progenies, the PCR technique was used to identify the F1 generation

bearing the predicted deletion, as well as the homozygosity of the allele mutation. The PCR

products from knockout animals were then sent to the Sanger sequencing (Macrogen, South

Korea). *wdr-31(*tur003*)II.* mutants have 1276-bp deletion covering a huge part of exon II (297 bp

out of 359 bp) and whole exon III, IV, and exon V. This is likely a null allele of *wdr-31.* sgRNA

sequences can be found in Table S1.

Generation of transgenic strains and strain list for *C. elegans*

To generate transgenic lines for localization, rescue experiments, and expression patterns, we

generated the following transgenic animals via microinjections.

OIK1042   turEx21[arl-13p::wdr-31   (T05A8.5)::GFP::unc-54   3'UTR   +rol-6};

T05A8.5(syb1568)II., elmd-1(syb630) II, rpi-2(K08D12.2)(ok1863) IV. (1 ng)

OIK1044 N2;turEx23[elmd-1p::GFP::elmd-1 (C56G7.3)::unc-54 3'UTR +rol-6} (5 ng)

OIK1045 N2;turEx24[arl-13p::GFP::elmd-1 (C56G7.3)::unc-54 3'UTR +rol-6} (5 ng)

OIK1046 N2;turEx25[elmd-1p(C56G7.3)::GFP::unc-54 3'UTR +rol-6} (50 ng)

OIK1047 N2;turEx26[wdr-31 (T05A8.5)p::GFP::unc-54 3'UTR +rol-6} (50 ng)

The rol-6 plasmid (50 ng/ µl plasmid pRF4) was co-injected as the co-transformation marker. In

brief, the plasmids were delivered by microinjections into the gonads of 1-day adult worms.

Worms were initially transferred onto a 2.5% agarose pad prior (Halocarbon oil, Sigma: 9002-83-

9), followed by microinjection. The microinjection was done using a Zeiss Axio Vert.A1 inverted

microscope with DIC optics coupled with a Narishige Micromanipulator MMO-4. We next

manually inspected the plates to find successful transgenics animals.

Wild Type and mutant alleles: N2; *FX30333; wdr-31(T05A8.5)(tm10423)II.; OIK393*

T05A8.5(tur003)II.; PHX1568 T05A8.5(syb1568)II.; nphp-4(tm925)V.; mks-6(gk674)I.;

PHX630, elmd-1(syb630) III.; RB1550 rpi-2(K08D12.2)(ok1863) IV.; MX52, bbs-8(nx77) V. We

obtained *wdr-31(T05A8.5)(tm10423)II. (FX30333)* mutant allele, which has a 160-bp deletion,

causing a frameshift, from the National Bioresource Project, Japan. The *wdr-*

31(T05A8.5)(tm10423)II. were outcrossed to wild type four times. The Caenorhabditis Genetics

Center (CGC), Minnesota, USA, provided the *RB1550 rpi-2(K08D12.2)(ok1863)* mutant, and the

rpi-2(ok1863) allele contains 1143-bp deletion that removes a large segment of exon III, exon IV,

and some portion of exon V. *T05A8.5(syb1568)II.* has 1888-bp deletion, deleting all exons except

exon I **(Supplementary Fig. 3B)**. Sunybiotech created an independent null allele of *elmd-1* via

CRISPR–Cas9. The *PHX630, elmd-1(syb630) III.* mutant contains 1784-bp deletion, where except

for exon I, all exons were removed **(Supplementary Fig. 2B)**. *elmd-1(syb630) III.* were outcrossed

to wild type two times.

Fluorescent transgenes for IFT proteins: *GOU2162   che-3(cas443[gfp::che-3]) I; xbx-*

1(cas502[xbx-1::tagRFP]) V ; GOU2362 ift-74(cas499[ift-74::gfp]) II.; EJP76   vuaSi15

[pBP36; Posm-6::osm-6::eGFP; cb-unc119(+)] I; unc-119(ed3) III; osm-6(p811) V;

N2;lqIs2[osm-6::gfp], N2;ejEx[osm-3::GFP + pRF4]; N2;ejEx[kap-1::gfp+pRF4]; EJP81

vuaSi24 [pBP43; Pift-140::ift-140::mCherry; cb-unc-119(+)]II; unc-119(ed3) III; ift-

140(tm3433) V; jhuEx [ift-140::GFP+pRF4]; Ex[rpi-2::GFP+ xbx-1::tdTomato+pRF4]; MX76

dpy-5(e907); nxEx(bbs-7::gfp+dpy-5 (+)).

Fluorescent transgenes for ciliary proteins: PHX1180, *wdr-31(syb1180 [wdr-*

31(T05A8.5)::GFP]); PHX4934 rpi-2(syb4934) [rpi-2::mCherry]; OIK1042 N2;turEx21[arl-

13p::wdr-31 (T05A8.5)::GFP::unc-54 3'UTR +rol-6} (5 ng); OIK1044   N2;turEx23[elmd-

1p::GFP::elmd-1 (C56G7.3)::unc-54 3'UTR +rol-6} (5 ng); OIK1045   N2;turEx24[arl-

13p::GFP::elmd-1 (C56G7.3)::unc-54 3'UTR +rol-6} (5 ng); OIK1046   N2;turEx25[elmd-

1p(C56G7.3)::GFP::unc-54   3'UTR   +rol-6}   (50   ng);   OIK1047   N2;turEx26[wdr-31

(T05A8.5)p::GFP::unc-54 3'UTR +rol-6} (50 ng); *N2;Ex[mks-2::GFP + tram-1::tdTOMATO +*

pRF4];   MX1409   N2;   nxEx785[tax-4:gfp+   Posm-5::xbx-1::tdTomato   +   rol-6(su1006)];

vuaSi21[pBP39;   Pmks-6::mks-6::mCherry;   cb-unc-119(+)]II;   MX2418   N2;nxEx1259[pbbs-

8::PLC-delta   PH::GFP;MKSR-2::tdTomato;   coel::GFP];   PY8847   oyIs65[str-1p::mcherry];

Ex[str-1p::nphp-4::gfp, unc122p::dsRed]. All PHX strains were generated using CRISPR/Cas9

by Sunybiotech. The list of extensive transgenic and mutant strains was provided in Table S2.

Fluorescence and Confocal microscopy for *C. elegans*

Fluorescence images (dye assay) and time-lapse movies (IFT movies) were captured using an epi-

fluorescence upright microscope (Leica DM6 B) (3 frames per second, and up to 120 frames). The

epi-fluorescence upright microscope was controlled with iQ3.6.2 Andor software and was attached

with an Andor iXon Ultra 897 EMCCD (an electron-multiplying charge-coupled device camera).

The LSM900 confocal microscope with Airyscan 2 (ZEN 3 Blue edition software) was used to

capture the high-resolution Z-stack images. On microscope slides, a drop of 2-3% agarose was

used to create an agarose pad, and worms were then moved to the agarose pad containing 1-3 μl

of 10 mM levamisole (an anesthetic agent). Images were collected at 0.14 m intervals with a Plan

ApoChromat 63x/1.40 NA target, then analyzed with Blue edition software ZEN 3 to create Z-

stacks, and processed with ImageJ (NIH) software (Schneider et al., 2012).

***In vivo* intraflagellar transport assay for IFT frequency and velocity**

The time-lapse movies of GFP-labelled IFT proteins were analyzed with the automatic kymograph

analyzing tools KymographClear and KymographDirect, both of which are ImageJ based

(Mangeol et al., 2016; Turan et al., 2022). KymographClear was used to produce kymographs from

time-lapse movies. We examined each produced kymograph and calculated IFT frequency or IFT

velocities for IFT-74::GFP (an endogenously tagged), GFP::CHE-3 (an endogenously tagged),

BBS-7::GFP (an overexpression transgene), OSM-6::GFP (a single copy transgene), OSM-3::GFP

(an overexpression transgene), and IFT-140::GFP (an overexpression transgen) in wild type, *wdr-*

31;elmd-1 double and *wdr-31;elmd-1;rpi-2* triple mutants. All worms were maintained at 20 °C

for IFT analysis. At least ten IFT videos were taken over three separate time periods for strains,

and IFT particles from at least ten kymographs were counted.

Whole-mount *in situ* hybridization in zebrafish

Whole-mount in situ hybridization was performed as previously described (Thisse et al., 2004).

The *wdr31* cDNA with T7 promoter was PCR amplified, with forward primer:

ATGGGGAAGCTACAGAGCAAGTTC and reverse primer:

TAATACGACTCACTATAGAAGCGAGCCACTTCAGTGATACTG, from a homemade

cDNA library of zebrafish embryos at 24 hpf. The antisense probe for *wdr31* was then transcribed

with digoxigenin-labeled UTPs and T7 RNA polymerases (Roche, Basel, Switzerland). The

stained embryos were dehydrated in glycerol and photographed with a Nikon SMZ1500

stereomicroscope (Nikon, Tokyo, Japan).

Mutagenesis of *wdr31* in zebrafish using the CRISPR/Cas9 technology

Generation of zebrafish mutants using the CRISPR/Cas9 system was carried out as previously

described (Chang et al., 2013). Briefly, two gRNAs targeting sequences in *wdr31* were chosen as

following: 5'- ACCCATGTGTGTTGGGTACC-3' and 5'-GAAGCCATCCAGGAGTTCAG-3'.

gRNA templates were PCR amplified and gRNAs were *in vitro* transcribed with T7 transcriptase

(NEB, cat# M0251S). gRNAs and Cas9 protein (NEB, cat# M0251S) were simultaneously injected

into the embryos at one-cell stage.

Immunofluorescence staining and microscopy

Immunostaining was performed as previously described (Xu et al., 2017). Briefly, embryos were

fixed in cold Dent's fixative (80% methanol: 20% dimethyl sulfoxide) at -20°C overnight and then

stored in methanol. Samples were permeabilized with 0.005% (m/v) trypsin for 30 minutes.

Samples were blocked with blocking buffer (10% [v/v] goat serum in PBST), followed by

incubation with primary antibody anti–acetylated tubulin (1:2000; Sigma-Aldrich, St. Louis, MO)

and then secondary antibodies conjugated with Alexa Fluor 568 (1:500; Invitrogen, Carlsbad, CA).

Samples were mounted on ProLong Gold Antifade Mountant with DAPI (Invitrogen) and images

were taken with a Nikon A1R confocal microscope.

Software and algorithms

Zen Blue   Zeiss   https://www.zeiss.com/corporate/int/home.html

Andor iQ3   Andor   https://andor.oxinst.com/

Fiji   ImageJ   https://fiji.sc

ImageJ   ImageJ   https://imagej.nih.gov/ij/

Illustrator (CS5.1) Adobe, USA   https://www.adobe.com/

KymographClear   Peterman Lab https://sites.google.com/site/kymographanalysis

| 277 | R | R Core Team | https://www.r-project.org/ |

Statistical analysis and Generating Figures

For statistical analysis of dye assay, Fisher's exact test (one tailed tests) was performed. For data involving continuous variables like IFT speed and cilia length, Mann-Whitney U-test or Welch's t-test was used depending on the distribution of data. Transgenic worms were kept at 15 °C for AWB cilia morphology analysis and AWB cilia length measurements, and only L4 stage animals were imaged for AWB cilia. For each strain, at least three independent microcopy analyses were performed, and 60-220 AWB cilia were examined. All statistical tests were performed using R software. The codes and files needed to generate figures and perform statistical analysis were openly shared, and the files and codes for making figures and performing statistical analysis may be accessed at https://github.com/thekaplanlab/WDR31-ELMOD-RP2.

Gateway Cloning (mammalian constructs)

Constructs for ELMOD3 (HsCD00288286) and WDR31 (HsCD00045652) were purchased from Harvard Medical School. LR Reaction (Invitrogen, USA) was performed to transfer the target sequence into destination vectors ((N)RFP, (N)CFP plasmids) with a subsequent transformation into *E. coli* DH5α. DNA was isolated according to Monarch® Plasmid Miniprep Kit (BioLabs, USA) and the PureYield® Plasmid Midiprep Protocol System (Promega, USA). Verification of successful cloning was done by sequencing (Eurofins, Luxembourg).

Immunofluorescence Staining (Mammals)

hTERT-RPE1 cells were transiently transfected according to Lipofectamine™ 3000 Reagent Protocol with ELMOD3-CFP and WDR31-RFP constructs followed by serum starvation for three

days to induce cilia formation. Cells were fixed with 4% PFA for 45min at 4°C, permeabilized for

5min with 0.3% PBST and blocked with 10% goat serum in PBST at 4°C overnight. A primary

antibody for ARL13B (1:50; Proteintech, Germany) and a secondary antibody conjugated to Alexa

Fluor 647 (1:350; Invitrogen, USA) were used. Cells were mounted using Fluoromount-G

(Invitrogen, USA).

Microscopy Setup (Mammals)

Images were captured using a Leica TCS SP8 scanning microscope (Leica Microsystems IR

GmbH, Germany). The setup includes 488 nm, 532 nm and 635 nm pulsed excitation lasers as

well as 100x oil immersion objective lens (NA 1.4) and a hybrid detector (HyD). Pixel number

was 1024 x 1024 and optimal pixel size was determined by Nyquist calculation resulting in a size

of 43×43 nm in XY. Additionally, Z-steps should not exceed 131 nm when acquiring a stack.

Laser intensity was adjusted for each sample and images were recorded with 2x frame averaging.

Image Processing (Mammals)

Acquired images were processed (LasX, Leica Microsystems), deconvoluted (Huygens Software,

SVI, Netherlands) and edited (FIJI software). The deconvolution was performed with a Classical

Maximum Likelihood Estimation (CMLE) algorithm under experimentally defined settings. The

background level was software estimated, the quality threshold was 0.001, the number of iterations

was 50 and signal-to-noise ratio was set to 20. All images were brightness-corrected for the

purpose of presentation.

Results

WDR31 and ELMOD are evolutionarily conserved constituents of cilia

As part of our ongoing effort to reveal novel cilia genes, using single-cell RNA-seq data, we

conducted comparative expression analysis in *C. elegans* and discovered that the WDR31 and

ELMOD orthologues, WDR-31 (T05A8.5) and ELMD-1 (C56G7.3), are likely expressed

exclusively in ciliated sensory neurons (manuscript in preparation). The head (amphid) and tails

(phasmid) of *C. elegans* contain a total of 60 ciliated sensory neurons, and the expression patterns

of genes can be reliably analyzed using fluorescence tagged promoter markers. Consistent with

our predictions, promoter-based GFP reporters revealed that both WDR-31 and ELMD-1 are

expressed in most of the ciliated sensory neurons **(Fig. S1A)**. To explore if both proteins are

concentrated within cilia, we investigated their subcellular localization in *C. elegans* and created

CRISPR/Cas9-mediated knock-in of WDR-31::GFP and a transgenic animal expressing GFP-

tagged ELMD-1. The endogenously expressed WDR-31::GFP is concentrated at the ciliary base

in both head and tails, where it colocalizes with the basal body marker γ-tubulin (TBG-1, the

ortholog of human TUBG1), and the IFT-140 (human IFT140) basal body signal **(Fig. 1A, B, C,**

and D), so our co-localization data suggests that WDR-31 is a new cilia-associated protein.

ELMD-1 is the sole orthologue of the three human ELMOD proteins (ELMOD1-3) that function

as GTPase-activating proteins (GAP). We found that GFP::ELMD-1 localizes to the periciliary

membrane compartment (PCMC) and the basal body (BB) (marked with IFT-140), but is proximal

to the MKS-6-labeled transition zone that is adjacent to the BB **(Fig. 1E, F, and G)** (Kaplan et al.,

2012).

To determine if both human WDR31 and ELMOD3 (also known RBED1, the top blastp hit for

elmd-1) are concentrated within cilia in mammalian cell lines, we generated hTERT-RPE1 cells

co-expressing WDR31-RFP and ELMOD3-CFP and stained them with a ciliary marker ARL13B.

Furthermore, WDR31-RFP or ELMOD3-CFP were transiently transfected into hTERT-RPE1

cells, and they were additionally stained for ARL13B. Our super-resolution microscopy analysis

revealed that both WDR31 and ELMOD3 are enriched in the cilium **(Fig. 2A, B and Fig. S2B**

and C). Taken together, our complementary approach demonstrates that both WDR31 and

ELMOD3 are evolutionarily conserved proteins associated with cilia.

Wdr31 regulates ciliogenesis in the Zebrafish ear

We turned our interest into zebrafish to develop an *in vivo* model for WDR31 to examine the role

of WDR31. We first used the whole-mount *in situ* hybridization (ISH) within zebrafish to visualize

the expression pattern of *wdr31* during zebrafish embryo development up to 1-day post-

fertilization. The dynamic and wide expression of *wdr31* during zebrafish embryonic development

(2 cell and 8 somite stages (SS)) has become restricted to the otic vesicle and brain region at 24

hours post-fertilization (24 hpf) **(Fig. 3A)**. We next knocked out (KO) *wdr31* in zebrafish with the

CRISPR/Cas9 system and the embryos showed heart edema and otolith malformation. As *wdr31*

is expressed in the otic vesicle and the otolith development requires cilia, we checked the cilia in

the otic vesicle. The cilia bundle in the lateral crista (LC) of the otic vesicle was stained with

acetylated tubulin in wild type and *wdr31* KO, and although we found that the length of LC cilia

was comparable with wild type, the width of the cilia bundle was reduced by 33% in LC of the

otic vesicle compared to that in the control embryos, suggesting the cilia number is decreased **(Fig.**

3B, C, and D). Our zebrafish work indicates that WDR31 plays a role in ciliogenesis in the

zebrafish ear.

Functional redundancy of WDR31, ELMOD, and RP2 for determining cilia morphology

We next wanted to gain mechanistic insight into the functions of *wdr-31* in cilia biogenesis. To

this end, we employed the nematode *C. elegans* and obtained/generated three *wdr-31* alleles:

tm10423 (160-bp frameshift causing deletion), syb1568 (1888-bp deletion removing all exons

except exon I), and tur003 (1276-bp deletion removing a large portion of exon II and exon III, IV,

and exon V) **(Fig. S3B)**. In *C. elegans,* the lipophilic fluorescent dye-uptake assay is employed to

indirectly evaluate the cilia structure, and our analysis revealed *wdr-31* mutants display wild type

level dye-uptake in both head (amphid) and tail (phasmid), suggesting cilia structures are likely

unaffected in these mutants **(Fig. 4B and Fig. S4A)**. To address whether loss of *wdr-31* leads to

subtle defects in cilia morphology, we expressed the fluorescence-based marker *str-1pro:mCherry,*

gcy-5pro::gfp, srb-6pro::gfp , which label the AWB, ASER and PHA/PHB cilia, respectively, in

wdr-31 mutants. The AWB dendritic tip extends the fork-like cilia in the wild type, and rod-like

cilia protrude from the dendritic endings of ASER and PHA/PHB sensory neurons. Our confocal

microscopy analysis revealed that the structure and length of ASER, PHA/PHB, and AWB cilia

are comparable with those of wild type cilia, suggesting, contrary to the importance of WDR31

for ciliogenesis in the zebrafish ear, the loss of *wdr-31* alone does not result in a severe defect in

cilia structure **(Fig. 4C, D, E, F, and G)**.

The lack of an apparent cilia phenotype in *C. elegans* may be attributed to a functional

redundancy for WDR-31 in cilia biogenesis, and functional redundancy is indeed a common

phenomenon in ciliopathy-related genes. For example, while the absence of individual ciliopathy

genes encoding MKS/MKSR proteins or NPHP proteins (NPHP-1, NPHP-4) does not result in

serious cilia defect, the loss of *nphp-4* (the ortholog of human NPHP4) in combination with *mks-*

6 (the ortholog of human CC2D2A) causes more severe cilia related defects (transition zone

membrane association defects) (Williams et al., 2011). We next explored genetic interaction

between *wdr-31* and *elmd-1*, since our results showed that both protein products are at the ciliary

base **(Fig. 1C, D, E, and F)**. We discovered that the *wdr-31(tm10423);elmd-1(syb603)* double

mutant (hereinafter referred to as "double mutant") has a partial Dyf defect in both head and tail

neurons, which was rescued by expression of a transgene containing the wild type *elmd-1* sequence

(Fig. 5B and Fig. S4A, $p < 0.0001$, Fisher's exact test**)**. Using the AWB, ASER and PHA/PHB

cilia fluorescence markers, we subsequently set out to visualize the ciliary structures in the double

mutant and single mutants. The AWB cilia in the double mutants have an extra projection in the

middle part of the cilia (7% (N: 100) and 52 % (N: 134) in WT and double mutant, respectively),

whereas ASER, AWB, and PHA/PHB cilia are normal length in WT and double mutant **(Fig. 4C**

and D, E, and F and Fig. S4B).

We explored genetic interaction between *wdr-31, elmd-1,* and *rpi-2* (the X-linked retinitis

pigmentosa protein RP2 and a GTPase-activating protein) since ELMD-1 and WDR-31

localizations are reminiscent of RPI-2 localization (the endogenously labeled RPI-2) in *C. elegans*

(Fig. 4A, and Fig. S5A) (Williams et al., 2011). We, therefore, first generated an *elmd-1;rpi-2*

double mutant, which revealed no Dyf phenotype **(Fig. 4B, and Fig. S4A)**. However, when we

created *wdr-31;elmd-1;rpi-2* triple mutants (hereinafter referred to as "triple mutant"), we

observed severe synthetic Dyf phenotype, which is significantly rescued by the introduction of a

wild-type copy of *wdr-31* or *elmd-1 or rpi-2* **(Fig. 4B, Fig. S3C, and Fig. S4A** $p < 0.0001$, Fisher's

exact test**)**.

We then sought to examine AWB, ASER and PHA/PHB cilia morphology in triple mutants

and compared them with the *wdr-31;elmd-1* and *rpi-2;elmd-1* double mutants. The Dyf defect of

triple mutants was indeed accompanied by significant changes in these cilia. The AWB cilia

displayed the ectopic projections, including a backward projection from the base of AWB cilia in

triple mutant and as wells as ectopic projections from the middle part of AWB cilia (0% backward

projection in wild type vs over 30% backward projection in triple mutant; $p < 0.0001$, Fisher's

exact test) (**Fig. 4F and G**). It is noteworthy that the similar backward projections in AWB cilia

were observed in two independent triple mutants generated with *wdr-31(syb1568)* and *wdr-*

31(tur003) (**Fig. S7A and B**). A backward projection from the ciliary base was observed for ASER

cilia, but not PHA/PHB cilia in triple mutants (**Fig. 4F**). We next measured the cilia lengths of

these cilia and discovered that ASER and PHA/PHB cilia are significantly shorter in triple mutants

(20% shorter for ASER in triple mutants, 13% shorter for PHA/PHB in triple mutants), whereas

the cilia length of AWB cilia was not significantly altered in triple mutants (**Fig. 4C, D, F, and**

Fig. S4B). This establishes a role for WDR-31 in controlling cilia length and morphology in a

subset of sensory neurons in a redundant manner with RPI-2 and ELMD-1.

WDR-31-ELMD-1-RPI-2 are needed for effective cilia entry and exit of IFT components.

We next aimed to investigate the mechanism by which WDR-31, ELMD-1, and RPI-2 regulate

cilia morphology in *C. elegans*. To this end, we investigated whether these genes affect the

localization of IFT proteins and IFT motors, including kinesin motors (Kinesin II and OSM-

3/KIF17), dyneins, and IFT components (IFT-A and IFT-B). The IFT complex and motor proteins

are critical for cilia construction and maintenance because they deliver ciliary constituents from

the cell body to the cilia. In *C. elegans,* two kinesin motors Kinesin II and OSM-3/KIF17

(heterotrimeric kinesin-II and homodimeric OSM-3) work cooperatively to carry the IFT complex

in the middle segment of the channel cilia in an anterograde direction, while OSM-3/KIF17

transports the entire IFT complex (IFT-A, IFT-B, and BBSome) in the distal part of channel cilia.

We generated single, double, and triple mutants expressing fluorescence tagged IFT and

motor proteins. Using single-copy transgenes, we found that IFT-B components predominantly

accumulate at cilia tips and/or middle of cilia in double mutants (96% ciliary tip accumulations for

IFT-74/IFT74, N: 26 and 82% ciliary tip accumulations for OSM-6/IFT52, N: 46) but the

localization of GFP::CHE-3 (human dynein heavy chain DYNC2H1) remains less affected in

double mutants (27% minor middle cilia accumulations for CHE-3; N: 48, see supplementary

Movie S1) **(Fig. 5A, C and D and Supplementary Movie S1, 2, 4)**. Furthermore, we next

investigated the localization of GFP::CHE-3 in triple mutants because the cilia morphology of

triple mutants is more severe than that of *wdr-31;elmd-1* or *wdr-31;rpi-2* or *elmd-1;rpi-2* double

mutants **(Fig. 5A).** Compared with wild type and double mutants, we noticed additional IFT

abnormalities in two independent triple mutants, including dim cilia staining with GFP::CHE-3

and accumulations of GFP::CHE-3 in the middle/distal part of cilia (81% middle cilia

accumulations, N: 96; 6.6% ciliary tip accumulations, N: 90) **(Fig. 5A and Supplementary Movie**

S1). Motor protein OSM-3/KIF17 accumulation within the ciliary tips was detected in double and

triple mutants **(Fig. 5B, and Supplementary Movie S3)**. Furthermore, in triple mutants, the dim

distal cilia staining was observed for IFT-140::GFP (IFT-A component) (63% dim distal cilia, N:

47) and XBX-1::mCherry (a dynein light intermediate chain) **(Fig. 5E and F and Supplementary**

Movie S5), but the ciliary distribution of KAP-1::GFP (Kinesin II) remains unaffected **(Fig. 5G)**.

Cilia accumulations of IFT-B components and OSM-3/KIF17 coupled with weak cilia

staining with IFT-A components in triple mutants forced us to better understand the role of WDR-

31-ELMD-1-RPI-2 in IFT defects, we therefore employed *in vivo* time-lapse video coupled with

kymography. For a subset of IFT components, we detected a significant decline in the quantity of

IFT particles traveling along the cilia in both directions in double and triple mutants **(Figure 6A,**

B, C, D, E, E, G, and H). We found significant decline in the IFT transport for kinesin motor

OSM-3::GFP in double and triple mutants in both anterograde and retrograde directions (the

anterograde and retrograde: 0.68 n/s and 0.61 n/s in wild type; 0.43 n/s and 0.32 n/s in double

mutants; 0.34 n/s and 0.23 n/s in triple mutants*,* $p < 0.0001$; the Mann–Whitney U test) **(Fig. 6A,**

and B), while our analysis revealed that the flux of cytoplasmic dynein motor protein CHE-3 along

the cilium was unchanged in both directions in both double and triple mutants (**data not shown**),

suggesting, indicating that the simultaneous elimination of these genes has an effect on the kinesin

motor OSM-3 but not on cytoplasmic dynein loading onto IFT, but their absence leads to both

ciliary cytoplasmic dynein accumulation and the kinesin motor OSM-3 (**Fig. 5A and B**). In wild

type, the average of IFT-74 particles moving in anterograde and retrograde directions is 0.58 (n/s:

number of particles/seconds) and 0.52 n/s, respectively, while in triple mutants, the anterograde

and retrograde IFT-74 particles are 0.27 n/s and 0.09 n/s. Comparable reductions were observed

for IFT-74::GFP (CRISPR tagged endogenous IFT-74) in *wdr-31;eldm-1* double mutant (0.36 n/s

and 0.21 n/s in *wdr-31;eldm-1;* $p < 0.0001$; the Mann–Whitney U test) (**Fig. 6G, and H**). For

OSM-6::GFP, the anterograde and retrograde IFT particles statistically differ between wild type

and triple mutants (the anterograde and retrograde: 0.52 n/s and 0.58 n/s in wild type, 0.36 and

0.30 in triple mutants*;* $p < 0.0001$; the Mann–Whitney U test) (**Fig. 6E, and F**). Furthermore, IFT-

A component IFT-140 transport declined in both directions both double and triple mutants (**Fig.**

6C, and D). Taken together, our fluorescent microscopy analysis reveals several IFT abnormalities

in triple mutants: first, in triple mutants, there is a significant decrease in the quantity of IFT

particles, including the OSM-3/KIF17 kinesin motor and both IFT-A and IFT-B components,

traveling in both anterograde and retrograde directions. Second, the ciliary accumulations of IFT-

B components and OSM-3/KIF17 kinesin motors in cilia together with weak distal cilia staining

of the IFT-A component and dynein (XBX-1) in triple mutants suggest that the defects in the return

of IFT from the ciliary tips. These defects might stem from a reduction in the loading of IFT

components and dynein onto IFT trains at the ciliary base.

To better investigate the role of WDR-31, ELMD-1, and RPI-2 in IFT, we measured the

anterograde (middle and distal segment) and retrograde IFT velocities in wild type and triple

mutants. When compared to wild-type IFT speeds, we found a significant increase in the average

anterograde IFT velocities of IFT-74::GFP, IFT-140::GFP, OSM-3::GFP, and OSM-6::GFP in the

middle segments of triple mutant (**Fig. S7A, B, C, and D**), implying that the integrity of

anterograde IFT components is likely compromised. In contrast, in the triple mutant, the retrograde

and distal anterograde IFT velocities are unaltered. Taken together, our findings suggest that with

the exception of heterotrimeric Kinesin II, all IFT proteins studied, including dynein CHE-3,

OSM-3 kinesin, IFT-B, and IFT-A components, display certain defects in *wdr-31* related mutants,

either ciliary accumulations for dynein CHE-3, OSM-3 and IFT-B component or dim distal cilia

staining for IFT-A component and dynein motors (CHE-3 and XBX-1) or changes in IFT

velocities.

WDR-31, ELMD-1, and RPI-2 restrict ciliary entry of non-ciliary proteins

Given that WDR-31, ELMD-1, and RPI-2 localize at the base of cilia, we wanted to examine the

role of these proteins in the ciliary gate, and we chose the TRAM-1 protein (the ortholog of human

TRAM1), which surrounds the PCMC in *C. elegans* but stays outside of cilia (Williams et al.,

2011), as well as the transition fiber protein DYF-19 (human FBF1) and the transition zone

proteins NPHP-1 (the human NPHP1) and MKS-2 (human TMEM216) (Wei et al., 2013). We

showed that neither *wdr-31, elmd-1*, or *rpi-2* deletion, nor double mutant combinations, result in

ciliary entry of TRAM-1 or MKS-2 (**Fig. 7A and B**), suggesting that neither of these alone affects

ciliary gating. In contrast, TRAM-1 protein leaks into cilia in all three independent *wdr-31* triple

mutants, but the majority of signal remains outside of cilia (**Fig. 7A and B**), whereas the

localizations of transition zone NPHP-1 and transition fiber protein DYF-19 remain unchanged

(**Fig. 7C**). In the *wdr-31;elmd-1;rpi-2* triple mutants, the PLC1-PH::GFP marker for

phosphatidylinositol 4,5-bisphosphate (PtdIns(4,5)P2) stays outside of cilia. In the wild type,

PLC1-PH::GFP stains PCMC membranes but does not penetrate cilia (**Fig. 7D**). Furthermore, we

explored the localization of TAX-4, a ciliary membrane protein, and it remains unaffected in the

triple mutants (**Fig. 6SB**). Taken together, our findings reveal a functionally redundant role for

WDR31, ELMD-1, and RPI-2 in restricting entry of non-ciliary proteins into cilia, despite the fact

that TZ protein localization remains unchanged.

WDR-31 and ELMD-1 are required for recruitment of BBS-7, a BBSome component, to cilia

Some of IFT defects (cilia accumulations of IFT-B complex, weak cilia staining of IFT-A

components, and increased in IFT speed) observed in the *wdr-31;elmd-1;rpi-2* triple mutants are

reminiscent of IFT defects observed in several *bbs* mutants (Blacque, 2004; Ou et al., 2005; Wei

et al., 2012; Xu et al., 2015;). We hypothesized that the IFT complex destabilization defects in

triple mutants might be due to defects in the BBSome. First, we found that ciliary tip accumulations

of IFT-74:GFP, an IFT-B component, in *bbs-8(nx77)* mutants were comparably similar to that of

double and triple mutants (**Fig. 9A**). Furthermore, there is no further increase of IFT

accumulations in the triple mutants *wdr-31;elmd-1;bbs-8*.

We then investigated whether the ciliary localization of BBS-7::GFP, a core member of

the BBSome, is controlled by WDR-31, ELMD-1, and RPI-2, because the ciliary localization of

other BBSome components is dependent on other BBSome subunits (Ou et al., 2007). We predict

that the ciliary localization of BBSome is more likely disturbed in triple mutants, contributing to

the observed IFT abnormalities. Consistent with expectations, BBS-7::GFP is lost or significantly

diminished in the cilia of double and triple mutants, but BBS-7::GFP localization persists at the

ciliary base in these mutants (**Fig. 9B**). We next performed the time-lapse movie analysis of BBS-

7::GFP with kymography, which revealed that BBS-7::GFP movements were undetectable in 27%

and 42% of cilia in the head in double and triple mutants, respectively (**Supplementary Movie**

S6). We were able to quantify the frequency of BBS-7::GFP even though the density of trajectories

(IFT particles) on kymographs was weak in double and triple mutants relative to wild type (**Fig.**

9C). Our analysis revealed a substantial decrease in the average of BBS-7::GFP particles

translocating in the anterograde and retrograde directions in the remaining worms that display IFT

(average anterograde and retrograde IFT particles: 0.55 n/s and 0.58 for wild type; 0.30 n/s and

0.33 n/s for double mutant; 0.07 n/s and 0.07 n/s for triple mutant, $p < 0.0001$; the Mann–Whitney

U test) (**Fig. 9C, D and Supplementary Movie S6**). We next used IFT-74::GFP to investigate IFT

speeds in *bbs-8* and its mutant combinations with *wdr-31* and *elmd-1*. The average velocities of

IFT-74::GFP in the middle segment of *bbs-8;wdr-31* and *bbs-8;wdr-31;elmd-1* were higher than

wild type, but were comparable to the *wdr-31;elmd-1;rpi-2* triple mutant (**Fig. S7A**). Taken

together, our results suggest that because the BBSome complex is unable to gain access to cilia in

double and triple mutants, IFT-B components and OSM-3/KIF17 likely accumulate excessively in

the distal portion of the cilia while IFT Complex A component staining the distal segment cilia

become dim (**Fig. 9E**).

Discussion

WDR31 is a new ciliary protein encoding gene required for cilia biogenesis

Clinical genomics, proteomics, functional genomics, and bioinformatics research have all

contributed to the expansion of the list of both ciliary and ciliopathy genes. Over 300 genes have

been identified as encoding ciliary proteins, including IFT-kinesin-dynein components, BSSome

components, structural ciliary components, signaling molecules, and IFT regulators, with many of

them being ciliopathy genes (Wheway et al., 2019; Vasquez et al., 2021). Our findings reveal the

roles of WDR-31, a member of the WD40-repeat protein (WDR) family, the ELMOD orthologue

(ELMD-1) and RP2 orthologue (retinitis pigmentosa 2; RPI-2) in cilia and IFT regulation. Several

members of the WD40-repeat protein family, including WDR34, WDR35, and WDR60, have

previously been connected to cilia, but the role of *WDR-31* in cilia has been unknown (Blacque et

al., 2006; Patel-King et al., 2013; Rompolas et al., 2007). Strikingly, we found that WDR31 is a

ciliary protein, and knocking it out in zebrafish using CRISPR/Cas9 causes ciliary defects,

including a decrease in the number of cilia in the LC. Furthermore, the link between WDR31 and

cilia was significantly strengthened by findings from *C. elegans*. First, the expression of the

WDR31 orthologue (WDR-31) is restricted to the ciliated sensory neurons, and WDR-31 is

localized to ciliary compartments (the ciliary base and cilia) in C. elegans and human TERT-RPE1

cells. The exclusive cilia expression pattern of *wdr-31* in *C. elegans* is similar to that of ciliary and

ciliopathy genes like ARL-13/ALR13B, IFT, and BBS (Blacque et al., 2005). Second,

CRISPR/Cas9 mediated knock out of *WDR-31* combination with the ELMOD orthologue (ELMD-

1) and RP2 orthologue (retinitis pigmentosa 2; RPI-2) results in an abnormality in cilia functions

and structure in *C. elegans*. Our findings reveal that WDR31 is a new ciliary protein required for

cilia morphology in both zebrafish and *C. elegans*, suggesting the evolutionarily conserved role of

WDR31 in cilia biology.

Functional redundancy between WDR-31, ELMD-1, and RPI-2

Our genetic analysis in *C. elegans* provides evidence for functional redundancy for *wdr-31*, *elmd-1*, and *rpi-2* in regulating cilia length, cilia morphology, and the trafficking of kinesin-IFT-BBSome complexes. Though cilia length and cilia morphology are disrupted in *wdr-31;elmd-1;rpi-2* triple mutants, with altered recruitment of kinesin, IFT, and BBS proteins and IFT dynamics, simultaneous elimination of *elmd-1* and *rpi-2* did not result in severe defects in cilia and IFT. Our interpretation for these data is that the additive defects in the triple mutants were likely due to *wdr-31*, and WDR-31 is a central player in controlling cilia morphology and IFT machinery. However, we cannot rule out the possibility of microtubule defects in cilia ultrastructure in *elmd-1;rpi-2* double mutants. Additional work is needed, and a transmission electron microscope (TEM) may be used to reveal further details in the ultrastructure of cilia in these double mutants.

How do these proteins work together to regulate cilia-related phenotypes? While WDR31 is a poorly characterized gene, the roles of human RP2 and ELMOD proteins in cilia biology are better understood. Human RP2 localizes to the cilium and the basal body and displays GAP activity toward two ARF family members, ARL2 and ARL3, both of which were linked to cilia biology (Evans et al., 2010; Schwarz et al., 2017; Wright et al., 2011), while the ELMOD protein family (ELMOD1-3) has GAP activity for ARL2 (ELMOD1 and ELMOD3) and ARF6 (ELMOD1) (Ivanova et al., 2014; Jaworek et al., 2013; Johnson et al., 2012; Miryounesi et al., 2019). ELMOD2 localizes to the basal body, and our study showed ELMOD3 localizes to cilium (Turn et al., 2021). Taking into account the fact that RPI-2 and ELMD-1 are GAPs, and loss of these two GAP proteins likely result in the overactivation of their target G proteins, how does WDR-31 function with these GAP proteins? One possibility is that WDR31 may function downstream of

regulatory GTPases, including ARL2, ARL3, or unidentified GTPases, activated by these GAP

proteins. One downside of this explanation is that *wdr-31* single mutants did not have significant

ciliary and IFT defects; but, if this were valid, we would expect to observe further anomalies in

cilia and IFT in *wdr-31* single mutants, close to the removal of these three genes. Alternatively,

the activity/function of overactive GTPases can be somehow regulated by WDR-31, maybe WDR-

31 may have a GAP activity for the regulatory GTPases or it may have a GTPase activity. While

WDR-31 does not seem to have a GAP domain, HHMER search revealed that human WDR31 has

a distant sequence resemblance to a nucleoside-triphosphatase (NTPase) domain (NACHT)

containing protein in *Penicillium camemberti* (Gabler et al., 2020). PSI-BLAST search confirmed

this result (unpublished data).

Furthermore, the ciliary roles of these two GAP proteins might be independent of their GAP

activities, and consistent with this idea, the recent study showed the ciliary roles of ELMOD2 are

partially independent of its GAP activity (Turn et al., 2021). However, a variety of evidence

suggests that WDR31 and ELMOD3 do not form a complex. First, mass spectrometry-based

proteomic analysis for either WDR31 or ELMOD3 showed that neither WDR31 nor ELMOD3

contains each other or RP2 (unpublished data). Second, the proper localization of any of these

three proteins was independent of each other (unpublished data). Further studies are needed to

understand a mechanistic link between WDR31, ELMOD, and RP2, and figure out the independent

contributions of WDR31 and both GAP proteins in cilia biology.

A potential function for WDR31-ELMOD-RP2 in determining the integrity of ciliary gate

Our findings that worm lacking WDR31-ELMOD-RP2 also display ciliary entrance of non-ciliary

proteins (TRAM-1) suggests that these proteins may play a function in the ciliary gate. Despite the

mechanism by which these proteins regulate the ciliary gate functions remains unknown, the

abnormality of IFT proteins (abnormal distribution and increased anterograde IFT velocities) in

triple mutant may provide some hint about the potential roles of these proteins in ciliary gate. One

possible explanation is that the IFT defect in triple mutant potentially results in compromised

ciliary gate function. Consistent with this possibility, it was previously reported that in *C. elegans,*

the elimination of IFT complex A components and cytoplasmic dynein motor CHE-3 result in a

defect in ciliary gate function (Jensen et al., 2018; Scheidel and Blacque, 2018).

The WDR31-ELMOD-RP2 is needed for efficient IFT entry and IFT dynamics

Our findings show that simultaneous disruption of *WDR-31*, *ELMD-1*, and *RPI-2* causes

differential effects on ciliary IFT protein localizations, including accumulation of IFT-B

subcomplex components (IFT-74 and OSM-6), dynein (CHE-3) and kinesin motor (OSM-3) or

dim cilia staining of IFT-A subcomplex component and a dynein component (CHE-11 and XBX-

1) or almost no cilia entry of a BBSome component (BBS-7). Furthermore, with the exception of

cytoplasmic dynein CHE-3, the triple mutant shows a significant decrease in moving IFT particles

along cilia in both directions for almost all IFT proteins, indicating that the combined elimination

of WDR-31, ELMD-1, and RPI-2 has a significant impact on ciliary recruitment of IFT/BBSome

components (IFT-74, OSM-6, CHE-11, and BBS-7) and OSM-3. Our *in vivo* time lapse movie

analysis indicated that anterograde IFT velocities for OSM-3, OSM-6, IFT-74, and CHE-11 in the

middle segment of cilia are considerably increased in the triple mutant, while IFT velocities in the

distal segment of cilia and retrograde direction are unaltered (**Fig. 8A, B, C and D**). This suggests

that in cilia lacking *WDR-31, ELMD-1*, or *RPI-2*, certain portions of these IFT proteins are not

effectively loaded onto anterograde IFT.

652   Based on our results, which involve increase in anterograde IFT speed, a decrease in

653 moving IFT particles in anterograde and retrograde transport, ciliary tip aggregation of OSM-

654 3/KIF17 and IFT-B, and failure of docking of BBSome to the IFT machinery, we propose that

655 some of these defects likely stem from the failure of BBSome to associate with moving IFT in

656 cilia in double and triple mutants. These proteins are likely regulator for IFT/BBSome. Consistent

657 with our proposal, the almost comparable IFT/BBSome phenotypes were reported with

658 hypomorphic mutations in *dyf-2* (WDR19 orthologue, an IFT-A component) and *bbs-1* (a BBSome

659 component), where they showed that while BBSome remains at the base of cilia and does not

660 undergo IFT, the failure of IFT-B reassociation with IFT-A at the ciliary tips results in IFT-B

661 ciliary tip accumulations, despite the fact the association of IFT-A and IFT-B components in

662 anterograde transport persists (Wei et al., 2012). They proposed that the BBSome controls the IFT

663 assembly at the ciliary base and IFT turnover at the ciliary tip. However, several of the

664 abnormalities identified in the triple mutant, such as diminished IFT frequencies and ectopic

665 projection from the ciliary base, are unlikely to be related to BBSome dysfunction in the triple

666 mutant, because our findings revealed that the IFT frequency in the *bbs-8* mutant stays intact. Plus,

667 we did not observe ectopic project from the ciliary base in the *bbs-8* mutant (Data not shown).

669   However, the IFT defects in the triple mutant are unlikely to be due to mislocalization of

670 transition fiber proteins such as DYF-19 (the FBF1 orthologue), which was implicated in

671 regulating the ciliary entry of IFT and BBSome, since our data showed that the localization of

672 DYF-19 was not impaired in the triple mutants (Wei et al., 2013).

673   On the other hand, the effect on IFT might be direct since RP2 was previously proposed to

674 control the intraflagellar transport protein IFT20 pool in the peri-basal body and trafficking of

675 Kif17 and Kif7 to the ciliary tip (Schwarz et al., 2017). However, several interesting questions

have yet to be resolved: what is the mechanism by which WDR-31-ELMD-1-RPI-2 regulates the

ciliary entry of the BBSome entry? How does WDR-31-ELMD-1-RPI-2 interact with proteins,

including ARL6, that regulate the BBSome recruitment (Jin et al., 2010)? Future study research

will help us to understand how these proteins regulate BBSome/IFT trafficking. In summary, this

study provides a mechanistic explanation of the function of WDR-31, a new ciliary protein,

ELMD-1, and RPI-2 in the regulation of cilia biogenesis and also contributes a new regulator for

IFT/BBSome.

Acknowledgments

We thank Oliver Blacque, Piali Sengupta, Micheal Leroux, the National BioResource Project

(NBRP) in Japan, and the CGC in the United States, which is financed by the NIH Office of

Research Infrastructure Programs (P40 OD010440) for sharing valuable strains. We thank Oliver

Blacque and Samuel Katz for critical reading of the manuscript. We thank the Abdullah Gul

University Scientific Research Project Coordination Unit (Project number: TOA-2018-110) for

providing the funding that initiated the project. M.U. was supported by the Tistou & Charlotte

Kerstan Stiftung. The Leica laser scanning microscope was funded by a grant from Deutsche

Forschungsgemeinschaft (INST 2388/62-1).

Figure 1: WDR-31/WDR31 and ELMD-1/ELMOD proteins are evolutionary conserved

ciliary proteins A, B) Shown are the representative drawing of the PHA/PHB sensory neuron

(phasmid neurons located in the tail). Cilia, dendrite, axon and cell soma (cell body) are depicted

in the drawing. Fluorescence images from the transgenic strain carrying WDR-31::GFP or

GFP::ELMD-1 were displayed in the PHA/PHB sensory neurons. Scale bars: 3 μm **C, D)** Co-

localization of WDR-31::GFP (Green) with the IFT-140::mCherry (Red, an IFT-A component, a

ciliary marker) or TBG-1:mKate (Red, γ-Tubulin, the basal body) in the tail (phasmids) and head

(amphid) sensory neurons. TZ and BB denote the transition zone and the basal body, respectively

E, F) GFP::ELMD-1 (Green) localizes to the BB and PCMC (the periciliary membrane)

compartment) proximal to the transition zone. Co-labelling of GFP::ELMD-1 with MKS-

6:mCherry marker (transition zone) or IFT-140::mCherry in the tail (phasmids) and head (amphid)

neurons. Scale bars: 2 μm **G)** Shown are representative localizations of WDR-31 and ELMD-1.

A)

MERGE

ARL13B

10 μm

WDR31 (N) RFP

ELMOD3 (N) CFP

10 μm

B)

ELMOD3 (N) CFP

WDR31 (N) RFP

ARL13B

MERGE

MERGE

Figure 2: **ELMOD3 and WDR31 localize to the primary cilium**. **A, B**) Shown are the staining

of WDR31 (tagged with cyan fluorescent protein) and ELMOD3 (tagged with red fluorescent

protein) together with a ciliary marker ARL13B and DAPI (nucleus) in hTERT-RPE1 cells.

hTERT-RPE1 cells were transiently transfected with 100 ng of WDR31:CFP and ELMOD3:RFP.

Scale bars: 10 μm (A) and 3 μm (B)

Figure 3: WDR31 regulates ciliogenesis in zebrafish A) Shown are the expression pattern

analysis of WDR31 in zebrafish embryos. *Wdr31* is ubiquitously expressed before the

segmentation stage (2 cell and 8 somite stages (SS). The expression of *Wdr31* becomes limited to

the otic vesicle and brain region at 24 hours post fertilization (24 hpf). **B)** Shown are cilia of lateral

crista (LC) of the otic vesicle, stained with acetylated-tubulin, in wild type and *Wdr31* knockout,

generated via CRISPR/Cas9. **C and D**) The length of cilia in the lateral crista (LC) of the otic

vesicle remains unaffected in zebrafish *Wdr31* knockout while the cilia number is decreased, as

shown with the measurement of width of cilia lateral crista (LC) of the otic vesicle.

Figure 4: WDR31-ELMOD-RP2 are needed for determining cilia length and morphology in

C. elegans

A) Shown are co-localizations of RPI-2::mCherry (red, the endogenously labeled RPI-2) with

either WDR-31::GFP (green, the endogenously labeled WDR-31) or GFP::ELMD-1 (green,

overexpressed) in tail (phasmid) sensory neurons in *C. elegans.* Scale bars: 1 μm. **B)** The fraction

of the dye uptake defects is presented in bar charts for wild type and the indicated mutants. Fisher's

exact test was performed for statistical analysis between indicated triple mutants and a rescue gene

for Dye assay. Brackets show statistical significance between two strains compared ($p < 0.0001$

and **** indicate statistical significance). **C, D, E)** Shown is the jitter plot for PHA/PHB cilia,

ASER cilia and AWB short cilia length (μm) for wild type and indicated mutant strains. Statistical

significance between wild type and triple mutants was shown with bracket. **** implies statistical

significance which means that p value is lower than $p < 0.0001$ while * means that p value is lower

than $p < 0.05$. **F)** Fluorescence images show the morphology of AWB cilia (fork-like structure

located in the head), ASER cilia (amphid channel cilia) and PHA/PHB cilia (phasmid channel

cilia) in wild type and indicated mutant backgrounds. The backward projection from the cilia is

shown with asterisks (*) while p indicates the ectopic projections from the middle parts of cilia.

Scale bars: 3 μm **G)** The percentage of the abnormality in AWB cilia morphology is depicted in

bar charts. Fisher's exact test was utilized for statistical analysis of AWB cilia morphology between

wild type and designated mutants, and **** denotes statistical significance.

Figure 5: IFT-B proteins and OSM-3/KIF17 accumulate at the ciliary tip. Shown are fluorescent images of PHA/PHB cilia (phasmid tail). **A)** Fluorescent images from a single copy GFP::CHE-3 (human dynein heavy chain DYNC2H1) in wild type and indicated mutant backgrounds are displayed. GFP::CHE-3 accumulations within cilia and dim distal cilia staining

were observed in two distinct triple mutants (*wdr-31(tm10423);elmd-1;rpi-2* and *wdr-*

31(syb1568);elmd-1;rpi-2). Scale bars: 3 μm **B, C, D, E, F)** Confocal microscopy analysis of IFT-

A (IFT-140::GFP) and IFT-B complex components (OSM-6/IFT52::GFP and IFT-74::GFP)

revealed differential abnormalities in the transport of IFT-A and IFT-B components in double

(w*dr-31;elmd-1*) and triple mutants. The localization of XBX-1::mCherry (Dynein subunit) in

wdr-31(syb1568);elmd-1;rpi-2 triple mutants phenocopies the dim distal cilia staining of IFT-A

(IFT-140::GFP) in the w*dr-31(tm10423);elmd-1;rpi-2* triple mutants. OSM-3/KIF17 Kinesin

motor accumulates at the ciliary tips in w*dr-31(tm10423);elmd-1* double and w*dr-*

31(tm10423);elmd-1;rpi-2 triple mutants. Compared to double mutants, the ciliary tip staining is

stronger in the triple mutants. Scale bars: 3 μm **G)** Fluorescent images from Kinesin II motor

(KAP-1::GFP) revealed that the restricted middle segment localization of KAP-1 remains

unchanged in w*dr-31(tm10423);elmd-1;rpi-2* triple mutants. Scale bars: 3 μm **H)** Shown are the

drawings of phasmid cilia (PHA/PHB sensory neurons in the tail) in wild type and mutants

showing ciliary accumulations.

Figure 6: Measurement of anterograde and retrograde IFT transport frequency A, C, E and G) Shown are representative kymographs of GFP tagged IFT proteins translocating in the tail cilia (PHA/PHB sensory neurons) of wild type and indicated mutants. Kymographs for anterograde, retrogrades and merged (Red & Green) were generated with ImageJ equipped with KymographClear. The trajectory represents a moving IFT particle, and the average number of moving IFT particles in wild type and indicated mutants (double, triple) was calculated by counting all trajectories in each kymograph. Travel time and distance are shown on kymographs. **B, D, F and H)** Box-and-Whisker charts with error bars were created to visualize the average number of IFT anterograde and retrograde particles between wild type and indicated mutants. The Mann–Whitney U test was used to measure statistical analysis and significance. The four and three

 asterisks (**** and ***) at the top of the brackets indicate that the p value between the two strains

is less than 0.0001 and 0.001, respectively, suggesting statistical significance. Ns stands for "not

significant."

Figure 7: A non-ciliary membrane protein TRAM-1 enters into cilia in *wdr-31;elmd-1;rpi-2*

triple mutants. A) Confocal fluorescent images exhibit the localization of tdTomato tagged

TRAM-1 (a PCMC marker) and MSK-2::GFP (a TZ marker) in wild type and indicated mutants.

TRAM-1 leaks into cilia in all three *wdr-31* triple mutants. Cilia, the periciliary membrane

compartment (PCMC), and transition zone (TZ) are depicted in the fluorescent image. Scale bars:

2 μm, **B)** TRAM-1 fluorescence intensities in cilia were measured in wild type and designated

mutants, and the results were shown in the plot. The four asterisks (****) indicate statistically

significant differences between the wild type and the identified triple mutants. **C)** The localization

of NPHP-1::GFP (a transition zone protein) and FBF-1::mCherry (a transition fiber protein) was

similar unaffected in *wdr-31;elmd-1;rpi-2* triple mutants **D)** The PLCδ1-PH::GFP (a marker for

monitoring phosphatidylinositol 4,5-bisphosphate (PtdIns(4,5)P2) in the plasma membrane)

decorates the membranes of PCMC and does not enter into cilia in wild type. MKSR-2 was used

to mark the transition zone. The PLCδ1-PH::GFP stays outside of cilia in the *wdr-31;elmd-1;rpi-*

2 triple mutants.

Figure 8: The average anterograde IFT velocities were significantly increased in *wdr-*
***13;elmd-1;rpi-2* triple mutants, A, B, C, and D)** The average anterograde (distal and middle

segments) and retrograde IFT velocities were calculated, and box-and-whisker plots with error

bars were used to depict them. Numbers shown at the top of bars represent the number of IFT

particles used for determining IFT velocities. To compare statistical analysis and significance

between wild type and specified mutations, the Mann–Whitney U test was performed. A p value

of less than 0.0001 indicates statistical significance between the two strains, as shown by the four

asterisks (****) at the top of the brackets.

Figure 9: WDR-31 and ELMD-1 regulate the recruitment of BBSome to cilia.

A) Shown are fluorescence images from the transgenic strain carrying IFT-74::GFP, an IFT-B component, in wild type, *wdr-31;elmd-1* double mutants, *wdr-31;elmd-1;rpi-2* and *wdr-31;elmd-1;bbs-8* triple mutants, and *bbs-8(nx77)*. The IFT-B subunit IFT-74:GFP accumulates at the ciliary

tips and cilia in the tail of all three mutants. **B)** Confocal fluorescence images showing the

localization of BBS-7:GFP, a BBSome subunit, in the heads and tails of wild type and *wdr-*

31(tm10423); wdr-31;elmd-1; *rpi-2;elmd-1* double and *wdr-31;elmd-1;rpi-2* triple mutants.

Fluorescence images showed absent or weak cilia staining of BBS-7::GFP in both the head and

tails of *wdr-31;elmd-1* and *wdr-31;elmd-1;rpi-2* triple mutants. **C)** Kymographs were created from

time-lapse BBS-7::GFP movies (PHA/PHB cilia) using KymographClear integrated into ImageJ.

Shown are representative kymographs for BBS-7::GFP translocating in wild type and indicated

mutants. Each trajectory in kymographs was counted. Travel time and distance are included on

kymograph **D)** The graph depicts the average number of BBS-7::GFP particles traveling around

cilia in both directions for wild type and indicated mutants. The Mann–Whitney U test revealed

statistical significance between the compared strains and that the p value was less than 0.0001

shown by the four asterisks (****) at the top of the brackets **E)** In wild type, the assembly of the

Kinesin-IFT-BBSome complex (Kinesin-II and OSM-3**,** IFT-B , IFT-A and BBSome) happens at

the base of cilia. In the middle segment of amphid and phasmid cilia in *C. elegans*, both

heterotrimeric Kinesin II and homodimeric OSM-3 transport the IFT-A- IFT-B- BBSome complex

in an anterograde direction. Heterotrimeric Kinesin II returns to the ciliary base when it reaches

the tip of the middle segment of amphid and phasmid cilia, whereas homodimeric OSM-3 is

responsible for the anterograde translocation of the IFT-BBSome complex in the distal segment of

amphid and phasmid cilia. When the OSM-3-IFT-BBSome complex reaches the ciliary tip,

cytoplasmic dynein transports them back to the ciliary base. In *wdr-31;elmd-1* double *wdr-*

31;elmd-1;rpi-2 triple mutants, the BBSome failed to enter into cilia, thus leading to accumulations

of OSM-3 and IFT-B components in the ciliary tips.

Supplementary Figure 1: Expression patterns of WDR-31 and ELMD-1. A) A total of 60 sensory neurons are distributed in the head (amphid) and tails (phasmid) of *C. elegans.* The ciliated sensory neurons are displayed in the schematic of *C. elegans.* The expression of *elmd-1promoter::gfp* (1000 kb) and *wdr-31promoter::gfp* (1000 kb) were shown in the fluorescence images. The cell of ciliated sensory neurons (cell soma) in the head and tails were displayed in brackets.

Supplementary Figure 2: Ciliary localization of WDR31 and ELMOD3 in hTERT-RPE1

cells A) Images showing immunostaining of a ciliary marker ARL13B and DAPI (nucleus) in

hTERT-RPE1 cells. NT represents no transfection of hTERT-RPE1 with WDR31 (tagged with

cyan fluorescent protein) and ELMOD3 (tagged with red fluorescent protein). Scale bars: 10 µm.

B) Images showing immunostaining of hTERT-RPE1 cells transfected with ELMOD3-CFP (cyan)

and also stained for ARL13B (yellow). The magenta channel only shows some background but no

localization to the cilium. Green regions in the "merge" picture show co-localization of both

ELMOD3 and ARL13B proteins. Scale bars: 3 µm. **C)** Fluorescence images displaying hTERT-

RPE1 Fluorescence images of immunostaining for ARL13B (yellow) and WDR31-RFP (magenta)

in hTERT-RPE1 cells transfected with WDR31-RFP (cyan). The cyan channel shows some

870   background, with no cilium localisation. WDR31 and ARL1B proteins are co-localized in the pink

871   sections of the "merge" image. Scale bars: 3 μm.

A

Human WDR31
1 — 367
NP_660284.1

WD 99-140 WD 142-183 WD 184-216 WD 274-319

C.elegans WDR-31
1 — 378
NP_494344.1

WD 139-179 WD 193-219 WD 266-310

B

250 bp

syb1568

tur003

tm10423

wdr-31

syb603

elmo-3

ok1863

rpi-2

C
wdr-31(tm10423);rpi-2;elmd-1

GFP::ELMD-1   Dye   Merge

D
RPI-2::GFP   Dye   Merge

RPI-2::GFP   Dye   Merge

Supplementary Figure 3: Rescue of dye uptake defects by ELMD-1 or RPI-2.

A) Shown are representative schematics of human and *C. elegans* WDR-31/WDR31. Human

WDR31 (NP_660284.1) has four WD domains while *C. elegans* WDR-31 (NP_494344.1) has

three WD domains (https://prosite.expasy.org/; Sigrist CJA et al 2012). **B)** Schematic diagrams of

wdr-31, *elmd-1* and *rpi-2* together with corresponding deletions are shown. Scale bars: 250 bp. **C**

and D) Fluorescence microscope images of mutant worms (head and tail) with red-fluorescent dye

uptake (texas red filter) and GFP-tagged ELMD-1 and RPI-2 (fluorescence filter set for GFP) are

shown. The head or tail in the dotted lines is indicated by numbers in the combined fluorescence

images. GFP expression in combination with RED fluorescence indicates the rescue of dye uptake

defects.

A

AWB long cilia length (μm)

Supplementary Figure 4: AWB cilia morphology in WDR-31-ELMD-1-RPI-2 triple mutants

A) Shown are the schematic representations of lipophilic fluorescent dye uptake in the head

(amphid) and tail (phasmid) sensory neurons in *C. elegans.* Red labelling indicates the normal dye

uptake while the failure of Dye uptake was shown in white. Fluorescence images show the dye

update in the head and tail neurons in the wild type and indicated mutant strains. No dye uptake

was observed in both head (amphid) and tail (phasmid) of *wdr-31(tm10423);elmd-1* double

mutants, *wdr-31(tm10423);elmd-1;rpi-2* and *wdr-31(syb1568);elmd-1;rpi-2* triple mutants.

Ex[elmd-1(+)] rescues the dye uptake defects of *wdr-31(tm10423);elmd-1* double and *wdr-*

31(tm10423);elmd-1;rpi-2 triple mutants, while *Ex[wdr-31(+)]* rescues the Dye uptake defects of

wdr-31(syb1568);elmd-1;rpi-2 triple mutants. Scale bars: 10 μm. **B)** Shown is the jitter plot for

AWB long cilia length (μm) in wild type and mutant strains.

Supplementary Figure 5: A) Shown is co-localization of RPI-2::GFP (human RP2) and XBX-1::mCherry (human DYNC2LI1) in the head and tail sensory neurons. XBX-1 stains the entire cilium and is used as a ciliary marker while RPI-2 is located at PCMC. PCMC denotes the periciliary membrane compartment.

Supplementary Figure 6: A) Confocal images (Z-stack) display the localization of OSM-6::GFP

(a single copy transgene, huma IFT52) in the head (a bunch of cilia) of wild type and indicated

mutants. Arrow points the ciliary tips while asterisks indicate the ciliary base. **B)** Confocal images

show colocalization of TAX-4 (a ciliary membrane protein) and XBX-1::tdTomato (a cilia

marker).

Supplementary Figure 7: A) The architecture of AWB cilia (a fork-like structure in the head) in

wild type and mutant backgrounds is shown in fluorescent images. In schematic illustrations, the

AWB cilia in wild type and altered mutants are illustrated. **B)** Shown are bar plots displaying

percentage of phenotypes in wild type and indicated triple mutants. Four asterisks (****) indicates

statistical significances between wild type and indicated triple mutants.

Supplementary Movies

For the IFT assay, time-lapse movies (3 frames per second) were generated with Leica DM6, and

were processed with Image J to generate GIFs (10 fps).

Supplementary Movies:

Supplementary Movie 1: IFT-74::GFP in wild type and indicated mutant backgrounds.

Supplementary Movie 2: OSM-6::GFP in wild type and indicated mutant backgrounds.

Supplementary Movie 3: OSM-3::GFP in wild type and indicated mutant backgrounds

Supplementary Movie 4: GFP::CHE-3 in wild type and indicated mutant backgrounds.

Supplementary Movie 5: IFT-140::GFP in wild type and indicated mutant backgrounds.

Supplementary Movie 6: BBS-7::GFP in wild type and indicated mutant backgrounds.

References

Ansley, S.J., Badano, J.L., Blacque, O.E., Hill, J., Hoskins, B.E., Leitch, C.C., Chul Kim, J.,
Ross, A.J., Eichers, E.R., Teslovich, T.M., Mah, A.K., Johnsen, R.C., Cavender, J.C.,
Alan Lewis, R., Leroux, M.R., Beales, P.L., Katsanis, N., 2003. Basal body dysfunction
is a likely cause of pleiotropic Bardet–Biedl syndrome. Nature 425, 628–633.
https://doi.org/10.1038/nature02030
930
931 Anvarian, Z., Mykytyn, K., Mukhopadhyay, S., Pedersen, L.B., Christensen, S.T., 2019. Cellular
932   signalling by primary cilia in development, organ function and disease. Nat. Rev.
933   Nephrol. 15, 199–219. https://doi.org/10.1038/s41581-019-0116-9
934
935 Arnaiz, O., Malinowska, A., Klotz, C., Sperling, L., Dadlez, M., Koll, F., Cohen, J., 2009. Cildb:
936   a knowledgebase for centrosomes and cilia. Database 2009.
937   https://doi.org/10.1093/database/bap022
938

939   Avidor-Reiss, T., Maer, A.M., Koundakjian, E., Polyanovsky, A., Keil, T., Subramaniam, S.,
940   Zuker, C.S., 2004. Decoding Cilia Function. Cell 117, 527–539.
941   https://doi.org/10.1016/S0092-8674(04)00412-X
942
943   Blacque, O., E., 2008. Intraflagellar transport: from molecular characterisation to mechanism.
944   Front. Biosci. 13, 2633. https://doi.org/10.2741/2871
945
946   Blacque, O.E., 2004. Loss of C. elegans BBS-7 and BBS-8 protein function results in cilia
947   defects and compromised intraflagellar transport. Genes Dev. 18, 1630–1642.
948   https://doi.org/10.1101/gad.1194004
949
950   Blacque, O.E., Li, C., Inglis, P.N., Esmail, M.A., Ou, G., Mah, A.K., Baillie, D.L., Scholey,
951   J.M., Leroux, M.R., 2006. The WD Repeat-containing Protein IFTA-1 Is Required for
952   Retrograde Intraflagellar Transport. Mol. Biol. Cell 17, 5053–5062.
953   https://doi.org/10.1091/mbc.e06-06-0571
954
955   Blacque, O.E., Perens, E.A., Boroevich, K.A., Inglis, P.N., Li, C., Warner, A., Khattra, J., Holt,
956   R.A., Ou, G., Mah, A.K., McKay, S.J., Huang, P., Swoboda, P., Jones, S.J.M., Marra,
957   M.A., Baillie, D.L., Moerman, D.G., Shaham, S., Leroux, M.R., 2005. Functional
958   Genomics of the Cilium, a Sensory Organelle. Curr. Biol. 15, 935–941.
959   https://doi.org/10.1016/j.cub.2005.04.059
960
961   Blacque, O.E., Sanders, A.A., 2014. Compartments within a compartment: What *C. elegans* can
962   tell us about ciliary subdomain composition, biogenesis, function, and disease.
963   Organogenesis 10, 126–137. https://doi.org/10.4161/org.28830
964
965   Bloodgood, R.A., 2009. From Central to Rudimentary to Primary: The History of an
966   Underappreciated Organelle Whose Time Has Come.The Primary Cilium, in: Methods in
967   Cell Biology. Elsevier, pp. 2–52. https://doi.org/10.1016/S0091-679X(08)94001-2
968
969   Brenner, S., 1974. The genetics of Caenorhabditis elegans. Genetics 77, 71–94.
970
971   Breslow, D.K., Hoogendoorn, S., Kopp, A.R., Morgens, D.W., Vu, B.K., Kennedy, M.C., Han,
972   K., Li, A., Hess, G.T., Bassik, M.C., Chen, J.K., Nachury, M.V., 2018. A CRISPR-based
973   screen for Hedgehog signaling provides insights into ciliary function and ciliopathies.
974   Nat. Genet. 50, 460–471. https://doi.org/10.1038/s41588-018-0054-7
975
976   Chang, N., Sun, C., Gao, L., Zhu, D., Xu, X., Zhu, X., Xiong, J.-W., Xi, J.J., 2013. Genome
977   editing with RNA-guided Cas9 nuclease in Zebrafish embryos. Cell Res. 23, 465–472.
978   https://doi.org/10.1038/cr.2013.45
979
980   Choksi, S.P., Babu, D., Lau, D., Yu, X., Roy, S., 2014. Systematic discovery of novel ciliary
981   genes through functional genomics in the zebrafish. Development 141, 3410–3419.
982   https://doi.org/10.1242/dev.108209
983
984   Dickinson, D.J., Ward, J.D., Reiner, D.J., Goldstein, B., 2013. Engineering the Caenorhabditis

985   elegans genome using Cas9-triggered homologous recombination. Nat. Methods 10,
986   1028–1034. https://doi.org/10.1038/nmeth.2641

988   Evans, R.J., Schwarz, N., Nagel-Wolfrum, K., Wolfrum, U., Hardcastle, A.J., Cheetham, M.E.,
989   2010. The retinitis pigmentosa protein RP2 links pericentriolar vesicle transport between
990   the Golgi and the primary cilium. Hum. Mol. Genet. 19, 1358–1367.
991   https://doi.org/10.1093/hmg/ddq012

993   Gabler, F., Nam, S., Till, S., Mirdita, M., Steinegger, M., Söding, J., Lupas, A.N., Alva, V.,
994   2020. Protein Sequence Analysis Using the MPI Bioinformatics Toolkit. Curr. Protoc.
995   Bioinforma. 72. https://doi.org/10.1002/cpbi.108

998   Herman, R.K., Hedgecock, E.M., 1990. Limitation of the size of the vulval primordium of
999   Caenorhabditis elegans by lin-15 expression in surrounding hypodermis. Nature 348,
169–171. https://doi.org/10.1038/348169a0

Ivanova, A.A., East, M.P., Yi, S.L., Kahn, R.A., 2014. Characterization of Recombinant
ELMOD (Cell Engulfment and Motility Domain) Proteins as GTPase-activating Proteins
(GAPs) for ARF Family GTPases. J. Biol. Chem. 289, 11111–11121.
https://doi.org/10.1074/jbc.M114.548529

Jaworek, T.J., Richard, E.M., Ivanova, A.A., Giese, A.P.J., Choo, D.I., Khan, S.N., Riazuddin,
Sheikh, Kahn, R.A., Riazuddin, Saima, 2013. An Alteration in ELMOD3, an Arl2
GTPase-Activating Protein, Is Associated with Hearing Impairment in Humans. PLoS
Genet. 9, e1003774. https://doi.org/10.1371/journal.pgen.1003774

Jensen, V.L., Carter, S., Sanders, A.A.W.M., Li, C., Kennedy, J., Timbers, T.A., Cai, J.,
Scheidel, N., Kennedy, B.N., Morin, R.D., Leroux, M.R., Blacque, O.E., 2016. Whole-
Organism Developmental Expression Profiling Identifies RAB-28 as a Novel Ciliary
GTPase Associated with the BBSome and Intraflagellar Transport. PLOS Genet. 12,
e1006469. https://doi.org/10.1371/journal.pgen.1006469

Jensen, V.L., Lambacher, N.J., Li, C., Mohan, S., Williams, C.L., Inglis, P.N., Yoder, B.K.,
Blacque, O.E., Leroux, M.R., 2018. Role for intraflagellar transport in building a
functional transition zone. EMBO Rep. 19. https://doi.org/10.15252/embr.201845862

Jin, H., White, S.R., Shida, T., Schulz, S., Aguiar, M., Gygi, S.P., Bazan, J.F., Nachury, M.V.,
2010. The Conserved Bardet-Biedl Syndrome Proteins Assemble a Coat that Traffics
Membrane Proteins to Cilia. Cell 141, 1208–1219.
https://doi.org/10.1016/j.cell.2010.05.015

Johnson, K.R., Longo-Guess, C.M., Gagnon, L.H., 2012. Mutations of the Mouse ELMO
Domain Containing 1 Gene (Elmod1) Link Small GTPase Signaling to Actin
Cytoskeleton Dynamics in Hair Cell Stereocilia. PLoS ONE 7, e36074.
https://doi.org/10.1371/journal.pone.0036074

1031
1032
1033 Kaplan, O.I., Doroquez, D.B., Cevik, S., Bowie, R.V., Clarke, L., Sanders, A.A.W.M., Kida, K.,
1034   Rappoport, J.Z., Sengupta, P., Blacque, O.E., 2012. Endocytosis Genes Facilitate Protein
1035   and Membrane Transport in C. elegans Sensory Cilia. Curr. Biol. 22, 451–460.
1036   https://doi.org/10.1016/j.cub.2012.01.060
1037
Lambacher, N.J., Bruel, A.-L., van Dam, T.J.P., Szymańska, K., Slaats, G.G., Kuhns, S.,
McManus, G.J., Kennedy, J.E., Gaff, K., Wu, K.M., van der Lee, R., Burglen, L.,
Doummar, D., Rivière, J.-B., Faivre, L., Attié-Bitach, T., Saunier, S., Curd, A., Peckham,
M., Giles, R.H., Johnson, C.A., Huynen, M.A., Thauvin-Robinet, C., Blacque, O.E.,
2016. TMEM107 recruits ciliopathy proteins to subdomains of the ciliary transition zone
and causes Joubert syndrome. Nat. Cell Biol. 18, 122–131.
https://doi.org/10.1038/ncb3273
1045
1046 Lechtreck, K.-F., Johnson, E.C., Sakai, T., Cochran, D., Ballif, B.A., Rush, J., Pazour, G.J.,
1047   Ikebe, M., Witman, G.B., 2009. The Chlamydomonas reinhardtii BBSome is an IFT
1048   cargo required for export of specific signaling proteins from flagella. J. Cell Biol. 187,
1049   1117–1132. https://doi.org/10.1083/jcb.200909183
1050
1051 Lee, E., Sivan-Loukianova, E., Eberl, D.F., Kernan, M.J., 2008. An IFT-A protein is required to
1052   delimit functionally distinct zones in mechanosensory cilia. Curr. Biol. CB 18, 1899–
1053   1906. https://doi.org/10.1016/j.cub.2008.11.020
1054
Li, J.B., Gerdes, J.M., Haycraft, C.J., Fan, Y., Teslovich, T.M., May-Simera, H., Li, H., Blacque,
O.E., Li, L., Leitch, C.C., Lewis, R.A., Green, J.S., Parfrey, P.S., Leroux, M.R.,
Davidson, W.S., Beales, P.L., Guay-Woodford, L.M., Yoder, B.K., Stormo, G.D.,
Katsanis, N., Dutcher, S.K., 2004. Comparative Genomics Identifies a Flagellar and
Basal Body Proteome that Includes the BBS5 Human Disease Gene. Cell 117, 541–552.
https://doi.org/10.1016/S0092-8674(04)00450-7
1061
1062 Liem, K.F., Ashe, A., He, M., Satir, P., Moran, J., Beier, D., Wicking, C., Anderson, K.V., 2012.
1063   The IFT-A complex regulates Shh signaling through cilia structure and membrane protein
1064   trafficking. J. Cell Biol. 197, 789–800. https://doi.org/10.1083/jcb.201110049
1065
1066 Loktev, A.V., Zhang, Q., Beck, J.S., Searby, C.C., Scheetz, T.E., Bazan, J.F., Slusarski, D.C.,
1067   Sheffield, V.C., Jackson, P.K., Nachury, M.V., 2008. A BBSome Subunit Links
1068   Ciliogenesis, Microtubule Stability, and Acetylation. Dev. Cell 15, 854–865.
1069   https://doi.org/10.1016/j.devcel.2008.11.001
1070
1071 Mangeol, P., Prevo, B., Peterman, E.J.G., 2016. KymographClear and KymographDirect: two
1072   tools for the automated quantitative analysis of molecular and cellular dynamics using
1073   kymographs. Mol. Biol. Cell 27, 1948–1957. https://doi.org/10.1091/mbc.E15-06-0404
1074
1075 Mick, D.U., Rodrigues, R.B., Leib, R.D., Adams, C.M., Chien, A.S., Gygi, S.P., Nachury, M.V.,
1076   2015. Proteomics of Primary Cilia by Proximity Labeling. Dev. Cell 35, 497–512.

1077   https://doi.org/10.1016/j.devcel.2015.10.015
1078
1079   Miryounesi, M., Bahari, S., Salehpour, S., Alipour, N., Ghafouri-Fard, S., 2019. ELMO Domain
1080   Containing 1 (ELMOD1) Gene Mutation Is Associated with Mental Retardation and
1081   Autism Spectrum Disorder. J. Mol. Neurosci. 69, 312–315.
1082   https://doi.org/10.1007/s12031-019-01359-z
1083
1084   Mukhopadhyay, S., Wen, X., Chih, B., Nelson, C.D., Lane, W.S., Scales, S.J., Jackson, P.K.,
1085   2010. TULP3 bridges the IFT-A complex and membrane phosphoinositides to promote
1086   trafficking of G protein-coupled receptors into primary cilia. Genes Dev. 24, 2180–2193.
1087   https://doi.org/10.1101/gad.1966210
1088
1089   Nachury, M.V., 2014. How do cilia organize signalling cascades? Philos. Trans. R. Soc. Lond.
1090   B. Biol. Sci. 369. https://doi.org/10.1098/rstb.2013.0465
1091
1092   Nachury, M.V., Loktev, A.V., Zhang, Q., Westlake, C.J., Peränen, J., Merdes, A., Slusarski,
1093   D.C., Scheller, R.H., Bazan, J.F., Sheffield, V.C., Jackson, P.K., 2007. A Core Complex
1094   of BBS Proteins Cooperates with the GTPase Rab8 to Promote Ciliary Membrane
1095   Biogenesis. Cell 129, 1201–1213. https://doi.org/10.1016/j.cell.2007.03.053
1096
1097   Nozaki, S., Castro Araya, R.F., Katoh, Y., Nakayama, K., 2019. Requirement of IFT-B–BBSome
1098   complex interaction in export of GPR161 from cilia. Biol. Open 8, bio043786.
1099   https://doi.org/10.1242/bio.043786
1100
1101   Ou, G., E. Blacque, O., Snow, J.J., Leroux, M.R., Scholey, J.M., 2005. Functional coordination
1102   of intraflagellar transport motors. Nature 436, 583–587.
1103   https://doi.org/10.1038/nature03818
1104
1105   Ou, G., Koga, M., Blacque, O.E., Murayama, T., Ohshima, Y., Schafer, J.C., Li, C., Yoder, B.K.,
1106   Leroux, M.R., Scholey, J.M., 2007. Sensory Ciliogenesis in *Caenorhabditis elegans* :
1107   Assignment of IFT Components into Distinct Modules Based on Transport and
1108   Phenotypic Profiles. Mol. Biol. Cell 18, 1554–1569. https://doi.org/10.1091/mbc.e06-09-
1109   0805
1110
1111   Patel-King, R.S., Gilberti, R.M., Hom, E.F.Y., King, S.M., 2013. WD60/FAP163 is a dynein
1112   intermediate chain required for retrograde intraflagellar transport in cilia. Mol. Biol. Cell
1113   24, 2668–2677. https://doi.org/10.1091/mbc.E13-05-0266
1114
1115   Pazour, G.J., Dickert, B.L., Vucica, Y., Seeley, E.S., Rosenbaum, J.L., Witman, G.B., Cole,
1116   D.G., 2000. Chlamydomonas IFT88 and its mouse homologue, polycystic kidney disease
1117   gene tg737, are required for assembly of cilia and flagella. J. Cell Biol. 151, 709–718.
1118   https://doi.org/10.1083/jcb.151.3.709
1119
1120   Piasecki, B.P., Burghoorn, J., Swoboda, P., 2010. Regulatory Factor X (RFX)-mediated
1121   transcriptional rewiring of ciliary genes in animals. Proc. Natl. Acad. Sci. 107, 12969–
1122   12974. https://doi.org/10.1073/pnas.0914241107

1123

1124 Prevo, B., Scholey, J.M., Peterman, E.J.G., 2017. Intraflagellar transport: mechanisms of motor
1125   action, cooperation, and cargo delivery. FEBS J. 284, 2905–2931.
1126   https://doi.org/10.1111/febs.14068

1127

1128 Reiter, J.F., Leroux, M.R., 2017. Genes and molecular pathways underpinning ciliopathies. Nat.
1129   Rev. Mol. Cell Biol. 18, 533–547. https://doi.org/10.1038/nrm.2017.60

1130

1131

1132 Rompolas, P., Pedersen, L.B., Patel-King, R.S., King, S.M., 2007. *Chlamydomonas* FAP133 is a
1133   dynein intermediate chain associated with the retrograde intraflagellar transport motor. J.
1134   Cell Sci. 120, 3653–3665. https://doi.org/10.1242/jcs.012773

1135

1136 Rosenbaum, J.L., Witman, G.B., 2002. Intraflagellar transport. Nat. Rev. Mol. Cell Biol. 3, 813–
1137   825. https://doi.org/10.1038/nrm952

1138

1139 Ruiz García, S., Deprez, M., Lebrigand, K., Cavard, A., Paquet, A., Arguel, M.-J., Magnone, V.,
1140   Truchi, M., Caballero, I., Leroy, S., Marquette, C.-H., Marcet, B., Barbry, P., Zaragosi,
1141   L.-E., 2019. Novel dynamics of human mucociliary differentiation revealed by single-cell
1142   RNA sequencing of nasal epithelial cultures. Development dev.177428.
1143   https://doi.org/10.1242/dev.177428

1144

1145 Satir, P., Christensen, S.T., 2007. Overview of Structure and Function of Mammalian Cilia.
1146   Annu. Rev. Physiol. 69, 377–400.
1147   https://doi.org/10.1146/annurev.physiol.69.040705.141236

1148

1149 Saudi Mendeliome Group, 2015. Comprehensive gene panels provide advantages over clinical
1150   exome sequencing for Mendelian diseases. Genome Biol. 16, 134.
1151   https://doi.org/10.1186/s13059-015-0693-2

1152

1153 Scheidel, N., Blacque, O.E., 2018. Intraflagellar Transport Complex A Genes Differentially
1154   Regulate Cilium Formation and Transition Zone Gating. Curr. Biol. 28, 3279-3287.e2.
1155   https://doi.org/10.1016/j.cub.2018.08.017

1156

1157 Schneider, C.A., Rasband, W.S., Eliceiri, K.W., 2012. NIH Image to ImageJ: 25 years of image
1158   analysis. Nat. Methods 9, 671–675. https://doi.org/10.1038/nmeth.2089

1159

1160 Scholey, J., 2007. The sensory cilia of Caenorhabditis elegans_Revised. WormBook.
1161   https://doi.org/10.1895/wormbook.1.126.2

1162

1163 Schwarz, N., Lane, A., Jovanovic, K., Parfitt, D.A., Aguila, M., Thompson, C.L., da Cruz, L.,
1164   Coffey, P.J., Chapple, J.P., Hardcastle, A.J., Cheetham, M.E., 2017. Arl3 and RP2
1165   regulate the trafficking of ciliary tip kinesins. Hum. Mol. Genet. 26, 2480–2492.
1166   https://doi.org/10.1093/hmg/ddx143

1167

Shaheen, R., Szymanska, K., Basu, B., Patel, N., Ewida, N., Faqeih, E., Al Hashem, A., Derar,

N., Alsharif, H., Aldahmesh, M.A., Alazami, A.M., Hashem, M., Ibrahim, N.,
Abdulwahab, F.M., Sonbul, R., Alkuraya, H., Alnemer, M., Al Tala, S., Al-Husain, M.,
Morsy, H., Seidahmed, M.Z., Meriki, N., Al-Owain, M., AlShahwan, S., Tabarki, B.,
Salih, M.A., Ciliopathy WorkingGroup, Faquih, T., El-Kalioby, M., Ueffing, M., Boldt,
K., Logan, C.V., Parry, D.A., Al Tassan, N., Monies, D., Megarbane, A., Abouelhoda,
M., Halees, A., Johnson, C.A., Alkuraya, F.S., 2016. Characterizing the morbid genome
of ciliopathies. Genome Biol. 17, 242. https://doi.org/10.1186/s13059-016-1099-5
1176

Shamseldin, H.E., Shaheen, R., Ewida, N., Bubshait, D.K., Alkuraya, H., Almardawi, E.,
Howaidi, A., Sabr, Y., Abdalla, E.M., Alfaifi, A.Y., Alghamdi, J.M., Alsagheir, A.,
Alfares, A., Morsy, H., Hussein, M.H., Al-Muhaizea, M.A., Shagrani, M., Al Sabban, E.,
Salih, M.A., Meriki, N., Khan, R., Almugbel, M., Qari, A., Tulba, M., Mahnashi, M.,
Alhazmi, K., Alsalamah, A.K., Nowilaty, S.R., Alhashem, A., Hashem, M., Abdulwahab,
F., Ibrahim, N., Alshidi, T., AlObeid, E., Alenazi, M.M., Alzaidan, H., Rahbeeni, Z., Al-
Owain, M., Sogaty, S., Seidahmed, M.Z., Alkuraya, F.S., 2020. The morbid genome of
ciliopathies: an update. Genet. Med. Off. J. Am. Coll. Med. Genet. 22, 1051–1060.
https://doi.org/10.1038/s41436-020-0761-1
1186

1187 Sigg, M.A., Menchen, T., Lee, C., Johnson, J., Jungnickel, M.K., Choksi, S.P., Garcia, G.,
1188   Busengdal, H., Dougherty, G.W., Pennekamp, P., Werner, C., Rentzsch, F., Florman,
1189   H.M., Krogan, N., Wallingford, J.B., Omran, H., Reiter, J.F., 2017. Evolutionary
1190   Proteomics Uncovers Ancient Associations of Cilia with Signaling Pathways. Dev. Cell
1191   43, 744-762.e11. https://doi.org/10.1016/j.devcel.2017.11.014
1192

1193 Silflow, C.D., Lefebvre, P.A., 2001. Assembly and Motility of Eukaryotic Cilia and Flagella.
1194   Lessons from *Chlamydomonas reinhardtii*. Plant Physiol. 127, 1500–1507.
1195   https://doi.org/10.1104/pp.010807
1196

1197 Sleigh, M.A., 1989. Adaptations of ciliary systems for the propulsion of water and mucus.
1198   Comp. Biochem. Physiol. A Physiol. 94, 359–364. https://doi.org/10.1016/0300-
1199   9629(89)90559-8
1200

1201 SYSCILIA Study Group, van Dam, T.J., Wheway, G., Slaats, G.G., Huynen, M.A., Giles, R.H.,
1202   2013. The SYSCILIA gold standard (SCGSv1) of known ciliary components and its
1203   applications within a systems biology consortium. Cilia 2, 7.
1204   https://doi.org/10.1186/2046-2530-2-7
1205

1206 Thisse, B., Heyer, V., Lux, A., Alunni, V., Degrave, A., Seiliez, I., Kirchner, J., Parkhill, J.-P.,
1207   Thisse, C., 2004. Spatial and Temporal Expression of the Zebrafish Genome by Large-
1208   Scale In Situ Hybridization Screening, in: Methods in Cell Biology. Elsevier, pp. 505–
1209   519. https://doi.org/10.1016/S0091-679X(04)77027-2
1210

1211 Turn, R.E., Linnert, J., Gigante, E.D., Wolfrum, U., Caspary, T., Kahn, R.A., 2021. Roles for
1212   ELMOD2 and Rootletin in ciliogenesis. Mol. Biol. Cell 32, 800–822.
1213   https://doi.org/10.1091/mbc.E20-10-0635
1214

1215   Turan MG, Kantarci H, Temtek SD, Cakici O, Cevik S, Kaplan OI. 2022., Protocol for
1216   determining the average speed and frequency of kinesin and dynein-driven intraflagellar
1217   transport (IFT) in C. elegans. STAR Protoc;3(3):101498. doi:
1218   10.1016/j.xpro.2022.101498.

1220   UK10K Rare Diseases Group, Boldt, K., van Reeuwijk, J., Lu, Q., Koutroumpas, K., Nguyen,
1221   T.-M.T., Texier, Y., van Beersum, S.E.C., Horn, N., Willer, J.R., Mans, D.A., Dougherty,
1222   G., Lamers, I.J.C., Coene, K.L.M., Arts, H.H., Betts, M.J., Beyer, T., Bolat, E.,
1223   Gloeckner, C.J., Haidari, K., Hetterschijt, L., Iaconis, D., Jenkins, D., Klose, F., Knapp,
1224   B., Latour, B., Letteboer, S.J.F., Marcelis, C.L., Mitic, D., Morleo, M., Oud, M.M.,
1225   Riemersma, M., Rix, S., Terhal, P.A., Toedt, G., van Dam, T.J.P., de Vrieze, E.,
1226   Wissinger, Y., Wu, K.M., Apic, G., Beales, P.L., Blacque, O.E., Gibson, T.J., Huynen,
1227   M.A., Katsanis, N., Kremer, H., Omran, H., van Wijk, E., Wolfrum, U., Kepes, F., Davis,
1228   E.E., Franco, B., Giles, R.H., Ueffing, M., Russell, R.B., Roepman, R., 2016. An
1229   organelle-specific protein landscape identifies novel diseases and molecular mechanisms.
1230   Nat. Commun. 7, 11491. https://doi.org/10.1038/ncomms11491

1232   van Dam, T.J.P., Kennedy, J., van der Lee, R., de Vrieze, E., Wunderlich, K.A., Rix, S.,
1233   Dougherty, G.W., Lambacher, N.J., Li, C., Jensen, V.L., Leroux, M.R., Hjeij, R., Horn,
1234   N., Texier, Y., Wissinger, Y., van Reeuwijk, J., Wheway, G., Knapp, B., Scheel, J.F.,
1235   Franco, B., Mans, D.A., van Wijk, E., Képès, F., Slaats, G.G., Toedt, G., Kremer, H.,
1236   Omran, H., Szymanska, K., Koutroumpas, K., Ueffing, M., Nguyen, T.-M.T., Letteboer,
1237   S.J.F., Oud, M.M., van Beersum, S.E.C., Schmidts, M., Beales, P.L., Lu, Q., Giles, R.H.,
1238   Szklarczyk, R., Russell, R.B., Gibson, T.J., Johnson, C.A., Blacque, O.E., Wolfrum, U.,
1239   Boldt, K., Roepman, R., Hernandez-Hernandez, V., Huynen, M.A., 2019. CiliaCarta: An
1240   integrated and validated compendium of ciliary genes. PLOS ONE 14, e0216705.
1241   https://doi.org/10.1371/journal.pone.0216705

1243   Vasquez SSV, van Dam J, Wheway G. 2021. An updated SYSCILIA gold standard (SCGSv2) of
1244   known ciliary genes, revealing the vast progress that has been made in the cilia research
1245   field. Mol Biol Cell. 2021 Dec 1;32(22):br13. doi: 10.1091/mbc.E21-05-0226.

1247   Wei, Q., Xu, Q., Zhang, Y., Li, Y., Zhang, Q., Hu, Z., Harris, P.C., Torres, V.E., Ling, K., Hu, J.,
1248   2013. Transition fibre protein FBF1 is required for the ciliary entry of assembled
1249   intraflagellar transport complexes. Nat. Commun. 4, 2750.
1250   https://doi.org/10.1038/ncomms3750

1252   Wei, Q., Zhang, Y., Li, Y., Zhang, Q., Ling, K., Hu, J., 2012. The BBSome controls IFT
1253   assembly and turnaround in cilia. Nat. Cell Biol. 14, 950–957.
1254   https://doi.org/10.1038/ncb2560

1256   Wheway, G., Genomics England Research Consortium, Mitchison, H.M., 2019. Opportunities
1257   and Challenges for Molecular Understanding of Ciliopathies–The 100,000 Genomes
1258   Project. Front. Genet. 10, 127. https://doi.org/10.3389/fgene.2019.00127

1260   Williams, C.L., Li, C., Kida, K., Inglis, P.N., Mohan, S., Semenec, L., Bialas, N.J., Stupay,

1261   R.M., Chen, N., Blacque, O.E., Yoder, B.K., Leroux, M.R., 2011. MKS and NPHP
1262   modules cooperate to establish basal body/transition zone membrane associations and
1263   ciliary gate function during ciliogenesis. J. Cell Biol. 192, 1023–1041.
1264   https://doi.org/10.1083/jcb.201012116
1265
1266   Williams, C.L., McIntyre, J.C., Norris, S.R., Jenkins, P.M., Zhang, L., Pei, Q., Verhey, K.,
1267   Martens, J.R., 2014. Direct evidence for BBSome-associated intraflagellar transport
1268   reveals distinct properties of native mammalian cilia. Nat. Commun. 5, 5813.
1269   https://doi.org/10.1038/ncomms6813
1270
1271   Wright, K.J., Baye, L.M., Olivier-Mason, A., Mukhopadhyay, S., Sang, L., Kwong, M., Wang,
1272   W., Pretorius, P.R., Sheffield, V.C., Sengupta, P., Slusarski, D.C., Jackson, P.K., 2011.
1273   An ARL3-UNC119-RP2 GTPase cycle targets myristoylated NPHP3 to the primary
1274   cilium. Genes Dev. 25, 2347–2360. https://doi.org/10.1101/gad.173443.111
1275
1276   Xu, Q., Zhang, Y., Wei, Q., Huang, Y., Li, Y., Ling, K., Hu, J., 2015. BBS4 and BBS5 show
1277   functional redundancy in the BBSome to regulate the degradative sorting of ciliary
1278   sensory receptors. Sci. Rep. 5, 11855. https://doi.org/10.1038/srep11855
1279
1280   Xu, W., Jin, M., Hu, R., Wang, H., Zhang, F., Yuan, S., Cao, Y., 2017. The Joubert Syndrome
1281   Protein Inpp5e Controls Ciliogenesis by Regulating Phosphoinositides at the Apical
1282   Membrane. J. Am. Soc. Nephrol. 28, 118–129. https://doi.org/10.1681/ASN.2015080906
1283
1284   Ye, F., Nager, A.R., Nachury, M.V., 2018. BBSome trains remove activated GPCRs from cilia
1285   by enabling passage through the transition zone. J. Cell Biol. 217, 1847–1868.
1286   https://doi.org/10.1083/jcb.201709041