1

Loss of Elp1 disrupts trigeminal ganglion neurodevelopment in a model of Familial Dysautonomia

Carrie E. Leonard[1], Frances Lefcort[2], Lisa A. Taneyhill[1]*

[1] Department of Avian and Animal Sciences, University of Maryland, College Park, MD 20742 USA

[2] Department of Cell Biology and Neuroscience, Montana State University, Bozeman, MT 59717 USA

Email:
ltaney@umd.edu
leonardc@umd.edu
flefcort@msu.edu

*Corresponding Author

Abstract

Familial Dysautonomia (FD) is a sensory and autonomic neuropathy caused by a mutation in *Elongator complex protein 1* (*ELP1*). FD patients have small trigeminal nerves and impaired perception of facial pain and temperature. These signals are relayed by nociceptive neurons in the trigeminal ganglion, a structure comprised of both neural crest- and placode-derived cells. Mice lacking *Elp1* in neural crest derivatives ("*Elp1 CKO*") are born with smaller trigeminal ganglia, suggesting Elp1 is important for trigeminal ganglion development, yet the function of Elp1 in this context is unknown. We demonstrate Elp1 expression in both neural crest- and placode-derived trigeminal neurons, which our data suggest give rise to primarily TrkA- and TrkB/C-expressing neurons, respectively. While Elp1 is not required for initial trigeminal ganglion formation, *Elp1 CKO* trigeminal neurons exhibit abnormal axon outgrowth and decreased target innervation. Developing nociceptors that express the receptor TrkA are especially vulnerable to Elp1 loss. TrkA expression is decreased in *Elp1 CKO* trigeminal nerve endings, coinciding with increased cell death. Subsequently, fewer TrkA neurons are present in the *Elp1 CKO* trigeminal ganglion, indicating Elp1 supports the target innervation and survival of trigeminal nociceptors. These findings explain the loss of facial pain and temperature sensation in FD.

Introduction

Hereditary sensory and autonomic neuropathies (HSANs) are a group of phenotypically similar, yet distinct, peripheral nervous system disorders that stem from unique mutations (Schwartzlow and Kazamel 2019). The most prevalent form is HSAN Type III, or Familial Dysautonomia (FD), which arises from an intronic mutation in the *Elongator Complex Protein 1 (ELP1)* gene (Anderson et al. 2001; Slaugenhaupt et al. 2001). This mutation causes tissue-specific mis-splicing of *ELP1*, subsequently reducing ELP1 protein in neurons (Slaugenhaupt et al. 2001; Hims et al. 2007). Elp1 is a requisite scaffolding protein of the six-subunit Elongator complex that functions in translation by modifying particular tRNAs (B. Huang, Johansson, and Byström 2005; B. Huang, Jian, and Byström 2008; Karlsborn, Tükenmez, Mahmud, et al. 2014; Xu et al. 2015; Esberg et al. 2006). Deletion of Elongator subunits has been shown to alter tRNA modifications and protein expression in several systems (C. Chen, Tuck, and Byström 2009; Esberg et al. 2006; Karlsborn, Tükenmez, Chen, et al. 2014; Goffena et al. 2018; Kojic et al. 2021). Consequently, loss or reduction of Elp1 protein leads directly and indirectly to the clinical phenotypes seen in FD, which include impaired pain and temperature sensation, feeding and swallowing difficulties, blood pressure instability, tachycardia, optic neuropathy, and gastrointestinal dysfunction, among other symptoms. The disease is fatal, with the majority of patients dying by age 50 (Gold-von Simson and Axelrod 2006). Autopsy reports and studies conducted in patients and animal models demonstrate FD phenotypes are caused by abnormal neurodevelopment, in addition to neurodegeneration across the lifespan of the affected individual (Pearson et al. 1978;

Hunnicutt et al. 2012; George et al. 2013; Jackson et al. 2014; Li, Gruner, and Tourellotte

2020; Won et al. 2019; Lefcort et al. 2017; Dietrich and Dragatsis 2016).

FD severely impacts the peripheral nervous system, composed of sensory and

autonomic neurons that are mostly derived from neural crest cells and some cranial

sensory neurons that arise from ectodermal placodes (Méndez-Maldonado et al. 2020;

Steventon, Mayor, and Streit 2014; Kameneva and Adameyko 2019). Since mice that are

null for *Elp1* are embryonic lethal (Y.-T. Chen et al. 2009; Dietrich et al. 2011), FD has

been modeled in mice using conditional knockout of *Elp1* in neural crest cells

(*Elp1fl/fl;Wnt1-Cre+*, abbreviated "*Elp1 CKO*"), which recapitulates several aspects of the

human disease, including significant loss of sensory and autonomic neurons (George et

al. 2013; Jackson et al. 2014). To date, these studies have focused on mechanisms of

Elp1 in sensory and sympathetic neurons in the trunk. However, FD patients also

experience cranial sensory deficits such as impaired sensation of facial pain and

temperature, neurogenic dysphagia, absent corneal reflexes, and reduced basal

lacrimation (Mendoza-Santiesteban et al. 2017; Palma, et al 2008; Barlow 2009; Geltzer

et al. 1964; Gutiérrez, Norcliffe-Kaufmann, and Kaufmann 2015; Palma et al. 2018).

These functions rely on input from the trigeminal nerve, the largest of the cranial nerves,

which contains axons of sensory neurons that reside in the trigeminal ganglion. A

quantitative MRI study revealed FD patients have smaller trigeminal nerves compared to

healthy, age-matched controls, but there are no indications of progressive trigeminal

nerve degeneration (Won et al. 2019). Moreover, *Elp1 CKO* mice are born with smaller

trigeminal ganglia compared to controls (Jackson et al. 2014). Together, these findings

suggest Elp1 may play an important role in the development of trigeminal sensory neurons, a critical aspect of FD that has yet to be explored.

While peripheral neurons in the trunk are exclusively neural crest-derived, cranial sensory neurons arise from two progenitor populations, neural crest cells and ectodermal placodes, which typically contribute neurons to spatially distinct regions of the cranial ganglia (York, Yuan, and McCauley 2020; Moody and Lamantia 2015; Park and Saint-Jeannet 2010; Steventon, Mayor, and Streit 2014). The trigeminal ganglion is unique in that it contains intermixed neural crest- and placode-derived neurons that rely on one another for proper migration, coalescence, and function during development (Blentic et al. 2011; D'amico-Martel 1982; D'Amico-Martel and Noden 1983; Hamburger 1961; Saint-Jeannet and Moody 2014; Shiau et al. 2008; Steventon, Mayor, and Streit 2014). Shortly after differentiation, trigeminal ganglion neuron subtypes are discernable by mutually exclusive expression of Tropomyosin receptor kinase (Trk) receptors, TrkA, TrkB, or TrkC, which are required for target innervation and long-term survival (E. J. Huang, Wilkinson, et al. 1999; Wilkinson et al. 1996; Scott-Solomon and Kuruvilla 2018; A. M. Davies 1997; Reichardt 2006; E. J. Huang, Zang, et al. 1999). Importantly, Trk expression generally correlates with the ultimate sensory modality encoded by a particular neuron; for example, small-diameter TrkA neurons are typically associated with pain and temperature perception (Mu et al. 1993; Genç, Ulupinar, and Erzurumlu 2005; d'Amico-Martel and Noden 1980; D'Amico-Martel and Noden 1983; A. Davies and Lumsden 1984). In dorsal root (sensory) and sympathetic (autonomic) ganglia, Elp1 is required for the generation and/or survival of TrkA and TrkB neurons, while TrkC neurons are spared during development (Jackson et al. 2014; George et al. 2013). In these contexts, neuronal

loss has been attributed to decreased neurogenesis resulting from early differentiation

and apoptosis of progenitors (George et al. 2013; Goffena et al. 2018), and to severe

target innervation defects and neuronal apoptosis due to insufficient neurotrophic support

(Jackson et al. 2014; Li, Gruner, and Tourtellotte 2020). Notably, other less common

forms of HSANs arise from germline mutations in the genes encoding either *TrkA* or its

high-affinity ligand, *nerve growth factor* (*NGF*) (Schwartzlow and Kazamel 2019). A

growing body of evidence from animal models suggests Elp1 regulates TrkA signaling as

well (Abashidze et al. 2014; Lefler et al. 2015; Naftelberg et al. 2016; Li, Gruner, and

Tourtellotte 2020), although the relationship between Elp1 and TrkA has not been

investigated in the cranial ganglia. While there are clear trigeminal sensory deficits in FD,

the function of Elp1 in the trigeminal ganglion and its nerves remains unexamined.

Moreover, it is unknown whether certain neuronal subtypes within the trigeminal ganglion

are more vulnerable to Elp1 loss than others, or whether trigeminal sensory phenotypes

in FD patients arise from defects in neural crest-derived neurons, placodal neurons, or

both.

Here, we describe the first comprehensive analysis of neurodevelopmental

changes in the trigeminal ganglion using an established *Elp1 CKO* mouse model of FD,

where *Elp1* is deleted from neural crest cells and their derivatives (George et al. 2013).

We demonstrate that Elp1 protein is enriched in the cytoplasm of differentiated neural

crest- and placode-derived neurons in the trigeminal ganglion. While initial formation of

the trigeminal ganglion is unaltered in *Elp1 CKO* mice, axon outgrowth, nerve formation,

and nerve branching are severely diminished as development proceeds. These changes

are accompanied by a drastic decrease in TrkA levels in the peripheral and central

projections of trigeminal ganglion neurons as well as a significant reduction in the number

of TrkA neurons within the ganglion. Finally, we demonstrate this loss of TrkA neurons

due to neural crest-specific deletion of *Elp1* is correlated with increased cell death.

Collectively, these findings indicate Elp1 is required for proper target innervation and

survival of TrkA neurons in the trigeminal ganglion. Our results complement previous

reports that demonstrate Elp1 modulates TrkA signaling, yet we now provide additional

insight into nuanced phenotypes that may arise due to differences between nerve

branches of the same ganglion. Moreover, by using a neural crest-specific knockout in

the trigeminal ganglion, which contains neurons of both neural crest and placodal origin,

we shed light on the neurodevelopmental dynamics and lineage of specific neuronal

subpopulations in the trigeminal ganglion. Importantly, our findings introduce a cellular

mechanism to explain the developmental defects occurring in the trigeminal ganglion,

which ultimately contribute to the etiology of facial sensory deficits experienced by

patients with FD.

148   Results

149   Elp1 protein is enriched in the cytoplasm of trigeminal ganglion sensory neurons

150   While ubiquitous Elp1 expression in rodent embryos has been previously reported

151   (Mezey et al. 2003; George et al. 2013), the spatio-temporal distribution of Elp1 in the

152   craniofacial complex has not been evaluated. To initially address this, we examined the

153   *Elp1:LacZ* (previously "*Ikbkap:LacZ*") reporter mouse that expresses β-galactosidase

154   from the *Elp1* locus (George et al. 2013). Whole-mount preparations revealed widespread

Figure 1. Elp1 protein is enriched in the cytoplasm of developing trigeminal ganglion neurons. (**A**) Lateral view of E10.5 *Elp1:LacZ* reporter mouse stained for β-galactosidase. Dashed line indicates the plane of section for the same embryo shown in **B**. (**B-C**) Representative horizontal section through embryo in **A** to reveal *Elp1* gene expression. Boxed region in **B** is magnified and shown in **C**. (**D-P**) Representative horizontal sections taken from wildtype E10.5 (**D-G**), E11.5 (**H-M**), or E12.5 (**N-P**) mouse embryos followed by fluorescent immunohistochemistry for Elp1 (**D-P**, green), Islet1 (**D**, **F**, **H**, **K**, "Isl1", purple), Neuropilin2 (**E**, "Nrp2", red), β-tubulin III (**I**, **L**, "Tubb3", red), and Sox10 (**N**, **P**, red). Boxed region in **D** and **E** is magnified in **F-G** and shows Elp1 (green), Isl1 (purple), and DAPI-stained nuclei (blue). (**J-M**) Higher magnification of box in **H** and **I**. (**O-P**) Higher magnification of box in **N**. Carets indicate Isl1-positive and/or Tubb3-positive neuronal cell bodies (**F**, **G**, **J-M**). Arrows identify axons (**J**, **L**, **M**, **O**, **P**), while arrowheads point to Sox10-positive glial progenitors (**O-P**). Abbreviations: e – eye, FN – frontonasal prominence, NT – neural tube, ov – otic vesicle, PA – pharyngeal arch, TG – trigeminal ganglion. Scale bars: 400µm (**A**), also applies to **B**; 50µm (**C**), applies to **D, E, H, I**; 100µm (**N**); 10µm (**F**), applies to **G**; 10µm (**J**), applies to **K, L, M, O, P**.

β-galactosidase staining in the head at embryonic day 10.5 (E10.5), with prominent

expression in the neural tube, regions of the developing cranial ganglia, the pharyngeal

arches, and the facial prominences (Fig. 1A). Horizontal sections through the head at

E10.5 revealed diffuse β-galactosidase expression in the trigeminal ganglion, which, at

this stage, is a newly condensed structure containing differentiated trigeminal placode-

derived neurons and undifferentiated cranial neural crest cells (Fig. 1B, C, Karpinski et al.

2016). β-galactosidase expression was also apparent in the neural tube, the eye, and

scattered throughout the cranial mesenchyme (Fig. 1B, C). Together, these results

indicate the *Elp1* gene is expressed in cranial neural tissues, similar to findings in the

trunk, and more specifically that *Elp1* is expressed in the newly formed trigeminal

ganglion.

Interestingly, detection of Elp1 via immunohistochemistry revealed an overlapping,

but more specific, expression pattern. At E10.5, Elp1 protein was abundant in the

cytoplasm of differentiated neurons identified by expression of the transcription factor

Islet1, and was not detected in the nucleus (Fig. 1D-G and Fig. 1–figure supplement 1A-

D). Since the vast majority of neurons in the trigeminal ganglion are placode-derived at

this stage, while surrounding neural crest cells have yet to differentiate (Karpinski et al.

2016), we conclude that Elp1 is initially expressed in trigeminal placode-derived neurons.

In contrast, condensed neural crest cells within the trigeminal ganglion, identified by

expression of the surface receptor Neuropilin2 or the transcription factor Pax3, and devoid

Figure 1–figure supplement 1. Elp1 is not expressed in Pax3-positive neural crest cells or glial progenitors in the developing trigeminal ganglion. (**A-H**) Fluorescent immunohistochemistry on representative horizontal sections from E10.5 (**A-D**) and E11.5 (**E-H**) wildtype mouse embryos demonstrating expression of Elp1 (**A-H**, green), Isl1 (**A-C**, **E**, **G**, purple) and Pax3 (**B**, **D**, **F**, **H**, red). Carets point to Isl1-positive neuron cell bodies (**C**, **D**, **G**, **H**). Arrows indicate axons (**G**, **H**), while arrowheads denote Pax3-positive neural crest cells (**C**, **D**) or glial progenitors (**G**, **H**). Scale bars: 50μm (**A**), applies to **B, E, F**; 20μm (**C**), applies to **D, G, H**.

of Islet1, expressed little to no Elp1 protein (Fig. 1E and Fig.1–figure supplement 1A-D).

Compared to the diffuse *Elp1:LacZ* signal in the trigeminal ganglion at the same stage

(Fig. 1A-C), this suggests *Elp1* may be post-transcriptionally regulated, leading to robust

Elp1 protein expression in neurons upon differentiation.

At E11.5, when neural crest-derived neurons are now present in the trigeminal

ganglion, Elp1 protein was observed in all differentiated neurons expressing β-tubulin III

(Tubb3) and Islet1, irrespective of their developmental origin, with the highest signal

detected in axons (Fig. 1H-M and Fig.1–figure supplement 1E-H). These findings are

similar to previously reported observations in other peripheral ganglia (Abashidze et al.

2014; Hunnicutt et al. 2012; George et al. 2013). By E12.5, enrichment of Elp1 protein in

the cytoplasm, especially axons, of Tubb3-positive neurons is still apparent, while Elp1 is

absent or expressed at relatively low levels in neural crest-derived Sox10-positive glia

within the ganglion and along the embryonic nerves (Fig. 1K-M). Collectively, these

results suggest important functions for Elp1 in developing sensory neurons of the

trigeminal ganglia.

Neural crest-specific deletion of *Elp1* causes progressive morphological

abnormalities within the trigeminal ganglion and its nerve branches

To understand the requirements for Elp1 during trigeminal ganglion development

and to gain insight into the etiology of FD phenotypes associated with trigeminal ganglion

dysfunction, an established mouse model was used, in which *Elp1* is conditionally deleted

from neural crest cells and their derivatives via Cre-mediated recombination under the

Wnt1 promoter (*Elp* *flox/flox*; *Wnt1-cre*+, "*Elp1 CKO*") (George et al. 2013). The anatomy

and morphology of the embryonic trigeminal ganglion and associated nerves of *Elp1 CKO*

and littermate controls were evaluated over developmental time. The normal progression

of trigeminal nerve development between E10.5 and E13.5 is depicted in Figure 2, while

Elp1 CKO trigeminal ganglion and nerve phenotypes relative to controls are summarized

in Table 1.

Figure 2. Schematic of normal mouse trigeminal ganglion and nerve anatomy over developmental time. (**A-D**) Diagram labels relevant ganglion and nerve anatomy in E10.5 (**A**), E11.5 (**B**), E12.5 (**C**), and E13 (**D**) mouse embryos for reference in **Figures 3-6**. Abbreviations: CR – central root, Fr – frontal nerve, Io – infraorbital nerve, MnV – mandibular nerve, MxV – maxillary nerve, Na – nasal nerve, NT – neural tube, OpV – ophthalmic nerve, TG – trigeminal ganglion.

When compared to littermate controls at E10.5, initial formation of the trigeminal

ganglion appeared normal in *Elp1 CKO*, as demonstrated by whole-mount Tubb3

immunohistochemistry, which labels all neurons (Fig. 3A, B). Gross anatomy was intact,

including a condensed ganglion and the emergence of the ophthalmic, maxillary, and

mandibular nerve branches (Fig. 3A, B). Immunohistochemistry on E10.5 tissue sections

revealed both Tubb3-positive placodal neurons and Sox10-positive undifferentiated

neural crest cells were present and appropriately distributed throughout the forming

ganglion (Fig. 3C, D). Altogether, these data suggest Elp1 is not required for cranial neural

crest cells to migrate to the trigeminal anlage and coalesce with placode-derived neurons

for ganglion assembly.

Table 1. Comparison of *Elp1 CKO* Trigeminal Ganglion and Nerve Phenotypes Relative to Control Across Developmental Time. Summary of observations of the trigeminal ganglion, ophthalmic nerve, maxillary nerve, mandibular nerve, and central nerve root in *Elp1 CKO* between E10.5 and E13.5, as compared to control littermates. *, mandibular nerve was difficult to visualize at these stages, so no observations were recorded.

	Gestational Age			
	E10.5	**E11.5**	**E12.5**	**E13-13.5**
Ganglion	No difference; Tubb3-positive placodal neurons and Sox10-positive neural crest cells are present/ distributed throughout forming ganglion	Slightly less compact	Less compact; increased cell death	Less compact; TrkA expression present but reduced; fewer TrkA-positive neurons present
Ophthalmic N.	Present; no difference in length	No difference	Axons straying from established fascicles; decreased branching complexity; nasal nerve absent	Decreased nerve complexity even more apparent; "dying back" of axons; nasal nerve still absent; reduction in TrkA expression
Maxillary N.	Present; no difference in length	Disorganized; many axons stray from established fascicles as they traverse toward the whisker pad	No change in whisker pad innervation	Decreased innervation of whisker pad; reduction in TrkA expression
Mandibular N.	Present; no difference in length	Disorganized and less compact; "hairy" appearance due to straying axons	*	*
Central root axons	No difference	Less compact	Smaller	Smaller; reduction in TrkA expression

Figure 3. Neural crest cell migration and initial trigeminal ganglion formation appear normal in *Elp1 CKO* at E10.5. (**A-B**) Lateral view of the trigeminal ganglion of Control (**A**, N = 3) and *Elp1 CKO* (**B**, N = 3) after Tubb3 whole-mount immunohistochemistry (white). (**C-D**) Fluorescent immunohistochemistry on representative sections within the trigeminal ganglion from Control (**C**, N = 2) or *Elp1 CKO* (**D**, N = 2) shows placodal neurons (Tubb3, red) and neural crest cells (Sox10, green). Scale bars: 400µm (**A**), also applies to **B**; 20µm (**C**), applies to **D**. Abbreviations: MnV – mandibular nerve, MxV – maxillary nerve, OpV – ophthalmic nerve, TG – trigeminal ganglion.

214   To capture the neurodevelopmental dynamics occurring in the trigeminal ganglion

215 in control versus *Elp1 CKO*, we visualized intact trigeminal ganglia and associated nerves

216 at intervals representing early (E11.5), middle (E12.5), and late (E13) stages of neural

217 crest neurogenesis, using whole-mount Tubb3 immunohistochemistry. By E11.5, control

218 and *Elp1 CKO* displayed robust outgrowth of axons from the trigeminal ganglion, which

219 contains both neural crest- and placode-derived neurons at this stage (Fig. 4A, D).

220 Although the ophthalmic, maxillary, and mandibular nerves were distinct in control and

221 *Elp1 CKO*, axons in *Elp1 CKO* nerves appeared less organized. In the maxillary nerve,

222 *Elp1 CKO* axons were observed straying from established axon bundles into the

223 surrounding mesenchyme or crossing to adjacent fascicles (Fig. 4B, E). Axons of the

224 mandibular nerve in *Elp1 CKO* also traveled away from the established nerve without

225 direction, compared to the compact mandibular nerve observed in controls (Fig. 4C, F).

226 By E12.5, all subdivisions of the ophthalmic nerve were severely deformed in *Elp1 CKO*,

227 yet there were nuanced changes depending on the specific nerve (Fig. 5A, D). For

Figure 4. Progressive trigeminal nerve abnormalities arise in *Elp1 CKO* starting at E11.5. (**A-F**) Maximum intensity projections of confocal Z-stacks through Control (**A-C**, N = 3) or *Elp1 CKO* (**D-F**, N = 4), which were processed for whole-mount immunohistochemistry to detect Tubb3 (white), followed by tissue clearing. (**B, C, E, F**) Higher magnification images of boxed regions shown in **A** and **D**. Arrows indicate less compact central root (**D**) and disorganized axons (**E, F**) in *Elp1 CKO*. Scale bar: 200µm (**A**), applies to **B**; also applies to **B, C, E, F** as 50µm. Abbreviations: MxV – maxillary nerve, MnV – mandibular nerve, OpV – ophthalmic nerve.

example, the developing frontal nerve was present in both control and *Elp1 CKO*;

however, the branching complexity of the nerve was decreased in *Elp1 CKO* (Fig. 5B, E),

as quantified by a modified Sholl analysis (Fig. 5G, H, Sholl 1953). Proximal to the

ganglion, stray axons were also observed growing away from established ophthalmic

branches (Fig. 5E). While the rostral extension of the ophthalmic nerve was present in

both control and *Elp1 CKO*, the medial and lateral nasal nerves that extend from this

branch were only established in control embryos at this stage (Fig. 5C, F). Interestingly,

though there was no obvious difference in the size of the central nerve root that contains

axons projecting to the brainstem at E11.5 (Fig. 4A, D), the nerve root was smaller in *Elp1*

CKO by E12.5, indicating the loss of some central axons during this developmental time

frame (Fig. 5A, D).

Figure 5. Trigeminal nerve branches are less complex or absent in *Elp1 CKO* at E12.5. (**A-F**) Maximum intensity projections of confocal Z-stacks through Control (**A-C**, N = 2) or *Elp1 CKO* (**D-F**, N = 4), which were processed for whole-mount immunohistochemistry to detect Tubb3 (white), followed by tissue clearing. (**B**, **C**, **E**, **F**) Higher magnification images of the frontal nerve division of the ophthalmic trigeminal branch around the eye (**B**, **E**) or the infraorbital nerve division of the maxillary branch at the developing whisker pad (**C**, **F**). Arrows indicate small central root and branch of ophthalmic nerve (**D**), disorganized axons (**E**), and the absence of the nasal nerve (ophthalmic division, **F**) in *Elp1 CKO*. (**G**) Diagram explaining Modified Sholl analysis, with concentric circles of increasing radii overlayed on representative traces (green) of the frontal nerve in Control (left) or *Elp1 CKO* (right). (**H**) Modified Sholl Analysis to quantify complexity of frontal nerve. Individual distributions are plotted in light blue (Control, N = 2) and light orange (*Elp1 CKO*, N = 4), while group averages are plotted in dark blue (Control) and dark orange (*Elp1 CKO*). Scale bar: 200μm (**A**), applies to **D**; 200μm (**B**), applies to **C, E, F**. Abbreviations: Fr - frontal nerve, Io – infraorbital nerve, MxV – maxillary nerve, OpV – ophthalmic nerve, TG – trigeminal ganglion.

Toward the end of the trigeminal ganglion neurogenic period, deficits in nerve

growth and branching were even more pronounced in *Elp1 CKO*. The frontal nerve

(ophthalmic) in *Elp1 CKO* was less complex at E13 than at E12.5, suggesting retraction

and/or dying of some *Elp1 CKO* ophthalmic axons after failure to properly innervate the

eye region (Fig. 6A, B). The lateral and medial nasal nerves (ophthalmic) were still absent

in *Elp1 CKO* yet clearly established in controls (Fig. 6C, D). That the nasal nerves were

not observed in *Elp1 CKOs* at any stage indicates some ophthalmic axons never reached

their distal target site. Additionally, the infraorbital nerve (maxillary), which innervates the

whisker pad, was less elaborate in *Elp1 CKO* compared to controls at E13 (Fig. 6C, D).

Together, these results provide a dynamic summary of the requirements for Elp1 in

axonal pathfinding and target innervation in trigeminal ganglion neurons. Importantly, our

data suggest that nerve defects caused by loss of Elp1 can differ, even within the same

ganglion, based on neuronal identity and/or specific target region.

Decreased TrkA correlates with increased cell death and loss of nociceptive

neurons in *Elp1 CKO* trigeminal ganglia

Given the finding that trigeminal ganglion target innervation is disrupted in *Elp1 CKO*,

combined with published studies from sympathetic neurons suggesting Elp1 is required

for TrkA signaling (Li, Gruner, and Tourtellotte 2020; Jackson et al. 2014; Naftelberg et

al. 2016), the distribution of TrkA expression was also examined via whole-mount

immunohistochemistry at E13. In control embryos, TrkA signal was strong throughout the

trigeminal ganglion and along all three major nerve divisions (Fig. 6E, G). In the frontal

(ophthalmic) and infraorbital (maxillary) nerves, TrkA was expressed ubiquitously,

Figure 6. TrkA expression is decreased with persistent innervation defects in *Elp1 CKO* at E13. (**A-H**) Maximum intensity projections of confocal Z-stacks through Control (**A, C, E, G**, N = 2) or *Elp1 CKO* (**B, D, F, H**, N = 3), which were processed for whole-mount immunohistochemistry to detect Tubb3 (**A-D**) and TrkA (**E-H**), followed by tissue clearing. Arrows indicate regions where nerves are absent or severely diminished in *Elp1 CKO* (**B**, **D**), while arrowheads point to areas with Tubb3-positive nerves but undetectable TrkA expression (**B**, **D**, **F**, **H**). Scale bar: 200µm (**A**) and applies to all images. Abbreviations: Fr - frontal nerve, Io – infraorbital nerve, MxV – maxillary nerve, OpV – ophthalmic nerve, TG – trigeminal ganglion.

including in the smallest, most distal branches (Fig. 6E, G). In *Elp1 CKO*, TrkA was

detected, but weaker, in the trigeminal ganglion and in the core of the ophthalmic and

maxillary branches (Fig. 6F). Strikingly, TrkA signal was drastically reduced in more distal

nerve branches at E13, including the frontal nerve endings, the rostral ophthalmic

projection, and the infraorbital axons of the whisker pad (Fig. 6F, H). Moreover, there was

significant loss of TrkA in the central root and brainstem axon tracts, indicating effects on

both central and peripheral projections at this stage (Figure 6-figure supplement 1).

Therefore, target innervation defects in *Elp1 CKO* trigeminal ganglion neurons are

associated with a loss of TrkA in the ganglion and nerves. These results are particularly

intriguing given that TrkA neurons typically function in nociception, and FD patients

experience a reduction or loss in the ability to sense facial pain and temperature.

Figure 6–figure supplement 1. TrkA expression is decreased in the central nerve root of the trigeminal ganglion in *Elp1 CKO* at E13. (**A-D**) Maximum intensity projections of confocal Z-stacks through Control (**A, C,** N = 2) or *Elp1 CKO* (**B, D**, N = 3), which were processed for whole-mount immunohistochemistry to detect Tubb3 (**A-B**) and TrkA (**C-D**), followed by tissue clearing. Arrows point to central root in *Elp1 CKO* (**B, D**). Scale bar: 200μm (**A**) and applies to all images. Abbreviations: Fr - frontal nerve, Io – infraorbital nerve, MnV – mandibular nerve, MxV – maxillary nerve, OpV – ophthalmic nerve.

Next, we determined whether TrkA neurons were devoid of Elp1 protein in *Elp1*

CKO trigeminal ganglia. To this end, co-labeling for Elp1 and TrkA was performed via

immunohistochemistry on control and *Elp1 CKO* tissue at E12.5. In control trigeminal

ganglia, Elp1 was present in all neurons, and therefore co-labeled TrkA neurons,

especially in axons (Fig. 7-figure supplement 1). In *Elp1 CKO* trigeminal ganglia, most

TrkA neurons and axons did not express Elp1 (Fig. 7-figure supplement 1), while the

remaining Elp1-positive neurons had large cell bodies characteristic of TrkB/C neurons

(d'Amico-Martel and Noden 1980; D'Amico-Martel and Noden 1983; E. J. Huang, Zang,

et al. 1999). To evaluate whether the loss of TrkA could be explained by neuronal cell

death, TUNEL assays were also performed at E12.5, with co-labeling for TrkA. Compared

to control embryos, which had occasional, scattered TUNEL-positive cells at this stage

(Fig. 7A-C), *Elp1 CKO* had a dramatic increase in the number of TUNEL-positive cells

distributed throughout the entire trigeminal ganglion, many of which also expressed TrkA

(Fig. 7D-I). To determine whether this increased cell death resulted in the loss of specific

Figure 7–figure supplement 1. Loss of Elp1 from TrkA neurons in *Elp1 CKO* trigeminal ganglia. (**A-F**) Fluorescent immunohistochemistry on representative horizontal sections from E12.5 Control (**A-C**, N = 4) or *Elp1 CKO* (**D-F**, N = 4) mouse embryos demonstrating expression of Elp1 (red) and TrkA (green). Arrows point to TrkA-positive axons. Scale bar: 20μm (**A**) and applies to all images.

neuronal subtypes, TrkA, TrkB, and TrkC neuron cell bodies were quantified in E13.5

trigeminal ganglia. While no differences were observed in the number of TrkB or TrkC

neurons at this stage, TrkA neurons were reduced by 23 percent in *Elp1 CKO* trigeminal

ganglia compared to controls (Fig. 7J, Control N=4; *Elp1 CKO* N=4; p = 0.016, unpaired

t-test corrected for multiple comparisons with Holm-Sidak method). Collectively, these

findings suggest that deletion of Elp1 from the neural crest lineage leads to detrimental

innervation defects in the head, and the selective loss of TrkA neurons, as a result of

widespread cell death within the trigeminal ganglion.

***Elp1 CKO* provides evidence for the developmental origins of Trk-expressing neuronal subtypes in the trigeminal ganglion**

Given the dual embryonic origin of the trigeminal ganglion, and our conditional

knockout of *Elp1* from neural crest but not placode-derived tissues, we inquired about the

ontogeny of Trk-expressing subpopulations to ask whether our findings could be due to

Figure 7. Loss of Elp1 from neural crest derivatives results in increased cell death and loss of TrkA neurons in the trigeminal ganglia. (**A-I**) Fluorescent immunohistochemistry on representative horizontal sections from E12.5 Control (**A-C**, N = 3) or *Elp1 CKO* (**D-I**, N = 3) revealing expression of TrkA (purple) with TUNEL staining (green). (**G-I**) Higher magnification of boxed regions in **D-F**. Arrowheads point to TrkA neurons that are TUNEL-positive (**G-I**). (**J**) Quantification of TrkA-, TrkB-, and TrkC-expressing neurons at E13.5 in Control (blue, N = 3) and *Elp1 CKO* (orange, N = 3) trigeminal ganglia. Values are mean ± SEM. *p = 0.016, unpaired t-test adjusted for multiple comparisons using Holm-Sidak method. Scale bars: 50μm (**A**), applies to **B-F**; 20μm (**G**), applies to **H-I**.

interactions between neural
crest and placode cells during
early trigeminal ganglion
development. First, we first explored the dynamics of normal TrkA neurogenesis in the
trigeminal ganglion. Consistent with reports that TrkB and TrkC neurons are born first in
the trigeminal ganglion (E9.5-11) followed by TrkA neurons (E11.5-13.5) (E. J. Huang,
Wilkinson, et al. 1999; E. J. Huang, Zang, et al. 1999), immunohistochemistry on
trigeminal ganglion sections revealed that all three Trk subtypes are present at E11.5
(Fig. 8A-F). At this stage, TrkC neurons comprise the largest proportion of neurons,
followed closely by TrkA, with TrkB neurons representing the smallest proportion of

trigeminal ganglion neurons (Fig. 8A-C). In accordance with previous findings (E. J.

Huang, Zang, et al. 1999), the number of TrkB and TrkC neurons did not appear to

increase between E11.5 and E12.5, but there was a robust increase in the number of

TrkA neurons, such that TrkA neurons became the vast majority of trigeminal ganglion

neurons by E12.5 (Fig. 8G-I). Together with our findings that Elp1 deletion from neural

crest specifically targets TrkA neurons, and the fact that placode-derived neurons

differentiate earlier than neural crest-derived neurons, we deduced that TrkA neurons in the

Figure 8. Dynamic expression of Six1 and Trk receptors occurs during trigeminal ganglion neurogenesis.

(**A-O**) Fluorescent immunohistochemistry on representative horizontal sections from E11.5 (**A-F**) and E12.5 (**G-O**) Control (**A-L**, N=2 each) or *Elp1 CKO* (**M-O**, N=2) mouse embryos demonstrating expression of TrkA (**A, D, G, J, M-O**, green), TrkB (**B, E, H, K**, green), TrkC (**C, F, I, L**, green), and Six1 (**D-F, J-L, N-O**, red). Arrowheads point to neurons that co-express Six1 with TrkA (**D, J, N, O**), TrkB (**E**), or TrkC (**F**). Scale bars: 100μm (**A**), applies to **B-C, G-I, M**; 20μm (**D**), applies to **E-F, J-L, N-O**.

trigeminal ganglion must be primarily neural crest-derived, while TrkB and TrkC neurons

are placode-derived – a concept that has been alluded to in the literature, but not explicitly

demonstrated.

We aimed to validate these assumptions by co-labeling for the transcription factor

Six1, which has historically been used as a marker of placode-derived neurons in the

trigeminal ganglion (Karpinski et al. 2016; Moody and Lamantia 2015). Unexpectedly, we

observed a large population of Six1-positive cells at E11.5, but these cells rarely

expressed TrkB or TrkC. Instead, most of the Six1-positive cells at E11.5 expressed TrkA

(presumptively neural crest-derived) or were devoid of any Trk receptor (Fig. 8D-F). At

E12.5, the few neurons that strongly expressed Six1 also expressed TrkA, but never TrkB

or TrkC (Fig. 8J-L). Given the timing of TrkA (E11.5-13.5) versus TrkB/TrkC (E9-11)

neurogenesis, these data suggest Six1 is a marker of recently differentiated neurons in

the trigeminal ganglion, rather than an exclusive marker of the placodal lineage. This

claim is consistent with reports of Six1 expression in placode-derived neurons at E10.5

(Karpinski et al. 2016), and with our findings of Six1 expression in neural crest-derived

TrkA neurons at E11.5-12.5. This conclusion is further supported by evidence that Six1

is transiently expressed in all newly differentiated, neural crest-derived neurons of the

dorsal root ganglia (Yajima et al. 2014). To our knowledge, Six1 expression had not been

examined in the trigeminal ganglion past E10.5, when the vast majority of neurons are

still only placode-derived, thus explaining the gap in knowledge. With this information in

hand, TrkA and Six1 expression was also examined in *Elp1 CKO* trigeminal ganglia. Six1-

and TrkA-double positive neurons were still abundant in *Elp1 CKO* trigeminal ganglia at

E12.5 (Fig. 8M-O). Altogether, these results suggest that neural crest-specific deletion of

Elp1 leads to selective effects on neural crest-derived TrkA nociceptive neurons during the examined stages of trigeminal ganglion development, and that these effects largely occur after neuronal differentiation, ultimately resulting in decreased innervation and survival of trigeminal nociceptors.

Discussion

Since the discovery that mutations in *ELP1* cause FD, several roles for Elp1 in neuronal development, function, and degeneration have been revealed (Slaugenhaupt et al. 2001; Anderson et al. 2001; Dietrich and Dragatsis 2016; Lefcort et al. 2017). To date, studies have emphasized defects in dorsal root ganglion and sympathetic neurons, which relay somatosensory and viscerosensory information from the trunk and limbs to the central nervous system (George et al. 2013; Jackson et al. 2014; Hunnicutt et al. 2012; Abashidze et al. 2014; Goffena et al. 2018; Li, Gruner, and Tourtellotte 2020; Morini et al. 2016). However, many questions remain regarding the cellular and molecular mechanisms underlying FD phenotypes, particularly since Elp1 has been proposed to mediate several cytoplasmic (e.g., protein trafficking, α-tubulin acetylation, stress signaling, exocytosis, and tRNA modification and translation of codon-biased transcripts) and nuclear (i.e., transcription) functions (Dalwadi and Yip 2018; Dietrich and Dragatsis 2016; Lefcort et al. 2017). Furthermore, while Elp1 knockdown in zebrafish leads to some defects in the formation of the trunk neural crest-derived enteric ganglia (Cheng et al. 2015), there is a dearth of information about Elp1 in the cranial ganglia, even when clinical deficits observed in FD clearly implicate trigeminal sensory dysfunction (Mendoza-

Santiesteban et al. 2017; Barlow 2009; Geltzer et al. 1964; Gutiérrez, Norcliffe-Kaufmann,

and Kaufmann 2015; Palma et al. 2018; Won et al. 2019). Perhaps this is due to the

complexity of cranial ganglion development; dorsal root and sympathetic ganglia neurons

are strictly neural crest-derived, whereas cranial ganglia contain neurons derived from

both neural crest cells and ectodermal placodes. The trigeminal ganglion is especially

unique, with intermingled placode- and neural crest-derived neurons (Blentic et al. 2011;

D'amico-Martel 1982; D'Amico-Martel and Noden 1983; Hamburger 1961; Saint-Jeannet

and Moody 2014; Steventon, Mayor, and Streit 2014). In this study, we provide the first

in-depth analyses of the morphological and cellular consequences of *Elp1* deletion from

the neural crest lineage during trigeminal ganglion development. Moreover, our data

reveal insights on the contributions of neural crest versus placode cells to specific neuron

populations and nerve branches emanating from the trigeminal ganglion.

Elp1 is enriched in trigeminal neurons during the course of neurogenesis and early

innervation

We first examined Elp1 expression in the developing trigeminal ganglion using an

Elp1:LacZ reporter mouse, in addition to immunohistochemistry on wildtype tissue (Fig. 1

and Fig. 1–figure supplement 1). While we observed diffuse β-galactosidase staining

throughout the neural tube, the trigeminal ganglia, and some cranial mesenchyme cells

(Fig. 1A-C), Elp1 protein expression was more discrete, with robust expression in the

cytoplasm of differentiated neurons (Fig. 1D-P, Fig. 1–figure supplement 1). These results

mirror previously reported patterns of Elp1 protein expression in differentiated trunk

sensory and sympathetic neurons (Hunnicutt et al. 2012; Jackson et al. 2014; Abashidze

et al. 2014). Notably, little to no Elp1 protein was detected in developing glial cells within

the trigeminal ganglion or in glia that line the embryonic nerves at the stages examined

(Fig. 1N-P, Fig. 1–figure supplement 1), which represent the period of normal trigeminal

ganglion neurogenesis and early innervation (E. J. Huang, Zang, et al. 1999; A. Davies

and Lumsden 1984; Wilkinson et al. 1996). While the anatomical distribution of Elp1

protein overlapped with β-galactosidase staining in *Elp1:LacZ* mice, the observation that

Elp1 protein is more selectively expressed in neurons suggests that Elp1 expression may

be controlled at the post-transcriptional or post-translational level, such that Elp1 protein

is upregulated upon neuronal differentiation. Altogether, our findings of highly enriched

Elp1 expression in axonal compartments indicate extranuclear functions for Elp1 in

developing trigeminal ganglion neurons, in agreement with previous studies that localize

Elp1 to axonal transport and synaptic vesicles in other neuronal systems (Abashidze et

al. 2014; Naftelberg et al. 2016; Tourtellotte 2016; Li, Gruner, and Tourtellotte 2020).

403　Elp1 is required for trigeminal ganglion nerve outgrowth and innervation of target

404　tissues

Next, we examined trigeminal ganglion development and trigeminal nerve deficits

associated with FD using an established mouse model in which *Elp1* is conditionally

deleted from neural crest cells and their derivatives (George et al. 2013). We found that

Elp1 is not necessary for the migration of cranial neural crest cells to the developing

ganglion, as the trigeminal ganglia in *Elp1 CKO* at E10.5 possessed appropriately

condensed placodal neurons and neural crest cells and the gross morphology of the

trigeminal ganglion was similar to controls (Fig. 3). These findings complement previously

published results showing Elp1 is not required for the migration of trunk neural crest cells to dorsal root or sympathetic ganglia (George et al., 2013; Jackson et al., 2014). Interestingly, knockdown of another Elongator complex member, Elp3, in *Xenopus* neural crest cells inhibits their migration, suggesting that these Elongator subunits may have distinct functions in the neural crest (Yang et al. 2016).

As development proceeds, however, progressive defects in trigeminal nerve outgrowth and innervation become apparent in *Elp1 CKO* mice. By collecting high-resolution images of the intact ganglion and nerves over several timepoints, we observed various axonal abnormalities that preceded this outcome. In the ophthalmic, maxillary, and mandibular divisions of the trigeminal nerve at E11.5, axons generally extended in the direction of their respective targets, but many deviated from established fascicles along the way, resulting in disorganized nerves with a "hairy" appearance and ectopic branches in some locations (Fig. 4). These data suggest Elp1 may be required for axonal fasciculation in embryonic nerves and/or that Elp1 may regulate axonal branching in some trigeminal sensory neurons. Neuronal Elp1 depletion has been reported to increase or decrease neurite branching in different contexts (Hunnicutt et al. 2012; Abashidze et al. 2014; Jackson et al. 2014; Ohlen et al. 2017). While these varying observations could be due to variations in developmental timing, they may also highlight genuine differences in Elp1 function, depending upon the identity and environment of individual neurons.

The notion of context-dependent functions for Elp1 is further supported by our observations of different neuronal subtype responses to the loss of Elp1 within the trigeminal ganglion. By E12.5, at least some axons overcame aforementioned pathfinding deficiencies to reach their targets (Fig. 5). In the ophthalmic division of *Elp1 CKO*, the

frontal nerve initially extended branches around the eye (albeit less elaborately than in

controls, Fig. 5B, E, H), but then retracted, resulting in an even less complex pattern of

innervation by E13 (Fig. 6A, B, E, F). Therefore, at least some frontal nerve axons in *Elp1*

CKO are able to reach their target site but are not maintained. These results suggest that

Elp1 has a role in trigeminal sensory axon maintenance and neuronal survival as seen in

other neuronal populations (Jackson et al. 2014; Li, Gruner, and Tourtellotte 2020;

Naftelberg et al. 2016; Abashidze et al. 2014). In contrast to the initial invasion and

subsequent retraction observed in frontal nerve axons, the medial and lateral nasal

nerves (also ophthalmic division) never formed in the *Elp1 CKO* (Fig. 5C, F; Fig. 6C, D,

G, H), indicating nasal nerve axons failed to navigate to their target destinations. Perhaps

in these neurons, Elp1 is required for axons to respond to long-distance guidance cues.

Further studies into the unique molecular, morphological, and functional identities of

different trigeminal ganglion subpopulations are needed to understand the requirement

for Elp1 in these different trigeminal neuronal subtypes.

450   Elp1 is required for survival of TrkA neurons

Our data demonstrate that Elp1 is required for axon maintenance in some

trigeminal ganglion neurons, indicating a potential role in neurotrophic signaling.

Therefore, we interrogated the expression of Trk receptors in *Elp1 CKO* and control

embryos. We found that loss of Elp1 from neural crest cells impacts TrkA neurons in the

trigeminal ganglion, which generally develop into small diameter nociceptive neurons that

relay pain and temperature information (Mu et al. 1993; Genç, Ulupinar, and Erzurumlu

2005; Reichardt 2006). Our results complement previous studies that demonstrate Elp1

is required for the survival of TrkA neurons in sympathetic and dorsal root ganglia (George

et al. 2013; Jackson et al. 2014; Naftelberg et al. 2016; Li, Gruner, and Tourtellotte 2020;

Ohlen et al. 2017). We observed a striking decrease in TrkA expression in the distal

branches of the ophthalmic, maxillary, and mandibular nerves, as well as decreased TrkA

signal in the central nerve root and the ganglion, itself (Fig. 6, Fig. 6–figure supplement

1). This could be explained by changes in the neurogenesis of TrkA-positive neurons,

reduction in TrkA expression or trafficking, retraction of TrkA-positive axons, and/or death

of TrkA neurons. While reduced TrkA numbers in dorsal root ganglia have been partially

attributed to early cell cycle exit and cell death of Pax3-positive TrkA progenitors due to

reduced DNA repair (George et al. 2013, Goffena et al., 2018), we did not observe Elp1

expression in Pax3-positive cells in the trigeminal ganglion (Fig. 1–figure supplement 1),

which have been demonstrated as glial, not neuronal, progenitors (Baker, Stark, and

Bronner-Fraser 2002). Moreover, Sox10-positive glia and glial precursors were present

and normally distributed in *Elp1 CKO*, indicating, qualitatively, the absence of any initial

effects on these cell populations (Fig. 3C-D).

Furthermore, we observe the initial presence of newly differentiated (Six1-

positive), TrkA-positive, Elp1-negative neurons in *Elp1 CKO* trigeminal ganglia (Fig. 8M-

O and Fig. 7–figure supplement 1). Therefore, our findings are most consistent with the

idea that loss of Elp1 has major effects after neuronal differentiation; however, our

analyses do not definitively rule out changes in neurogenesis in *Elp1 CKO*. In *Elp1 CKO*,

we also observe that the majority of distal Tubb3-positive trigeminal nerve branches lack

TrkA, whereas TrkA is robustly expressed into the smallest visible branches of the

ophthalmic, maxillary, and mandibular nerves in controls at E13 (Fig. 6). Given the

observation of widespread cell death in *Elp1 CKO* trigeminal ganglia at E12.5 (Fig. 7A-I)

and the specific loss of TrkA neurons by E13.5 (Fig. 7J), it is unlikely that the remaining

nerve endings in *Elp1 CKO* at E13 are from TrkA neurons. Collectively, our findings

suggest Elp1 is required for trigeminal ganglion axons to grow, properly invade target

tissues, and receive neurotrophic feedback that would maintain projections and prevent

apoptosis, which are consistent with reports that severe innervation defects and

subsequent apoptosis account for the majority of post-mitotic TrkA neuron loss in the

trigeminal ganglia (Jackson et al. 2014; Abashidze et al. 2014; Li, Gruner, and Tourtellotte

2020).

In sympathetic neurons, Elp1 is reported to modulate TrkA/NGF retrograde

signaling by regulating the phosphorylation of TrkA receptors in signaling endosomes (Li,

Gruner, and Tourtellotte 2020). Our findings neither confirm nor negate the possibility of

a similar mechanism for Elp1 in some trigeminal ganglion neurons. For axons that

successfully navigate to their intended targets, deficient TrkA retrograde signaling could

result in dying axons and subsequent cell death. Indeed, the retraction of axons that we

note in the frontal nerve could indicate an inability to receive and/or appropriately respond

to target-derived neurotrophic support. The observation that TrkA expression is severely

diminished in distal axons (Fig. 6) could potentially contradict this mechanism or others

that implicate deficiencies in TrkA retrograde signaling, since the phosphorylation state

of TrkA receptors and/or retrograde transport speed is secondary to the requirement of

the presence of adequate TrkA-positive receptors on distal axon terminals (Li, Gruner,

and Tourtellotte 2020; Naftelberg et al. 2016). However, examination of earlier timepoints

may reveal the initial presence of TrkA in Elp1-deficient nerve endings, followed by a

decrease over time due to failed innervation and cell death. Detailed spatio-temporal studies are needed to gain a better understanding of the dynamics of TrkA expression and trafficking in trigeminal ganglion neurons and the role of Elp1 in these processes. Alternatively, Elp1 may be required for axonal outgrowth through other mechanisms, including anterograde trafficking of critical receptors for growth or guidance factors and/or regulation of local protein synthesis, all of which are required within growing axons (Scott-Solomon and Kuruvilla 2018; Ascaño et al. 2009; Cioni et al. 2019; Korsak et al. 2016; Batista and Hengst 2016; Kang and Schuman 1996). It is important to remember that these functions of Elp1 could be direct or indirect via its essential role as part of the Elongator complex in modifying tRNAs (B. Huang, Johansson, and Byström 2005; B. Huang, Jian, and Byström 2008; Y.-T. Chen et al. 2009; C. Chen, Tuck, and Byström 2009; Karlsborn, Tükenmez, Mahmud, et al. 2014; Karlsborn, Tükenmez, Chen, et al. 2014; Goffena et al. 2018). Indeed, the dorsal root ganglia proteome in *Elp1 CKO* exhibits substantial changes (compared to control) across a wide range of cellular functions, including axon guidance and pathfinding (Goffena et al. 2018), but whether similar proteome alterations occur in the trigeminal ganglion remain to be explored. Collectively, these diverse functions for Elp1 could explain, in part, the varying outcomes we observe across the trigeminal nerves, given the complexities associated with different target tissues, long-range signals, and neurotrophic factors required by different trigeminal neuron subpopulations. Moreover, the dual origin of the trigeminal ganglion makes it difficult to know whether these phenotypes are due to intrinsic differences in neural crest-versus placode-derived neurons, which will be critical to parse out in future studies.

Insights into neural crest versus placodal cell dynamics in the developing trigeminal ganglion

During the course of this study, we made several observations on the temporal sequence of events during neurogenesis and axon growth in the developing trigeminal ganglion. Although the ganglion contains a mixture of neural crest- and placode-derived neurons, which we demonstrate both express Elp1 (Fig. 1), only neural crest-derived cells were targeted for *Elp1* deletion in these experiments. Since the vast majority of trigeminal ganglion neurons are placode-derived at E10.5 (Karpinski et al. 2016), it was not surprising that Tubb3 whole-mount immunohistochemistry, and trigeminal ganglion development, were similar between *Elp1 CKOs* and controls at this stage (Fig. 3). Striking alterations in the trigeminal ganglion and nerves were only noted in *Elp1 CKO* embryos after E10.5, once neural crest cells have begun differentiating into neurons. Neurons within the trigeminal ganglion will eventually express a distinct Trk receptor, which is typically correlated with a specific sensory modality (Patapoutian and Reichardt 2001). In *Elp1 CKO* embryos, TrkA neurons are vulnerable to loss of Elp1, with no impact on TrkB and TrkC neurons at the examined developmental stages.

The sequence of neural crest- versus placode-derived neuron differentiation may also clarify the developmental origin of Trk receptor-expressing neuronal subpopulations in the trigeminal ganglion. Previous studies demonstrated that, in the mouse trigeminal ganglion, TrkB and TrkC neurons are born before E11, whereas TrkA neurons are born on/after E11 (E. J. Huang, Wilkinson, et al. 1999; E. J. Huang, Zang, et al. 1999). As this timing coincides with placode-derived neurogenesis (starts at E9) preceding neural crest-derived neurogenesis (starts after E10.5, Karpinski et al. 2016), we propose that TrkB

and TrkC neurons are likely placode-derived while TrkA neurons are primarily neural

crest-derived. This notion is supported by the selective loss of TrkA neurons in neural

crest targeted-*Elp1 CKO* trigeminal ganglia, the absence of initial effects on TrkB and

TrkC neurons (Fig. 7J), and our findings that Six1, a marker of recently differentiated

sensory neurons, co-localizes to TrkA neurons, and not TrkB or TrkC neurons, at E11.5

and E12.5 (Fig. 8). While extensive studies have defined Trk receptor expression and the

timing of neural crest- vs. placode-derived neuron differentiation in the mouse trigeminal

ganglion (E. J. Huang, Wilkinson, et al. 1999; E. J. Huang, Zang, et al. 1999; Wilkinson

et al. 1996; Karpinski et al. 2016), this *Elp1 CKO* mouse has serendipitously allowed us

to deduce the cellular origin of Trk-expressing neuronal subtypes. Although no differences

in the number of TrkB or TrkC neurons were observed in our study, it is possible that

(presumptive placode-derived) TrkB or TrkC neurons will be indirectly affected later in

development, in response to the neural crest-specific loss of Elp1, given the important

reciprocal interactions between the two lineages (Shiau et al. 2008; Shiau and Bronner-

Fraser 2009; Wu and Taneyhill 2019; Steventon, Mayor, and Streit 2014; Wu et al. 2014).

It would be informative to selectively delete Elp1 from placode-derived neurons only, or

in addition to neural crest derivatives, in order to better understand the effects of cell type-

specific loss of Elp1 on trigeminal ganglion development. Unfortunately, there are no

trigeminal placode-specific Cre drivers available at this time to answer those questions.

Remarkably, the pattern of nerve deficits in *Elp1 CKO* embryos provides indirect

evidence that certain branches of trigeminal sensory nerves may be specifically neural

crest- or placode-derived. For instance, innervation of the whisker pad is initiated by

E11.5 in both controls and *Elp1 CKO* (Fig. 4A, D); however, the innervation field does not

increase substantially in the *Elp1 CKO* (Fig. 5C, F; Fig. 6C, D), and surviving nerve

terminals are TrkA-negative, whereas control embryos develop a more expansive, TrkA-

abundant innervation pattern by E13.5 (Fig. 6). This suggests that placode-derived

neurons may reach the whisker pad first, and act as scaffolds upon which later-

differentiating TrkA axons can traverse. A similar phenomenon is observed in the frontal

nerve of the ophthalmic division in *Elp1 CKO* embryos, where the main branch is

established by E11.5 (Fig. 4) and maintained at E13, but further elaboration and

innervation are not supported over developmental time (Fig. 5D, E, H; Fig. 6A, B). In

contrast, some trigeminal nerve branches never form in the *Elp1 CKO*, including the

medial and lateral nasal nerves of the ophthalmic division (Fig. 5F, Fig. 6D), begging the

question of whether these nerves are exclusively comprised of axons from neural crest-

derived neurons. More in-depth studies are required to understand the diversity of

interactions between neural crest- versus placode-derived cells during trigeminal

ganglion neurogenesis and innervation, and to understand the nuanced roles that Elp1

has in such a complex system.

FD is characterized by a wide range of sensory and autonomic phenotypes,

including those that implicate the involvement of the cranial trigeminal ganglion and its

associated nerves. Through studies in a conditional knockout mouse model that

eliminates Elp1 from the neural crest lineage, we have uncovered distinct effects on the

development of trigeminal sensory neurons, specifically those expressing TrkA that

generally function in nociception. Our results introduce a mechanistic basis for the

neuronal deficits that contribute to the loss of facial pain and temperature sensation

experienced by FD patients.

Materials & Methods

Animal Husbandry

All animal care and use described herein was in accordance with federal and institutional guidelines and approved by Montana State University's and University of Maryland's IACUC, under protocols #2018-81 (MSU) and #R-MAR-20-15 (UMD). The generation of the *Elp1* conditional knockout mouse was described previously by Dr. Frances Lefcort (Montana State University), who generously provided the embryos for this work (George et al. 2013). All mice were created and maintained on a C57BL/6 background. $Elp1^{fl/fl}$ mice have *LoxP* sites flanking the coding region of the fourth exon of *Elp1* (previously *Ikbkap*). When the floxed region is excised via Cre recombinase, the resulting truncated *Elp1* transcript is eliminated from cells by nonsense-mediated decay. The role of Elp1 in Wnt1-expressing neural crest cells and derivatives was examined by crossing homozygous $Elp1^{fl/fl}$ mice with hemizygous $Wnt1\text{-}Cre^{+/-}$ mice (The Jackson Laboratory, stock no. 003829) to create $Elp^{fl/+}$;Wnt1-Cre$^{+/-}$ males, which were then crossed with Elp1$^{fl/fl}$ females to generate *Elp1* conditional knockout mice ($Elp1^{fl/fl}$; $Wnt1\text{-}Cre^{+}$, abbreviated "*Elp1 CKO*" throughout). For all analyses, *Elp1 CKO* mice were compared to littermate controls ($Elp1^{fl/+}$; $Wnt1\text{-}Cre^{-}$) and at least 2 litters were examined per experiment. Genotyping was performed via PCR using the primer sequences listed below. For timed breeds, the day of the vaginal plug was considered embryonic day 0.5 (E0.5). Samples sizes were determined based on previously published studies using this mouse model or similar *Elp1* conditional knockouts mouse models (George et al. 2013; Jackson et al. 2014; Li, Gruner, and Tourtellotte 2020).

Genotyping

Genomic DNA was extracted using the Extracta DNA Prep for PCR kit (Quantabio) according to manufacturer instructions. *Elp1* CKO and wildtype alleles were detected with the following primers: 5'-GCACCTTCACTCCTCAGCAT-3' (forward) and 5'-AGTAGGGCCAGGAGAGAACC-3' (reverse). The *Wnt1-Cre* allele was detected with the following primers: 5'-GCCAATCTATCTGTGACGGC-3' (forward) and 5'-CCTCTATCGAACAAGCATGCG-3' (reverse). PCR mixtures were prepared using DreamTaq Green PCR Master Mix (Thermo Fisher) according to the manufacturer's protocol.

Tissue Collection and Preparation

Timed pregnant females were euthanized via CO_2 asphyxiation followed by cervical dislocation. Embryos were collected and placed in ice-cold 1X phosphate-buffered saline (1X PBS). A hindlimb bud was collected from each embryo for genotyping. Embryos were fixed via submersion and gentle shaking in 4% paraformaldehyde/1X PBS for between 20 minutes (E10.5) to 90 minutes (E13.5) at room temperature, then rinsed three times in 1X PBS for 20 minutes each. Fixed embryos were stored in 1X PBS with 0.02% sodium azide at 4°C until further analysis. For sectioning, embryos were rinsed twice with 1X PBS, then submerged in 15% sucrose (w/v) in 1X PBS at 4°C overnight, or until tissue sank, followed by submersion in 30% sucrose at 4°C until tissue sank. Embryos were first equilibrated in a 1:1 solution of 30% sucrose/1X PBS and Tissue-Tek Optimal Cutting Temperature compound (OCT, Fisher) for 2 hours at 4°C and then in 100% OCT at 4°C for 2 hours. Embryos were embedded in 100% OCT using liquid nitrogen vapor and stored at -80°C,

followed by sectioning at 12μm on a cryostat (Leica) and collection of tissue sections on

Superfrost Plus charged slides (VWR).

Immunohistochemistry

Tissue Sections

A hydrophobic boundary was drawn around tissue sections using an ImmEdge Pen

(Vector Labs). Tissue sections were rehydrated with 1X PBS for 5 minutes, then

permeabilized with 1X PBS/0.5% Triton X-100 (Tx-100) for 5 minutes at room temperature.

Tissue was blocked with 5% Bovine Serum Albumin (BSA, Fisher Scientific) (w/v) in PBS-

Tx (1X PBS, 0.1% Tx-100) for approximately 1 hour at room temperature, then rinsed once

in PBS-Tx. Primary antibodies were diluted in PBS-Tx plus 1% BSA and applied overnight

at 4°C in a humidified chamber. Unbound primary antibodies were washed off with 4 PBS-

Tx rinses for 5 minutes each at room temperature. Sections were then incubated with

secondary antibodies, diluted in PBS-Tx plus 1% BSA, for 1 hour at room temperature in

a humidified chamber. Sections were rinsed 3 times in PBS-Tx for 5 minutes each,

followed by 2 rinses in 1X PBS for 5 minutes each, all at room temperature. Coverslips

were mounted with DAPI Fluoromount-G Mounting Medium (Southern Biotech) and

allowed to dry in the dark at room temperature overnight before imaging.

Whole-mount

Fixed embryos were rinsed twice with 1X PBS for 5 minutes per rinse, then dehydrated

through a series of increasingly concentrated methanol (MeOH) washes (50:50, 80:20,

100:0 MeOH:PBS) for 30 minutes each at room temperature. Embryos were incubated in

Dent's Bleach (4:1:1 MeOH:DMSO:30% H_2O_2) for 6 hours at room temperature with gentle

shaking, then rehydrated through a series of decreasingly concentrated methanol washes

(50:50, 20:80, 0:100 MeOH:PBS) for 30 minutes each at room temperature. For blocking,

embryos were incubated in antibody dilution solution (1X PBS, 0.1% Tx-100, 5% BSA) for

at least 2 hours at room temperature. Next, embryos were incubated with fresh antibody

dilution solution containing primary antibodies for 2 days (E10.5) up to 4 days (E13.5) at

4°C with gentle shaking. Embryos were washed 4 times for ~1 hour each at room

temperature with PBS-Tx, then incubated in fresh dilution solution with secondary

antibodies for 1 day (E10.5) up to 3 days (E13.5) at 4°C with gentle shaking. Embryos

were washed 3 times for ~1 hour each at room temperature with PBS-Tx, followed by 2

washes with 1X PBS for 30 minutes each at room temperature. E10.5 embryos were

imaged at this step, whereas older embryos were cleared before imaging, as described

below.

Antibodies

Primary antibodies used included the following: Elp1 (Sigma #SAB2701068, 1:500), β-

tubulin III (Abcam #ab78078, 1:1000 for sections, 1:300 for whole-mount), Sox10 (R&D

#AF2864, 1:200 or GeneTex #GTX128374, 1:500), TrkA (R&D #AF1056, 1:500 for

sections, 1:200 for whole-mount), TrkB (R&D #AF1494, 1:300), TrkC (R&D #AF1404,

1:300), Six1 (Sigma #HPA001893, 1:500), Islet1 (DSHB #PCRP-ISL1-1A9, 1:500),

Neuropilin2 (R&D cat. AF567, 1:500), and Pax3 (DSHB, "Pax3", 1:200). All species/

isotype-specific, Alexafluor-conjugated secondary antibodies were purchased from

Thermo Scientific and used at a dilution of 1:500 on sections, or 1:300 in whole-mount.

FRUIT Clearing

After whole-mount immunohistochemistry, embryos were subjected to FRUIT clearing

(Hou et al. 2015). Briefly, embryos were moved through a series of aqueous FRUIT

buffers, containing 8M urea (Sigma), 0.5% (v/v) α-thioglycerol (TCI America), and

increasing concentrations of fructose (Fisher). Embryos were incubated at room

temperature with gentle rocking in 35% FRUIT for 6 hours, followed by 40% FRUIT

overnight, 60% FRUIT for 8 hours, and 80% FRUIT overnight. Embryos were stored in

80% FRUIT at 4°C until imaging in this buffer.

TUNEL Staining

TUNEL staining was performed on tissue sections after immunohistochemistry using the

In Situ Cell Death Detection Kit, TMR Red (Roche) according to the manufacturer's

instructions. After washing off unbound secondary antibodies, slides were post-fixed with

4% PFA in 1X PBS for 5 minutes at room temperature, then washed twice with 1X PBS

for 5 minutes each at room temperature. Sections were incubated with TUNEL reaction

mixture for 60 minutes at 37°C in the dark, followed by 3 washes in 1X PBS for 5 minutes

each at room temperature. Coverslips were mounted using DAPI Fluoromount-G

Mounting Medium (Southern Biotech) and allowed to dry in the dark at room temperature

overnight before imaging.

Imaging

E10.5 embryos that underwent whole-mount β-galactosidase staining or fluorescent

immunohistochemistry were imaged on a Zeiss SteREO Discovery V8 Pentafluor

fluorescent microscope using AxioVision software (Zeiss). Embryos E11.5 and older that

were processed for whole-mount immunohistochemistry were imaged in 80% FRUIT

buffer on a Zeiss LSM 800 confocal microscope. Z-stacks were collected at 5μm intervals

using 5X or 10X air objectives. Fluorescent immunohistochemistry on tissue sections was

also visualized on the Zeiss confocal microscope using 10X and 20X air objectives, or the

63X oil objective. For all applications, laser power, gain, offset, and digital zoom were

identical when imaging comparable regions of interest in Control versus *Elp1 CKO*

embryos and the pinhole was set to 1 airy scan unit at all times. CZI files were processed

in Zen software (Blue edition 2.0, Zeiss). For Z-stacks, CZI files were processed in

ImageJ, where Maximum Intensity Projections were created using the Z-Project function

in Hyperstack mode.

Nerve Tracing & Sholl Analysis

Tracing of the frontal nerve was performed on TIFF images of maximum intensity Z-stack

projections, described above, using the Simple Neurite Tracer plug-in in Image J (NIH).

After tracing, the Sholl Analysis function within the Simple Neurite Tracer plug-in was

used to quantify nerve branching complexity under the following settings: use standard

axes, no normalization of intersections, and 10μm radius step size. The center point was

set on the primary frontal nerve branch, just below the first branch point. Individual

distributions and group average distributions were plotted together in Microsoft Excel (N

= 2 Control, N = 4 *Elp1 CKO*, from two different litters).

Quantification of Trk-expressing neurons

10X images were acquired that encompassed the entire trigeminal ganglion within each

horizontal section, where tissue was co-labeled for Tubb3 and TrkA, TrkB, or TrkC. The

boundary of the trigeminal ganglion was determined in each section by Tubb3 staining,

which marked all neurons. Using the Cell Counter plug-in in ImageJ, each TrkA, B, or C

neuron was counted within the pre-determined trigeminal ganglion boundary. Neurons

were counted across 5 serial sections per animal, at the level of the maxillary nerve, and

the average number of neurons per section was calculated and presented as mean ±

standard error. The number of TrkA, B, or C neurons per section in *Elp1 CKO* was

compared to Control using an unpaired t-test, adjusted for multiple comparisons using the

Holm-Sidak method (Graphpad, N = 4 Control, N = 4 *Elp1 CKO* embryos, from two

different litters).

Acknowledgements

We thank Marta Chaverra for assistance with collecting embryos, Lynn George for

guidance regarding genetics, Vickie Riojas for help with animal husbandry, and Jenn

Lafrican for technical assistance.

Competing Interest Statement

745 Dr. Lefcort is the Co-Chair of the Scientific Advisory Board of the Familial Dysautonomia

746 Foundation, Inc.

References Cited

Abashidze, Anastasia, Veronica Gold, Yaron Anavi, Hayit Greenspan, and Miguel Weil. 2014. "Involvement of IKAP in Peripheral Target Innervation and in Specific JNK and NGF Signaling in Developing PNS Neurons." *PLoS ONE* 9 (11): 113428. https://doi.org/10.1371/journal.pone.0113428.

Anderson, Sylvia L, Rocco Coli, Ira W Daly, Elizabeth A Kichula, Matthew J Rork, Sabrina A Volpi, Josef Ekstein, and Berish Y Rubin. 2001. "Familial Dysautonomia Is Caused by Mutations of the IKAP Gene." *American Journal of Human Genetics* 68 (3): 753–58. https://doi.org/10.1086/318808.

Ascaño, Maria, Alissa Richmond, Philip Borden, and Rejji Kuruvilla. 2009. "Axonal Targeting of Trk Receptors via Transcytosis Regulates Sensitivity to Neurotrophin Responses." *Journal of Neuroscience* 29 (37): 11674–85. https://doi.org/10.1523/JNEUROSCI.1542-09.2009.

Baker, Clare V.H., Michael R. Stark, and Marianne Bronner-Fraser. 2002. "Pax3-Expressing Trigeminal Placode Cells Can Localize to Trunk Neural Crest Sites but Are Committed to a Cutaneous Sensory Neuron Fate." *Developmental Biology* 249 (2): 219–36. https://doi.org/10.1006/dbio.2002.0767.

Barlow, Steven M. 2009. "Central Pattern Generation Involved in Oral and Respiratory Control for Feeding in the Term Infant." *Current Opinion in Otolaryngology and Head and Neck Surgery* 17 (3): 187–93. https://doi.org/10.1097/MOO.0b013e32832b312a.

Batista, Andreia Filipa Rodrigues, and Ulrich Hengst. 2016. "Intra-Axonal Protein Synthesis in Development and Beyond." *International Journal of Developmental Neuroscience* 55: 140–49. https://doi.org/10.1016/j.ijdevneu.2016.03.004.

Blentic, Aida, David Chambers, Adam Skinner, Jo Begbie, and Anthony Graham. 2011. "The Formation of the Cranial Ganglia by Placodally-Derived Sensory Neuronal Precursors." *Molecular and Cellular Neuroscience* 46: 452–59. https://doi.org/10.1016/j.mcn.2010.11.010.

Chen, Changchun, Simon Tuck, and Anders S. Byström. 2009. "Defects in TRNA Modification Associated with Neurological and Developmental Dysfunctions in Caenorhabditis Elegans Elongator Mutants." *PLoS Genetics* 5 (7). https://doi.org/10.1371/journal.pgen.1000561.

Chen, Yei-Tsung, Matthew M. Hims, Ranjit S. Shetty, James Mull, Lijuan Liu, Maire Leyne, and Susan A. Slaugenhaupt. 2009. "Loss of Mouse Ikbkap , a Subunit of Elongator, Leads to Transcriptional Deficits and Embryonic Lethality That Can Be Rescued by Human IKBKAP." *Molecular and Cellular Biology* 29 (3): 736–44. https://doi.org/10.1128/mcb.01313-08.

Cheng, William Wai Chun, Clara Sze Man Tang, Hong Sheng Gui, Man Ting So, Vincent Chi Hang Lui, Paul Kwong Hang Tam, and Maria Mercè Garcia-Barcelo.

2015. "Depletion of the IKBKAP Ortholog in Zebrafish Leads to Hirschsprung Disease-like Phenotype." *World Journal of Gastroenterology* 21 (7): 2040–46. https://doi.org/10.3748/wjg.v21.i7.2040.

Cioni, Jean Michel, Julie Qiaojin Lin, Anne V. Holtermann, Max Koppers, Maximilian A.H. Jakobs, Afnan Azizi, Benita Turner-Bridger, et al. 2019. "Late Endosomes Act as MRNA Translation Platforms and Sustain Mitochondria in Axons." *Cell* 176 (1–2): 56-72.e15. https://doi.org/10.1016/j.cell.2018.11.030.

d'Amico-Martel, A, and D M Noden. 1980. "An Autoradiographic Analysis of the Development of the Chick Trigeminal Ganglion." *Journal of Embryology and Experimental Morphology* 55 (February): 167–82.

D'Amico-Martel, A, and D M Noden. 1983. "Contributions of Placodal and Neural Crest Cells to Avian Cranial Peripheral Ganglia." *The American Journal of Anatomy* 166 (4): 445–68. https://doi.org/10.1002/aja.1001660406.

D'amico-Martel, Adele. 1982. "Temporal Patterns of Neurogenesis in Avian Cranial Sensory and Autonomic Ganglia." *American Journal of Anatomy* 163 (4): 351–72. https://doi.org/https://doi.org/10.1002/aja.1001630407.

Dalwadi, Udit, and Calvin K Yip. 2018. "Structural Insights into the Function of Elongator." *Cellular and Molecular Life Sciences*. https://doi.org/10.1007/s00018-018-2747-6.

Davies, Alun, and Andrew Lumsden. 1984. "Relation of Target Encounter and Neuronal Death to Nerve Growth Factor Responsiveness in the Developing Mouse Trigeminal Ganglion." *Journal of Comparative Neurology* 223 (1): 124–37. https://doi.org/doi:10.1002/cne.902230110.

Davies, Alun M. 1997. "Studies of Neurotrophin Biology in the Developing Trigeminal System." *Journal of Anatomy* 191 (4): 483–91. https://doi.org/10.1017/S002187829700229X.

Dietrich, Paula, and Ioannis Dragatsis. 2016. "Familial Dysautonomia: Mechanisms and Models." *Genetics and Molecular Biology*. https://doi.org/10.1590/1678-4685-GMB-2015-0335.

Dietrich, Paula, Junming Yue, E. Shuyu, and Ioannis Dragatsis. 2011. "Deletion of Exon 20 of the Familial Dysautonomia Gene Ikbkap in Mice Causes Developmental Delay, Cardiovascular Defects, and Early Embryonic Lethality." *PLoS ONE* 6 (10): 27015. https://doi.org/10.1371/journal.pone.0027015.

Esberg, Anders, Bo Huang, Marcus J.O. Johansson, and Anders S. Byström. 2006. "Elevated Levels of Two TRNA Species Bypass the Requirement for Elongator Complex in Transcription and Exocytosis." *Molecular Cell* 24 (1): 139–48. https://doi.org/10.1016/j.molcel.2006.07.031.

Geltzer, Arthur I, Louis Gluck, Norman S Talner, and Herbert F Polesky. 1964. "Familial Dysautonomia." *New England Journal of Medicine* 271 (9): 436–40. https://doi.org/10.1056/NEJM196408272710903.

Genç, Bariş, Emel Ulupinar, and Reha S Erzurumlu. 2005. "Differential Trk Expression
in Explant and Dissociated Trigeminal Ganglion Cell Cultures."
https://doi.org/10.1002/neu.20134.

George, Lynn, Marta Chaverra, Lindsey Wolfe, Julian Thorne, Mattheson Close-Davis,
Amy Eibs, Vickie Riojas, et al. 2013. "Familial Dysautonomia Model Reveals Ikbkap
Deletion Causes Apoptosis of Pax3+ Progenitors and Peripheral Neurons."
Proceedings of the National Academy of Sciences of the United States of America
110 (46): 18698–703. https://doi.org/10.1073/pnas.1308596110.

Goffena, Joy, Frances Lefcort, Yongqing Zhang, Elin Lehrmann, Marta Chaverra,
Jehremy Felig, Joseph Walters, Richard Buksch, Kevin G Becker, and Lynn
George. 2018. "Elongator and Codon Bias Regulate Protein Levels in Mammalian
Peripheral Neurons." *Nature Communications* 9 (1). https://doi.org/10.1038/s41467-
018-03221-z.

Gold-von Simson, Gabrielle, and Felicia B. Axelrod. 2006. "Familial Dysautonomia:
Update and Recent Advances." *Current Problems in Pediatric and Adolescent*
Health Care 36 (6): 218–37. https://doi.org/10.1016/j.cppeds.2005.12.001.

Gutiérrez, Joel V., Lucy Norcliffe-Kaufmann, and Horacio Kaufmann. 2015. "Brainstem
Reflexes in Patients with Familial Dysautonomia." *Clinical Neurophysiology* 126 (3):
626–33. https://doi.org/10.1016/j.clinph.2014.06.028.

Hamburger, Viktor. 1961. "Experimental Analysis of the Dual Origin of the Trigeminal
Ganglion in the Chick Embryo." *Journal of Experimental Zoology* 148 (2): 91–123.
https://doi.org/https://doi.org/10.1002/jez.1401480202.

Hims, Matthew M, Ranjit S Shetty, James Pickel, James Mull, Maire Leyne, Lijuan Liu,
James F Gusella, and Susan A Slaugenhaupt. 2007. "A Humanized IKBKAP
Transgenic Mouse Models a Tissue-Specific Human Splicing Defect." *Genomics* 90
(3): 389–96. https://doi.org/10.1016/j.ygeno.2007.05.012.

Hou, Bing, Dan Zhang, Shan Zhao, Mengping Wei, Zaifu Yang, Shaoxia Wang, Jiarui
Wang, et al. 2015. "Scalable and DiI-Compatible Optical Clearance of the
Mammalian Brain." *Frontiers in Neuroanatomy* 9 (FEB): 1–11.
https://doi.org/10.3389/fnana.2015.00019.

Huang, Bo, Lu Jian, and Anders S. Byström. 2008. "A Genome-Wide Screen Identifies
Genes Required for Formation of the Wobble Nucleoside 5-
Methoxycarbonylmethyl-2-Thiouridine in Saccharomyces Cerevisiae." *Rna* 14 (10):
2183–94. https://doi.org/10.1261/rna.1184108.

Huang, Bo, Marcus J.O. Johansson, and Anders S. Byström. 2005. "An Early Step in
Wobble Uridine TRNA Modification Requires the Elongator Complex." *Rna* 11 (4):
424–36. https://doi.org/10.1261/rna.7247705.

Huang, Eric J., Keling Zang, Andrea Schmidt, Asta Saulys, Mengqing Xiang, and Louis
F. Reichardt. 1999. "POU Domain Factor Brn-3a Controls the Differentiation and
Survival of Trigeminal Neurons by Regulating Trk Receptor Expression."
Development 126 (13): 2869–82. https://pubmed.ncbi.nlm.nih.gov/10357931.

Huang, Eric J, George A Wilkinson, Isabel Fariñas, Carey Backus, Keling Zang, Sharyl
L Wong, and Louis F Reichardt. 1999. "Expression of Trk Receptors in the
Developing Mouse Trigeminal Ganglion: In Vivo Evidence for NT-3 Activation of
TrkA and TrkB in Addition to TrkC." *Development*. Vol. 126.
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2710120/pdf/nihms112680.pdf.

Hunnicutt, Barbara J, Marta Chaverra, Lynn George, and Frances Lefcort. 2012.
"Ikap/Elp1 Is Required in Vivo for Neurogenesis and Neuronal Survival, but Not for
Neural Crest Migration." *PLoS ONE* 7 (2).
https://doi.org/10.1371/journal.pone.0032050.

Jackson, Marisa Z, Katherine A Gruner, Charles Qin, and Warren G Tourtellotte. 2014.
"A Neuron Autonomous Role for the Familial Dysautonomia Gene ELP1 in
Sympathetic and Sensory Target Tissue Innervation." *Development (Cambridge)*
141 (12): 2452–61. https://doi.org/10.1242/dev.107797.

Jose-Alberto Palma, Lucy Kaufmann, Cristina Fuente, Leila Percival, Carlos Mendoza,
Horacio Kaufmann. 2008. "Current Treatments in Familial Dysautonomia." *Bone* 23
(1): 1–7. https://doi.org/10.1517/14656566.2014.970530.Current.

Kameneva, Polina, and Igor Adameyko. 2019. "Recent Advances in Our Understanding
of Central and Peripheral Nervous System Progenitors." *Current Opinion in Cell
Biology* 61: 24–30. https://doi.org/10.1016/j.ceb.2019.07.003.

Kang, H, and E M Schuman. 1996. "A Requirement for Local Protein Synthesis in
Neurotrophin-Induced Hippocampal Synaptic Plasticity." *Science (New York, N.Y.)*
273 (5280): 1402–6.

Karlsborn, Tony, Hasan Tükenmez, Changchun Chen, and Anders S. Byström. 2014.
"Familial Dysautonomia (FD) Patients Have Reduced Levels of the Modified
Wobble Nucleoside Mcm5s2U in TRNA." *Biochemical and Biophysical Research
Communications* 454 (3): 441–45. https://doi.org/10.1016/j.bbrc.2014.10.116.

Karlsborn, Tony, Hasan Tükenmez, A. K.M.Firoj Mahmud, Fu Xu, Hao Xu, and Anders
S. Byström. 2014. "Elongator, a Conserved Complex Required for Wobble Uridine
Modifications in Eukaryotes." *RNA Biology* 11 (12): 1519–28.
https://doi.org/10.4161/15476286.2014.992276.

Karpinski, Beverly A., Corey A. Bryan, Elizabeth M. Paronett, Jennifer L. Baker,
Alejandra Fernandez, Anelia Horvath, Thomas M. Maynard, Sally A. Moody, and
Anthony S. LaMantia. 2016. "A Cellular and Molecular Mosaic Establishes Growth
and Differentiation States for Cranial Sensory Neurons." *Developmental Biology*
415 (2): 228–41. https://doi.org/10.1016/j.ydbio.2016.03.015.

Kojic, Marija, Tomasz Gawda, Monika Gaik, Alexander Begg, Anna Salerno-Kochan,
Nyoman D Kurniawan, Alun Jones, et al. 2021. "Elp2 Mutations Perturb the
Epitranscriptome and Lead to a Complex Neurodevelopmental Phenotype." *Nature
Communications* 12 (1): 19. https://doi.org/10.1038/s41467-021-22888-5.

Korsak, Lulu, Molly Mitchell, Katerine Shepard, and Michael Akins. 2016. "Regulation of
Neuronal Gene Expression by Local Axonal Translation." *Physiology & Behavior*

176 (1): 100–106. https://doi.org/10.1007/s40142-016-0085-2.Regulation.

Lefcort, Frances, Marc Mergy, Sarah B Ohlen, Yumi Ueki, and Lynn George. 2017.
"Animal and Cellular Models of Familial Dysautonomia." *Clinical Autonomic*
Research. https://doi.org/10.1007/s10286-017-0438-2.

Lefler, Sharon, Malkiel A. Cohen, Gal Kantor, David Cheishvili, Aviel Even, Anastasya
Birger, Tikva Turetsky, et al. 2015. "Familial Dysautonomia (FD) Human Embryonic
Stem Cell Derived PNS Neurons Reveal That Synaptic Vesicular and Neuronal
Transport Genes Are Directly or Indirectly Affected by IKBKAP Downregulation."
PLoS ONE 10 (10): 1–28. https://doi.org/10.1371/journal.pone.0138807.

Li, Lin, Katherine Gruner, and Warren G Tourtellotte. 2020. "Retrograde Nerve Growth
Factor Signaling Abnormalities in Familial Dysautonomia." *The Journal of Clinical*
Investigation 130 (5). https://doi.org/10.1172/JCI130401.

Méndez-Maldonado, Karla, Guillermo A. Vega-López, Manuel J. Aybar, and Iván
Velasco. 2020. "Neurogenesis From Neural Crest Cells: Molecular Mechanisms in
the Formation of Cranial Nerves and Ganglia." *Frontiers in Cell and Developmental*
Biology 8 (August). https://doi.org/10.3389/fcell.2020.00635.

Mendoza-Santiesteban, Carlos E., Jose Alberto Palma, Lucy Norcliffe-Kaufmann, and
Horacio Kaufmann. 2017. "Familial Dysautonomia: A Disease with Hidden Tears."
Journal of Neurology 264 (6): 1290–91. https://doi.org/10.1007/s00415-017-8486-z.

Mezey, Éva, Alissa Parmalee, Ildiko Szalayova, Sandra P Gill, Math P Cuajungco,
Maire Leyne, Susan A Slaugenhaupt, and Michael J Brownstein. 2003. "Of Splice
and Men: What Does the Distribution of IKAP MRNA in the Rat Tell Us about the
Pathogenesis of Familial Dysautonomia?" *Brain Research* 983 (1–2): 209–14.
https://doi.org/10.1016/S0006-8993(03)03090-7.

Moody, Sally A, and Anthony-Samuel Lamantia. 2015. "Transcriptional Regulation of
Cranial Sensory Placode Development HHS Public Access." *Curr Top Dev Biol*
111: 301–50. https://doi.org/10.1016/bs.ctdb.2014.11.009.

Morini, Elisabetta, Paula Dietrich, Monica Salani, Heather M Downs, Gregory R
Wojtkiewicz, Shanta Alli, Anthony Brenner, et al. 2016. "Sensory and Autonomic
Deficits in a New Humanized Mouse Model of Familial Dysautonomia."
https://doi.org/10.1093/hmg/ddv634.

Mu, Xiaojun, Iva Inmaculada Silos-Santiago, Steven L Carroll, and William D Snider.
1993. "Neurotrophin Receptor Genes Are Expressed in Distinct Patterns in
Developing Dorsal Root Ganglia." *The Journal of Neuroscience*. Vol. 73.

Naftelberg, Shiran, Ziv Abramovitch, Shani Gluska, Sivan Yannai, Yuvraj Joshi, Maya
Donyo, Keren Ben-Yaakov, et al. 2016. "Phosphatidylserine Ameliorates
Neurodegenerative Symptoms and Enhances Axonal Transport in a Mouse Model
of Familial Dysautonomia." *PLoS Genetics* 12 (12): 1–27.
https://doi.org/10.1371/journal.pgen.1006486.

Ohlen, Sarah B, Magdalena L Russell, Michael J Brownstein, and Frances Lefcort.

2017. "BGP-15 Prevents the Death of Neurons in a Mouse Model of Familial
Dysautonomia." *Proceedings of the National Academy of Sciences of the United*
States of America 114 (19): 5035–40. https://doi.org/10.1073/pnas.1620212114.

Palma, Jose Alberto, Christy Spalink, Erin P. Barnes, Lucy Norcliffe-Kaufmann, and
Horacio Kaufmann. 2018. "Neurogenic Dysphagia with Undigested Macaroni and
Megaesophagus in Familial Dysautonomia." *Clinical Autonomic Research* 28 (1):
125–26. https://doi.org/10.1007/s10286-017-0487-6.

Park, Byung-Yong, and Jean-Pierre Saint-Jeannet. 2010. *Induction and Segregation of*
the Vertebrate Cranial Placodes. San Rafael (CA).
https://doi.org/10.4199/C00014ED1V01Y201007DEB003.

Patapoutian, Ardem, and Louis F. Reichardt. 2001. "Trk Receptors: Mediators of
Neurotrophin Action." *Current Opinion in Neurobiology.*
https://doi.org/10.1016/S0959-4388(00)00208-7.

Pearson, John, Barbara A. Pytel, Nicola Grover-Johnson, Felicia Axelrod, and Joseph
Dancis. 1978. "Quantitative Studies of Dorsal Root Ganglia and Neuropathologic
Observations on Spinal Cords in Familial Dysautonomia." *Journal of the*
Neurological Sciences 35 (1): 77–92. https://doi.org/10.1016/0022-510X(78)90103-
X.

Reichardt, Louis F. 2006. "Neurotrophin-Regulated Signalling Pathways." *Philosophical*
Transactions of the Royal Society of London. Series B, Biological Sciences 361
(1473): 1545–64. https://doi.org/10.1098/rstb.2006.1894.

Saint-Jeannet, Jean Pierre, and Sally A. Moody. 2014. "Establishing the Pre-Placodal
Region and Breaking It into Placodes with Distinct Identities." *Developmental*
Biology 389 (1): 13–27. https://doi.org/10.1016/j.ydbio.2014.02.011.

Schwartzlow, Coreen, and Mohamed Kazamel. 2019. "Hereditary Sensory and
Autonomic Neuropathies: Adding More to the Classification." *Current Neurology*
and Neuroscience Reports 19 (8). https://doi.org/10.1007/s11910-019-0974-3.

Scott-Solomon, Emily, and Rejji Kuruvilla. 2018. "Mechanisms of Neurotrophin
Trafficking via Trk Receptors." *Molecular and Cellular Neuroscience* 91 (March):
25–33. https://doi.org/10.1016/j.mcn.2018.03.013.

Shiau, Celia E., and Marianne Bronner-Fraser. 2009. "N-Cadherin Acts in Concert with
Slit1-Robo2 Signaling in Regulating Aggregation of Placode-Derived Cranial
Sensory Neurons." *Development* 136 (24): 4155–64.
https://doi.org/10.1242/dev.034355.

Shiau, Celia E, Peter Y Lwigale, Raman M Das, Stuart A Wilson, and Marianne
Bronner-Fraser. 2008. "Robo2-Slit1 Dependent Cell-Cell Interactions Mediate
Assembly of the Trigeminal Ganglion." *NATURE NEUROSCIENCE VOLUME* 11.
https://doi.org/10.1038/nn2051.

Sholl, D A. 1953. "Dendritic Organization in the Neurons of the Visual and Motor
Cortices of the Cat." *Journal of Anatomy* 87 (4): 387–406.

Slaugenhaupt, Susan A, Anat Blumenfeld, Sandra P Gill, Maire Leyne, James Mull, Math P Cuajungco, Christopher B Liebert, et al. 2001. "Tissue-Specific Expression of a Splicing Mutation in the IKBKAP Gene Causes Familial Dysautonomia." *American Journal of Human Genetics* 68 (3): 598–605. https://doi.org/10.1086/318810.

Steventon, Ben, Roberto Mayor, and Andrea Streit. 2014. "Neural Crest and Placode Interaction during the Development of the Cranial Sensory System." *Developmental Biology* 389 (1): 28–38. https://doi.org/10.1016/j.ydbio.2014.01.021.

Tourtellotte, Warren G. 2016. "Axon Transport and Neuropathy: Relevant Perspectives on the Etiopathogenesis of Familial Dysautonomia." *American Journal of Pathology* 186 (3): 489–99. https://doi.org/10.1016/j.ajpath.2015.10.022.

Wilkinson, George A, Isabel Fariñ, Carey Backus, Cathleen K Yoshida, and Louis F Reichardt. 1996. "Neurotrophin-3 Is a Survival Factor In Vivo for Early Mouse Trigeminal Neurons."

Won, Eugene, Jose Alberto Palma, Horacio Kaufmann, Sarah S. Milla, Benjamin Cohen, Lucy Norcliffe-Kaufmann, James S. Babb, and Yvonne W. Lui. 2019. "Quantitative Magnetic Resonance Evaluation of the Trigeminal Nerve in Familial Dysautonomia." *Clinical Autonomic Research* 29 (4): 469–73. https://doi.org/10.1007/s10286-019-00593-0.

Wu, Chyong Yi, Rachel M. Hooper, Kyeong Han, and Lisa A. Taneyhill. 2014. "Migratory Neural Crest Cell AN-Catenin Impacts Chick Trigeminal Ganglia Formation." *Developmental Biology* 392 (2): 295–307. https://doi.org/10.1016/j.ydbio.2014.05.016.

Wu, Chyong Yi, and Lisa A. Taneyhill. 2019. "Cadherin-7 Mediates Proper Neural Crest Cell–Placodal Neuron Interactions during Trigeminal Ganglion Assembly." *Genesis* 57 (1): 1–14. https://doi.org/10.1002/dvg.23264.

Xu, Huisha, Zhijie Lin, Fengzhi Li, Wentao Diao, Chunming Dong, Hao Zhou, Xingqiao Xie, Zheng Wang, Yuequan Shen, and Jiafu Long. 2015. "Dimerization of Elongator Protein 1 Is Essential for Elongator Complex Assembly." *Proceedings of the National Academy of Sciences of the United States of America* 112 (34): 10697–702. https://doi.org/10.1073/pnas.1502597112.

Yajima, Hiroshi, Makoto Suzuki, Haruki Ochi, Keiko Ikeda, Shigeru Sato, Ken ichi Yamamura, Hajime Ogino, Naoto Ueno, and Kiyoshi Kawakami. 2014. "Six1 Is a Key Regulator of the Developmental and Evolutionary Architecture of Sensory Neurons in Craniates." *BMC Biology* 12 (1): 1–18. https://doi.org/10.1186/1741-7007-12-40.

Yang, Xiangcai, Jiejing Li, Wanli Zeng, Chaocui Li, and Bingyu Mao. 2016. "Elongator Protein 3 (Elp3) Stabilizes Snail1 and Regulates Neural Crest Migration in Xenopus OPEN." https://doi.org/10.1038/srep26238.

York, Joshua R, Tian Yuan, and David W. McCauley. 2020. "Evolutionary and Developmental Associations of Neural Crest and Placodes in the Vertebrate Head:

1031   Insights From Jawless Vertebrates." *Frontiers in Physiology.*
1032   https://doi.org/10.3389/fphys.2020.00986.

1033