The dynamic properties of a nuclear coactivator binding domain are evolutionarily conserved

Elin Karlsson[1], Frieda A. Sorgenfrei[1,†], Eva Andersson[1], Jakob Dogan[1], Per Jemth[1,*], and Celestine N. Chi[1,2,*]

[1]Department of Medical Biochemistry and Microbiology, Uppsala University, BMC Box 582, SE-75123 Uppsala, Sweden.

[2]Department of Pharmaceutical Biosciences, Uppsala University, BMC Box 582, SE-75123 Uppsala, Sweden

[†]Present address: acib GmH, Krenngasse 37, 8010 Graz c/o University of Graz, Institute of Chemistry, NAWI Graz, BioTechMed Graz, Heinrichstrasse 28, 8010 Graz, Austria

*Correspondence to

Celestine Chi, Celestine.Chi@imbim.uu.se

Per Jemth, Per.Jemth@imbim.uu.se

ABSTRACT

Evolution of proteins is constrained by their structure and function. While there is a consensus that the plasticity of intrinsically disordered proteins relaxes the structural constraints on evolution there is a paucity of data on the molecular details of these processes. The Nuclear co-activator binding domain (NCBD) from CREB-binding protein is a protein-protein interaction domain, which contains a hydrophobic core but is not behaving as a typical globular domain, and has been desribed as 'molten-globule like'. The highly dynamic properties of NCBD makes it an interesting model system for evolutionary structure-function investigation of intrinsically disordered proteins. We have here compared the structure and biophysical properties of an ancient version of NCBD present in a bilaterian animal ancestor living around 600 million years ago with extant human NCBD. Using a combination of NMR spectroscopy, circular dichroism and kinetic methods we show that although NCBD has increased its thermodynamic stability, it has retained its dynamic biophysical properties in the ligand-free state in the evolutionary lineage leading from the last common bilaterian ancestor to humans. Our findings suggest that the dynamic properties of NCBD have been maintained by purifying selection and thus are important for its function, which includes mediating several distinct protein-protein interactions.

  Introduction

Evolution has shaped proteins into a wide spectrum of structure, stability and dynamics,

with fully disordered proteins at one end of the scale and well-folded globular proteins

at the other one (*1*). It is becoming clear that dynamic properties *per se* are important

for protein function (*1–3*) but it is not trivial to prove that the dynamics are essential

for biological function and not merely a general property of proteins. The Nuclear

Coactivator Binding Domain (NCBD) is a small (approximately 50 residues) domain

from CREB-binding protein (CBP, also called CREBBP), which is a transcriptional

coactivator with histone acetylase activity and present in all animals (*4*). The NCBD

domain has a hydrophobic core, but is very dynamic and even verging on being

intrinsically disordered (*5–7*). It has therefore been described as "molten-globule-like"

(*8–11*). The fact that NCBD possesses properties of both intrinsically disordered and

globular proteins makes it an interesting system for assessing the role of dynamics in

protein structure and how it modulates function. We have previously subjected NCBD

to "evolutionary biochemistry" (*12, 13*). Thus, we resurrected and characterized

ancestral versions of NCBD and its protein ligand CBP-Interacting Domain (CID) from

the NCOA1, 2 and 3 protein family (with three members in human called Src1, Tif2

and ACTR, respectively) to address the evolution of affinity in this protein-protein

interaction. The oldest "maximum likelihood" (ML) NCBD variant we could resurrect

was present before the Cambrian period, some 540-600 million years ago, in the

common ancestor of all present-day animals with bilaterian symmetry. These animals

are divided into deuterostomes and protostomes (D/P) and the ancestral NCBD domain

is denoted $NCBD_{D/P}{}^{ML}$. It was shown to bind the CID domain with relatively low

affinity ($K_d \sim$ 1-5 μM) whereas more recent variants of NCBD in the vertebrate lineage

(ca. 440 million years ago) had acquired present-day affinity for CID ($K_d \sim$ 0.1-0.2 µM).

Structural characterization of ancestral and present-day CID/NCBD complexes showed

that the increase in affinity observed for more modern NCBD variants was due to a

combination of factors. These include several new interactions driving structural and

dynamic changes in the complex, such as formation of a third α helix in CID upon

interaction (*13*). In the present paper, we investigate the structural and biophysical

properties of the ligand-free state of the ancestral Cambrian-like low-affinity

$NCBD_{D/P}^{ML}$ and compare it to that of high-affinity present-day $NCBD_{Human}$ to track the

evolution of structure and dynamics and their relation to function. The comparison

reveals that the overall properties of the ancestral $NCBD_{D/P}^{ML}$ domain are very similar

to those of the present-day $NCBD_{Human}$. In general, conservation implies function. Thus,

this conservation of a highly dynamic structure and molten-globule like properties

suggest that these features are subject to purifying selection and hence that dynamics

are indeed important for the biological function of NCBD.

Results

The structures of ancient $NCBD_{D/P}$ and extant $NCBD_{Human}$ are similar but not

identical. Because of its dynamic properties it is challenging to solve the structure of

ligand-free NCBD (*8*). First, we performed experiments with ^{15}N labeled $NCBD_{D/P}^{ML}$.

We discovered by serendipity that low pH significantly improved the quality of the

$^{1}H^{15}N$ heteronuclear single quantum coherence (HSQC) spectrum resulting in well

dispersed peaks for $NCBD_{D/P}^{ML}$ (Fig. 1), and allowing for a near complete assignment

of the backbone and side-chain residues from NMR 3D experiments using ^{13}C and ^{15}N

labeled samples. The NMR structure was thus determined at pH 2.4 by measuring 3D

1H-1H ^{13}C/^{15}N resolved NOESY experiments for distance restraints determination and

$^{3}J_{HNHA}$ for dihedral angle determination. The structure of NCBD$_{D/P}^{ML}$ was found to be

overall consistent with that of the free (8) and bound NCBD$_{Human}$ (6) and bound

NCBD$_{D/P}^{ML}$ determined at pH 6.8 (13) (Fig. 1). However, a detailed comparison

between NCBD$_{D/P}^{ML}$ and NCBD$_{Human}$ shows that the orientation of helix 1 (α1) and

helix 3 (α3) differ between the structures. Upon structural alignment of NCBD$_{Human}$

and NCBD$_{D/P}^{ML}$, we observed that both the α1 and α3 from NCBD$_{D/P}^{ML}$ are slightly

tilted away relative to that of NCBD$_{Human}$. We also observed that for CID-bound

NCBD$_{D/P}^{ML}$ and free NCBD$_{D/P}^{ML}$ the α3 displayed structural rearrangements (Fig. 2).

The amino acid sequence of NCBD$_{D/P}^{ML}$ was reconstructed using phylogenetic

methods[4]. The probability is low that the resulting maximum likelihood sequence is

identical to the actual sequence present in the ancestor. Thus, NCBD$_{D/P}^{ML}$ is rather one

of a large number of likely ancestral variants with similar properties. We wondered if

errors in the sequence of NCBD$_{D/P}^{ML}$ resulted in the differences we observe in helix α1

and α3 compared to NCBD$_{Human}$. To investigate this further, and to test how structurally

robust the dynamic NCBD domain is, we performed a control experiment where we

expressed, purified and characterized an alternative NCBD$_{D/P}$ variant denoted

NCBD$_{D/P}^{AltAll}$. In this NCBD$_{D/P}^{AltAll}$ variant, all residues with a posterior probability

lower than around 0.9 (12) were replaced with the second most likely residue at that

position. For example, residue 2107 is His with 87% probability and Gln with 12%

probability in the ancestral NCBD$_{D/P}$. Thus, NCBD$_{D/P}^{ML}$ has a His in position 2107 and

NCBD$_{D/P}^{AltAll}$ a Gln residue. NCBD$_{D/P}^{AltAll}$ can be considered a "worst case scenario"

variant and a good control of the robustness of conclusions drawn from resurrection

experiments. If ML and AltAll variants display similar properties it is very likely that

the actual ancestral protein shares these properties as well. Because of this, an AltAll

variant is a convenient alternative to combinations of point mutations (*14*).

$NCBD_{D/P}{}^{AltAll}$ contains six substitutions as compared to $NCBD_{D/P}{}^{ML}$ (Fig. 1). Three of

the differences between $NCBD_{D/P}{}^{ML}$ and $NCBD_{D/P}{}^{AltAll}$ are in the N-terminus, but only

one of these is in a structured region, a Gln2065→Pro at the beginning of α1. The other

three differences are Ser2078→Asn in the loop between α1 and α2, a solvent exposed

Gln2088→His in α2 and His2108→Gln at the C-terminus. We solved the NMR

structure of $NCBD_{D/P}{}^{AltAll}$ (Fig. S1) and found that it folds into a similar structure as the

other NCBD domains. The overall RMSD between $NCBD_{D/P}{}^{AltAll}$ and $NCBD_{D/P}{}^{ML}$ was

7.9 Å. As a comparison, the overall RMSD between $NCBD_{Human}$ and $NCBD_{D/P}{}^{AltAll}$ or

$NCBD_{D/P}{}^{ML}$ were 4.5 Å and 7.0 Å, respectively. Detailed inspection of the structures

revealed differences in the backbone between $NCBD_{D/P}{}^{ML}$ and $NCBD_{D/P}{}^{AltAll}$,

especially in the orientations of the helices. An overlay of the structures of

$NCBD_{D/P}{}^{AltAll}$, $NCBD_{D/P}{}^{ML}$, and $NCBD_{Human,}$ shows differences between α1, α2 and α3

(Fig. S1). One reason for the differences is that there are several long-range NOEs in

$NCBD_{D/P}{}^{AltAll}$ that are not present in $NCBD_{D/P}{}^{ML}$, between the following pairs of

residues: 2068/2082, 2069/2080, 2070/2083 and 2071/2099. There is also an NOE

between Ile2062 Hγ, and Pro2065 Hα, two of the substituted residues in $NCBD_{D/P}{}^{AltAll}$.

However, there appears to be a consistency in the hydrophobic core of both $NCBD_{D/P}{}^{ML}$

and $NCBD_{D/P}{}^{AltAll}$ since the single Phe residue at position 2100 is packed and buried in

a similar fashion in the two variants. Phe2100 is stabilized by Leu2090 in $NCBD_{D/P}{}^{ML}$,

and both Ile2089 and Leu2090 in $NCBD_{D/P}{}^{AltAll}$ (Fig. 3). Interestingly, this is the same

kind of packing that was observed in the free $NCBD_{human}$ domain (*8*), suggesting

conserved features in the hydrophobic core.

Despite the differences discussed in the previous paragraph, the overall biophysical

properties of $NCBD_{D/P}^{ML}$ and $NCBD_{D/P}^{AltAll}$ were found to be similar (Fig. S2)

corroborating the overall similarity seen with the NMR structures. Specifically, to test

the affinity of $NCBD_{D/P}^{AltAll}$ for CID, we expressed and purified ML and AltAll

versions of its ligand CID_{1R} based on our previous reconstruction (*8*). K_d values of all

four combinations ($NCBD_{D/P}^{ML}$ and $NCBD_{D/P}^{AltAll}$ with CID_{1R}^{ML} and CID_{1R}^{AltAll}) were

between 2-9 μM (Fig. S2a), which is in the same range as previous estimates of affinity

of the ancestral complex (*12*, *13*). Furthermore, the stability as monitored by urea

denaturation (Fig. S2b) and thermal denaturation (Fig. S2c-d) were similar for

$NCBD_{D/P}^{ML}$ and $NCBD_{D/P}^{AltAll}$. Thus, based on the highly similar biophysical

properties of $NCBD_{D/P}^{ML}$ and $NCBD_{D/P}^{AltAll}$ we conclude that our conclusions are

robust with regard to the uncertainty in the reconstructed sequence.

Ancient $NCBD_{D/P}$ is less thermodynamically stable than extant $NCBD_{Human}$ but

display a similar pH dependence with regard to structure and stability. Since the

$NCBD_{D/P}^{ML}$ structure was determined at low pH, we next investigated how pH

influences the fold and stability. First, we performed ^{1}H-^{15}N HSQC-monitored pH

titrations of $NCBD_{D/P}^{ML}$ and compared them to those of $NCBD_{Human}$ at different pH

values (Fig. 4). The NMR titration experiments showed that $NCBD_{D/P}^{ML}$ is as folded at

low pH as it is at high pH values. However, the peaks (^{1}H-^{15}N) are much sharper

(reduced line-width) at lower pH values indicating less conformational heterogeneity.

Second, the CD-monitored urea denaturation experiments showed identical stability

pattern at two pH values (3.0 and 7.4) for both ancient $NCBD_{D/P}^{ML}$ and extant

$NCBD_{Human}$ (Fig. 4c and d, Supplementary table S1). This indicates that while at lower

pH the $NCBD_{D/P}^{ML}$ still exhibits the same stability as at high pH values, there is a shift

in conformational heterogeneity towards a single state at low pH. Furthermore, whereas

the apparent thermodynamic stability differs by approximately 1 kcal mol^{-1} for ancient

$NCBD_{D/P}^{ML}$ and extant $NCBD_{Human}$, both are largely unaffected by a drop in pH to 3.0

(Supplementary Table S1). The observed difference in stability between $NCBD_{D/P}^{ML}$

and $NCBD_{Human}$ appears to be a robust result. Previous experiments suggested that the

ancestral $NCBD_{D/P}^{ML}$ was slightly less stable than younger and extant NCBD variants

as judged by urea denaturation experiments at pH 7.5 (*12*). However, the broad

unfolding transition of NCBD resulting from its small hydrophobic core leads to a large

error in the m_{D-N} value, which makes it difficult to unequivocally determine the

thermodynamic stability of the domain. However, all present results (including

experiments on the stabilized $NCBD_{D/P}^{ML}$ variant $NCBD_{D/P}^{T2073W}$ (Supplementary

Table S1) agree well with the previous experiments and we can therefore conclude with

greater confidence that the ancestral $NCBD_{D/P}^{ML}$ was less thermodynamically stable

than evolutionarily younger variants. Interestingly, with some exceptions (*15, 16*),

ancestral reconstructions often yield proteins with higher thermostability, which

correlates with thermodynamic stability, than extant proteins, either due to a bias for

stability in the reconstruction or due to higher temperatures in past times (*17–21*), like

during Cambrian (*22, 23*). In any case, ancestral NCBD appears to have populated the

native state to a lesser extent than present day human NCBD.

Ancient $NCBD_{D/P}^{ML}$ and extant $NCBD_{Human}$ display similar temperature-

dependent structural changes. Previous experiments demonstrated that $NCBD_{Human}$

shows an apparent non-cooperative unfolding behavior when subjected to increasing

temperatures, as monitored by CD (*6, 8*). This behavior may result from a low ΔH_{D-N}

value rather than a true non-cooperative unfolding. (The urea denaturation experiments

are consistent with a cooperative two-state unfolding.) We compared ^1H-^{15}N HSQC

and CD spectra, respectively, of $NCBD_{D/P}^{ML}$ and $NCBD_{Human}$ at different temperatures

(Fig. 5). In agreement with the previous data we observed that both variants unfold non-

cooperatively with temperature as monitored by the increase in molar ellipticity at 222

nm suggesting loss of α-helix as the temperature is increased. However, for both

$NCBD_{D/P}^{ML}$ and $NCBD_{Human}$ the signal is not completely lost even at 363 K (90°C)

underscoring that this is not a typical native to highly-disordered-denatured state

transition. Furthermore, ^1H-^{15}N HSQC spectra of $NCBD_{D/P}^{ML}$ at temperatures up to 333

K (60°C) show well dispersed peaks indicating a presumably globular, collapsed

architecture (Fig. S3) and corroborating the notion that NCBD retains structure at high

temperatures.

Rate constants for (un)folding of ancient $NCBD_{D/P}^{ML}$ and extant $NCBD_{Human}$. The

folding of human NCBD is complex, and, based on kinetic experiments on a human

NCBD variant, we proposed a three-state system involving 'native' NCBD (*i.e.*, the

solved structure), an intermediate state with unknown structure and a denatured state,

which likely retains significant structure under native conditions (*24*). Rate constants

in the same order was determined by relaxation dispersion NMR experiments by

Kjaergaard et al. (*10*) It is clear from the NMR experiments that low pH promotes the

native state of NCBD and that pH does not affect the thermodynamic stability as

monitored by urea denaturation experiments to any significant extent, neither for

$NCBD_{D/P}^{ML}$ nor $NCBD_{Human}$. Previous folding experiments on $NCBD_{Human}$ (*24*)

involving jumps from low to neutral pH thus resulted in small shifts in the equilibria

between the native, intermediate and denatured states yielding the observed kinetic

transients. In fact, a likely explanation for the better NMR spectra at low pH is that

there is less of the intermediate state under these conditions.

To further compare the biophysical properties of ancient $NCBD_{D/P}^{ML}$ and extant

$NCBD_{Human}$ we performed kinetic folding experiments using stopped-flow as well as

temperature-jump fluorescence spectroscopy. To achieve this we introduced a Trp at

position 2073 in $NCBD_{D/P}^{ML}$ as previously done in the folding studies of $NCBD_{Human}$

(*24*). This modification resulted in a thermodynamic stabilization of around 1 kcal mol$^-$

^{1} for both $NCBD_{D/P}^{T2073W}$ and $NCBD_{Human}^{T2073W}$ (Fig. 6a-b, Supplementary Table S1).

Previous folding experiments were performed on a slightly longer $NCBD_{Human}$ variant

(residues 2058-2116), which was used in the original studies of CID/NCBD (*6*). In our

resurrection studies, we used a shorter version of $NCBD_{Human}$, only containing the

evolutionarily conserved region (residues 2062-2109). Therefore, we first repeated the

temperature-jump folding experiments for $NCBD_{Human}^{T2073W}$ and showed that the

shorter construct used in our evolutionary studies displayed similar kinetics as the

longer construct (Fig. 6c). We then performed folding experiments with

$NCBD_{D/P}^{T2073W}$. The kinetic transients of $NCBD_{D/P}^{T2073W}$ from temperature jump

experiments had lower signal-to-noise than those of $NCBD_{Human}^{T2073W}$, but observed

rate constants were in the same range (Fig. 6d). To investigate the folding kinetics

further we used stopped-flow fluorimetry, which has a lower time resolution but can be

used to compare folding rate constants for $NCBD_{D/P}^{T2073W}$ and $NCBD_{Human}^{T2073W}$ at low

temperature (Fig. 6e-f). The stopped-flow folding experiments were conducted by

making pH jumps from a 5 mM HCl solution (pH ~ 2) to a buffer solution of 20 mM

sodium phosphate (pH 7.4, 150 mM NaCl, 1 M TMAO) and monitoring the relaxation

through a 330 nm band-pass filter at 4°C. The low temperature is necessary to reduce

the observed rate constant such that the folding or other conformational transitions take

place in the millisecond time window. We observed two kinetic phases for

$NCBD_{Human}^{T2073W}$ (one fast phase with a negative amplitude and a second slower phase

with a positive amplitude) but only one kinetic phase for $NCBD_{D/P}^{T2073W}$ (with a

positive amplitude). The values of the observed rate constants for $NCBD_{Human}^{T2073W}$

($\sim$300 s^{-1} and $\sim$70 s^{-1}, respectively) were similar to those observed previously for the

longer construct (*24*). The single value obtained for the $NCBD_{D/P}^{T2073W}$ was $\sim$140 s^{-1}.

Trp fluorescence is a very sensitive but crude structural probe. Thus, the number of

kinetic phases in folding experiments could be dependent on small structural

rearrangements around the Trp. Nevertheless, the observed rate constant for

$NCBD_{D/P}^{T2073W}$ in the stopped flow pH jump experiment is in the same range as the two

rate constants observed under the same conditions for human NCBD.

Discussion

The biophysics and molecular evolution of NCBD is interesting for several reasons,

which all relate to the dynamic properties of NCBD. Firstly, while it has a hydrophobic

core and globular shape, NCBD is a very dynamic protein domain with characteristics

of an IDP (*5*, *6*, *8–11*, *25*). Secondly, NCBD has several binding partners including the

three NCOA transcriptional co-regulator paralogs (*5*, *26*), the transcription factors p53

(*27*), Ets-2 (*28*) and interferon factor-3 (*29*), and viral proteins such as Tax (*30*) from

T-cell leukemia virus. Intriguingly, NCBD displays great conformational plasticity as

shown by its complex with interferon factor-3 where the helices of NCBD adopt a

distinct conformation as compared to the complex with CID (*31*). Thirdly, evolutionary

snapshots of the complex between CID and NCBD suggest that several new

intermolecular contacts and conformational plasticity increased the affinity when going

from the Cambrian-like $NCBD_{D/P}{}^{ML}$ ($K_d \sim$ 1-5 μM) to younger variants ($K_d \sim$ 0.2 μM)

(*12*, *13*, *32*). Fourthly, while affinity and structural order of modern NCBD/CID

complexes were increased, the transition state of the folding-induced binding displayed

more conformational heterogeneity as compared to the ancestral complex (*33*). Our

present results, how the free, unbound state of NCBD has evolved during the transition

from a low-affinity to a high-affinity CID binder relate to all four points.

The most striking finding is that $NCBD_{D/P}{}^{ML}$ displays very similar molten-globule like

properties as the previously characterized human NCBD (*8*, *10*). This conservation of

properties, borderline between a globular domain and an IDP, suggests that the

dynamics of NCBD is preserved by purifying selection and therefore of functional

importance. In general, structural plasticity is regarded as one main functional feature

of IDPs allowing for multiple protein-protein interactions (*34*). NCBD interacts with

several protein domains, both folded and disordered ones, and the dynamic nature of

NCBD likely increases its propensity to adapt to different partners. Moreover, NCBD

interactions result in coupled folding and binding. Many such reactions have recently

been shown to occur via a mechanism denoted templated folding, where the binding

partner influences the folding pathway (*35–37*). Indeed, this was observed for the

interaction between human NCBD and CID (*33*, *38*, *39*). Thus, our present findings

suggest that templated folding may be a general feature of binding reactions with

NCBD where the conserved dynamics facilitate sampling of different free states,

heterogeneity of bound states, as well as the pathway between free and bound states.

Going into details of the NMR model, we note that while overall structure is robust to

errors in the predicted sequence as shown by the comparison of the NMR models of

free $\text{NCBD}_{\text{D/P}}^{\text{ML}}$ and $\text{NCBD}_{\text{D/P}}^{\text{AltAll}}$, there are differences in for example helix

orientations. Globular domains with a sequence identity as high as these NCBD variants

(77-88%) would typically have identical structures due to a highly funneled energy

landscape (*40, 41*), while proteins with larger structural heterogeneity have less

funneled landscapes (*1*). Indeed, the differences between the structures of $\text{NCBD}_{\text{Human}}$,

$\text{NCBD}_{\text{D/P}}^{\text{ML}}$ and $\text{NCBD}_{\text{D/P}}^{\text{AltAll}}$, in particular regarding the angles of the helices in

relation to each other, are consistent with a less funneled energy landscape resulting in

conformational heterogeneity (*11*). This heterogeneity is also consistent with the

conserved dynamic properties of NCBD. In conclusion, our data suggest evolutionarily

conserved dynamics of NCBD that underlie functional plasticity in ground state

conformations, templated binding-and-folding and a propensity to interact with several

binding partners.

Materials and methods

Protein sequences, expression and purification. The reconstruction of ancestral

sequences was previously published (*12*). The AltAll versions of $\text{NCBD}_{\text{D/P}}$ (Fig. 1) and

CID_{1R} used in the present paper was based on posterior probabilities from the

reconstruction with a cutoff of 91%. One position in $\text{NCBD}_{\text{D/P}}$ (Thr2062) is very

uncertain. A Thr residue (15%) was originally chosen at this position in $\text{NCBD}_{\text{D/P}}^{\text{ML}}$ to

avoid a hydrophobic residue, although Ile (19%) has a higher posterior probability. In

the current $\text{NCBD}_{\text{D/P}}^{\text{AltAll}}$ residue 2062 is Ile. NCBD and CID variants were expressed

in *E. coli* from a pRSETA plasmid and purified as previously described (*33*)

Concentrations were determined by absorbance at 280 nm for NCBD (using calculated

extinction coefficients based on amino acid composition) or 205 nm for CID using an

extinction coefficient of 250,000 $M^{-1}cm^{-1}$. Purity was checked by SDS-PAGE and

identity of the purified proteins was confirmed by MALDI-TOF mass spectrometry.

Circular dichroism. Far-UV CD experiments were performed using a JASCO J-1500

CD spectrophotometer with a Peltier temperature control system. The protein

concentration in all CD experiments was 18-20 μM. The buffer solution was either 20

mM sodium phosphate (pH 7.4), 150 mM NaCl or 50 mM potassium formate (pH 3.0),

150 mM NaCl, and CD spectra were recorded at 277 K. The scanning speed was 50

nm/min, bandpass 1 nm and integration time 1 second. Each reported spectrum is an

average of 3-4 recorded spectra. In the equilibrium thermal denaturation experiments

the proteins were dissolved in 50 mM potassium formate buffer (pH 3.0), 150 mM NaCl

and the sample was heated from 277 to 368 K. The heating rate was 1 K/min, waiting

time 5 seconds and the denaturation was monitored at 222 nm. During the heating,

spectra (average of three measurements) were taken at 283, 303, 323, 343 and 363 K,

respectively. In the equilibrium urea denaturation experiments the proteins were

denatured in 0-6.6 M urea at 277 K and the unfolding was monitored at 222 nm. The

data was analyzed in GraphPad Prism and fitted by non-linear regression to a two-state

equilibrium model (*42*). Because of the broad unfolding transition and lack of native

baseline for several variants, four parameters were shared among all urea denaturation

data sets in the curve fitting (Fig. S4, Supplementary Table 1). One of these parameters

was the molar ellipticity of the native state, with the reasonable assumption that the

secondary structures of the native states are similar for all variants. The other

parameters were the slopes of the denatured and native baselines as well as the m_{D-N}

value (assuming a similar change in solvent accessible surface area upon denaturation

for all variants).

Temperature jump experiments. Temperature jump experiments were conducted

using 100-200 μM protein on a TJ-64 temperature jump system (TgK Scientific). The

buffer solution was 20 mM sodium phosphate (pH 7.4), 150 mM NaCl. The proteins

were subjected to temperature jumps of 2-8.5 K to different target temperatures. The

data were fitted in the Kinetic Studio software (TgK scientific) to a double exponential

function, to account both for the fast phase (relaxation time ~ 80 μs), which corresponds

to the heating time of the instrument, and to the slow phase, which corresponds to the

conformational transition in the protein. A Trp variant of $NCBD_{D/P}^{ML}$ ($NCBD_{D/P}^{T2073W}$)

was chosen for kinetic experiments since a Trp in this position was previously used for

human NCBD. We engineered a Trp at other positions in $NCBD_{D/P}^{ML}$ and obtained rate

constants in the same range for all variants (Supplementary Table S2).

Stopped flow experiments. The stopped flow experiments were performed using an

upgraded SX-17MV stopped flow spectrophotometer (Applied Photophysics) and the

measurements were conducted at 277 K. Proteins were dissolved in 5 mM HCl (pH ~

2) and rapidly mixed 1:1 with buffer solutions complemented with trimethylamine N-

oxide (TMAO) to a final concentration of 20 mM sodium phosphate (pH 7.4), 150 mM

NaCl, 1 M TMAO. The final protein concentration was 5 μM. The excitation

wavelength was 280 nm and the resulting fluorescence emission was detected after

passage through a 330 nm bandpass filter. The transients were analyzed in GraphPad

Prism and fitted to a single or double exponential function.

NMR experiments. NMR experiments were acquired on Bruker 600, 700 and 900 MHz spectrometers equipped with triple resonance cryogenic temperature probes at 298 K except otherwise stated. The final NMR samples contained 500 μM protein to which 0.01% NaN_3 and 5% D_2O was added. Experiments for assignment and subsequent structure determination were done at pH 2.4 (HCl adjusted with NaOH) for $NCBD_{D/P}^{ML}$ and pH 6.8 for $NCBD_{D/P}^{AltAll}$. The following NMR experiments were recorded for assignment and subsequent structure determination: standard 2D ^1H-^{15}N HSQC, 3D HNCACB ^{15}N-resolved [^1H-^1H]-NOESY-HSQC, ^{15}N-TOCSY-HSQC ^{13}C-resolved [^1H-^1H]-NOESY-HSQC, HCCH-TOCSY and ^{13}C-resolved [^1H-^1H]-TOCSY-HSQC. Phi-angle restraining $^3J_{HNHA}$ couplings were determined from 3D HNHA type experiments using quantitative $-J$ intensity modulated experiments (*43*). The temperature variation experiments were performed in 20 mM sodium formate, pH 3.0. All experiments were processed with nmrpipe (*44*) and analysed with CCPnmr (*45*).

Structure calculation. Structure calculations were done using the CYANA 3.97 (*46*) package as follows: Initially, cross peaks were converted into upper distance restraints following an automated process in CYANA. These distance restraints together with φ/ψ dihedral angles determined from C^α-chemical shifts and $^3J_{HNH}\alpha$ (measured) were used as input for the initial structure calculations. The structures were calculated with 200,000 torsion angle dynamics steps for 100 conformers starting from random torsion angles by simulated annealing. 20 conformers with the lowest target function values were selected and analyzed. The structural statistics together with all input data for the structure calculations are presented in Supplementary Table S3. Assignments and

structural coordinates have been deposited in the RCSB protein data bank with pdb

code: 7OSR (accession number BMRB ID 34635) ($NCBD_{D/P}^{ML}$) and 7OSW (accession

number BMRB ID 34636) ($NCBD_{D/P}^{AltAll}$).

Acknowledgements

This work was supported by the Wenner-Gren foundation WG-17 returning grants to

C. N. C., and by the Swedish Research Council (2020-04395 to P.J.) and Knut and

Alice Wallenberg foundation (Evolution of new genes and proteins, to P.J.). We used

the NMR Uppsala infrastructure, which is funded by the Department of Chemistry -

BMC and the Disciplinary Domain of Medicine and Pharmacy. We thank Magnus

Kjaergaard for valuable comments on the NMR experiments.

References

1. S. Gianni, M. I. Freiberger, P. Jemth, D. U. Ferreiro, P. G. Wolynes, M. Fuxreiter, Fuzziness and Frustration in the Energy Landscape of Protein Folding, Function, and Assembly. *Acc Chem Res*. **54**, 1251–1259 (2021).

2. O. F. Lange, N.-A. Lakomek, C. Farès, G. F. Schröder, K. F. A. Walter, S. Becker, J. Meiler, H. Grubmüller, C. Griesinger, B. L. de Groot, Recognition dynamics up to microseconds revealed from an RDC-derived ubiquitin ensemble in solution. *Science*. **320**, 1471–1475 (2008).

3. S. J. Wodak, E. Paci, N. V. Dokholyan, I. N. Berezovsky, A. Horovitz, J. Li, V. J. Hilser, I. Bahar, J. Karanicolas, G. Stock, P. Hamm, R. H. Stote, J. Eberhardt, Y. Chebaro, A. Dejaegere, M. Cecchini, J.-P. Changeux, P. G. Bolhuis, J. Vreede, P. Faccioli, S. Orioli, R. Ravasio, L. Yan, C. Brito, M. Wyart, P. Gkeka, I. Rivalta, G. Palermo, J. A. McCammon, J. Panecka-Hofman, R. C. Wade, A. Di Pizio, M. Y. Niv, R. Nussinov, C.-J. Tsai, H. Jang, D. Padhorny, D. Kozakov, T. McLeish, Allostery in Its Many Disguises: From Theory to Applications. *Structure*. **27**, 566–578 (2019).

4. H. J. Dyson, P. E. Wright, Role of Intrinsic Protein Disorder in the Function and Interactions of the Transcriptional Coactivators CREB-binding Protein (CBP) and p300. *J. Biol. Chem.* **291**, 6714–22 (2016).

5. C. H. Lin, B. J. Hare, G. Wagner, S. C. Harrison, T. Maniatis, E. Fraenkel, A small domain of CBP/p300 binds diverse proteins: solution structure and functional studies. *Mol Cell*. **8**, 581–590 (2001).

6. S. J. Demarest, M. Martinez-Yamout, J. Chung, H. Chen, W. Xu, H. J. Dyson, R. M. Evans, P. E. Wright, Mutual synergistic folding in recruitment of CBP/p300 by p160 nuclear receptor coactivators. *Nature*. **415**, 549–553 (2002).

7. M.-O. Ebert, S.-H. Bae, H. J. Dyson, P. E. Wright, NMR relaxation study of the complex formed between CBP and the activation domain of the nuclear hormone receptor coactivator ACTR. *Biochemistry*. **47**, 1299–308 (2008).

8. M. Kjaergaard, K. Teilum, F. M. Poulsen, Conformational selection in the molten globule state of the nuclear coactivator binding domain of CBP. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 12535–12540 (2010).

9. M. Kjaergaard, F. M. Poulsen, K. Teilum, Is a malleable protein necessarily highly dynamic? The hydrophobic core of the nuclear coactivator binding domain is well ordered. *Biophys J*. **102**, 1627–1635 (2012).

10. M. Kjaergaard, L. Andersen, L. D. Nielsen, K. Teilum, A folded excited state of ligand-free nuclear coactivator binding domain (NCBD) underlies plasticity in ligand recognition. *Biochemistry*. **52**, 1686–1693 (2013).

11. E. Papaleo, C. Camilloni, K. Teilum, M. Vendruscolo, K. Lindorff-Larsen,
Molecular dynamics ensemble refinement of the heterogeneous native state of
NCBD using chemical shifts and NOEs. *PeerJ*. **6**, e5125 (2018).

12. G. Hultqvist, E. Åberg, C. Camilloni, G. N. Sundell, E. Andersson, J. Dogan, C.
N. Chi, M. Vendruscolo, P. Jemth, Emergence and evolution of an interaction
between intrinsically disordered proteins. *Elife*. **6**, e16059 (2017).

13. P. Jemth, E. Karlsson, B. Vögeli, B. Guzovsky, E. Andersson, G. Hultqvist, J.
Dogan, P. Güntert, R. Riek, C. N. Chi, Structure and dynamics conspire in the
evolution of affinity between intrinsically disordered proteins. *Sci. Adv.* **4**,
eaau4130 (2018).

14. G. N. Eick, J. T. Bridgham, D. P. Anderson, M. J. Harms, J. W. Thornton,
Robustness of Reconstructed Ancestral Protein Functions to Statistical
Uncertainty. *Mol Biol Evol*. msw223 (2016).

15. K. M. Hart, M. J. Harms, B. H. Schmidt, C. Elya, J. W. Thornton, S. Marqusee,
Thermodynamic system drift in protein evolution. *PLoS Biol*. **12**, e1001994
(2014).

16. L. Laursen, J. Čalyševa, T. J. Gibson, P. Jemth, Divergent Evolution of a Protein-
Protein Interaction Revealed through Ancestral Sequence Reconstruction and
Resurrection. *Mol Biol Evol*. **38**, 152–167 (2021).

17. P. D. Williams, D. D. Pollock, B. P. Blackburne, R. A. Goldstein, Assessing the
accuracy of ancestral protein reconstruction methods. *PLoS Comput Biol*. **2**, e69
(2006).

18. V. A. Risso, J. A. Gavira, E. A. Gaucher, J. M. Sanchez-Ruiz, Phenotypic
comparisons of consensus variants versus laboratory resurrections of
Precambrian proteins. *Proteins*. **82**, 887–896 (2014).

19. M. L. Romero-Romero, V. A. Risso, S. Martinez-Rodriguez, B. Ibarra-Molero, J.
M. Sanchez-Ruiz, Engineering ancestral protein hyperstability. *Biochem J*. **473**,
3611–3620 (2016).

20. D. L. Trudeau, M. Kaltenbach, D. S. Tawfik, On the Potential Origins of the High
Stability of Reconstructed Ancestral Proteins. *Mol Biol Evol*. **33**, 2633–2641
(2016).

21. L. C. Wheeler, S. A. Lim, S. Marqusee, M. J. Harms, The thermostability and
specificity of ancient proteins. *Curr Opin Struct Biol*. **38**, 37–43 (2016).

22. T. W. Hearing, T. H. P. Harvey, M. Williams, M. J. Leng, A. L. Lamb, P. R.
Wilby, S. E. Gabbott, A. Pohl, Y. Donnadieu, An early Cambrian greenhouse
climate. *Sci Adv*. **4**, eaar5690 (2018).

23. T. Wotte, C. B. Skovsted, M. J. Whitehouse, A. Kouchinsky, Isotopic evidence
for temperate oceans during the Cambrian Explosion. *Sci Rep*. **9**, 6330 (2019).

24. J. Dogan, A. Toto, E. Andersson, S. Gianni, P. Jemth, Activation Barrier-Limited
Folding and Conformational Sampling of a Dynamic Protein Domain.
Biochemistry. **55**, 5289–5295 (2016).

25. S. J. Demarest, S. Deechongkit, H. J. Dyson, R. M. Evans, P. E. Wright, Packing,
specificity, and mutability at the binding interface between the p160 coactivator
and CREB-binding protein. *Protein Sci.* **13**, 203–210 (2004).

26. J. Xu, Q. Li, Review of the in vivo functions of the p160 steroid receptor
coactivator family. *Mol. Endocrinol.* **17**, 1681–1692 (2003).

27. J. A. Livengood, K. E. S. Scoggin, K. Van Orden, S. J. McBryant, R. S.
Edayathumangalam, P. J. Laybourn, J. K. Nyborg, p53 Transcriptional activity is
mediated through the SRC1-interacting domain of CBP/p300. *J Biol Chem.* **277**,
9054–9061 (2002).

28. G. Jayaraman, R. Srinivas, C. Duggan, E. Ferreira, S. Swaminathan, K.
Somasundaram, J. Williams, C. Hauser, M. Kurkinen, R. Dhar, S. Weitzman, G.
Buttice, B. Thimmapaya, p300/cAMP-responsive element-binding protein
interactions with ets-1 and ets-2 in the transcriptional activation of the human
stromelysin promoter. *J Biol Chem.* **274**, 17342–17352 (1999).

29. S. Matsuda, J. C. Harries, M. Viskaduraki, P. J. F. Troke, K. B. Kindle, C. Ryan,
D. M. Heery, A Conserved alpha-helical motif mediates the binding of diverse
nuclear proteins to the SRC1 interaction domain of CBP. *J Biol Chem.* **279**,
14055–14064 (2004).

30. K. E. Scoggin, A. Ulloa, J. K. Nyborg, The oncoprotein Tax binds the SRC-1-
interacting domain of CBP/p300 to mediate transcriptional activation. *Mol Cell
Biol.* **21**, 5520–5530 (2001).

31. B. Y. Qin, C. Liu, H. Srinath, S. S. Lam, J. J. Correia, R. Derynck, K. Lin, Crystal
structure of IRF-3 in complex with CBP. *Structure.* **13**, 1269–1277 (2005).

32. E. Karlsson, A. Lindberg, E. Andersson, P. Jemth, High affinity between
CREBBP/p300 and NCOA evolved in vertebrates. *Protein Sci.* **29**, 1687–1691
(2020).

33. E. Karlsson, C. Paissoni, A. M. Erkelens, Z. A. Tehranizadeh, F. A. Sorgenfrei, E.
Andersson, W. Ye, C. Camilloni, P. Jemth, Mapping the transition state for a
binding reaction between ancient intrinsically disordered proteins. *J Biol Chem.*
295, 17698–17712 (2020).

34. P. E. Wright, H. J. Dyson, Intrinsically disordered proteins in cellular signalling
and regulation. *Nat Rev Mol Cell Biol.* **16**, 18–29 (2015).

35. A. Toto, C. Camilloni, R. Giri, M. Brunori, M. Vendruscolo, S. Gianni, T. A, C.
C, G. R, B. M, V. M, G. S., Molecular Recognition by Templated Folding of an
Intrinsically Disordered Protein. *Sci. Rep.* **6**, 21994 (2016).

36. A. Toto, F. Malagrinò, L. Visconti, F. Troilo, L. Pagano, M. Brunori, P. Jemth, S.
Gianni, Templated folding of intrinsically disordered proteins. *J. Biol. Chem.*
295, 6586–6593 (2020).

37. D. Wu, H.-X. Zhou, W. D, Z. HX., Designed Mutations Alter the Binding
Pathways of an Intrinsically Disordered Protein. *Sci. Rep.* **9**, 6172 (2019).

38. P. Jemth, X. Mu, Å. Engström, J. Dogan, A frustrated binding interface for
intrinsically disordered proteins. *J. Biol. Chem.* **289**, 5528–5533 (2014).

39. E. Karlsson, E. Andersson, J. Dogan, S. Gianni, P. Jemth, C. Camilloni, A
structurally heterogeneous transition state underlies coupled binding and folding
of disordered proteins. *J. Biol. Chem.* **294**, 1230–1239 (2019).

40. C. Chothia, A. M. Lesk, The relation between the divergence of sequence and
structure in proteins. *EMBO J.* **5**, 823–826 (1986).

41. M. Oliveberg, P. G. Wolynes, The experimental survey of protein-folding energy
landscapes. *Q. Rev. Biophys.* **38**, 245–288 (2005).

42. A. Fersht, *Structure and mechanism in protein science: a guide to enzyme*
catalysis and protein folding. (Macmillan, 1999).

43. N.-A. Lakomek, J. Ying, A. Bax, Measurement of ^{15}N relaxation rates in
perdeuterated proteins by TROSY-based methods. *J Biomol NMR.* **53**, 209–221
(2012).

44. F. Delaglio, S. Grzesiek, G. W. Vuister, G. Zhu, J. Pfeifer, A. Bax, NMRPipe: a
multidimensional spectral processing system based on UNIX pipes. *J Biomol*
NMR. **6**, 277–293 (1995).

45. W. F. Vranken, W. Boucher, T. J. Stevens, R. H. Fogh, A. Pajon, M. Llinas, E. L.
Ulrich, J. L. Markley, J. Ionides, E. D. Laue, The CCPN data model for NMR
spectroscopy: development of a software pipeline. *Proteins.* **59**, 687–696 (2005).

46. P. Güntert, C. Mumenthaler, K. Wüthrich, Torsion angle dynamics for NMR
structure calculation with the new program Dyana. *J. Mol. Biol.* **273**, 283–298
(1997).

Figures

Figure 1. Structures of ancestral and human NCBD domains. (a) ^{1}H-^{15}N HSQC

spectrum of NCBD$_{D/P}^{ML}$ at pH 2.4 with assigned residues. All assignments and the

structural coordinates have been deposited in the pdb data bank with accession

number BMRB ID 34635 and pdb code: 7OSR. (b) A schematic tree depicting the

relationship between the deuterostome/protostome (D/P) ancestor and modern

species. The times on the x-axis are approximate, in particular the divergence of

protostomes and deuterostomes. (c) Sequence alignment for NCBD$_{Human}$, NCBD$_{D/P}^{ML}$

NCBD$_{D/P}^{AltAlt}$. The residues differing between NCBD$_{D/P}^{ML}$ and NCBD$_{D/P}^{AltAlt}$ are

marked in red while additional differences to NCBD$_{Human}$ are marked in yellow. (d)

Comparison of the NMR structure of ancestral NCBD$_{D/P}^{ML}$ solved at low pH (red, pdb

code: 7OSR), and the previously determined human NCBD (magenta, 2KKJ) (*8*).

Note that the structure of human NCBD was solved using a slightly longer construct.

(e) Overlay of the 20 lowest energy structures of the $NCBD_{D/P}^{ML}$ domain with those

of $NCBD_{human}$.

Figure 2. Comparison of the ligand-free and bound ancestral NCBD$_{D/P}^{ML}$ and extant NCBD$_{Human}$. The ligand-bound conformations (pdb codes 6ES5 and 6ES7) and the structure of free human NCBD (pdb code: 2KKJ) (*8*) were previously determined. (a) Overlay of the ^{1}H-^{15}N HSQC spectra between free NCBD$_{D/P}^{ML}$ (red) and free NCBD$_{Human}$ (grey). (b) Overlay of the ^{1}H-^{15}N HSQC spectra between CID-bound NCBD$_{D/P}^{ML}$ (magenta) and CID-bound NCBD$_{Human}$ (grey). Structural alignment between (c) free NCBD$_{D/P}^{ML}$ (red, 7OSR) and free NCBD$_{Human}$ (grey, 2KKJ), and (d) CID-bound NCBD$_{D/P}^{ML}$ (magenta, , 6ES5) and CID-bound NCBD$_{Human}$ (grey, 6ES7).

Figure 3. Phe2100 makes NOEs to Ile2089 and Leu2090 in NCBD$_{D/P}^{ML}$ and NCBD$_{D/P}^{AltAlt}$. (a) ^{1}H-^{13}C HSQC spectra of the methyl region of NCBD$_{D/P}^{ML}$, showing the Hδ1-Cδ1 shifts of L2090 and I2089. (b) ^{1}H-^{13}C HSQC spectra of the methyl region of NCBD$_{D/P}^{AltAlt}$, showing the Hδ1-Cδ1 shifts of L2090 and I2089. (c) Strip from a ^{1}H-^{1}H ^{13}C resolved NOESY showing NOEs from the Hβ1/2 atoms of Phe2100 and the associated NOEs to Leu2090Hδ1 for NCBD$_{D/P}^{ML}$. (d) Strip from a ^{1}H-^{1}H ^{13}C resolved NOESY showing NOEs from the Hβ1/2 atoms of Phe2100 and the associated NOEs to Leu2090Hδ1 and Ile2089Hδ1 for NCBD$_{D/P}^{AltAlt}$. These signature NOEs implies that both proteins form a hydrophobic core similar to that of NCBD$_{Human}$.

Figure 4. Comparison of ancestral NCBD$_{D/P}$^{ML} and extant NCBD$_{Human}$ at two

different pH values. a) Overlay of the ^{1}H-^{15}N HSQC spectra of NCBD$_{D/P}$^{ML} at pH 2.4

(red) and at pH 6.8 (cyan). b) Overlay of the ^{1}H-^{15}N HSQC spectra of NCBD$_{Human}$ at

pH 2.4 (blue) and at pH 6.8 (yellow). c) Urea denaturation experiments for NCBD$_{D/P}$^{ML}

at pH 3.0 (red) and at pH 7.4 (cyan). d) Urea denaturation experiments for NCBD$_{Human}$

at pH 3.0 (yellow) and at pH 7.4 (blue).

Figure 5. Temperature dependence for ancestral NCBD$_{D/P}^{ML}$ and extant human

NCBD. a) Overlay of ^{1}H-^{15}N HSQC spectra of NCBD$_{D/P}^{ML}$ at 283 K (blue) and at 303

K (cyan). b) Overlay of ^{1}H-^{15}N HSQC spectra of NCBD$_{Human}$ at 283 K (blue) and at 283

K (cyan). Circular dichroism spectra at a range of temperatures (283-363 K) for (c)

NCBD$_{D/P}^{ML}$ and (d) NCBD$_{Human}$. Molecular ellipticity at 222 nm (probing α-helix

content) as a function of temperature for (e) NCBD$_{D/P}^{ML}$ and (f) NCBD$_{Human}$.

Figure 6. Stability and temperature dependence of kinetic rate constants for

NCBD$_{D/P}$^{ML} and extant human NCBD. (a) Stability measurement of NCBD$_{Human}$^{WT}

(open circles) and NCBD$_{Human}$^{T2073W} (filled circles) measured by urea denaturation.

(b) Stability measurement of NCBD$_{D/P}$^{ML} (open circles) and NCBD$_{D/P}$^{T2073W} (filled

circles). Parameters from the curve fitting are shown in Supplementary Table S1. (c)

Temperature jump kinetic experiments for a longer human NCBD construct used in

previous studies (*24*) (open circles) and the shorter human NCBD variant used in this

study (filled circles). Both variants contained the T2073W mutation. (d) A

comparison of observed rate constants from temperature-jump experiments for

NCBD$_{Human}$^{T2073W} (blue) and NCBD$_{D/P}$^{T2073W} (red). All experiments were performed

in 20 mM sodium phosphate pH 7.4, 150 mM NaCl. (e-f) Kinetic (un)folding of

NCBD$_{Human}$^{T2073W} and NCBD$_{D/P}$^{T2073W} measured by stopped flow fluorescence

spectroscopy. (e) (Un)folding kinetic transient of NCBD$_{Human}$^{T2073W} (blue) fitted to a

double exponential function (black line). Below each transient are the residuals from

the respective fits. (f) Stopped flow (un)folding kinetic transient of NCBD$_{D/P}$^{T2073W}

(red) fitted to a single exponential function (black line). The experiments were

performed by jumping from 5 mM HCl (pH ~ 2) to a final buffer concentration of 20

mM sodium phosphate (pH 7.4), 150 mM NaCl, 1 M TMAO. The relaxation was

monitored using a 330 band-pass emission filter. Fitting a double exponential function

resulted in k_{obs} values of 290 ± 30 and 67 ± 4 s^{-1} for NCBD$_{Human}^{T2073W}$. Fitting a single

exponential function yielded a k_{obs} value of 139 ± 7 s^{-1} for NCBD$_{D/P}^{T2073W}$.

Supplementary information

The dynamic properties of a nuclear coactivator binding domain are evolutionarily conserved

Elin Karlsson[1], Frieda A. Sorgenfrei[1,†], Eva Andersson[1], Jakob Dogan[1], Per Jemth[1,*], and Celestine N. Chi[1,2,*]

[1]Department of Medical Biochemistry and Microbiology, Uppsala University, BMC Box 582, SE-75123 Uppsala, Sweden.

[2]Department of Pharmaceutical Biosciences, Uppsala University, BMC Box 582, SE-75123 Uppsala, Sweden

[†]Present address: acib GmbH, Krenngasse 37, 8010 Graz c/o University of Graz, Institute of Chemistry, NAWI Graz, BioTechMed Graz, Heinrichstrasse 28, 8010 Graz, Austria

*Correspondence to

Celestine Chi, Celestine.Chi@imbim.uu.se

Per Jemth, Per.Jemth@imbim.uu.se

Supplementary figures

Figure S1. Overlay of the structures of NCBD$_{D/P}{}^{ML}$, NCBD$_{D/P}{}^{AltAlt}$ and NCBD$_{Human}$. (a) An overlay of the structures determined from NMR for NCBD$_{D/P}{}^{ML}$ (red, pdb code: 7OSR), NCBD$_{D/P}{}^{AltAlt}$ (green, 7OSW), and the previously determined NCBD$_{Human}$ (magenta, 2KKJ). **(b)** Overlay of NCBD$_{D/P}{}^{AltAlt}$ (green, 7OSW), and the previously determined CID-bound NCBD$_{D/P}{}^{ML}$ (grey, 6ES5). **(c)** Overlay of NCBD$_{D/P}{}^{AltAlt}$ (green, 7OSW), and the previously determined NCBD$_{Human}$ (magenta, 2KKJ). **(d)** Overlay of NCBD$_{D/P}{}^{AltAlt}$ (green, 7OSW) and NCBD$_{D/P}{}^{ML}$ (red, 7OSR).

Figure S2. The biophysical properties of NCBD$_{D/P}^{ML}$ are robust to errors in the sequence. a) Binding affinities of NCBD$_{D/P}^{ML}$ or NCBD$_{D/P}^{AltAlt}$ and CID$_{1R}^{ML}$ or CID$_{1R}^{AltAll}$ measured with ITC. The buffer solution was 20 mM sodium phosphate pH 7.4, 150 mM NaCl. (b) Stability of NCBD$_{D/P}^{AltAll}$ measured by urea denaturation in 50 mM formate (pH 3.0), 150 mM NaCl. The CD signal at 222 nm was monitored and the data was fitted to a two-state function. A reliable [urea]$_{50\%}$ value could not be obtained, although qualitative comparison with NCBD$_{D/P}^{ML}$ indicates that the stability of the two variants is similar. (c-d) Temperature stability for NCBD$_{D/P}^{AltAll}$. The CD signal at 222 nm was monitored and spectra were taken at temperatures 283-363 K.

Figure S3. ^{15}N-^{1}H HSQC spectra of NCBD$_{D/P}^{ML}$ at different temperatures. The experiments were performed from 293 K (blue) to 333 K (red).

Figure S4. Equilibrium urea denaturation experiments. All data sets were fitted simultaneously to a two-state model in GraphPad Prism with shared parameters for the native state molar ellipticity, the denatured and native baseline slopes and the m_{D-N} value. Experiments were performed in either 50 mM potassium formate buffer (pH 3.0), 150 mM NaCl or 20 mM sodium phosphate buffer (pH 7.4), 150 mM NaCl. The shared m_{D-N} value was fitted as 0.74 ± 0.07 kcal mol^{-1}M^{-1}.

Supplementary Tables

Supplementary Table S1. Stabilities of NCBD variants expressed as the urea concentration at 50% denaturation. The proteins were denatured with urea and the structural transition was monitored with CD at 222 nm at 277 K. The solutions were either 50 mM potassium formate buffer (pH 3.0), 150 mM NaCl or 20 mM sodium phosphate buffer (pH 7.4), 150 mM NaCl. The data were fitted to a two-state model in GraphPad Prism. The parameters for the signal of the native state, the denatured and native baseline slopes, and the m_{D-N} value were shared among the data sets in the curve fitting to allow fitting of the low stability $NCBD_{D/P}{}^{ML}$ and $NCBD_{D/P}{}^{AltAll}$ variants. The shared m_{D-N} value was fitted as 0.82 ± 0.025 kcal mol^{-1}M^{-1}. The errors are the standard error from the curve fitting and likely a low estimate of.

Variant	pH	[Urea]$_{50\%}$ (M)	ΔG_{D-N} (kcal/mol)
$NCBD_{D/P}{}^{AltAll}$	3.0	0.22 ± 0.04	0.18 ± 0.03
$NCBD_{D/P}{}^{ML}$	3.0	0.38 ± 0.04	0.31 ± 0.03
$NCBD_{D/P}{}^{ML}$	7.4	0.68 ± 0.04	0.55 ± 0.04
$NCBD_{D/P}{}^{T2073W}$	3.0	2.13 ± 0.04	1.73 ± 0.06
$NCBD_{D/P}{}^{T2073W}$	7.4	2.08 ± 0.04	1.70 ± 0.06
$NCBD_{Human}$	3.0	1.61 ± 0.04	1.31 ± 0.05
$NCBD_{Human}$	7.4	1.94 ± 0.05	1.58 ± 0.06
$NCBD_{Human}{}^{T2073W}$	3.0	3.24 ± 0.07	2.64 ± 0.10
$NCBD_{Human}{}^{T2073W}$	7.4	3.46 ± 0.08	2.82 ± 0.11

Supplementary Table 2. Rate constants and kinetic amplitudes of folding for different D/P NCBD Trp variants measured in temperature jump experiments. The experiments were conducted using 200 μM protein in 20 mM sodium phosphate (pH 7.4), 150 mM NaCl. The jump in temperature was from 277 to 285.5 K.

Variant	Amplitude	k_{obs} (s^{-1})
NCBD$_{D/P}^{T2073W}$	1.06 ± 0.06	2400 ± 200
NCBD$_{D/P}^{L2067W}$	0.91 ± 0.02	1530 ± 50
NCBD$_{D/P}^{S2078W}$	0.35 ± 0.01	700 ± 60
NCBD$_{D/P}^{H2107W}$	0.47 ± 0.02	820 ± 80
NCBD$_{D/P}^{Q2108W}$	0.73 ± 0.02	960 ± 60

Supplementary Table 3

	NCBD$_{D/P}^{ML}$	NCBD$_{D/P}^{AltAlt}$
NMR distance and dihedral restraints		
Distance restraints		
Total NOEs	329	347
Intra-residue	113	145
Sequential ($|i - j| = 1$)	122	111
Medium-range ($1 < |i - j| < 4$)	76	76
Long-range ($|i - j| > 5$)	18	39
Total dihedral angle restraints	76	63
$^3J_{HN\alpha}$ scalar couplings	39	43
$^{13}C^{\alpha}$ chemical shifts	100	100
Structure statistics		
Average CYANA target function value (Å^2)	2.5 ± 0.2	2.8 ± 0.3
Violations		
Distance constraints (>0.5 Å)	0	0
Dihedral angle constraints (>5°)	0	0
Deviations from idealized geometry		
Bond lengths (Å)	0	0
Bond angles (º)	0	0
Impropers (º)	0	0
Average pairwise r.m.s. deviation (Å)		
Backbone residues	2.2 ± 1.0	0.5 ± 0.2
Heavy atoms residues	2.7 ± 1.0	1.0 ± 0.3