TITLE

A CRISPR toolbox for generating intersectional genetic mice for functional, molecular, and

anatomical circuit mapping

AUTHOR NAMES AND AFFILIATIONS

Savannah J. Lusk[1], Andrew McKinney[1], Patrick J. Hunt[3], Paul G. Fahey[1], Jay Patel[1], Jenny J.

Sun[1], Vena K. Martinez[2], Ping Jun Zhu[1], Jeremy R. Egbert[4], Xiaolong Jiang[1], Benjamin R.

Arenkiel[3,5], Andreas S. Tolias[1], Mauro Costa-Mattioli[1], Russell S. Ray[1,5,6]

[1]Department of Neuroscience, Baylor College of Medicine, Houston TX

[2]Department of Pharmacology, Baylor College of Medicine, Houston TX

[3]Department of Molecular and Human Genetics, Baylor College of Medicine, Houston TX

[4]Department of Cell Biology, University of Connecticut, Farmington CT

[5]McNair Medical Institute, Houston Tx

[6]Corresponding author, russell.ray@bcm.edu

ABSTRACT

Background

A full understanding of circuits and cellular mechanisms governing health and disease requires

the dissection and multi-faceted study of discrete cell subtypes in developing and adult animal

models. Recombinase-driven expression of transgenic response alleles represents a significant

and powerful approach to delineate cell populations for functional, molecular, and anatomical

study. In addition to single recombinase systems, the expression of two recombinases in distinct,

but partially overlapping, populations allows for more defined target expression. Although the

application of this method is becoming increasingly popular, the expense and difficulty associated

with production of customized intersectional mouse lines have limited widespread application to

more common allele manipulations that are often commercially produced at great expense.

Results

We present a simplified CRISPR toolkit for rapid, inexpensive, and facile intersectional allele

production. Briefly, we produced 7 intersectional mouse lines using a dual recombinase system,

one mouse line with a single recombinase system, and three embryonic stem (ES) cell lines that

are designed to study how functional, molecular, and anatomical features relate to each other in

building circuits that underlie physiology and behavior. As a proof-of-principle, we applied three

of these lines to different neuronal populations for anatomical mapping and functional *in vivo*

investigation of respiratory control. We also generated a mouse line with a single recombinase-

responsive allele that controls the expression of the calcium sensor Twitch-2B. This mouse line

was applied globally to study the effects of follicle stimulating hormone (FSH) and luteinizing

hormone (LH) on calcium release in the ovarian follicle.

Conclusions

Lines presented here are representative examples of outcomes possible with the successful

application of our genetic toolkit for the facile development of diverse, modifiable animal models.

This toolkit will allow labs to create single or dual recombinase effector lines easily for any cell

population or subpopulation of interest when paired with the appropriate Cre and FLP

recombinase mouse lines or viral vectors. We have made our tools and derivative intersectional

mouse and ES cell lines openly available for non-commercial use through publicly curated

repositories for plasmid DNA, ES cells, and transgenic mouse lines.

KEY WORDS

Intersectional genetics, mouse models, CRISPR toolkit

BACKGROUND

Targeted expression of effector molecules, like fluorescent markers, calcium reporters,

optogenetic actuators, or exogenous ligand-responsive receptors (DREADDs) (1), are

increasingly applied in a variety of fields for greater precision, quantitative expression level

control, and reduced side effects compared to previous methods for labeling and manipulation.

For example, the Cre/LoxP system is used to achieve permanent cell-type restriction by using a

promoter or enhancer to express the site-specific recombinase Cre (2). In such examples, cre

transgenes are paired with a constitutively active, but conditional, allele where expression of an

effector molecule, such as eGFP, is interrupted by a LoxP-Stop-LoxP cassette that is recombined

out by Cre, which enables expression in targeted, Cre-expressing cells. The use of various

effector molecules in this paradigm enables fluorescent marking, neuronal perturbation, molecular

affinity pull-downs, activity tracking, and other studies (1,3–28).

In some fields, it is becoming increasingly clear that recombinase expression based on a single

gene does not offer the resolution needed for a variety of developmental or targeting applications,

like circuit mapping (29–31). Intersectional genetics adds needed resolution by employing a dual

recombinase system using both Cre recombinase as well as a second recombinase, FLP, to

activate a conditional effector allele only in cells where both recombinases have been expressed

in the same cell (though not necessarily concurrently) (20,32–39). With new methodologies being

developed to use Cre and FLP not only as traditional genetic markers, but also as activity and

connectivity markers, unique combinatorial cell type definitions become possible (40–44). Thus, a

resource consisting of multiple dual recombinase intersectional alleles that each express different

effector molecules would add significant value and needed resolution to our ability to deconstruct

neural circuits on multiple levels. This technology could also be applied to a multitude of other

fields where intersectionally defined subpopulations of target cells exist and may play different

roles in the measured outcomes.

Although mouse intersectional technology provides relatively benign access to otherwise

inaccessible populations of cells, few laboratories have generated single transgenic or

intersectional genetic mouse lines in house for a number of reasons. The complexity and size of

the final targeting vectors puts them beyond present (cost-effective) commercial DNA synthesis

capabilities, thus requiring some level of recombinant DNA cloning and precluding straightforward

production of large, high fidelity ssDNA donors that can facilitate pro-nuclear CRISPR-mediated

targeting (45). While intersectional targeting vectors are available from the Addgene plasmid

repository, they are finished vectors that require significant reverse engineering or modifications

for use in a new context or approach. To our knowledge, there are no modular intersectional

targeting vectors that are publicly available for facile and rapid production of new targeting alleles

for the generation of intersectional mouse models. Furthermore, vector stability and other *in vitro*

difficulties combined with the expense and time associated with target vector insertion and mouse

line production limit the number of intersectional mouse lines available for public use. Thus, the

production of intersectional genetic mouse lines has been largely limited to a small number of

pioneering labs or resource rich institutions such as the Howard Hughes Medical Institute and the

Allen Institute for Brain Science (38,46).

To address these pitfalls and make the production of intersectional genetic mouse models more

widely feasible, we aimed to produce a freely available resource toolbox consisting of several

intersectional and single-recombinase responsive *Rosa26* targeting vectors for rapid, facile, and

cost-effective generation of complex mouse lines through the use of CRISPR/Cas9-mediated

homologous recombination in mouse embryonic stem cells and early oocytes. In oocytes,

genomic insertions/deletions (in/dels) and short targeted insertions are readily produced whereas

large construct targeting in oocytes has shown more limited but growing success. Additionally,

requisite equipment and facilities are difficult to access and out of reach for many investigators.

ES cell approaches are well established and widely available, allowing for rapid screening and

identification, and, if using morula aggregation, require much simpler methodologies and

equipment for mouse generation.

Towards this, we produced 7 intersectional mouse lines using a dual recombinase system; one

mouse line with a single recombinase system, and three additional ES cell lines to study how

functional, molecular, and anatomical features relate to each other in building the circuits that

underlie physiology and behavior. As a proof-of-principle, we applied three of these lines to

different neuronal populations for anatomical mapping and functional *in vivo* characterization in

respiratory control. Next, we globally applied the single recombinase-responsive line, which

controls the expression of the calcium sensor Twitch-2B. Twitch-2B was expressed globally in the

generated mouse line to study the effects of follicle stimulating hormone (FSH) and luteinizing

hormone (LH) on calcium release in the ovarian follicle. The publication and availability of this

technology will allow for the seamless production of a highly diverse group of mouse lines that

can be used to generate animal models of human disease, label specific cell populations for

developmental or connectivity studies, or modulate cellular activity in established disease model

lines, among other possibilities. All reagents and vectors used or generated in this study are now

openly available for not-for-profit research.

RESULTS

Vector design and optimization: For each of the targeting vectors generated, the intersectional

or Dre-responsive cassette was knocked into a well-established site in the *Rosa26* locus (**Fig. 1)**

(47,48). For positive ES cell clone selection, the targeting vector was simplified and improved in

several ways compared to earlier approaches (49). A simplified neomycin resistance cassette

was integrated into the intersectional cassette before the second LoxP-flanked stop cassette to

utilize the *CAG* promoter and polyadenylation (*pA*) sequences of the FRT flanked stop cassette

eliminating prior use of an additional PGK promoter and *Bovine Growth Hormone polyA* (BGHpA*)*

signal. To use CRISPR/Cas9, we cloned an sgRNA that was close to the 5'/3' homology junction

into the px330 vector (50), which expresses both Cas9 and the subcloned sgRNA

(*px330_Rosa26_sgRNA*). A Woodchuck Hepatitis Virus (WHV) Posttranscriptional Regulatory

Element (WPRE) and BGHpA were added at the end of the expression cassette to enhance

effector molecule expression and limit reliance on the disrupted Rosa Locus for transcript

termination (49). The homology arms were significantly shortened to 1Kb to remove repetitive

genomic sequences and stabilize the vector for prokaryotic propagation. Lastly, the terminal non-

homology Diphtheria Toxin A (DTA) chain negative selection cassette was removed, allowing for

the complete targeting vector to be functionally tested via cell culture or *in utero* electroporation,

which was not possible with the presence of the terminal DTA selection cassette without an

additional subcloning step. Redesigned vectors and their Addgene ID numbers are outlined in

Table 1 for public distribution.

Embryonic stem cell electroporation: We optimized traditional but widespread ES cell targeting

by applying CRISPR/Cas9-mediated HDR (51). Our initial electroporation (EP) experiments using

an earlier, more complex version of the *Rosa26* intersectional targeting vector (containing the

longer 4.2 kb 3' homology arm, DTA, and no CRISPR/Cas9) showed targeting rates ranging from

6-14% (over the course of four electroporations we saw targeting rates of 6/48, 3/48, 7/48, and

3/48 clones, mouse strains listed in **Table 2**). To determine the effect of CRISPR/Cas9 on

targeting efficiency, we co-electroporated the *px330_Rosa26_sgRNA* vector expressing Cas9

and a *Rosa26* specific sgRNA with the optimized *RR5* targeting vector at four different molar

ratios (0:1, 0.5:1, 1:1, and 10:1), proportionally decreasing the amount of donor vector to

accommodate the increasing *px330_Rosa26_sgRNA* vector per EP (18-20µg total DNA per EP

for $1x10^7$ ES cells) (**Fig 1C)**. Under the 0:1 *px330* vector:donor vector condition, we saw a 6%

targeting rate similar to those seen in electroporations without Cas9 and with a much longer 3'

homology arm and negative selection, suggesting that the shortening did not have a large impact

on targeting efficiency under our EP conditions. Under the 0.5:1, 1:1, and 10:1 conditions, we

observed 33%, 27%, and 58% targeting, respectively, suggesting that higher ratios of the Cas9

vector resulted in increased targeting efficiency (**Fig. 1D**).

Since the 10:1 ratio had the highest targeting efficiency, despite significantly less donor DNA, we

used this ratio in subsequent EPs. Because the targeting efficiency was notably high, we next

attempted a single electroporation with multiple vectors targeting the same locus, but containing

equimolar amounts of five different cassettes, while keeping the overall *px330_Rosa26_sgRNA*

vector to total donor vector ratio at 10:1. The five vectors consisted of: 1) a Cre/FLP responsive

mutant methionyl-tRNA synthetase (*RR6*) for selective labeling of newly synthesized proteins; 2)

a Cre/FLP responsive modified G-protein coupled receptor (*RR7*); 3) a Cre/FLP responsive

bicistronic reporter with H2B-TagBFP and sfGFP separated by a p2a element, and tdTomato

expressed in the subtractive population (cells that express FLP but not Cre) (*RR9*); 4) a Dre-

responsive tricistronic reporter with H2B-TagBFP, sfGFP, and synaptophysin-tdTomato

separated by p2a elements (*RR11*); and 5) a Dre-responsive bicistronic reporter with H2B-

TagBFP and sfGFP separated by a p2a element (*RR10*). We saw a 52% targeting efficiency and

successful targeting of all five cassettes at varying efficiencies (**Table 3**). Due to the lower

targeting efficiency of *RR7* (2%), we attempted another 10:1 electroporation using the *RR7* donor

vector alone and obtained a 63% targeting efficiency (genotyping results shown in **Fig. 1**). Thus,

our results show that we are able to target and recover as many as five *Rosa* alleles in a single

EP, significantly increasing efficiency and reducing costs toward intersectional mouse generation.

Oocyte targeting: Given the high efficiency of targeting, we also attempted to create a founder

line (*RR8*) through direct oocyte injection of the optimized *Rosa26* targeting vector with a Cre-

responsive calcium indicator, Twitch (52) (total cassette size without homology arms, 6.5kb),

using pro-nuclear injection of Cas9 protein (30 ng/µl), sgRNA (20 ng/µl), and double-stranded

DNA plasmid (2 ng/µl). A total of 1266 embryos were injected yielding 129 mice, of which 7

genotyped positive for the calcium indicator but only one targeted successfully (<1% targeting

efficiency of mice born), suggesting that direct oocyte injection is inefficient with this system under

the specified parameters.

Off-target analysis: Genomic sequences that are similar to the sgRNA used for targeted double

stranded breaks may cause unintended gene mutations or editing at off-target sites. To account

for this possibility, we predicted the off-target sites for each sgRNA using the crispr.mit.edu tool

and selected the top 5 sites for follow up. We PCR amplified and sequenced these loci from three

correctly targeted ES cell clones and did not detect any genetic changes (**Fig. 2**). While we cannot

rule out off-target effects in other loci, these data suggest that off-target effects are not prevalent

in this setting.

Mouse line derivation: The goal of our efforts was to produce freely available genetic tools that

delineate and access distinct populations for multi-faceted circuit mapping or functional

characterization studies. With this toolkit in hand, we collectively produced 11 intersectional or

Dre-responsive alleles available as either mouse lines (8) or ES cells (3) **(Table 2)**. To perturb

neuron function, we produced three lines that express metabotropic DREADD receptors, hM4D

(*RR1*), hM3D (*RR2*), and Rs-EGFP (*RR7*), and a fourth line expressing the ionotropic PSAM

receptor (*RR4*) (24,53,54). Two lines enable molecular characterization: one expresses the

EGFP-L10A fusion protein for ribosomal affinity purification and capture of translating mRNAs

(*RR3*) (55), while the second line expresses methionyl-tRNA synthetase (Met-RS) that

incorporates an artificial amino acid into nascent peptides (*RR6*) (13). One line enables ratiometric

calcium imaging via expression of Twitch2B (*RR8*). Our neuro-anatomical mapping lines

described below also incorporate a tagged histone (H2B), offering the possibility of chromatin

affinity isolations. We created four alleles of differing Cre, FLP, and Dre recombinase responsive

configurations that fluorescently mark distinct cellular compartments for unambiguous cell counts,

morphological characterizations, and projection mapping (*RR5, RR9, RR10, RR11)*. The

derivation of lines *RR5*, *RR6*, and *RR7* demonstrated that ES pluripotency was maintained, and

that germline transmission was not diminished in our Cas9-mediated single and multiplexed

electroporations after correctly targeted clones were selected and injected into blastocysts for

chimera generation.

Select mouse line characterization: Upon dual recombinase expression, the *RR5* tricistronic

multi-color reporter allele expresses three spectrally separated and modified fluorescent proteins

to highlight the nucleus in blue, fill the neuron in green, and emphasize pre-synaptic contacts in

red. At the site of dual recombinase expression, targeted cells brightly label with TagBFP, sfGFP,

and tdTomato where TagBFP fluorescence is constrained to the nucleus by an H2B fusion, sfGFP

is unmodified so that fluorescence is seen throughout the cell body, and tdTomato is fused to

synaptophysin (56) so that fluorescence is excluded from nuclear areas and primarily seen in

projection areas. Individual nuclei can be resolved using blue fluorescence and co-localization

with sfGFP-labeled somata.

We used *RR5* to evaluate the functional activity and specificity *in vivo* of our CRISPR/Cas9

approach to generating mouse lines in three distinct contexts; 1) germline recombinase

expression, 2) viral recombinase expression, and 3) combinatorial retrograde viral and germline

recombinase expression to target single gene defined neurons by a specific projection field. First,

to demonstrate genetically-restricted, germline expression of recombinases in the intersectional

RR5 line, we bred the *RR5* line to a double *Dopamine-Beta-Hydroxylase (DBH)^{p2aFLPo}*(51); *Bactin-*

Cre recombinase driver to express the tricistronic fluorescent cassette in *DBH*-positive

noradrenergic neurons in the brainstem **(Fig. 3 A-H)**. In both the locus coeruleus (**Fig. 3 A-D)** and

the A5 nucleus (**Fig. 3 E-H)** as well as all other noradrenergic nuclei (not shown), we could cleanly

resolve blue nuclei, green cells, and red puncta without the need for antibody enhancement.

Additionally, we bred an intersection of double *Vgat-Cre; Vglut2-FLPo* recombinase expression

to the *RR5* line and found the entopeduncular nucleus labeled green with local projections labeled

red and distal projections in the lateral habenula labeled red (DAPI was applied to help delineate

the target field, obscuring the genetic TagBFP signal) (**Fig. 3 J-K**).

Second, we verified viral expression of Cre and FLP recombinase where adult *RR5* mice were

stereotaxically injected with equal titer amounts of AAV viruses expressing Cre and FLP into the

dentate gyrus **(Fig. 4 A-H)**, amygdala (**Fig. 4 I-P**), and olfactory bulb (**Fig. 4 Q-T**). Clear

expression of all three fluorescent proteins were resolved at the site of injection to the appropriate

cellular locations. Third, defining properties of the subpopulations can be extended to include any

combination of gene expression or anatomical location and projection target when injection of

retrograde viral vectors is applied. To utilize this option, we genetically marked a neuronal subtype

defined by the partial overlap of a genetic selector and projection target by injecting canine

adenovirus 2 (CAV2)-Cre virus (57,58) into the basolateral amygdala of *DBH^{p2aFLPo}; RR5* mice

that express only FLP in all noradrenergic neurons. CAV2-Cre virus efficiently transduces axon

terminals, thus genetically marking presynaptic neurons. In this context, only noradrenergic

neurons expressing DBH that project to the amygdala express both Cre and FLP and the resulting

tricistronic fluorescent cassette (**Fig. 5 A**). We only observed recombination in the brainstem

noradrenergic nuclei and in the locus coeruleus, primarily on the ipsilateral side (**Fig. 5 B-E**) with

some sparse marking on the contralateral side (**Fig. 5 F-I**), in agreement with a prior study (59).

Overlapping red and green puncta (with no blue) could be seen in the injected amygdala,

suggesting that the marked neurons project to the amygdala, as expected given the nature of the

CAV2-Cre virus (**Fig. 5 J-M**). We also observed collateral projections to several additional areas

in the mid and forebrain, including the dorsal raphé (**Fig. 5 N-Q**), reticulotegmental pontine nuclei

(**Fig. 5 R-U**), dentate gyrus (**Fig. 5 V-Y**), and olfactory bulb (**Fig. 5 Z-CC**). This restriction by

retrograde selection was also achieved in other regions of the brain with a second viral vector.

Vglut2_Cre; RR5 mice were injected with retro-AAV-Ef1a-FLPo into the lateral hypothalamus

(**Fig. 6 A**) where tricistronic expression was clearly visualized in the cingulate gyrus (**Fig. 6 B-E**),

piriform cortex (**Fig. 6 F-**I), and medial habenula (**Fig. 6 J-M**), indicating these regions as pre-

synaptic inputs to the lateral hypothalamus. Together, these results are in agreeance with

previous connectivity studies (59).

Before *in vivo* characterization of the cell autonomous DREADD systems coded by *RR1* and *RR2*,

we first examined the ability of the *RR1 and RR2* lines to modulate *Tg(Dbh-cre)KH212Gsat*

(TgDBH-Cre)-defined NA neuron activity at the cellular level. We conducted whole-cell recordings

from P30-P60 locus coeruleus neurons from Cre-only responsive alleles of mouse lines. These

lines were developed by germline deletion of the FRT flanked stop cassettes by crossing to the

Bact_FLPe allele(60) (*RR1P, RR2P)* or using the previously published *RC::FPDi* (*RC::PDi),* each

bred to the *RC::FrePe*(61) (also previously bred to the *Bact_FLPe* mouse line to produce the Cre

only responsive *RC::epe)* line to express both GFP and a DREADD in *TgDBH-Cre*-defined cells

for acute brain slice visualization of DREADD expressing neurons. *RC::PDi*(49) was included as

it represents a similarly constructed hM4D intersectional allele, but lacks the modifications and

optimizations made in our targeting system (**Fig. 1 A**). For example*, RC::PDi* contains an

additional PGK Promoter and pA signal sequence flanking neomycin in the first stop cassette and

lacks the WPRE element and pA found in *RR1* and *RR2*. We hypothesized that the addition of a

Woodchuck Hepatitis Virus (WHV) Posttranscriptional Regulatory Element (WPRE) would

enhance DREADD effectiveness. After bath application of 10 µM CNO, for *RR2P; RC::epe* we

observed a change in depolarization from pre-CNO average = -62.26±1.53 to a post-CNO

average = -55.09±2.16 mV (p = 0.0034) (n=19 neurons across 3 mice). After bath application of

10 µM CNO, for *RR1P; RC::epe* we observed a change in hyperpolarization from pre-CNO

average = -56.98±3.50 mV to post-CNO average = -65.3137±3.60 mV (p = 0.046) (n=10 neurons

across 3 mice). After bath application of 10 µM CNO, for *RC::PDi; RC::epe* we observed a change

in hyperpolarization from pre-CNO average = -53.71±2.93 mV to post-CNO average = -

56.72±3.42 mV (p = 0.027) (n=10 neurons across 3 mice) (**Fig. 7 A-C**).

We net performed whole-body plethysmography to assess changes in respiratory function caused

by CNO-DREADD mediated perturbation of targeted neurons *in vivo*. *DBH-Cre* defined NA

neurons were inhibited by crossing mouse line *RR1P* to *Tg(Dbh-cre)KH212Gsat (TgDBH-Cre)*

mice. We observed no changes under room air conditions (**Fig. 8 A)**. However, under hypercapnic

conditions, we saw a reduction in ventilation (**Fig. 8 B)**, with decreases in V_T (-26.6%, p=0.0021,

Fig. 8 D), $\dot{V}_E$ (-26.8%, p=0.026, **Fig. 8 E)**, and $\dot{V}_E/\dot{V}_{O2}$ (-28.7%, p<0.0001, **Fig. 8 G)**. No significant

differences were observed in body temperature between experimental animals and sibling

controls (**Fig. 8 H)**. Notably, sibling controls showed no difference in respiratory parameters pre-

and post-CNO administration.

We then tested the applicability of the *RR2P* line *in vivo.* After CNO-DREADD mediated

stimulation of *DBH-Cre* defined neurons, we saw significant increases under room air ventilation

(**Fig. 9 B)** in V_f (+28.1%, p=0.0076, **Fig. 9 D)**, V_T (+43.3%, p=0.00077, **Fig. 9 E)**, $\dot{V}_E$ (+83.6%,

p<0.0001, **Fig. 9 F)**, and $\dot{V}_{O2}$ (+50.9%, p<0.0001, **Fig. 9 G)**. The increase in $\dot{V}_E$ was proportional

to the increase in $\dot{V}_{O2}$, however, so $\dot{V}_E/\dot{V}_{O2}$ remained constant **(Fig. 9 H)**. Under hypercapnic

conditions, as compared to pre-CNO values, $\dot{V}_{O2}$ was increased (+37.2%, p=0.0015, **Fig. 9 G)**,

resulting in a reduced $\dot{V}_E/\dot{V}_{O2}$ (-24.2%, p=0.04, **Fig. 9 H)**. We also observed a deficit in body

temperature in *DBH-Cre; RR2P* animals 30 minutes after the end of the assay at room

temperature (~1.5 hrs after CNO injection) as compared to sibling controls (**Fig. 9 I)**, but not

immediately at exit from the respiratory chamber, which is kept at a thermoneutral temperature,

30-32°C**.** Sibling controls showed no difference in respiratory parameters pre- or post-CNO

administration. The respiratory phenotypes seen here clearly demonstrate an acute and cell

autonomous involvement of the whole NA system in CO_2 chemosensitivity and baseline

metabolism while avoiding the confounds of developmental compensatory events or off-target

effects. Collectively, these data highlight how our intersectional lines can be used to functionally

assess dispersed and difficult to access neuronal populations, setting the stage for further

characterization and dissection of a variety of neuronal systems in a given measured outcome

through intersectional subdivision by genetic or viral methods.

Finally, we characterized a fourth line, produced by oocyte CRISPR-mediated recombination to

express the Twitch2B ratiometric calcium indicator upon Cre recombinase expression. Calcium

signaling was recorded in outer granulosa cells surrounding an oocyte taken from a germline

recombined (*RR8; Bact_Cre*) mouse line (**Fig. 10 A**). Upon application of 10µM ionomycin in the

presence of 10mM $CaCl_2$, the YFP/CFP ratio increased more than 3-fold (**Fig. 10 B**). The data

demonstrate how the targeting system here can be used to build conditional mouse lines that are

useful beyond the nervous system and throughout the body (62).

DISCUSSION

Vector design and optimization: Our tricistronic mouse line vector (*RR5*) was a complex vector

to design. Thus, the discussion of its construction applies to the other simpler vectors and best

describes choices made to optimize the vectors. Our initial goal was to create an intersectional

mouse strain expressing three fluorescent components to highlight the nucleus, whole neuron,

and presynaptic contacts, enabling facile study of cell counts, cellular morphology, and axonal

and projection targets (*RR5* consisting of H2B-TagBFP, sfGFP, and synaptophysin-tdTomato

separated by p2a elements) without the need for signal amplification using antibodies. However,

likely due to the high number of repetitive elements in the *Rosa26* 3' homology arm and the high

complexity and repetitive components of the intersectional cassette (i.e. two stop cassettes

consisting of multiple SV40 polyadenylation sequences, tdTomato protein dimer, double loxP and

FRT sites, and two p2a self-cleaving peptides separating the elements), we had no success in

stably subcloning the intersectional cassette into a low copy vector (p15a ori) containing the

original *Rosa26* homology arms (1.1kb 5' homology arm and 4.2 kb 3' homology arm) and DTA

negative selection gene (49). Thus, we sought to make a number of improvements on the prior

approaches used to build intersectional genetic mice. First, we sought to stabilize the vector by

shortening the homology arms and eliminating repetitive genomic elements found at the *Rosa26*

locus. Second, we removed the DTA negative selection terminal non-homology to eliminate the

need to validate intersectional cassettes separately before placing the cassette in the targeting

vector. Third, we reduced the size of the selection cassette. Prior intersectional constructs carried

an additional PGK promoter and BGHpA tail for neomycin expression. We eliminated both

promoter and BGHpA as the functions of these elements can easily be served by the CAG

promoter and SV40pA in the stop cassette. With smaller homology arms, it was possible that our

targeting efficiency would fall. Therefore, we introduced a CRISPR/Cas9 targeting strategy to

enhance homologous recombination in either ES cells or oocytes. Surprisingly, in ES cells, the

shorter homology arms performed equally well as the full-length homology arms (see discussion

below). The resulting strategy offered a simplified, smaller, and highly stable vector that remains

efficient at targeting the *Rosa26* locus. Through this optimization process we built a set of baseline

vectors that can be used to rapidly and inexpensively build any intersectional genetic targeting

allele by simply cloning in a cDNA of interest into one or both of the multiple cloning sites for both

intersectional and subtractive expression.

Embryonic stem cell electroporation: While CRISPR strategies are widespread for knockout

mutations and small deletions and insertions in mouse zygotes, consistent knock-in of large

targeting cassettes (>4kb) still remains a challenge with fewer but increasing successes (63–65).

To address this limitation, our new multiplexed methodology enabled us to: **1)** simplify our base

targeting vector to less than 5kb in total length; **2)** see successful targeting of added cassettes up

to 11kb; **3)** increase the rate of targeting by 5-10 fold over previous *Rosa26* targeting attempts

(that used a significantly longer 3' homology arm) under our ES cell strain and conditions; and **4)**

further reduce cost by co-electroporating five different targeting vectors in a single ES cell

electroporation that were easily resolved to produce new mouse lines. While targeting rates for

the *Rosa26* locus can vary greatly in the literature, we compare only to our own experiments, as

significant variability in targeting efficiency from lab to lab or facility to facility can arise from the

several factors in ES cell electroporation ranging from electroporation conditions (e.g. buffer ionic

strength, adjuvants, field strength, field duration, field shape, square wave vs exponential decay,

etc.) to culture conditions (ES cell strain and genetic background, feeder type, media composition,

inclusion of LIF, etc.). Notably, when we co-electroporated a CRISPR/Cas9 plasmid to facilitate

targeting, there was a proportional increase in targeting efficiency despite reduced amounts of

targeting vector (total DNA in the EP was capped at 18-20 µg per EP). At the highest efficiencies,

we could further subdivide the small amount of targeting vector across multiple plasmids, which

allowed us to introduce and recover as many as five vectors in a single EP. This multiplexed

approach significantly reduced the costs of mouse production via ES cells. With these goals met,

multiplexed ES cell targeting enables the rapid and high throughput production of intersectional

alleles that can be readily distributed throughout the mouse research community for further

studies.

Oocyte targeting: Oocyte injection was only attempted with one line, *RR8*. Less than 1% of the

embryos contained a properly targeted vector, which in our experience suggests that direct oocyte

injection is not efficient. Future optimization and the development of new technologies may

increase the efficiency of targeting for more rapid generation of transgenic mice; for example, a

study showed 10-20% targeting of 8-11kb cassettes to the *Rosa26* locus in oocytes using longer

homology arms, while another showed successful targeting of a 5.5kb cassette into the *Rosa26*

locus in rats with co-injection of two ssDNA oligos (65,66). As noted above, there can be

significant variability in targeting across labs based upon a variety of factors including background

strain (vs strain from which the homology arms are derived), the use of Cas9 mRNA or protein,

and site of injection (pro-nucleus vs cytoplasm, single vs 2-cell stage). Thus, our outcomes here

may reflect our core facility capabilities as much as inherent limitations in consistent Cas9

mediated oocyte targeting of large constructs. The successful outcomes here and in a prior

publication demonstrate the use of the system to develop lines capable of functional imaging (62).

Select mouse line characterization: Intersectional expression of fluorescent proteins (FPs) has

been used to great effect for anatomical characterizations of targeted neuron populations (61,67).

Characterizations aimed at counting or highlighting single cells or cell populations made with a

single FP can be made difficult, however, in cases where cell, axon, or dendrite density (neuropil)

is high. Usually, one or two FPs are deployed to either fill the cell or highlight specific features,

which necessitates the use of multiple reporter alleles to clearly observe cell number, morphology,

and projections. To ameliorate these issues, we sought to create a mouse line (*RR5*) that

highlights three specific features in an intersectionally defined neuron. Overall, this novel three-

color mouse line enables intersectional and simultaneous labeling of three subcellular

compartments. The polycistronic fluorescent labeling cassette gives unambiguous cell counts and

clear visualization of cell morphology, axonal projections, and synaptic contacts. The strength of

expression of all three colors is likely due to the use of p2a elements, rather than IRES elements,

which cause expression level reduction. Reduced expression from an IRES may stem from cryptic

splice elements in the IRES that could interact with the intron in the CAG promoter or splice

acceptors in some stop cassettes (68). Our use of P2A appears to bypass those issues as all

three fluorescent proteins could be readily visualized without additional enhancement as

compared to other approaches (35).

We also show that combinations of germline, viral, and retrograde viral expression of

recombinases in the *RR5* line can all be used to clearly map the shared innervation of multiple

regions by a small subset of genetic, projection, and/ or intersectionally defined neurons. In

contrast to prior intersectional studies using two genetic delimiters to define a subpopulation with

a single fluorescent protein (with or without an inert retrograde marker injected into separate sites

in the brain), our approach enables facile collateral mapping across the whole nervous system

with higher resolution that is growing in popularity (33,59,61). These results demonstrate that

despite the high complexity and repetitive nature of the *RR5* cassette, we could efficiently target

this allele, the allele remains stable through germline transmission, and it is responsive to Cre

and FLP recombinase expression. The restriction of expression to anatomically defined NA

neurons as a whole or as a projection subset further establishes the recombinase specificity of

our targeting schematic. This approach should also translate equally well to further anatomical,

functional, and molecular characterizations of NA and other subtypes defined by their projection

patterns with the use of the additional lines reported here.

Noradrenergic (NA) neurons are strongly implicated in control of respiratory homeostasis and

chemosensitivity (8,69–75), but many prior studies bear methodological caveats such as lack of

resolution due to overly broad lesions or injections (76), developmental and non-cell-autonomous

compensation after gene mutations, or restraint or anesthesia *in vivo* (77). Here we show in

conscious, unrestrained mice that inhibition of *TgDBH-Cre*-defined noradrenergic neurons led to

significantly reduced tidal volume and minute ventilation following hypercapnic exposure. On the

other hand, activation of *TgDBH-Cre*-defined noradrenergic neurons led to significantly increased

respiratory rate, tidal volume, and minute ventilation in the room air breathing condition. These

data represent modulation of the noradrenergic system that significantly deviates from previously

published studies interrogating this system and provides an alternative to those investigating

control of breathing that reduces the influence of possible side effects or unintended effects.

In conclusion, the methods presented here represent a significant improvement in feasibility and

affordability for labs to generate their own intersectional mouse models for investigating circuit

development. Our vector design and application allow for large cassettes to be targeted to the

Rosa26 locus with high throughput, increased efficiency, germline transmission, and preserved

pluripotency. These protocols and techniques can be used to investigate most, if not all, neural

circuits in the context of development and/ or disease. Implementing these techniques also serve

as an alternative to cost-prohibitive commercially available vectors and mouse lines. Together,

we present a facile, cost-effective method for producing gene specific intersectional mouse lines

for neural circuit investigation.

CONCLUSIONS

During development, many distinct neuronal and glial subtypes arise and intercalate to create the complex neural circuits that govern our behavior and physiology. Key to understanding this organization and how it may be perturbed in neurological diseases is the ability to identify and access discrete cell subtypes in the developing and adult mouse nervous systems for multi-faceted studies. It is now clear that even within narrowly defined cell types once thought as homogeneous, significant diversity is found at multiple levels including genetic and molecular signatures, activity patterns, synaptic connectivity, and projection patterns. Given this complexity, it is often difficult to define and access specific cellular populations for study, particularly during embryogenesis, where *in utero* development makes access problematic.

The power of studying combinatorially defined neuronal populations increases significantly when multiple distinct neuronal features, such as birthdate, collateral projection targets, molecular profiles, or a functional requirement in a given physiological process or behavior can be examined in parallel to reveal deeper insights into the mechanisms involved in the establishment and maintenance of neural circuit organization. However, the ability to carry out such studies is restricted by the limited number of publicly available intersectional mouse lines as well as the inherent difficulty in developing new intersectional alleles.

We present a novel CRISPR/Cas9-mediated system for consistently targeting large cassettes to the *Rosa26* locus for high throughput and parallel generation of intersectional and conditional mouse lines. These studies demonstrate that CRISPR/Cas9 can be used to increase targeting efficiency at the *Rosa26* locus in mouse embryonic stem cells with shortened 1-1.2 kb homology arms while preserving pluripotency and the ability to transmit alleles via the germline. This technology allows for the opportunity to generate mouse lines that express a wide range of effector molecules in neural cell types, enabling the developmental, anatomical, molecular, and

functional characterization of neural circuit organization in behavior and physiology. To

democratize the generation of intersectional and conditional targeted ES cells and mouse lines,

we have generated a publicly available vector toolkit for single-step cloning into *Rosa26*-targeting

vectors for double and single-recombinase responsive cassettes (**Table 1).** We have also

generated several mouse lines available, without restriction, to the not-for-profit research

community enabling high specificity intersectional access of neuronal and other populations for

cross correlative mapping approaches using functional neuronal perturbation, molecular profiling,

and anatomical characterizations for multifaceted studies (**Table 2**). Overall, these resources and

the modular nature of the intersectional approach make rapid and low-cost production of large

numbers of intersectional alleles possible that can be efficiently distributed as ES cells or mouse

lines throughout the mouse research community. All vectors (Addgene; 97007-97012, 99142),

mouse lines (MMRRC; 043513-043519), and ES cells (BCM ES Cell Core Facility) have been

made publicly and unconditionally available to the academic research community.

METHODS

Construction of targeting and Cas9 vectors: The simplified *Rosa26* targeting vector was

derived using standard cloning procedures from a targeting vector used in previous studies (49)

consisting of a p15a origin of replication, kanamycin selection cassette, DTA negative selection

cassette, and *Rosa26* homology (1081 bp 5' homology arm and 4342 bp 3' homology arm). The

total vector size was 9084 bps. The DTA was removed and the 5' homology arm was shortened

to 1025 bp fragment (deleting the sgRNA target sequence) while the 3' homology arm was

shortened to 1231 bps, reducing the total vector length to 4694 bps and eliminating a number of

repetitive regions that often resulted in vector instability when constructing intersectional targeting

vectors.

For the Cas9 expressing vector, we selected an sgRNA target sequence that was close to the 5'

and 3' junction of the *Rosa26* gene locus (ACTGGAGTTGCAGATCACGA with PAM motif GGG).

The selected sgRNA was cloned into the BbsI sites of the px330 vector that expresses Cas9 (final

vector is called *px330_Rosa26_sgRNA*).

Knock-in cassettes were assembled in either 1) a previously used intersectional template plasmid

(*RR1-4, RR6-7*); 2) a newly constructed intersectional template plasmid with neomycin promoter

and pA elements removed (*RR5* and *RR8*); or 3) a constructed Dre-responsive template plasmid

(*RR9* and *RR10*). The intersectional plasmid consisted of a ubiquitous *CAG* promoter sequence;

an *FRT*-flanked stop cassette consisting of a PGK-neo sequence (for positive selection of

targeted ES cells) and three *SV40 pA* sequences; a *LoxP*-flanked stop cassette containing

mCherry and a PBS302 stop cassette; and a cloning site for insertion followed by a *WPRE* and

BGHpA sequence. The new intersectional template consisted of a ubiquitous *CAG* promoter

sequence; an *FRT* flanked stop cassette consisting of a neomycin sequence and *His3_SV40 pA*;

a *LoxP*-flanked subtractive cloning site stop cassette consisting of a *His3_SV40 pA*; and a cloning

site for insertion followed by *WPRE* and *BGHpA* sequences. The Dre-responsive template

consisted of a *Rox*-flanked stop cassette consisting of a neomycin sequence and PBS302 stop

cassette, and a cloning site followed by a *WPRE* and *BGHpA* sequence.

For assembly of complete targeting vectors, the cDNAs of interest were PCR amplified and cloned

into the corresponding vector. The intersectional or Dre-responsive cassettes were then cut out

with PacI or PacI/AscI and cloned into the shortened *Rosa26* targeting vector.

To facilitate single step cloning of complete targeting vectors, we generated a vector toolkit (**Table**

1) by constructing and moving empty *CAG* cassettes in the Rosa26 targeting vector. We

generated four template vectors with neomycin resistance: 1) a Cre/FLP responsive targeting

vector; 2) Cre only; 3) FLP only; and 4) Dre-responsive targeting vector. The intersectional

template vector consists of the *CAG* promoter, an *FRT*-flanked neomycin resistance gene and

stop cassette, a subtractive cloning site (EcoRV) and stop cassette flanked by *LoxP* sites, an

intersectional cloning site (PmeI), and *WPRE* and *BHGpA* elements. Each single template vector

consists of the *CAG* promoter, stop cassette(s) and neomycin resistance cassette flanked by

recombinase sites, cloning site (SwaI), and *WPRE* and *bgh polyA* elements. Additionally, for

possible direct oocyte injections we also generated a Cre/FLP responsive targeting vector that

does not contain a neomycin resistance gene, with a subtractive cloning site (EcoRV) and

intersectional cloning site (SwaI). All cloning sites are blunt restriction enzyme sites for cloning

facilitation.

Sequences of the template vectors and *px330* vector and plasmids are available through Addgene

(provisional ID #s 97007-97012, 99142).

Generation of knock-in mice: Embryonic stem (ES) cells (AB2.2) were electroporated with 15-

20 µg of varying ratios of the *px330_Rosa26_sgRNA* vector to the targeting vector. Neomycin

selected clones were screened for homologous recombination using PCR genotyping. Targeted

clones were identified using PCR genotyping for 5' and 3' targeting from outside the homology

arm and were considered to be successful targeting events if positive for both 5' and 3'

genotyping. We used pairs 5'CGCCTAAAGAAGAGGCTGTG (Rosa26-F) and

5'GGGCGTACTTGGCATATGAT (CAG-R), producing a 1450 bp band; and

5'AATCAACCTCTGGATTACAAAATTT (WPRE-F) and 5'TGGCTCCTCTGTCCACAGTT

(Rosa26-R), producing a 2472 bp band.

Select targeted clones were microinjected into C57B1/J6 blastocysts and chimeric males were

bred to wildtype C57B1/J6 females to achieve germline transmission.

Pronuclear injection: We constructed a targeting vector by cloning a calcium indicator, Twitch-

2B (52), into the Cre responsive *Rosa26* targeting vector. The Baylor College of Medicine

Embryonic Stem (ES) Cell Core generated the *Rosa26*-specific sgRNA and Cas9 protein for

injection. Pronuclear injections were performed by the Baylor College of Medicine Genetically

Engineered Mouse (GEM) Core using the following parameters: 30 ng/µl Cas9 protein, 20 ng/µl

sgRNA, and 2 ng/µl dsDNA plasmid. Potential founders were screened for targeting as described

above.

Breeding, genetic background, and maintenance of mice: We maintained colonies of select

mouse strains by backcrossing to C57BL/6J mice. For routine genotyping, we carried out PCR

amplification of DNA from ear punch preparations using the boiling alkaline lysis procedure.

Rosa26 specific primers for the mice were 5'-GCACTTGCTCTCCCAAAGTC, 5'-

GGGCGTACTTGGCATATGAT, and 5'-CTTTAAGCCTGCCCAGAAGA, and yield a 495 bp band

(targeted) and 330 bp band (wt). For histology experiments, *RR5* mice were bred to

DBH^{p2aFLPo}(51)*; B6N.FVB-Tmem163$^{Tg(ACTB-cre)2Mrt}$/CjDswJ (Bactin-Cre)* and *DBH^{p2aFLPo}* mice. For

plethysmography experiments, *B6.FVB(Cg)-Tg(Dbh-Cre)KH212Gsat/Mmucd (DBH-Cre)* mice

were mated with homozygous Cre-only responsive *RR1P* and *RR2P* mice (after crossing

intersectional alleles to *B6;SJL-Tg(ACTFLPe)9205Dym/J* [JAX 003800] mice to recombine out

the FLP-responsive stop cassette followed by homozygosity) to derive animals in which all mice

carried the Cre-responsive hM4D or hM3D allele. Sibling animals that did not inherit the *Cre* allele

were used as control animals to the *Cre* positive offspring. For electrophysiology experiments,

DBH-Cre; RR2P mice were mated to *RC::ePe* mice expressing a floxed eGFP. Cre-specific

primers were 5'-ATCGCCATCTTCCAGCAGGCGCACCATTGCCC and 5'-

GCATTTCTGGGGATTGCTTA and yielded a 550 bp band if positive. FLPo-specific primers were

5'CACGCCCAGGTACTTGTTCT and 5'CCACAGCAAGAAGATGCTGA and yielded a 226 bp

band if positive.

Established mouse lines reside at the Mutant Mouse Regional Resource Center (MMRRC) and

will be unconditionally available to the academic research community (MMRRC 043513-043519).

Targeted ES cells have been archived and are available upon request under the condition that

any new mouse lines are deposited in a public repository.

All animal experiments were performed with the approval of IACUC.

Off-target analysis: Off-target sequences were identified using the Optimized CRISPR Design

tool (crispr.mit.edu). The top 5 sequences for each sgRNA were amplified from selected targeted

ES cells and sequenced to determine if any mutations or changes occurred in off-target sites.

Viral injection: To test the functionality of the *RR5* mouse strain, adult *RR5* mice were injected

with equal titers of AAV9-hSyn-FLPo and AAV1-hSyn-Cre viruses (obtained from M. Xue at BCM,

250 nl at 1.12×10^{12} GC/mL) into the dentate gyrus (coordinates from bregma AP=-2.70, DV=-

2.12, ML=1.84) and with equal titers of AAV-EF1a-Cre-WPRE and AAV-EF1a-FLPo-WPRE (UNC

Vector Core, 250 nl at 5.06×10^{11} GC/mL) into the amygdala (coordinates from bregma AP=-1.06,

DV=-4.61, ML=-2.86) and allowed to incubate for 7-14 days. For expression in the olfactory bulb,

adult *RR5* mice were injected with AAV-CAG-Cre (obtained from Neuroconnectivity Core at Jan

and Dan Duncan Neurological Research Institute, AAV2/9 690 μL at 5.37x10^{13} pp/mL) and AAV-

Ef1a-Flp (obtained from Neuroconnectivity Core at Jan and Dan Duncan Neurological Research

Institute, AAV2/9 690 μL at 7.16x10^{11} pp/mL) into the core of the olfactory bulb (coordinates from

bregma AP=4.5mm, DV=-2.25mm, ML=$\pm$0.8mm) and allowed to incubate for 14 days. For the

CAV2-Cre virus experiments, *RR5; DBH^{p2aFLP}* mice were injected with CAV2-Cre virus (IGMM

Viral Core, 500 nL at 6.0x10^{12} pp/mL) into the amygdala (coordinates -2.86, -1.06, -4.61) and

allowed to incubate for 17 days. For the Retro-AAV-Ef1a-FLPo virus experiments, *RR5;*

Vglut2_Cre mice were injected with Retro-FLPo virus (obtained from Neuroconnectivity Core at

Jan and Dan Duncan Neurological Research Institute, 690 μL at 1.49x10^{12} pp/mL) into the lateral

hypothalamic area (coordinates from bregma AP=-1.22mm, DV=-5.12mm, ML=$\pm$0.97mm) and

allowed to incubate for 21 days.

Histology: For *RR5* expression, animals were sacrificed and transcardially perfused with 0.1M

phosphate-buffered saline (PBS) then with 4% paraformaldehyde (PFA) in PBS. Brains were

dissected out and drop fixed for two hours in 4% PFA before a PBS rinse and equilibration in 20%

sucrose in PBS. Brains were sectioned into 30-40 μm coronal sections and mounted on slides.

Images were collected on a Zeiss confocal LSM780 microscope or Leica TCS SPE confocal

microscope.

Electrophysiology

Slice preparation

Slice preparation from the mouse LC follows an N-Methyl-D-glucamine (NMDG) slicing protocol

(78,79). Briefly, animals were deeply anesthetized using 3% isoflurane. After decapitation, the

brain was removed and placed into cold (0−4 °C) oxygenated NMDG solution containing 93 mM

NMDG, 93 mM HCl, 2.5 mM KCl, 1.2 mM NaH$_2$PO$_4$, 30 mM NaHCO$_3$, 20 mM HEPES, 25 mM

glucose, 5 mM sodium ascorbate, 2 mM Thiourea, 3 mM sodium pyruvate, 10mM MgSO$_4$ and 0.5

581 mM $CaCl_2$, pH 7.35 (all from SIGMA-ALDRICH). Horizontal slices were prepared using a

582 vibratome (200 μm thick) using zirconia blades. The brain slices were kept at 37.0 ± 0.5 °C in

583 oxygenated NMDG solution for 10 minutes. They were then transferred to an artificial

584 cerebrospinal fluid (ACSF) containing 125 mM NaCl, 2.5 mM KCl, 1.25 mM NaH_2PO_4, 25 mM

585 $NaHCO_3$, 1 mM $MgCl_2$, 25 mM glucose, and 2 mM $CaCl_2$ (pH 7.4) for at least 1 hour prior to the

586 beginning of recordings. During the recording sessions, the slices were submerged in a custom

587 chamber and were stabilized with a fine nylon net attached to a custom-designed platinum ring.

588 This recording chamber was continuously perfused with oxygenated physiological solution

589 throughout the recording session.

590 ***Recordings***

591 Whole-cell recordings were performed as described previously (79–81). Briefly, patch pipettes (2-

592 8 MΩ) were filled with an internal solution containing 120 mM potassium gluconate, 10 mM

593 HEPES, 4 mM KCl, 4 mM MgATP, 0.3 mM Na_3GTP, 10 mM sodium phosphocreatine and 0.5%

594 biocytin (pH 7.25). Whole-cell recordings from up to 8 LC neurons were performed using two

595 Quadro EPC 10 amplifiers (HEKA Electronic, Germany). PatchMaster (HEKA) and custom-written

596 Matlab-based programs (Mathworks) were used to operate the recording system and perform

597 online and offline data analysis. In current-clamp recordings, neurons were first current clamped

598 at ~-40pA to prevent spontaneous firing. Prior to investigating the effect of drugs, we calculated

599 spike thresholds and recorded firing patterns in response to sustained depolarizing currents by

600 injecting increasing current steps (+10pA). Continuous recordings were obtained from LC neurons

601 current clamped at -40 to 0 pA during drug wash-on experiments. We also calculated other

602 intrinsic electrophysiological parameters, such as the input resistance, membrane time constant,

603 spike amplitude, after-hyperpolarization (AHP) etc. (79–81).

604 **Plethysmography**

Plethysmography on conscious, unrestrained mice was carried out as described on 6-12 week

old adult animals (49,82). Mice were subjected to a five-day habituation protocol with each day

consisting of several minutes of handling, temperature taken by rectal probe, intraperitoneal saline

injection, and 30 minutes in the plethysmography chamber. Mice were then tested within one

week of the last day of conditioning.

On the day of testing, mice were taken from their home cage, weighed, and rectal temperature

was taken. Animals were then placed into an airtight, temperature controlled (~32°C)

plethysmography chamber and allowed to acclimate for at least 20 minutes in room air (21%

O_2/79% N_2) conditions. After acclimation and measurement under room air, the chamber gas was

switched to a hypercapnic mixture of 5% CO_2/21% O_2/74% N_2 for 20 minutes. Chamber gas was

then switched back to room air for 20 minutes. The mice were briefly removed for rectal

temperature measurement and intra-peritoneal injection of clozapine-N-oxide (CNO, National

Institute of Mental Health Chemical Synthesis and Drug Supply Program) dissolved in saline (0.1

mg/mL) for an effective concentration 1 mg/kg. The animal was returned to the chamber for

another 20 minutes of room air, 20 minutes of hypercapnia, and 20 minutes of room air. The

animal was then removed from the chamber and rectal temperature was taken immediately

afterwards and again 30 minutes after the termination of the experiment. The animal was placed

in its own cage during these 30 minutes at the ambient room temperature (~23°C).

Plethysmography data collection and analysis

Plethysmography pressure changes were measured using a Validyne DP45 differential pressure

transducer and CD15 carrier demodulator in comparison to a reference chamber and recorded

with LabChartPro in real time. Waveforms were analyzed offline using LabChart Pro and custom

written MATLAB code to determine respiratory rate (V_f), tidal volume (V_T) (49), minute ventilation

($\dot{V}_E$), and pattern analysis. Respiratory waveforms were collected offline during periods when the

animal was at rest and readings were free from movement artifacts. A minimum of 1-minute

cumulative data compiled from traces at least 10 seconds long from the last 10 minutes of a given

experimental condition were analyzed. O_2 consumption was determined by comparing the gas

composition between calibration in an empty chamber and live breathing using an AEI oxygen

sensor and analyzer. Chamber temperature was constantly monitored and recorded using a

ThermoWorks MicroThermo 2 and probe and was recorded with LabChartPro in real time.

Plethysmography statistics

Results (V_f, V_T, $\dot{V}_E$, $\dot{V}_{O2}$, $\dot{V}_E/\dot{V}_{O2}$) for room air and hypercapnic data were compared between all

cohorts using a linear mixed-effects regression model with animal type (experimental or control)

and CNO administration (pre- or post-injection) as fixed effects and animal ID as a random effect.

Temperature data was compared using a linear mixed-effects regression model with animal type

(experimental or control) as a fixed effect. The p-values reported correspond to statistical

significance of the conditional interaction between animal type and CNO administration. A p-value

of <0.05 was used to indicate statistical significance and standard error of the mean is shown on

all charts.

Imaging of Twitch 2 in mouse preovulatory follicles

Preovulatory follicles were isolated and imaged as previously described (83). Briefly, antral

follicles were dissected from 23- to 26-d-old *RR8; Bact_Cre mice*. Follicles were cultured for 24–

30 h on organotypic membranes (Millipore) in the presence of follicle-stimulating hormone. The

follicle was held in a perfusion slide consisting of a plastic slide (ibidi) and a glass coverslip and

assembled using silicon grease. The slide was constructed such that medium containing ovine

LH (National Hormone and Peptide Program; 10 µg/mL) could be perfused through a 200-µm-

deep channel holding the follicle. Temperature was maintained at 30–34 °C, by use of a warm air

blower (Nevtek). Preovulatory Follicles were imaged using a Zeiss Pascal confocal microscope

with a 40X/1.2 NA objective. Images were collected every 10 sec. Measurements were corrected

for autofluorescence and for spectral bleed-through of CFP into the YFP channel. Ratios were

calculated by dividing the mean CFP intensity in each region of interest by the mean YFP intensity.

Data analysis was done using ImageJ and Excel software. Data is representative of 4 follicles.

Declarations

Ethics approval and consent to participate

All experiments reported herein were conducted with explicit approval of and oversight by both Baylor College of Medicine Institutional Animal Care and Use Committee (IACUC) and University of Connecticut IACUC and abide by all state and national regulations regarding animal research for each work site.

Consent for publication

Not applicable.

Availability of data and materials

All datasets, animals, and materials generated and/ or used in this study are publicly available (Addgene and Mutant Mouse Resource and Research Centers Supported by NIH (MMRRC)) or available from the corresponding author on reasonable request.

Competing interests

The authors declare no competing interests.

Funding

R01: H1130249 (RR)

R21: OD025327 (RR)

R37: HD014939 (JRE and Laurinda A. Jaffe)

Authors' contributions

SJL completed experiments, analyzed data, and prepared the manuscript.

AM completed experiments, analyzed data, and prepared the manuscript.

JJS completed experiments, analyzed data, and prepared the manuscript.

PJH completed experiments, analyzed data, and prepared the manuscript.

PGF completed experiments, analyzed data and prepared the manuscript.

JP completed experiments, analyzed data, and prepared the manuscript.

VKM assisted in the completion of experiments.

JRE completed experiments, analyzed data, and prepared the manuscript.

PJZ completed experiments, analyzed data and prepared the manuscript

XJ completed experiments, analyzed data, and prepared the manuscript.

BRA completed experiments, analyzed data, and prepared the manuscript.

AST completed experiments, analyzed data and prepared the manuscript

M C-M assisted with experimental design and prepared the manuscript

RR conceptualized the study, performed experiments, analyzed data, and prepared the

manuscript.

All authors read and approved the final manuscript.

Acknowledgements

We thank M. Xue for providing the Cre and FLPo AAV viruses used in the dentate gyrus injection

and Ronda Kram for technical assistance with cloning, mouse husbandry, and maintenance. We

thank J. Dougherty, E. Schuman, J. Wess, B. Roth, and S. Sternson for providing plasmid

templates for the EGFP-L10A, MetRS, G_s-D, hM4D and hM3D, and PSAM cDNA, respectively.

We thank Laurinda A. Jaffe for collaborating with us and author JRE with the Twitch studies. The

Baylor College of Medicine Embryonic Stem (ES) Cell Core performed the ES cell

electroporations and were exceptionally helpful in our experiments and the Baylor College of

Medicine Genetically Engineered Mouse (GEM) Core performed the blastocyst injections.

Imaging was carried out at the Baylor College of Medicine Optical Imaging and Vital Microscopy

Core.

703 **Table 1. Publicly available modular targeting vectors for rapid generation of recombinase-**

704 **responsive ES cells and mouse lines.**

705   **Table 2. *Rosa26* knock-in alleles generated with or without CRISPR/Cas9 methods.** Shown

706   are targeted *Rosa26* alleles, function, length of cassette, current status, and expected MGI name

707   (to be determined) in grey rows.

708   **Table 3. Targeting efficiencies for a multiplexed five cassette ES cell electroporation for**

709   **the *Rosa26* locus.**

Figure 1. Generation of intersectional *Rosa26* mouse lines. **A)** Targeting schematic showing the modular targeting vector containing a 1kb 5' homology arm, *CAG* promoter, *FRT*-flanked neomycin and stop cassette, *LoxP*-flanked (optional) subtractive gene of interest (GOI-S) and stop cassette, intersectional gene of interest (GOI-I), *WPRE*, *bgh poly A* element, and 1.2kb 3' homology arm. The full intersectional cassette was knocked into the *Rosa26* locus, with a 22 bp deletion of the CRISPR sgRNA. **B)** PCR genotyping of neomycin selected ES cell clones. Targeting knock-in was determined using PCR primers that spanned from outside the *Rosa26* homology arms to either the *CAG* promoter (5' end) or *WPRE* (3' end). Amplification of a band indicates targeting. Shown are results from a targeting event with over 60% targeting efficiency (*RR7*). **C)** Four conditions with different Cas9:targeting vector ratios were used in our initial study: a 0:1 with no Cas9, 0.5:1, 1:1, and 10:1 **D)** Targeting efficiency results from the different ratios. The 10:1 Cas9:targeting vector ratio showed a 5-10 fold increase over an electroporation with no Cas9. Shown in red are results from previous electroporations using the traditional Rosa26 targeting vector with more commonly used longer homology arms.

Figure 2. Analysis of *Rosa26* sgRNA off-target sites. No mutations were seen in the top 5

potential off-target sites. **A)** List of top 10 potential off-target sites as determined by the Optimized

CRISPR tool, base pair mismatches, and location in the genome. **B-F)** Sequence chromatograms

of each off-target site, showing the correct sequence for each of the 3 selected clones that were

injected into blastocysts as compared to the wildtype sequence.

Figure 3. Germline Cre/FLP expression in *RR5* adult mice to fluorescently label three

cellular compartments in targeted cells. A-H) *RR5* mice bred to *DBH^{p2aFLP}; Bactin-Cre* mice

express Cre ubiquitously and FLP in all noradrenergic (NA) neurons, thus marking only NA

neurons, including the locus coeruleus (**A-D**) and A5 nucleus (**E-H**). **I-P)** *RR5* mice bred to

Vgat_Cre; Vglut2-FLPo mice with dual recombinase expression imaged in the entopeduncular

nucleus (**I-L**) and lateral habenula (**M-P**). In targeted cells that express both Cre and FLP, cell

nuclei are labeled with TagBFP, cell soma and processes are labeled with sfGFP, and pre-

synaptic contacts are labeled with tdTomato.

Figure 4. Viral-mediated expression of Cre or FLP recombinase in *RR5* adult mice to

simultaneously, fluorescently label three cellular components. *RR5* mice injected with

equal titers of AAV-Cre and AAV-FLP viruses into the dentate gyrus (**A-H**), basolateral

amygdala (**I-P**), and olfactory bulb (**Q-T**).

Figure 5. Retrograde-viral Cre and germline FLP mediated expression of dual

recombinases for tricistronic FP expression on *RR5* background. A) *RR5; DBH^{p2aFLPo}* mice

were injected with CAV2-Cre virus into the basolateral amygdala. All *DBH* noradrenergic neurons

express FLP, but only those noradrenergic neurons projecting to the injected amygdala will also

express Cre. Marked double Cre/FLP positive cells will express H2B-TagBFP highlighting the

nucleus in blue, sfGFP filling the cell including the axon, and synaptophysin-tdTomato labeling

presynaptic contacts. BLA: Basolateral amygdala. DG: Dentate gyrus. DR: Dorsal raphé. LC:

Locus coeruleus. OB: Olfactory bulb. RtTg: Reticulotegmental pontine nucleus. *DBH* positive

neurons that project to the amygdala arise primarily from the ipsilateral locus coeruleus (**B-E**) with

some sparse labeling in the contralateral locus coeruleus (**F-I**). Red puncta overlapping with green

but lacking blue marked nuclei (indicative of projections from the marked population) can be seen

in a variety of mid- and forebrain areas, including the injected amygdala (**J-M**), the raphé nucleus

(**N-Q**), reticulotegmental pontine nuclei (**R-U**), dentate gyrus (**V-Y**), and olfactory bulb (**Z-CC**).

Figure 6. Retrograde-viral FLP and germline Cre mediated expression of dual

recombinases for tricistronic FP expression on *RR5* background. A) *RR5; Vglut2^{Cre}* mice

were injected with retro-AAV-Ef1a-FLPo virus into the lateral hypothalamus. All *Vglut2* neurons

express Cre, but only those Vglut2 neurons projecting to the injected LHA will also express FLP.

Marked double Cre/FLP positive cells will express H2B-TagBFP highlighting the nucleus in blue,

sfGFP filling the cell including the axon, and synaptophysin-tdTomato labeling presynaptic

contacts. LHA: Lateral hypothalamic area. Vglut2 positive neurons that project to the LHA arise

from the cingulate gyrus (**B-E**), piriform cortex (**F-I**), and medial habenula (**J-M**).

Figure 7. Electrophysiological characterization of CNO-DREADD mediated responses in

lines *RR1(P)* *RR2(P)* and *RC::FPDi* *(P)* in noradrenergic *DBH-Cre* Locus Coeruleus

neurons. **(A)** Representative wash-on responses for each *RR1(P) (n=19*

neurons), RR2(P) (n=10 neurons), and *RC::FPDi*(P) *(n=10 neurons)* in response to DREADD

agonists*. **(B)** Individual membrane potential plots of pre-CNO vs post-CNO values. **(C)** Summary

plots of absolute voltage change for each DREADD line tested following CNO treatment. *p<0.05;

**p<0.01; ***p<0.001.

Figure 8. CNO-hM4D mediated perturbation of noradrenergic *DBH-Cre* neurons results in a reduced hypercapnic response. A) Representative breathing traces from a *DBH-Cre; RR1P* animal before and after CNO administration under hypercapnic conditions ($5\%CO_2/21\%O_2/74\%N_2$). **B-F)** Quantification of respiratory and metabolic parameters under room air and hypercapnic conditions in *DBH-Cre; RR1P* animals and sibling controls. Measured values include respiratory rate (**B**), tidal volume (**C**), minute ventilation **(D)**, oxygen consumption **(E)**, and minute ventilation normalized to oxygen consumption **(G)**. No difference was seen under room air conditions but *DBH-Cre; RR1P* animals showed a deficit in volume, minute ventilation, and minute ventilation normalized to oxygen consumption under hypercapnic conditions. **G)** No difference in temperature was seen between *DBH-Cre; RR1P* animals and sibling controls. $*p<0.05$; $**p<0.01$; $***p<0.001$.

Figure 9. CNO-hM3D mediated perturbation of noradrenergic *DBH-Cre* neurons results in

enhanced room air ventilation and a reduced hypercapnic response. A) Representative

breathing traces from a *DBH-Cre; RR2P* animal before and after CNO administration under

room air conditions ($21\%O_2/79\%N_2$). **B)** Representative breathing traces from a *DBH-Cre;*

RR2P animal before and after CNO administration under hypercapnic conditions

($5\%CO_2/21\%O_2/74\%N_2$). **C-G)** Quantification of respiratory and metabolic parameters under

room air and hypercapnic conditions in *DBH-Cre; RR2P* animals and sibling controls. Measured

values include respiratory rate (**C**), tidal volume (**D**), minute ventilation **(E)**, oxygen consumption

(F), and minute ventilation normalized to oxygen consumption **(G)**. After CNO administration,

DBH-Cre; RR2P animals showed increase rate, volume, minute ventilation, and oxygen

consumption under room air conditions. Under hypercapnic conditions, *DBH-Cre; RR2P* animals

showed increased oxygen consumption resulting in a reduced minute ventilation to oxygen

consumption value. **I)** *DBH-Cre; RR2P* animals showed a significant deficit in temperature 30

minutes after the end of the assay. $*p<0.05$; $**p<0.01$; $***p<0.001$.

Figure 10. Expression and function of Twitch-2B in the granulosa cells of mouse ovarian

follicles. (**A**) A follicle from a mouse expressing Twitch-2B (homozygote). A follicle consists of

multiple layers of granulosa cells surrounding an oocyte in the center. Twitch-2B is uniformly

expressed in the granulosa cells but is not detectable in the oocyte. A layer of theca cells,

adhering to the outside of the follicle, expresses Twitch-2B at a higher level. Scale bar = 50 μm.

(**B**) ~6-fold increase in YFP/CFP ratio in the outer granulosa cells after application of 10 μM

ionomycin in the presence of 10 mM $CaCl_2$. Representative of 4 follicles.

REFERENCES

1. Armbruster BN, Li X, Pausch MH, Herlitze S, Roth BL. Evolving the lock to fit the key to create a family of G protein-coupled receptors potently activated by an inert ligand. Proc Natl Acad Sci USA. 2007 Mar 20;104(12):5163–8.

2. Sauer B. Manipulation of transgenes by site-specific recombination: Use of cre recombinase. Guide to techniques in mouse development. Elsevier; 1993. p. 890–900.

3. Chuong AS, Miri ML, Busskamp V, Matthews GAC, Acker LC, Sørensen AT, et al. Noninvasive optical inhibition with a red-shifted microbial rhodopsin. Nat Neurosci. 2014 Aug;17(8):1123–9.

4. Rowitch DH, S-Jacques B, Lee SM, Flax JD, Snyder EY, McMahon AP. Sonic hedgehog regulates proliferation and inhibits differentiation of CNS precursor cells. J Neurosci. 1999 Oct 15;19(20):8954–65.

5. Urlinger S, Baron U, Thellmann M, Hasan MT, Bujard H, Hillen W. Exploring the sequence space for tetracycline-dependent transcriptional activators: novel mutations yield expanded range and sensitivity. Proc Natl Acad Sci USA. 2000 Jul 5;97(14):7963–8.

6. Sander JD, Joung JK. CRISPR-Cas systems for editing, regulating and targeting genomes. Nat Biotechnol. 2014 Apr;32(4):347–55.

7. Platt RJ, Chen S, Zhou Y, Yim MJ, Swiech L, Kempton HR, et al. CRISPR-Cas9 knockin mice for genome editing and cancer modeling. Cell. 2014 Oct 9;159(2):440–55.

8. Dubreuil V, Ramanantsoa N, Trochet D, Vaubourg V, Amiel J, Gallego J, et al. A human mutation in Phox2b causes lack of CO_2 chemosensitivity, fatal central apnea, and specific loss of parafacial neurons. Proc Natl Acad Sci USA. 2008 Jan 22;105(3):1067–72.

9. Zucker SN, Bancroft TA, Place DE, Des Soye B, Bagati A, Berezney R. A dominant negative Cx43 mutant differentially affects tumorigenic and invasive properties in human metastatic melanoma cells. J Cell Physiol. 2013 Apr;228(4):853–9.

10. LeMaster AM, Krimm RF, Davis BM, Noel T, Forbes ME, Johnson JE, et al. Overexpression of brain-derived neurotrophic factor enhances sensory innervation and selectively increases neuron number. J Neurosci. 1999 Jul 15;19(14):5919–31.

11. Collins AL, Levenson JM, Vilaythong AP, Richman R, Armstrong DL, Noebels JL, et al. Mild overexpression of MeCP2 causes a progressive neurological disorder in mice. Hum Mol Genet. 2004 Nov 1;13(21):2679–89.

12. Ellis T, Smyth I, Riley E, Bowles J, Adolphe C, Rothnagel JA, et al. Overexpression of Sonic Hedgehog suppresses embryonic hair follicle morphogenesis. Dev Biol. 2003 Nov 15;263(2):203–15.

13. Ngo JT, Champion JA, Mahdavi A, Tanrikulu IC, Beatty KE, Connor RE, et al. Cell-selective metabolic labeling of proteins. Nat Chem Biol. 2009 Oct;5(10):715–7.

14. Doyle JP, Dougherty JD, Heiman M, Schmidt EF, Stevens TR, Ma G, et al. Application of a translational profiling approach for the comparative analysis of CNS cell types. Cell. 2008 Nov 14;135(4):749–62.

15. Marvin JS, Borghuis BG, Tian L, Cichon J, Harnett MT, Akerboom J, et al. An optimized fluorescent probe for visualizing glutamate neurotransmission. Nat Methods. 2013 Feb;10(2):162–70.

16. Akemann W, Mutoh H, Perron A, Park YK, Iwamoto Y, Knöpfel T. Imaging neural circuit dynamics with a voltage-sensitive fluorescent protein. J Neurophysiol. 2012 Oct;108(8):2323–37.

17. Lin C-W, Sim S, Ainsworth A, Okada M, Kelsch W, Lois C. Genetically increased cell-intrinsic excitability enhances neuronal integration into adult brain circuits. Neuron. 2010 Jan 14;65(1):32–9.

18. Johns DC, Marx R, Mains RE, O'Rourke B, Marbán E. Inducible genetic suppression of neuronal excitability. J Neurosci. 1999 Mar 1;19(5):1691–7.

19. Ibañez-Tallon I, Wen H, Miwa JM, Xing J, Tekinay AB, Ono F, et al. Tethering naturally occurring peptide toxins for cell-autonomous modulation of ion channels and receptors in vivo. Neuron. 2004 Aug 5;43(3):305–11.

852   20.   Duan B, Cheng L, Bourane S, Britz O, Padilla C, Garcia-Campmany L, et al. Identification of
853   spinal circuits transmitting and gating mechanical pain. Cell. 2014 Dec 4;159(6):1417–32.

854   21.   Kim JC, Cook MN, Carey MR, Shen C, Regehr WG, Dymecki SM. Linking genetically defined
855   neurons to behavior through a broadly applicable silencing allele. Neuron. 2009 Aug 13;63(3):305–
856   15.

857   22.   Conklin BR, Hsiao EC, Claeysen S, Dumuis A, Srinivasan S, Forsayeth JR, et al. Engineering
858   GPCR signaling pathways with RASSLs. Nat Methods. 2008 Aug;5(8):673–8.

859   23.   Beier KT, Samson MES, Matsuda T, Cepko CL. Conditional expression of the TVA receptor
860   allows clonal analysis of descendents from Cre-expressing progenitor cells. Dev Biol. 2011 May
861   15;353(2):309–20.

862   24.   Magnus CJ, Lee PH, Atasoy D, Su HH, Looger LL, Sternson SM. Chemical and genetic
863   engineering of selective ion channel-ligand interactions. Science. 2011 Sep 2;333(6047):1292–6.

864   25.   Vardy E, Robinson JE, Li C, Olsen RHJ, DiBerto JF, Giguere PM, et al. A new DREADD
865   facilitates the multiplexed chemogenetic interrogation of behavior. Neuron. 2015 May
866   20;86(4):936–46.

867   26.   Zhao R, Grunke SD, Keralapurath MM, Yetman MJ, Lam A, Lee T-C, et al. Impaired Recall of
868   Positional Memory following Chemogenetic Disruption of Place Field Stability. Cell Rep. 2016 Jul
869   19;16(3):793–804.

870   27.   Beier KT, Saunders A, Oldenburg IA, Miyamichi K, Akhtar N, Luo L, et al. Anterograde or
871   retrograde transsynaptic labeling of CNS neurons with vesicular stomatitis virus vectors. Proc Natl
872   Acad Sci USA. 2011 Sep 13;108(37):15414–9.

873   28.   Wickersham IR, Finke S, Conzelmann K-K, Callaway EM. Retrograde neuronal tracing with a
874   deletion-mutant rabies virus. Nat Methods. 2007 Jan;4(1):47–9.

875   29.   Zeisel A, Muñoz-Manchado AB, Codeluppi S, Lönnerberg P, La Manno G, Juréus A, et al. Brain
876   structure. Cell types in the mouse cortex and hippocampus revealed by single-cell RNA-seq.
877   Science. 2015 Mar 6;347(6226):1138–42.

878   30.   Fuzik J, Zeisel A, Máté Z, Calvigioni D, Yanagawa Y, Szabó G, et al. Integration of
879   electrophysiological recordings with single-cell RNA-seq data identifies neuronal subtypes. Nat
880   Biotechnol. 2016 Feb;34(2):175–83.

881   31.   Cadwell CR, Palasantza A, Jiang X, Berens P, Deng Q, Yilmaz M, et al. Electrophysiological,
882   transcriptomic and morphologic profiling of single neurons using Patch-seq. Nat Biotechnol. 2016
883   Feb;34(2):199–203.

884   32.   Dymecki SM, Ray RS, Kim JC. Mapping Cell Fate and Function Using Recombinase-Based
885   Intersectional Strategies. Guide to Techniques in Mouse Development, Part B: Mouse Molecular
886   Genetics, 2nd Edition. Elsevier; 2010. p. 183–213.

887   33.   Robertson SD, Plummer NW, de Marchena J, Jensen P. Developmental origins of central
888   norepinephrine neuron diversity. Nat Neurosci. 2013 Aug;16(8):1016–23.

889   34.   Brust RD, Corcoran AE, Richerson GB, Nattie E, Dymecki SM. Functional and developmental
890   identification of a molecular subtype of brain serotonergic neuron specialized to regulate breathing
891   dynamics. Cell Rep. 2014 Dec 24;9(6):2152–65.

892   35.   Niederkofler V, Asher TE, Okaty BW, Rood BD, Narayan A, Hwa LS, et al. Identification of
893   Serotonergic Neuronal Modules that Affect Aggressive Behavior. Cell Rep. 2016 Nov
894   15;17(8):1934–49.

895   36.   Ruffault P-L, D'Autréaux F, Hayes JA, Nomaksteinsky M, Autran S, Fujiyama T, et al. The
896   retrotrapezoid nucleus neurons expressing Atoh1 and Phox2b are essential for the respiratory
897   response to CO_2. Elife. 2015 Apr 13;4.

898   37.   He M, Tucciarone J, Lee S, Nigro MJ, Kim Y, Levine JM, et al. Strategies and tools for
899   combinatorial targeting of gabaergic neurons in mouse cerebral cortex. Neuron. 2016 Sep
900   21;91(6):1228–43.

901   38.   Madisen L, Garner AR, Shimaoka D, Chuong AS, Klapoetke NC, Li L, et al. Transgenic mice for
902   intersectional targeting of neural sensors and effectors with high specificity and performance.

Neuron. 2015 Mar 4;85(5):942–58.

39. Chen Y-W, Das M, Oyarzabal EA, Cheng Q, Plummer NW, Smith KG, et al. Genetic identification of a population of noradrenergic neurons implicated in attenuation of stress-related responses. Mol Psychiatry. 2019;24(5):710–25.

40. Tervo DGR, Hwang B-Y, Viswanathan S, Gaj T, Lavzin M, Ritola KD, et al. A designer AAV variant permits efficient retrograde access to projection neurons. Neuron. 2016 Oct 19;92(2):372–82.

41. Zingg B, Chou X-L, Zhang Z-G, Mesik L, Liang F, Tao HW, et al. AAV-Mediated Anterograde Transsynaptic Tagging: Mapping Corticocollicular Input-Defined Neural Pathways for Defense Behaviors. Neuron. 2017 Jan 4;93(1):33–47.

42. Kawano F, Okazaki R, Yazawa M, Sato M. A photoactivatable Cre-loxP recombination system for optogenetic genome engineering. Nat Chem Biol. 2016 Dec;12(12):1059–64.

43. Tang JCY, Rudolph S, Dhande OS, Abraira VE, Choi S, Lapan SW, et al. Cell type-specific manipulation with GFP-dependent Cre recombinase. Nat Neurosci. 2015 Sep;18(9):1334–41.

44. DeNardo LA, Liu CD, Allen WE, Adams EL, Friedmann D, Fu L, et al. Temporal evolution of cortical ensembles promoting remote memory retrieval. Nat Neurosci. 2019 Jan 28;22(3):460–9.

45. Quadros RM, Miura H, Harms DW, Akatsuka H, Sato T, Aida T, et al. Easi-CRISPR: a robust method for one-step generation of mice carrying conditional and insertion alleles using long ssDNA donors and CRISPR ribonucleoproteins. Genome Biol. 2017 May 17;18(1):92.

46. Allen WE, Luo L. Intersectional illumination of neural circuit function. Neuron. 2015 Mar 4;85(5):889–92.

47. Zambrowicz BP, Imamoto A, Fiering S, Herzenberg LA, Kerr WG, Soriano P. Disruption of overlapping transcripts in the ROSA beta geo 26 gene trap strain leads to widespread expression of beta-galactosidase in mouse embryos and hematopoietic cells. Proc Natl Acad Sci USA. 1997 Apr 15;94(8):3789–94.

48. Soriano P. Generalized lacZ expression with the ROSA26 Cre reporter strain. Nat Genet. 1999 Jan;21(1):70–1.

49. Ray RS, Corcoran AE, Brust RD, Kim JC, Richerson GB, Nattie E, et al. Impaired respiratory and body temperature control upon acute serotonergic neuron inhibition. Science. 2011 Jul 29;333(6042):637–42.

50. Cong L, Ran FA, Cox D, Lin S, Barretto R, Habib N, et al. Multiplex genome engineering using CRISPR/Cas systems. Science. 2013 Feb 15;339(6121):819–23.

51. Sun JJ, Ray R. Generation of Two Noradrenergic-Specific Dopamine-Beta-Hydroxylase-FLPo Knock-In Mice Using CRISPR/Cas9-Mediated Targeting in Embryonic Stem Cells. PLoS One. 2016 Jul 21;11(7):e0159474.

52. Thestrup T, Litzlbauer J, Bartholomäus I, Mues M, Russo L, Dana H, et al. Optimized ratiometric calcium sensors for functional in vivo imaging of neurons and T lymphocytes. Nat Methods. 2014 Feb;11(2):175–82.

53. Alexander GM, Rogan SC, Abbas AI, Armbruster BN, Pei Y, Allen JA, et al. Remote control of neuronal activity in transgenic mice expressing evolved G protein-coupled receptors. Neuron. 2009 Jul 16;63(1):27–39.

54. Guettier J-M, Gautam D, Scarselli M, Ruiz de Azua I, Li JH, Rosemond E, et al. A chemical-genetic approach to study G protein regulation of beta cell function in vivo. Proc Natl Acad Sci USA. 2009 Nov 10;106(45):19197–202.

55. Heiman M, Kulicke R, Fenster RJ, Greengard P, Heintz N. Cell type-specific mRNA purification by translating ribosome affinity purification (TRAP). Nat Protoc. 2014 May 8;9(6):1282–91.

56. Li L, Tasic B, Micheva KD, Ivanov VM, Spletter ML, Smith SJ, et al. Visualizing the distribution of synapses from individual neurons in the mouse brain. PLoS One. 2010 Jul 9;5(7):e11503.

57. Hnasko TS, Perez FA, Scouras AD, Stoll EA, Gale SD, Luquet S, et al. Cre recombinase-mediated restoration of nigrostriatal dopamine in dopamine-deficient mice reverses hypophagia and bradykinesia. Proc Natl Acad Sci USA. 2006 Jun 6;103(23):8858–63.

58.   Soudais C, Laplace-Builhe C, Kissa K, Kremer EJ. Preferential transduction of neurons by canine
adenovirus vectors and their efficient retrograde transport in vivo. FASEB J. 2001
Oct;15(12):2283–5.

59.   Plummer NW, Chandler DJ, Powell JM, Scappini EL, Waterhouse BD, Jensen P. An Intersectional
Viral-Genetic Method for Fluorescent Tracing of Axon Collaterals Reveals Details of
Noradrenergic Locus Coeruleus Structure. Eneuro. 2020 May 15;7(3).

60.   Rodríguez CI, Buchholz F, Galloway J, Sequerra R, Kasper J, Ayala R, et al. High-efficiency
deleter mice show that FLPe is an alternative to Cre-loxP. Nat Genet. 2000 Jun;25(2):139–40.

61.   Jensen P, Farago AF, Awatramani RB, Scott MM, Deneris ES, Dymecki SM. Redefining the
serotonergic system by genetic lineage. Nat Neurosci. 2008 Apr;11(4):417–9.

62.   Egbert JR, Fahey PG, Reimer J, Owen CM, Evsikov AV, Nikolaev VO, et al. Follicle-stimulating
hormone and luteinizing hormone increase Ca^{2+} in the granulosa cells of mouse ovarian follicles†.
Biol Reprod. 2019 May 14;

63.   Quadros RM, Miura H, Harms DW, Akatsuka H, Sato T, Aida T, et al. Easi-CRISPR: a robust
method for one-step generation of mice carrying conditional and insertion alleles using long
ssDNA donors and CRISPR ribonucleoproteins. Genome Biol. 2017 May 17;18(1):92.

64.   Wang B, Li K, Wang A, Reiser M, Saunders T, Lockey RF, et al. Highly efficient CRISPR/HDR-
mediated knock-in for mouse embryonic stem cells and zygotes. BioTechniques. 2015 Oct
1;59(4):201–2, 204, 206.

65.   Chu VT, Weber T, Graf R, Sommermann T, Petsch K, Sack U, et al. Efficient generation of Rosa26
knock-in mice using CRISPR/Cas9 in C57BL/6 zygotes. BMC Biotechnol. 2016 Jan 16;16:4.

66.   Yoshimi K, Kunihiro Y, Kaneko T, Nagahora H, Voigt B, Mashimo T. ssODN-mediated knock-in
with CRISPR-Cas for large genomic regions in zygotes. Nat Commun. 2016 Jan 20;7:10431.

67.   Bang SJ, Jensen P, Dymecki SM, Commons KG. Projections and interconnections of genetically
defined serotonin neurons in mice. Eur J Neurosci. 2012 Jan;35(1):85–96.

68.   Bochkov YA, Palmenberg AC. Translational efficiency of EMCV IRES in bicistronic vectors is
dependent upon IRES sequence and gene location. BioTechniques. 2006 Sep;41(3):283–4, 286,
288 passim.

69.   Viemari J-C, Roux J-C, Tryba AK, Saywell V, Burnet H, Peña F, et al. Mecp2 deficiency disrupts
norepinephrine and respiratory systems in mice. J Neurosci. 2005 Dec 14;25(50):11521–30.

70.   Viemari J-C, Ramirez J-M. Norepinephrine differentially modulates different types of respiratory
pacemaker and nonpacemaker neurons. J Neurophysiol. 2006 Apr;95(4):2070–82.

71.   Li A, Emond L, Nattie E. Brainstem catecholaminergic neurons modulate both respiratory and
cardiovascular function. Adv Exp Med Biol. 2008;605:371–6.

72.   Biancardi V, Bícego KC, Almeida MC, Gargaglioni LH. Locus coeruleus noradrenergic neurons
and CO2 drive to breathing. Pflugers Arch. 2008 Mar;455(6):1119–28.

73.   Viemari J-C. Noradrenergic modulation of the respiratory neural network. Respir Physiol
Neurobiol. 2008 Dec 10;164(1–2):123–30.

74.   Gargaglioni LH, Hartzler LK, Putnam RW. The locus coeruleus and central chemosensitivity.
Respir Physiol Neurobiol. 2010 Oct 31;173(3):264–73.

75.   Kuo F-S, Falquetto B, Chen D, Oliveira LM, Takakura AC, Mulkey DK. In vitro characterization
of noradrenergic modulation of chemosensitive neurons in the retrotrapezoid nucleus. J
Neurophysiol. 2016 Sep 1;116(3):1024–35.

76.   Engelbrecht AH, Russell V, Carstens ME, De Villiers AS, Searson A, Jaffer A, et al. Evidence that
noradrenergic neurons in the A1 and A2 nuclei are lesioned by low doses of 6-OHDA injected into
the locus coeruleus. J Neurosci Methods. 1994 Apr;52(1):57–60.

77.   Massey CA, Iceman KE, Johansen SL, Wu Y, Harris MB, Richerson GB. Isoflurane abolishes
spontaneous firing of serotonin neurons and masks their pH/CO_2 chemosensitivity. J Neurophysiol.
2015 Apr 1;113(7):2879–88.

78.   Ting JT, Daigle TL, Chen Q, Feng G. Acute brain slice methods for adult and aging animals:
application of targeted patch clamp analysis and optogenetics. Patch.

1005 79. Jiang X, Shen S, Cadwell CR, Berens P, Sinz F, Ecker AS, et al. Principles of connectivity among
1006   morphologically defined cell types in adult neocortex. Science. 2015 Nov 27;350(6264):aac9462.
1007 80. Scala F, Kobak D, Shan S, Bernaerts Y, Laturnus S, Cadwell CR, et al. Layer 4 of mouse neocortex
1008   differs in cell types and circuit organization between sensory areas. Nat Commun. 2019 Sep
1009   13;10(1):4174.
1010 81. Scala F, Kobak D, Bernabucci M, Bernaerts Y, Cadwell CR, Castro JR, et al. Phenotypic variation
1011   of transcriptomic cell types in mouse motor cortex. Nature. 2020 Nov 12;
1012 82. Martinez VK, Saldana-Morales F, Sun JJ, Zhu PJ, Costa-Mattioli M, Ray RS. Off-Target Effects of
1013   Clozapine-N-Oxide on the Chemosensory Reflex Are Masked by High Stress Levels. Front
1014   Physiol. 2019 May 22;10:521.
1015 83. Shuhaibar LC, Egbert JR, Norris RP, Lampe PD, Nikolaev VO, Thunemann M, et al. Intercellular
1016   signaling via cyclic GMP diffusion through gap junctions restarts meiosis in mouse ovarian
1017   follicles. Proc Natl Acad Sci USA. 2015 Apr 28;112(17):5527–32.
1018

Figure 1

Figure 2

A

	Sequence	Score	Mut	Locus
	ACTGGAGTTGCAGATCACGA GGG			
Off1:	.T.T.............A. CA.	1.1	3MM	chr15:-9941110
Off2:	C...TG...T......... T..	0.8	4MM	chr14:-25087434
Off3:	C......C.TG......... TA.	0.7	4MM	chr9:+122854629
Off4:	.T..A....TT......... .A.	0.7	4MM	chr6:-97126315
Off5:	TA..C.............T AA.	0.7	4MM	chr16:-4397663
Off6:	.T......G.........A. AA.	0.7	3MM	chr10:+35384261
Off7:	.A..A..A........T. A..	0.5	4MM	chr11:-82755370
Off8:	.........AT.C....... AA.	0.5	3MM	chr6:-124885038
Off9:	..G.T....T.......T. T..	0.5	4MM	chr6:-41845311
Off10:	......C..C....A...... CA.	0.5	3MM	chr12:+103941450
Off11:	C.....AC.........C... ...	0.4	4MM	chr13:+43290788
Off12:	.G.....AA........C... T..	0.4	4MM	chr8:+95318409
Off13:	....CT..........A.. .A.	0.4	3MM	chr1:+11497683
Off14:	.A..T...G.........A. TA.	0.3	4MM	chr12:+13635268
Off15:	.A.A...G.........G.. ...	0.3	4MM	chr6:+41114906

Figure 3

Scale bar 50 μm

Figure 4

Scale bar 50 μm

Figure 5

Scale bar 50 μm

Figure 6

Scale bar 100 μm

Figure 7

A Representative Traces

B

Maximum Depolarization
RR2-hM3D

Maximum Hyperpolarization
RR2-hM3D

RC::FPDi-hM3D

C Mean ΔVm after CNO

Figure 8

Figure 9

Figure 10

A

Transmitted light   CFP fluorescence   YFP fluorescence

theca

outer granulosa

oocyte

B

$10\ mM\ Ca^{2+} + 10\ \mu M$ Ionomycin

YFP / CFP

Control

Ionomycin-induced maximum

Time (min)

Table 1

Vector	Function	Addgene ID Number
	Cas9 (px330) vector with sgRNA	97007
	FLP and Cre responsive cloning vector	97012
	FLP and Cre responsive cloning vector (no neo)	99142
	Cre responsive cloning vector	97009
	FLP responsive cloning vector	97010
	Dre responsive cloning vector	97011
	Empty *Rosa26* targeting vector	97008

*CS: Cloning site

Table 2

Short Name	MMRRC ID	Subtractive gene	Intersectional gene	Function	Line status
RR1*	43516	mCherry	hM4D	Metabotropic neuron perturbation	Established and functional
Gt(ROSA)26Sor^{tm#(CAG-mCherry,-hM4D)Rray}					
RR2*	43515	mCherry	hM3D	Metabotropic neuron stimulation	Established and functional
Gt(ROSA)26Sor^{tm#(CAG-mCherry,-hM3D)Rray}					
RR3*	43518	mCherry	EGFP-L10A	Translating mRNA pulldown	Established
Gt(ROSA)26Sor^{tm#(CAG-mCherry,-EGFP-L10A)Rray}					
RR4*	43519	mCherry	PSAM	Ionotropic neuron perturbation	Established
Gt(ROSA)26Sor^{tm#(CAG-mCherry,-PSAM)Rray}					
RR5	43513	None	H2B-TagBFP; sfGFP; synaptophysin-tdTomato	Intersectional fluorescent marking	Established and functional
Gt(ROSA)26Sor^{tm#(CAG-H2B-TagBFP-sfGFP-synaptophysin-tdTomato)Rray}					
RR6	43517	mCherry	Met-RS	Synthesized protein pulldown	Established
Gt(ROSA)26Sor^{tm#(CAG-mCherry,-MetRS)Rray}					
RR7	43514	mCherry	Rs-EGFP	Metabotropic Gs excitation	Established
Gt(ROSA)26Sor^{tm#(CAG-mCherry,-Rs-eGFP)Rray}					
RR8	TBD	None	Twitch2B	Ratiometric calcium imaging	Established and functional
Gt(ROSA)26Sor^{tm#(CAG-Twitch2B)Rray}					
RR9		tdTomato	H2B-TagBFP; sfGFP	Subtractive and intersectional fluorescent marking	Targeted ES cells
Gt(ROSA)26Sor^{tm#(CAG-tdTomato,-H2B-TagBFP-sfGFP)Rray}					
RR10		None	H2B-TagBFP; sfGFP	Dre-responsive fluorescent marking	Targeted ES cells
Gt(ROSA)26Sor^{tm#(CAG-H2B-TagBFP-sfGFP)Rray}					
RR11		None	H2B-TagBFP; sfGFP; synaptophysin-tdTomato	Dre-responsive fluorescent marking	Targeted ES cells
Gt(ROSA)26Sor^{tm#(CAG-H2B-TagBFP-sfGFP-synaptophysin-tdTomato)Rray}					

Grey rows highlight Mouse Genome Informatics names; (tm#) are currently being determined
*Made using an earlier Rosa26 targeting vector without CRISPR/Cas9

Table 3

Knock-in Cassette	Short name	Length	Efficiency
CAG F S F P mCherry S P Met-RS	RR6	11037 bps	6/48 (13%)
CAG F S F P mCherry S P Rs-EGFP	RR7	9534 bps	1/48 (2%)
CAG F S F P tdTomato S P H2B-TagBFP sfGFP	RR9	9597 bps	6/48 (13%)
CAG R S R H2B-TagBFP sfGFP synaptophysin-tdTomato	RR11	9202 bps	5/48 (10%)
CAG R S R H2B-TagBFP sfGFP	RR10	6732 bps	9/48 (19%)
		Total	25/48 (52%)