Growth allometry and dental topography in Upper Triassic conodonts

support trophic differentiation and molar-like element function

Valentin Kelz[1], Pauline Guenser*[1], Manuel Rigo[2], Emilia Jarochowska[1, 3]

[1] Fachgruppe Paläoumwelt, GeoZentrum Nordbayern, Friedrich-Alexander-Universität

Erlangen-Nürnberg, Erlangen, Germany. valentin.kelz@fau.de; pauline.guenser@gmail.com

[2] Department of Geosciences, University of Padova, Padova, Veneto, Italy.

manuel.rigo@unipd.it

[3] Department of Earth Sciences, Utrecht University, Utrecht, Netherlands.

e.b.jarochowska@uu.nl

*Corresponding Author

mailto: pauline.guenser@gmail.com

Abstract. — Conodont elements have high rates of morphological evolution, but the drivers of

this disparity are debated. Positive allometric relationships between dimensions of food-

processing surfaces and entire P_1 elements have been used in to argue that these elements

performed mechanical digestion. If involved in food processing, the surface of the element

should grow at a rate proportional to the increase in energy requirements of the animal. This

inference of function relies on the assumption that the energy requirements of the animal grew

faster ($\cong$ mass$^{0.75}$) than the tooth area ($\cong$ mass$^{0.67}$). We re-evaluate this assumption based on

metabolic rates across animals, and calculate the allometry in platform-bearing P_1 elements of

Late Triassic co-occurring taxa, *Metapolygnathus communisti* and *Epigondolella rigoi,* using 3D

models of ontogenetic series. Positive allometry is found in platform and element dimensions in

both species, supporting a grasping-tooth hypothesis, based on the assumption that metabolic rate

in conodonts scaled with body mass similarly to that in fish and ectoterms. We also calculate the

curvature of P_1 platform surface using the Dirichlet Normal Energy (DNE) as a proxy for diet.

DNE values increase with body mass, supporting the assumption that conodont metabolic rates

increased faster than mass$^{0.67}$. We finally find that adults in both taxa differ in their food bases,

which supports trophic diversification as an important driver of the remarkable disparity of

conodont elements.

Competing interests

36   The authors declare no competing interests.

Introduction

Conodonts are extinct, eel-shaped marine animals that lived from the mid-Cambrian to Early

Jurassic (Du et al. 2020). They are early vertebrates, either stem-gnathostomes (Donoghue et al.

1998) or stem-cyclostomes (Miyashita et al. 2019), and they are distinguished by an extensive

fossil record (Foote and Sepkoski 1999; Donoghue 2001*a*). The majority of the conodont fossil

record consists of the phosphatic elements forming the feeding apparatus of the animal (Schmidt

1934; Scott 1934; Purnell et al. 2000). Conodont elements were retained throughout the life of an

individual (Donoghue and Purnell 1999*a*), recording periodic growth through the apposition of

crown tissue lamellae (Zhang et al. 1997; Dzik 2008). Soft tissues are rarely found and have, so

far, not revealed a great diversity of body forms. Conodont taxonomy and functional morphology

are thus based on their feeding elements (Mazza et al. 2012*b* for the Upper Triassic and

references therein). Patterns interpreted to be produced by elements shearing on the surface and

repaired during the animal's growth suggest macrophagy (Purnell 1995; Donoghue and Purnell

1999*a*; Shirley et al. 2018). An active, predatory lifestyle is supported by the discovery of a

conodont specimen with preserved extrinsic eye musculature (Gabbott et al. 1995), which was

interpreted as indicative of conodonts having pattern vision (Purnell 1994). Calcium isotope

analyses indicated that Late Devonian conodonts were first-level zooplanktivore and piscivore

consumers (Balter et al. 2019), suggesting that in this period, conodonts did not live a purely

predatory lifestyle. However, despite several publications related to this topic, the diet of

conodonts and its evolution are far from resolved. Conodonts changed their apparatus structure

and disparity across their stratigraphic range (Dzik 1991, 2015), possibly reflecting their

evolving niches as marine ecosystems increased in complexity (Klug et al. 2010; Ginot and

Goudemand 2019). Under the assumption that conodont element morphology is an adaptation to

their diet (Jones et al. 2012*a*; Ginot and Goudemand 2019; Guenser et al. 2019; Petryshen et al.

2020), the disparity suggests changing trophic position throughout the existence of the lineage.

Calcium isotope analysis indicated that trophic niches overlapped, suggesting competition

between some taxa (Balter et al. 2019). On the other hand, Sr/Ca ratios in Silurian conodont

assemblages indicate differences between species (Terrill et al. 2022), possibly reflecting trophic

niche differentiation through disruptive selection. Since there is no direct evidence of conodont

food base, trophic diversity of conodonts must be inferred from proxies, e.g. by evaluating

morphological and functional diversity of food-processing elements.

Here we revisit the test between two hypotheses on the function of conodont apparatus: the

suspension-feeding model (Nicoll 1985, 1987) *versus* the grasping-tooth model (Aldridge et al.

1987; Purnell and von Bitter 1992). In the grasping-tooth model and its subsequent

modifications, S and M elements, positioned in the anterior part of the mouth, are interpreted to

perform a grasping function, whereas P elements, placed in the posterior part of the apparatus, in

the pharynx of the animal, have a function similar to molars of mammals (Purnell and von Bitter

1992; Purnell 1994; Donoghue and Purnell 1999*b*; Goudemand et al. 2011). In the suspension

feeder model by Nicoll (1987), elements would be covered in tissue and S and M elements

would filter particles and create current, whereas P elements would only lightly mash food.

Several analytical methods have been proposed in favor of the grasping-tooth model: (1)

observation of microwear patterns and damage of conodont elements, produced *in vivo* (Purnell

1995; Martínez-Pérez et al. 2014*a*; Shirley et al. 2018); (2) occlusion models (Donoghue and

Purnell 1999*b*; Jones et al. 2012*a*; Martínez-Pérez et al. 2014*b*, *a*); (3) Finite Element Analysis

(FEA) (Jones et al. 2012*b*; Martínez-Pérez et al. 2014*a*, 2016); (4) histological adaptation

(Donoghue 2001*b*; Jones et al. 2012*a*); and (5) growth allometry (Purnell 1993, 1994), which is

central in this study.

Allometry describes proportional relationships of body parts, usually of the size of an organ

relative to the total size of the organism. Proportional growth, whereby the growth of an organ

and the size of the animal increase at the same rate, is called isometry. Positive allometry then

describes the organ growing at a faster rate than the rest of the animal. Negative allometry,

conversely, describes the organ growing at a slower rate than the rest of the body (e.g., Gould

1966; Alberch et al. 1979; Klingenberg 1996). Methods of investigation listed in the previous

paragraph have focused on elements termed P_1. Those elements are typically the largest and the

most robust in the apparatus and, thus, inferred to have the most pronounced dental function.

They also have the highest morphological evolutionary rate and, thus, present the most diverse

shapes within the conodont apparatus. Among the possible P_1 morphologies, some of them bear a

platform, which is inferred to be a food-processing surface (Purnell 1995). Positive allometry of

the platform area relative to the total length of the P_1 element in Carboniferous taxa

Idiognathodus sp. and *Gnathodus bilineatus* has been used in seminal studies by Purnell (1993,

1994) to test between the two hypotheses on feeding in conodonts. The test used by Purnell

(1993, 1994) was devised based on the understanding of growth allometry at the time, which led

to the following assumptions: (1) a tooth's ability to process food is proportional to its area; (2)

any tooth linear dimension increases as any other tooth linear dimension (isometric slope = 1);

(3) tooth area increases as the square of its linear dimensions (isometric slope = 2) or as its

volume to the power of 0.67; (4) the animal's metabolic rate and thus energy requirements

increase as its volume to the power of 0.75 (Gould 1966, 1975). Based on these assumptions, the

dental function of P_1 conodont elements, analogously to molar teeth in mammals, should

manifest itself in positive allometry of the food-processing area *versus* the element's linear

dimensions to meet the food requirements (Figure 1).

In the present study we first apply the same allometric evaluation of P_1 elements as used by

Purnell (1993, 1994) to two Late Triassic platform-bearing conodont species *Metapolygnathus*

communisti Hayashi, 1968 (Hayashi 1968) and *Epigondolella rigoi* Kozur, 2007 (Noyan and

Kozur 2007). However, we also revisit assumption 3 (previous paragraph) based on the research

progress since the original studies by Purnell (1993, 1994) and examine the implications when

they are violated (see Discussion). Assumption 3 is that tooth area can be approximated by a

two-dimensional surface, such as when analyzing photographs of teeth in a plane view. An

alternative way of increasing the surface is introducing topography – a solution found in

mammals (Santana et al. 2011; Pérez-Ramos et al. 2020). Conodonts have developed intricate

topographies on the platform surface of their P_1 elements, which would not be identified as

increased areas when approximating the total area with an ellipse described by the element's

linear dimensions (Purnell 1993, 1994). Therefore, here we use 3D meshes to calculate the three-

dimensional platform area and thus account for the topography. To evaluate the importance of

dental topography in conodonts, we calculate Dirichlet Normal Energy (DNE), which measures

the curvature and morphological irregularity of a surface. Essentially, DNE measures how much

a given surface differs from a plane (Bunn et al. 2011). Surface topography is an important

feature of teeth because it helps break down food to satisfy the demand for energy (Bunn et al.

2011). Then, a particular DNE value of the occlusal surface of a tooth (or tooth-like buccal elements in conodonts) should reflect a particular diet. However, DNE has, until now, been only used to analyze skeletal parts of mammals, especially primate teeth (Bunn et al. 2011; Godfrey et al. 2012; Winchester et al. 2014; Prufrock et al. 2016; Berthaume and Schroer 2017; López-Torres et al. 2018; Pampush et al. 2019; Fulwood 2020; Li et al. 2020; Cuesta-Torralvo et al. 2021), but also those of marsupials (Lang et al. 2022), carnivorans (hyenas, bears) (Pérez-Ramos et al. 2020; de Vries et al. 2021; Lang et al. 2022), scandentians (tree shrews) (Selig et al. 2019), rodents (Prufrock et al. 2016; Renaud and Ledevin 2017; Vermeer 2019; de Vries et al. 2021), chiropterans (Pellegrom 2019; López-Aguirre et al. 2022; Villalobos-Chaves and Santana 2022), multituberculates (Robson 2018), artiodactyls (suids) (Rannikko et al. 2020), eulipotyphles (hedgehogs) (Vitek et al. 2021) and teeth of one mammal stem group (Harper et al. 2019). DNE is defined in such a way that it is independent of scale. Thus, any change with the linear dimensions of the animal might be interpreted as a functional change and not a scaling effect (Pampush et al., 2022). DNE values of platform surface of P_1 elements have the potential to decipher the diet of a conodont species. Preliminary dental topographic analyses, different than DNE, have been applied to conodont elements suggested that conodonts might had different diets through their evolution (Purnell and Evans 2009; Stockey et al. 2021, 2022). These recent findings support the potential of dental topographic methods for resolving conodont diets and DNE is a novel method to test functional hypotheses.

The aims of this study are to:

1. Test the null hypothesis of isometric growth against the alternative of positive allometric growth of P_1 elements in a new set of taxa, to evaluate whether the relationship previously observed in Carboniferous taxa by Purnell (1993, 1994) holds more widely in the class Conodonta;

2. Test the null hypothesis of an absence of relationship between DNE values of P_1 platform surface and P_1 element dimensions (i.e. element length, platform length and platform area). If such a relationship exists, by fitting the data with a power law, it is discussed in the light of the energy requirement of the animal;

3. Revise the assumption that the metabolic rate in conodonts increased faster with body mass than tooth area and evaluate how violation of this assumption would affect the outcome of the allometric analysis by Purnell (1993, 1994) and carried out here;

4. Test the hypothesis that differential morphologies in P_1 elements of the co-occurring taxa *M. communisti* and *E. rigoi* reflect different food bases within the ecosystem by comparing DNE values between P_1 platformsurfaces in adult specimens.

FIGURE 1

Materials & Methods

Material

We studied two growth series of the ozarkodinid conodont species *Metapolygnathus communisti* and *Epigondolella rigoi* from the Pizzo Mondello section in western Sicily, Italy. They were collected from a section of 430 m thick marine limestone dated to the upper Carnian to upper Norian (Mazza et al. 2012*a*). Twenty-seven P_1 elements of *M. communisti* and 23 P_1 elements of *E. rigoi* were used, separated into six growth stages (GS) based on the maturity of morphological characters of the platform (Mazza and Martínez-Pérez 2015). The six growth stages are GS1 – early juvenile, GS2 – juvenile, GS3 – late juvenile, GS4 – early adult, GS5 – adult and GS6 – late adult (Mazza and Martínez-Pérez 2015) (Table 1). At Pizzo Mondello, *M. communisti* occurs from the upper Carnian to the lower Norian (from c.a. -227.5 Ma to c.a. -226.5 Ma) (Mazza et al. 2012*a*, 2018; Ogg et al. 2020). The specimens range from late juvenile to late adult (i.e., from GS3 to GS6), though mature elements are more abundant (Table 1). The stratigraphic range of *E. rigoi* at Pizzo Mondello is from the lower Norian to the middle Norian (from c.a. -227 Ma to c.a. -216 Ma) (Mazza et al. 2010, 2012*a*; Ogg et al. 2020), a longer interval than *M. communisti*. Elements range from GS2 to GS5: earlier ontogenetic stages are sparse (Table 1). These specimens have an average colour alteration index (CAI) of 1, suggesting minimal post-depositional heating (Epstein et al. 1977; Nicora et al. 2007; Mazza et al. 2012*a*). The studied elements are housed in the collection of the Dipartimento di Scienze della Terra "A. Desio" of the Università degli Studi di Milano. The whole conodont collection from the Pizzo Mondello section is housed in Milan and Padova (Department of Geosciences, University of Padova).

TABLE 1

Methods

Scanning

The specimens were scanned with a resolution of 1 µm with a microtomograph nanotomS (General Electric) of the AniRA-ImmOs platform, SFR Biosciences (UMS 3444), Ecole Normale Supérieure de Lyon, France. Amira© software was used for the 3D reconstruction (Guenser et al. 2019; Figure 2). The meshes are available on MorphoBank:
http://morphobank.org/permalink/?P4048

FIGURE 2

Growth allometry

The length of the element was used as a proxy for the size of the conodont animal, as was done in previous studies (Purnell 1993, 1994; Zhang et al. 2018; Ginot and Goudemand 2019). The element length, the platform length and the platform area were measured using the 3D software MeshLab 2020.12 (Cignoni et al. 2008). The length of the element was measured from the anteriormost point of the blade in a straight line to the middle of the posterior edge of the element's platform (Figure 3). As the platform is not equally long on the two sides of the blade, its length was measured in two ways. First, as most elements are curved, the convex side of the platform was measured. In *Metapolygnathus communisti*, the convex size tends to be the longer side of the platform, though not always. In *Epigondolella rigoi* the convex side was almost exclusively the longer side. Alternatively, the longest side of the platform was measured, regardless of curvature (Figure 3, see also Figure S1 in Supplementary Information). In *M. communisti*, in both instances, the platform was measured from the most anterior part of the platform to its posterior end in a line parallel to the imagined symmetrical axis of the platform (Figure 3). In *E. rigoi*, the platform was measured from the geniculation point to the platform's posterior end. This measure was chosen because the anterior trough margin in this species, though reduced, reaches quite far up the blade, especially in more mature growth stages (see Mazza et al. 2012*a* for details about the taxonomic characters).

FIGURE 3

The measured area of the platform includes the platform, the cusp, and any postcarinal nodes

(Figure 3). In *M. communisti*, specimens of earlier growth stages tend to exhibit only one

postcarinal node, already present in GS1 (Mazza and Martínez-Pérez 2015). From GS3 on, a

second posterior node may appear (Mazza and Martínez-Pérez 2015). Adult specimens show

three or four posterior carinal nodes (Mazza et al. 2012*a*). However, our measurements

consistently included only two or three postcarinal nodes in adult specimens. In *E. rigoi*, the cusp

is always followed by a single larger postcarinal node (Mazza et al. 2012*a*). These parts of P_1

elements were added to the measurements of the platform area, even though they are

taxonomically not part of the platform because they likely played a similar part in the processing

of food as the platform itself. In *E. rigoi*, the anterior trough margin was not included in the

measurements of the platform area (Figure 3). The anterior trough margin is not present in *M.*

communisti (Mazza et al. 2012*a*).

Reduced major axis regression (RMA) was calculated using the R package "smatr" 3.4-8

(Warton et al. 2012; R Core Team 2021) to examine the relationship between the length of the

platform and the length of the element, as well as the platform area and the length of the element.

All measurements were log-transformed to assess allometric relationships. RMA was chosen as a

method because both variables are mutually dependent. Slopes obtained with *E. rigoi* and *M.*

communisti data were compared with the "slope.com" function of the "smatr" R package. The

same function was used to compare slopes related to the convex side and the longer side of the

platform within a species. Slopes between both species and isometry were compared with

"slope.test" function of "smart" R package. Isometry was modeled with a slope coefficient of 1

when testing the platform length *vs.* element length; a slope coefficient of 2 when testing the

platform area *vs.* element length. In the case slopes do not differ significantly, we compare the

intercepts and their confidence intervals showed with the "sma" function. A differentiation in

trophic niche could be indeed reflected in different intercepts values when slopes are similar

(Lumer et al. 1942; Gould 1979).

Purnell (1994) used a Z-test (Hayami and Matsukuma 1970) to test whether slope coefficients

of *Idiognathodus* sp. and *Gnathodus bilineatus* differed significantly from isometric growth. A

Z-index higher than 1.96 means that the relationship differs from isometry significantly. We

consider this index comparable to the p-values we obtained for *M. communisti* and *E. rigoi* when

comparing their slope coefficients with slopes expected under isometry. To compare slope

coefficients from this study with those provided by Purnell (1994), 95% confidence intervals

(95% CI) of slope coefficients of *Idiognathodus* sp. and *G. bilineatus* were calculated according

to the following formula:

$$\beta_1 = b_1 \pm t_{0.025, n-2} \times SE(b_1),$$

Where b_1 is the slope coefficient, *n* is the number of measurements, *t* is the t distribution and

SE – standard error. Measurements for the platform length of *G. bilineatus* were unavailable

(Purnell 1994).

Quantitative topographic analysis (DNE)

Dirichlet Normal Energy (DNE) measures a surface's curvature (Bunn et al. 2011). The DNE

of an object is independent of its size and orientation. Its equation is commonly written as

follows:

$$DNE = \sum e(p) \times area(p)$$

Here, *e(p)* is the Dirichlet Energy Density at a point *p*. The sum of the areas of all points *p*

(however small) on a surface is equal to the total area of the surface. Flat planes have DNE

values of 0. Therefore, a higher DNE value expresses the elements' complexity and the average

"sharpness" of a surface (Bunn et al. 2011).

We applied the DNE on P_1 element meshes with the R package "molaR" 5.0 (Pampush et al.

2016). To ensure consistency of DNE calculation, all meshes were simplified to 10 000 faces

(Spradley et al. 2017) using Quadric Edge Collapse Decimation in MeshLab. They were then

rescaled so that each platform area equaled 0.1 mm^2. Meshes were then smoothed in Avizo using

the Avizo smoothing function with lambda = 0.6 and 25 iterations. Twenty to thirty smoothing

iterations, a conservative amount when compared to other approaches (Bunn et al. 2011;

Winchester et al. 2014; Spradley et al. 2017), are recommended because they eliminate scanning

noise while capturing fine-scale features and avoiding the creation of artificial dimples that can

be caused by over smoothing (Spradley et al. 2017). Different numbers of iterations (5, 10, 15,

20, 25, 30, 40, 50, 60, 70, 80, 90, and 100) were tested on a single specimen of *M. communisti* to

determine the impact of the number of smoothing iterations (Figure S2). The DNE appears stable

from ~20 to 25 iterations.

The meshes were then manually cut to keep the occluding surface of the platform, the cusp

and the postcarinal nodes (Figure 4). Additionally, we cut out the aboral part of the platform

because it does not take an active part in food processing. The meshes were finally imported into

R Software as binary *ply* files. Individual pieces of meshes created by the smoothing operation

were removed to prevent them from affecting the DNE calculation. DNE was calculated with an

included boundary exclusion criterion (BoundaryDiscard="vertex"), as advised by Spradley et al.

(2017). The total surface DNE is the mean of all DNE values for individual faces of a surface

(Pampush et al. 2016). Because a DNE value is not a traditional measurement, we did not log-

transformed the data. We rather fitted the relationship between DNE values and element

dimensions (i.e element length, platform length and platform area) with a power law using the

"aomisc" R package (Onofri 2020). In addition, Growth stage 5 (GS5) was chosen to compare

DNE values between species because this stage comprised enough specimens from both species

to allow statistical analysis (Table 1). GS5, representing adults, also allows for interpretations of

the diet. The medians of DNE distributions were compared with a Kruskal-Wallis test in R

(Hollander and Wolfe 1973).

The R code and data used for investigating allometric patterns and DNE are available on OSF

at this address: https://osf.io/283wq/?view_only=6f22274998134eb99cbe43749c6e3e7e (Kelz et

al. 2022).

FIGURE 4

Results

Growth allometry

In all examined relationships, high coefficients of determination ($R^2 \geq 0.89$) indicated that

linear regression captured the relationships between the variables sufficiently (Table 2). In

Metapolygnathus communisti, linear regression slopes of the platform length over the P_1 element

length and the platform area over the P_1 element length showed positive allometry (Figure 5A-

B). In both cases, the slopes differed significantly from values corresponding to isometry, i.e.

one and two, respectively (Table 2). No differences between the two ways the platform length

was measured could be detected (p=0.23; Figure S1A).

In *Epigondolella rigoi*, both regression slopes (platform length over element length and

platform area over element length) indicated positive allometry and were significantly higher

than predicted under the null hypothesis of isometric growth (Table 2; Figure 5 Morphological

characters follow Mazza et al. (2012*a*, *b*).

Figure 4: Log-transformed DNE values across growth stages GS3 to GS6 of *Metapolygnathus*

communisti and across growth stages GS2 to GS5 of *Epigondolella rigoi*. Names of the

specimens are specified in Figure 2.

Figure 5C-D). No significant difference between the two ways the platform length was

measured could be detected (p=0.83; Figure S1B).

TABLE 2

Slope coefficients of the platform length over element length did not differ significantly

between *M. communisti* and *E. rigoi* (p=0.166), but in both cases, they were significantly higher

than that of *Idiognathodus* sp. as their confidence intervals did not overlap (Table 2, Figure 5E,

G). Platform areas in all four species showed positive allometry over element length (slope

coefficients between 2.149 and 2.561) but the four 95% confident intervals overlapped, and no

significant difference was detected between *M. communisti* and *E. rigoi* (p=0.479). No

differences in intercepts could be detected for any of the examined relationships between *E. rigoi*

and *M. communisti* (Table 2).

FIGURE 5

Dental topography (DNE)

In *M. communisti*, DNE values ranged between 99.93 and 279.84; in *E. rigoi*, between 117.59

and 353.71 (Figure 6). Specimens classified as GS5 of *E. rigoi* showed higher DNE values than

GS5 specimens of *M. communisti* (Kruskal-Wallis test, p=0.015; Figure 6A).

The relationship between DNE values and P_1 dimensions show a modest fit to the power law

($0.42 \leq R^2 \leq 0.59$; Table 3, Figure 6B-D).Figure 6 Between *E. rigoi* and *M. communisti*, the

scaling exponents ("b" in Table 3) are respectively 0.66 and 0.72 in DNE *vs.* element length;

0.52 and 0.48 in DNE *vs.* platform length; and 0.29 and 0.30 in DNE *vs.* platform area (Table 3).

No difference in exponents is detected for any of the examined relationships (Figure 6E).

336  FIGURE 6

337  TABLE 3

Discussion

339  The null hypothesis of an isometric growth of P_1 elements in *M. communisti* and *E. rigoi*

340 could be rejected based on positive growth allometry of the platform length and platform area

341 over element length. These results support previous findings of positive growth allometry in

342 these organs in much older, Carboniferous, ozarkodinid taxa *G. bilineatus* and *Idiognathodus* sp.

343 (Purnell 1993, 1994). Under the assumption that positive allometry reflects the organism's

344 energy demand growing at a higher pace than the surface of a (molar) tooth (Gould 1966, 1975),

345 these results are consistent with the interpretation of P_1 elements as organs used for mechanical

346 slicing and grinding of food, as previously proposed based on microwear (Purnell 1995;

347 Martínez-Pérez et al. 2014*b*), enamel-like ultrastructure of conodont lamellar crown tissue

348 (Donoghue 2001*b*), and finite element analysis (Jones et al. 2012*a*). However, the assumptions

349 of the original studies by Purnell (1993, 1994) need to be revisited in the context of the studies

350 which had been made in the field of metabolic ecology over the last 20 years. The premise of the

351 argument based on tooth positive allometry was that an animal's metabolic rate, and thus energy

352 requirements, increase as its volume to the power of 0.75. This scaling exponent has been

353 predicted by several theories (Kleiber 1932; Feldman and McMahon 1983; West et al. 1997;

354 Savage et al. 2004) but an exponent of 0.67 has been equally reported in mammals and derived

355 from theoretical considerations (Heusner 1982; Dodds et al. 2001; White and Seymour 2003), so

356 the same exponent that in assumption 3 developed in the Introduction. This disagreement has

357 important implications for the reasoning in the studies by Purnell (1993, 1994): if the metabolic

358 rate in conodonts scaled as body mass to the power of 0.67 and not 0.75, then an isometric

359 relationship between the food acquisition surface and the animal's dimensions might well be

360 expected. In such case, it would be impossible to distinguish whether positive allometry reflects

361 differences in scaling between energy demand and food-processing tooth surface or whether the

362 two variables are not related. The latter possibility would preclude the use of growth allometry in

363 inferring trophic ecology in fossil organisms. We attempt, therefore, to extrapolate the

364 relationship in conodonts by comparing with related organisms. The average scaling exponent in

fish has been reported as 0.79 (Clarke and Johnston 1999)or even 0.99 (Glazier 2009). In

ectoterms, exponents higher than 0.75 have been reported across all groups compiled by Glazier

(2007). Based on analogy, a scaling exponent of metabolic rate over body mass or volume can be

in conodonts expected to be higher than 0.67. It should be, however, kept in mind that this is a

strong and untestable assumption, as the scaling exponents vary extremely between animals and

even within taxa, depending on their interactions with other organisms (Bokma 2004; Glazier

2009; Sieg et al. 2009). The scaling exponents of DNE over element length lie between 0.66 and

0.72, thus closer to the 0.67 exponent than to 0.75. This might serve as an indication that in

conodonts, energy demand grew in a more mammal-like fashion (closer to $M^{0.67}$) than in a fish or

reptile fashion ($M^{>0.67}$), in line with the idea of conodont P_1 elements functioning analogously to

molars (Donoghue and Purnell 1999*b*). But scaling of DNE with tooth size has only been

recently identified apes (Berthaume and Schroer 2017) and the interpretation of the relationships

between this metric, body size and metabolism are little known, therefore the scaling observed

here requires further comparisons with extant organisms. Here we tentatively uphold the

reasoning and conclusions by Purnell (1993, 1994) and extend them to the new taxa examined

here: if the metabolic rate in conodonts scaled as $M^{>0.67}$, then the positive allometry of conodont

element dimensions supports the grasping tooth model in all four taxa and the test may be used

in other conodont element morphologies.

Functional differences between conodont species

The allometric relationship between platform area and element length did not differ

significantly between the two Triassic species examined here and Carboniferous taxa

Idiognathodus sp. and *G. bilineatus* (Figure 5G2). However, there is a significant difference

between Triassic species and *Idiognathodus* sp. when considering the growth of the platform

length (Figure 5G1). Perhaps this difference in slope coefficient resulted from the

methodological differences in how platform measurements were taken here and in Purnell (1993,

1994). Measurements based on pictures often suffer from distortions from projecting a 3D

structure onto a plane, where differences in levelling the photographed specimens might affect

the results (Mullin and Taylor 2002; Collins and Gazley 2017). Proper 3D measurements of P_1

platform length and area using our methodological protocol should be investigated in

Idiognathodus sp. and *G. bilineatus* to verify any difference in allometric slope coefficients.

However, as platform length showed a stronger positive allometry in Late Triassic taxa than in

Idiognathodus sp., it suggests that the increase of functional surface was primarily achieved by

platform elongation, rather than growth in width.

Our hypothesis that *M. communisti* and *E. rigoi* differed in the growth allometry of their P_1

elements could not be rejected (Table 2). Similar growth allometry might indicate similar energy

demand, as well as similar efficiency in food acquisition and assimilation.

Conodont diet inferred by platform curviness

Previous investigations showed that dental topography methods could be applied to non-

homologous dental tools to track dietary differences between distantly related clades (Stockey et

al. 2021). Conodont elements in this study showed similar curviness to complex mammal teeth.

This similarity does not mean that direct dietary associations can be made between conodonts

and mammals, but dental topography allows comparisons between taxa and ontogenetic stages

and helps in constraining conodont ecology (Purnell and Evans 2009). In *M. communisti* and *E.*

rigoi, DNE values differed significantly at the adult growth stage (i.e., GS5). We tested the

differences in DNE values only at GS5 because ontogenetically younger growth stages were

represented by fewer specimens in both species (Table 1), which made observations of these

stages less conclusive. Differences in GS5 DNE values between *M. communisti* and *E. rigoi*

allow rejecting the hypothesis that adult specimens of both species shared the same diet. More

DNE analyses on conodonts are needed to understand the scope of DNE values in this group and

to confidently suggest that a discrepancy between DNE values of different species reflects

different dietary niches. Indeed, DNE values of GS5 specimens of *M. communisti* are similar to

those reported for folivores or omnivores; insectivores or folivores in the case of *E. rigoi*, but

these diets based on DNE values stem from studies on primates (Bunn et al. 2011; Winchester et

al. 2014). Moreover, conodont elements are more than ten times smaller than primate teeth,

complicating dietary comparison as conodonts could not eat the same objects. Though primate

dietary classifications do not apply to conodonts, they may offer a general reference point for the

methods of breaking down different food types. Insectivores rely on sharp cusps to apply

maximal force to a small surface area to pierce hard insect chitin, and folivores also use steeply

sloped cusps to shear tough cellulose-rich leaves (Lucas 1979; Strait 1997). Therefore, it is

possible that conodont element platforms evolved to break down food types with similar

mechanical properties. For instance, conodonts may have punctured arthropod larvae (Dzik

2021), a diet consistent with the DNE values observed in *M. communisti*.

Current limits on DNE and advices for further investigations

The definition of the platform for DNE measurement is worth discussing because it is

somewhat subjective. In this work, we decided to include the cusp and all postcarinal nodes

because, in many cases, the cusp marked a notable transition between sharper denticles of the

blade and flatter nodes on the platform. In *M. communisti*, the number of postcarinal nodes

increases through growth (Mazza et al. 2012*a*; Mazza and Martínez-Pérez 2015), impacting DNE

values through the ontogeny of this species. In *E. rigoi*, the number of posterior nodes stays the

same (Mazza et al. 2012*a*), so variations of DNE values are more related to the growth of prior

postcarinal nodes or the addition of new nodes on the edge of the platform. This difference in

node location on the platform might affect the way conodonts broke down food but it should be

further investigated by assessing occlusal kinematics of *M. communisti* and *E. rigoi* P_1 elements,

which is currently impossible because no clusters (i.e., a conodont apparatus with elements found

in connection) are currently known for these species.

There are several challenges when applying DNE to conodonts. As DNE is a comparatively

new tool, reference values and understanding of variability (e.g. intraspecific, ontogenetic and

taphonomic) are limited. So far, DNE research has focused on mammals, whose tooth function

largely relies on jaws acting as levers. The different mechanics of feeding in jawed organisms

necessitates that the comparisons drawn here between conodont elements and primate molars

must be viewed as extremely hypothetical. Marine environment, evolutionary distance and the

lack of jaws in conodonts make tooth function likely not completely analogous between the two.

Even more importantly, conodonts are unique among vertebrates in repairing their teeth by

periodic apposition of new growth layers on top of the ones previously used for food processing

(Shirley et al. 2018). This mechanism is both used to repair damage and restore sharpness

(Donoghue, 1998), as well as to generate topographic complexity (Müller and Nogami 1971).

Thus, DNE values in conodont elements will increase as the elements grow and so the animal

gets older. This is not the case in mammals due to their lack of tooth repairing ability: it has been

showed with mice's molars that DNE values decrease with the age of the animal because of teeth

abrasion (Savriama et al. 2022). That is why typical applications of DNE on mammals do not

consider ontogenetic development, authors rather investigate DNE variation between young

adults from different species, or between different teeth within one specimen. For example,

Pérez-Ramos et al. (2020) compared DNE values between premolars and molars in cave bears,

and they suggested that "increasing upper tooth surface areas also increase the values of both

topographic variables, and probably, also their chewing efficiency".

A consistent protocol of mesh preparation before DNE calculation is also needed to get a

better reproducibility of the method. For example, the scan resolution and the number of

smoothing iterations vary in current literature, which can impact DNE values (Spradley et al.

2017; Assemat et al. 2022). This variation in processing is important because comparisons are

most conclusive when drawn between data with similar preparation. Furthermore, the scan

resolution depends on the size of the studied object. We cannot expect a 1 µm resolution (as for

conodont P_1 elements from the Pizzo Mondello (Guenser et al. 2019)) for mammal teeth of

several millimetres in length. However, we can set a standardized scan resolution for conodont

elements of 1 µm (even lower would be better), which would allow investigating ontogenetic

patterns by including P_1 elements of less than 400 µm in length (i.e., juveniles). About the post-

scanning preparation, we second Spradley et al.'s (2017) recommendations: a conservative

number of smoothing iterations (20-30) using non-Laplace-based smoothing operators, such as

that implemented in Avizo, and mesh simplification to a fixed number of faces.

Conclusions

We used 3D meshes to test the null hypothesis that Late Triassic conodont P_1 elements grew

isometrically, thus revisiting Purnell's (1993, 1994) test of conodont element function. We tested

this hypothesis against the alternative that the elements showed positive allometry. Positive

allometry had been proposed to be a test of the grasping-tooth hypothesis (Aldridge et al. 1987;

Purnell and von Bitter 1992). However, the test requires knowledge of the scaling exponent of

the metabolic rate to body mass (volume). We followed Purnell's protocol (1993, 1994),

analyzing the growth allometry of the platform length and area *vs.* total length of the element,

originally performed on 2D projections of conodont elements. Platform length and platform area

showed positive allometry relative to element length, allowing us to reject the null hypothesis. If

metabolic rate in conodonts scaled with body mass similarly to that in fish and ectoterms, the test

holds and our results support the grasping-tooth hypothesis. Slope coefficients nor intercepts did

not differ between *M. communisti* and *E. rigoi*. However, for platform length over element

length, slope coefficients were higher than those reported for Carboniferous conodonts
Idiognathodus sp. (Purnell 1994), which may suggest that the energy demand (and thus
metabolic rate) increased faster in this species as the organism grew. A more precise
measurement of surface area in 3D models compared to 2D projections used in *Idiognathodus*
sp. did not result in a significantly different relationship. In contrast, length- rather than area-
based measurements allowed detecting significant differences between Late Triassic taxa and
Idiognathodus sp.

We added dental topographic analysis of the platforms using DNE to test the hypothesis that
co-occurring species *M. communisti* and *E. rigoi* shared the same food base. *E. rigoi* showed
significantly higher DNE values than *M. communisti* when comparing adult growth stages.
Although DNE is a size-independent measure, its values increased with the size of adult
specimens in both Triassic taxa. This increase is interpreted here as the reflection of energy
demand growing faster than the area of the element and thus potentially reflecting the scaling of
the metabolic rate consistent with the assumptions of the studies by Purnell (1993, 1994). Scaling
exponents of DNE over element length in an allometric model were closer to those reported for
mammals than for fish and other ectoterms. As these exponents did not differ substantially
between *M. communisti* and *E. rigoi*, the differences in DNE are not likely to arise from differing
metabolic rates and can be attributed to different food bases.

Our study revisited one of the main arguments used in interpreting the function of conodont
elements: their positive growth allometry. We argue that a key assumption of this interpretation,
the scaling of metabolic rate as mass (volume) to the power of 0.75, cannot be made for
conodonts automatically, as there is substantial variation between known animals. We also show
that co-occurring taxa differed in their diets, as reported previously in Silurian conodont
communities (Terrill et al. 2022), which supports trophic diversification as an important driver of
the remarkable disparity of their elements.

Acknowledgements

511 **Acknowledgements**
512   We thank Bryan Shirley for support in using Aviso and Wyatt Petryshen for advice on
513 cleaning and saving meshes. We also thank Michele Mazza for their advice and field sampling.
514 We finally thank Nicolas Goudemand for supporting the project (French ANR grant, ACHN
515 project EvoDevOdonto). PG was supported by Visiting Scholarship awarded by Friedrich-

Alexander-Universität Erlangen-Nürnberg. EJ was supported by Deutsche

Forschungsgemeinschaft (project no JA 2718/3-1). MR was supported by DOR2054230/20 by

University of Padova. We are grateful to Viktor Karádi, Nicolas Campione and Paleobiology

Editor James Crampton for constructive comments, which improved the manuscript, and to Craig

White for sharing data on the scaling of metabolic rates in various taxa.

Literature cited

Alberch, P., S. J. Gould, G. F. Oster, and D. B. Wake. 1979: Size and shape in ontogeny and
        phylogeny. Paleobiology 5:296–317.
Aldridge, R. J., M. P. Smith, R. D. Norby, and D. E. G. Briggs. 1987: The architecture and
        function of Carboniferous polygnathacean conodont apparatuses. Palaeobiology of
        conodonts:63–75.
Assemat, A., G. Thiery, T. Lieffroy, and C. Girard. 2022: 3D topography as tool for shape
        discrimination of conodont elements. Fifth International Conodont Symposium (ICOS5).
Balter, V., J. E. Martin, T. Tacail, G. Suan, S. Renaud, and C. Girard. 2019: Calcium stable
        isotopes place Devonian conodonts as first level consumers. Geochemical Perspectives
        Letters 10:36–39.
Berthaume, M. A., and K. Schroer. 2017: Extant ape dental topography and its implications for
        reconstructing the emergence of early Homo. Journal of Human Evolution 112:15–29.
Bokma, F. 2004: Evidence against Universal Metabolic Allometry. Functional Ecology 18:184–
        187.
Bunn, J. M., D. M. Boyer, Y. Lipman, E. M. St. Clair, J. Jernvall, and I. Daubechies. 2011:
        Comparing Dirichlet normal surface energy of tooth crowns, a new technique of molar
        shape quantification for dietary inference, with previous methods in isolation and in
        combination. American Journal of Physical Anthropology 145:247–261.
Cignoni, P., M. Callieri, M. Corsini, M. Dellepiane, F. Ganovelli, and G. Ranzuglia. 2008:
        Meshlab: an open-source mesh processing tool. Eurographics Italian Chapter Conference
        2008:129–136.
Clarke, A., and N. M. Johnston. 1999: Scaling of metabolic rate with body mass and temperature
        in teleost fish. Journal of Animal Ecology 68:893–905.
Collins, K. S., and M. F. Gazley. 2017: Does my posterior look big in this? The effect of
        photographic distortion on morphometric analyses. Paleobiology 43:508–520.
Cuesta-Torralvo, E., D. Pacheco, L. M. Martínez, A. Romero, C. Umbelino, Y. Avià, and A.
        Pérez-Pérez. 2021: Three-dimensional proxies to dental wear characterization in a known
        age-at-death skeletal collection. Journal of Archaeological Method and Theory 28:1261–
        1275.
Dodds, P. S., D. H. Rothman, and J. S. Weitz. 2001: Re-examination of the "3/4-law" of
        Metabolism. Journal of Theoretical Biology 209:9–27.
Donoghue, P. C. J. 2001a: Conodonts meet cladistics: recovering relationships and assessing the
        completeness of the conodont fossil record. Palaeontology 44:65–93.
———. 2001b: Microstructural variation in conodont enamel is a functional adaptation.
        Proceedings of the Royal Society of London. Series B: Biological Sciences 268:1691–
        1698.

Donoghue, P. C. J., and M. A. Purnell. 1999a: Growth, function, and the conodont fossil record.
Geology 27:251–254.
———. 1999b: Mammal-like occlusion in conodonts. Paleobiology 25:58–74.
Donoghue, P. C. J., M. A. Purnell, and R. J. Aldridge. 1998: Conodont anatomy, chordate
phylogeny and vertebrate classification. Lethaia 31:211–219.
Du, Y., M. Chiari, V. Karádi, A. Nicora, T. Onoue, J. Pálfy, G. Roghi, Y. Tomimatsu, and M.
Rigo. 2020: The asynchronous disappearance of conodonts: New constraints from
Triassic-Jurassic boundary sections in the Tethys and Panthalassa. Earth-Science
Reviews 203:103176.
Dzik, J. 1991: Evolution of oral apparatuses in the conodont chordates. Acta Palaeontologica
Polonica 36:265–323.
———. 2008: Evolution of morphogenesis in 360-million-year-old conodont chordates
calibrated in days. Evolution & Development 10:769–777.
———. 2015: Evolutionary roots of the conodonts with increased number of elements in the
apparatus. Earth and Environmental Science Transactions of the Royal Society of
Edinburgh 106:29–53.
———. 2021: Protaspis larva of an aglaspidid-like arthropod from the Ordovician of Siberia and
its habitat. Arthropod Structure & Development 61:101026.
Epstein, A. G., J. B. Epstein, and L. D. Harris. 1977: Conodont color alteration: an index to
organic metamorphism. United States Geological Survey Professional Paper 995:1–27.
Feldman, H. A., and T. A. McMahon. 1983: The 3/4 mass exponent for energy metabolism is not
a statistical artifact. Respiration Physiology 52:149–163.
Foote, M., and J. J. Sepkoski. 1999: Absolute measures of the completeness of the fossil record.
Nature 398:415–417.
Fulwood, E. L. 2020: Ecometric modelling of tooth shape and precipitation gradients among
lemurs on Madagascar. Biological Journal of the Linnean Society 129:26–40.
Gabbott, S. E., R. J. Aldridge, and J. N. Theron. 1995: A giant conodont with preserved muscle
tissue from the Upper Ordovician of South Africa. Nature 374:800–803.
Ginot, S., and N. Goudemand. 2019: Conodont size, trophic level, and the evolution of platform
elements. Paleobiology 45:458–468.
Glazier, D. S. 2007: 22.1. Ecology of metabolic scaling in animals and plants. Seventh
International Congress of Comparative Physiology and Biochemistry, Salvador, Bahia,
Brazil, August 12–16, 2007 148:S99.
Glazier, D. S. 2009: Activity affects intraspecific body-size scaling of metabolic rate in
ectothermic animals. Journal of Comparative Physiology B 179:821–828.
Godfrey, L. R., J. M. Winchester, S. J. King, D. M. Boyer, and J. Jernvall. 2012: Dental
topography indicates ecological contraction of lemur communities. American Journal of
Physical Anthropology 148:215–227.
Goudemand, N., M. J. Orchard, S. Urdy, H. Bucher, and P. Tafforeau. 2011: Synchrotron-aided
reconstruction of the conodont feeding apparatus and implications for the mouth of the
first vertebrates. Proceedings of the National Academy of Sciences 108:8720–8724.
Gould, S. J. 1966: Allometry and size in ontogeny and phylogeny. Biological Reviews 41:587–
638.
———. 1975: On the Scaling of Tooth Size in Mammals. American Zoologist 15:353–362.
———. 1979: An Allometric Interpretation of Species-Area Curves: The Meaning of the
Coefficient. The American Naturalist 114:335–343.

Guenser, P., L. Souquet, S. Dolédec, M. Mazza, M. Rigo, and N. Goudemand. 2019: Deciphering the roles of environment and development in the evolution of a Late Triassic assemblage of conodont elements. Paleobiology 45:440–457.

Harper, T., A. Parras, and G. W. Rougier. 2019: *Reigitherium* (Meridiolestida, Mesungulatoidea) an enigmatic Late Cretaceous mammal from Patagonia, Argentina: morphology, affinities, and dental evolution. Journal of Mammalian Evolution 26:447–478.

Hayami, I., and A. Matsukuma. 1970: Variation of bivariate characters from the standpoint of allometry. Palaeontology 13:588–605.

Hayashi, S. 1968: The Permian conodonts in chert of the Adoyama Formation, Ashio Mountains, central Japan. Earth Science 2:63–77.

Heusner, A. A. 1982: Energy metabolism and body size I. Is the 0.75 mass exponent of Kleiber's equation a statistical artifact? Respiration Physiology 48:1–12.

Hollander, M., and D. A. Wolfe. 1973: Nonparametric statistical methods. Vol. 751. John Wiley & Sons, p.

Jones, D., A. R. Evans, E. J. Rayfield, K. K. W. Siu, and P. C. J. Donoghue. 2012a: Testing microstructural adaptation in the earliest dental tools. Biology Letters 8:952–955.

Jones, D., A. R. Evans, K. K. W. Siu, E. J. Rayfield, and P. C. J. Donoghue. 2012b: The sharpest tools in the box? Quantitative analysis of conodont element functional morphology. Proceedings of the Royal Society B: Biological Sciences 279:2849–2854.

Kelz, V., P. Guenser, M. Rigo, and E. Jarochowska. 2022: Growth allometry and dental topography in Upper Triassic conodonts support trophic differentiation and molar-like element function. Dataset retrieved from osf.io/283wq. .

Kleiber, M. 1932: Body size and metabolism. Hilgardia 6:315–353.

Klingenberg, C. P. 1996: Multivariate Allometry. Pp.23–49 *in* L. F. Marcus, M. Corti, A. Loy, G. J. P. Naylor, and D. E. Slice, eds. Advances in Morphometrics. Springer US, Boston, MA.

Klug, C., B. Kröger, W. Kiessling, G. L. Mullins, T. Servais, J. Fryda, D. Korn, and S. Turner. 2010: The Devonian nekton revolution. Lethaia 43:465–477.

Lang, A. J., T. Engler, and T. Martin. 2022: Dental topographic and three-dimensional geometric morphometric analysis of carnassialization in different clades of carnivorous mammals (Dasyuromorphia, Carnivora, Hyaenodonta). Journal of Morphology 283:91–108.

Li, P., P. E. Morse, and R. F. Kay. 2020: Dental topographic change with macrowear and dietary inference in *Homunculus patagonicus*. Journal of Human Evolution 144:102786.

López-Aguirre, C., S. J. Hand, N. B. Simmons, and M. T. Silcox. 2022: Untangling the ecological signal in the dental morphology in the bat superfamily Noctilionoidea. Journal of Mammalian Evolution.

López-Torres, S., K. R. Selig, K. A. Prufrock, D. Lin, and M. T. Silcox. 2018: Dental topographic analysis of paromomyid (Plesiadapiformes, Primates) cheek teeth: more than 15 million years of changing surfaces and shifting ecologies. Historical Biology 30:76–88.

Lucas, P. W. 1979: The Dental-Dietary Adaptations of Mammals. Neues Jahrbuch für Geologie und Paläontologie Monatshefte 8:486–512.

Lumer, H., B. G. Anderson, and A. H. Hersh. 1942: On the Significance of the Constant b in the Law of Allometry y = bxa. The American Naturalist 76:364–375.

Martínez-Pérez, C., E. J. Rayfield, M. A. Purnell, and P. C. J. Donoghue. 2014a: Finite element,
occlusal, microwear and microstructural analyses indicate that conodont microstructure is
adapted to dental function. Palaeontology 57:1059–1066.
Martínez-Pérez, C., E. J. Rayfield, H. Botella, and P. C. J. Donoghue. 2016: Translating
taxonomy into the evolution of conodont feeding ecology. Geology 44:247–250.
Martínez-Pérez, C., P. Plasencia, D. Jones, T. Kolar-Jurkovšek, J. Sha, H. Botella, and P. C. J.
Donoghue. 2014b: There is no general model for occlusal kinematics in conodonts.
Lethaia 47:547–555.
Mazza, M., and C. Martínez-Pérez. 2015: Unravelling conodont (Conodonta) ontogenetic
processes in the Late Triassic through growth series reconstructions and X-ray
microtomography. Bollettino della Società Paleontologica Italiana 54:161–186.
Mazza, M., M. Rigo, and M. Gullo. 2012a: Taxonomy and biostratigraphic record of the Upper
Triassic conodonts of the Pizzo Mondello section (western Sicily, Italy), GSSP candidate
for the base of the Norian. Rivista Italiana di Paleontologia e Stratigrafia 118:85–130.
Mazza, M., A. Cau, and M. Rigo. 2012b: Application of numerical cladistic analyses to the
Carnian–Norian conodonts: a new approach for phylogenetic interpretations. Journal of
Systematic Palaeontology 10:401–422.
Mazza, M., A. Nicora, and M. Rigo. 2018: *Metapolygnathus parvus* Kozur, 1972 (Conodonta): a
potential primary marker for the Norian GSSP (Upper Triassic). Bollettino della Società
Paleontologica Italiana.
Mazza, M., S. Furin, C. Spötl, and M. Rigo. 2010: Generic turnovers of Carnian/Norian
conodonts: Climatic control or competition? Palaeogeography, Palaeoclimatology,
Palaeoecology 290:120–137.
Miyashita, T., Coates Michael I., Farrar Robert, Larson Peter, Manning Phillip L., Wogelius Roy
A., Edwards Nicholas P., Anné Jennifer, Bergmann Uwe, Palmer A. Richard, and Currie
Philip J. 2019: Hagfish from the Cretaceous Tethys Sea and a reconciliation of the
morphological–molecular conflict in early vertebrate phylogeny. Proceedings of the
National Academy of Sciences 116:2146–2151.
Müller, K. J., and Y. Nogami. 1971: Über den Feinbau der Conodonten. Memoirs of the Faculty
of Science, Kyoto University. Series of geology and mineralogy 38:1–87.
Mullin, S. K., and P. J. Taylor. 2002: The effects of parallax on geometric morphometric data.
Computers in Biology and Medicine 32:455–464.
Nicoll, R. S. 1985: Multielement composition of the conodont species Polygnathus xylus xylus
Stauffer, 1940 and Ozarkodina brevis (Bischoff and Ziegler, 1957) from the Upper
Devonian of the Canning Basin, Western Australia. Bureau of Mineral Resources Journal
of Australian Geology and Geophysics 9:133–147.
———. 1987: Form and function of the Pa element in the conodont animal. Pp.77–90 *in* R. J.
Aldridge, ed. Palaeobiology of Conodonts. Ellis Horwood, Chichester.
Nicora, A., M. Balini, A. Bellanca, A. Bertinelli, S. A. Bowring, P. Di Stefano, P. Dumitrica, C.
Guaiumi, M. Gullo, A. Hungerbuehler, and others. 2007: The Carnian/Norian boundary
interval at Pizzo Mondello (Sicani Mountains, Sicily) and its bearing for the definition of
the GSSP of the Norian Stage. Albertiana 36:102–129.
Noyan, Ö. F., and H. W. Kozur. 2007: Revision of the late Carnian–early Norian conodonts from
the Stefanion section (Argolis, Greece) and their palaeobiogeographic implications.
Neues Jahrbuch für Geologie und Paläontologie-Abhandlungen 245:159–178.

Ogg, J. G., Z.-Q. Chen, M. J. Orchard, and H. S. Jiang. 2020: Chapter 25 - The Triassic Period.
Pp.903–953 *in* F. M. Gradstein, J. G. Ogg, M. D. Schmitz, and G. M. Ogg, eds. Geologic
Time Scale 2020. Elsevier.
Onofri, A. 2020: Aomisc: Statistical methods for the agricultural sciences. .
Pampush, J. D., J. M. Winchester, P. E. Morse, A. Q. Vining, D. M. Boyer, and R. F. Kay. 2016:
Introducing molaR: a New R Package for Quantitative Topographic Analysis of Teeth
(and Other Topographic Surfaces). Journal of Mammalian Evolution 23:397–412.
Pampush, J. D., J. Crowell, A. Karme, S. A. Macrae, R. F. Kay, and P. S. Ungar. 2019:
Technical note: Comparing dental topography software using platyrrhine molars.
American Journal of Physical Anthropology 169:179–185.
Pellegrom, C. 2019: Dental topographic analysis of maxillary and mandibular phyllostomid bat
dentitions: implications for dietary prediction in the fossil record. Masters Theses.
Pérez-Ramos, A., A. Romero, E. Rodriguez, and B. Figueirido. 2020: Three-dimensional dental
topography and feeding ecology in the extinct cave bear. Biology Letters 16:20200792.
Petryshen, W., C. M. Henderson, K. De Baets, and E. Jarochowska. 2020: Evidence of parallel
evolution in the dental elements of Sweetognathus conodonts. Proceedings of the Royal
Society B: Biological Sciences 287:20201922.
Prufrock, K. A., S. López-Torres, M. T. Silcox, and D. M. Boyer. 2016: Surfaces and spaces:
troubleshooting the study of dietary niche space overlap between North American stem
primates and rodents. Surface Topography: Metrology and Properties 4:024005.
Purnell, M. A. 1993: Feeding mechanisms in conodonts and the function of the earliest
vertebrate hard tissues. Geology 21:375–377.
———. 1994: Skeletal ontogeny and feeding mechanisms in conodonts. Lethaia 27:129–138.
———. 1995: Microwear on conodont elements and macrophagy in the first vertebrates. Nature
374:798–800.
Purnell, M. A., and P. H. von Bitter. 1992: Blade-shaped conodont elements functioned as
cutting teeth. Nature 359:629–631.
Purnell, M. A., and A. Evans. 2009: Conodont tooth complexity: quantification, convergence
with mammals, and implications for dietary analysis. 53:39–40.
Purnell, M. A., P. C. J. Donoghue, and R. J. Aldridge. 2000: Orientation and anatomical notation
in conodonts. Journal of Paleontology 74:113–122.
R Core Team. 2021: R: a language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.
Rannikko, J., H. Adhikari, A. Karme, I. Žliobaitė, and M. Fortelius. 2020: The case of the grass-
eating suids in the Plio-Pleistocene Turkana Basin: 3D dental topography in relation to
diet in extant and fossil pigs. Journal of Morphology 281:348–364.
Renaud, S., and R. Ledevin. 2017: Impact of wear and diet on molar row geometry and
topography in the house mouse. Archives of Oral Biology 81:31–40.
Robson, S. V. 2018: An analysis of North American taeniolabidoid multituberculate (Mammalia,
Allotheria) dentitions using mammalian dietary proxies.Masters Theses, University of
Calgary, Calgary, Albertapp.
Santana, S. E., S. Strait, and E. R. Dumont. 2011: The better to eat you with: functional
correlates of tooth structure in bats. Functional Ecology 25:839–847.
Savage, V. M., J. F. Gillooly, W. H. Woodruff, G. B. West, A. P. Allen, B. J. Enquist, and J. H.
Brown. 2004: The predominance of quarter-power scaling in biology. Functional Ecology
18:257–282.

Savriama, Y., C. Romestaing, A. Clair, L. Averty, J. Ulmann, R. Ledevin, and S. Renaud. 2022:
Wild versus lab house mice: Effects of age, diet, and genetics on molar geometry and
topography. Journal of Anatomy 240:66–83.
Schmidt, H. 1934: Conodonten-Funde in ursprünglichem Zusammenhang. Palaeontologische
Zeitschrift 16:76–85.
Scott, H. W. 1934: The zoological relationships of the conodonts. Journal of Paleontology
8:448–455.
Selig, K. R., S. López-Torres, E. J. Sargis, and M. T. Silcox. 2019: First 3D dental topographic
analysis of the enamel-dentine junction in non-primate euarchontans: contribution of the
enamel-dentine junction to molar morphology. Journal of Mammalian Evolution 26:587–
598.
Shirley, B., M. Grohganz, M. Bestmann, and E. Jarochowska. 2018: Wear, tear and systematic
repair: testing models of growth dynamics in conodonts with high-resolution imaging.
Proceedings of the Royal Society B: Biological Sciences 285:20181614.
Sieg, A. E., M. P. O'Connor, J. N. McNair, B. W. Grant, S. J. Agosta, and A. E. Dunham. 2009:
Mammalian Metabolic Allometry: Do Intraspecific Variation, Phylogeny, and Regression
Models Matter? The American Naturalist 174:720–733.
Spradley, J. P., J. D. Pampush, P. E. Morse, and R. F. Kay. 2017: Smooth operator: The effects
of different 3D mesh retriangulation protocols on the computation of Dirichlet normal
energy. American Journal of Physical Anthropology 163:94–109.
Stockey, C., P. C. J. Donoghue, T. H. P. Harvey, D. J. E. Murdock, and M. A. Purnell. 2021:
Multivariate dental topographic metrics demonstrate the dietary breadth and specialisms
of conodonts. :46–47.
Stockey, C., P. C. . J. Donoghue, T. H. P. Harvey, D. J. E. Murdock, and M. A. Purnell. 2022:
Multivariate dental topographic metrics demonstrate the dietary breadth and specialisms
of conodonts. .
Strait, S. G. 1997: Tooth use and the physical properties of food. Evolutionary Anthropology:
Issues, News, and Reviews 5:199–211.
Terrill, D. F., E. Jarochowska, C. M. Henderson, B. Shirley, and O. Bremer. 2022: Sr/Ca and
Ba/Ca ratios support trophic partitioning within a Silurian conodont community from
Gotland, Sweden. Paleobiology:1–21.
Vermeer, H. 2019: The association between dietary niche variation in rodents and climate
change across the Paleocene-Eocene Thermal Maximum.Masters Theses, Grand Valley
State Universitypp.
Villalobos-Chaves, D., and S. E. Santana. 2022: Craniodental traits predict feeding performance
and dietary hardness in a community of Neotropical free-tailed bats (Chiroptera:
Molossidae). Functional Ecology 36:1690–1699.
Vitek, N. S., P. E. Morse, D. M. Boyer, S. G. Strait, and J. I. Bloch. 2021: Evaluating the
responses of three closely related small mammal lineages to climate change across the
Paleocene–Eocene thermal maximum. Paleobiology 47:464–486.
de Vries, D., S. Heritage, M. R. Borths, H. M. Sallam, and E. R. Seiffert. 2021: Widespread loss
of mammalian lineage and dietary diversity in the early Oligocene of Afro-Arabia.
Communications Biology 4:1172.
Warton, D. I., R. A. Duursma, D. S. Falster, and S. Taskinen. 2012: smatr 3– an R package for
estimation and inference about allometric lines. Methods in Ecology and Evolution
3:257–259.

West, G. B., J. H. Brown, and B. J. Enquist. 1997: A General Model for the Origin of Allometric
Scaling Laws in Biology. Science 276:122–126.
White, C. R., and R. S. Seymour. 2003: Mammalian basal metabolic rate is proportional to body
mass2/3. Proceedings of the National Academy of Sciences 100:4046–4049.
Winchester, J. M., D. M. Boyer, E. M. St. Clair, A. D. Gosselin-Ildari, S. B. Cooke, and J. A.
Ledogar. 2014: Dental topography of platyrrhines and prosimians: Convergence and
contrasts. American Journal of Physical Anthropology 153:29–44.
Zhang, S., R. J. Aldridge, and P. C. J. Donoghue. 1997: An Early Triassic conodont with
periodic growth? J. Micropalaeontol. 16:65–72.
Zhang, Z. T., Y. D. Sun, P. B. Wignall, J. L. Fu, H. X. Li, M. Y. Wang, and X. L. Lai. 2018:
Conodont size reduction and diversity losses during the Carnian Humid Episode in SW
China. Journal of the Geological Society 175:1027–1031.

Table captions

Table 1: Numbers of conodont P1 element specimens by growth stages used for the study.

Abbreviations: GS – Growth stage.

Table 2: Linear regressions for platform length and platform area over element length for

Metapolygnathus communisti, *Epigondolella rigoi*, *Idiognathodus* sp. and *Gnathodus bilineatus*.

Data for *Idiognathodus* sp. and *G. bilineatus* were extracted from Purnell (1994) except for the

95% confidence intervals (CI). Slope coefficients and R^2 result from Reduced Major Axis

method. 95% CI values resulting from "sma" function calculation for *M. communisti* and

E. rigoi; from calculation for *Idiognathodus* sp. and *G. bilineatus*. The p-value results from the

"slope.test" function that compared the coefficients between species and isometry for

M. communisti and *E. rigoi*; from Z test (Hayami and Matsukuma 1970) for *Idiognathodus* sp.

and *G. bilineatus*.

Table 3: Parameters of the allometric equation $y = aX^b$ fitted to DNE values over element

dimensions for each studied conodont species. All values have been rounded to the second

significant digit.

Figure captions

815   Figure 1: Theoretical framework of the present study. Data from birds, mammals, reptiles,

816 fishes and amphibians were extracted from Glazier (2009); dashed lines represent 95% CI.

817  Figure 2: Growth stages of *Metapolygnathus communisti* and *Epigondolella rigoi*. For

818 *M. communisti*: **GS3** – specimen NA37_18; **GS4** – specimen NA37_12; **GS5** – specimen

819 NA37_01; **GS6** – specimen NA37_24. For *E. rigoi*: **GS2** – specimen NA59_20; **GS3** – specimen

820 NA59_19; **GS4** – specimen NA59_01; **GS5** – specimen NA59_05.

821  Figure 3: Measurements and morphological characters of P$_1$ elements illustrated on

822 *Metapolygnathus communisti* specimen NA37_24 and *Epigondolella rigoi* specimen NA59_05.

823 In most cases the convex side is also the long side. Morphological characters follow Mazza et al.

824 (2012*a*, *b*).

825  Figure 4: Log-transformed DNE values across growth stages GS3 to GS6 of *Metapolygnathus*

826 *communisti* and across growth stages GS2 to GS5 of *Epigondolella rigoi*. Names of the

827 specimens are specified in Figure 2.

828  Figure 5: **A-D.** Allometric growth of P$_1$ platform length and P$_1$ platform area for

829 *Metapolygnathus communisti* (n=27) and *Epigondolella rigoi* (n=23); each dot represents an

830 element and the colours are related to the growth stage. **A.** Platform length over element length

831 for *M. communisti*; convex platform length only. **B.** Platform area over element length for *M.*

832 *communisti*. **C.** Platform length over element length for *E. rigoi*; convex platform length only. **D.**

833 Platform area over element length for *E. rigoi*. **E-F.** Comparison of the allometric growth

834 between *Gnathodus bilineatus* (Purnell 1994), *Idiognathodus* sp. (Purnell 1993, 1994), *E. rigoi*

835 and *M. communisti*. **E.** Platform length over element length. Platform length in *M. communisti*

836 and *E. rigoi* was measured on the convex side. No data were available for *G. bilineatus* (Purnell

837 1994). **F.** Platform area over element length. **G.** Comparison of slope coefficients between

838 species; G1, Platform length over element length; G2, platform area over element length.

839  Figure 6: **A.** Distribution of DNE values across growth stages in *E. rigoi* and *M. communisti*.

840 **B.** DNE values over element length with fitting to the allometric equation $y = aX^b$. **C.** DNE

841 values over platform length with fitting to the allometric equation $y = aX^b$. **D.** DNE values over

842 platform area with fitting to the allometric equation $y = aX^b$. **E.** Comparison of scaling exponents

843 between species; E1, DNE values over element length; E2, DNE values over platform length;

844 E3, DNE values over platform area.

What scaling of metabolic rate can we assume in conodonts?

Analogy with other taxa

Metabolic rate

Body mass

Birds
Mammals
Reptiles
Fish and amphibians

Energy demand estimated from DNE

DNE

$y=x^{0.75}$

$y=x^{0.67}$

Element dimension

POSITIVE ALLOMETRY (Purnell 1994, 1995)

Metabolic Rate = Body Mass$^{> 0.67}$

log(platform length)

log(element length)

log(platform area)

log (element length)

Two possible scenarios under different assumptions

Metabolic rate

$y= x^{0.75}$

$y= x^{0.67}$

0.1   1   10   100   1000

Body mass

ISOMETRY

Metabolic Rate = Body Mass$^{0.67}$

log(platform length)

log(element length)

log(platform area)

log(element length)

0,1 mm

Metapolygnathus communisti

Epigondolella rigoi

GS2   GS3   GS4   GS5   GS6

Metapolygnathus communisti

Epigondolella rigoi

anterior

denticles

blade

anterior through margin

geniculation point

carina

platform nodes

cusp

platform

postcarinal nodes

platform area

platform length on the long side

platform length on the convex side

posterior

Metapolygnathus communisti

Scaled
Log DNE Value
Per Face

0.6000
0.1368
0.0312
0.0071
0.0016
0.0004
0.0001
0.0000
0.0000
0.0000

Edges

Outliers

0,1 mm

Epigondolella rigoi

GS2   GS3   GS4   GS5   GS6

Table 1: Numbers of conodont P1 element specimens by growth stages used for the study. Abbreviations: GS – Growth stage.

	Metapolygnathus communisti	*Epigondolella rigoi*
GS2	0	1
GS3	3	1
GS4	2	6
GS5	7	15
GS6	15	0

Table 2: Linear regressions for platform length and platform area over element length for each studied conodont species. Slope coefficients and R² result from the Reduced Major Axis method. The p-value results from the "slope.test" function that compared the coefficients between species and isometry.

Regression model	Taxon	Slope	Slope CI lower value	Slope CI upper value	R²	p-value (h0 = isometric growth)	Intercept	Intercept CI lower value	Intercept CI upper value
Platform length over element length	*Metapolygnathus communisti*	1.580	1.402	1.781	0.92	1.74×10^{-8}	-0.271	-0.361	-0.180
	Epigondolella rigoi	1.403	1.238	1.590	0.92	1.10×10^{-5}	-0.428	-0.538	-0.318
	Idiognathodus sp.	1.089	1.011	1.167	0.98	2.345 (Z test)			
Platform area over element length	*Metapolygnathus communisti*	2.561	2.230	2.940	0.89	1.039×10^{-3}	-1.376	-1.545	-1.206
	Epigondolella rigoi	2.392	2.080	2.750	0.90	1.443×10^{-2}	-1.229	-1.439	-1.019
	Idiognathodus sp.	2.149	2.002	2.296	0.98	2.060 (Z test)			
	Gnathodus bilineatus	2.164	2.020	2.308	0.98	2.329 (Z test)			

Table 3: Parameters of the allometric equation $y = aX^b$ fitted to DNE values over element dimensions for each studied conodont species. All values have been rounded to the second significant digit.

Regression model	Species	a	SE(a)	b	SE(b)	R²
DNE over element length	*Metapolygnathus communisti*	255.25	15.54	0.72	0.14	0.54
	Epigondolella rigoi	324.71	30.92	0.66	0.17	0.42
DNE over platform length	*Metapolygnathus communisti*	296.15	23.56	0.48	0.09	0.59
	Epigondolella rigoi	420.53	55.22	0.52	0.11	0.51
DNE over platform area	*Metapolygnathus communisti*	392.06	49.79	0.30	0.05	0.59
	Epigondolella rigoi	474.07	78.79	0.29	0.07	0.48