Towards rainy high Arctic winters: how ice-encasement impacts tundra plant phenology, productivity and reproduction

Mathilde Le Moullec[1], Anna-Lena Hendel[2,3], Matteo Petit Bon[4,5], Ingibjörg Svala Jónsdóttir[4,6], Øystein Varpe[4,7,8], René van der Wal[9], Larissa Teresa Beumer[1,10], Kate Layton-Matthews[1,8], Ketil Isaksen[11] and Brage Bremset Hansen[1,8].

1 Norwegian University of Science and Technology, Dept. of Biology, Centre for Biodiversity Dynamics; 2 University of Bayreuth, Faculty of Biology Chemistry & Earth Sciences; 3 University of Freiburg, Dept. of Wildlife Ecology and Management;4 The University Centre in Svalbard, Dept. of Arctic Biology; 5 Utah State University, Dept. of Wildland Resources; 6 University of Iceland, Institute of Life and Environmental Sciences; 7 University of Bergen, Dept. of Biological Sciences; 8 Norwegian Institute for Nature Research; 9 Agricultural University of Sweden, Dept. of Ecology; 10; Aarhus University, Dept. of Bioscience; 11 Norwegian Meteorological Institute.

Abstract

The Arctic is warming rapidly, with winters warming up to seven times as fast as summers in some regions. Warm spells in winter lead to more frequent extreme rain-on-snow events that alter snowpack conditions and can encapsulate tundra vegetation in 'basal ice' ('icing') for several months. However, tundra climate change studies have mainly focused on summer warming. Here, we investigate icing effects on vascular plant phenology, productivity, and reproduction in a pioneer field experiment in high Arctic Svalbard, simulating rain-on-snow and resultant icing in five consecutive winters, assessing vascular plant responses throughout each subsequent growing season. We also tested whether icing responses were modified by experimentally increased summer temperatures. Icing alone delayed early phenology of the dominant shrub, *Salix polaris*, but with evidence for a 'catch-up' (through shortened developmental phases and increased community-level primary production) later in the growing season. This compensatory response occurred at the expense of delayed seed maturation and reduced community-level inflorescence production. Both the phenological delay and allocation trade-offs were associated with icing-induced lags in spring thawing and warming of the soil, crucial to regulating plant nutrient availability and acquisition.

Experimental summer warming modified icing effects by advancing and accelerating plant

phenology (leaf and seed development), thus increasing primary productivity already early in

the growing season, and partially offsetting negative icing effects on reproduction. Thus,

winter and summer warming must be considered simultaneously to predict tundra plant

climate change responses. Our findings demonstrate that winter warm spells can shape high

Arctic plant communities to a similar level as summer warming. However, the absence of

accumulated effects over the years reveals an overall resistant community which contrasts

with earlier studies documenting major die-off. As rain-on-snow events will be rule rather

than exception in most Arctic regions, we call for similar experiments in coordinated

circumpolar monitoring programmes across tundra plant communities.

Keywords: basal ice, compensatory growth, extreme events, open top chamber, Svalbard,

Salix polaris, seasonality, rain-on-snow, trade-offs.

1. Introduction

Global warming comes with stronger and more frequent extreme climate events, prompting research activity aimed at determining their ecological and evolutionary consequences for terrestrial biota (Grant et al., 2017; Harris et al., 2018; IPCC, 2021; Robinson 2022; Trisos et al., 2020). In the Arctic, temperatures are rising three times as fast as the global average and the particularly pronounced warming trends during winter have resulted in increasingly frequent and widespread winter warm spells (AMAP, 2021; Bintanja & Andry, 2017; Graham et al., 2017; Vikhamar-Schuler et al., 2016; You et al., 2021). Despite evidence for possible severe consequences of these warm spells on vegetation and carbon balance (Bjerke et al., 2017; Bokhorst et al. 2008, 2009, 2011, 2012, 2022; Phoenix & Bjerke, 2016; Treharne et al., 2019), studies have almost exclusively focused on the effects of rising summer temperatures and changes in snow depth (Bjorkman et al., 2020; Collins et al., 2021; Elmendorf et al., 2012; Prevéy et al., 2019; Rixen et al., 2022; Wipf and Rixen, 2010). Recently, studies comparing winter to summer warming effects on plant traits stress the neglected importance of winter temperatures for fine-scale regional patterns and traits sensitive to cold involved in resource acquisition and conservation (Bjorkman et al., 2018; Niittynen et al., 2020). Given the key role that tundra ecosystems play for global carbon cycling and climate feedbacks (Chapin et al., 2005; Zhang et al., 2020), it is now urgent to better understand how the addition of extreme warm spells outside the growing season shape Arctic plant communities.

Warm spells in winter often come with "rain-on-snow" events, i.e., when precipitation falls as rain instead of snow, often transforming the entire snowpack (AMAP, 2021; Bokhorst et al., 2016; Pan et al., 2018; Rasmus et al., 2018). Under such mild conditions, the snowpack can melt completely, exposing plants to thaw-freeze cycles that may directly cause vegetation damage (Bokhorst et al., 2009, 2011). Alternatively, and especially common in the high Arctic, rain and resulting meltwater can collect and freeze on the ground, releasing latent

heath transferred to the soil and forms a layer of basal ice (hereafter referred to as 'icing') up

to several decimetres thick (Peeters et al., 2019; Putkonen & Roe, 2003). Low-growing Arctic

vegetation can remain entirely encapsulated in this ice layer for several months, until the onset

of spring. Icing has the potential for ecosystem-wide consequences by directly or indirectly

affecting several trophic levels, e.g. by reducing soil arthropod abundances (Coulson et al.,

2000), disrupting snow conditions for small mammals' survival and reproduction (Kausrud et

al., 2008), and causing starvation-induced die-offs in large herbivores, which in turn can

influence predators and scavengers (Hansen et al., 2013; Sokolov et al., 2016). However, the

immediate and lasting effects of icing on high Arctic plants' productivity, reproduction, and

phenology – fuelling the tundra food web – remain largely unknown.

The increased occurrence of regional Arctic vegetation 'browning' (i.e., decrease in

expected 'greening' under warmer climate) has been associated to several factors, including

winter warm spells associated to thaw-freeze cycles (Bokhorst et al., 2008, 2009; Phoenix &

Bjerke, 2016), but also to icing (Bjerke et al., 2017, Milner et al., 2016, Vickers et al., 2016).

This contrasts with the large-scale 'Arctic greening' pattern associated to shrub expansion and

improved plant growing conditions (Epstein et al., 2012; Frost et al., 2019; Myers-Smith et

al., 2011; Walker et al., 2006). However, changes in winter conditions can also exacerbate the

effect of summer warming alone (Frei and Henry, 2021), thus adding complexity and spatial

variability for vegetation productivity trends (Berner et al., 2020; Frost et al., 2019; Kelsey et

al., 2021; Myers-Smith et al., 2020), pressing needs for teasing appart effects of icing.

Physiologically, icing can lead to plant cell death, either through frost damage or

stress-induced anoxia metabolite accumulation from anaerobic respiration (Bokhorst et al.,

2010; Crawford et al., 1994; Preece & Phoenix, 2014). Studies investigating the effects of

icing described that evergreen shrubs and lichens were negatively affected, either through

altered leaf physiology and/or shoot and individual mortality (Bjerke, 2011, 2017, 2018a,

2018b; Milner et al., 2016; Preece et al., 2012). In comparison, deciduous shrubs and

graminoids showed higher tolerance, yet, there was still evidence of damage and biomass

reduction (Bjerke et al., 2018b; Preece & Phoenix, 2014). In addition, icing can indirectly

affect the vegetation by altering soil thermal properties (Putkonen & Roe, 2003), delaying

plant phenology as seen under late snow-melt (Assmann et al., 2019; Frei and Henry, 2021;

Semenchuk et al., 2016), or alternatively, icing may protect plants against winter herbivory.

Plants facing sub-optimal conditions may channel more energy into either growth or

reproduction, modifying existing energy allocation trade-offs (Bazzaz et al., 1987; Jónsdóttir,

2011). Induced stress can enhance vegetative growth, thereby constraining or delaying other

reproductive responses, i.e., flower phenology and/or number (Chapin, 1991; Orcutt & Nilsen,

2000). Such responses are found in Arctic plants coping with shortened growing seasons (e.g.

late snowmelt, Cooper et al., 2011; Wipf & Rixen, 2010), damages from herbivory

(Jónsdóttir, 1991; Skarpe & Van der Wal, 2002), thaw-freeze events (Bokhorst et al., 2011),

as well as icing (Milner et al., 2016; Preece et al., 2012). In these latter studies simulating

icing, Milner et al. (2016) showed high mortality of apical meristems and entire shoots in the

evergreen shrub *Cassiope tetragona* in the high Arctic, but increased growth of surviving

shoots, at the cost of reduced flower production (Milner et al., 2016). Along these lines,

Preece et al. (2012) expected delayed leaf phenology following icing in the low Arctic, but

instead found accelerated leaf emergence in spring in *Vaccinium myrtillus* and *V. vitis-idaea*,

suggesting a reallocation of resources to the remaining living shoots, yet, the cost of flowering

was unclear. Species like *Empetrum nigrum* were resistante to icing (Preece et al., 2012).

Warmer summers may counteract expected negative effects of icing through greater

activation of growth hormones fluxes and enhanced nutrient cycling and availability (Chapin,

1991; Frei and Henry, 2021; Gornall et al., 2011; Sundberg et al., 2000). Again, responses

may vary across species and tissue types, i.e., if the tissues are directly or indirectly exposed

to icing in addition to summer warming. For instance, high Arctic perennial species

commonly produce flower buds the previous summer (Arft et al., 1999, Barrett and Hollister

2016), thus directly experiencing icing conditions, while vegetative tissues grow generally in

direct response to June-July temperatures (Van der Wal & Stien, 2014). Possible delay in

phenology under icing could be altered by summer warming, advancing more reproductive

than vegetative, as well as early than late phenophases (Collins et al., 2021), with greater

effects on early than late flowering species (Prevéy et al. 2019). However, regardless of

summer warming, the duration of phenophases tend to be fix (Barrett and Hollister 2016;

Collins et al., 2021; Semenchuk et al., 2016).

Plot-level field experiments are powerful tools to investigate fine-scale plant response to

simulated climate change scenarios (Bjorkman et al., 2020; Bjorkman et al., 2018; Elmendorf

et al., 2015). Here, we assessed how experimental winter rain-on-snow events and subsequent

icing affect productivity, reproduction, and phenology of a high Arctic plant community. To

achieve this, we simulated plot-level rain-on-snow and ice formation in winter over five

consecutive years. Using a full factorial design, we also increased summer temperatures via

open top chambers, following the protocol of the Arctic and Alpine International Tundra

Experiment (ITEX, Henry & Molau, 1997). During each subsequent growing season, we

determined plant species and community responses to treatments. We expected icing to cause

overall negative effects, i.e., to 1) delay the plants' phenology due to a delaying effect of icing

on thawing processes; 2) reduce the overall plant productivity ('browning'); and 3) reduce the

flower production due to re-allocation of resources to growth. We anticipated the addition of

summer warming to overall reduce negative effects of icing, while the net effects are difficult

to predict *a priori*, given the scarcity of research in this field, the many mechanisms involved,

and the likely variation across species and tissue types.

2. Methods

Study site

The high Arctic Svalbard (78°17`N, 16°02`E) is a region warming twice faster than the Arctic average and five-seven times faster than the global average (Isaksen et al. 2022b). The experiment was located in the valley of Adventdalen and lasted for five consecutive years (January 2016–August 2020). Near the study site, the annual mean air temperature recorded at Svalbard Airport (15 km away) was -5.9 °C for the reference period 1971-2000 (Hanssen-Bauer et al., 2019), and -2.2 °C for the study period. Local winter (December–February) temperature rise five to seven times faster than in summer (June–August; Isaksen et al., 2022), thus at a greater rate than at the circumarctic scale. The interior fjord and valley areas around Adventdalen are some of the driest in Svalbard, with an average annual precipitation of 196 mm for the reference period (Hanssen-Bauer et al., 2019) and 223 mm for the study period. In comparison to the reference period, median RCP8.5 projections for Svalbard anticipate an increase in annual temperature and precipitation of about 7-9 °C and 20-40%, respectively, towards 2100 (Hanssen-Bauer et al., 2019). The frequency of rain-on-snow events has increased dramatically since a regime shift around the year 2000 (Hansen et al 2014, Peters et al. 2019), and Svalbard currently experience conditions soon expected across large parts of the Arctic (AMAP, 2021; Bintanja & Andry, 2017).

Soil thermal conditions on Svalbard are linked to the underlying permafrost which warms at a rate of 1.4 °C per decade at the permafrost table (2 m depth), observed 6 km away from our study site between 2002–2018 (Etzelmüller et al., 2020). At the study site, the sub-surface (5 cm depth) soil temperature was on average 4.7 °C in June and 7.5 °C in July (see methodology below, Table S1a), the two most important months for plant growth (Van der Wal & Stien, 2014). The volumetric water content (i.e., soil moisture) is relatively low for mesic communities, drying throughout the summer season from about 45% in early June to

25% mid-August at 10 cm depth (Table S1a). Soil moisture and sub-surface temperature are

strongly negatively related (Table S1b).

The tundra plant growing season starts immediately after the soil-thaw onset (Descals

et al., 2020). Adventdalen, which is in the bioclimatic subzone C 'middle Arctic tundra'

(CAVM Team, 2003), has a growing season of approximately three months, starting in mid-

late May (Table 1). The plant community studied here was a mesic grass and moss-rich dwarf

shrub heath, dominated by the prostrate deciduous dwarf shrub *Salix polaris* and the grass

Alopecurus borealis. Other abundant vascular plants were the horsetail *Equisetum arvense*,

the rush *Luzula confusa*, the forb *Bistorta vivipara* and the grass *Poa arctica* (Fig. S1). The

most abundant bryophytes were *Sanionia uncinata*, *Tomentypnum nitens* and *Polytrichastrum*

spp. Nomenclature follows http://panarcticflora.org/ for vascular plants and Frisvoll and

Elvebakk (1996) for bryophytes.

Experimental design

The study design followed a full-factorial randomized block design with four

treatment combinations (2 levels of icing × 2 levels of warming), replicated three times within

each of the three blocks (Fig. 1). In summer 2015, the three blocks (150–780m apart) were

selected in homogenous mesic tundra. Within each block, we chose 12 plots of 60 × 60 cm

with similar communities (n = 36 plots, Fig. 1). The four treatments, icing (*I*), summer

warming (*W*), a combined winter icing and summer warming treatment (*IW*), and control (*C*),

were randomly assigned to plots. Plots received the same treatment each year, starting in

winter 2015–2016 (Table S2). This allowed us to investigate both short-term (one year) as

well as longer-term (up to five years) plant responses. However, not all traits and parameters

were measured in all years, hence some results are presented based only on a subset of years.

<u>Winter icing</u>: The experimentally formed ice was applied to *I* and *IW* plots within a

period of 2–3 days in January–February (Fig. 1a). Snow was carefully removed from all plots

to apply the same level of disturbance, and any occurrence of natural ice was recorded.

Natural ice was particularly common in 2017, with partial ice coverage occurring in all plots

(3 cm average thickness). Snow was immediately placed back on non-icing (*C* and *W*) plots,

while for *I* and *IW*, a 13 cm high wooden frame (60×60 cm) was placed around the plot and

gradually filled up with cold water (for 1-3 days) until it was full of solid ice (Fig. 1a),

following Milner et al. (2016). Frames had no influence on the snow accumulation pattern

later in winter and they were removed at the beginning of snowmelt. Snow- and ice-melt

occurred almost simultaneously during spring meltwater floods.

Summer warming: Right after snowmelt (Table S2), hexagonal open top chambers

(1.4×1.4 m basal diagonal) were deployed over the *W* and *IW* plots following the

specifications given by the ITEX protocol (Henry & Molau, 1997). To avoid confounding

factors, all plots were excluded from grazing herbivores during the snow-free season using

metal net cages (*I* and *C* plots) or nets on top of the open top chambers (*W* and *IW* plots;

mesh-size 1.9 cm $\times$ 1.9 cm, Fig. 1b). Open top chambers and nets were all deployed and

removed on the same day in the start and end of the growing season.

Field measurements

Soil temperatures were recorded with iButton loggers (type DS1921G-F5, $\pm$ 1.0 °C accuracy,

0.5 °C resolution) every 120 minutes in summer and 240 minutes in winter. Loggers were

placed in all plots at depths of both 2 cm and 5 cm. In 2018, we also recorded temperatures at

10 cm and 20 cm depth (not in *IW*). We estimated the onset of soil-thaw in spring for each

plot based on logger data at 5 cm depth; if soil sub-surface temperatures were $\geq$ 0 °C for a

minimum of 10 consecutive days, the first day of this period defined the soil-thaw onset. We

also monitored surface air temperature (5 cm high) with HOBO loggers (type U23-003/UA-

001; $\pm$ 0.2°C accuracy) in a limited number of plots in *C* and *W* from 15[th] of June to 1[st] of

September 2016-2018, every 30 min (Table S1a).

To assess treatment effects on plant production, we monitored the vegetation weekly

during the growing season by measuring Normalized Difference Vegetation Index (NDVI, a

proxy for plant productivity with a value from -1 to 1; Pettorelli et al., 2005). This was done at

the central part of each plot (30 cm diameter) with a Skye SpectroSense2+ handheld device.

NDVI typically increases in the first part of the growing season until reaching a maximum, to

then drop progressively in the second part of the growing season. We considered the peak

growing season to be the average day C plots reached the maximum NDVI (Table 1). As an

additional measure of plant productivity, we measured community-level vascular plant

species abundance (hereafter referred to as 'relative abundance') in the period after the peak

growing season (end of July, Table 1), in 2016–2019 (Table S2). To estimate relative

abundance, we used the point intercept methodology (Bråthen & Hagberg, 2004), with a 50 ×

50 cm frame elevated above the canopy (~20 cm high) and with 25 evenly distributed points.

The points were marked by crossings of double strings with the frame to give a 90º projection.

At each point, a wooden pin of 3 mm diameter was lowered down onto the moss layer and all

'hits' of vascular plants were recorded.

To further study vegetative responses of the dominant vascular species, we collected *S.*

polaris leaves (before leaf senescence, Table S2) to measure leaf area, leaf dry weight and

specific leaf area (SLA, the ratio of leaf area to dry mass). The protocol and sample size of

leaf collection varied between years (Table S2), preventing direct comparison of leaf-size

traits between years, but still enabling within year comparisons of *I*, *IW* and *W* effect sizes

relative to *C*. Leaves were kept moist after collection until scanned. Leaf area was measured

in imageJ 1.48 (Schindelin et al., 2015). Subsequently, leaves were oven-dried at 60 °C for

four days and weighed on an XS204 METTLER TOLEDO scale to the nearest 0.01 mg.

To assess treatment effects on flowering frequency, the total number of inflorescences

were counted for the species *S. polaris* (males and females [catkins])*, B. vivipara, A. borealis,*

L. confusa and P. arctica in mid-July each year (Table S2). We counted inforescences in a 50

× 50 cm frame (Fig. 1c), subdivided into 16 sub-squares.

To assess treatment effects on the phenology of the dominant vascular plant, we

measured the most advanced vegetative and reproductive phenological stages ('phenophases')

of *S. polaris* in each of 16 sub-squares per plot of the 50 × 50 cm frame. We used the

following vegetative phenophases: 1) leaves starting to unfurl, 2) leaves fully expanded, 3)

start of senescence, and 4) leaves fully senesced. The reproductive phenophases were 1)

distinct inflorescence buds visible, 2) buds recognisable as female or male, 3) receptive

stigmas (female) and open anther releasing pollen (male), 4) stigma and anthers withered, and

5) seed dispersal. The highest number of phenological monitoring rounds took place in 2018,

whereas no phenological monitoring was done in 2020 (Table S2).

Statistical analysis

We derived different metrics from the NDVI data by fitting a generalized additive

model (GAM) to each plot for each year, using the repeated measurements over a growing

season. We employed the function 'gam' from the mgvc package (Wood et al., 2015) in

software R-3.6.3 (R Core team, 2020), fitted with restricted maximum likelihood. Model fit

was good (visual interpretation and k-indices > 1), except for summer 2017 when

measurements stopped before the peak growing season, i.e., they were still in the increasing

phase (Table S2). Thus, we removed 2017 from all NDVI analyses. For all other years, we

predicted daily NDVI values for each plot and derived 1) the maximum NDVI value, and 2)

the day-of-year when the maximum NDVI was reached. We also computed 3) the cumulative

NDVI in the first part of the growing season (hereafter 'cumulative start') by summing daily

values from the first day of measurements to the day when *C* plot, on average, reached the

maximum NDVI, 4) the cumulative NDVI in the second part of the growing season (hereafter

'cumulative end') by summing daily values from the day when *C* plot, on average, reached

the maximum NDVI to the last measurement day of the season, and 5) the cumulative NDVI

across the growing season (hereafter 'cumulative total') by summing daily values from the

first to the last day of measurements. These cumulative NDVI metrics were then standardized

by the number of days over which the integral was computed. This was necessary because the

field season length differed between years, but not between treatments within year.

We used (generalized) linear mixed-effect models as the main tool to analyse our data

and account for the hierarchical spatial and temporal structures of our study design, using the

functions 'lmer' and 'glmer' from the lme4 R package (Bates et al., 2015). We built a separate

model for each response variable, i.e., soil temperature, onset of soil-thaw, NDVI (different

metrics), plant relative abundance, leaf size traits, inflorescence number, and phenophases.

We conducted four different models for each response variable. 1) We computed the

treatment effect sizes or predicted mean over years (treatment [category] was set as fixed

effect) and the temporal replication was accounted for by including year as a random

intercept, or day-of-year nested within year when measurements were also replicated within a

given year (e.g., NDVI). 2) We calculated the annual effect sizes or predicted means

('treatment [category] × year [factor]' interaction as fixed effects). 3) We tested for an overall

icing-warming interaction effect over years, (icing [0,1] × warming [0,1] interaction as fixed

effects) with 'year' or 'day-of-year' nested within 'year' as random intercepts. 4) We tested

for an annual icing-warming interaction effect, (icing [0,1] × warming [0,1] × 'year [factor]

interaction as fixed effects. For all models, the random intercept structure (i.e., variation in

means between replicated units) always included the nested plot (n = 36) structure within

blocks (n = 3). It also included sub-squares nested within plots (16 sub-squares per plot, n =

576) for measurements of leaf traits (in 2019 and 2020, Table S2), inflorescence count, and

phenology. In 2018, leaf traits included information on leaves nested within shoots (Table

S2). Statistical significance can be interpreted as when the 95% confidence interval (CI) of

estimates did not overlap the predicted mean of *C*.

Data from air and soil temperature and NDVI metrics were best summarized by a

normal distribution and count data from plant relative abundance and inflorescence counts by

a Poisson distribution. Leaf size traits were log-transformed to follow a normal distribution.

Each phenophase was converted to a binary data format to fit a logistic regression from a

binomial distribution. Percentage change reported were calculated for back-transformed

predicted means, in comparison to controls.

3. Results

Soil temperature and soil-thaw onset

The liquid water added during the simulated rain-on-snow on both icing (*I*) and icing $\times$ warming (*IW*) plots released latent heat, increasing sub-surface soil temperatures for three days, by on average 2.9 °C [2.3;3.5] (results are reported as mean effect sizes [95% CI] in comparison to controls [*C*] over 2016–2020, unless specified differently) (Fig. S2).

During spring snowmelt, sub-surface soil-thaw onset was delayed by 6.0 [2.2;9.9] days in *I* and 4.4 [0.4;8.5] days in IW, while it remained unaffected in warming (*W*) (-0.9 [-5.0;3.1] days, Fig. 2a). However, the sub-surface soil-thaw onset, and the period over which it occurred, varied considerably between years (Table 1). For instance, the thawing onset of *C* occurred on average on 30[th] of April in 2019, ranging over a period of 32 days, while it occurred on the 21[st] of Mai over a period of 6 days in 2020 (Table 1). At 20 cm depth (data only from 2018), soil-thaw onset was delayed by 9.0 [1.3;16.7] days in *I*, after staying at a constant temperature of -0.5 °C for about two weeks (i.e., 'the zero curtain period', Outcalt et al., 1990; Fig. 2b), resulting in lagged soil temperature increases in *I* (i.e., significantly lower during entire June, Fig. 2b).

The presence of open top chambers increased soil sub-surface temperatures by 0.9 [0.7;1.2] °C in *W* and 0.8 [0.6;1.1] in *IW* (at 2 cm depth in June-August, over 2016, 2018 and 2019) and surface air temperature (5 cm high, 2016-2018) by 0.8 [0.7;0.9] °C in *W* (Table S1a). The soil volumetric water content did not vary between treatments but varied between years (Table S1a). The first and last year of the experiment were the warmest summers (2016 and 2020), while 2017 was the coldest (Table 1). Soil was the wettest in 2020 and the driest in 2019 (Table S1).

Vegetative responses: NDVI metrics, relative abundance, leaf size traits and phenology

Right after soil-thaw onset, the NDVI curves increased rapidly until reaching a

maximum in mid-July in *C* (15th of July on average, Table 1 and Fig. 3a) and decreased

slowly thereafter (Fig. S3). The maximum NDVI timing was delayed by on average 4 [2;7]

days in *I*, and consistently so across years (Fig. 3i). This delay was reduced in *IW* in most

years (2 [-1;4]) while it was -3 [-5;0] days earlier in *W* (Fig. 3i and Table S4). Maximum

NDVI values themselves were higher in *I* and *IW* in the first year of treatment (2016), while

this effect vanish in later years (Fig. 3j and Table S4). The last year of treatment (2020) had a

high maximum NDVI compared to previous years (Fig. 3b and Table 1), where even *C*

reached values similar to those in *I* and *IW* in other years (Fig. S3, Table S4). Maximum

NDVI values were unaffected by *W* (Fig. 3j and Table S4).

The shape of the NDVI curves varied among treatments, particularly after the

maximum NDVI. This effect was captured by the NDVI cumulative metrics. For the first part

of the growing season, only *IW* increased the NDVI 'cumulative start', and remained high for

the second part of the growing season (NDVI 'cumulative end'), resulting in an overall

increased NDVI ('cumulative total') across the growing season (Fig. 3k-m, Table S4). In the

second part of the growing season, *I* caught-up with *IW*, and thus also exceeded the NDVI

'cumulative end'. NDVI cumulative metrics of *W* did not differ (Fig. 3b). The treatment effect

sizes of NDVI metrics decreased over the years (except for the timing of the maximum

NDVI, Table S4), but still were of comparable magnitude as annual variation in *C* (Fig. 3).

Vascular plants' relative abundance was unchanged under *I* (3% [-11;19] change, Fig.

3n and Table S5), with a slight tendency for higher abundance of *S. polaris* (21% [-17;75]).

However, under *IW*, the relative abundance was on average 26% [9;46] higher, with large

effects on the first treatment year (2016, 69% [43;99] change, Fig. 3n and Table S5). These

patterns were largely consistent with the NDVI metric for the entire growing season

('cumulative total') and driven by the two most abundant vascular plants, the shrub *S. polaris*

and the graminoid *A. borealis* (Fig. S4), which relative abundance had a large influence on the

NDVI patterns (based on among-plot correlations, Table S3). *A. borealis* was even three times

more abundant in *IW* on the first treatment year (Fig. S4 and Table S5). Under *W*, the relative

abundance tended to increase by 13% [-2;31], a change not related to *S. polaris* which

decreased by -32% [-54;-1] (Figs. S3 and Table S5). The vascular plant' species community

composition remained overall unchanged across the study period (Fig. S5).

The SLA of *S. polaris* was higher in *I*, and this difference significantly increased over

the years (Fig. 3o and Table S8), being particularly pronounced from 2018 onwards. In those

years (2018-2020) dry weight decreased more than the area, with a respective change of -19%

[-28;-1] and -14% [-28;4] (Table S6). Although the leaf area and dry weight tended to be

lower in all treatments and all years, their ratio (i.e., SLA) was not different in *IW* and *W*, with

large inter-annual variation (Figs. 3o, S6 and Table S6).

No significant interaction effect was found between *I* and *W* in vegetative traits, except

for SLA in 2020 and early leaf phenology (see below), i.e., there were no strong indications of

an overall modifying effect of summer warming on the response to icing (Tables S4-S6).

Reproductive response: inflorescence production

The total number of vascular plant inflorescences in this mesic community was significantly

reduced across all treatments, and this particularly strongly over the years with all types of

measurements (2016, 2018 and 2019), showing a reduction by one third (Fig. 3p and Table

S7). Yet, *I* had no effect in the first and last treatment year (i.e. 2016 and 2020, Fig. 3p, Table

S7). Although *S. polaris* produced the highest number of inflorescences (catkins and male

flowers), and drove this overall pattern of annual variation, inflorescences of the forb *B.*

vivipara and graminoids were severely impacted by *I* (and *IW*) in all years (respectively by -

52% [-72;-27] and -67% [-95;-11] reduction, Fig. S7 and Table S7). Under *W*, *S. polaris*

inflorescences were also strongly reduced (-70% [-81;-56]), while graminoid inflorescences

were enhanced (146% [46;225]). Note that in 2020, there was a record number of

inflorescences in the community, with more than twice the amount compared to any other

study year (Fig. 3h, S7 and Table 1). For instance, even if the greatest reduction of

inflorescences under *W*, occurred in 2020 (Fig. 3p), after a progressive decrease over time

(Table S8), it was still the year with the greatest inflorescence production under this treatment

(Fig. S7 and Table S7).

The interaction effect of *I* and *W* was positive, in other words, there were more

inflorescences under *IW* than expected if the effects of *I* and *W* were additive (Table S7).

Phenological responses

Early phenophases of *S. polaris* leaves developed later in *I*, while they developed

earlier in *IW* and *W* (i.e., when 50% of the *C* reached the phenophase of unfurled leaf, 10% in

I and 90% in *IW* and *W* did so, Fig. 5a). Hence, the unfurled stage was reached 4 [1;8] days

later in *I* and -4 [-8;-1] days earlier in *IW* and *W* (Fig. S8). Despite this delay in *I*, the duration

to unfold leaves (from unfurled to fully expanded) was shortened by one third compared to *C*

and the other treatments (-3 [-4;-1] days shorter, Fig. 5a-b and Table S10). This resulted in

leaves being fully expanded with no apparent delay under *I* (2 [-2;5] days), but -5 [-8;-2] days

earlier in *IW* and *W* (i.e., when less than 10% of the *C* and *I* reached the fully expanded stage,

all plots had already reached it *IW* and *W*, Figs. 5b, S8, S9, Table S9 and S10). However, the

onset of leaf senescence happened almost simultaneously across treatments, and this pattern

occurred consistently across years (Figs. 5c-d, S8 and S9). Yet, this timing of leave

senescence varied among years with more than 10 days in *C* (Fig. S8a). The duration of

senesced was unchanged in *I*, while in *IW* and *W* it was 30% faster (-2 [-3;-1] days shorter).

Note that in contrast, *IW* and *W* leaves stayed 20% longer time fully expanded (6 [2;8] days;

Table S10).

Several *S. polaris* reproductive phenophases were reached later in *I* but earlier in *IW* and

W compared to *C* and these differences persisted throughout years, being particularly

pronounced in 2017 (Figs. 5 and S8b-c). The difference in timing was largest during seed

dispersal. In 2018, it occurred slightly later in *I* (5 [-2;12] days), but -11 [-17;-6] and -5 [-

12;2] days earlier in I*W* and *W* respectively (i.e., when 50% on the *C* reach seed dispersal,

19% in *I,* 96% in *IW* and 86% in *W* had reached it; Figs. 5 and S9). The ealrier seed dispersal

in *IW* coincided with a 30% shorter duration for seeds to mature (-8 [-14;-1] days shorter),

while the stage of stigma being receptive to pollen was extended by 50% compared to *C* (2

[0;5] days longer; Table S10).

Male flower anthers were visible and tended to released pollen later in *I* (although non-

significantly in 2018, 2 [-3;7] days, but see Fig. S8c for other years), earlier in *IW* (-5 [-10;-

0.4] days) and tended to be earlier in *W* (-4 [-10;2] days) (i.e., when 50% of C reached the

pollen released stage, 27% in *I*, 94% in *IW* and 67% in *W* had reached it; Figs. 5j-k, S8c and

Table S4), However, anther senescence was rather synchronous across treatments in 2018

(Fig. 5l). This suggested a double longer duration of pollen release in *IW* (5 [1;10] days

longer), but a 40% shorter duration in *I* (-2 [-3;0] days shorter; Table S10).

There was evidence of a positive interaction effect between *I* and *W* for the phenophases

of early leaf development, seed dispersal and anther development. The addition of warming

reversed the effect of *I,* advancing phenophases even more than under *W* alone (Table S9).

4. Discussion:

This high Arctic tundra experiment has demonstrated how icing can initially delay, but eventually enhance plant growth of a mesic community later in the growing season (Fig. 3i), while affecting reproduction (i.e., inflorescence production, Fig. 3p). This was confirmed at the species level, where the vegetative phenology of the dominant vascular plant, the dwarf shrub *S. polaris*, was delayed in early summer (Fig. 4a-b), but this seemed compensated for through an accelerated leaf development, increased SLA (Fig. 3o) and, to some extent, higher relative abundance. Seed maturation was however delayed (Fig. 4i). These observed species- and community-level delays in phenology and productivity due to icing appeared to result from later spring thawing processes, with a month-long lag in soil warming in the upper active layer (Fig. 2). The addition of summer warming partly counteracted icing effects (i.e., additive effects), but could also reverse the delayed phenology of icing by advancing it even more than under warming alone (i.e., interaction effects). This resulted in a greater increase of primary production, while also markedly advancing e.g., seed dispersal. This first multiyear tundra icing experiment also indicates a rather resistant vegetation community to a warming climate, with the absence of dramatic responses or amplified treatment effects across years.

Delayed but increased primary production: evidence of compensatory growth?

The enhanced above-ground community-level productivity and apparent absence of major die-off due to icing contrasts with the documented 'Arctic browning' following extreme winter warm spells, a phenomenon related to damage and mortality in evergreen shrubs, and associated reductions in measured primary production (Bjerke et al., 2017; Frost et al., 2019; Phoenix & Bjerke, 2016). These differences may be explained by contrasting overwintering strategies of the dominant growth forms in the different studies. For instance, evergreen shrubs, virtually lacking in our study plots, maintain long-lived leaves and dormant buds well above-ground, making them more dependent on insulating snow cover and more exposed to

direct damage from icing than deciduous shrubs (Givnish, 2002). Dead shoots of evergreen

shrubs will remain for several years, and even if growth may be enhanced in remaining shoots

(Milner et al. 2016), the net result is likely a decrease in NDVI (Bjerke et al., 2017; Treharne

et al., 2020). In contrast, in our experiment dominated by the deciduous shrub *S. polaris*, there

was no evident sign of dead shoots, likely allowing compensatory growth to enhance NDVI

later in the growing season. Hence, despite the severity of the consecutive treatments

experienced, the vegetation community studied here seemed rather resistant to the increasing

frequency of icing, possibly even enhancing 'Arctic greening'.

In the 'race' for high primary production within the reduced time window available in

icing plots (i.e., due to delayed soil processes), *S. polaris* appears to compensate for delayed

leaves phenology by accelerating the rate of leaves development while reducing leaf dry

weight and area. Because the dry weight decreased more than the area, leaf SLA (i.e., the

inversed ratio) was increased over time. Leaf SLA is one of the leaf resource capture and

retention traits sensitive to winter temperatures (Bjorkman et al., 2017), also used as indicator

of leaf growth-rate (Pérez-Harguindeguy et al., 2013). Our findings may reveal increased

growth rate from direct icing stress on the overwintering buds or due to the compressed

growing season. These results agree with a laboratory icing experiment on *S. polaris* that also

found reduced leaf sizes, as well as increased leaf numbers (not measured here), while

photosynthetic capacity remained unchanged (Bjerke et al., 2018b). The possible increased

number of leaves supports that this shrub has a remarkable ability to activate dormant buds to

compensate for damages, as also shown in response to herbivory (Skarpe & Van der Wal,

2002). Furthermore, *Salix polaris* has most of its structure and reserves below-ground,

accumulated in late summer as secondary growth (i.e. ring growth). Rain-on-snow events

were found to reduce ring growth in coastal areas of Svalbard frequently exposed to icing (Le

Moullec et al., 2020). Therefore, greater primary growth may come at the cost of reduced

secondary growth, which could alter the otherwise strong correlation between primary and

secondary growth in this species (Le Moullec et al., 2019).

The observed phenological delay under icing was likely caused by the extensive icing-induced lag in the warming of the upper soil layers, despite an almost simultaneous melting of ice and snow across treatments. This can be explained by infiltration and refreezing of meltwater in the underlying soil, resulting in higher ice content in near-surface soils. The thermal soil properties are strongly dependent on ice and water content. Ice-rich/wet soils require more accumulated energy to warm up than quickly thawed/dry soils, especially with presence of underlying permafrost (Isaksen et al., 2022a). Thereafter, the addition of summer warming shortened the lag in soil temperature increased found under icing.

As we expected, additive effects of summer warming (partly) counteracted negative effects of icing alone, but also with evidence of positive interaction effects. For instance, leaf phenology of *S. polaris* was instead advanced, tending to increase primary production already early in the growing season. Results from the low Arctic, where summer temperatures are much higher, match those found for our combined icing-warming treatment, i.e., simulated icing also advanced shrubs' early leaf emergence, explained by compensatory mechanisms from frost damaged ramets (Preece et al., 2012; Preece et al., 2014). Later in the growing season, the primary production in this treatment remained high. This is likely occurring because leaves from *S. polaris* stayed fully expanded 20% longer time, and the relative abundance of the dominant graminoid *A. borealis* increased, compared to controls. In another high Arctic experiment, grasses were the growth form most rapidly responding to experimentally increased soil temperature, increasing their total live plant biomass (Brooker & van der Wal, 2003). Although there was no change in primary production under icing-warming in 2020, this can be caused by NDVI reaching saturation across treatments and controls, as expected with values approaching 0.8 (Myers-Smith et al., 2020).

492   The greater primary production in the second part of the growing season under icing

493 and icing-warming combined occurred despite of a synchronous leaf senescence across

494 treatments. This synchronicity may in part be driven by cues fixed in time, e.g., photoperiod.

495 However, the annual variation in senescence time suggests the presence of interactions

496 between fix cues and other variables shared across treatments, e.g., snow/ice-melt timing

497 (Arft et al., 1999; Bjorkman et al., 2015; Cooper et al., 2011; Kelsey et al., 2021; Wipf &

498 Rixen, 2010). A slight delay in leaf senescence under simulated warming alone was also

499 detected in other Arctic sites, yet the main circumpolar pattern was the absence of change

500 (Bjorkman et al., 2020; Collins et al., 2021).

501 *Negative effects on reproduction: a result of trade-offs?*

502 Compensatory vegetative growth may come at the cost of reproduction, which has been

503 observed after thaw-freeze cycles (Bokhorst et al., 2011), snow manipulation (Cooper et al.,

504 2011), nutrient addition (Petraglia et al., 2013), and icing of evergreen shrubs (Milner et al

505 2016). The reduced inflorescence numbers due to icing (halved in some years) could result

506 from prioritized allocation to vegetative growth, if not from direct winter damages. In the first

507 experimental year, no inflorescence reduction occurred for *S. polaris*, suggesting that

508 inflorescence buds formed the previous summer survived a winter of icing. Other species,

509 e.g., *B. vivipara* and *A. boralis,* exhibited a strong reduction in inflorescence numbers without

510 a lag. In an experiment conducted in the same valley, a record low number of inflorescences

511 was associated with extreme natural rain-on-snow events in winter 2012 (Semenchuk et al.,

512 2013). Furthermore, shorter time between green-up and flowering (i.e., green-up being more

513 delayed than flowering) can lead to lower investment into flowering (Gougherty and

514 Gougherty, 2018; Collins et al., 2021). Hence, similarly to other Svalbard studies, this

515 resulted in delayed flower phenology and seed dispersal, as well as reduced flowering

516 (Cooper et al., 2011; Karlsen et al., 2014; Semenchuk et al., 2013; Semenchuk et al., 2016).

The addition of summer warming to icing reduced the negative effects of icing on

reproductive traits to a similar level or even beyond the rates obtained from warming alone.

Advancing flowering and fruiting due to warming are consistent with results of simulated

summer warming at a circumpolar scale (Bjorkman et al., 2020; Collins et al., 2021; Prevéy et

al., 2019). Thus, like Collins et al. (2021), we found a larger shift in reproductive than

vegetative phenophases, resulting in a shorter period between leaf emergence and seed

dispersal. However, contrary to their overall result and to a neighbour snow manipulation

experiment (Semenchuk et al., 2016), the duration of phenophases were not fixed. The

duration of inflorescence and seed development were both shortened under summer warming

(regardless of icing). Yet, the duration of the critical phases for reproduction, i.e., pollen

release and stigmas receptive to pollen, were doubled, and this, only under combined icing-

warming. Phenology studies rarely separate this short, but critical stage from the flowering

time, while our results suggested important implications for pollination success (Song and

Saavedra, 2018). Combining icing-warming thus indicates a high energy investment to

reproduction, despite the also high investment to vegetative growth. Exploring warming

effects through annual variation supported this, i.e., during the warm summer of 2020, we

observed simultaneously a record high number of inflorescences and NDVI maximum.

Evolutionary implications

Evolutionary consequences can sometimes be expected from changing trade-offs

influencing sexual reproductive structure. However, the plant species composing Arctic mesic

communities are generally perennial, allowing them to wait for favourable conditions to

reproduce sexually (Bazzaz et al., 1987). In unfavourable years, loss of those cohorts can be

buffered by seed banks, although for instance seeds from *S. polaris* are short-lived (Cooper et

al., 2004) and often need to germinate within the same year they are produced. Nevertheless,

S. polaris, as most Arctic species, also reproduces asexually by vegetative clonal growth

(Jónsdóttir, 2011), and if clone size would increase, the subsequent self-pollination increase

could limit evolutionary consequences of decreased seed production from environmental

changes (Barrett, 2015). Furthermore, the lack of accentuated effects after five consecutive

years of extreme winter conditions indicates that species in such communities have evolved

strategies to cope with these conditions. Local adaptation should not be underestimated, as

shown by a high Arctic transplant experiment where endogenous population were better

adapted to local conditions than foreign populations, regardless of the drastic ~ 3°C increase

of the summer warming treatment (Bjorkman et al., 2017).

This overall resistant and resilient mesic community studied may already have adapted

to icing. In the oceanic climate of Svalbard, winter rain-on-snow events have indeed been

occurring at least for decades (Vikhamar-Schuler et al., 2016). However, their frequency and

extent have increased dramatically in recent years (Hansen et al., 2014; Peeters et al., 2019),

and the selection acting upon life-history traits might change when acute stress episodes from

occasional icy winters become 'chronic', as expected in the near future (Bintanja & Andry,

2017). Regions that have not yet encountered rain-on-snow events but are predicted to in the

near future (e.g., Canadian high Arctic), may experience more drastic effects, especially since

there would be no possible introgression of adapted genes from nearby habitat (Bjorkman et

al., 2017). Comparative 'space-for-time' studies including Arctic regions where rain-on-snow

events are still rare could shed light on plants' adaptive capacities to icing.

Ecological implications

The effect sizes of winter icing *versus* summer warming were of similar magnitude across

most traits and within the range of natural inter-annual variation, which is typically large in

the high Arctic. Although the use of open top chambers only increased summer surface air

and sub-surface soil temperatures by 0.8 and 0.9 °C, respectively, this increase was similar to

the decadal trend found in this region (Etzelmüller et al., 2020, Isaksen et al., 2022). It was

also comparable to other ITEX sites in the high Arctic, where irradiance is low, restricting the

warming effect (Bokhorst et al., 2013). There was no evidence of exacerbated treatments

effects over the years, as decadal simulated summer warming also indicated (Barrett and

Hollister, 2016). The observed primary production increase under warming was therefore

limited. Other tundra studies/experiments that have investigated effects of altered winter

conditions, notably snow accumulation, also found similar or larger effect sizes of the winter

than the summer season (Cooper et al., 2011; Niittynen et al., 2020; Wahren et al., 2005). Our

study thus supports the overall insight that conditions during the 'dormant season' can have a

substantial effect on the growth of tundra vegetation.

In contrast to evergreen shrubs communities, the community studied here represent

important resources for Arctic herbivores (Chapin III et al., 1996). Thus, the affected

phenology and unexpected positive effect of icing on primary production may have potential

implications for the whole ecosystem, such as through plant-herbivore interactions (e.g.

phenological mismatches, Gauthier et al., 2013; Post & Forchhammer, 2008; Post et al.,

2009), carbon fluxes (Schuur et al., 2015) and vegetation-climate feedbacks (Zhang et al.,

2020). Tundra icing may thus have counteracting effects on herbivores, directly by limiting

food accessibility during winter (Forbes et al., 2016; Hansen et al., 2013), but also indirectly

by affecting – and possibly increasing – the food quantity during summer.

Conclusion and outlooks.

High Arctic environments are naturally variable, and plants have developed elaborate

strategies to persist under these extreme conditions. Nevertheless, this study has shown that

winter rain-on-snow, and associated tundra icing, can have similarly large impacts as summer

warming. On the occasion of the 30-year anniversary of ITEX experiments conducted across

the poles, providing critical knowledge regarding tundra responses to summer warming

(Bjorkman et al., 2018; Collins et al., 2021; Prevéy et al., 2019), we call for the

implementation of similar large-scale experiments investigating effects of winter warming,

and rain-on-snow events in particular. Further exploration can be numerous: across

communities, plant functional types, and species, as well as modulating the timing, severity,

and duration of these events (Bokhorst et al., 2011; Preece & Phoenix, 2014), in interaction

with different levels of summer warming, snow/ice-melt timing (Kelsey et al., 2021) and

herbivory (Gornall et al., 2009). Such a coordinated effort on long-term effects is required to

gain a holistic understanding of future changes in the tundra biome and its possible resistance

and resilience to both winter and summer warming (Bjorkman et al., 2020). This is especially

urgent given that winter warming is by far outpacing summer warming (Isaksen et al., 2022b),

with potentially ecosystem-wide consequences such as alteration of below-ground vegetation

productivity, microorganism activity, and carbon flux exchange, and cascading effects across

the food web. This pioneer long-term icing experiment sets the stage for forthcoming studies

to fill this fundamental knowledge gap.

Acknowledgements

This project was funded by the Research Council Norway (FRIPRO 276080 and SFF-III

223257) and Svalbard Environmental Protection Fund (project 16/113). We thank UNIS

logistics for logistical support and Longyearbyen Lokalstyre for permission to perform the

experiment (RiS ID: 10484). We are also deeply grateful for all the help in the field from

Solvei B. Hovdal, Martha Grotheim, Hanne K. Haraldsen, Julia Greulich, Karrín Björnsdóttir,

Kristine Valøen, Robin Zweigel, Ådne Nafstad, Marit Arneberg, Kristiane Midhaug, Marte

Søreng, Marianne Angård, Hamish Burnett, Lia Lechler, Charlotte Karlsen, Kjerstin

Hilmarsen, Lukas Tietgen, and Svea Zimmermann.

Table 1. Overview of annual mean biotic and abiotic measurements. The daily average air temperatures were recorded at the Svalbard Airport

(available on https://seklima.met.no/), while the soil-thaw onset day, maximum Normalized Difference Vegetation Index (max. NDVI) metrics,

and the community inflorescence counts (Flower count) were recorded in control plots. Predicted means are presented with their 95% confidence

intervals (CI), or range of values. The 'time of max. NDVI' is interpreted as the peak growing season. DOY = Day-of-year (i.e. Julian day, where

1 = 1 January).

	2016	2017	2018	2019	2020
Soil-thaw onset day (range)	13th May (8th–22rd May)	21th May (3rd May–1st June)	10th May (6th–14th May)	30th Apr (18th Apr–20th May)	18th May (16th–22th May)
Air T°C June (range)	5.0 (3.6–6.7)	4.6 (3.2–6.5)	4.0 (2.3–6)	4.8 (3.3–6.8)	4.5 (2.8–6.8)
July (range)	9.0 (7.4–11.1)	6.9 (5.4–8.9)	7.2 (5.6–9.5)	8.4 (6.5–10.6)	9.8 (7.7–12.2)
Time of max. NDVI, DOY [CI]	16th July 197 [189;205]	NA	19th July 199 [191;207]	12th July 192 [184;200]	15th July 196 [188;204]
Max. NDVI [CI]	0.74 [0.72;0.77]	NA	0.76 [0.73;0.78]	0.76 [0.73;0.78]	0.79 [0.76;0.81]
Flower count m^2 [CI]	263 [181;345]	229 [151;288]	314 [218;409]	295 [204;386]	491 [341;642]

Fig. 1. The experimental set up in the Adventdalen valley, Svalbard. Overview of the field

site (a) in the polar night, when applying the icing treatment by filling a wooden frame with

water, and (b) in summer with the open top chambers (plexiglass hexagons) simulating

warming and net cages to protect against herbivory. (c) Experimental full-factorial

randomized block design with the vegetation frame fitting the plot size. C = control, I = icing,

IW = icing × warming, W = warming. Picture credits: Ø. Varpe and M. Le Moullec.

Fig. 2. Daily estimates of soil temperatures during the soil-thawing period from mid-May to

end of June 2018, at (a) 5 cm depth (i.e., sub-surface), and (b) 20 cm depth. Shaded areas

represent 95% CIs and the orange vertical dashed lines shows the day the warming treatments

started.

Fig. 3. Effects of experimental treatments on several Normalized Difference Vegetation Index

(NDVI) metrics, relative abundance (abundance) of vascular plants, specific leaf area (SLA)

of *Salix polaris* and the community inflorescence count (flower count). (a-h) Model

predictions (black dots) and their 95% confidence intervals (CIs) for unmanipulated control

plots, separately for each year. The horizontal full and dashed lines represent the among-year

model predictions and their 95% CIs, which are only displayed for variables comparable

between years (i.e., with comparable sampling periods and/or design). Dots in the background

show the raw data at the plot (n = 9) or sample replication within plot (n range [109-214])

level, which were jittered for display purposes. Model predictions and their CIs were back-

transformed on the response scale prior presentation. (i-p) Effect sizes and their 95% CIs for

the effect of treatments, displayed separately for each year (left panel - year is used as a fixed-

effect in interaction with treatment in the model) and across all years (right panel - year is

used as a random-effect in the model). In the left panel, the reference level at 0 effect size

represents within-year unmanipulated controls (corresponding to the model predictions in a-h)

and effect sizes refer to within-year treatment effects. In the right panel, the reference level at

0 effect size represents across-year unmanipulated controls (corresponding to the horizontal

lines in a-h) and effect sizes refer to across-year treatment effects. Abbreviations as

follows: NDVI Time max. = Time of maximum NDVI as day-of-year; NDVI max. =

Maximum NDVI values from -1 to 1; NDVI cum. start = NDVI cumulative start. NDVI cum.

end = NDVI cumulative end. NDVI cum. total = NDVI cumulative total (see Material and

Methods for details).

Fig. 4. Probability curves of *Salix polaris* phenophases (i.e. the estimated probability of

reaching a specific phase) for vegetative phases (a-d) and reproductive phases (e-i: female

flowers; j-l: male flowers). Phenophases correspond to (a) leaves unfurled, (b) leaves fully

expanded, (c) leaves started senescing, (d) leaves fully senesced, (e) flower visible (female or

male), (f) stigma visible, (g) stigma receptive, (h) stigma senesced, (i) seed dispersed, (j)

anther visible, (k) pollen released, and (l) anther senesced. Estimates are shown for 2018, i.e.

the year when all phases could be estimated after the most intensive monitoring effort (for

other years see Fig. S8). The predicted estimates were computed with generalized linear

mixed-effect models, accounting for the replicated structure of the study design (i.e., random

effects), and the shaded areas represent the associated 95% CIs to the fixed-effects. Red tick

marks on the x-axis represent sampling days. See Fig. S9 and Table S9 for estimates of the

day at which each phenophases was reached (i.e., proportion of 1).

5. References

AMAP. (2021). *Arctic Climate Change Update 2021: Key Trends and Impacts. Summary for Policy-makers. Arctic Monitoring and Assessment Programme (AMAP)* , Tromsø, Norway. 16 pp.

Arft, A.M., Walker, M.D., Gurevitch, J., Alatalo, J.M., Bret-Harte, M.S., Dale, . . . Wookey, P.A. (1999), Responses of tundra plants to experimental warming: meta-analysis of the International Tundra Experiment. *Ecological Monographs*, 69: 491-511. https://doi.org/10.1890/0012-9615(1999)069[0491:ROTPTE]2.0.CO;2

Assmann, J. J., Myers-Smith, I. H., Phillimore, A. B., Bjorkman, A. D., Ennos, R. E., Prevéy, J. S., . . . Hollister, R. D. (2019). Local snow melt and temperature—but not regional sea ice—explain variation in spring phenology in coastal Arctic tundra. *Global Change Biology*. 2019; 25: 2258– 2274. https://doi.org/10.1111/gcb.14639

Barrett SC (2015) Influences of clonality on plant sexual reproduction. *Proceedings of the National Academy of Sciences, 112*(29), 8859–8866. https://doi.org/10.1073/pnas.1501712112

Barrett R. T., & Hollister R. D. (2016). Arctic plants are capable of sustained responses to long-term warming. *Polar Research*, 35. https://doi.org/10.3402/polar.v35.25405

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software, 67*(1), 1-48. https://doi.org/doi:10.18637/jss.v067.i01

Bazzaz, F. A., Chiariello, N. R., Coley, P. D., & Pitelka, L. F. (1987). Allocating Resources to Reproduction and Defense. *Bioscience, 37*(1), 58-67. https://doi.org/10.2307/1310178

Berner, L. T., Massey, R., Jantz, P., Forbes, B. C., Macias-Fauria, M., Myers-Smith, I., . . . Goetz, S. J. (2020). Summer warming explains widespread but not uniform greening in the Arctic tundra biome. *Nature Communications, 11*(1), 4621. https://doi.org/10.1038/s41467-020-18479-5

Bintanja, R., & Andry, O. (2017). Towards a rain-dominated Arctic. *Nature Climate Change, 7*(4), 263-267. https://doi.org/10.1038/nclimate3240

Bjerke, J. W. (2011). Winter climate change: Ice encapsulation at mild subfreezing temperatures kills freeze-tolerant lichens. *Environmental and Experimental Botany, 72*(3), 404-408. https://doi.org/https://doi.org/10.1016/j.envexpbot.2010.05.014

Bjerke, J. W., Elvebakk, A., & Tømmervik, H. (2018a). Alpine garden plants from six continents show high vulnerability to ice encasement. *Norsk Geografisk Tidsskrift - Norwegian Journal of Geography, 72*(1), 57-64. https://doi.org/10.1080/00291951.2017.1391876

Bjerke, J. W., Elverland, E., Jaakola, L., Lund, L., Zagajewski, B., Bochenek, Z., . . . Tømmervik, H. (2018b). High tolerance of a high-arctic willow and graminoid to simulated ice encasement. *Boreal Environment Research, 23*, 329–338.

Bjerke, J. W., Treharne, R., Vikhamar-Schuler, D., Karlsen, S. R., Ravolainen, V., Bokhorst, S., . . . Tommervik, H. (2017). Understanding the drivers of extensive plant damage in boreal and arctic ecosystems: Insights from field surveys in the aftermath of damage. *Science of the Total Environment, 599-600*, 1965-1976. https://doi.org/10.1016/j.scitotenv.2017.05.050

Bjorkman, A. D., García Criado, M., Myers-Smith, I. H., Ravolainen, V., Jónsdóttir, I. S., Westergaard, K. B., . . . Normand, S. (2020). Status and trends in Arctic vegetation: Evidence from experimental warming and long-term monitoring. *Ambio, 49*(3), 678-692. https://doi.org/10.1007/s13280-019-01161-6

Bjorkman, A. D., Myers-Smith, I. H., Elmendorf, S. C., Normand, S., Rüger, N., Beck, P. S. A., . . . Weiher, E. (2018). Plant functional trait change across a warming tundra biome. *Nature, 562*(7725), 57-62. https://doi.org/10.1038/s41586-018-0563-7

Bjorkman, A. D., Vellend, M., Frei, E. R., Henry, G. H. R. (2017). Climate adaptation is not enough: warming does not facilitate success of southern tundra plant populations in the high Arctic. *Global Change Biology*, *23*, 1540–1551, https://doi.org/10.1111/gcb.13417

Bokhorst, S., Bjerke, J. W., Bowles, F. W., Melillo, J., Callaghan, T. V., & Phoenix, G. K. (2008). Impacts of extreme winter warming in the sub-Arctic: growing season responses of dwarf shrub heathland. *Global Change Biology, 14*, 2603–2612. https://doi.org/10.1111/j.1365-2486.2008.01689.x

Bokhorst, S. F., Bjerke, J. W., Tømmervik, H., Callaghan, T. V., & Phoenix, G. K. (2009). Winter warming events damage sub-Arctic vegetation: consistent evidence from an experimental manipulation and a natural event. *Journal of Ecology, 97*(6), 1408-1415. https://doi.org/10.1111/j.1365-2745.2009.01554.x

Bokhorst, S., Bjerke, J. W., Davey, M. P., Taulavuori, K., Taulavuori, E., Laine, K., . . . Phoenix, G. K. (2010). Impacts of extreme winter warming events on plant physiology in a sub-Arctic heath community. *Physiologia Plantarum, 140*(2), 128-140. https://doi.org/10.1111/j.1399-3054.2010.01386.x

Bokhorst, S., Bjerke, J. W., Street, L. E., Callaghan, T. V., & Phoenix, G. K. (2011). Impacts of multiple extreme winter warming events on sub-Arctic heathland: phenology, reproduction, growth, and CO2 flux responses. *Global Change Biology, 17*(9), 2817-2830. https://doi.org/10.1111/j.1365-2486.2011.02424.x

Bokhorst, S., Bjerke, J. W., Tommervik, H., Preece, C. & Phoenix, G. K. Ecosystem response to climatic change: the importance of the cold season. *Ambio* 41, 246–255 (2012).

Bokhorst, S., Huiskes, A., Aerts, R., Convey, P., Cooper, E. J., Dalen, L., . . . Dorrepaal, E. (2013). Variable temperature effects of Open Top Chambers at polar and alpine sites explained by irradiance and snow depth. *Global Change Biology, 19*(1), 64-74. https://doi.org/10.1111/gcb.12028

Bokhorst, S., Pedersen, S. H., Brucker, L., Anisimov, O., Bjerke, J. W., Brown, R. D., . . . Callaghan, T. V. (2016). Changing Arctic snow cover: A review of recent developments and assessment of future needs for observations, modelling, and impacts. *Ambio, 45*(5), 516-537. https://doi.org/10.1007/s13280-016-0770-0

Bokhorst, S., Cornelissen, J. H. C., & Veraverbeke, S. (2022). Long-term legacies of seasonal extremes in Arctic ecosystem functioning. *Global Change Biology, 28*, 3161-3162. https://doi.org/10.1111/gcb.16078

Bråthen, K. A., & Hagberg, O. (2004). More efficient estimation of plant biomass. *Journal of Vegetation Science, 15*(5), 653-660. https://doi.org/10.1111/j.1654-1103.2004.tb02307.x

Brooker, R., & van der Wal, R. (2003). Can soil temperature direct the composition of high arctic plant communities? *Journal of Vegetation Science, 14*(4), 535-542. https://doi.org/https://doi.org/10.1111/j.1654-1103.2003.tb02180.x

CAVM Team. (2003). *Circumpolar Arctic Vegetation Map. (1:7,500,000 scale), Conservation of Arctic Flora and Fauna (CAFF) Map No. 1. .* U.S. Fish and Wildlife Service.

Chapin III, F. S. (1991). Integrated Responses of Plants to Stress. *Bioscience, 41*(1), 29-36. https://doi.org/10.2307/1311538

Chapin III, F. S., Bret-Harte, M. S., Hobbie, S. E., & Zhong, H. (1996). Plant functional types as predictors of transient responses of arctic vegetation to global change. *Journal of Vegetation Science, 7*(3), 347-358. https://doi.org/https://doi.org/10.2307/3236278

Chapin III, F.S., M. Sturm, M.C. Serreze, J.P. McFadden, J.R. Key, A.H. Lloyd, A.D. McGuire, T.S. Rupp, et al. 2005. Role of landsurface changes in Arctic summer warming. *Science* 310: 657–660. https://doi.org/10.1126/science.1117368.

Collins, C. G., Elmendorf, S. C., Hollister, R. D., Henry, G. H. R., Clark, K., Bjorkman, A. D., . . . Suding, K. N. (2021). Experimental warming differentially affects vegetative and reproductive phenology of tundra plants. *Nature Communications, 12*(1), 3442. https://doi.org/10.1038/s41467-021-23841-2

Cooper, E. J., Alsos, I. G., Hagen, D., Smith, F. M., Coulson, S. J., & Hodkinson, I. D. (2004). Plant recruitment in the High Arctic: Seed bank and seedling emergence on Svalbard. *Journal of Vegetation Science, 15*(1), 115-124. https://doi.org/10.1111/j.1654-1103.2004.tb02244.x

Cooper, E. J., Dullinger, S., & Semenchuk, P. (2011). Late snowmelt delays plant development and results in lower reproductive success in the High Arctic. *Plant Science, 180*(1), 157-167. https://doi.org/https://doi.org/10.1016/j.plantsci.2010.09.005

Coulson, S. J., Leinaas, H. P., Ims, R. A., & Søvik, G. (2000). Experimental manipulation of the winter surface ice layer: the effects on a High Arctic soil microarthropod community. *Ecography, 23*(3), 299-306. https://doi.org/10.1111/j.1600-0587.2000.tb00285.x

Crawford, R. M. M., Chapman, H. M., & Hodge, H. (1994). Anoxia Tolerance in High Arctic Vegetation. *Arctic and Alpine Research, 26*(3), 308-312.

Descals, A., Verger, A., Filella, I., Baldocchi, D., Janssens, I. A., Fu, Y. H., . . . Peñuelas, J. (2020). Soil thawing regulates the spring growth onset in tundra and alpine biomes. *Science of the Total Environment, 742*, 140637. https://doi.org/https://doi.org/10.1016/j.scitotenv.2020.140637

Elmendorf, S. C., Henry, G. H. R., Hollister, R. D., Björk, R. G., Bjorkman, A. D., Callaghan, T. V., . . . Wookey, P. A. (2012). Global assessment of experimental climate warming on tundra vegetation: heterogeneity over space and time. *Ecology Letters, 15*(2), 164-175. https://doi.org/doi:10.1111/j.1461-0248.2011.01716.x

Elmendorf, S. C., Henry, G. H. R., Hollister, R. D., Fosaa, A. M., Gould, W. A., Hermanutz, L., . . . Walker, M. D. (2015). Experiment, monitoring, and gradient methods used to infer climate change effects on plant communities yield consistent patterns. *Proceedings of the National Academy of Sciences, 112*(2), 448. https://doi.org/10.1073/pnas.1410088112

Epstein, H. E., Raynolds, M. K., Walker, D. A., Bhatt, U. S., Tucker, C. J., & Pinzon, J. E. (2012). Dynamics of aboveground phytomass of the circumpolar arctic tundra during the past three decades. *Environmental Research Letters, 7*(1), 015506. https://doi.org/10.1088/1748-9326/7/1/015506

Epstein, H. E., Walker, M. D., Chapin Iii, F. S., & Starfield, A. M. (2000). A transient, nutrient-based model of arctic plant community response to climatic warming. *Ecological Applications, 10*(3), 824-841. https://doi.org/https://doi.org/10.1890/1051-0761(2000)010[0824:ATNBMO]2.0.CO;2

Etzelmüller, B., Guglielmin, M., Hauck, C., Hilbich, C., Hoelzle, M., Isaksen, K., . . . Ramos, M. (2020). Twenty years of European mountain permafrost dynamics—the PACE legacy. *Environmental Research Letters, 15*(10), 104070. https://doi.org/10.1088/1748-9326/abae9d

Forbes, B. C., Kumpula, T., Meschtyb, N., Laptander, R., Macias-Fauria, M., Zetterberg, P., . . . Bartsch,
A. (2016). Sea ice, rain-on-snow and tundra reindeer nomadism in Arctic Russia. *Biology
Letters, 12*(11). https://doi.org/10.1098/rsbl.2016.0466

Frei, E. R. and Henry. G. H. R. (2021)Long-term effects of snowmelt timing and climate warming on
phenology, growth, and reproductive effort of Arctic tundra plant species. *Arctic Science*. e-
First https://doi.org/10.1139/as-2021-0028

Frisvoll, A. A., & Elvebakk, A. (1996). Part 2. Bryophytes. In A. Elvebakk & P. Presttrud (Eds.), *Svalbard
plants, fungi, algae and cyanobateria*. Norsk Polarinstitutt, Skrifter nr. 198.

Frost, G. V., Bhatt, U. S., Epstein, H. E., Walker, D. A., Raynolds, M. K., & Berner, L. T., et al. (2019).
Tundra greenness. (NOAA Arctic Report Card 2019, Issue.

Gauthier, G., Bêty, J., Cadieux, M.-C., Legagneux, P., Doiron, M., Chevallier, C., . . . Berteaux, D.
(2013). Long-term monitoring at multiple trophic levels suggests heterogeneity in responses
to climate change in the Canadian Arctic tundra. *Philosophical Transactions of the Royal
Society B: Biological Sciences, 368*(1624), 20120482. https://doi.org/10.1098/rstb.2012.0482

Gougherty, A. V. & Gougherty, S. W. Sequence of flower and leaf emergence in deciduous trees is
linked to ecological traits, phylogenetics, and climate. *New Phytologist, 220*, 121–131 (2018).

Givnish, T. (2002). Adaptive significance of evergreen vs. deciduous leaves: solving the triple paradox.
Silva Fennica, 36(3), 535. https://doi.org/doi:10.14214/sf.535

Gornall, J., Woodin, S., Jónsdóttir, I., & van der Wal, R. (2009). Herbivore impacts to the moss layer
determine tundra ecosystem response to grazing and warming. *Oecologia, 161*, pages 747–
758. https://doi.org/10.1007/s00442-009-1427-5

Gornall, J., Woodin, S., Jónsdóttir, I., & van der Wal, R. (2011). Balancing positive and negative plant
interactions: how mosses structure vascular plant communities. *Oecologia, 166*(3), 769-782.
https://doi.org/10.1007/s00442-011-1911-6

Graham, R. M., Cohen, L., Petty, A. A., Boisvert, L. N., Rinke, A., Hudson, S. R., . . . Granskog, M. A.
(2017). Increasing frequency and duration of Arctic winter warming events. *Geophysical
Research Letters, 44*(13), 6974-6983. https://doi.org/https://doi.org/10.1002/2017GL073395

Grant, P. R., Grant, B. R., Huey, R. B., Johnson, M. T. J., Knoll, A. H., & Schmitt, J. (2017). Evolution
caused by extreme events. *Philosophical Transactions of the Royal Society B: Biological
Sciences, 372*(1723), 20160146. https://doi.org/10.1098/rstb.2016.0146

Hansen, B. B., Grotan, V., Aanes, R., Saether, B. E., Stien, A., Fuglei, E., . . . Pedersen, A. O. (2013).
Climate events synchronize the dynamics of a resident vertebrate community in the High
Arctic. *Science, 339*(6117), 313-315. https://doi.org/10.1126/science.1226766

Hansen, B. B., Isaksen, K., Benestad, R. E., Kohler, J., Pedersen, Å. Ø., Loe, L. E., . . . Varpe, Ø. (2014).
Warmer and wetter winters: characteristics and implications of an extreme weather event in
the High Arctic. *Environmental Research Letters, 9*, 114021. https://doi.org/10.1088/1748-
9326/9/11/114021

Hanssen-Bauer, I., Førland, E. J., Hisdal, H., Mayer, S., Sandø, A. B., & Sorteberg, A. (2019). *Climate in
Svalbard 2100 – a knowledge base for climate adaptation (NCCS Report No. 1/2019).*
https://klimaservicesenter.no/kss/rapporter/rapporter-og-publikasjoner_2

Harris, R. M. B., Beaumont, L. J., Vance, T. R., Tozer, C. R., Remenyi, T. A., Perkins-Kirkpatrick, S. E., . . .
Bowman, D. M. J. S. (2018). Biological responses to the press and pulse of climate trends and

extreme events. *Nature Climate Change, 8*(7), 579-587. https://doi.org/10.1038/s41558-018-
0187-9

Henry, G. H. R., & Molau, U. (1997). Tundra plants and climate change: the International Tundra
Experiment (ITEX). *Global Change Biology, 3*(S1), 1-9. https://doi.org/10.1111/j.1365-
2486.1997.gcb132.x

IPCC. (2021). Regional fact sheet–Polar regions. In V. Masson-Delmotte, P. Zhai, A. Pirani, et al. (Eds.),
Climate change 2021: The physical science basis. Contribution of Working Group I to the Sixth
Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University
Press. https://www.ipcc.ch/repor t/ar6/wg1/resou rces/factsheets

Isaksen, K., Lutz, J., Sørensen, A. M., Godøy, Ø., Ferrighi, L., Eastwood, S., & Aaboe, S. (2022a)
Advances in operational permafrost monitoring on Svalbard and in Norway. *Environment*
Research Letters, 17, 095012. https://doi.org/10.1088/1748-9326/ac8e1c

Isaksen, K., Nordli, Ø., Ivanov, B., Køltzow, M. A. Ø., Aaboa, S., . . . Karandasheva, T. (2022b)
Exceptional warming over the Barents area. *Scientific Reports, 12,* 9371.
https://doi.org/10.1038/s41598-022-13568-5Jónsdóttir, I. S. (1991). Effects of Grazing on
Tiller Size and Population Dynamics in a Clonal Sedge (Carex bigelowii). *Oikos, 62*(2), 177-188.
https://doi.org/10.2307/3545263

Jónsdóttir, I. S. (2011). Diversity of plant life histories in the Arctic. *Preslia, 83*(3), 281-300.

Karlsen, S., Elvebakk, A., Høgda, K., & Grydeland, T. (2014). Spatial and temporal variability in the
onset of the growing season on Svalbard, Arctic Norway — Measured by MODIS-NDVI
satellite data. *Remote Sensing, 6*(9), 8088-8106. https://doi.org/10.3390/rs6098088

Kelsey, K. C., Pedersen, S. H., Leffler, A. J., Sexton, J. O., Feng, M., Welker, J. M. (2021) Winter snow
and spring temperature have differential effects on vegetation phenology and productivity
across Arctic plant communities. *Global Change Biology*, *27*, 1572-
1586. https://doi.org/10.1111/gcb.15505

Kausrud, K. L., Mysterud, A., Steen, H., Vik, J. O., Østbye, E., Cazelles, B., . . . Stenseth, N. C. (2008).
Linking climate change to lemming cycles. *Nature, 456*(7218), 93-97.
https://doi.org/10.1038/nature07442

Le Moullec, M., Buchwal, A., van der Wal, R., Sandal, L., & Hansen, B. B. (2019). Annual ring growth of
a widespread high-arctic shrub reflects past fluctuations in community-level plant biomass.
Journal of Ecology, 107(1), 436-451. https://doi.org/10.1111/1365-2745.13036

Le Moullec, M., Sandal, L., Grøtan, V., Buchwal, A., & Hansen, B. B. (2020). Climate synchronises
shrub growth across a high-arctic archipelago: contrasting implications of summer and winter
warming. *Oikos, 129*(7), 1012-1027. https://doi.org/10.1111/oik.07059

Meredith, M., Sommerkorn, M., Cassotta, S., Derksen, D., Ekaykin, A., Hollowed, A., . . . Schuur, E. A.
G. (2019). Polar Regions. In H.-O. Pörtner, D. C. Roberts, V. Masson-Delmotte, P. Zhai, M.
Tignor, E. Poloczanska, K. Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama, &
N. M. Weyer (Eds.), *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*.

Milner, J. M., Varpe, Ø., Van der Wal, R., & Hansen, B. B. (2016). Experimental icing affects growth,
mortality, and flowering in a high arctic dwarf shrub. *Ecology and Evolution, 6*(7), 2139-2148.
https://doi.org/10.1002/ece3.2023

Myers-Smith, I. H., Forbes, B. C., Wilmking, M., Hallinger, M., Lantz, T., Blok, D., . . . Hik, D. S. (2011).
Shrub expansion in tundra ecosystems: dynamics, impacts and research priorities.

Environmental Research Letters, 6(4), 045509. https://doi.org/10.1088/1748-
9326/6/4/045509

Myers-Smith, I. H., Kerby, J. T., Phoenix, G. K., Bjerke, J. W., Epstein, H. E., Assmann, J. J., . . . Wipf, S.
(2020). Complexity revealed in the greening of the Arctic. *Nature Climate Change, 10*(2), 106-
117. https://doi.org/10.1038/s41558-019-0688-1

Niittynen, P., Heikkinen, R. K., Aalto, J., Guisan, A., Kemppinen, J., & Luoto, M. (2020). Fine-scale
tundra vegetation patterns are strongly related to winter thermal conditions. *Nature Climate
Change*. https://doi.org/10.1038/s41558-020-00916-4

Orcutt, D. M., & Nilsen, E. T. (2000). *The physiology of plants under stress: soil and biotic factors*. John
Wiley and Sons.

Outcalt, S. I., Nelson, F. E., & Hinkel, K. M. (1990). The zero-curtain effect: Heat and mass transfer
across an isothermal region in freezing soil. *Water Resources Research, 26*(7), 1509-1516.
https://doi.org/https://doi.org/10.1029/WR026i007p01509

Pan, C. G., Kirchner, B. P., Kimball, S. J., Kim, Y., & Du, J. (2018). Rain-on-snow events in Alaska, their
frequency and distribution from satellite observations. *Environmental Research Letters,
13*(7), 075004. http://stacks.iop.org/1748-9326/13/i=7/a=075004

Peeters, B., Pedersen, A. O., Loe, L. E., Isaksen, K., Veiberg, V., Stien, A., . . . Hansen, B. B. (2019).
Spatiotemporal patterns of rain-on-snow and basal ice in high Arctic Svalbard: detection of a
climate-cryosphere regime shift. *Environmental Research Letters, 14*(1), 015002.
https://doi.org/10.1088/1748-9326/aaefb3

Pérez-Harguindeguy, N., Díaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., . . .
Cornelissen, J. H. C. (2013). New handbook for standardised measurement of plant functional
traits worldwide. *Australian Journal of Botany, 61*(3), 167-234.
https://doi.org/10.1071/BT12225

Petraglia, A., Carbognani, M., & Tomaselli, M. (2013). Effects of nutrient amendments on modular
growth, flowering effort and reproduction of snowbed plants. *Plant Ecology & Diversity, 6*(3-
4), 475-486. https://doi.org/10.1080/17550874.2013.795628

Pettorelli, N., Vik, J. O., Mysterud, A., Gaillard, J.-M., Tucker, C. J., & Stenseth, N. C. (2005). Using the
satellite-derived NDVI to assess ecological responses to environmental change. *Trends in
Ecology & Evolution, 20*(9), 503-510.
https://doi.org/https://doi.org/10.1016/j.tree.2005.05.011

Phoenix, G. K., & Bjerke, J. W. (2016). Arctic browning: extreme events and trends reversing arctic
greening. *Global Change Biology, 22*(9), 2960-2962. https://doi.org/10.1111/gcb.13261

Post, E., & Forchhammer, M. C. (2008). Climate change reduces reproductive success of an Arctic
herbivore through trophic mismatch. *Philosophical Transactions of the Royal Society B:
Biological Sciences, 363*(1501), 2367-2373. https://doi.org/10.1098/rstb.2007.2207

Post, E., Forchhammer, M. C., Bret-Harte, M. S., Callaghan, T. V., Christensen, T. R., Elberling, B., . . .
Aastrup, P. (2009). Ecological dynamics across the Arctic associated with recent climate
change. *Science, 325*(5946), 1355-1358. https://doi.org/10.1126/science.1173113

Preece, C., Callaghan, T. V., & Phoenix, G. K. (2012). Impacts of winter icing events on the growth,
phenology and physiology of sub-arctic dwarf shrubs. *Physiologia Plantarum, 146*(4), 460-
472. https://doi.org/10.1111/j.1399-3054.2012.01640.x

Preece, C., & Phoenix, G. K. (2014). Impact of early and late winter icing events on sub-arctic dwarf
shrubs. *Plant Biology, 16*(1), 125-132. https://doi.org/10.1111/plb.12015

Prevéy, J. S., Rixen, C., Rüger, N., Høye, T. T., Bjorkman, A. D., Myers-Smith, I. H., . . . Wipf, S. (2019).
Warming shortens flowering seasons of tundra plant communities. *Nature Ecology &*
Evolution, 3(1), 45-52. https://doi.org/10.1038/s41559-018-0745-6

Putkonen, J., & Roe, G. (2003). Rain-on-snow events impact soil temperatures and affect ungulate
survival. *Geophysical Research Letters, 30*(4). https://doi.org/10.1029/2002GL016326

R Core team. (2020). *R: a language and environment for statistical computing*. R Foundation for
Statistical Computing.

Rasmus, S., Kivinen, S., & Irannezhad, M. (2018). Basal ice formation in snow cover in Northern
Finland between 1948 and 2016. *Environmental Research Letters, 13*(11), 114009.
https://doi.org/10.1088/1748-9326/aae541

Rixen, C., Høye, T. T., Macek, P., Aerts, R., Alatalo, J. M., Anderson, J. T., Arnold, P. A., … Zong, S.
(2022). Winters are changing: snow effects on Arctic and alpine tundra ecosystems. *Arctic*
Science. e-First. https://doi.org/10.1139/as-2020-0058

Robinson, S. A. (2022) Climate change and extreme events are changing the biology of Polar Regions.
Global Change Biology, 28(20), 5861-5864. https://doi.org/10.1111/gcb.16309

Schindelin, J., Rueden, C. T., Hiner, M. C., & Eliceiri, K. W. (2015). The ImageJ ecosystem: an open
platform for biomedical image analysis. *Molecular Reproduction and Development, 82*(7-8),
518-529. https://doi.org/10.1002/mrd.22489

Schuur, E. A. G., McGuire, A. D., Schädel, C., Grosse, G., Harden, J. W., Hayes, D. J., . . . Vonk, J. E.
(2015). Climate change and the permafrost carbon feedback. *Nature, 520*(7546), 171-179.
https://doi.org/10.1038/nature14338

Semenchuk, P. R., Elberling, B., & Cooper, E. J. (2013). Snow cover and extreme winter warming
events control flower abundance of some, but not all species in high arctic Svalbard. *Ecology*
and Evolution, 3(8), 2586-2599. https://doi.org/10.1002/ece3.648

Semenchuk, P. R., Gillespie, M. A. K., Rumpf, S. B., Baggesen, N., Elberling, B., & Cooper, E. J. (2016).
High Arctic plant phenology is determined by snowmelt patterns but duration of
phenological periods is fixed: an example of periodicity. *Environmental Research Letters,*
11(12), 125006. https://doi.org/10.1088/1748-9326/11/12/125006

Skarpe, C., & Van der Wal, R. (2002). Effects of simulated browsing and length of growing season on
leaf characteristics and flowering in a deciduous arctic shrub, *Salix polaris*. *Arctic, Antarctic,*
and Alpine Research, 34(3), 282-286.

Sokolov, A. A., Sokolova, N. A., Ims, R. A., Brucker, L., & Ehrich, D. (2016). Emergent rainy winter
warm spells may promote boreal predator expansion into the Arctic. *Arctic, 69*(2), 121-129.
https://doi.org/https://doi.org/10.14430/arctic4559

Song, C. & Saavedra, S. (2018) Structural stability as a consistent predictor of phenological events.
Proc eedings of the Royal Society B. 285, 20180767.

Sundberg, B., Uggla, C., & Tuominen, H. (2000). Cambial growth and auxin gradients. In R. A. Savidge,
J. R. Barnett, & R. Napier (Eds.), *Cell and molecular biology of wood formation* (pp. 169‑188).
Oxford, UK: BIOS Scientific Publishers.

Treharne, R., Bjerke, J. W., Tømmervik, H., & Phoenix, G. K. (2020). Development of new metrics to
assess and quantify climatic drivers of extreme event driven Arctic browning. *Remote Sensing*
of Environment, 243, 111749. https://doi.org/https://doi.org/10.1016/j.rse.2020.111749

Treharne, R., Bjerke, J. W., Tømmervik, H., Stendardi, L., & Phoenix, G. K. (2019). Arctic browning:
Impacts of extreme climatic events on heathland ecosystem CO2 fluxes. *Global Change*
Biology, 25(2), 489-503. https://doi.org/10.1111/gcb.14500

Trisos, C. H., Merow, C., & Pigot, A. L. (2020). The projected timing of abrupt ecological disruption
from climate change. *Nature, 580*(7804), 496-501. https://doi.org/10.1038/s41586-020-
2189-9

Van der Wal, R., & Stien, A. (2014). High-arctic plants like it hot: a long-term investigation of
between-year variability in plant biomass. *Ecology, 95*(12), 3414-3427.
https://doi.org/10.1890/14-0533.1

Vickers, H., Høgda, K. A., Solbø, S., Karlsen, S., Tømmervik, H., Aanes, R., & Hansen, B. B. (2016).
Changes in greening in the high arctic: insights from a 30 year AVHRR max NDVI dataset for
Svalbard. *Environmental Research Letters, 11*, 105004. https://doi.org/10.1088/1748-
9326/11/10/105004

Vikhamar-Schuler, D., Isaksen, K., Haugen, J. E., Tømmervik, H., Luks, B., Schuler, T. V., & Bjerke, J. W.
(2016). Changes in Winter Warming Events in the Nordic Arctic Region. *Journal of Climate,*
29(17), 6223-6244. https://doi.org/10.1175/JCLI-D-15-0763.1

Wahren, C. H., Walker, M., & Bret-Harte, M. (2005). Vegetation responses in Alaskan arctic tundra
after 8 years of a summer warming and winter snow manipulation experiment. *Global*
Change Biology, 11(4), 537-552.

Walker, M. D., Wahren, C. H., Hollister, R. D., Henry, G. H. R., Ahlquist, L. E., Alatalo, J. M., . . .
Wookey, P. A. (2006). Plant community responses to experimental warming across the
tundra biome. *Proceedings of the National Academy of Sciences, 103*(5), 1342.
https://doi.org/10.1073/pnas.0503198103

Wipf, S., & Rixen, C. (2010). A review of snow manipulation experiments in Arctic and alpine tundra
ecosystems. *Polar Research, 29*(1), 95-109. https://doi.org/10.3402/polar.v29i1.6054

Wood, S. N., Goude, Y., & Shaw, S. (2015). Generalized additive models for large data sets. *Journal of*
the Royal Statistical Society: Series C (Applied Statistics), 64(1), 139-155.
https://doi.org/10.1111/rssc.12068

You, Q., Cai, Z., Pepin, N., Chen, D., Ahrens, B., Jiang, Z., . . . Zhang, Y. (2021). Warming amplification
over the Arctic Pole and Third Pole: Trends, mechanisms and consequences. *Earth-Science*
Reviews, 217, 103625. https://doi.org/https://doi.org/10.1016/j.earscirev.2021.103625

Zhang, W., Döscher, R., Koenigk, T., Miller, P. A., Jansson, C., Samuelsson, P., . . . Smith, B. (2020). The
Interplay of Recent Vegetation and Sea Ice Dynamics—Results From a Regional Earth System
Model Over the Arctic. *Geophysical Research Letters, 47*(6), e2019GL085982.
https://doi.org/10.1029/2019GL085982

1008