Towards rainy Arctic winters: experimental icing impacts tundra plant productivity and reproduction

Mathilde Le Moullec[1], Anna-Lena Hendel[2], Matteo Petit Bon[3-4-5], Ingibjörg Svala Jónsdóttir[3,6], Øystein Varpe[3,7], René van der Wal[8], Larissa Teresa Beumer[1,9], Kate Layton-Matthews[1,10], Ketil Isaksen[11] and Brage Bremset Hansen[1].

1 Biology Department, Norwegian University of Science and Technology, Dept. of Biology, Centre for Biodiversity Dynamics, 2 Universität Bayreuth, 3 The University Centre in Svalbard (UNIS), 4 Arctic University of Norway, 5 Institute of Botany of the Czech Academy of Sciences, 6 University of Iceland, 7 Department of Biological Sciences, University of Bergen, 8 Agricultural University of Sweden, 9 Arctic Research Centre, Aarhus University, 10 Norwegian Institute for Nature Research, 11 Norwegian Meteorological Institute.

Abstract

The Arctic is the most rapidly warming region on Earth, but our understanding of ecosystem impacts is still poor. For instance, warming occurs more than twice as fast in winter than in summer, yet long-term tundra vegetation studies have almost exclusively focused on effects of the latter. In winter, more frequent extreme warm spells and associated rain-on-snow events can dramatically alter snowpack conditions and even encapsulate the vegetation in 'basal ice' for several months. Such icing effects on plant phenology, productivity and reproduction remain largely unexplored. We performed a novel five-year-long field experiment in which we simulated every winter a rain-on-snow event resulting in ice encasement of the vegetation, and assessed vascular plant responses in each subsequent growing season. We also tested whether these responses could be modified by summer warming (open top chambers). Icing delayed the dominant shrub's phenology, in particular early leaf development and seed maturation, increased community-level primary production in the second half of the growing season, but also reduced flower production. These delays and allocation trade-offs were associated with a delay in sub-surface soil thawing and soil warming in spring/summer, conditions known to influence plant root activity and nutrient availability at high latitudes. Interestingly, summer warming mitigated effects of icing alone and the combined treatment showed advanced phenology, increased primary production

across the growing season and reduced the negative effects on reproduction. In general, effect sizes of icing were as large as those observed for experimental summer warming alone. The community-level increase in primary production following icing is in sharp contrast with the recently proposed 'arctic browning' effect of extreme winter warm spells in evergreen shrub-dominated tundra communities. Yet, the negative effect on reproductive traits can become critical for species viability if these events become chronic. As we are heading towards a rain-dominated Arctic, our findings highlight the need for coordinated monitoring of the effects of warmer and rainier winters across tundra plant communities and growth forms across the Arctic.

1. Introduction

Global warming comes with more frequent extreme climate events, such as droughts, hurricanes and flooding, spurring research activity aimed at determining the short- and long-term ecological and evolutionary consequences for terrestrial biota (Bailey & van de Pol, 2016; Holmgren et al., 2006; Parmesan et al., 2000). In the Arctic, temperatures rise twice as fast as the global average and particularly markedly during winters (You et al., 2021). This has resulted in increasingly frequent and widespread winter warm spells (AMAP, 2017; Bintanja & Andry, 2017; Graham et al., 2017; Vikhamar-Schuler et al., 2016). Despite evidence for severe consequences of extreme climatic events for tundra biota (Post et al., 2009) and ecosystem processes (Treharne et al., 2019), the effects of winter warm spells remain poorly understood and yet most long-term tundra vegetation studies have exclusively focused on effects of rising summer temperatures. This has left a knowledge gap on how changing conditions outside the growing season may contribute to shape the net impacts of global warming on arctic vegetation, especially considering the key role tundra vegetation plays for the global climate-feedbacks (Zhang et al., 2020).

Climate warming has stimulated plant growth in the Arctic resulting in large scale 'greening', which has mainly been attributed to shrub expansion and improved growing conditions for plants (Epstein et al., 2012; Frost et al., 2019; Ju & Masek, 2016; Myers-Smith et al., 2011; Walker et al., 2006). Recently, however, the overall trend of arctic 'greening' seems to have decelerated (Berner et al., 2020; Frost et al., 2019; Myers-Smith et al., 2020; Vickers et al., 2016), with extreme warm spells in winter as one of the proposed causes, potentially even leading to vegetation 'browning' (Bjerke et al., 2017; Bokhorst et al., 2008; Bokhorst et al., 2009; Phoenix & Bjerke, 2016).

During warm spells in winter, increased proportion of precipitation falls as rain instead of snow, often leading to extensive rain-on-snow events influencing the heat budget of the entire snowpack (AMAP, 2017; Bintanja & Andry, 2017; Bokhorst et al., 2016; Pan et al.,

2018; Rasmus et al., 2018). On the one hand, mild weather and rain can lead to a complete

snowmelt, exposing plants to thaw-freeze cycles that may directly cause vegetation damage

(Bokhorst et al., 2011; Bokhorst et al., 2009). On the other hand, as typical for the high

Arctic, rain and the resulting meltwater can also collect on the deeply frozen ground, where it

freezes, leading to the formation of a layer of basal ice (hereafter referred to as 'icing'), up to

several decimetres thick (Peeters et al., 2019; Putkonen & Roe, 2003). Low-growing arctic

vegetation may be entirely encapsulated in this layer of ice for several months, with

ecosystem-wide consequences by affecting soil arthropod abundance (Coulson et al., 2000)

and blocking access to forage for overwintering herbivores, sometimes resulting in major

starvation-induced die-offs (Forbes et al., 2016; Kohler & Aanes, 2004; Miller & Gunn, 2003;

Rennert et al., 2009) which in turn fuel the arctic fox *Vulpes lagopus* populations (Hansen et

al., 2013). Nonetheless, little is known about the effects of icing on the vegetation itself.

Physiologically, icing can lead to plant cell death, either through frost damage or

stress-induced anoxia metabolite accumulation from anaerobic respiration (Bokhorst et al.,

2010; Crawford et al., 1994; Preece & Phoenix, 2014). In addition, icing could indirectly

affect the vegetation through altered soil thermal properties (Putkonen & Roe, 2003) and

delayed phenology, as found after late snow melt (Semenchuk et al., 2016). A handful of

studies investigated effects of icing in the low Arctic (Bjerke, 2011; Bjerke et al., 2018a;

Preece et al., 2012; Preece & Phoenix, 2014), or on the short-term in the high Arctic (Bjerke

et al., 2018b; Bjerke et al., 2017; Milner et al., 2016). Evergreen shrub and lichens were

strongly negatively affected by icing, with evidence of altered leaf physiology, shoot and

individual mortality (Bjerke, 2011; Bjerke et al., 2017; Milner et al., 2016; Preece et al.,

2012). In comparison, deciduous shrubs and graminoids showed high-tolerance, yet, with

evidence of damage and biomass reduction (Bjerke et al., 2018b; Preece & Phoenix, 2014).

When organisms face sub-optimal environmental conditions, they may channel more

energy either to growth or reproduction causing pronounced energy allocation trade-offs

(Bazzaz et al., 1987; Jónsdóttir, 2011). Induced plant stress can enhance vegetative growth,

thereby constraining or delaying (i.e. lagged effects) other reproductive trait responses

(Chapin, 1991; Orcutt & Nilsen, 2000), such as flower phenology and number. Such

allocation trade-offs are found for arctic plants coping with shortened growing seasons (e.g.

late snow melt, Cooper et al., 2011; Wipf & Rixen, 2010), damages from e.g. herbivory

(Jónsdóttir, 1991; Skarpe & Van der Wal, 2002), thaw-freeze (Bokhorst et al., 2011) or,

possibly, icing. For example, in a one-year high arctic field experiment, icing of the evergreen

shrub *Cassiope tetragona* led to high mortality of apical meristems and entire shoots (Milner

et al., 2016). As a result, this shrub invested more in the growth of surviving shoots the

following summer, at the cost of reproductive structures, i.e., reduced flower production.

Plot-level field experiments are powerful tools to investigate fine-scale changes in plant

response to simulated climate change scenarios (Bjorkman et al., 2020; Bjorkman et al., 2018;

Elmendorf et al., 2012; Elmendorf et al., 2015). Here, we assess how experimental winter

rain-on-snow events and subsequent icing affect productivity, reproduction, and phenology of

an high arctic tundra plant community. We also tested whether these responses could be

modified by summer warming. To achieve this, over five consecutive years, we simulated

plot-level rain-on-snow and icing formation in winter. In full factorial design, we also

increased summer temperature via open top chambers, following the protocol of the Arctic

and Alpine International Tundra Experiment (ITEX, Henry & Molau, 1997). During each

growing seasons we determined plant species and community responses to our icing and

summer warming treatments. We expected icing to delay plant phenology due to later thawing

out of the soil, reduce plant productivity due to shoot mortality ('browning') and,

consequently, reduced flowering due to reallocation of plant resources to regrowth. We

expected summer warming to mitigate the icing effects.

2. Methods

Study site

The experiment was located in high Arctic Svalbard (78°17`N, 16°02`E), in the valley of

Adventdalen, and lasted for five years (Jan. 2016 – Aug. 2020). The annual mean air

temperature recorded at Svalbard Airport (Longyearbyen), 15 km from the study site, was -

5.9 °C for the reference period 1971-2000 (Hanssen-Bauer et al., 2019), and -2.2 °C for the

study period. During the reference period, winter (December-February) temperature rose four

to five times faster than in summer (June-August) (Hanssen-Bauer et al., 2019). The interior

fjord and valley areas around Longyearbyen are among the driest in Svalbard with average

annual precipitation of 196 mm for the reference period (Hanssen-Bauer et al., 2019), and 223

mm for the study period. In comparison to the reference period, median RCP8.5 projections

for Svalbard anticipate an increase in temperature and precipitation of about 9-13 °C and 20-

40%, respectively, towards 2100 (Hanssen-Bauer et al., 2019). The frequency of rain-on-snow

events has increased dramatically, with a regime shift around year 2000 (Hansen et al 2014,

Peters et al. 2019). In that sense, contemporary Svalbard reflects conditions soon expected

across the Arctic (Bintanja & Andry, 2017).

Soil thermal conditions on Svalbard are linked to the underlying permafrost which

warms at a rate of 1.4 °C per decade at the permafrost table (2.0 m depth), observed 6 km

away from our study site, over 2002-2018 (Etzelmüller et al., 2020). At the study site, the soil

sub-surface (2 cm depth) temperature was on average 4.7°C in June and 7.5°C in July (see

methodology below), the two most important months for plant growth (Van der Wal & Stien,

2014). The volumetric water content (i.e., soil moisture) is relatively low for mesic

communities, drying throughout the summer season from about 45% to 25% at 10 cm depth

(Table S1a). Across the growing season, soil moisture and sub-surface temperature are

strongly negatively related (Table S1b).

The plant growing season in the tundra starts immediately after the soil thaw onset

(Descals et al., 2020). Adventdalen, which is in the bioclimatic subzone C ('middle arctic

tundra' (CAVM Team, 2003), has a the growing season of approximately three months long,

starting in mid-late May (Table 1). The plant community studied here was a mesic grass and

moss-rich dwarf shrub heath, dominated by the prostrate deciduous dwarf shrub *Salix polaris*

and the grass *Alopecurus borealis*. Other abundant vascular plants were the horsetail

Equisetum arvense, the rush *Luzula confusa*, the forb *Bistorta vivipara* and the grass *Poa*

arctica (Figure S1). *The most abundant bryophytes were Sanionia uncinata, Tomentypnum*

nitens and *Polytrichastrum* spp. Nomenclature follows http://panarcticflora.org/ for vascular

plants and Frisvoll and Elvebakk (1996) for bryophytes.

Experimental design

The study design followed a full-factorial randomized block design with four

treatment combinations (2 levels of icing x 2 levels of warming) replicated three times within

each of three blocks (Figure 1). In summer 2015, the three blocks (150-780m apart) were

selected in homogenous mesic tundra. Within each block we chose 12 similar plots of 60×60

cm (Figure 1). The four treatments, icing (*I*), summer warming (*W*), a combined winter icing

and summer warming treatment (*IW*), and control (*C*), were randomly assigned to plots. Plots

received the same treatment each year, starting in winter 2015-2016 (Table S2). This allowed

us to investigate plant responses from the short-term (one year) to the longer-term (up to five

years). However, not all traits and parameters were measured in all years, hence some results

are presented based only on subsets of years.

Winter icing: The experimentally formed ice was applied to *I* and *IW* plots within a

period of 2-3 days in January-February (Figure 1a). Snow was carefully removed from all

plots to apply the same level of disturbance, and any occurrence of natural ice was recorded.

Natural ice was particularly common in 2017, with partial ice coverage occurring in all plots

(3 cm average thickness). Snow was immediately placed back on non-icing (*C* and *W*) plots,

while for *I* and *IW*, a 13 cm high wooden frame (60×60 cm) was placed around the plot and

gradually filled up with cold water (for 1-3 days) until it was full of solid ice (Figure 1a),

following Milner et al. (2016). Frames had no influence on the snow accumulation pattern

later in winter and were removed at the beginning of snowmelt.

Summer warming: Right after snowmelt (Table S2), hexagonal open top chambers (1.4

$\times$ 1.4 m basal diagonal) were deployed over the *W* and *IW* plots following the specifications

given by the ITEX protocol (Henry & Molau, 1997). To avoid confounding factors, all plots

were excluded from grazing herbivores during the snow-free season using metal net cages or

nets on top of the open top chambers (mesh-size 1.9 cm $\times$ 1.9 cm)(Figure 1b).

Field measurements

Soil temperatures were recorded with iButton loggers (type DS1921G-F5, $\pm$ 1.0 °C accuracy,

0.5 °C resolution) every 120 minutes in summer and 240 minutes in winter. Loggers were

placed in all plots at depths of both 2 cm and 5 cm. In 2018, we also recorded temperatures at

10 cm and 20 cm depth, in half of the plots (not in *IW*). We estimated the onset of soil-thaw in

spring for each plot based on logger data at 5 cm depth , thereby taking the start of period for

which soil sub-surface temperature had been ≥ 0 °C for a minimum of 10 consecutive days.

At high latitudes, plant growth commences almost immediately upon the onset of soil thaw

(Descals et al., 2020).

To assess treatment effects on plant production we monitored the vegetation weekly

during the growing season by measuring NDVI (a proxy for plant productivity with a value

between 0-1). This was done at the central part of each plot (30 cm diameter) with a Skye

SpectroSense2+ handheld device. NDVI typically increase until in the first part of the

growing season, reaching a maximum, which we considered as the day of peak growing

season (Table 1), to then drop progressively in the second part of the growing season. As an

additional measure of plant productivity, we measured community-level vascular plant

species abundance in the period after the peak growing season (end of July, Table 1), in 2016-

2019 (Table S2). We used a point intercept methodology, hereafter referred to as 'relative

abundance' (Bråthen & Hagberg, 2004), with a 50×50 cm frame elevated above the canopy

(~20 cm high) and with 25 evenly distributed points. The points were marked by crossings of

double strings with the frame to give a 90º projection. At each point, a wooden pin of 3 mm

diameter was lowered down onto the moss layer and all 'hits' of vascular plants were

recorded.

To further study vegetative responses of the dominant vascular species, we collected

S. polaris leaves (before leaf senescence, Table S2) to measure area, leaf dry weight and

specific leaf area (SLA, the ratio of leaf area to dry mass, a frequently used proxy for relative

growth rate (Pérez-Harguindeguy et al., 2013)). The protocol and sample size of leaf

collection varied between year (Table S2), preventing direct comparison of leaf-size traits

between years, but still enabling between-year comparison of *I*, *IW* and *W* effect sizes relative

to C. Leaves were kept moist after collection, and scanned. Leaf area was measured with the

program imageJ 1.48 software (Schindelin et al., 2015). Then, leaves were oven-dried at 60°C

for four days and weighed on an XS204 METTLER TOLEDO scale to the nearest 0.01 mg.

To assess treatment effects on flowering frequency, the total number of flowers for the

species *S. polaris, B. vivipara, A. borealis, L. confusa and P. arctica* were counted in mid-

July each year (Table S2). We counted flowers in a 50×50 cm frame (Figure 1c), subdivided

in 16 sub-squares.

To assess treatment effects on phenology of the dominant vascular plant, we measured

the most advanced vegetative and reproductive phenological stages (phenophases) of *S.*

polaris in each of 16 sub-squares per plot of the 50×50 cm frame. We used the following

vegetative phenophases: 1) leaves starting to unfurl, 2) leaves fully expanded, 3) start of

senescence and 4) leaves fully senesced. The reproductive phenophases were. 1) distinct

flower buds visible, 2) buds recognisable as female or male, 3) receptive stigmas (female) and

open anther releasing pollen (male), 4) stigma and anthers withered, and 5) seed dispersal.

The greatest phenological monitoring effort took place in 2018, and no phenological

monitoring was done in 2020 (Table S2).

Statistical analysis

All data were analysed in R-3.6.3 (R Core team, 2020).

We derived different metrics from the NDVI data by fitting a generalized additive

model (GAM) to each plot for each year, using the repeated measurements over a growing

season. We employed the function 'gam' from the mgvc package (Wood et al., 2015), fitted

with restricted maximum likelihood. Model fit was good (visual interpretation and k-indices >

1), except for summer 2017 when measurements stopped before reaching the peak growing

season, i.e., they were still in the increasing phase (Table S2). Thus, we removed 2017 from

all NDVI analyses. For all other years, we then predicted daily NDVI values for each plot and

derived: 1) the maximum NDVI value and 2) the day-of-year when this was reached; 3) the

cumulative NDVI in the first part of the growing season (hereafter 'cumulative start'), by

summing daily values from the first day of measurements to the day when the average control

plot reached the maximum NDVI; 4) the cumulative NDVI in the second part of the growing

season (hereafter 'cumulative end'), by summing daily values from the day when the average

control plot reached the maximum NDVI to the last measurement day of the season; 5) the

cumulative NDVI across the growing season (hereafter 'cumulative total'), by summing daily

values from first to last day of measurements.

We used (generalized) linear mixed-effect models as the main tool to analyse our data,

to account for the hierarchical spatial and temporal structures of our study design, using the

functions 'lmer' and 'glmer' from the lme4 R-package (Bates et al., 2015). We built a

separate model for each response variable of interest, i.e., soil temperature, onset of soil thaw,

NDVI (different metrics), plant relative abundance, leaf size traits, flower number, and

phenophases. The fixed effects included varied depending on whether the analysis was to

estimate treatment effect sizes (treatment was set as fixed effect), annual means ('treatment ×

year' interaction was set as fixed effect), or the average of a response over a certain time

period ('treatment × day-of-year' interaction were set as fixed effects). The random intercept

structure (i.e. variation in means between replicated units) always included the nested

structure of plot within block. It also included sub-square nested within plot or leaf nested

within shoot where this level of replication was relevant. Temporal replication was accounted

for by including year as a random intercept (except when already set as fixed effect), and day-

of-year nested within year a random intercept when measurements were also replicated within

year.

Data from soil temperature and NDVI metrics followed a normal distribution, data

from plant relative abundance and flower counts followed a Poisson distribution. Leaf size

traits were log-transformed. Each phenophase was converted to a binary data format to fit a

logistic regression from a binomial distribution.

3. Results

Soil temperature and thawing onset

The added liquid water during the simulated rain-on-snow (i.e. icing treatment) released latent

heat, temporarily increasing sub-surface (i.e. 2 cm depth) soil temperature for three days by

on average 2.9 °C [2.3,3.5] (mean [95% CI], 2016-2020) in plots being ice-encased (i.e., icing

[*I*] and icing × warming [*IW*] plots), compared to control plots (*C*). During snowmelt in

spring, sub-surface soil thaw onset was delayed by 6.0 [2.2,9.9] days in I and 4.4 [0.4,8.5]

days in IW, while it remained unaffected in warming (*W*) compared to C (-0.9 [-5.0,3.1] days;

open top chambers were mounted after snowmelt, Figure 2a). However, the sub-surface soil-

thaw onset, and the period over which it occurred (i.e., its day-of-year range), varied

considerably between years (Table 1). At 20 cm depth (data from 2018), soil thaw onset was

delayed by 9.0 [1.3,16.7] days in *I*, after staying at a constant temperature of -0.5 °C for about

two weeks (i.e., 'the zero curtain period', (Outcalt et al., 1990), Figure 2b). This resulted in a

lag in soil temperature increase, i.e. a month-long slower increase in spring soil temperature in

I versus *C* (i.e. entire June, Figure 2b). Open top chambers presence increased soil sub-surface

temperatures (mid-May to mid-September) by 0.8 ± 0.2 (mean ± SE) and 0.6 ± 0.2 °C in *W*

and *IW*, respectively.

Vegetative responses: NDVI metrics, relative abundance, leaf size traits

The NDVI curves increased rapidly until reaching a maximum in mid-July (Table 1)

and decreased slowly thereafter (Figure 3a, Figure S2). The shape of the NDVI curves varied

among treatments, particularly after the maximum NDVI. This effect was captured by the

NDVI cumulative metrics. For the first part of the growing season, only *IW* increased the

NDVI 'cumulative start', and remained high for the second part of the growing season (NDVI

'cumulative end'), resulting in an overall increased NDVI ('cumulative total') across the

growing season, compared to C (Figure 3b). In the second part of the growing season, *I* catch-

up with *IW*, and thus exceeded NDVI 'cumulative end' compared to *C*. NDVI cumulative

metrics of *W* did not differ from *C* (Figure 3b).

The maximum NDVI timing was delayed on average by 4 [2;7] days in *I*, advanced by

on average -3 [-5;0] days in *IW*, and unchanged in *W* (2 [2-7] days) compared to *C* (Figure

4a). Maximum NDVI values themselves were higher in *I* and *IW* versus *C* in the first year of

treatment (2016), while this effect seemed to vanish in later years (Figure 4b). In 2020 for

instance, even *C* had high maximum NDVI in comparison to other years (Table 1), reaching

similar values to what *I* and *IW* reached other years (Figure S2). Maximum NDVI values were

unaffected by *W* (Figure 4b).

Vascular plants' relative abundance was higher under *IW* compared to *C*, largely

consistent with the NDVI cumulative patterns for the entire growing season ('cumulative

total', see above) (Figure 3c). This pattern was driven by the two most abundant vascular

plants, the shrub *S. polaris* and the graminoid *A. borealis* (Figure S3), which apparently had a

large influence on the NDVI pattern based on among-plot correlations (Table S3). Among-

species, the relative abundance did not change markedly across the study period, i.e. the

community composition remained unchanged (Figure S4).

The leaf specific area (i.e., SLA a proxy for leaf growth rate,) of *S. polaris* was higher

in *I* compared to *C*, and this difference became greater over the years (Figure 4c). Overall,

SLA was not different in *IW* and *W* versus *C*, although variation in effect size among years

was considerable (Figure 4c). Both the leaf area and dry weight tended to decrease in all

treatments compare to C, and this consistently in *I* from 2018 and onwards (Figure S5).

Reproductive response: flower production

The total number of vascular plant flowers in this mesic community was strongly reduced in I

plots, but to some extent also in IW and W, compared to C. In 2017-2019, icing (I) reduced

the flower production to about half of that in C, but it had no effect in the first and last years

of treatment (i.e. 2016 and 2020, Figure 4d and Figure S6). At the species level, *S. polaris*

represented the greatest proportion of flowers and drove this overall pattern of annual

variation in treatment effect, while the forb *B. vivipara* and graminoid inflorescences showed

reduced flower production in all years under I only (Figure S6). Note that in 2020, there was a

record number of flowers in the community, with more than twice the amount of any other

year of the study (Table 1, Figure S6).

 Phenology responses

Leaves' early phenophases (i.e., unfurling) of *S. polaris* were significantly delayed by

I, while the timing of both leaves unfurling and fully expanding were significantly advanced

(by a week) in *IW* and *W* compared to *C* (Table S4, Figure 5a-b). However, the onset of leaf

senescence happened simultaneously across treatments (Table S4, Figure 5c-d), and this

consistently each years (Figure S7a).

Several *S. polaris* flowers's phenophases were delayed by *I* and advanced by *IW* and *W*,

compared to *C*. This difference in timing was largest for the phenophase of seed dispersal

(Table S4, Figure 54e-l). Seed dispersal was delayed by 5 (range at the plot-level: -2:12) days

with *I*, advanced by -6 (-13:1) days with *W* and advanced by as much as -13 (-19:-6) days

with *IW* (estimates for 2018, Figure 5i). Similar patterns were observed among years (Figure

S7b). Furthermore, development of male flower anthers and pollen release phenophases were

advanced by -4 (-7:3) days in *IW* plots (but not in *I*) compared to *C* (Figure 5j-k, Table S4),

while anther senescence was rather synchronous across treatments (Figure 5l).

4. Discussion:

This five-year field experiment demonstrates how extreme winter warm spells, resulting in tundra icing, affect seasonal plant growth and reproduction in a high arctic tundra community. Icing caused delayed, but eventually enhanced, above-ground community-level productivity, opposite to what we expected (Figure 3). This was confirmed at the species level, where the vegetative phenology (Figure 5) of the dominant dwarf shrub *S. polaris* was delayed in early summer, but more than compensated for this through increased leaf relative growth rate (i.e., SLA, Figure 4) and relative abundance (Figure S3). These observed species- and community-level delays in phenology and productivity due to simulated rain-on-snow and icing appeared to result from later soil spring thawing and delayed soil warming in the upper active layer, compared to control plots (Figure 2). Importantly, although the delaying effects of icing seem to be compensated during summer by increased growth, icing had negative effects on reproductive traits (Figure 3 and 4i), suggesting changing resource allocation trade-offs with life history consequences. Interestingly, the addition of summer warming mitigated the effects of icing alone, while still increasing the primary production. These results highlight the potentially large implications of winter climate change for arctic tundra vegetation, so far largely overlooked in both observational and experimental studies.

4.2. Icing causes delayed but increased primary production

The icing-induced delay in the thawing of upper soil layers expectedly caused the delay in early-developmental stages of the dominant vascular plant, the dwarf shrub *S. polaris*. However, the tendency for the vegetation after icing to catch-up the delay by having greater community-wide primary production later in the growing season contrasts with the documented 'arctic browning' following winter warm spells (Frost et al., 2019; Phoenix & Bjerke, 2016). These different observations might be explained by differences in the overwintering strategies of plants. Such 'browning' has mainly been associated with damage

on evergreen shrubs, which maintain long-lived leaves and dormant buds well above-ground, making them more dependent on insulating snow cover and more exposed to direct damage from icing than deciduous shrubs (Givnish, 2002). The dead above-ground shoots of evergreen shrubs can mask compensatory growth (Milner et al. 2016), resulting in NDVI decrease (Bjerke et al., 2017; Treharne et al., 2020). In this experiment, without evidence of dead shoots on the dominant deciduous shrub, the compensatory growth appear to enhance NDVI.

Indirect effects of the altered soil thermal dynamics represent a likely underlying mechanism for the compensatory vegetative growth observed, both in *S. polaris* and the vascular plant community as a whole. Soil temperature influence the nutrient availability like nitrogen and phosphorus, which are limiting resources for plant growth in arctic soils (Chapin, 1991; Epstein et al., 2000). In a similar mesic community, soil temperature had a large influence on microbial biomass and activity, as well as plant-available nutrients, thus influencing the energy transfer to the plant (Gornall et al., 2011). In the 'race' for high primary production within the short time window available in icing plots (due to the delayed soil processes), *S. polaris* appears to compensate delayed leaf development by increasing leaf relative growth rates (i.e. high SLA and a faster transition between phenophases from unfolding to fully expanded leaves), reducing dry weight and area but with overall increasing abundance. These results agree with a laboratory icing experiment on *S. polaris* also finding reduced leaf size and increased leaf number, while maintaining high photosynthetic capacity (Bjerke et al., 2018b). This shrub has a remarkable capacity to activate dormant buds to compensate damage from e.g., herbivory (Skarpe & Van der Wal, 2002). The increased primary production after the peak growing season occurred despite a synchronous start of *S. polaris* leaf senescence across treatments (Figure S7a), which can be driven by cues fixed in time (e.g., photoperiod). However the rate of senescence can interacting with other

environmental factors such as temperature (Fracheboud et al., 2009; Paus et al., 1986).

Regarding icing effects on below-ground (secondary) growth of *S. polaris,* annual ring growth

was reduced in summers following heavy rain-on-snow in coastal areas frequently

characterised by icing (Le Moullec et al., 2020). Therefore, larger primary growth may come

at the cost of reduced secondary growth, which could alter the otherwise strong correlation

between primary and secondary growth in this species (Le Moullec et al., 2019).

Although winter temperatures are rising much faster, summer temperatures are still

increasing at an average rate of 0.4°C per decade across Svalbard (Hanssen-Bauer et al.,

2019). The corresponding combination of icing and warming treatments, unexpectedly

magnified primary production already early in the growing season so that icing and warming

appeared additive. This can be explained by the advanced leaf phenology of *S. polaris*, but

also by the increased relative abundance of the dominant graminoid *A. boralis.* Grasses were

found to be the growth form most rapidly responding to experimentally increased soil

temperature by increasing their total live plant biomass (Brooker & van der Wal, 2003). The

extremely high summer temperature in 2020 may reflect a 'contemporary threshold' in

primary production of this community, beyond which it is possibly limited by nutrient

availability (Chapin, 1991; Epstein et al., 2000). In that year, the estimated maximum NDVI

was very high across treatments, i.e., at the same level as in icing plots in other years (Figure

S2). In addition to the heat waves that occurred in July, the soil thaw onset occurred

synchronous across blocks and plots (for instance, all control plots thawed over four days

instead of e.g., over one month the year before, Table 1).

In contrast to evergreen shrubs, deciduous shrubs represent the main nutritional plane

of arctic herbivores in arctic regions (Chapin III et al., 1996). Thus, the delaying effect and

unexpected positive effect of icing on primary production have potential implications for the

whole ecosystem, at least regarding plant-herbivore interactions (Gauthier et al., 2013; Post &

Forchhammer, 2008; Post et al., 2009), carbon fluxes (Schuur et al., 2015) and vegetation-

climate feedbacks (Zhang et al., 2020). For instance, the late summer/autumn season is when

arctic and temperate large herbivores accumulate body fat, essential for surviving the

following winter (Hurley et al., 2014; Loe et al., 2021). Tundra icing may thus have

counteracting effects for the herbivores, limiting food availability over the winter leading to

starvation, but also increasing the food quantity available in late summer.

4.3. Negative effects on reproduction: a result of trade-offs?

Compensatory vegetative growth may come at the cost of reproduction and has been observed

after thaw-freeze cycles (Bokhorst et al., 2011), snow manipulation (Cooper et al., 2011),

nutrient addition (Petraglia et al., 2013), and icing of evergreen shrubs (Milner et al 2016).

The reduced flower numbers in our experiment (by about half in some years) could result

from prioritized energy allocation to vegetative growth, if not from direct winter damage. In

the first experimental year, no flower reduction occurred for *S. polaris* suggesting that flower

buds formed the previous summer survived a winter of icing. Other species e.g. *B. vivipara*

and *A. boralis* exhibited a strong reduction in flower number without lag (Figure S6). In an

experiment conducted in the same valley, a record low number of flowers was associated with

extreme rain-on-snow events in winter 2012 (Semenchuk et al., 2013). The delay in flower

phenology and in particular seed dispersal of *S. polaris* were likely exacerbated by a

compressed growing season due to icing. For instance, delayed spring onset by late snow melt

also showed delayed flower development and seed dispersal, as well as reduced flowering

(Cooper et al., 2011; Karlsen et al., 2014; Semenchuk et al., 2013; Semenchuk et al., 2016).

Interestingly, summer warming mitigated the energy allocation trade-off found in

icing alone. Under such conditions, plants invested simultaneously in vegetative growth and

reproduction, demonstrated by the advanced flower phenology compare to controls. In

particular, the timing of flowering in male *S. polaris* advanced remarkably, yet was still

followed by a synchronous senescence of the anthers, suggesting an extended period of pollen

release of about 40% compared to untreated plots (Figure 5k-i). Our data for the extremely

warm summer of 2020 also support that warming enhanced both the investment in high

vegetative and reproductive production of control plots, as visible by a record high number of

flowers (Figure S6). However, the reduction of flowers found in warming and icing $\times$

warming plots may still reflect nutrient limitation of the system under which vegetative

growth is prioritized (Petraglia et al., 2013).

Evolutionary consequences are expected from trade-offs influencing sexual

reproductive structure. However, the plant species composing arctic mesic communities are

generally perennial, allowing them to wait for years for favourable conditions, to reproduce

sexually (Bazzaz et al., 1987). In unfavourable years, loss of a cohort can be buffered by seed

banks, yet, seeds from *S. polaris* may be short-lived as they showed low germination rates

from seed banks (Cooper et al., 2004). Nevertheless, *S. polaris*, as most arctic species, also

reproduces asexually by clonal growth (Jónsdóttir, 2011), limiting possible evolutionary

consequences of environmental changes. This also indicates that such communities have

developed adaptive strategies to cope with these extreme conditions. In the oceanic climate of

Svalbard, rain-on-snow events have indeed been occurring at least for decades (Vikhamar-

Schuler et al., 2016), but their frequency and extent have increased dramatically in recent

years (Hansen et al., 2014; Peeters et al., 2019). However, the selection acting upon life-

history traits might change when acute stress episodes from occasional icy winters become

'chronic', i.e. repeated, as expected in the near future (Bintanja & Andry, 2017). Comparative

studies from arctic regions where rain-on-snow events are still rare would shed light on

plants' adaptive capacities to icing.

4.5. The relative impacts of winter versus summer warming.

The effect sizes of winter icing and summer warming treatments were similar across most traits and in the range of natural inter-annual variation, which is typically large in the high arctic (Table 1). Although the use of open top chamber only marginally increased mean summer soil temperatures, the increase was comparable to other ITEX sites in the high Arctic, where irradiance is low, restricting the warming effect (Bokhorst et al., 2013). The primary production increase under warming was therefore limited. Other experiments altering winter conditions, such as snow accumulation, also found similar or larger effect sizes of the respective winter treatment than summer warming by open top chamber (Cooper et al., 2011; Wahren et al., 2005). Our study thus supports the overall insights that the tundra vegetation growth implications of variation in conditions during the 'dormant season' can be substantial.

Climate scenarios suggest a further acceleration of winter and summer warming (Meredith et al., 2019) but their impacts will likely be heterogeneously distributed depending on landscape structure (e.g., aspect). The expected complexity of conditions shaping vegetation growth patterns is supported by circumpolar studies documenting a mosaic of tundra greening and browning (Buchwal et al., 2020; Myers-Smith et al., 2020). Here, we have gained new insights by revealing some of the likely scenarios and underlying mechanisms of winter warming effects. We expect further complexity related to the severity of these events, such as the duration of ice encasement, repeated icing events (Preece & Phoenix, 2014) with possible inter-periods of thaw-freeze events (Bokhorst et al., 2011), and varying intensity of summer warming.

4.6. Conclusion and outlooks.

High arctic environments are naturally variable, and plants have developed elaborate strategies to persist under extreme conditions. Still, this study has shown that winter rain-on-snow, and associated tundra icing, can have similarly large impacts as summer warming on

high arctic plant communities. On the occasion of the 30-year anniversary of ITEX experiments conducted across the poles, which have provided critical knowledge regarding tundra responses to summer warming (Bjorkman et al., 2018; Prevéy et al., 2019), we call for the implementation of similar large-scale experiments investigating effects of winter warming, notably rain-on-snow and icing events. Such a coordinated effort on long-term effects across communities, plant functional types and species, is required to gain a holistic understanding of the tundra biome's future changes and its possible resistance and resilience to both winter and summer warming (Bjorkman et al., 2020). This is especially urgent given that winter warming is outpacing summer warming by far, which may have ecosystem-wide consequences, including alteration of below-ground vegetation productivity, microorganism activity and carbon flux exchanges, and cascading effects across the food web. Our pioneer long-term high arctic icing experiment sets the stage for forthcoming studies to fill this fundamental knowledge gap.

5. Acknowledgements

This project was funded by the Research Council Norway (FRIPRO 276080 and SFF-III 223257) and Svalbard Environmental Protection Fund (project 16/113). We thank UNIS logistics for logistical support and Longyearbyen Lokalstyre for permission to perform the experiment (RiS ID: 10484). We are also deeply grateful for all the help in the field from Solvei B. Hovdal, Martha Grotheim, Hanne K. Haraldsen, Julia Greulich, Karrín Björnsdóttir, Kristine Valøen, Robin Zweigel, Ådne Nafstad, Marit Arneberg, Kristiane Midhaug, Marte Søreng, Marianne Angård, Hamish Burnett, Lia Lechler, Charlotte Karlsen, Kjerstin Hilmarsen, Lukas Tietgen, Svea Zimmermann.

Table 1. Overview of annual mean biotic and abiotic measurements. The daily average air

temperatures were recorded at the Longyearbyen-Svalbard Airport (available on

https://seklima.met.no/), while the soil thaw onset day, maximum Normalized Difference

Vegetation Index (max. NDVI) metrics and flower counts were recorded in control plots. The

time of max. NDVI is interpreted as the peak growing season. DOY = Day of year (1 = 1

January).

	2016	2017	2018	2019	2020
Soil thaw onset day (range)	13th May (8th–22rd May)	21th May (3rd May–1st June)	10th May (6th–14th May)	30th Apr (18th Apr–20th May)	18th May (16th–22th May)
Air T°C June (range)	5.0 (3.6–6.7)	4.6 (3.2–6.5)	4.0 (2.3–6)	4.8 (3.3–6.8)	4.5 (2.8–6.8)
July (range)	9.0 (7.4–11.1)	6.9 (5.4–8.9)	7.2 (5.6–9.5)	8.4 (6.5–10.6)	9.8 (7.7–12.2)
Time of max. NDVI, DOY [CI]	16th July 197 [189;205]	NA	19th July 199 [191;207]	12th July 192 [184;200]	15th July 196 [188;204]
Max. NDVI [CI]	0.74 [0.72;0.77]	NA	0.76 [0.73;0.78]	0.76 [0.73;0.78]	0.79 [0.76;0.81]
Flower count m^2 [CI]	263 [181;345]	229 [151;288]	314 [218;409]	295 [204;386]	491 [341;642]

Figure 1. The experimental set up in Adventdalen, Svalbard. Overview of the field site (a) in the polar night, when applying the icing treatment by filling a wooden frame with water, and (b) in summer with the Open Top Chambers (plexiglass hexagons) simulating warming and net cages to protect against herbivory. (c) Experimental full-factorial randomized block design with the vegetation frame fitting the plot size. C = control, I = icing, IW = icing × warming, W = warming.

Figure 2. Daily estimate of soil temperature during the soil thawing period of mid-May to end

of June 2018, (a) at the sub-surface (5 cm depth), and (b) at 20 cm depth. Shaded areas

represent the 95% CI and the orange vertical dashed lines shows the day the warming

treatments started.

Figure 3. Experimental effects on primary productivity proxies of the community (2016, 2018-2019). (a) Estimated Normalized Difference Vegetation Index (NDVI) curves and 95% CI across the growing season. (b) Effect size of treatments in contrast to control plots (i.e., the reference level marked by the zero line). NDVI cum. start = NDVI cumulative start. NDVI cum. end = NDVI cumulative end. NDVI cum. total = NDVI cumulative total. (c) Relative abundance of the vascular plants assessed by point intercept methods close to the peak growing season. See Figure S2 and S3 respectively for the NDVI curves in 2020 and for relative abundance estimates of 2017.

Figure 4. Effect size of treatments in comparison to controls (i.e., the reference level marked by the zero line) for annual (a) the time of maximum Normalized Difference Vegetation Index (Time of max. NDVI), in number of days; (b) the max. NDVI (values from 0 to 1); (c) the leaf specific area (SLA) of *Salix polaris* (log-scale); (d) flower count per sub-square (logit-scale). The bars shows the 95% CI.

Figure 5. Probability curves of *Salix polaris* phenophases (i.e. the estimated probability of

reaching a specific phase) for vegetative phases (a-d) and reproductive phases (female

flowers; e-i, male flowers; j-l). Phenophases correspond to (a) leaves unfurled, (b) leaves fully

expanded, (c) leaves started senescing, (d) leaves fully senesced, (e) flower visible (female or

male), (f) stigma visible, (g) stigma receptive, (h) stigma senesced, (i) seed dispersed, (j)

anther visible, (k) pollen released, (l) anther senesced. Estimates are shown for 2018, i.e.

when all phases could be estimated after an intensive monitoring effort (for additional

estimates, see Figure S5). The predicted estimates were computed with generalized linear

mixed-effect models, accounting for the replicated structure of the study design (i.e., random

effects), and the shaded areas represent the associated 95% CI to the fixed-effects. Red tick

marks on the x-axis represent sampling days. C = control, I = icing, IW = icing × warming, W

= warming.

6. References

AMAP. (2017). Snow, Water, Ice and Permafrost in the Arctic (SWIPA). In *Arctic Monitoring and Assessment Programme (AMAP)*.

Bailey, L. D., & van de Pol, M. (2016). Tackling extremes: challenges for ecological and evolutionary research on extreme climatic events [https://doi.org/10.1111/1365-2656.12451]. *Journal of Animal Ecology, 85*(1), 85-96. https://doi.org/https://doi.org/10.1111/1365-2656.12451

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software, 67*(1), 1-48. https://doi.org/doi:10.18637/jss.v067.i01

Bazzaz, F. A., Chiariello, N. R., Coley, P. D., & Pitelka, L. F. (1987). Allocating Resources to Reproduction and Defense. *Bioscience, 37*(1), 58-67. https://doi.org/10.2307/1310178

Berner, L. T., Massey, R., Jantz, P., Forbes, B. C., Macias-Fauria, M., Myers-Smith, I., . . . Goetz, S. J. (2020). Summer warming explains widespread but not uniform greening in the Arctic tundra biome. *Nature Communications, 11*(1), 4621. https://doi.org/10.1038/s41467-020-18479-5

Bintanja, R., & Andry, O. (2017). Towards a rain-dominated Arctic. *Nature Climate Change, 7*(4), 263-267. https://doi.org/10.1038/nclimate3240

Bjerke, J. W. (2011). Winter climate change: Ice encapsulation at mild subfreezing temperatures kills freeze-tolerant lichens. *Environmental and Experimental Botany, 72*(3), 404-408. https://doi.org/https://doi.org/10.1016/j.envexpbot.2010.05.014

Bjerke, J. W., Elvebakk, A., & Tømmervik, H. (2018a). Alpine garden plants from six continents show high vulnerability to ice encasement. *Norsk Geografisk Tidsskrift - Norwegian Journal of Geography, 72*(1), 57-64. https://doi.org/10.1080/00291951.2017.1391876

Bjerke, J. W., Elverland, E., Jaakola, L., Lund, L., Zagajewski, B., Bochenek, Z., . . . Tømmervik, H. (2018b). High tolerance of a high-arctic willow and graminoid to simulated ice encasement. *Boreal Environment Research, 23*, 329–338.

Bjerke, J. W., Treharne, R., Vikhamar-Schuler, D., Karlsen, S. R., Ravolainen, V., Bokhorst, S., . . . Tommervik, H. (2017). Understanding the drivers of extensive plant damage in boreal and arctic ecosystems: Insights from field surveys in the aftermath of damage. *Science of the Total Environment, 599-600*, 1965-1976. https://doi.org/10.1016/j.scitotenv.2017.05.050

Bjorkman, A. D., García Criado, M., Myers-Smith, I. H., Ravolainen, V., Jónsdóttir, I. S., Westergaard, K. B., . . . Normand, S. (2020). Status and trends in Arctic vegetation: Evidence from experimental warming and long-term monitoring. *Ambio, 49*(3), 678-692. https://doi.org/10.1007/s13280-019-01161-6

601 Bjorkman, A. D., Myers-Smith, I. H., Elmendorf, S. C., Normand, S., Rüger, N., Beck, P. S. A., . . .
602   Weiher, E. (2018). Plant functional trait change across a warming tundra biome. *Nature,*
603   *562*(7725), 57-62. https://doi.org/10.1038/s41586-018-0563-7

604
605 Bokhorst, S., Bjerke, J. W., Bowles, F. W., Melillo, J., Callaghan, T. V., & Phoenix, G. K. (2008). Impacts
606   of extreme winter warming in the sub-Arctic: growing season responses of dwarf shrub
607   heathland. *Global Change Biology, 14*, 2603–2612. https://doi.org/10.1111/j.1365-
608   2486.2008.01689.x

609
610 Bokhorst, S., Bjerke, J. W., Davey, M. P., Taulavuori, K., Taulavuori, E., Laine, K., . . . Phoenix, G. K.
611   (2010). Impacts of extreme winter warming events on plant physiology in a sub-Arctic heath
612   community. *Physiologia Plantarum, 140*(2), 128-140. https://doi.org/10.1111/j.1399-
613   3054.2010.01386.x

614
615 Bokhorst, S., Bjerke, J. W., Street, L. E., Callaghan, T. V., & Phoenix, G. K. (2011). Impacts of multiple
616   extreme winter warming events on sub-Arctic heathland: phenology, reproduction, growth,
617   and CO2 flux responses. *Global Change Biology, 17*(9), 2817-2830.
618   https://doi.org/10.1111/j.1365-2486.2011.02424.x

619
620 Bokhorst, S., Huiskes, A., Aerts, R., Convey, P., Cooper, E. J., Dalen, L., . . . Dorrepaal, E. (2013).
621   Variable temperature effects of Open Top Chambers at polar and alpine sites explained by
622   irradiance and snow depth. *Global Change Biology, 19*(1), 64-74.
623   https://doi.org/10.1111/gcb.12028

624
625 Bokhorst, S., Pedersen, S. H., Brucker, L., Anisimov, O., Bjerke, J. W., Brown, R. D., . . . Callaghan, T. V.
626   (2016). Changing Arctic snow cover: A review of recent developments and assessment of
627   future needs for observations, modelling, and impacts. *Ambio, 45*(5), 516-537.
628   https://doi.org/10.1007/s13280-016-0770-0

629
630 Bokhorst, S. F., Bjerke, J. W., Tømmervik, H., Callaghan, T. V., & Phoenix, G. K. (2009). Winter
631   warming events damage sub-Arctic vegetation: consistent evidence from an experimental
632   manipulation and a natural event. *Journal of Ecology, 97*(6), 1408-1415.
633   https://doi.org/10.1111/j.1365-2745.2009.01554.x

634
635 Bråthen, K. A., & Hagberg, O. (2004). More efficient estimation of plant biomass. *Journal of*
636   *Vegetation Science, 15*(5), 653-660. https://doi.org/10.1111/j.1654-1103.2004.tb02307.x

637
638 Brooker, R., & van der Wal, R. (2003). Can soil temperature direct the composition of high arctic plant
639   communities? *Journal of Vegetation Science, 14*(4), 535-542.
640   https://doi.org/https://doi.org/10.1111/j.1654-1103.2003.tb02180.x

641
Buchwal, A., Sullivan, P. F., Macias-Fauria, M., Post, E., Myers-Smith, I. H., Stroeve, J. C., . . . Welker, J.
M. (2020). Divergence of Arctic shrub growth associated with sea ice decline. *Proceedings of*
the National Academy of Sciences, 117(52), 33334.
https://doi.org/10.1073/pnas.2013311117

CAVM Team. (2003). *Circumpolar Arctic Vegetation Map. (1:7,500,000 scale), Conservation of Arctic*
Flora and Fauna (CAFF) Map No. 1. . U.S. Fish and Wildlife Service.

Chapin, F. S. (1991). Integrated Responses of Plants to Stress. *Bioscience, 41*(1), 29-36.
https://doi.org/10.2307/1311538

Chapin III, F. S., Bret-Harte, M. S., Hobbie, S. E., & Zhong, H. (1996). Plant functional types as
predictors of transient responses of arctic vegetation to global change. *Journal of Vegetation*
Science, 7(3), 347-358. https://doi.org/https://doi.org/10.2307/3236278

Cooper, E. J., Alsos, I. G., Hagen, D., Smith, F. M., Coulson, S. J., & Hodkinson, I. D. (2004). Plant
recruitment in the High Arctic: Seed bank and seedling emergence on Svalbard. *Journal of*
Vegetation Science, 15(1), 115-124. https://doi.org/10.1111/j.1654-1103.2004.tb02244.x

Cooper, E. J., Dullinger, S., & Semenchuk, P. (2011). Late snowmelt delays plant development and
results in lower reproductive success in the High Arctic. *Plant Science, 180*(1), 157-167.
https://doi.org/https://doi.org/10.1016/j.plantsci.2010.09.005

Coulson, S. J., Leinaas, H. P., Ims, R. A., & Søvik, G. (2000). Experimental manipulation of the winter
surface ice layer: the effects on a High Arctic soil microarthropod community. *Ecography,*
23(3), 299-306. https://doi.org/10.1111/j.1600-0587.2000.tb00285.x

Crawford, R. M. M., Chapman, H. M., & Hodge, H. (1994). Anoxia Tolerance in High Arctic Vegetation.
Arctic and Alpine Research, 26(3), 308-312.

Descals, A., Verger, A., Filella, I., Baldocchi, D., Janssens, I. A., Fu, Y. H., . . . Peñuelas, J. (2020). Soil
thawing regulates the spring growth onset in tundra and alpine biomes. *Science of the Total*
Environment, 742, 140637. https://doi.org/https://doi.org/10.1016/j.scitotenv.2020.140637

Elmendorf, S. C., Henry, G. H. R., Hollister, R. D., Björk, R. G., Bjorkman, A. D., Callaghan, T. V., . . .
Wookey, P. A. (2012). Global assessment of experimental climate warming on tundra
vegetation: heterogeneity over space and time. *Ecology Letters, 15*(2), 164-175.
https://doi.org/doi:10.1111/j.1461-0248.2011.01716.x

Elmendorf, S. C., Henry, G. H. R., Hollister, R. D., Fosaa, A. M., Gould, W. A., Hermanutz, L., . . .
Walker, M. D. (2015). Experiment, monitoring, and gradient methods used to infer climate
change effects on plant communities yield consistent patterns. *Proceedings of the National*
Academy of Sciences, 112(2), 448. https://doi.org/10.1073/pnas.1410088112

Epstein, H. E., Raynolds, M. K., Walker, D. A., Bhatt, U. S., Tucker, C. J., & Pinzon, J. E. (2012).
Dynamics of aboveground phytomass of the circumpolar arctic tundra during the past three
decades. *Environmental Research Letters, 7*(1), 015506. https://doi.org/10.1088/1748-
9326/7/1/015506

Epstein, H. E., Walker, M. D., Chapin Iii, F. S., & Starfield, A. M. (2000). A transient, nutrient-based
model of arctic plant community response to climatic warming. *Ecological Applications,*
10(3), 824-841. https://doi.org/https://doi.org/10.1890/1051-
0761(2000)010[0824:ATNBMO]2.0.CO;2

695
696   Etzelmüller, B., Guglielmin, M., Hauck, C., Hilbich, C., Hoelzle, M., Isaksen, K., . . . Ramos, M. (2020).
697   Twenty years of European mountain permafrost dynamics—the PACE legacy. *Environmental*
698   *Research Letters, 15*(10), 104070. https://doi.org/10.1088/1748-9326/abae9d

699
700   Forbes, B. C., Kumpula, T., Meschtyb, N., Laptander, R., Macias-Fauria, M., Zetterberg, P., . . . Bartsch,
701   A. (2016). Sea ice, rain-on-snow and tundra reindeer nomadism in Arctic Russia. *Biology*
702   *Letters, 12*(11). https://doi.org/10.1098/rsbl.2016.0466

703
704   Fracheboud, Y., Luquez, V., Björkén, L., Sjödin, A., Tuominen, H., & Jansson, S. (2009). The Control of
705   Autumn Senescence in European Aspen. *Plant Physiology, 149*(4), 1982.
706   https://doi.org/10.1104/pp.108.133249

707
708   Frisvoll, A. A., & Elvebakk, A. (1996). Part 2. Bryophytes. In A. Elvebakk & P. Presttrud (Eds.), *Svalbard*
709   *plants, fungi, algae and cyanobateria*. Norsk Polarinstitutt, Skrifter nr. 198.

710
711   Frost, G. V., Bhatt, U. S., Epstein, H. E., Walker, D. A., Raynolds, M. K., & Berner, L. T., et al. (2019).
712   *Tundra greenness.* (NOAA Arctic Report Card 2019, Issue.

713
714   Gauthier, G., Bêty, J., Cadieux, M.-C., Legagneux, P., Doiron, M., Chevallier, C., . . . Berteaux, D.
715   (2013). Long-term monitoring at multiple trophic levels suggests heterogeneity in responses
716   to climate change in the Canadian Arctic tundra. *Philosophical Transactions of the Royal*
717   *Society B: Biological Sciences, 368*(1624), 20120482. https://doi.org/10.1098/rstb.2012.0482

718
719   Givnish, T. (2002). Adaptive significance of evergreen vs. deciduous leaves: solving the triple paradox.
720   *Silva Fennica, 36*(3), 535. https://doi.org/doi:10.14214/sf.535

721
722   Gornall, J., Woodin, S., Jónsdóttir, I., & van der Wal, R. (2011). Balancing positive and negative plant
723   interactions: how mosses structure vascular plant communities. *Oecologia, 166*(3), 769-782.
724   https://doi.org/10.1007/s00442-011-1911-6

725
726   Graham, R. M., Cohen, L., Petty, A. A., Boisvert, L. N., Rinke, A., Hudson, S. R., . . . Granskog, M. A.
727   (2017). Increasing frequency and duration of Arctic winter warming events. *Geophysical*
728   *Research Letters, 44*(13), 6974-6983. https://doi.org/https://doi.org/10.1002/2017GL073395

729
730   Hansen, B. B., Grotan, V., Aanes, R., Saether, B. E., Stien, A., Fuglei, E., . . . Pedersen, A. O. (2013).
731   Climate events synchronize the dynamics of a resident vertebrate community in the High
732   Arctic. *Science, 339*(6117), 313-315. https://doi.org/10.1126/science.1226766

733
734   Hansen, B. B., Isaksen, K., Benestad, R. E., Kohler, J., Pedersen, Å. Ø., Loe, L. E., . . . Varpe, Ø. (2014).
735   Warmer and wetter winters: characteristics and implications of an extreme weather event in

the High Arctic. *Environmental Research Letters, 9*, 114021. https://doi.org/10.1088/1748-9326/9/11/114021

Hanssen-Bauer, I., Førland, E. J., Hisdal, H., Mayer, S., Sandø, A. B., & Sorteberg, A. (2019). *Climate in Svalbard 2100 – a knowledge base for climate adaptation (NCCS Report No. 1/2019).* https://klimaservicesenter.no/kss/rapporter/rapporter-og-publikasjoner_2

Henry, G. H. R., & Molau, U. (1997). Tundra plants and climate change: the International Tundra Experiment (ITEX). *Global Change Biology, 3*(S1), 1-9. https://doi.org/10.1111/j.1365-2486.1997.gcb132.x

Holmgren, M., Stapp, P., Dickman, C. R., Gracia, C., Graham, S., Gutiérrez, J. R., . . . Squeo, F. A. (2006). Extreme climatic events shape arid and semiarid ecosystems. *Frontiers in Ecology and the Environment, 4*(2), 87-95. https://doi.org/10.1890/1540-9295(2006)004[0087:ECESAA]2.0.CO;2

Hurley, M. A., Hebblewhite, M., Gaillard, J.-M., Dray, S., Taylor, K. A., Smith, W. K., . . . Bonenfant, C. (2014). Functional analysis of Normalized Difference Vegetation Index curves reveals overwinter mule deer survival is driven by both spring and autumn phenology. *Philosophical Transactions of the Royal Society B: Biological Sciences, 369*(1643), 20130196. https://doi.org/10.1098/rstb.2013.0196

Jónsdóttir, I. S. (1991). Effects of Grazing on Tiller Size and Population Dynamics in a Clonal Sedge (Carex bigelowii). *Oikos, 62*(2), 177-188. https://doi.org/10.2307/3545263

Jónsdóttir, I. S. (2011). Diversity of plant life histories in the Arctic. *Preslia, 83*(3), 281-300.

Ju, J., & Masek, J. G. (2016). The vegetation greenness trend in Canada and US Alaska from 1984–2012 Landsat data. *Remote Sensing of Environment, 176*, 1-16. https://doi.org/10.1016/j.rse.2016.01.001

Karlsen, S., Elvebakk, A., Høgda, K., & Grydeland, T. (2014). Spatial and temporal variability in the onset of the growing season on Svalbard, Arctic Norway — Measured by MODIS-NDVI satellite data. *Remote Sensing, 6*(9), 8088-8106. https://doi.org/10.3390/rs6098088

Kohler, J., & Aanes, R. (2004). Effect of winter snow and ground-icing on a svalbard reindeer population: results of a simple snowpack model. *Arctic, Antarctic, and Alpine Research, 36*(3), 333-341.

Le Moullec, M., Buchwal, A., van der Wal, R., Sandal, L., & Hansen, B. B. (2019). Annual ring growth of a widespread high-arctic shrub reflects past fluctuations in community-level plant biomass. *Journal of Ecology, 107*(1), 436-451. https://doi.org/10.1111/1365-2745.13036

Le Moullec, M., Sandal, L., Grøtan, V., Buchwal, A., & Hansen, B. B. (2020). Climate synchronises
shrub growth across a high-arctic archipelago: contrasting implications of summer and winter
warming. *Oikos, 129*(7), 1012-1027. https://doi.org/10.1111/oik.07059

Loe, L. E., Liston, G. E., Pigeon, G., Barker, K., Horvitz, N., Stien, A., . . . Albon, S. D. (2021). The
neglected season: Warmer autumns counteract harsher winters and promote population
growth in Arctic reindeer. *Global Change Biology, 27*, 993-1002.
https://doi.org/https://doi.org/10.1111/gcb.15458

Meredith, M., Sommerkorn, M., Cassotta, S., Derksen, D., Ekaykin, A., Hollowed, A., . . . Schuur, E. A.
G. (2019). Polar Regions. In H.-O. Pörtner, D. C. Roberts, V. Masson-Delmotte, P. Zhai, M.
Tignor, E. Poloczanska, K. Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama, &
N. M. Weyer (Eds.), *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*.

Miller, F. L., & Gunn, A. (2003). Catastrophic die-off of Peary caribou on the Western Queen Elizabeth
Islands, Canadian High Arctic. *Arctic, 56*(4), 381-390.

Milner, J. M., Varpe, Ø., Van der Wal, R., & Hansen, B. B. (2016). Experimental icing affects growth,
mortality, and flowering in a high arctic dwarf shrub. *Ecology and Evolution, 6*(7), 2139-2148.
https://doi.org/10.1002/ece3.2023

Myers-Smith, I. H., Forbes, B. C., Wilmking, M., Hallinger, M., Lantz, T., Blok, D., . . . Hik, D. S. (2011).
Shrub expansion in tundra ecosystems: dynamics, impacts and research priorities.
Environmental Research Letters, 6(4), 045509. https://doi.org/10.1088/1748-
9326/6/4/045509

Myers-Smith, I. H., Kerby, J. T., Phoenix, G. K., Bjerke, J. W., Epstein, H. E., Assmann, J. J., . . . Wipf, S.
(2020). Complexity revealed in the greening of the Arctic. *Nature Climate Change, 10*(2), 106-
117. https://doi.org/10.1038/s41558-019-0688-1

Orcutt, D. M., & Nilsen, E. T. (2000). *The physiology of plants under stress: soil and biotic factors*. John
Wiley and Sons.

Outcalt, S. I., Nelson, F. E., & Hinkel, K. M. (1990). The zero-curtain effect: Heat and mass transfer
across an isothermal region in freezing soil. *Water Resources Research, 26*(7), 1509-1516.
https://doi.org/https://doi.org/10.1029/WR026i007p01509

Pan, C. G., Kirchner, B. P., Kimball, S. J., Kim, Y., & Du, J. (2018). Rain-on-snow events in Alaska, their
frequency and distribution from satellite observations. *Environmental Research Letters,*
13(7), 075004. http://stacks.iop.org/1748-9326/13/i=7/a=075004

Parmesan, C., Root, T. L., & Willig, M. R. (2000). Impacts of Extreme Weather and Climate on
Terrestrial Biota. *Bulletin of the American Meteorological Society, 81*(3), 443-450.
https://doi.org/10.1175/1520-0477(2000)081<0443:IOEWAC>2.3.CO;2

Paus, E., Nilsen, J., & Junttila, O. (1986). Bud dormancy and vegetative growth in Salix polaris as
825   affected by temperature and photoperiod. *Polar Biology, 6*(2), 91-95.
826   https://doi.org/10.1007/BF00258258

827
828   Peeters, B., Pedersen, A. O., Loe, L. E., Isaksen, K., Veiberg, V., Stien, A., . . . Hansen, B. B. (2019).
829   Spatiotemporal patterns of rain-on-snow and basal ice in high Arctic Svalbard: detection of a
830   climate-cryosphere regime shift. *Environmental Research Letters, 14*(1), 015002.
831   https://doi.org/10.1088/1748-9326/aaefb3

832
833   Pérez-Harguindeguy, N., Díaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., . . .
834   Cornelissen, J. H. C. (2013). New handbook for standardised measurement of plant functional
835   traits worldwide. *Australian Journal of Botany, 61*(3), 167-234.
836   https://doi.org/10.1071/BT12225

837
838   Petraglia, A., Carbognani, M., & Tomaselli, M. (2013). Effects of nutrient amendments on modular
839   growth, flowering effort and reproduction of snowbed plants. *Plant Ecology & Diversity, 6*(3-
840   4), 475-486. https://doi.org/10.1080/17550874.2013.795628

841
842   Phoenix, G. K., & Bjerke, J. W. (2016). Arctic browning: extreme events and trends reversing arctic
843   greening. *Global Change Biology, 22*(9), 2960-2962. https://doi.org/10.1111/gcb.13261

844
845   Post, E., & Forchhammer, M. C. (2008). Climate change reduces reproductive success of an Arctic
846   herbivore through trophic mismatch. *Philosophical Transactions of the Royal Society B:*
847   *Biological Sciences, 363*(1501), 2367-2373. https://doi.org/10.1098/rstb.2007.2207

848
849   Post, E., Forchhammer, M. C., Bret-Harte, M. S., Callaghan, T. V., Christensen, T. R., Elberling, B., . . .
850   Aastrup, P. (2009). Ecological dynamics across the Arctic associated with recent climate
851   change. *Science, 325*(5946), 1355-1358. https://doi.org/10.1126/science.1173113

852
853   Preece, C., Callaghan, T. V., & Phoenix, G. K. (2012). Impacts of winter icing events on the growth,
854   phenology and physiology of sub-arctic dwarf shrubs. *Physiologia Plantarum, 146*(4), 460-
855   472. https://doi.org/10.1111/j.1399-3054.2012.01640.x

856
857   Preece, C., & Phoenix, G. K. (2014). Impact of early and late winter icing events on sub-arctic dwarf
858   shrubs. *Plant Biology, 16*(1), 125-132. https://doi.org/10.1111/plb.12015

859
860   Prevéy, J. S., Rixen, C., Rüger, N., Høye, T. T., Bjorkman, A. D., Myers-Smith, I. H., . . . Wipf, S. (2019).
861   Warming shortens flowering seasons of tundra plant communities. *Nature Ecology &*
862   *Evolution, 3*(1), 45-52. https://doi.org/10.1038/s41559-018-0745-6

863
864   Putkonen, J., & Roe, G. (2003). Rain-on-snow events impact soil temperatures and affect ungulate
865   survival. *Geophysical Research Letters, 30*(4). https://doi.org/10.1029/2002GL016326

866
867   R Core team. (2020). *R: a language and environment for statistical computing*. R Foundation for
868   Statistical Computing.

Rasmus, S., Kivinen, S., & Irannezhad, M. (2018). Basal ice formation in snow cover in Northern
Finland between 1948 and 2016. *Environmental Research Letters, 13*(11), 114009.
https://doi.org/10.1088/1748-9326/aae541

Rennert, K. J., Roe, G., Putkonen, J., & Bitz, C. M. (2009). Soil thermal and ecological impacts of rain
on snow events in the circumpolar Arctic. *Journal of Climate, 22*(9), 2302-2315.
https://doi.org/10.1175/2008JCLI2117.1

Schindelin, J., Rueden, C. T., Hiner, M. C., & Eliceiri, K. W. (2015). The ImageJ ecosystem: an open
platform for biomedical image analysis. *Molecular Reproduction and Development, 82*(7-8),
518-529. https://doi.org/10.1002/mrd.22489

Schuur, E. A. G., McGuire, A. D., Schädel, C., Grosse, G., Harden, J. W., Hayes, D. J., . . . Vonk, J. E.
(2015). Climate change and the permafrost carbon feedback. *Nature, 520*(7546), 171-179.
https://doi.org/10.1038/nature14338

Semenchuk, P. R., Elberling, B., & Cooper, E. J. (2013). Snow cover and extreme winter warming
events control flower abundance of some, but not all species in high arctic Svalbard. *Ecology
and Evolution, 3*(8), 2586-2599. https://doi.org/10.1002/ece3.648

Semenchuk, P. R., Gillespie, M. A. K., Rumpf, S. B., Baggesen, N., Elberling, B., & Cooper, E. J. (2016).
High Arctic plant phenology is determined by snowmelt patterns but duration of
phenological periods is fixed: an example of periodicity. *Environmental Research Letters,
11*(12), 125006. https://doi.org/10.1088/1748-9326/11/12/125006

Skarpe, C., & Van der Wal, R. (2002). Effects of simulated browsing and length of growing season on
leaf characteristics and flowering in a deciduous arctic shrub, *Salix polaris*. *Arctic, Antarctic,
and Alpine Research, 34*(3), 282-286.

Treharne, R., Bjerke, J. W., Tømmervik, H., & Phoenix, G. K. (2020). Development of new metrics to
assess and quantify climatic drivers of extreme event driven Arctic browning. *Remote Sensing
of Environment, 243*, 111749. https://doi.org/https://doi.org/10.1016/j.rse.2020.111749

Treharne, R., Bjerke, J. W., Tømmervik, H., Stendardi, L., & Phoenix, G. K. (2019). Arctic browning:
Impacts of extreme climatic events on heathland ecosystem CO2 fluxes. *Global Change
Biology, 25*(2), 489-503. https://doi.org/10.1111/gcb.14500

Van der Wal, R., & Stien, A. (2014). High-arctic plants like it hot: a long-term investigation of
between-year variability in plant biomass. *Ecology, 95*(12), 3414-3427.
https://doi.org/10.1890/14-0533.1

Vickers, H., Høgda, K. A., Solbø, S., Karlsen, S., Tømmervik, H., Aanes, R., & Hansen, B. B. (2016).
Changes in greening in the high arctic: insights from a 30 year AVHRR max NDVI dataset for
Svalbard. *Environmental Research Letters, 11*, 105004. https://doi.org/10.1088/1748-
9326/11/10/105004

Vikhamar-Schuler, D., Isaksen, K., Haugen, J. E., Tømmervik, H., Luks, B., Schuler, T. V., & Bjerke, J. W.
917   (2016). Changes in Winter Warming Events in the Nordic Arctic Region. *Journal of Climate,*
918   *29*(17), 6223-6244. https://doi.org/10.1175/JCLI-D-15-0763.1

920 Wahren, C. H., Walker, M., & Bret-Harte, M. (2005). Vegetation responses in Alaskan arctic tundra
921   after 8 years of a summer warming and winter snow manipulation experiment. *Global*
922   *Change Biology, 11*(4), 537-552.

924 Walker, M. D., Wahren, C. H., Hollister, R. D., Henry, G. H. R., Ahlquist, L. E., Alatalo, J. M., . . .
925   Wookey, P. A. (2006). Plant community responses to experimental warming across the
926   tundra biome. *Proceedings of the National Academy of Sciences, 103*(5), 1342.
927   https://doi.org/10.1073/pnas.0503198103

929 Wipf, S., & Rixen, C. (2010). A review of snow manipulation experiments in Arctic and alpine tundra
930   ecosystems. *Polar Research, 29*(1), 95-109. https://doi.org/10.3402/polar.v29i1.6054

932 Wood, S. N., Goude, Y., & Shaw, S. (2015). Generalized additive models for large data sets. *Journal of*
933   *the Royal Statistical Society: Series C (Applied Statistics), 64*(1), 139-155.
934   https://doi.org/10.1111/rssc.12068

936 You, Q., Cai, Z., Pepin, N., Chen, D., Ahrens, B., Jiang, Z., . . . Zhang, Y. (2021). Warming amplification
937   over the Arctic Pole and Third Pole: Trends, mechanisms and consequences. *Earth-Science*
938   *Reviews, 217*, 103625. https://doi.org/https://doi.org/10.1016/j.earscirev.2021.103625

940 Zhang, W., Döscher, R., Koenigk, T., Miller, P. A., Jansson, C., Samuelsson, P., . . . Smith, B. (2020). The
941   Interplay of Recent Vegetation and Sea Ice Dynamics—Results From a Regional Earth System
942   Model Over the Arctic. *Geophysical Research Letters, 47*(6), e2019GL085982.
943   https://doi.org/10.1029/2019GL085982