Including imprecisely georeferenced specimens improves accuracy of species distribution
models and estimates of niche breadth: Don't let the perfect be the enemy of the good

Running title: Using imprecisely geolocated occurrences

Authors:

Adam B. Smith[1*] [ORCiD: 0000-0002-6420-1659]
Stephen J. Murphy[1,2] [ORCiD: 0000-0001-8679-2851]
David Henderson[3] [ORCiD: 0000-0003-3523-0175]
Kelley D. Erickson[1] [ORCiD: 0000-0001-7980-9985]

[1] Center for Conservation and Sustainable Development, Missouri Botanical Garden, 4344 Shaw
Boulevard, Saint Louis MO 63110 USA

[2] Present address: The Ohio State University, Department of Evolution, Ecology, and
Organismal Biology, 318 W 12th Avenue, Columbus, OH 43210 USA, murphy.1132@osu.edu

[3] Department of Biology, Washington University in Saint Louis, 1 Bookings Drive, Saint Louis
MO 63130 USA, hdavid@wustl.edu

* Corresponding author

Abstract [300 words]

Museum and herbarium specimen records are frequently used to assess species' conservation

status and responses to climate change. Typically, records of occurrence with imprecise

geolocality information are discarded because they cannot be matched confidently to

environmental conditions, and are thus expected to increase uncertainty in downstream analyses.

However, using only those records that can be precisely georeferenced risks undersampling of

species' environmental and geographic distributions. Using simulated and real species, we

compared the effect of discarding versus retaining imprecise records on the accuracy of

ecological niche models (ENMs), and estimates of niche breadth and extent of occurrence.

Imprecise records were assigned to locations or climate values using a conservative approach.

For simulated species, including imprecisely-georeferenced records alongside precisely-

georeferenced records improved the accuracy of ENMs projected to the present and the future.

Improvements were especially notable for species with <~20 precise occurrences, but accuracy

continued to improve and eventually matched the accuracy of ENMs using all records with no

georeferencing error. Using only precise records underestimated loss in suitable habitat and

overestimated the amount of suitable habitat in both the present and future. Including imprecise

records also improved univariate and multivariate estimates of niche breadth, and of species'

extent of occurrence. An analysis of 44 species in the genus *Asclepias* (milkweeds) revealed

similarly large differences between cases using only precise records and those using precise plus

imprecise records. Although the inclusion of imprecisely-georeferenced occurrence records may

not always be appropriate, careful consideration must be made regarding the trade-offs between

fidelity of the match between occurrences and the environment, versus sample size and how well

precisely-geolocated records sample species' environmental and geographic distributions. The

preponderance of imprecisely georeferences specimens in museums and herbaria could be

gainfully employed to address known shortfalls in sampling of species' distributions and climatic

niche tolerances.

Keywords

climate change vulnerability, coordinate uncertainty, georeferencing, niche breadth, natural

history museum specimen records, niche truncation, rare species

Introduction

Accurate estimation of species' environmental tolerances and distributions is key to addressing

many pressing issues in ecology, evolution, and conservation (e.g., Fisher-Reid et al. 2012;

Foden et al. 2013; Quintero & Wiens 2013a and b). Information on species' environmental

tolerances and distributions is commonly inferred from occurrence records obtained from natural

history museums and herbaria. These records can be matched to associated environmental

conditions and then used to calculate the range of environments inhabited by a species (e.g.,

Foden et al. 2013). Similarly, ecological niche models (ENMs; also known as species

distribution models) can be employed to estimate niche breadth and predict the extent of suitable

habitat (Dawson et al. 2011; Pacifici et al. 2015; Foden & Young 2016; Miller et al. 2012;

Young et al. 2016; IUCN 2019). Conservation assessments also frequently utilize the extent of

occurrence, or area encompassed by occurrence records, to appraise species' potential exposure

to broad-scale threats (IUCN 2019; Faber-Langendoen et al. 2012). Methods utilizing

opportunistic occurrence records have thus become an important component of biodiversity and

conservation research (Heberling 2020). Unfortunately, a large portion of these records is

imprecisely geolocated (Moudrý & Devillers 2020), which poses challenges for estimating

niches and distributions (Moudrý & Šímová 2012).

For occurrence data to be useful for estimating niche limits and distributions, two key conditions

are desirable. First, environmental information assigned to an occurrence record should reflect

the conditions that were actually experienced by the species at the location where it was

observed (Graham et al. 2008). When records can only be geolocated imprecisely (i.e., to a large

region or geopolitical unit), it becomes difficult to confidently assign a specific environmental

datum to each record (Feeley & Silman 2010). Second, when the goal is to estimate

environmental tolerances, the set of records used to estimate niche limits should encompass as

much of the species' fundamental niche as possible (Thuiller et al. 2004; Peterson et al. 2018). If

available occurrences represent only a portion of the niche, then the species' actual

environmental tolerances will be underestimated (Thuiller et al. 2004; Qiao et al. 2019).

The desire for spatial precision and representative sampling of occurrences leads to a critical

trade-off. On the one hand, if records can only be imprecisely geolocated, using them risks

introducing uncertainty and bias into estimates of environmental tolerances (Graham et al. 2008;

Fernandez et al. 2009; Osborne & Leitão 2009; Feeley & Silman 2010; Tulowiecki et al. 2015;

Gábor et al. 2020; Mitchell et al. 2016; Collins et al. 2017; Cheng et al. 2021). On the other

hand, discarding records risks under-representing the true geographic and environmental range of

a species, even if the location of occurrences is uncertain (Graham et al. 2008). These risks are

especially great for rare species, which tend to be represented by just a few records (Lomba et al.

2010; Sheth et al. 2012; Zizka et al. 2018).

To date, these trade-offs have been almost exclusively managed by discarding imprecisely

georeferenced records (Moudrý & Šímová 2012). Indeed, in a literature survey of publications

using museum or herbarium specimens with ENMs, we found that half of studies that included

any description of their data cleaning process discarded spatially imprecise records, and no

studies reported purposefully retaining them (Supplementary material Appendix 1 Fig. A2). In

other words, the risk of including inaccurate climate data has been assumed more serious than

undersampling of the realized niche. However, the justification for discarding spatially

imprecise records is often based on studies that do not necessarily reflect real-world use cases.

Instead, these studies evaluate the effects of coordinate imprecision on the accuracy of ENMs by

artificially adding spatial error to otherwise precise records, and then comparing results between

"fuzzed" and accurate records (e.g., Graham et al. 2008; Fernandez et al. 2009; Osborne &

Leitão 2009; Gueta & Carmel 2016; Mitchell et al. 2016; Hefley et al. 2017; Soultan & Safi

2017; Tulowiecki et al. 2015; Gábor et al. 2020). This approach, however, is not reflective of the

common situation where an assessor starts with a mix of relatively precisely- and imprecisely-

geolocated records and must decide how to delineate the two groups and whether or not to

discard the imprecise ones.

Here we reexamine the trade-offs between retaining versus discarding spatially imprecise records

using virtual and real species. We designated records as "precise" if they had spatial uncertainty

small enough to match them with confidence to environmental data, and "imprecise" if they

could not. Imprecision in record coordinates can be represented in a variety of ways, including

through the use of user-defined polygons (Wieczorek et al. 2004) or by assignment of records to

the smallest geopolitical unit encompassing the area of likely collection (e.g., county, parish,

state, etc.; Park & Davis 2017). We emphasize that our definition of an imprecise record does not

include records with locations appearing to be outside the range of the species (i.e., geographic

outliers; Feeley & Silman 2010) or specimens that do not pass quality-assurance checks

(Chapman 2005). We compared ENMs, niche breadth, and the spatial extents of occurrence

estimated using only precise records to estimates based on precise plus imprecise records. We

hypothesized that the inclusion of imprecise records alongside precise records would improve the

accuracy of predicted distributions, estimates of univariate and multivariate niche breadth, and

extents of occurrence. We evaluated the accuracy of each of these metrics as a function of how

many imprecise records were included, and compared them to benchmark estimates based on

"omniscient" records where all populations of a species could be georeferenced without error.

Methods

Study region

For both virtual and real species, the region of analysis encompassed North America (i.e.,

Canada, the United States, and Mexico, excluding distant islands such as the Hawaiian Islands).

However, we calibrated and evaluated ENMs within species-specific regions as noted below.

Virtual species

We used virtual species, for which we could obtain "true" distributions and environmental

relationships, to explore the effects of including versus excluding imprecisely geolocated

specimens. Full details are found in Fig. 1 and Supplementary material Appendix 2, so are

briefly described here. We first calculated a principal component (PC) analysis of all present-day

climatic conditions for North America using all 19 BIOCLIM variables (Nix 1986) from

WorldClim Version 2.1 at 10 arcmin resolution, which represents average conditions across

1970-2000 (Fick & Hijmans 2017). Species' niches were generated using a trivariate Gaussian

distribution in the first three PC axes, then projected to North America (Fig. 1, steps). We

interpreted these values as probabilities of presence.

We then created occurrence data for each species by sampling raster cells according to their

probability of occurrence times cell area. We first drew a set of "omniscient" records, which

reflect an ideal case where a collector is able to georeference every single population of a species

with negligible error. We generated species with 20, 40, 80, 160, and 320 omniscient records.

From the omniscient records we randomly sampled a subset of 5, 10, 15, 20, 25, or 30 to

represent "precise" records (so long as the number of precise records was less than the number of

omniscient records). Precise records represent a realistic case where a collector has access to a

subset of the species' occurrences georeferenced with little error. To represent "imprecise"

records that could only be georeferenced to a county, parish, or equivalent, from the remainder of

omniscient records we sampled a subset of 1, 2, 4, 8, …, records, up to the total number of

omniscient records minus the number of precise records. We then assigned these records to the

county in which they fell.

For each combination of number of omniscient, precise, and imprecise records, we generated 200

species. We then constructed ENMs and estimated climatically suitable area and niche breadth

for each species in three separate ways. First, we calibrated an ENM and niche breadth metrics

using all omniscient occurrences to represent a baseline against which to assess the other two

sets of models and metrics. Second, we calculated a "precise-only" ENM and metrics using just

the precise records, recreating the common practice of discarding imprecise records before

biogeographic analysis. Third, we combined the precise and imprecise records to generate

"precise + imprecise" ENMs and metrics.

ENMs: We modeled each species' realized niche using MaxEnt (Phillips et al. 2006; Phillips &

Dudík 2008) with linear, quadratic, and interaction terms, with the optimal set of terms and

master regularization parameter identified using AIC_c (Warren & Siefert 2011). Models were

projected to the present and to a future climate scenario defined by the average across five earth

system models for 2061-2080 from CMIP6 under Representative Concentration Pathway 8.5

(achievable under Shared Socioeconomic Pathway 5-8.5; O'Neill et al. 2015). We assessed

calibration accuracy (the degree to which model output reflects the true probability of presence)

of the model predictions in the present and the future using the Pearson correlation coefficient

between the ENM predictions and the actual probability of presence. Background sites for model

calibration and assessment were drawn from a 300-km buffer around the minimum convex

polygon constructed around all omniscient points. We also assessed area of climatically suitable

habitat within this same region by applying a threshold such that training sensitivity was 0.9. We

calculated the area of current and future favorable habitat, and the area of habitat that was lost,

gained, or remained favorable. To determine how inclusion of imprecise records affected model

complexity, we calculated the number of non-zero coefficients (ignoring the intercept) for each

type of model.

Niche breadth and extent of occurrence: We also calculated a set of metrics broadly used in

ecology, evolution, and conservation: (a) univariate niche breadth in mean annual temperature

(MAT) and total annual precipitation (TAP), estimated from the range of the inner 90% of values

of each variable across each set of occurrences; (b) multivariate niche breadth, represented by the

volume and surface area of a minimum convex hull around each set of occurrences in three-

dimensional environmental space; and (c) extent of occurrence (EOO), which is the area of the

minimum convex polygon encompassing all records in geographic space (IUCN 2019). EOO is

used in conservation assessments as an index of the propensity for multiple populations to

experience the same broad scale threat (IUCN 2019).

Assigning imprecise records locations and environments: Since the imprecise records could not,

by design, be geolocated with confidence to a single cell, we used a simple yet conservative

approach to assign climatic conditions to these records. For the ENMs and multivariate niche

breadth, we used the PC score of the cell in each imprecise record's county that was closest in

PC space to the mean of the precise records' PC scores. To calculate univariate niche breadth, we

used the value of MAT or TAP from across each imprecise record's county that was closest to

the mean across the precise records (Fig. 2b and c). To calculate extent of occurrence, we

assigned each imprecise occurrence to the point in the county where it was located closest to the

centroid of the precise records (Fig. 2a). For example, the actual location of an imprecise record

in a county was almost always further from the precise records' centroid than the closest point in

the county.

We did not formally compare results between data types (omniscient, precise, precise plus

imprecise) using statistical tests since these are inappropriate for simulations where the existence

of differences is known a priori (White et al. 2014). Rather, we compared the inner 90th-

percentile distribution of each metric (calibration accuracy, EOO, etc.) for each case.

Real species

We also evaluated 44 species of *Asclepias* (milkweeds; family Apocynaceae) native to North

America, which display a range distributions from narrowly endemic to those covering

approximately one-third of the continent. Records were obtained from the Global Biodiversity

Information Facility (www.gbif.org), and detailed procedures for data cleaning and modeling are

described in Supplementary material Appendices 3 and 4. For this analysis we used only

herbarium specimens and species with ≥5 precise and unique (i.e., non-duplicate) records. We

evaluated the same set of metrics as for the virtual species and created ENMs following the same

procedures. However, because we did not have omniscient records for the real species, we

estimated climatically suitable habitat area within the region defined by a 300-km buffer around

the minimum convex polygon surrounding all available records. We tested for differences in

EOO and in univariate and multivariate niche breadth calculated with or without imprecise

records using a paired Wilcoxon signed-rank test.

Reproducibility

The analysis relied primarily on the sp (Bivand et al. 2013), rgeos (Bivand & Rundel 2020),

geosphere (Hijmans 2019), dismo (Hijmans et al. 2017), raster (Hijmans 2021), and enmSdm

(Smith 2021) packages for R Version 4.10 (R Core Team 2021). Scripts used in this analysis are

available on the GitHub repository https://github.com/adamlilith/enms_impreciseRecords.

Results

Virtual species

For brevity, we focus on results for species with 40 and 320 omniscient records, each with 5 or

20 precise records (see Supplementary material Appendix 2 for all results). We found that adding

imprecise records caused the greatest improvements in metrics when the number of precise

records was <15-20, although the accuracy of some measures continued to improve as imprecise

records were added. To reflect real-world use cases where assessors may have a set of records

that can only be located to a geopolitical unit but not otherwise know if they represent truly

geographically unique specimens, we plotted the change in each metric (ENM accuracy, niche

breadth, etc.) against the number of county-level records accrued as imprecise records were

added. Hence, hereafter we refer to changes in metrics as a function of the number of counties

encompassing the imprecise records (counties are only counted once, regardless of the number of

imprecise records they contain).

ENMs: Compared to precise-only models, precise + imprecise models were more accurate when

projected to both present and future climate scenarios (Figs. 3a and b, and Appendix 2 Figs. A1

and A2). Accuracy increased most rapidly when the number of precise records in precise +

imprecise models was ≤15, but continued to rise thereafter. For example, for present-day climate

and a species with 40 total occurrences and only 5 precise occurrences, the median correlation

between the true probability of presence and predictions from precise-only ENMs was 0.15 (90%

inner quantiles: -0.02 to 0.58). Adding county records increased the correlation up to 0.63 (0.30-

0.86), which was equivalent to the accuracy of omniscient ENMs (median: 0.60, 90% quantile:

0.27-0.81). Similar results were obtained for species with a greater number of total occurrences.

When the number of precise records was ≥20 or when >~30 county records were included,

increases in accuracy of precise + imprecise models were more gradual (Fig. 3a). However, for

species with ≥160 total occurrences, there was a second notable increase in accuracy when the

number of county records surpassed about half of the total number of omniscient records (Fig. 3a

and b). Accuracy decreased when ENMs were projected to the future, although precise +

imprecise models still outperformed precise-only models when a sufficient number of county

records was included (Fig. 3b). Precise-only models were simpler (fewer non-zero coefficients)

than omniscient and precise + imprecise models (Supplementary material Appendix 2 Fig. A13).

Compared to estimates based on omniscient records, precise-only ENMs overestimated the area

of climatically suitable habitat by up to >300% (median value) for the present (Fig. 3c) and by

>500% for the future (Fig. 3d and Supplementary material Appendix Figs. A6 and A7).

Overestimation was greatest when the number of precise occurrences was <15. Adding just ~10-

30 county-level records reduced bias to ~0 relative to omniscient models.

When the number of precise records was ≤15, precise-only ENMs underestimated loss of

suitable area by 90% or more (median; Fig. 3e and f, Supplementary material Appendix 2 Fig.

A3). Adding 10-30 county records eliminated this bias. Gains in suitable area were on average

unbiased (Supplementary material Appendix 2 Fig. A4). Precise-only ENMs overestimated area

that remained climatically suitable through time (Supplementary material Appendix 2 Fig. A5).

Niche breadth: Using just precise records underestimated univariate niche breadth and

multivariate niche volume and surface area (Fig. 4b-e and Supplementary material Appendix 2

Figs. A8 to A11). For example, for a species with 40 total occurrences of which just 5 could be

geolocated precisely, precise-only niche breadth in MAT was only 29% (6-71%) of omniscient

niche breadth. However, precise + imprecise niche breadth in temperature increased up to ~90%

(~70-110%) of omniscient breadth as county records were added. For species with ≥20 precise

occurrences, improvements in estimates of niche breadth were less dramatic. Similar trends were

observed in niche breadth in TAP, and in multivariate niche volume and surface area (Fig. 4c-d).

Occasionally, adding imprecise data led to overestimation, especially for niche breadth in MAT

and niche volume. For MAT, across all species and scenarios, 4.5% of precise + imprecise

estimates were larger than omniscient values, with the median of overestimated values 5% larger

than the true value. For niche volume, 9% of precise + imprecise estimates were larger than

omniscient values, with the median of overestimated values 27% larger than the true value. For

MAT, overestimation was more common when the number of precise records was large and

additional county records was small (Fig. 4b), but for niche volume overestimation was more

likely as county records increased (Fig. 4d).

Extent of occurrence: EOO was consistently underestimated when only precise records were

used (Fig. 4a and Supplementary material Appendix 2 Fig. A12). Underestimation was worse for

species with few precise records and a large number of occurrences. In these cases, median

estimated EOO was as small as 12% of the actual value (inner 90% quantile range: 2-37%).

Adding county records improved estimates in nearly every case, though the rate of improvement

declined as more county records were added. EOO was occasionally overestimated when county

records were included. Across all scenarios, 1.7% of precise/imprecise estimates were larger than

omniscient values, with the median of overestimate values 47% larger than the true value.

Asclepias

The data obtained from GBIF comprised 53,623 herbarium records. Following data cleaning,

removal of observational and duplicate records, and elimination of species with fewer than 5

geographically unique precise records, we were left with just 16% of the original records (8,480)

and only 32% of the species (44 of 137). Imprecise records were the most abundant, comprising

on average 70% of all usable records for a species (range: 44-97%; Supplementary material

Appendix 3 Table A1). Thirty-six percent (16 of 44) of species had ≤20 precise records and

twenty percent (9 of 44) had ≤10.

ENMs: Relative to precise-only ENMs, precise + imprecise models predicted greater climatically

suitable area for 63% of species (28 of 44) in the present and 71% of species (31 of 44) in the

future (Fig. 5a and b, Supplementary material Appendix A7 Figs. A3). Gains and losses in

suitable area were also larger when imprecise records were included (median gain 16% ranging

from -80 to 3952%; median loss 30% ranging from -89 to 1652%; Supplementary material

Appendix A7 Fig. A5).

Niche breadth: Including imprecise records increased median univariate niche breadth in MAT

and TAP by 25% across species (range: 0 to 353%) and 28% (0 to 292%), respectively (Fig. 5d;

Wilcoxon V ~0 and $P<10^{-6}$ in both cases). Including imprecise records increased multivariate

niche volume by a median value of 175% (8 to 13,909%; $P<10^{-12}$, Wilcoxon V~0) and niche

surface area by 79% (3 to 1515%; $P<10^{-12}$, Wilcoxon V~0; Fig. 5e). Species with the fewest

records had the greatest increase in niche breadth, volume, and area when imprecise records were

included.

Extent of occurrence: Precise + imprecise EOO was 86% larger (median; range 0-2011%) than

precise-only EOO (Fig. 5c). Including imprecise records at least doubled EOO for 34% of

species (15 of 44), and at least tripled EOO for 27% of species (12 of 44). Species with fewer

precise records tended head greater increases in EOO when imprecise records were included.

Discussion

Spatially imprecise occurrences are commonly discarded prior to biogeographic analyses

(Appendix 1 Fig. S2). Although the justification for this decision is due to the potential for

spatial error to propagate through an analysis (Moudrý & Šímová 2012), discarding records also

risks introducing error by undersampling geographic and environmental space. We found that

including geospatially-imprecise records increased the accuracy of niche models projected to

both present and future climate conditions (Fig. 3a and b). Including these records also led to

more accurately estimated climatically suitable area (Fig. 3c-e) and improved estimates of niche

breadth and extent of occurrence (EOO; Fig. 4). Sometimes estimates using just precise records

were orders-of-magnitude different from those based on "omniscient" records, but adding

imprecise records helped close this gap.

A critical distinction of our work is that we show the effects of adding imprecise records to

precise records, whereas most studies on coordinate uncertainty demonstrate the effects of

making precise records spatially imprecise (e.g., Graham et al. 2008; Fernandez et al. 2009;

Soultan & Safi 2017; Gábor et al. 2020). Likewise, studies that have examined the effects of

adding additional imprecise records to ENMs have typically not employed a conservative

method for assigning environmental values or locations to imprecise records (e.g., Bloom et al.

2018; Collins et al. 2017; Cheng et al. 2021), or have focused on mean values (versus extremes,

which define environmental limits; Pender et al. 2019). Our findings have direct implications for

studies in ecology, evolution, and conservation that use occurrences to estimate species'

relationships to the environment. More broadly, the uncertainty inherent in the decision over

whether to use or discard imprecise records should be reflected in the outcome of analyses

relying on occurrence data.

Why including imprecise records improves niche models and estimates of niche breadth

The effect of imprecise records on ENM accuracy and niche breadth depends on the abundance

of precise records and how well they sample geographic and niche space (Moudrý & Šímová

2012; Tulowiecki et al. 2015; Soultan & Safi 2017). We found that including a sufficient number

of imprecise records could fully compensate for a lack of precise records even when there were

as few as 5 precise occurrences. Gains in accuracy were most notable for species with <~20

precise records, but even species >20 precise occurrences experienced improvements from

adding imprecise records. Sample size is one of the largest influences on niche model accuracy

(Santini et al. 2021), with minimum recommended sizes ranging from <10 to several hundred

(Wisz et al. 2008; van Proossdij et al. 2016; Santini et al. 2021; see also Rivers et al. 2011).

Given that many species are known from just a handful of records (Zizka et al. 2018), including

imprecise records could be especially helpful when sample sizes are small. Adding imprecise

records can improve the sampling of geographic and environmental space (e.g., Fig. 2a).

Implications for studies in evolution, ecology, and conservation

Although the amount of biodiversity data is growing at an astounding rate, most species'

distributions remain poorly characterized (Meyer et al. 2016). For North American *Asclepias*,

even with >53,000 specimen records we had enough data to analyze only a third species found in

North America. Of those we did analyze, 70% of their records would have normally been

discarded due to spatial uncertainty in their locations. Similar rates of spatial uncertainty are

common in biodiversity databases (Moudrý & Devillers 2020).

Spatially imprecise records, if used carefully, have great potential to address the "Wallacean

shortfall," the lack of information on species' distributions (Hortal et al. 2015), and the

"Hutchinsonian" shortfall, the lack of information on species' environmental tolerances

(Cosentino & Maiorano 2021). Answers to many key questions in ecology and evolution are

susceptible to these shortfalls, and so could be informed by addition of erstwhile "unusable"

imprecise records. For example, inadequate representation of environmental conditions

inhabitable by a species can bias estimates of the rate of climatic niche evolution (Saupe et al.

2018) and alter species' response curves along environmental gradients (Hannemann et al. 2016).

Likewise, investigations of relationships between niche breadth, geographic range size and

environmental variation (Quintero & Wiens 2013a and b), and measurements of niche overlap

(Warren et al. 2008) will be inherently sensitive to the degree to which realized niches are

adequately sampled. The many studies that rely on ENMs for reconstructing species' past,

present, and potential future distributions are especially sensitive to sample size (Santini et al.

2021) and uneven sampling intensity among inhabitable environments (Raes 2012; Perret & Sax

2022).

Discarding imprecise records has the potential to overestimate species' vulnerability and thus

bias conservation assessments. For example, under IUCN Red List criterion B1, species qualify

as threatened if they have an EOO <100 km^2 (in addition to other criteria; IUCN 2019; see also

Young et al. 2016). While none of the 44 species of *Asclepias* in our analysis had an EOO <100

km^2 when using just precise records, it is certainly possible that this threshold could be crossed

by some of the other 93 species in our original data that we did not analyze because they had <5

geographically unique precise records. Similarly, estimates of species' adaptive capacities (Cang

et al. 2016), rates of community thermophilization (Feeley et al. 2020), and exposure to

anticipated climate change (Fig. 3) could be misrepresented by undersampling due to removal of

imprecise specimens.

Our intent is to provoke a reconsideration of the benefits and costs of discarding spatially

imprecise records when assessing a species' response to environmental gradients and geographic

distributions. These trade-offs must be assessed within the goals and philosophical approach of

an analysis. For example, many conservation assessments adopt a precautionary strategy that errs

on the side of assuming a species is more vulnerable than it may actually be (Moyle 2005;

Huntley et al. 2016a; IUCN 2019). In contrast, an evidentiary approach aims to classify species

as vulnerable only if there is strong evidence to support such a designation (IUCN 2019).

Discarding imprecise records decreased estimates of niche breadth and EOO (Fig. 4), so aligns

with a precautionary approach because the species appear more vulnerable then they may be.

However, using just precise records did overestimate climatically suitable area for the virtual

species (Fig. 3c and 3d). In contrast, niche breadth and EOO were occasionally overestimated

when imprecise records were included (Fig. 4). Hence, whether or not a decision to retain versus

keep imprecise records is precautionary or evidentiary depends on the metric used to assess

vulnerability.

In this context, we did find some differences between virtual and real species. Specifically,

including imprecise records reduced the estimated climatically suitable area in the present and

future for virtual species, but increased it for real species (Fig. 3c and d versus Fig. 5d and f; cf.

Collins et al. 2017). Inspection of the individual ENMs for virtual species revealed that models

that overestimated suitable habitat were: a) largely relegated to cases with small sample size

(total number of precise or precise plus imprecise records) and small EOO; and b) were much

simpler (i.e., had only a single term or were intercept-only models; Supplementary material

Appendix 2 Fig. A13). Small sample size favors simpler models because available information

does not justify complex responses (Phillips et al. 2006; Warren & Siefert 2011; Merow et al.

2014; Vignali et al 2020). However, simple models containing, say, just linear terms cannot

represent unimodal responses along environmental gradients (Whittaker 1953). Rather, they

predict that suitability increases along a gradient, even if conditions eventually become worse

again. As a result, and somewhat ironically, overly simple models tended to predict that suitable

habitat for species with few occurrences is larger than for the same species with more

occurrences, which allowed for more complex models (Brun et al. 2020). We did not see this

effect in the real species, but 70% of our real species (31 of 44) had >15 precise records, so

models were less likely to be simple.

Uncertainty

Analyses are most informative when they account for all relevant aspects of uncertainty (Huntley

et al. 2016b; IUCN 2019). The decisions over whether to retain or discard imprecise records,

and indeed, over what constitutes an "imprecise" record, represent key aspects of uncertainty.

However, across all of the articles that we reviewed, none of them evaluated the consequences of

discarding imprecise records (Supplementary material Appendix 1 Fig. A2). Ignoring the

subjectivity inherent in these decisions reduces apparent uncertainty in the final assessment of

species' distributions and environmental tolerances. Acknowledging uncertainty is important

even in cases where the exclusion of imprecise records would not change a species' overall

vulnerability status, since increased uncertainty reduces the reliability of returns on conservation

investment (Smith et al. 2016).

Conclusions

We advocate for a re-consideration over whether spatially imprecise occurrence records should

be excluded from biogeographic analyses. Using only precise records reduces niche model

accuracy, and can underestimate niche breadth and extent of occurrence. The decision over how

to define imprecise records and whether or not to use them is an important contributor to the

overall uncertainty in inherent in the outcome of analyses relying on specimen data. Discarding

imprecise records ignores a critical aspect of uncertainty and risks undersampling species'

realized environmental and geographic distributions. Practitioners need to consider the trade-offs

between using versus discarding imprecise records, especially given the preponderance of

imprecise records available in specimen databases and the Wallacean and Hutchinsonian

shortfalls that beset our sampling of the distribution of life on Earth.

Acknowledgements

DH was supported by the Department of Biology of Washington University in Saint Louis. SJM

and KDE were supported by the Institute of Museum and Library Services (FAIN MG-30-15-

0094-15) to ABS. This research was also supported by the Alan Graham Fund in Global Change.

Conflict of Interest Statement

The authors declare no conflict of interest.

Biosketch

The Global Change Conservation Laboratory at the Missouri Botanical Garden identifies

solutions to pressing environmental problems by leveraging precise and the imprecise

information on biodiversity. We are inspired by the cumulative person-millennia of field work

and curatorial attention devoted to amassing collection records of Earth's species and the

exigency of using this data to its fullest potential.

Literature cited

Bivand, R. and Rundel, C. 2020. rgeos: Interface to Geometry Engine: Open Source ('GEOS').
Version 0.5-5. https://CRAN.R-project.org/package=rgeos
Bivand, R., Pebesma, E., and Gómez-Rubio, V. 2013. Applied Spatial Data Analysis with R.
Springer-Verlag, New York.
Bloom, T.D.S., Flower, A., and DeChaine, E.G. 2018. Why georeferencing matters:
Introducing a practical protocol to prepare species occurrence records for spatial analysis.
Ecology and Evolution 8:765-777.
Brun, P., Thuiller, W., Chauvier, Y., Pellissier, L., Wüest, R.O., Wang, Z., and Zimmermann,
N.E. 2020. Model complexity affects species distribution projections under climate change.
Journal of Biogeography 47:130-142.
Cang, F.A., Wilson, A.A., and Wiens, J.J. 2016. Climate change is projected to outpace rates of
niche change in grasses. Biology Letters 12: 20160368.
Chapman, A.D. 2005. Principles and Methods of Data Cleaning: Primary Species and Species-
Occurrence Data, Version 1.0. Report for the Global Biodiversity Information Facility,
Copenhagen.
Cheng, Y., Tjaden, N.B., Jaeschke, A., Thomas, S.M, and Beierkuhnlein, C. 2021. Using
centroids of spatial units in ecological niche modeling: Effects on model performance in the
context of environmental data grain size. Global Ecology and Biogeography 30:611-621.

Collins, S.D., Abbott, J.C., and McIntyre, N.E. 2017. Quantifying the degree of bias from using

county-scale data in species distribution modeling: Can increasing sample size or using

county-averaged environmental data reduce distributional overprediction? Ecology and

Evolution 7:6012-6022.

Cosentino, F. and Laiorano, L. 2021. Is geographic sampling bias representative in

environmental space? Ecological Informatics 64:101369.

Dawson, T.P., Jackson, S.T., House, J.I., Prentice, I.C., and Mace, G.M. 2011. Beyond

predictions: Biodiversity conservation in a changing climate. Science 332:53-58.

Faber-Langendoen, D., J. Nichols, L. Master, K. Snow, A. Tomaino, R. Bittman, G. Hammerson,

B. Heidel, L. Ramsay, A. Teucher, and B. Young. 2012. NatureServe Conservation Status

Assessments: Methodology for Assigning Ranks. NatureServe, Arlington, VA.

Feeley, K.J. and Silman, M.R. 2010. Modelling the responses of Andean and Amazonian plant

species to climate change: The effects of georeferencing errors and the importance of data

filtering. Journal of Biogeography 37:733-740.

Feeley, K.J., Bravo-Avila, C., Fadrique, B., Perez, T.M., and Zuleta, D. 2020. Climate-driven

changes in the composition of New World plant communities. Nature Climate Change

10:965-970.

Fernandez, M.A., S.D. Blum, S. Reichle, Q. Guo, B. Holzman, and H. Hamilton. 2009. Locality

uncertainty and the differential performance of four common niche-based modeling

techniques. Biodiversity Informatics 6:36-52.

Fick, S.E. and Hijmans, R.J. 2017. WorldClim 2: New 1-km spatial resolution climate surfaces

for global land areas. International Journal of Climatology 37:4302-4315.

Fisher-Reid, M.C., Kozak, K.H., and Wiens, J.J. 2012. How is the rate of climatic-niche

evolution related to climatic-niche breadth? Evolution 66:3836-3851.

Foden, W.B. and Young, B.E. (eds.) 2016. IUCN SSC Guidelines for Assessing Species'

Vulnerability to Climate Change. Version 1.0. Occasional Paper of the IUCN Species

Survival Commission No. 59. Cambridge, UK and Gland, Switzerland: IUCN Species

Survival Commission. x+114 pp.

Foden, W.B., Mace, G.M., and Butchart, S.H.M. 2013. Indicators of climate change impacts on

biodiversity. Pp. 120-137 in Collin, B., Pettorelli, N., Baillie, J.E.M., and Durant, S.M.

(eds.) Biodiversity Monitoring and Conservation: Bridging the Gap between Global

Commitment and Local Action, 1st ed. John Wiley and Sons, Indianapolis.

Gábor, L., Moudrý, V., Lecours, V., Malavasi, M., Barták, V., Fogl, M., Šímová, P., Rocchini,

D., and Václavík, T. 2020. The effect of positional error on fine scale species distribution

models increases for specialist species. Ecography 43:256-269.

Graham, C.H., Elith, J., Hijmans, R.J., Guisan, A., Peterson, A.T., Loiselle, B.A., and the

NCEAS Predicting Species Distributions Working Group. 2008. The influence of spatial

errors in species occurrence data used in distribution models. Journal of Applied Ecology

45:239-247.

Gueta, T. and Carmel, Y. 2016. Quantifying the value of user-level cleaning for big data: A

case study using mammal distribution models. Ecological Informatics 34:139-145.

Hannemann, H., Willis, K.J., and Macias-Fauria, M. 2016. The devil is in the details: Unstable

response functions in species distribution models challenge bulk ensemble modeling. Global

Ecology and Biogeography 25:26-35.

Heberling, J.M. 2020. Global change biology: Museum specimens are more than meet the eye.

Current Biology 30: R1368-R1370.

Hefley, T.J., Brost, B.M., and Hooten, M.B. 2017. Bias correction of bounded location errors in

presence-only data. Methods in Ecology and Evolution 8:1566-1573.

Hijmans, R.J. 2016. raster: Geographic Data Analysis and Modeling. R package version 2.5-8.

https://CRAN.R-project.org/package=raster.

Hijmans, R.J. 2019. geosphere: Spherical Trigonometry. R package version 1.5-10.

https://CRAN.R-project.org/package=geosphere.

Hijmans, R.J., Phillips, S.J., Leathwick, J., and Elith, J. 2020. dismo: Species Distribution

Modeling. R package version 1.3-3. https://CRAN.R-project.org/package=dismo.

Hortal, J., de Bello, F., Diniz-Filho, J.A.F., Lewinson, T.M., Lobo, J.M., and Ladle, R.J. 2015.

Seven shortfalls that beset large-scale knowledge of biodiversity. Annual Review of Ecology,

Evolution, and Systematics 46:523-549.

Huntley, B., Foden, W.B., Pearce-Higgins, J., and Smith, A.B. 2016. Chapter 6. Understanding

and working with uncertainty. In W.B. Foden and B.E. Young, editors. IUCN SSC

Guidelines for Assessing Species' Vulnerability to Climate Change. Version 1.0. Occasional

Paper of the IUCN Species Survival Commission No. 59. Gland, Switzerland and

Cambridge, UK. pp 49-56.

Huntley, B., Foden, W.B., Smith, A.B., Platts, P., Watson, J. and Garcia, R.A. 2016. Chapter 5.

Using CCVAs and interpreting their results. In W.B. Foden and B.E. Young, editors. IUCN

SSC Guidelines for Assessing Species' Vulnerability to Climate Change. Version 1.0.

Occasional Paper of the IUCN Species Survival Commission No. 59. Gland, Switzerland and

Cambridge, UK. pp 33-48.

IUCN Standards and Petitions Committee. 2019. Guidelines for Using the IUCN Red List

Categories and Criteria. Version 14. Prepared by the Standards and Petitions Committee.

Downloadable from http://www.iucnredlist.org/documents/RedListGuidelines.pdf (2021-05-

24).

Lomba, A., L. Pellissier, C. Randin, J. Vicente, J. Horondo, and A. Guisan. 2010. Overcoming

the rare species modeling complex: A novel hierarchical framework applied to an Iberian

endemic plant. Biological Conservation 143:2647-2657.

Merow, C., Allen, J.M., Aiello-Lammens, M., and Silander, Jr., J.A. 2016. Improving niche and

range estimates with Maxent and point process models by integrating spatially explicit

information. Global Ecography and Biogeography 25:1022-1036.

Merow, C., Smith, M.J., Edwards, Jr., T.C., Guisan, A., McMahon, S.M., Normand, S., Thuiller,

W., Wüest, R.O., Zimmermann, N.E., and Elith, J. 2014. What do we gain from simplicity

versus complexity in species distribution models? Ecography 37:1267-1281.

Miller, J.S., Porter-Morgan, H.A., Stevens, H., Boom, B., Krupnick, G.A., Acevedo-Rodríguez,

P., Fleming, J., and Gensler, M. 2012. Addressing targets two of the Global Strategy for

Plant Conservation by rapidly identifying plants at risk. Biodiversity Conservation 21:1877-

1887.

Mitchell, P.J., Monk, J., and Laurenson, L. 2016. Sensitivity of fine-scale species distribution

models to locational uncertainty in occurrence data across sample sizes. Methods in Ecology

and Evolution 8:12-21.

Moudrý, V. and Devillers, R. 2020. Quality and usability challenges of global marine

biodiversity databases: An example for marine mammal data. Ecological Informatics

56:101051.

Moudrý, V. and Šímová, P. 2012. Influence of positional accuracy, sample size and scale on
modeling species distributions: A review. International Journal of Geographic Information
Science 26:2083-2095.

Moyle, B. 2005. Making the Precautionary Principle work for biodiversity: Avoiding perverse
outcomes in decision-making under uncertainty. Pp. 159-172 in Cooney, R. and B. Dickson
(eds.) Biodiversity and the Precautionary Principle: Risk and Uncertainty in Conservation
and Sustainable Use. Earthscan, London. 314 pp.

Nix, H.A. 1986. A biogeographic analysis of Australian elapid snakes. Atlas of elapid snakes of
Australia: Australian flora and fauna series 7 (ed. by R. Longmore), pp. 4-15. Bureau of
Flora and Fauna, Canberra.

O'Neill, B.C., Kreigler, E., Ebi, K.L., Kemp-Benedict, E., Riahi, K., Rothman, D.S., van
Ruijven, B., van Vuuren, D.P., Birkman, J., Kok, K., Levy, M., and Solecki, W. 2017. The
roads ahead: Narratives for shared socioeconomic pathways describing world futures in the
21st century. Global Environmental Change 42:169-180.

Osborne, P.E. and Leitão, P.J. 2009. Effects of species and habitat positional errors on the
performance and interpretation of species distribution models. Diversity and Distributions
15:671-681.

Pacifici, M., Foden, W.B., Visconti, P., Watson, J.E.M., Butchart, S.H.M., Kovacs, K.M.,
Scheffers, B.R., Hole, D.G., Martin, T.G., Akçakaya, H.R., Corlett, R.T., Huntley, B.,
Brickford, D., Carr, J.A., Hoffmann, A.A., Midgley, G.F., Pearce-Kelly, P. Pearson, R.G.,
Williams, S.E., Willis, S.G., Yoing, B., and Rondinini, C. 2015. Assessing species
vulnerability to climate change. Nature Climate Change 5:215-225.

Park, D.S. and Davis, C.C. 2017. Implications and alternatives of assigning climate data to
geographical centroids. Journal of Biogeography 44:2188-2198.

Pender, J.E., Hipp, A.L., Hahn, M., Kartesz, J., Nishino, M., and Starr, J.R. 2019. How
sensitive are climatic niche inferences to distribution data sampling? A comparison of Biota
of North America Program (BONAP) and Global Biodiversity Information Facility (GBIF)
datasets. Ecological Informatics 54:100991.

Perret, D.L. and Sax, D.F. 2022. Evaluating a study design for optimal sampling of species'
climatic niches. Ecography 2022:e06014.

Peterson, A.T., Cobos, M.E., and Jiménez-García, D. 2018. Major challenges for correlational
ecological niche model projections to future climate conditions. Annals of the New York
Academy of Sciences 1429:66-77.

Phillips, S.J. and Dudík, M. 2008. Modeling species distributions with Maxent: New extensions
and a comprehensive evaluation. Ecography 31:161-175.

Phillips, S.J., Anderson, R.P., and Schapire, R.E. 2006. Maximum entropy modeling of species
geographic distributions. Ecological Modelling 190:231-259.

Qiao, H., Feng, X., Escobar, L.E., Peterson, A.T., Soberón, J., Zhu, G., and Papeş. 2019. An
evaluation of transferability of ecological niche models. Ecography 42:521-534.

Quintero, I. and Wiens, J.J. 2013a. What determines the climatic niche width of a species? The
role of spatial and temporal climatic variation in three vertebrate clades. Global Ecology and
Biogeography 22:422-432.

Quintero, I. and Wiens, J.J. 2013b. Rates of projected climate change dramatically exceed past
rates of climate niche evolution among vertebrate species. Ecology Letters 16:1095-1103.

R Core Team. 2021. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Raes, N. 2012. Partial versus full species distribution models. Natureza and Conservação
10:127-138.

Rivers, M.C., Taylor, L., Brummitt, N.A., Meagher, T.R., Roberts, D.L., and Lughadha, E.N.
2011. How many herbarium specimens are needed to detect threatened species? Biological
Conservation 144:2541-2547.

Santini, L., Benítez-López, Maiorano L., Čengič, M., and Huijbreghts, M.A. 2021. Assessing
the reliability of species distribution projections in climate change research. Diversity and
Distributions 27:1035-1050.

Saupe, E.E. Barve, N., Owens, H.L., Cooper, J.C., Hosner, P.A., and Peterson, A.T. 2018.
Reconstructing niche evolution when niches are incompletely characterized. Systematic
Biology 67:428-438.

Sheth, S.N., L.G. Lohmann, T. Distler, and I. Jiménez. 2012. Understanding bias in geographic
range size estimates. Global Ecology and Biogeography 21:732-742.

Smith, A.B. 2021. enmSdm: Tools for modeling niches and distributions of species. R package
version 0.7.0. http://github.com/adamlilith/enmSdm

Smith, A.B., Long, Q.G., and Albrecht, M.A. 2016. Shifting targets: spatial priorities for ex situ

plant conservation depend on interactions between current threats, climate change, and

uncertainty. Biodiversity and Conservation 25:905-922.

Soultan, A. and Safi, K. 2017. The interplay of various sources of noise and reliability on

species distribution models hinges on ecological specialization. Public Library of Science

ONE 12:e0187906.

Thuiller, W., Araújo, M.B., Pearson, R.G., Whittaker, R.J., Brotons, L., and Lavorel, S. 2004.

Uncertainty in predictions of extinction risk. Nature 430:33.

Tulowiecki, S.J., Larsen, C.P.S., and Wang, Y-C. 2015. Effects of positional error on modeling

species distributions: A perspective using presettlement land survey records. Plant Ecology

216:67-85.

van Proossdij, A.S.J., Sosef, M.S.M., Wieringa, J.J. and Raes, N. 2016. Minimum number of

specimen records to develop accurate species distribution models. Ecography 39:542-552.

Vignali, S., Barras, A.G., Arlettaz, R. and Braunisch, V. 2020. SDMtune: An R package to tune

and evaluate species distribution models. Ecology and Evolution 10:11488-11506.

Warren, D.L. and S.N. Siefert. 2011. Ecological niche modeling in Maxent: The importance of

model complexity and the performance of model selection criteria. Ecological Applications

21:335-342.

Warren, D.L., Glor, R.E., and Turelli, M. 2008. Environmental niche equivalency versus

conservatism: Quantitative approaches to niche evolution. Evolution 62:2868-2883.

White, J.W., Rassweiler, A., Samhouri, J.F., Stier, A.C., and White, C. 2014. Ecologists should

not use statistical significance tests to interpret simulation model results. Oikos 123:385-388.

Whittaker, R.H. 1953. A consideration of climax theory: the climax as a population and a

pattern. Ecological Monographs 23:41-78.

Wieczorek, J., Guo, Q., and Hijmans, R.J. 2004. The point-radius method for georeferencing

locality descriptions and calculating associated uncertainty. International Journal for

Geographical Information Science 18:745-767.

Wisz, M.S., Hijmans, R.J., Peterson, A.T., Graham, C.H., Guisan, A., and NCEAS Predicting

Species Distributions Working Group. 2008. Effects of sample size on the performance of

species distribution models. Diversity and Distributions 14:763-773.

Young, B.E., Byers, E., Hammerson, G., Frances, A., Oliver, L., and Treher, A. 2016.

Guidelines for Using the NatureServe Climate Change Vulnerability Index, Version 3.02.

NatureServe, Arlington.

Zizka, A., ter Steege, H., Pessoa, M. de C.R, and Antonello, A. 2018. Finding needles in the

haystack: Where to look for rare species in the American tropics. Ecography 41:321-330.

Figure 1. The process of generating and analyzing a virtual species starting with generation of

the fundamental niche (step 1), projection to geographic space (2), and generation of omniscient

(3), precise (4), and imprecise (5) occurrences. Suitable climate area is analyzed within a

buffered region surrounding omniscient records (bottom row). The process was repeated 200

times for each combination of number of omniscient, precise, and imprecise records.

Figure 2. An example of including versus discarding imprecise records using *Asclepias viridiflora*. a) Extent of occurrence estimated using either just precise or precise plus imprecise records from the minimum convex polygon (MCP). Imprecise records are represented by the smallest geopolitical unit to which a record can be located, or by circles representing area of likely collection. Projection: Albers conic, equal-area. b) and c) Difference in climatic niche breadth. The values of temperature or precipitation at precise records are represented by points (green). The distributions of temperature or precipitation across all locations encompassed by imprecise records are represented by smoothed density kernels, one per record (orange). Niche breadths estimated using just precise versus precise plus imprecise are shown.

Figure 3. Adding imprecise records (here, reflected as the number of counties they occupy) improves niche model accuracy for the present (a) and future (b). Calibration accuracy is the correlation between the real probability of presence and model output. Current (c) and future (d) climatically suitable area are overestimated when only precise records are included. Values are the ratio between estimates using only precise or precise + imprecise records to estimates from models using all "omniscient" occurrences of a species. Note the log scales. Each point represents a species. Violins encompass the inner 90% quantile of values. Thick trendlines represent the median trend, and thin trendlines encompass the inner 90% quantile of values.

a) Extent of occurrence

b) Univariate niche breadth in temperature

c) Univariate niche breadth in precipitation

d) Multivariate niche volume

e) Multivariate niche surface area

Figure 4. Adding imprecise records (here, reflected as the number of counties they occupy) increases the accuracy of estimates of extent of occurrence (a), univariate niche breadth in mean annual temperature (b) and total annual precipitation (c), and multivariate niche volume (d) and surface area (e). In each case, values represent the ratio of estimates using only precise or precise + imprecise records to estimates calculated using all "omniscient" occurrences of a species. Each point represents a species. Violins encompass the inner 90% quantile of values. Thick trendlines represent the median trend, and thin trendlines encompass the inner 90% quantile of values. To aid visualization for univariate niche breadth and niche volume, the y-axis limits encompass only the lower 99.9% of values.

Figure 5. Effects of including imprecise records on niche breadth and climatically suitable area
for *Asclepias*. Differences in climatically suitable area in the present (a) and future (b) under
RCP8.5. (c) Change in extent of occurrence. (d) Change in univariate niche breadth in mean
annual temperature and mean annual precipitation. (e) Change in multivariate niche volume and
surface area of this volume. In panels c-e, species are sorted from top to bottom from most to
least number of precise records. Also note the log scale along the x-axis.

34