A putative *de novo* evolved gene required for spermatid chromatin condensation in *Drosophila melanogaster*

Emily L. Rivard[1,4,*], Andrew G. Ludwig[1,*], Prajal H. Patel[1,*], Anna Grandchamp[2], Sarah E. Arnold[1,5], Alina Berger[2], Emilie M. Scott[1], Brendan J. Kelly[1,6], Grace C. Mascha[1], Erich Bornberg-Bauer[2,3], Geoffrey D. Findlay[1,**]

1. Department of Biology, College of the Holy Cross, Worcester, MA, USA
2. Institute for Evolution and Biodiversity, University of Münster, Münster, Germany
3. Department of Protein Evolution, Max Planck Institute for Developmental Biology, Tübingen, Germany
4. Current address: Department of Molecular and Cellular Biology, Harvard University, Cambridge, MA, USA
5. Current address: Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA, USA
6. Current address: Department of Entomology, The Ohio State University, Columbus, OH, USA

*equal contribution
**Correspondence to: gfindlay@holycross.edu

Abstract

Comparative genomics has enabled the identification of genes that potentially evolved *de novo* from non-coding sequences. Many such genes are expressed in male reproductive tissues, but their functions remain poorly understood. To address this, we conducted a functional genetic screen of over 40 putative *de novo* genes with testis-enriched expression in *Drosophila melanogaster* and identified one gene, *atlas,* required for male fertility. Detailed genetic and cytological analyses show that *atlas* is required for proper chromatin condensation during the final stages of spermatogenesis. Atlas protein is expressed in spermatid nuclei and facilitates the transition from histone- to protamine-based chromatin packaging. Complementary evolutionary analyses revealed the complex evolutionary history of *atlas*. The protein-coding portion of the gene likely arose at the base of the *Drosophila* genus on the X chromosome but was unlikely to be essential, as it was then lost in several independent lineages. Within the last ~15 million years, however, the gene moved to an autosome, where it fused with a conserved non-coding RNA and evolved a non-redundant role in male fertility. Altogether, this study provides insight into the integration of novel genes into biological processes, the links between genomic innovation and functional evolution, and the genetic control of a fundamental developmental process, gametogenesis.

44   **<u>Introduction</u>**
45
The evolution of new genes is integral to the extensive genotypic and phenotypic
diversity observed across species. The best-characterized mechanism of novel gene
emergence is gene duplication (Zhang 2003; Lipinski et al. 2011); however, rapid expansion in
high-quality genomic resources has provided mounting evidence of lineage-specific sequences
and the existence of alternative modes of new gene origination. One such mechanism is *de*
novo evolution, the birth of new genes from previously non-genic or intronic regions, which is
now a widely acknowledged source of protein-coding and RNA genes (McLysaght and Hurst
2016; Bornberg-Bauer and Schmitz 2017; Van Oss and Carvunis 2019). Although *de novo*
origination was once considered an unlikely event, catalogs of *de novo* genes have now been
published for an expansive range of species (Zhao et al. 2014; Guerzoni and McLysaght 2016;
Li et al. 2016; Ruiz-Orera et al. 2016; Lu et al. 2017; Zhang et al. 2019; Chamakura et al. 2020;
Puntambekar et al. 2020). Multiple models explain how protein-coding *de novo* genes may
acquire both an open reading frame (ORF) and regulatory sequences permitting transcription
(Carvunis et al. 2012; Schlötterer 2015; Wilson et al. 2017; Schmitz et al. 2018). Interrogation
of the biochemical and biophysical properties of the proteins encoded by *de novo* genes has
offered initial insight into the mechanisms of emergence and functional potential of these genes
(Schmitz et al. 2018; Vakirlis et al. 2018; Heames et al. 2020; Lange et al. 2021).
The capacity of protein-coding *de novo* genes to evolve important functions is a topic of
interest from evolutionary, physiological and molecular perspectives (Keeling et al. 2019). In the
last couple of decades, the products of *de novo* genes have been shown to play diverse roles in
a variety of organisms. For example, *de novo* genes function in fundamental molecular
processes in yeast, such as *BSC4*, a gene implicated in DNA repair, and *MDF1,* which mediates
crosstalk between reproduction and growth (Cai et al. 2008; Li et al. 2014). *De novo* genes also
evolve roles in organismal responses to disease and changing environmental factors. A
putatively *de novo* evolved gene in rice regulates the plant's pathogen resistance response to
strains causing bacterial blight (Xiao et al. 2009). Antifreeze glycoprotein genes, essential for
survival in frigid ocean temperatures, evolved *de novo* in the ancestor of Arctic codfishes to
coincide with cooling oceans in the Northern Hemisphere (Baalsrud et al. 2018; Zhuang et al.
2019). *De novo* genes are additionally implicated in the development and physiology of
mammals. In house mice, a *de novo* evolved gene expressed in the oviduct functions in female
fertility by regulating pregnancy cycles (Xie et al. 2019). A *de novo* gene found in humans and
chimpanzees regulates the oncogenesis and growth of neuroblastoma, revealing the relevance
of novel genes to human disease (Suenaga et al. 2014). These studies have started to
demonstrate the significance of *de novo* genes, thereby challenging previous assumptions that
only ancient, highly conserved genes can be essential.
Across multicellular animals, male reproductive tissues serve as hubs for new gene
emergence via numerous mechanisms, including *de novo* evolution (Marques et al. 2005;
Levine et al. 2006; Begun et al. 2007; Kaessmann 2010; Baker et al. 2012; Cui et al. 2015;
Ruiz-Orera et al. 2016). Proposed causes of this "out of the testis" phenomenon include the
high level of promiscuous transcription in testis cells (Soumillon et al. 2013; Necsulea and
Kaessmann 2014), the relative simplicity of promoter regions driving expression in the testis
(Sorourian et al. 2014), and preferential retention of novel genes with male-biased expression
(Palmieri et al. 2014). Sexual selection also drives rapid evolution of reproductive proteins
(Wilburn and Swanson 2016) and could drive the emergence of new genes as a mechanism of
improving male reproductive ability (Levine et al. 2006). The testis-biased expression of novel
genes, combined with growing evidence for new genes acting across a variety of tissue
contexts, suggests that many novel genes may function in male reproduction. For example, a
pair of young duplicate genes in *Drosophila, apollo* and *artemis,* are essential for male and
female fertility, respectively (VanKuren and Long 2018). Continued efforts to identify and

characterize testis-expressed novel genes are consequently critical for understanding the
genetic basis of male reproductive phenotypes.
Drosophila serves as an ideal system for interrogating the prevalence, sequence
attributes, expression patterns, and functions of testis-expressed *de novo* genes. The
availability of well-annotated genomes for numerous *Drosophila* species, the tractability of flies
to molecular genetics techniques, and our thorough understanding of *Drosophila* reproductive
processes facilitate comprehensive analyses of novel fly reproductive proteins (Demarco et al.
2014; Hales et al. 2015). As observed in other biological systems, *Drosophila de novo* genes
retained by selection demonstrate enriched expression in the testis (Levine et al. 2006; Begun
et al. 2007; Zhao et al. 2014; Heames et al. 2020). The expression patterns of emerging *de*
novo genes in the *Drosophila* testis were recently analyzed at single cell resolution (Witt et al.
2019), thereby providing insight into the dynamics of novel gene expression throughout
spermatogenesis. In addition to bioinformatic screens that have started to identify *de novo*
genes and large-scale expression analyses of testis-expressed genes, RNAi (Reinhardt et al.
2013) and CRISPR/Cas9-based (Kondo et al. 2017) functional screens have identified putative,
testis-expressed *de novo* genes required for fertility. However, a need remains for in-depth
experimental and evolutionary characterization of the genes identified in such screens. Detailed
examination of the function of *de novo* proteins will enable us to understand how these proteins
might integrate into existing gene networks and become essential.
We previously conducted a pilot functional screen of *de novo* genes with testis-enriched
expression in *D. melanogaster* and identified two novel genes, *goddard* and *saturn*, that are
required for full fertility (Gubala et al. 2017). *Goddard* knockdown males failed to produce any
sperm. *Saturn* knockdown males produced fewer sperm, which were inefficient at migrating to
female sperm storage organs. Subsequent characterization of Goddard using null deletion
alleles and a biochemically tagged rescue construct showed that the protein localizes to
elongating sperm axonemes and that, in its absence, individualization complexes associate less
efficiently with spermatid nuclei and do not successfully progress along sperm tails (Lange et al.
2021). These data suggested that putative *de novo* genes can evolve essential roles in a
rapidly evolving reproductive process, spermatogenesis.
Here, we expand this functional screen by evaluating whether any of 42 putative *de novo*
genes that show testis-enriched expression in *D. melanogaster* are required for male fertility.
We identified one gene, which we named *atlas,* whose knockdown or knockout results in nearly
complete male sterility. We show that *atlas* encodes a transition protein that facilitates
spermatid chromatin condensation. The *atlas* gene in *D. melanogaster* arose when a likely *de*
novo evolved protein-coding sequence moved off of the X chromosome and was inserted
upstream of a well-conserved non-coding RNA. While the *atlas* protein-coding sequence has
undergone multiple, independent gene loss events since its apparent origin at the base of the
Drosophila genus, the gene has evolved a critical function in *D. melanogaster*. These results
underscore the importance of detailed functional and evolutionary characterization in
understanding the origins of new protein-coding genes and the selective forces that affect their
subsequent evolution.
136

137   **Results**
138
139   *An RNAi screen identifies a putative* de novo *gene essential for* Drosophila *male fertility*
140
141   A previous pilot screen of 11 putative *de novo* evolved, testis-expressed genes identified
two genes that are critical for male fertility in *Drosophila melanogaster* (Gubala et al. 2017).
This result, and other recent work (e.g., Abrusán 2013; Zhang et al. 2015; VanKuren and Long
2018), suggested that lineage-specific, newly evolved genes can rapidly become important for
fertility, perhaps by gaining interactions with existing protein networks. To determine more
comprehensively the frequency with which potential *de novo* evolved genes become essential
for fertility, we identified *de novo* or putative *de novo* evolved genes with testis-biased
expression. A previous computational analysis identified genes that are detectable only within
the *Drosophila* genus, lack identifiable protein domains, and show no homology to other known
proteins through BLASTP and TBLASTN searches (Heames et al. 2020). We filtered these
genes to identify those expressed exclusively or predominantly in the testis, a common site of
de novo gene expression in animal species (Begun et al. 2007; Wu et al. 2011; Palmieri et al.
2014; Xie et al. 2019). This resulted in a set of 96 target genes.
154

155
Figure 1. An RNAi screen of putative *de novo* **genes identifies** *CG13541* **as a major contributor to**
***Drosophila melanogaster* male fertility.** A) All RNAi lines that showed at least partial knockdown of the
target gene were screened in group fertility assays (see Materials and Methods). Relative fertility was
calculated by dividing the average number of progeny produced per female mated to knockdown males
by the average number of progeny produced per female mated to control males in a contemporaneous
experiment. Relative fertility measurements lack error bars because each gene was tested in only 1-2
replicates. Knockdown of *goddard* was used as a positive control. B) A single-mating, single-pair fertility
assay confirms the observed defect when males are knocked down for *CG13541*, as knockdown males
showed significantly reduced fertility (control fertility (mean ± SEM): 109.0 ± 5.3; knockdown fertility: 0.2 ±
0.1; two-sample *t*-test assuming unequal variances, *p* = 5.6 x 10^{-13}).
166
167   We used testis-specific RNA interference to screen these genes for roles in male fertility.
We obtained RNAi lines from the Vienna *Drosophila* Resource Center (VDRC) and the
Transgenic RNAi Project (TRiP) and constructed additional lines using the TRiP-style
pValium20 vector (Ni et al. 2011), which is optimized for male germline expression. We
obtained and tested an RNAi line for 57 genes and induced knockdown for each with the *Bam*-
GAL4 driver, which is expressed in the male germline, and enhanced with a copy of UAS-
Dicer2. RT-PCR confirmed at least partial knockdown in lines representing 42 genes (see
example in Fig. S1). We then screened knockdown males for fertility by allowing groups of 7
knockdown males to mate with 5 wild-type females for 2 days. Progeny counts were
standardized to the number of progeny produced by concurrently mated groups of 7 control

males and 5 wild-type females. The results are shown in Fig. 1A. This initial screen identified
CG13541, whose knockdown severely reduced male fertility. We confirmed the result for
CG13541 by performing single-pair mating fertility assays (Fig. 1B). Knockdown of *CG43072*
and *CG33284* also showed consistent fertility defects, but these results did not replicate upon
testing with CRISPR-generated null alleles for each gene, so we do not consider them further
here. Consistent with our previous convention of naming testis-expressed genes after American
rocketry (Gubala et al. 2017), we will from here on refer to *CG13541* as *atlas*.
184
185 *CRISPR-mediated gene mutation validates* atlas *RNAi results*
186
We validated the observed fertility defect by using CRISPR/Cas9-based genome editing
to construct putative loss-of-function alleles for *atlas* (Fig. S2). The principal allele we used for
validation and further functional studies (described below) was a null allele that completely
deleted *atlas* from the genome. This allele was generated by targeting each end of the locus
with a gRNA. We made three additional frameshift alleles by inducing double-stranded breaks
at a gRNA target site just downstream of the *atlas* start codon, which induced non-homologous
end joining. Males homozygous for the *atlas* deletion allele have the same fertility defect as
knockdown males (Fig. 2A). Males homozygous for any of three frameshift alleles showed
significantly reduced, but non-zero, fertility (Fig. 2B). It is possible that residual *atlas* function
may be present in these animals, perhaps due to translation initiation at a downstream start
codon to create a shorter protein with partial function. Finally, we constructed a genomic rescue
construct carrying both the *atlas* coding region and its native regulatory sequences. *Atlas* null
males that carried a single copy of the rescue construct had fully restored fertility (Fig. S3).
Overall, these data demonstrate that *atlas* loss, and not an RNAi or CRISPR off-target, causes
nearly complete male sterility.
202

203
204
Figure 2. CRISPR-generated deletion and frameshift alleles of *atlas* confirm the gene's
requirement for male fertility. A) Single-pair fertility assay for males homozygous for the null (Δ*atlas*)
allele or heterozygous controls (Δ*atlas*/SM6). Flies homozygous for the deletion had significantly reduced
fertility (control fertility: 82.9 ± 4.5; null fertility: 0.3 ± 0.6; two-sample t-test assuming unequal variances, *p*
= 5.4 x 10^{-18}). B) Single-pair fertility assays for males homozygous or heterozygous for three frameshift
alleles of *atlas* generated by imprecise non-homologous end joining at a CRSIRP/Cas9 target site just
downstream of the start codon: *atlas*^{52} (control fertility: 104.7 ± 3.8, mutant fertility: 11.2 ± 4.6; two-sample
t-test assuming unequal variances: *p* = 8 x 10^{-12}), *atlas*^{62} (control fertility: 96.2 ± 5.7; mutant fertility: 8.4 ±
2.9; two-sample *t*-test assuming unequal variances: *p* = 6.1 x 10^{-10}) and *atlas*^{86} (control fertility: 67.5 ± 6.6;
mutant fertility: 17.3 ± 5.8; two-sample *t*-test assuming unequal variances: *p* = 9.5 x 10^{-6}).
215

216   Atlas *is required for proper spermatid nuclear condensation*
217
We next examined how *atlas* loss of function impacted male fertility at the cellular level.
Dissection and phase-contrast imaging of *atlas* deletion null or knockdown male reproductive
tracts revealed that while the pre-meiotic and meiotic stages of spermatogenesis appeared
normal, sperm accumulated at the basal end of the testes, rather than in the seminal vesicles
(SVs), over the first week of adulthood (Fig. 3A and Fig. S4). To further characterize the fertility
defects in the absence of *atlas*, we examined the Mst35Bb-GFP ("protamine"-GFP) marker in
null or knockdown backgrounds (Manier et al. 2010). Mst35Bb encodes one of two protamine-
like proteins (highly similar paralogs of each other) that bind DNA in mature sperm. Its GFP
fusion construct thus allows visualization of nuclei in late stage spermiogenesis and mature
sperm. Consistent with the observed conglomeration of sperm tails at the basal testes, SVs
from either *atlas* null or knockdown males contain fewer mature sperm (Fig. 3B, S4C). The
nuclei of sperm from null males also appeared wider and less elongated than those of controls.
Together, these data suggest that *atlas* is required after meiosis, as developing spermatids take
on their final structures.
We next examined two post-meiotic processes: individualization of 64-cell spermatid
cysts into mature sperm, and spermatid nuclear condensation. Individualization initiates when
an actin-rich individualization complex (IC) associates with the bundle of spermatid nuclei. The
IC then proceeds down the sperm tails, expelling cytoplasmic waste and remodeling cell
membranes to form 64 individual sperm. We visualized this process in males 0-1 days old,
when spermatogenesis occurs at high levels, by staining whole mount testes for actin (Fig. 3C-
D). Although ICs could associate with nuclear bundles present at the basal end of the testes in
both control and *atlas* null males, we observed significantly fewer nuclear bundle-associated ICs
in nulls (Fig. 3C-D). While control testes typically had several ICs progressing down sperm tails,
we saw a significantly reduced proportion of progressed bundles in nulls (Fig. 3C-D). In some
null testes, we also observed individual investment cones dissociated away from progressing
ICs (Fig. 3C).
The ability of ICs to assemble at nuclear bundles and progress down sperm tails may be
reduced if nuclear condensation is aberrant (reviewed in Steinhauer 2015). During *Drosophila*
spermiogenesis, round spermatid nuclei undergo a series of stepwise, morphological changes
that are the product of two distinct, but related processes: changes in the chromatin packaging
of DNA, and changes in nuclear shape (Rathke et al. 2007; Fabian and Brill 2012; Rathke et al.
2014). The end result is thin, condensed nuclei. We quantified this process in testes dissected
from newly eclosed wild-type and *atlas* null males expressing Mst35Bb-GFP, which marks the
final stages of condensation. We shredded the post-meiotic region of the testes in the presence
of a fixative and counted the number of nuclear bundles that exhibited each of five stages of
condensation (Rathke et al. 2007): round nuclei, early canoe-stage (unmarked with Mst35Bb-
GFP), late canoe-stage (marked with Mst35Bb-GFP), elongated nuclei, and fully condensed
nuclei (Fig. 4). Condensation of the nuclear bundles in *atlas* null testes progressed at similar
rates to controls through the late canoe stage (Table S1). However, in *atlas* null males, all
nuclear bundles that progressed past the canoe stage (which included ~60% [range: 26-100%]
of all observed bundles) showed an aberrant "curled" phenotype (Fig. 4; Table S1). These data
suggest that Atlas protein is required during the later stages of nuclear condensation and are
consistent with the idea that the loss of *atlas* affects nuclear shape in a way that reduces IC
assembly and sperm individualization (see Fig. 3C).
262

264
Figure 3. Cytological investigations of the *atlas* mutant fertility defect. A) Phase contrast
microscopy of male reproductive tracts dissected from 7-day-old, unmated control (w^{1118}) or *atlas* null
males. Control males show the expected accumulation of sperm in the seminal vesicle (SV), which
appears here as a darker brown shading, while null male have an aberrant accumulation of sperm tails at
the basal end of the testis (T). B) Visualization of Mst35Bb-GFP in 4-day old control and *atlas* null testes.
While Mst35Bb is expressed in spermatid nuclei in the absence of *atlas*, the nuclei appear shorter and
much less numerous in the outlined SV. C) Representative images from phalloidin staining of w^{1118} and
atlas null testes used to assess the association of individualization complexes (ICs) with nuclear bundles
and the progression of ICs down the length of sperm tails. D) At top, number of nuclear bundles with ICs
associated in control and *atlas* null testes. Significantly more ICs were observed in control testes (control:
N = 17, median = 14; mutant: N = 13, median = 7; Wilcoxon rank-sum test W = 34, p = 0.0014). At
bottom, proportion of all observed ICs that were intact and that had progressed away from nuclear
bundles. Three mutant testes with no observed ICs were excluded from the analysis. A significantly
higher proportion of ICs progressed in control testes (control: N = 17, median = 0.27; control: N = 10,
median = 0; Wilcoxon rank-sum test W = 28, p = 0.0038).

280
281

282
Figure 4. *Atlas* null males show aberrant nuclear shaping at and beyond the elongated stage of
spermatid nuclear condensation. Early and late canoe stages were distinguished by the absence or
presence of Mst35Bb-GFP, respectively. Elongated and condensed stages were distinguished by the
presence or absence of GFP-positive puncta, respectively. As shown in Table S1, nuclear bundles from
atlas null testes consistently took on a curved shape after the canoe stage, though the degree of
curvature was variable, as exemplified above.

289
That condensing spermatid nuclei are misshapen in the absence of *atlas* suggests the
possibility that Atlas protein is critical for nuclear condensation. This idea is further supported
by its predicted biochemical properties. Previously characterized spermatid chromatin binding
proteins are small and highly basic (Jayaramaiah Raja and Renkawitz-Pohl 2005; Rathke et al.
2007; Kanippayoor et al. 2013), as the excess of positively charged amino acid side chains
facilitates ionic interactions with negatively charged DNA. Many such proteins (i.e., Tpl94D,
Mst35Ba, Mst35Bb, Prtl99C and Mst77F) also contain a conserved protein domain, the high-
mobility-group box (HMG-box) domain (Dorus et al. 2008; Alvi et al. 2013; Rathke et al. 2014;
Eren-Ghiani et al. 2015; Gärtner et al. 2015; Alvi et al. 2016; Gärtner et al. 2019), suggesting
that this variety of chromatin binding proteins could have originated through gene duplication
and divergence. Consistent with its putative *de novo* origin, Atlas lacks a detectable HMG-box
domain. However, Atlas is otherwise similar to these other sperm chromatin binding proteins:
the ~20 kDa protein has a highly basic predicted isoelectric point of 10.7, and its primary
sequence contains the sequence KRDK, which matches the canonical consensus sequence for
nuclear import, K(K/R)X(K/R) (Lange et al. 2007). To test the hypothesis that Atlas is nuclear
localized, and could thus bind DNA, we generated an *atlas-GFP* transgene under UAS control

and expressed it ubiquitously using *tubulin*-GAL4 and in the early male germline using *Bam*-
GAL4. In both larval salivary glands and early male germline cells, atlas-GFP appeared to be
nuclear localized (Fig. S5).
While these results were consistent with Atlas protein localizing to the nucleus, they did
not allow us to visualize Atlas in the cells in which it is normally expressed. To do so, we used
CRISPR/Cas9-induced homology directed repair (https://flycrispr.org/scarless-gene-editing/)
(Bruckner et al. 2017; Bier et al. 2018; Hill et al. 2019) to create an *atlas*-GFP fusion at the
endogenous *atlas* locus (see Fig. S6 and Materials and Methods). We first confirmed the
functionality of the knock-in allele by showing that males with the *atlas* locus genotype *atlas*-
GFP/Δ*atlas* had equivalent fertility to males of genotype *atlas*+/Δ*atlas* (Fig. 5A). We then
visualized Atlas-GFP fusion protein in whole-mount testes in conjunction with phalloidin-stained
actin (Fig. 5C). Atlas-GFP was absent from seminal vesicles, consistent with its absence from
the proteome of mature *D. melanogaster* sperm (Dorus et al. 2006; Wasbrough et al. 2010).
Instead, Atlas-GFP colocalized with condensing nuclear bundles near the basal end of the
testes (Fig. 5C). Actin-based ICs were also observed in the basal testes, but generally did not
co-localize with Atlas-GFP, suggesting that Atlas-GFP is present in condensing nuclei before IC
association (Fig. 5C). This result, taken together with the aberrant nuclear condensation in the
absence of *atlas* (Fig. 4), is consistent with the idea that Atlas is a transition protein. Transition
proteins are chromatin components that act transiently during spermatid nuclear condensation.
A series of transition proteins first replace histones as the primary DNA binding proteins in the
nucleus and then give way to protamines, the proteins that package chromatin in mature sperm
(Rathke et al. 2007; Rathke et al. 2014; Gärtner et al. 2015).
To further elucidate the role of *atlas* in nuclear condensation, we next examined Atlas-
GFP localization in the presence of either an early spermatid nuclear marker, histone H2Av-
RFP (Schuh et al. 2007; Rathke et al. 2014), or Mst35Bb-dsRed (Manier et al. 2010), a marker
of nuclei from the late canoe stage through final condensation. Atlas-GFP showed no co-
localization with H2Av-RFP, suggesting that Atlas functions after histone removal (Fig. 5B). In
contrast, some GFP-positive bundles co-localized with Mst35Bb-dsRed, but others did not (Fig.
5D). These data suggest that Atlas may be one of the final transition proteins used in nuclear
condensation before the chromatin becomes fully condensed with protamines.
336

337
338
Figure 5. An *atlas*-GFP allele generated at the endogenous *atlas* locus rescues the fertility defect
of *atlas* null flies and encodes a protein that localizes to condensing spermatid nuclei. A) A single
copy of the *atlas*-GFP allele completely rescues the fertility defect caused by the Δ*atlas* allele and shows
equivalent fertility to males heterozygous for the wild-type *atlas* allele (Δ*atlas*/*atlas*-GFP fertility: 86.0 ±
3.2; Δ*atlas*/+ fertility: 84.9 ± 4.1; two-sample t-test assuming unequal variances, *p* = 0.85). B) Atlas-GFP
does not co-localize with histone H2Av-RFP, a marker of the initial stages of spermatid nuclear
condensation. C) Visualization of Atlas-GFP in whole-mount testes from *atlas-GFP* homozygotes shows
that the fusion protein co-localizes with a subset of condensing spermatid nuclear bundles. While actin
associates with fully condensed nuclei at the basal testis, Atlas-GFP does not overlap and is also absent
from the seminal vesicle. Some Atlas-GFP is observed near progressing individualization complexes
toward the apical testis in the merged image (marked with asterisks; see also Fig. 6D). D) Atlas-GFP
partially colocalizes with Mst35Bb-dsRed, a marker of the final stage of nuclear condensation. Open
arrow: example of co-localization. Filled arrowhead: example of Atlas-GFP that does not co-localize with
Mst35Bb-dsRed. Collectively, these data suggest that atlas may serve as a transition protein involved in
the final stages of nuclear condensation.
354
To determine the stage(s) of nuclear condensation at which *atlas* functions, we analyzed
the shape of fixed nuclear bundles from shredded testes isolated from *atlas*-GFP males on the
day of eclosion. Based on the stage of the defect in *atlas* null males (Fig. 3-4) and the pattern
of Atlas-GFP-positive bundles in whole-mount testes (Fig. 5), we hypothesized that Atlas-GFP
would localize to the later stages of nuclear condensation. Consistent with this hypothesis, we

did not detect Atlas-GFP in round or early canoe stage bundles (Fig. 6A-B). Atlas-GFP co-
localized with DNA in late canoe stage bundles (Fig. 6C). Interestingly, when nuclei elongated
further, GFP was detected not in the nucleus, but as puncta basal to the nuclei (Fig. 6D; see
also Fig. 5C). Since Atlas-GFP is not observed in mature sperm in the SV (Fig. 5C), these data
suggest that Atlas may function as a transition protein that facilitates the condensation of
spermatid nuclei from histone-based DNA packaging to protamine-like-based DNA packaging
(Rathke et al. 2007) and is then removed from nuclei once protamines bind DNA. Indeed, the
appearance of Atlas in nuclei during the late canoe stage of condensation is similar to the
pattern observed for a previously characterized transition protein, Tpl94D (Rathke et al. 2007).
We hypothesize that the failure of *atlas* null sperm to form needle-like nuclei can be explained
by the absence of Atlas from the late canoe nucleus. It is also possible that the apparent
removal of Atlas-GFP from nuclei (Fig. 6D) represents a mechanism for removing transition
proteins from the nucleus after they exert their functions. We observed above that some
Mst35Bb-GFP also appears to be removed from the nucleus in puncta during the elongation
stage of nuclear condensation (see elongated stage of control nuclear bundles, Fig. 4), even
though other Mst35Bb-GFP molecules ultimately package DNA in mature, individualized sperm.
This could occur if Mst35Bb-GFP is present in excess of what is needed to package DNA.
377

378
379
380   **Figure 6. Atlas-GFP is present in late canoe-stage spermatid nuclei and then appears to leave the**
381   **nucleus in puncta.** Staging of condensing spermatid nuclei fixed in paraformaldehyde from *atlas*-GFP
382   males stained with TO-PRO-3 DNA stain. Atlas-GFP is not detectable in (A) round stage or (B) early
383   canoe stage nuclei. Atlas-GFP is nuclear localized in the late canoe stage (C). When nuclei become fully
384   elongated (D), puncta of Atlas-GFP appear to be removed from the nucleus.

385
386

387   *Evolutionary origins of* atlas
388
To better understand the evolutionary origin of *atlas* and its evolution since emergence,
we used a combination of BLAST- and synteny-based approaches to identify *atlas* orthologs
throughout the genus (Gubala et al. 2017; Rele et al. 2020). One notable feature of this two-
exon gene is that the protein-coding region (519 nucleotides) is contained entirely within the first
exon (622 nt); the longer, second exon (910 nt) appears to be entirely non-coding (Fig. 7).
Surprisingly, the second exon is more widely conserved. BLASTN detected significant matches
to this region (range of hit length: 185-864 nt) in the same genomic location on Muller element C
(see Schaeffer et al. 2008 for explanation of Muller elements), as assessed by synteny, in all
Drosophila species examined, including distantly related species such as *D. virilis* and *D.
grimshawi* (Fig. 7 and S7). The protein-coding first exon showed a more limited phylogenetic
distribution. In most members of the *melanogaster* group of *Drosophila* (gray box in Fig. 7), this
exon is found in a conserved position, adjacent to the non-coding region on the equivalent of *D.
melanogaster* chromosome 2R (Table S2). In *D. ananassae*, however, the protein-coding
region is found on the X chromosome (Muller element A). A putative ortholog for the protein-
coding sequence is detectable by BLASTP in a partially syntenic region on the same Muller
element in *D. virilis* (Table S2, Fig. S8). These data suggest that the *atlas* protein-coding
sequence initially arose on Muller element A and then moved to Muller element C, giving rise to
the gene structure observed in extant *D. melanogaster* and its sister species.
407

408
Figure 7. Molecular evolution and gene expression of *atlas* across the *Drosophila* genus. The
gene structure of *atlas* in D. melanogaster is shown at top left. The predicted protein-coding sequence is
contained entirely within exon 1, while exon 2 encodes the presumed 3' UTR. The gene is located on
chromosome 2R, equivalent to Muller element C. The phylogeny shows BLAST- and synteny-based
detection of sequences orthologous to the protein-coding sequence and the 3' UTR sequence across
Drosophila species. Sex-specific adult RNA-seq data were used to assess male expression across
species, with RT-PCR verification performed in species marked with asterisks. RNA-seq data for the
syntenic region of the 3' UTR in *D. virilis* were ambiguous; see Figs. S7 and S9.
417
To confirm the lack of *atlas* protein-coding sequences identifiable by BLASTP or
TBLASTN in most non-*melanogaster* group species, we identified the regions syntenic to those
containing *atlas* in *D. ananassae* and *D. virilis* in 11 additional *Drosophila* species and used

more sensitive methods to search for potential orthologs (Rele et al. 2020). Specifically, we: a)
relaxed the BLAST cut-offs for detection, since default parameters can cause false-negative
results when searching for potential *de novo* genes in divergent species (Weisman et al. 2020);
b) used adult male RNA-seq data to detect transcribed areas within each syntenic region that
did not match annotated genes; and, c) predicted the isoelectric point of the potential proteins
encoded, under the hypothesis that Atlas orthologs would have conserved, DNA-binding
functions. The results are summarized in Fig. S8 and Table S2. These searches detected no
evidence for *atlas* orthologs in the following *Drosophila* species: *obscura*, *miranda, willistoni,*
hydei, arizonae, mojavensis, navojoa, and *grimshawi*. Some of these species had unannotated,
male-expressed transcripts in the regions syntenic to the Muller A location of *atlas* in *D.*
ananassae and *D. virilis*, but when each was compared with BLASTP to the *D. melanogaster*
proteome, all matched proteins other than Atlas, suggesting they may be lineage-specific
paralogs of other genes (Fig. S8). In sister species *D. pseudoobscura* and *D. persimilis*, we
detected a male-expressed transcript predicted to encode a protein with a pI > 10 in the region
syntenic to the location of *atlas* in *D. virilis*, but the predicted protein sequences showed no
significant BLASTP similarity to *atlas* orthologs (Table S2). While this predicted protein may
represent a divergent *atlas* ortholog, the abSENSE method predicts low probabilities of BLASTP
detection failure when searching for Atlas protein in these species (0.02 and 0.04, respectively),
so we favor the hypothesis of a lineage-specific, newly evolved gene in the region. Conversely,
in *D. busckii*, we detected, in the region syntenic to the *D. virilis atlas* locus, a male-expressed
transcript predicted to encode a protein with significant BLASTP identity to *D. melanogaster*
Atlas (e = 4e-8), but with a predicted pI of 5.1 and a ~50 percent shorter open-reading frame
(Table S2). The ortholog status of this predicted protein is also unclear, but because of its
dramatically altered size and pI, it is unlikely to have a functional role equivalent to that of *D.*
melanogaster Atlas.
To investigate whether the protein-coding region may have reproductive functions in
other species, we used sex-specific RNA-seq data from numerous *Drosophila* species curated
by the Genomics Education Partnership (Rele et al. 2020; thegep.org) and verified several of
these results by RT-PCR (Fig. 7, Fig. S9). In all species in which *atlas* was detected, the
protein-coding region was expressed specifically in males regardless of its genomic location
(Fig. 7). Interestingly, the non-coding region showed male-specific expression in species
lacking an unambiguous, orthologous coding region, such as *D. pseudoobscura* and *D.*
mojavensis. Conversely, while *D. yakuba* and *D. erecta* express the protein-coding region
robustly, we found no RNA-seq evidence to support expression of the non-coding second exon,
in spite of its sequence conservation (Fig. S7). Based on its high level of sequence
conservation, consistent genomic location and expression in a variety of species, it is possible
that what we now consider to be the 3' untranslated region of *atlas* from *D. melanogaster* was,
ancestrally, a non-coding RNA.
The FlyBase database reports two transcript isoforms of *atlas* in *D. melanogaster*: the
atlas-RA isoform is 986 nucleotides, while the *atlas*-RB isoform is 1528 nt. These isoforms
differ in how much of the second, non-coding exon is included in the transcript. We used RT-
PCR of whole male cDNA to assess the presence of these isoforms and their relative
abundances. Primers designed to amplify a region present in both isoforms produced products
that appeared more abundant than primers designed to amplify only the long isoform, even
though both primer pairs appeared to amplify genomic DNA with equal efficiency. Based on
RT-PCR band intensities and controlling for product size and genomic PCR band intensities, we
estimated that the short isoform is about 3-fold more abundant. This difference in abundance is
mirrored in available RNA-seq data, which show approximately 3- to 4-fold higher levels of
expression in the upstream part of exon 2 (Fig. S7), a pattern that also appears in *D. simulans*
and *D. sechellia*. Evaluating the potential significance of this finding awaits functional
characterization of the non-coding region.

As we have observed for other putative *de novo* genes with essential male reproductive
functions (Gubala et al. 2017), the pattern of *atlas* protein-coding sequence presence/absence
across the phylogeny is difficult to explain parsimoniously. If we assume that gene birth events
are less frequent than gene deaths, since the latter can occur through many possible mutational
events and can happen separately along multiple phylogenetic lineages, our data support the
hypothesis of a single origin of the protein-coding sequence at the base of the genus, followed
by independent loss events on the lineages leading to *D. grimshawi*, *D. mojavensis* and *D.*
willistoni, and potentially also *D. pseudoobscura/persimilis* (Table S2). We summarize these
findings for 12 representative species of *Drosophila* in Fig. 7. The general patterns of loss do
not change when all species of Table S2 are considered, though an additional loss in the
melanogaster group is likely due to the absence of a detectable ortholog in *D. kikkiwai* and *D.*
serrata. As noted above, the pattern of gene loss can also appear due to orthology detection
failure (Weisman et al. 2020), for which we tried to account with our additional search methods
described above. We also note, however, that the probability of BLASTP-based ortholog
detection failure is relatively low for some *Drosophila* species that lack *atlas*, including *D.*
pseudoobscura (probability of non-detection due to divergence = 0.02), *D. persimilis* (p = 0.04)
and *D. willistoni* (p = 0.06). The probability is higher for other species, *D. mojavensis* (p = 0.33)
and *D. grimshawi* (p = 0.66), underscoring the importance of our additional search strategies.
Overall, our data support the hypothesis of multiple, independent loss events within *Drosophila*.
AbSENSE produces a 1.00 probability of BLASTP-based atlas ortholog detection failure
outside of *Drosophila*, reflecting the protein's short length and relatively rapid divergence (see
below). Indeed, the protein-coding and non-coding transcriptomes from each species showed
no matches to Atlas protein or cDNA sequences by BLAST. We thus used another synteny-
based approach, summarized in Fig. S10, to look for the protein-coding gene in other Dipterans
with well-resolved genomes: *Musca domestica*, *Glossina morsitans*, *Lucilia cuprina*, *Aedes*
aegypti, *Anopheles darlingi*, *Anopheles gambiae*, *Culex quinquefasciatus* and *Mayetiola*
destructor. In none of these species was a putative homolog found in any potential syntenic
region.
Recognizing the limitation of even this approach, we also used HMMER (Potter et al.
2018) to search iteratively either all genomes in ENSEMBL, or all metazoan genomes in
ENSEMBL, for annotated proteins with identity to Atlas from *D. melanogaster* or *D. virilis*.
These searches initially identified significant hits to the Atlas orthologs we identified above from
other *Drosophila* species. When these collections of orthologs were used as queries, no further
proteins outside of *Drosophila* were a significant match. As a control, we performed the same
search strategy with *D. melanogaster* Mst35Bb, a protein whose length, amino acid
composition, and function are similar to Atlas. These searches readily identified orthologs
throughout Diptera, consistent with predictions of its conservation from the OrthoDB database
(Zdobnov et al. 2017). Thus, we conclude that *atlas* is a putative *de novo* evolved gene that is
limited phylogenetically to the *Drosophila* genus.
Finally, we used standard tests of molecular evolution to examine the selective
pressures that have shaped Atlas protein within the *melanogaster* group. We aligned the *atlas*
protein-coding sequences from 12 species and used PAML to ask whether a model (M8_
allowing for positive selection, as well as neutral evolution and purifying selection, explained the
data better than models (M7 and M8a) that allowed only neutral evolution and purifying
selection (Yang 2007; McGeary and Findlay 2020). These data showed that while the *atlas*
protein-coding sequence's rate of evolution was accelerated relative to most *Drosophila* proteins
(whole-gene estimated d_N/d_S, ω = 0.41 by PAML model M0), there was no significant evidence
for positive selection acting to recurrently diversify a subset of sites within the protein (Table 2).
520
521

522 **Table 2. PAML sites tests for positive selection acting on atlas in the *melanogaster***
523 **group.**
524
Model M0 (uniform w across all sites): ω = 0.41, ln L = -4032.24, np = 23
Model M7 (10 site classes, each with $0 \leq \omega \leq 1$): ln L = -3961.28, np = 24
Model M8 (10 site classes as in M7, plus one class with $\omega \geq 1$): ln L = -3960.41, np = 26, ω for
extra class of sites = 1.39 (9.1% of sites)
Model M8a (10 site classes as in M7, plus one class with ω = 1): ln L = -3961.02, np = 25,
16.6% of sites in the ω = 1 class
531
532 M7 vs. M8 likelihood ratio test: χ^2 = 1.74, df = 2, *p* = 0.42
533 M8 vs. M8a likelihood ratio test: χ^2 = 1.22, df = 1, *p* = 0.27
534
535
536

537   **Discussion**
538
Across taxa, many *de novo* evolved genes are expressed in the male reproductive
system (Levine et al. 2006; Begun et al. 2007; Cui et al. 2015; Ruiz-Orera et al. 2016).
Identifying those genes that have evolved essential roles in reproduction will provide insight into
how newly evolved genes integrate with existing cellular networks (Abrusán 2013) and how
evolutionary novelties permit adaptation in the face of sexual selection. Here, we screened 42
putatively *de novo* evolved genes for major effects on male *D. melanogaster* reproduction. Our
primary screen identified three genes whose knockdown caused an apparent reduction in male
fertility. However, subsequent CRISPR-mediated gene deletion revealed that only one of these
genes, *atlas*, was truly essential. This result underscores the importance of validating genes
identified in RNAi screens through traditional loss-of-function genetics and other approaches.
Using such genetic tools, we then showed that *atlas* loss of function reduces fertility by
affecting mature sperm production. During spermiogenesis, *atlas* mutants show aberrant
nuclear condensation and an inability to individualize spermatid bundles successfully. GFP-
tagged Atlas protein localizes to condensing spermatid nuclei in the basal testis and partially co-
localizes with Mst35Bb, a protamine around which DNA is wrapped in mature, individualized
sperm. Evolutionary analysis showed that the *atlas* protein-coding sequence likely arose at the
base of the *Drosophila* genus but was unlikely to have played an essential role immediately
upon birth, as the gene was subsequently lost along several independent lineages. Within the
melanogaster group of *Drosophila*, however, the gene moved from the X chromosome to an
autosome, where it formed a single transcriptional unit with a conserved, non-coding sequence.
Since this point, the gene has encoded a protein with a conserved length, isoelectric point and
male-specific expression pattern, suggesting potential functional conversation over the last ~15
million years.
Several lines of evidence suggest that Atlas is a transition protein that facilitates the
change from histone-based to protamine-based chromatin packaging in spermatid nuclei. Atlas
localizes throughout spermatid nuclei (Fig. 6) and has biochemical properties consistent with
direct DNA interaction. The protein appears specifically at the late canoe stage of nuclear
compaction (Fig. 5B-D). Its lack of overlap with testis-specific histones (Fig. 5C), partial overlap
with Mst35Bb (Fig. 5D), likely removal from needle-stage nuclei (Fig. 6) and absence from
mature sperm (Figs. 5-6) are all consistent with the expression profile of a transition protein.
Several other transition proteins have been characterized in *D. melanogaster*, including Tpl94D,
thmg-1, thmg-2, and Mst84B (Rathke et al. 2007; Gärtner et al. 2015; Gärtner et al. 2019).
Collectively, the transition proteins vary in the stage of nuclear condensation at which they first
appear and the range of nuclear shapes over which they are found (Hundertmark et al. 2018),
but otherwise match Atlas in their biochemical properties, transient roles, and localization
throughout the nucleus. Compared to these other transition proteins, Atlas is present over a
fairly narrow range of nuclear condensation stages and reaches its peak expression just prior to
the onset of individualization. *Atlas* is also the only transition protein gene characterized to date
whose removal disrupts fertility, as *Tpl94D*, *thmg-1*, *thmg-2* and *Mst84B* mutants are all fertile
(Rathke et al. 2007; Gärtner et al. 2015; Gärtner et al. 2019). This may reflect the relatively
later timing of Atlas's expression in spermatid nuclei, reduced functional redundancy between
DNA-binding proteins at the later stages of condensation, a potential interaction between Atlas
and an essential protamine-like protein, and/or a more stringent requirement for DNA binding at
these stages.
Transition proteins give way in spermatid nuclei to protamine-like proteins, which bind
DNA in mature sperm and persist through fertilization. In this way, protamine-like proteins
function analogously to vertebrate protamines, though they are believed to be evolutionarily
independent (Jayaramaiah Raja and Renkawitz-Pohl 2005; Doyen et al. 2015). In *D.*
melanogaster, protamine-like proteins include Mst35Ba, Mst35Bb, Prtl99C and Mst77F (Rathke

et al. 2010; Eren-Ghiani et al. 2015; Kimura and Loppin 2016). Interestingly, while all
characterized protamine-like proteins are present in mature sperm, only some are essential for
fertility. Knockouts of *Mst35Ba*, *Mst35Bb*, or both show occasional nuclear shaping defects, but
male fertility is normal (Rathke et al. 2010; Tirmarche et al. 2014). In contrast, mutants of
Prtl99C or *Mst77F* are sterile. Prtl99C and Mst35Ba/b bind condensed DNA independently of
each other and contribute additively to the shortening of needle-stage nuclei, but Prtl99C's
effect is ~3x greater (Doyen et al. 2015; Eren-Ghiani et al. 2015). This difference is apparently
great enough to reduce fertility only in *Prtl99C* mutants. In contrast, *Mst77F* and *Mst35Ba/b*
show a genetic interaction, as *Mst35Ba/b* null flies become nearly sterile in an *Mst77F*
heterozygous background (Kimura and Loppin 2016). Furthermore, while Mst35Bb-GFP is
expressed in *Mst77F* nulls, these flies show deformed spermatid nuclei that do not reach a
recognizable needle-like stage. Because *atlas* nulls show considerable phenotypic similarity to
Mst77F nulls, but not *Prtl99C* nulls, we hypothesize that Atlas may act in a pathway with
Mst77F. Our observation of inefficient IC movement down sperm tails in *atlas* null testes is
reminiscent of a similar phenotype in *Mst77F* nulls (Kimura and Loppin 2016), providing further
evidence that these proteins may act in a common pathway. In both cases, ICs can form at
misshapen canoe-stage nuclei, but fail at a subsequent step. While the exact relationship
between nuclear abnormalities in the late canoe stage and individualization is not entirely
understood, it is possible that nuclear shape and the organization of nuclear bundles impact the
ability of IC association and IC progression, as is also observed in mutants of another gene,
dPSMG1, which controls nuclear shape (Gärtner et al. 2019).
Because *de novo* genes emerge from non-coding sequences, they typically encode
proteins that are short and lack complex structure (Schlötterer 2015; Van Oss and Carvunis
2019). Indeed, expanding the length of the protein-coding region and evolving higher-level
protein structures are hypothesized to be among the final stages of new gene evolution
(Bornberg-Bauer and Schmitz 2017). In light of these constraints, what kinds of cellular
functions might be available to newly evolved proteins? Vakirlis et al. (2020) overexpressed
emerging proto-genes in *S. cerevisiae* and found that those encoding proteins with predicted
transmembrane (TM) domains were more likely to be adaptive, as assessed by the effect of
proto-gene overexpression on growth rate. Such proteins may arise when thymine-rich
intergenic regions undergo mutations that allow protein-coding gene birth and expression, since
many codons with multiple U nucleotides encode amino acids commonly found in TM domains
(Vakirlis et al. 2020). Our imaging data, in addition to the prediction tools employed by Vakirlis
et al. (2020), suggest that Atlas does not contain a TM domain. However, just as the amino
acid compositional requirements of a TM domain are not overly complex, neither are those of
DNA binding proteins. In essence, these proteins must simply be small, have a high
concentration of positively charged residues, and contain a nuclear localization signal, which
itself requires a small patch of positively charged residues (Lange et al. 2007). Thus, DNA
binding proteins may be a relatively easy class of protein to evolve *de novo*.
While many putative *de novo* genes are expressed in the *D. melanogaster* testis (Fig. 1A
and Heames et al. 2020), *atlas* was the only verified hit from our screen that was essential for
male fertility. This result raises two related questions. First, why has selection maintained the
expression of the other potential *de novo* genes in our screen? In general, it is common for the
knockdown of protein-coding genes expressed in reproductive tissues in *D. melanogaster* to
result in no detectable fertility defects (Ravi Ram and Wolfner 2007; Schnakenberg et al. 2011;
Findlay et al. 2014; Gubala et al. 2017). One hypothesis to explain this pattern is that while the
loss of function of such genes may cause small reductions in fertility that would be subject to
strong negative selection in nature, the conditions used to assay such knockdown animals in the
primary screens are rarely tailored to detect differences of this magnitude. Another possibility,
not mutually exclusive, is that while the genes may be expendable in non-competitive, non-
exhaustive mating conditions, their absence may result in lower fitness in sperm competitive

environment, environments in which males mate several times in quick succession, or
environments in which sperm must persist in storage for longer intervals or during less optimal
conditions (Wong et al. 2008; Yeh et al. 2012; Civetta and Finn 2014).
A second question raised by our finding that *atlas* encodes an essential transition protein
is: how might *atlas* have evolved to become essential for fertility in *D. melanogaster*, particularly
when other transition proteins appear functionally redundant? Other proteins involved in
spermatid chromatin compaction show variable levels of conservation across *Drosophila*. For
example, the protamines around which DNA is wrapped in mature sperm (Jayaramaiah Raja
and Renkawitz-Pohl 2005) are found across all sequenced *Drosophila* species (Alvi et al. 2013),
and orthologs are also reported in FlyBase from other Dipteran and non-Dipteran insects (Larkin
et al. 2021). However, transition protein Tpl94D is reported to be restricted to species ranging
from *D. melanogaster* to *D. pseudoobscura* (Alvi et al. 2016), as are the related proteins tHMG1
and tHMG2 with high-mobility group domains (Gärtner et al. 2015; Larkin et al. 2021). Results
like these suggest that while some protamine-like proteins (i.e., Mst35Ba and Mst35Bb) have
consistently been among the final chromatin-packaging proteins, the specific proteins facilitating
the transition from histones to protamines have likely varied over evolutionary time. Against this
backdrop, and based on our analyses of the protein's presence/absence, biochemical
properties, and expression patterns in extant species, we hypothesize that while the Atlas
protein likely had some DNA-binding ability and male-specific expression upon its origin, it was
only one of several proteins involved in spermatid chromatin compaction. Since *atlas* was lost
independently in several lineages after its birth (Fig. 7), Atlas was likely non-essential at its
outset, but rather evolved an essential function within the *melanogaster* group of species. Such
evolution of essentiality could have occurred because of the loss of a protein with a
complementary function and/or changes in the process of spermatogenesis that thrust Atlas into
a functionally unique role. It is also worth noting that species that have evidently lost *atlas* might
have undergone other compensatory changes in their repertoires of spermatid DNA binding
proteins. For example, *D. willistoni* lacks *atlas* but appears to have several additional paralogs
of the protamines found only in duplicate in *D. melanogaster*, which could have evolved
transition-protein-like roles.
While our study cannot establish whether the movement of the *atlas* protein-coding
sequence off of the X chromosome onto an autosome early in the evolution of the *melanogaster*
group (Fig. 7) affected the gene's essentiality, such movement remains noteworthy. Prior work
has found a significant dearth of testis-expressed genes on the X chromosome in *Drosophila*
(Parisi et al. 2003; Parisi et al. 2004; Dorus et al. 2006; Vibranovski, Zhang, et al. 2009) and
other species (Emerson et al. 2004; Reinke et al. 2004). Furthermore, *Drosophila* exhibit
suppression of X-linked testis-expressed genes, and transfer of such genes from the X
chromosome to autosomal loci results in higher expression levels (Kemkemer et al. 2014;
Argyridou et al. 2017; Argyridou and Parsch 2018). One of several proposed mechanisms for
both the paucity of X-linked testis-expressed genes and the suppression of their expression is
meiotic sex chromosome inactivation (MSCI), in which the X chromosome becomes
transcriptionally silenced earlier than autosomes (Vibranovski, Lopes, et al. 2009; Zhang,
Vibranovski, Krinsky, et al. 2010; Zhang, Vibranovski, Landback, et al. 2010; Vibranovski et al.
2012; Gao et al. 2014; Mahadevaraju et al. 2021). Thus, genes that affect meiotic or post-
meiotic processes, as *atlas* does, could exert beneficial effects more strongly and/or for a longer
period of time if they become encoded autosomally. While the *atlas* protein-coding sequence
appears to show male-specific expression regardless of its chromosomal location, it is possible
that the movement of *atlas* to chromosome 2 allowed it to evolve a broader or different
expression pattern that expanded or modified its role in spermiogenesis. The complex
molecular bases of both X suppression and "escape" from the X chromosome in *Drosophila*
continue to be actively investigated and debated (Meiklejohn et al. 2011; Mikhaylova and
Nurminsky 2011; Vibranovski et al. 2012; Gao et al. 2014; Vibranovski 2014; Landeen et al.

2016; Mahadevaraju et al. 2021), but continued research in this area might inform further
interrogation of the forces driving *atlas* off of the X chromosome.
The movement of the *atlas* protein-coding sequence to chromosome 2 also created the
two-exon gene observed in *D. melanogaster*, in which the longer second exon appears to be
entirely non-coding. This second exon is highly conserved across the genus in both sequence
and genomic location, and it shows male-specific expression in several species that lack the
protein-coding sequence upstream (Fig. 7 and Fig. S7). These patterns of conservation
suggest that the second exon might originally have been a non-coding RNA, a class of molecule
whose importance in *Drosophila* male reproduction has recently become recognized (Wen et al.
2016; Bouska and Bai 2021). While these previous examples of functional ncRNAs in
spermatogenesis have generally acted in *trans* to regulate other genes or affect the functions of
other proteins, it is also possible that the long 3' UTR of *atlas* in *D. melanogaster* could affect
the translation of *atlas* transcripts. Many genes functioning in spermatid differentiation are
transcribed early in spermatogenesis but translationally repressed until later in spermiogenesis,
a phenomenon that relies on various forms of post-transcriptional regulation (White-Cooper
2010; Lim et al. 2012). Future studies of the *atlas* protein-coding sequence in the absence of its
3' UTR, the expression patterns of Atlas protein in species in which it is encoded from the X
chromosome, or the genetic ablation of the conserved region in species lacking the protein-
coding sequence will provide additional insights.
A final issue raised by our results is the exact timing and mechanism of origin for the
atlas protein-coding sequence. The bioinformatic screen (Heames et al. 2020) that identified
atlas and the other genes tested in Fig. 1 was designed to identify both "*de novo*" genes,
defined as protein-coding regions in *Drosophila* that had recognizable, but non-ORF-
maintaining, TBLASTN hits in outgroup species, and "putative *de novo*" genes, which had no
TBLASTN hits in outgroup species. (Importantly, the screen also eliminated any protein with an
identifiable protein domain, thus reducing the chances of identifying divergent members of gene
families.) The vast majority of the genes we tested with RNAi, including *atlas*, fell into the
putative *de novo* category. The bioinformatic screen's criteria were reasonable for a high-
throughput analysis, but BLAST-based methods have known limitations for detecting
orthologous sequences in diverged species (Moyers and Zhang 2015, 2018; Weisman et al.
2020). The lack of identifiable *atlas* protein-coding genes in several *Drosophila* species (e.g., *D.*
pseudoobscura and *D. willistoni*) is unlikely to be due to BLAST homology detection failure, and
extensive synteny-based searches confirmed the gene's absence (Fig. S9). BLAST and
synteny-based searches for orthologs in non-*Drosophila* species also did not detect an ortholog,
though BLAST searches are not predicted to have adequate sensitivity, for a protein of this size
and evolutionary rate, at this level of species divergence (Weisman et al. 2020). Hence, in
addition to using synteny to search for orthologs, we used HMMER, which employs hidden
Markov models and builds a sequence profile of the target protein using information from
multiple orthologs. Since HMMER also did not detect orthologs outside of *Drosophila*, we
hypothesize that *atlas* evolved *de novo* at the base of the genus. However, since we remain
unable to identify the non-protein-coding sequence from which *atlas* arose, we continue to refer
to *atlas* as a putative *de novo* gene (McLysaght and Hurst 2016).
Overall, we find that while many putative *de novo* evolved genes are expressed in the *D.*
melanogaster testes, few have major, non-redundant effects on fertility. However, several such
genes have acquired critical roles, acting at distinct stages of spermatogenesis and sperm
function. We showed previously that the putative *de novo* gene *saturn* is required for maximal
sperm production, as well as for the ability of transferred sperm to migrate successfully to sperm
storage organs in females (Gubala et al. 2017). Another putative *de novo* gene, *goddard*, is
required for sperm production and encodes a cytoplasmic protein that appears to localize to
elongating axonemes (Gubala et al. 2017; Lange et al. 2021). Loss of *goddard* impairs the
individualization of spermatid bundles (Lange et al. 2021), thus exerting an effect that appears

to be upstream of those observed for *saturn* and *atlas*. Here, we report another novel function
for a putative *de novo* gene: encoding an essential transition protein that is necessary for proper
nuclear condensation in spermiogenesis. Taken together, these results demonstrate that while
many *de novo* genes may play subtle roles or share functional redundancy with other genes, *de*
novo genes can also become essential players in complex cellular processes that mediate
successful reproduction.
747

748   **Materials and Methods**
749
750   *RNA Interference Screen*
751
De novo and putative *de novo* genes inferred to be no older than the *Drosophila* genus
were identified previously (Heames et al. 2020). We filtered these genes with publicly available
RNA-seq data (Brown et al. 2014) to identify those expressed predominantly in the testes
(>50% of RPKM sum deriving from the testes from ModENCODE data; Brown et al. 2014),
giving a total of 96 genes. To assess each of these candidates for effects on male fertility, we
induced knockdown in the male germline by crossing UAS-RNAi flies to Bam-GAL4, UAS-
Dicer2 flies (White-Cooper 2012; Gubala et al. 2017). Control flies were generated by crossing
the genetic background into which UAS-RNAi was inserted crossed to the same GAL4 line.
Flies carrying UAS-RNAi were of two types. Roughly half of the genes had publicly available
lines from the Vienna *Drosophila* Resource Center (Dietzl et al. 2007) or the Transgenic RNAi
Project (Ni et al. 2011). For the other genes, no publicly available RNAi stock was available, so
we constructed TRiP-style stocks in the pValium20 vector as previously described (Findlay et al.
2014). These constructs were integrated into an AttP site in stock BL 25709 (y^1 v^1 P{nos-
phiC31\int.NLS}X; P{CaryP}attP40) from the Bloomington *Drosophila* Stock Center (injections
by Genetivision; Houston, TX, USA) and crossed into a *y v* background to screen for v^+. We
attempted at least two rounds of transgenic production for each gene. In total, we were able to
obtain and test RNAi lines for 57 of the 96 identified genes.
We initially screened males knocked down for each candidate gene for major fertility
defects by crossing groups of 7 knockdown or control males to 5 virgin Canton S females,
letting the adults lay eggs for ~48 hours, and then discarding adults and quantifying the resulting
progeny by counting the pupal cases, as previously described (Gubala et al. 2017). To assess
the degree of knockdown achieved, 10 whole males of each line were homogenized in TRIzol
reagent (Life Technologies, Carlsbad, CA). RNA isolation, DNAse treatment, cDNA synthesis
and semi-quantitative RT-PCR with gene-specific primers were performed as previously
described; amplification of *RpL32* was used as a positive control (Gubala et al. 2017). Any
gene that did not show at least partial knockdown was discarded from further analysis, leaving a
total of 42 genes successfully screened. Table S3 lists all lines used and results of tests for
effective target gene knockdown.
780
781   *CRISPR Genome Editing*
782
To validate RNAi results for *atlas*, *CG43072* and *CG33284*, we used CRISPR/Cas9
genome-editing to generate null alleles that could be used for further analysis, as described
previously (Lange et al. 2021). Briefly, our general strategy was to design gRNAs in the pU6.3
vector (Drosophila Genome Resource Center (DGRC) #1362) that targeted each end of a locus.
These plasmids, along with plasmids encoding gRNAs that targeted the w+ locus, were co-
injected by Rainbow Transgenics (Camarillo, CA) into embryos laid by *vasa*-Cas9 females in a
w+ background, Bloomington stock #51323 (Ge et al. 2016). G_0 animals were crossed to *w-*
flies, and members of G_1 broods with a higher-than-expected fraction of *w-* progeny were
individually crossed to an appropriate balancer line and then PCR-screened for the desired
deletion of the targeted locus.
We also constructed three frameshift, expected loss-of-function alleles for *atlas* by using
CRISPR to induce non-homologous end joining at a single PAM site just downstream of the
atlas start codon. *Vasa*-Cas9 embryos were co-injected and screened for w- progeny as
described above. We then used squish preps to isolate DNA from G1 flies and used a PCR-
RFLP assay to detect mutations. PCR products spanning the gRNA-targeted site were digested
with *Bfa*I (New England Biolabs (NEB), Ipswich, MA); undigested products in which the

expected *Bfa*I site was lost indicated a mutation, which was balanced and then confirmed by
PCR and sequencing of homozygous mutant lines.
We used scarless CRISPR editing and homology-directed repair (HDR) to insert the
GFP protein-coding sequence in-frame at the end of the *atlas* protein sequence (see Fig. S6;
(https://flycrispr.org/scarless-gene-editing/) (Bruckner et al. 2017; Bier et al. 2018; Hill et al.
2019). We first generated an *atlas*-GFP DNA construct by cloning the *atlas* protein-coding
sequence into pENTR and using LR Clonase II (Thermo Fisher Scientific, Waltham, MA) to
recombine the sequence with pTWG (DGRC #1076; T. Murphy), generating a C-terminally
tagged *atlas*:GFP construct. We amplified the *atlas* fragment from *vasa*-Cas9 strain #51323
genomic DNA. Once *atlas*-GFP was obtained in a plasmid, we amplified it with primers that
contained 5' *Esp*3I sites and overhangs designed for Golden Gate Assembly (GGA) and that, in
the case of the reverse primer, also added on 42 nucleotides downstream of the *atlas* stop
codon to reach a PAM site identified by FlyCRISPR TargetFinder (Gratz et al. 2014) as being
optimal for Cas9/gRNA recognition and cleavage. The primer also introduced a mutation in the
PAM site so that insertion of the designed piece of DNA into the genome *in vivo* would not be
subject to re-cutting. We also used the NEB Q5 Site-Directed Mutagenesis kit to introduce a
silent mutation into the *atlas* protein-coding sequence to eliminate an internal *Esp*3I site. The
resulting construct was used as the "left" homology arm for homology-directed repair (HDR)
editing. We constructed a "right" homology arm by using NEB Q5 PCR to amplify a 982-bp
fragment downstream of the PAM site, using primers modified to contain *Esp*3I sites and
overhangs compatible with GGA. We performed GGA by combining these left and right arms, a
plasmid containing a PiggyBac transposase-excisable 3xP3-dsRed flanked by *Esp*3I sites, and
backbone plasmid pXZ13, with *Esp*3I and T4 DNA ligase (NEB). A combination of colony PCR,
restriction digestion and sequencing identified properly assembled plasmids suitable for HDR.
Vasa-Cas9 embryos were co-injected with the assembled plasmid and a pU6.3 plasmid
encoding a gRNA targeting the region just downstream of the *atlas* stop codon. G0 flies were
crossed to w^{1118} adults, and G1 flies were screened for red fluorescent eyes using the
NIGHTSEA system (NIGHTSEA LLC, Lexington, MA). Six balanced lines from two independent
G1 broods were established. To remove the dsRed from the *atlas* locus, we crossed these lines
to a PiggyBac transposase line (BDSC #8285) and then selected against pBac and dsRed in
the following generation. PCR and sequencing confirmed the expected "scarless" insertion of
GFP at the *atlas* locus.
831
832 Atlas *Rescue Line*
833
We constructed an HA-tagged *atlas* rescue line that contained the *atlas* gene flanked by
1345 bp of sequence upstream of the start codon (but excluding the coding sequence of
upstream gene CG3124) and 3000 bp of sequence downstream of the stop codon (including the
full 3' UTR) as follows. Genomic sequences were PCR amplified using Q5 High fidelity
Polymerase (NEB), purified Canton S genomic DNA (Gentra Puregene Tissue Kit, Qiagen,
Germantown, MD), and the atlas rescue F1/R1 and atlas rescue F3/R3 primer sets (see Table
S4). The 3x-HA tag was likewise amplified from pTWH (DGRC 1100; T. Murphy) using atlas
rescue F2/R2 primers. These DNA fragments were subsequently assembled into a XbaI/AscI-
linearized w+attB plasmid (Addgene, Watertown, MA, plasmid 30326, deposited by J.
Sekelsky). The assembled construct was then phiC31 integrated into the PBac{y^+-attP-
3B}VK00037 (Bloomington *Drosophila* Stock Center (BDSC) stock #24872) docking site
(Rainbow Transgenics) and crossed into the *atlas* null background to assess rescue.
846
847 *Fertility Assays and Sperm Visualization*
848

To validate the finding of reduced fertility for *atlas* knockdown males in the group fertility
assay described above, we performed single-pair fertility assays in which knockdown or mutant
males or their controls were mated individually to Canton S virgin females. Based on previous
experience analyzing genes that resulted in sterility or near-sterility (Gubala et al. 2017; Lange
et al. 2021), we designed assays with *N* = 20-30 flies per male genotype. Matings were
observed, and males were discarded after copulation. Females were allowed to lay eggs into
the vials for 4 days and then discarded. Pupal cases were counted as a measure of fertility.
Crosses to generate and mating assays involving RNAi flies were maintained at 25º to optimize
knockdown. Before all assays, flies were reared to sexual maturity (3-7 days) in single-sex
groups on cornmeal-molasses food supplemented with dry yeast grains (Gubala et al. 2017).
To assess the ability of *atlas*-GFP to rescue the fertility defect of the Δ*atlas* allele, we
crossed atlas-GFP and w1118 flies to Δ*atlas*/SM6. Males with genotypes atlas-GFP/Δ*atlas* and
+/Δ*atlas* were compared using the single-pair fertility assay described above.
To observe the production of sperm in knockdown or mutant males, we introduced the
Mst35Bb-GFP marker into these males, which labels mature sperm and late-stage spermatid
nuclei with GFP (Manier et al. 2010). Samples were prepared, imaged and analyzed as
described previously (Gubala et al. 2017); because the large differences in testis shape and
sperm production observed in initial phase contrast imaging would be obvious to any
experimenter, we were not blind to male genotypes.
868
869　*Atlas-GFP Ectopic Expression*
870
We used the Gateway cloning system (Thermo) to construct an *atlas*-GFP transgene
expressed under UAS control (primers in Table S4). The *atlas* protein-coding sequence in
pENTR was recombined with pTWG (*Drosophila* Genomics Resource Center, T. Murphy) as
described above. The resulting plasmid was then inserted into w- flies using P-element-
mediated transposition (Rainbow Transgenics), w+ G1s were selected, and several
independent insertions were balanced. We crossed male UAS-*atlas*:GFP flies to females from
two different driver lines: *tubulin*-GAL4 (to drive ubiquitous expression) and *Bam*-GAL4 (to drive
expression in the early germline). We dissected larval salivary glands of the *tub>atlas*:GFP
males, since these cells are exceptionally large and ideal for visualizing subcellular localization.
We then dissected the testes of *Bam>atlas*:GFP males to evaluate whether the localization
pattern observed in the salivary gland was consistent in testis tissue, albeit not the same cells in
which endogenous atlas appears to be expressed. Protein localization was visualized by
fluorescence confocal microscopy on a Leica SP5 microscope (Leica Microsystems, Wetzlar,
Germany) and images were captured with LASAF as described previously (Lange et al. 2021).
885
886　*Imaging Spermatogenesis and Spermatid Nuclear Condensation*
887
We used phase-contrast microscopy to examine the stages of spermatogenesis in whole
mount testes (White-Cooper 2004). To assess the processes of nuclear condensation and
individualization of 64-cell cysts of spermatids in the post-meiotic stages of spermatogenesis,
we used fluorescence and confocal microscopy to visualize actin-based individualization
complexes and nuclei. Samples were processed, and actin and nuclear DNA were visualized
with TRITC-phalloidin (Molecular Probes, Eugene, OR) and TOPRO-3 iodine (Thermo),
respectively, as described previously (Lange et al. 2021). The final stages of nuclear
condensation were visualized with the Mst35Bb-GFP marker described above, as well as an
equivalent marker, Mst35Bb-dsRed (Manier et al. 2010). Earlier nuclear stages were visualized
with histone H2AvD-RFP (BDSC stock #23651), which is present in round spermatid nuclei and
the earliest stages of nuclear elongation (Clarkson and Saint 1999; Rathke et al. 2007). Images

with H2AvD-RFP were obtained with epifluorescence microscopy, since we lack an appropriate
confocal laser for RFP.
To examine spermatid nuclei at various stages of condensation, we visualized nuclear
bundles using TOPRO-3. Testes of newly eclosed (<1 day old) *atlas* null and control males
were dissected in PBS. Testes were then transferred to a droplet of 4% paraformaldehyde on
poly-L-lysine treated glass slides and were gently shredded in the post-meiotic region to release
sperm bundles. Testes were gently squashed beneath coverslips coated in Sigmacote (Sigma
Aldrich, St. Louis, MO). We then froze slides in liquid nitrogen for a few seconds and popped off
of the siliconized coverslip with a razor. Slides were incubated in Coplin jars filled with 95%
ethanol at -20°C for 30 minutes and then mounted in VECTASHIELD (Vector Laboratories,
Burlingame, CA). Nuclear staging was performed by examining the shape of the nuclei. Early
and late canoe stages of condensation were distinguished by the absence or presence of
Mst35Bb-GFP, respectively. Elongated and condensed stages were distinguished by the
presence or absence, respectively, of vesicles of GFP-tagged nuclear proteins (Atlas-GFP or
Mst35Bb-GFP) located basal to the nuclei. Examples of stages are given in Fig. 4. Confocal
stacks were taken on a Leica SP5 microscope, images were captured by LASAF, and ImageJ
was used to flatten stacks into a single, two-dimensional image. All intact nuclear bundles were
counted for each dissection.
For the experiments measuring nuclear condensation stage (Table S1), a sample size of
N = 10 for each genotype was selected based on the magnitude of the *atlas* null phenotype and
the consistent differences observed in previous dissections of these genotypes with Mst35Bb-
GFP. Likewise, for the IC-nuclear bundle association and IC progression analysis (Fig. 3C-D),
we selected sample sizes of N = ~15 per genotype based on pilot experiments showing that
aberrant actin phenotypes were highly consistent in null testes and previous experience with
such quantification (Lange et al. 2021).
924
925 *Evolutionary and Gene Expression Analysis of* atlas
926
We searched for orthologs of the *D. melanogaster* Atlas protein in the original
sequenced *Drosophila* species with BLASTP searches in FlyBase (Clark et al. 2007). We also
used TBLASTN searches to identify orthologs in species lacking complete protein annotations.
We identified syntenic regions for each species by looking for conserved neighboring genes,
such as *ord* and *CG3124*. In addition to analyzing the *atlas* coding region, we conducted
separate BLASTN searches for the sequence of the *D. melanogaster* 3'UTR across *Drosophila*
species since it has a different conservation pattern than the coding sequence.
To test for sex-specific expression biases for both the ORF and the 3' UTR sequences,
we used adult male- and female-specific RNA-seq data from numerous *Drosophila* species
accessed through the Genomics Education Partnership version of the UCSC Genome Browser
(http://gander.wustl.edu/) and initially collected by Brown et al. (2014) and Chen et al. (2014).
We also confirmed these findings experimentally in several species by performing RT-PCR on
cDNA isolated from whole males and whole females, as previously described (Gubala et al.
2017).
To search for *atlas* orthologs in non-*Drosophila* Dipterans, we obtained from ENSEMBL
Metazoa the genomes of *Musca domestica, Glossina morsitans, Lucilia cuprina, Aedes aegypti,*
Anopheles darlingi, Anopheles gambiae, Culex quinquefasciatus and *Mayetiola destructor*. We
performed a synteny search (summarized in Fig. S10) in each species by identifying the nearest
neighbors of *atlas* in the *D. ananassae* and *D. virilis* genomes that had an identifiable homolog
in each species. In all cases, the homologs of the nearest neighbors on each side of atlas were
found on different contigs, suggesting synteny breakdown. We obtained up to 1 Mb of
sequence on each side of each identified homolog and queried it with BLASTN, TBLASTN, and
Exonerate (Slater and Birney 2005) for regions with significantly similarity to any portion of the

Atlas protein or cDNA sequences. No significant hits, and no hits better than what could be
found in other parts of the genome, were found. Finally, we used HMMER to search for
orthologs in all annotated proteomes and all metazoan proteomes. We first queried the
database with Atlas from either *D. melanogaster* or *D. virilis* and accepted hits that fell below an
e-value cutoff of 0.01 and a minimum hit length of 3%. These hits were then included iteratively
in subsequent searches until no new significant hits were found.
956   We analyzed the molecular evolution of the *atlas* protein-coding sequence by obtaining
orthologous protein-coding sequences from *melanogaster* group species. (Analysis out of this
group was not performed due to high sequence divergence and poor alignment quality.) We
used BLASTP to identify these sequences from GenBank and then extracted the coding DNA
sequence for each. Sequences were aligned, checked for recombination, used to construct a
gene tree, and analyzed with the PAML sites test as described previously (McGeary and Findlay
2020), except that alignment positions that included gaps were masked from the PAML
analysis. We initially analyzed a set of 13 species (*melanogaster*, *simulans*, *sechellia*, *yakuba*,
erecta, *suzukii*, *takahashii*, *biarmipes*, *rhopaloa*, *ficusphila*, *elegans*, *eugracilis* and *ananassae*);
we excluded an ortholog detected in *D. bipectinata* due to poor alignment. This initial analysis
detected a class of sites with significant evidence of positive selection, but closer inspection of
the alignment revealed that the site with the strongest evidence of selection, corresponding to
D. melanogaster residue 31R, may have been driven by a questionable alignment due to an
insertion in that region that was unique to *D. takahashii*. The results reported derive from an
analysis with *D. takahashii* excluded, which produced a more reliable alignment and showed no
evidence for any sites under positive selection.
972

<u>Acknowledgements</u>

We thank Dr. Alexis Hill for assistance with developing the *atlas*-GFP transgene; Gynesis
Vance, Elvis Perez, Ishanpepe Jagusah, Emily Gualdino, and students in the Biology 261 Lab at
Holy Cross for assistance with RNAi fertility screens; Dr. Rob Bellin for assistance with confocal
microscopy; Dr. Justin McAlister for use of the NIGHTSEA system; the Bloomington Stock
Center and the Vienna *Drosophila* Resource Center for fly strains; and Dr. Mariana Wolfner and
members of the Findlay and Bornberg-Bauer labs for helpful discussions about the project and
the manuscript. This work was supported by NSF CAREER Award #1652013 (to GDF) and a
Humboldt Fellowship (to AG). GDF and EBB are also grateful to the Evolution Think Tank at
the University of Münster, which funded GDF as a visiting fellow for in-person collaboration.

References

Abrusán G. 2013. Integration of New Genes into Cellular Networks, and Their Structural
        Maturation. Genetics 195:1407-1417.
Alvi ZA, Chu T-C, Schawaroch V, Klaus AV. 2016. Genomic and expression analysis of
        transition proteins in *Drosophila*. Spermatogenesis 5:e1178518.
Alvi ZA, Chu T-C, Schawaroch V, Klaus AV. 2013. Protamine-like proteins in 12 sequenced
        species of Drosophila. Protein and peptide letters 20:17-35.
Argyridou E, Huylmans AK, Königer A, Parsch J. 2017. X-linkage is not a general inhibitor of
        tissue-specific gene expression in *Drosophila melanogaster*. Heredity 119:27-34.
Argyridou E, Parsch J. 2018. Regulation of the X Chromosome in the Germline and Soma of
        Drosophila melanogaster Males. Genes 9:242.
Baalsrud HT, Tørresen OK, Solbakken MH, Salzburger W, Hanel R, Jakobsen KS, Jentoft S.
        2018. *De novo* gene evolution of antifreeze glycoproteins in codfishes revealed by whole
        genome sequence data. Molecular Biology and Evolution 35:593-606.
Baker RH, Narechania A, Johns PM, Wilkinson GS. 2012. Gene duplication, tissue-specific
        gene expression and sexual conflict in stalk-eyed flies (Diopsidae). Philosophical
        Transactions of the Royal Society of London. Series B, Biological Sciences 367:2357-
        2375.
Begun DJ, Lindfors HA, Kern AD, Jones CD. 2007. Evidence for *de Novo* Evolution of Testis-
        Expressed Genes in the *Drosophila yakuba/Drosophila erecta* Clade. Genetics
        176:1131-1137.
Bier E, Harrison MM, O'Connor-Giles KM, Wildonger J. 2018. Advances in Engineering the Fly
        Genome with the CRISPR-Cas System. Genetics 208:1-18.
Bornberg-Bauer E, Schmitz JF. 2017. Fact or fiction: Updates on how protein-coding genes
        might emerge *de novo* from previously non-coding DNA. F1000Research 6:57.
Bouska MJ, Bai H. 2021. Long noncoding RNA regulation of spermatogenesis via the spectrin
        cytoskeleton in Drosophila. G3 Genes|Genomes|Genetics.
Brown JB, Boley N, Eisman R, May GE, Stoiber MH, Duff MO, Booth BW, Wen J, Park S,
        Suzuki AM, et al. 2014. Diversity and dynamics of the Drosophila transcriptome. Nature
        512:393-399.
Bruckner JJ, Zhan H, Gratz SJ, Rao M, Ukken F, Zilberg G, O'Connor-Giles KM. 2017. Fife
        organizes synaptic vesicles and calcium channels for high-probability neurotransmitter
        release. Journal of Cell Biology 216:231-246.
Cai J, Zhao R, Jiang H, Wang W. 2008. *De novo* origination of a new protein-coding gene in
        Saccharomyces cerevisiae. Genetics 179:487-496.
Carvunis A-R, Rolland T, Wapinski I, Calderwood MA, Yildirim MA, Simonis N, Charloteaux B,
        Hidalgo C, Barbette J, Santhanam B, et al. 2012. Proto-genes and *de novo* gene birth.
        Nature 487:370-374.
Chamakura KR, Tran JS, O'Leary C, Lisciandro HG, Antillon SF, Garza KD, Tran E, Min L,
        Young R. 2020. Rapid de novo evolution of lysis genes in single-stranded RNA phages.
        Nature Communications 11.
Chen Z-X, Sturgill D, Qu J, Jiang H, Park S, Boley N, Suzuki AM, Fletcher AR, Plachetzki DC,
        FitzGerald PC, et al. 2014. Comparative validation of the *D. melanogaster* modENCODE
        transcriptome annotation. Genome Research 24:1209-1223.
Civetta A, Finn S. 2014. Do candidate genes mediating conspecific sperm precedence affect
        sperm competitive ability within species? A test case in *Drosophila*. G3 (Bethesda, Md.)
        4:1701-1707.
Clark AG, Eisen MB, Smith DR, Bergman CM, Oliver B, Markow TA, Kaufman TC, Kellis M,
        Gelbart W, Iyer VN, et al. 2007. Evolution of genes and genomes on the Drosophila
        phylogeny. Nature 450:203-218.

1036   Clarkson M, Saint R. 1999. A His2AvDGFP fusion gene complements a lethal His2AvD mutant
1037   allele and provides an in vivo marker for Drosophila chromosome behavior. DNA and
1038   Cell Biology 18:457-462.
1039   Cui X, Lv Y, Chen M, Nikoloski Z, Twell D, Zhang D. 2015. Young genes out of the male: an
1040   insight from evolutionary age analysis of the pollen transcriptome. Molecular Plant
1041   8:935-945.
1042   Demarco RS, Eikenes TH, Haglund K, Jones DL. 2014. Investigating spermatogenesis in
1043   *Drosophila melanogaster*. Methods 68:218-227.
1044   Dietzl G, Chen D, Schnorrer F, Su K-C, Barinova Y, Fellner M, Gasser B, Kinsey K, Oppel S,
1045   Scheiblauer S, et al. 2007. A genome-wide transgenic RNAi library for conditional gene
1046   inactivation in *Drosophila*. Nature 448:151-156.
1047   Dorus S, Busby SA, Gerike U, Shabanowitz J, Hunt DF, Karr TL. 2006. Genomic and functional
1048   evolution of the *Drosophila melanogaster* sperm proteome. Nature Genetics 38:1440-
1049   1445.
1050   Dorus S, Freeman ZN, Parker ER, Heath BD, Karr TL. 2008. Recent Origins of Sperm Genes in
1051   Drosophila. Molecular Biology and Evolution 25:2157-2166.
1052   Doyen CM, Chalkley GE, Voets O, Bezstarosti K, Demmers JA, Moshkin YM, Verrijzer CP.
1053   2015. A Testis-Specific Chaperone and the Chromatin Remodeler ISWI Mediate
1054   Repackaging of the Paternal Genome. Cell Reports 13:1310-1318.
1055   Emerson JJ, Kaessmann H, Betrán E, Long M. 2004. Extensive gene traffic on the mammalian
1056   X chromosome. Science 303:537-540.
1057   Eren-Ghiani Z, Rathke C, Theofel I, Renkawitz-Pohl R. 2015. Prtl99C Acts Together with
1058   Protamines and Safeguards Male Fertility in *Drosophila*. Cell Reports 13:2327-2335.
1059   Fabian L, Brill JA. 2012. Drosophila spermiogenesis. Spermatogenesis 2:197-212.
1060   Findlay GD, Sitnik JL, Wang W, Aquadro CF, Clark NL, Wolfner MF. 2014. Evolutionary rate
1061   covariation identifies new members of a protein network required for *Drosophila
1062   melanogaster* female post-mating responses. PLoS Genetics 10:e1004108.
1063   Gao G, Vibranovski MD, Zhang L, Li Z, Liu M, Zhang YE, Li X, Zhang W, Fan Q, Vankuren NW,
1064   et al. 2014. A long-term demasculinization of X-linked intergenic noncoding RNAs in
1065   *Drosophila melanogaster*. Genome Research 24:629-638.
1066   Gärtner SMK, Hundertmark T, Nolte H, Theofel I, Eren-Ghiani Z, Tetzner C, Duchow TB,
1067   Rathke C, Krüger M, Renkawitz-Pohl R. 2019. Stage-specific testes proteomics of
1068   *Drosophila melanogaster* identifies essential proteins for male fertility. European Journal
1069   of Cell Biology 98:103-115.
1070   Gärtner SMK, Rothenbusch S, Buxa MK, Theofel I, Renkawitz R, Rathke C, Renkawitz-Pohl R.
1071   2015. The HMG-box-containing proteins tHMG-1 and tHMG-2 interact during the
1072   histone-to-protamine transition in *Drosophila* spermatogenesis. European Journal of Cell
1073   Biology 94:46-59.
1074   Ge DT, Tipping C, Brodsky MH, Zamore PD. 2016. Rapid screening for CRISPR-directed
1075   editing of the *Drosophila* genome using *white* coconversion. G3 (Bethesda, Md.) 6:3197-
1076   3206.
1077   Gratz SJ, Ukken FP, Rubinstein CD, Thiede G, Donohue LK, Cummings AM, O'Connor-Giles
1078   KM. 2014. Highly specific and efficient CRISPR/Cas9-catalyzed homology-directed
1079   repair in Drosophila. Genetics 196:961-971.
1080   Gubala AM, Schmitz JF, Kearns MJ, Vinh TT, Bornberg-Bauer E, Wolfner MF, Findlay GD.
1081   2017. The *goddard* and *saturn* genes are essential for *Drosophila* male fertility and may
1082   have arisen *de novo*. Molecular Biology and Evolution 34:1066-1082.
1083   Guerzoni D, McLysaght A. 2016. *De novo* genes arise at a slow but steady rate along the
1084   primate lineage and have been subject to incomplete lineage sorting. Genome biology
1085   and evolution 8:1222-1232.

1086   Hales KG, Korey CA, Larracuente AM, Roberts DM. 2015. Genetics on the fly: a primer on the
1087   *Drosophila* model system. Genetics 201:815-842.
1088   Heames B, Schmitz J, Bornberg-Bauer E. 2020. A continuum of evolving de novo genes drives
1089   protein-coding novelty in *Drosophila*. Journal of Molecular Evolution 88:382-398.
1090   Hill A, Jain P, Ben-Shahar Y. 2019. The *Drosophila* ERG channel seizure plays a role in the
1091   neuronal homeostatic stress response. PLoS Genetics 15:e1008288.
1092   Hundertmark T, Gärtner SMK, Rathke C, Renkawitz-Pohl R. 2018. Nejire/dCBP-mediated
1093   histone H3 acetylation during spermatogenesis is essential for male fertility in *Drosophila*
1094   *melanogaster*. PLOS ONE 13:e0203622.
1095   Jayaramaiah Raja S, Renkawitz-Pohl R. 2005. Replacement by *Drosophila melanogaster*
1096   protamines and Mst77F of histones during chromatin condensation in late spermatids
1097   and role of Sesame in the removal of these proteins from the male pronucleus.
1098   Molecular and Cellular Biology 25:6165-6177.
1099   Kaessmann H. 2010. Origins, evolution, and phenotypic impact of new genes. Genome
1100   Research 20:1313-1326.
1101   Kanippayoor RL, Alpern JHM, Moehring AJ. 2013. Protamines and spermatogenesis in
1102   *Drosophila* and *Homo sapiens*. Spermatogenesis 3:e24376.
1103   Keeling DM, Garza P, Nartey CM, Carvunis A-R. 2019. The meanings of 'function' in biology
1104   and the problematic case of de novo gene emergence. eLife 8:e47014.
1105   Kemkemer C, Catalán A, Parsch J. 2014. 'Escaping' the X chromosome leads to increased
1106   gene expression in the male germline of *Drosophila melanogaster*. Heredity 112:149-
1107   155.
1108   Kimura S, Loppin B. 2016. The Drosophila chromosomal protein Mst77F is processed to
1109   generate an essential component of mature sperm chromatin. Open Biology 6:160207.
1110   Kondo S, Vedanayagam J, Mohammed J, Eizadshenass S, Kan L, Pang N, Aradhya R, Siepel
1111   A, Steinhauer J, Lai EC. 2017. New genes often acquire male-specific functions but
1112   rarely become essential in *Drosophila*. Genes & development 31:1841-1846.
1113   Landeen EL, Muirhead CA, Wright L, Meiklejohn CD, Presgraves DC. 2016. Sex Chromosome-
1114   wide Transcriptional Suppression and Compensatory *Cis*-Regulatory Evolution Mediate
1115   Gene Expression in the *Drosophila* Male Germline. PLOS Biology 14:e1002499.
1116   Lange A, Mills RE, Lange CJ, Stewart M, Devine SE, Corbett AH. 2007. Classical nuclear
1117   localization signals: definition, function, and interaction with Importin α. Journal of
1118   Biological Chemistry 282:5101-5105.
1119   Lange A, Patel PH, Heames B, Damry AM, Saenger T, Jackson CJ, Findlay GD, Bornberg-
1120   Bauer E. 2021. Structural and functional characterization of a putative de novo gene in
1121   *Drosophila*. Nature Communications 12:1667.
1122   Larkin A, Marygold SJ, Antonazzo G, Attrill H, Gilberto, Garapati PV, Joshua, Millburn G,
1123   Strelets VB, Tabone CJ, et al. 2021. FlyBase: updates to the *Drosophila melanogaster*
1124   knowledge base. Nucleic acids research 49:D899-D907.
1125   Levine MT, Jones CD, Kern AD, Lindfors HA, Begun DJ. 2006. Novel genes derived from
1126   noncoding DNA in *Drosophila melanogaster* are frequently X-linked and exhibit testis-
1127   biased expression. Proceedings of the National Academy of Sciences 103:9935-9939.
1128   Li D, Yan Z, Lu L, Jiang H, Wang W. 2014. Pleiotropy of the *de novo*-originated gene *MDF1*.
1129   Scientific reports 4:7280.
1130   Li Z-W, Chen X, Wu Q, Hagmann J, Han T-S, Zou Y-P, Ge S, Guo Y-L. 2016. On the origin of
1131   *de novo* genes in *Arabidopsis thaliana* populations. Genome biology and evolution
1132   8:2190-2202.
1133   Lim C, Tarayrah L, Chen X. 2012. Transcriptional regulation during Drosophila
1134   spermatogenesis. Spermatogenesis 2:158-166.

1135   Lipinski KJ, Farslow JC, Fitzpatrick KA, Lynch M, Katju V, Bergthorsson U. 2011. High
1136   spontaneous rate of gene duplication in Caenorhabditis elegans. Current biology : CB
1137   21:306-310.
1138   Lu T-C, Leu J-Y, Lin W-C. 2017. A comprehensive analysis of transcript-supported de novo
1139   genes in *Saccharomyces sensu stricto* yeasts. Molecular Biology and Evolution 34:2823-
1140   2838.
1141   Mahadevaraju S, Fear JM, Akeju M, Galletta BJ, Pinheiro MMLS, Avelino CC, Cabral-De-Mello
1142   DC, Conlon K, Dell'Orso S, Demere Z, et al. 2021. Dynamic sex chromosome
1143   expression in Drosophila male germ cells. Nature Communications 12.
1144   Manier MK, Belote JM, Berben KS, Novikov D, Stuart WT, Pitnick S. 2010. Resolving
1145   mechanisms of competitive fertilization success in *Drosophila melanogaster*. Science
1146   328:354-357.
1147   Marques AC, Dupanloup I, Vinckenbosch N, Reymond A, Kaessmann H. 2005. Emergence of
1148   young human genes after a burst of retroposition in primates. PLOS Biology 3:e357.
1149   McGeary MK, Findlay GD. 2020. Molecular evolution of the sex peptide network in *Drosophila*.
1150   Journal of Evolutionary Biology 33:629-641.
1151   McLysaght A, Hurst LD. 2016. Open questions in the study of *de novo* genes: what, how and
1152   why. Nature Reviews Genetics 17:567.
1153   Meiklejohn CD, Landeen EL, Cook JM, Kingan SB, Presgraves DC. 2011. Sex Chromosome-
1154   Specific Regulation in the *Drosophila* Male Germline But Little Evidence for
1155   Chromosomal Dosage Compensation or Meiotic Inactivation. PLOS Biology 9:e1001126.
1156   Mikhaylova LM, Nurminsky DI. 2011. Lack of global meiotic sex chromosome inactivation, and
1157   paucity of tissue-specific gene expression on the Drosophila X chromosome. BMC
1158   Biology 9:29.
1159   Moyers BA, Zhang J. 2015. Phylostratigraphic bias creates spurious patterns of genome
1160   evolution. Molecular Biology and Evolution 32:258-267.
1161   Moyers BA, Zhang J. 2018. Toward reducing phylostratigraphic errors and biases. Genome
1162   biology and evolution 10:2037-2048.
1163   Necsulea A, Kaessmann H. 2014. Evolutionary dynamics of coding and non-coding
1164   transcriptomes. Nature Reviews Genetics 15:734-748.
1165   Ni J-Q, Zhou R, Czech B, Liu L-P, Holderbaum L, Yang-Zhou D, Shim H-S, Tao R, Handler D,
1166   Karpowicz P, et al. 2011. A genome-scale shRNA resource for transgenic RNAi in
1167   *Drosophila*. Nature Methods 8:405-407.
1168   Palmieri N, Kosiol C, Schlötterer C. 2014. The life cycle of *Drosophila* orphan genes. eLife
1169   3:e01311.
1170   Parisi M, Nuttall R, Edwards P, Minor J, Naiman D, Lü J, Doctolero M, Vainer M, Chan C,
1171   Malley J, et al. 2004. A survey of ovary-, testis-, and soma-biased gene expression in
1172   *Drosophila melanogaster* adults. Genome biology 5:R40.
1173   Parisi M, Nuttall R, Naiman D, Bouffard G, Malley J, Andrews J, Eastman S, Oliver B. 2003.
1174   Paucity of genes on the *Drosophila* X chromosome showing male-biased expression.
1175   Science 299:697-700.
1176   Potter SC, Luciani A, Eddy SR, Park Y, Lopez R, Finn RD. 2018. HMMER web server: 2018
1177   update. Nucleic acids research 46:W200-W204.
1178   Puntambekar S, Newhouse R, Navas JSM, Chauhan R, Vernaz G, Willis T, Wayland MT,
1179   Umrania Y, Miska EA, Prabakaran S. 2020. Evolutionary divergence of novel open
1180   reading frames in cichlids speciation. Scientific reports 10.
1181   Rathke C, Baarends WM, Awe S, Renkawitz-Pohl R. 2014. Chromatin dynamics during
1182   spermiogenesis. In. Chromatin and epigenetic regulation of animal development. p. 155-
1183   168.
1184   Rathke C, Baarends WM, Jayaramaiah-Raja S, Bartkuhn M, Renkawitz R, Renkawitz-Pohl R.
1185   2007. Transition from a nucleosome-based to a protamine-based chromatin

configuration during spermiogenesis in *Drosophila*. Journal of cell science 120:1689-
1700.
Rathke C, Barckmann B, Burkhard S, Jayaramaiah-Raja S, Roote J, Renkawitz-Pohl R. 2010.
Distinct functions of Mst77F and protamines in nuclear shaping and chromatin
condensation during *Drosophila* spermiogenesis. European Journal of Cell Biology
89:326-338.
Ravi Ram K, Wolfner MF. 2007. Sustained post-mating response in *Drosophila melanogaster*
requires multiple seminal fluid proteins. PLoS Genetics 3:e238.
Reinhardt JA, Wanjiru BM, Brant AT, Saelao P, Begun DJ, Jones CD. 2013. *De novo* ORFs in
Drosophila are important to organismal fitness and evolved rapidly from previously non-
coding sequences. PLoS Genetics 9:e1003860.
Reinke V, San Gil I, Ward S, Kazmer K. 2004. Genome-wide germline-enriched and sex-biased
expression profiles in *Caenorhabditis elegans*. Development 131:311-323.
Rele CP, Sandlin KM, Leung W, Reed LK. 2020. Manual Annotation of Genes within *Drosophila*
Species: the Genomics Education Partnership protocol. bioRxiv 12.10.420521.
Ruiz-Orera J, Hernandez-Rodriguez J, Chiva C, Sabidó E, Kondova I, Bontrop R, Marqués-
Bonet T, Albà MM. 2016. Origins of *de novo* genes in human and chimpanzee. PLoS
Genetics 11:e1005721.
Schaeffer SW, Bhutkar A, McAllister BF, Matsuda M, Matzkin LM, O'Grady PM, Rohde C,
Valente VLS, Aguadé M, Anderson WW, et al. 2008. Polytene Chromosomal Maps of 11
Drosophila Species: The Order of Genomic Scaffolds Inferred From Genetic and
Physical Maps. Genetics 179:1601-1655.
Schlötterer C. 2015. Genes from scratch – the evolutionary fate of *de novo* genes. Trends in
Genetics 31:215-219.
Schmitz JF, Ullrich KK, Bornberg-Bauer E. 2018. Incipient *de novo* genes can evolve from
frozen accidents that escaped rapid transcript turnover. Nature Ecology & Evolution
2:1626-1632.
Schnakenberg SL, Matias WR, Siegal ML. 2011. Sperm-storage defects and live birth in
Drosophila females lacking spermathecal secretory cells. PLOS Biology 9:e1001192.
Schuh M, Lehner CF, Heidmann S. 2007. Incorporation of *Drosophila* CID/CENP-A and CENP-
C into Centromeres during Early Embryonic Anaphase. Current Biology 17:237-243.
Slater G, Birney E. 2005. Automated generation of heuristics for biological sequence
comparison. BMC Bioinformatics 6:31.
Sorourian M, Kunte MM, Domingues S, Gallach M, Özdil F, Río J, Betrán E. 2014. Relocation
facilitates the acquisition of short *cis*-regulatory regions that drive the expression of
retrogenes during spermatogenesis in *Drosophila*. Molecular Biology and Evolution
31:2170-2180.
Soumillon M, Necsulea A, Weier M, Brawand D, Zhang X, Gu H, Barthès P, Kokkinaki M, Nef S,
Gnirke A, et al. 2013. Cellular source and mechanisms of high transcriptome complexity
in the mammalian testis. Cell Reports 3:2179-2190.
Steinhauer J. 2015. Separating from the pack: molecular mechanisms of Drosophila spermatid
individualization. Spermatogenesis 5:e1041345.
Suenaga Y, Islam SMR, Alagu J, Kaneko Y, Kato M, Tanaka Y, Kawana H, Hossain S,
Matsumoto D, Yamamoto M, et al. 2014. NCYM, a *cis*-antisense gene of MYCN,
encodes a *de novo* evolved protein that inhibits GSK3β resulting in the stabilization of
MYCN in human neuroblastomas. PLoS Genetics 10:e1003996.
Tirmarche S, Kimura S, Sapey-Triomphe L, Sullivan W, Landmann F, Loppin B. 2014.
Drosophila Protamine-Like Mst35Ba and Mst35Bb Are Required for Proper Sperm
Nuclear Morphology but Are Dispensable for Male Fertility. G3
Genes|Genomes|Genetics 4:2241-2245.

Vakirlis N, Acar O, Hsu B, Castilho Coelho N, Van Oss SB, Wacholder A, Medetgul-Ernar K,
        Bowman RW, Hines CP, Iannotta J, et al. 2020. De novo emergence of adaptive
        membrane proteins from thymine-rich genomic sequences. Nature Communications 11.
Vakirlis N, Hebert AS, Opulente DA, Achaz G, Hittinger CT, Fischer G, Coon JJ, Lafontaine I.
        2018. A molecular portrait of de novo genes in yeasts. Molecular Biology and Evolution
        35:631-645.
Van Oss SB, Carvunis A-R. 2019. *De novo* gene birth. PLoS Genetics 15:e1008160.
VanKuren NW, Long M. 2018. Gene duplicates resolving sexual conflict rapidly evolved
        essential gametogenesis functions. Nature Ecology & Evolution 2:705-712.
Vibranovski MD. 2014. Meiotic Sex Chromosome Inactivation in *Drosophila*. Journal of
        Genomics 2:104-117.
Vibranovski MD, Lopes HF, Karr TL, Long M. 2009. Stage-specific expression profiling of
        Drosophila spermatogenesis suggests that meiotic sex chromosome inactivation drives
        genomic relocation of testis-expressed genes. PLoS Genetics 5:e1000731.
Vibranovski MD, Zhang Y, Long M. 2009. General gene movement off the X chromosome in the
        Drosophila genus. Genome Research 19:897-903.
Vibranovski MD, Zhang YE, Kemkemer C, Lopes HF, Karr TL, Long M. 2012. Re-analysis of the
        larval testis data on meiotic sex chromosome inactivation revealed evidence for tissue-
        specific gene expression related to the drosophila X chromosome. BMC Biology 10:49.
Wasbrough ER, Dorus S, Hester S, Howard-Murkin J, Lilley K, Wilkin E, Polpitiya A, Petritis K,
        Karr TL. 2010. The *Drosophila melanogaster* sperm proteome-II (DmSP-II). Journal of
        Proteomics 73:2171-2185.
Weisman CM, Murray AW, Eddy SR. 2020. Many, but not all, lineage-specific genes can be
        explained by homology detection failure. PLOS Biology 18:e3000862.
Wen K, Yang L, Xiong T, Di C, Ma D, Wu M, Xue Z, Zhang X, Long L, Zhang W, et al. 2016.
        Critical roles of long noncoding RNAs in *Drosophila* Spermatogenesis. Genome
        Research 26:1233-1244.
White-Cooper H. 2010. Molecular mechanisms of gene regulation during Drosophila
        spermatogenesis. REPRODUCTION 139:11-21.
White-Cooper H. 2004. Spermatogenesis: analysis of meiosis and morphogenesis. In:
        Henderson D, editor. Drosophila Cytogenetics Protocols. Totowa, NJ: Humana Press. p.
        45-75.
White-Cooper H. 2012. Tissue, cell type and stage-specific ectopic gene expression and RNAi
        induction in the Drosophila testis. Spermatogenesis 2:11-22.
Wilburn DB, Swanson WJ. 2016. From molecules to mating: Rapid evolution and biochemical
        studies of reproductive proteins. Journal of Proteomics 135:12-25.
Wilson BA, Foy SG, Neme R, Masel J. 2017. Young genes are highly disordered as predicted
        by the preadaptation hypothesis of *de novo* gene birth. Nature Ecology & Evolution
        1:0146.
Witt E, Benjamin S, Svetec N, Zhao L. 2019. Testis single-cell RNA-seq reveals the dynamics of
        de novo gene transcription and germline mutational bias in *Drosophila*. eLife 8.
Wong A, Albright SN, Giebel JD, Ram KR, Ji S, Fiumera AC, Wolfner MF. 2008. A role for
        Acp29AB, a predicted seminal fluid lectin, in female sperm storage in *Drosophila
        melanogaster*. Genetics 180:921-931.
Wu DD, Irwin DM, Zhang YP. 2011. De novo origin of human protein-coding genes. PLoS
        Genetics 7.
Xiao W, Liu H, Li Y, Li X, Xu C, Long M, Wang S. 2009. A rice gene of *de novo* origin negatively
        regulates pathogen-induced defense response. PLOS ONE 4:e4603.
Xie C, Bekpen C, Künzel S, Keshavarz M, Krebs-Wheaton R, Skrabar N, Ullrich KK, Tautz D.
        2019. A de novo evolved gene in the house mouse regulates female pregnancy cycles.
        eLife 8.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. Molecular Biology and
Evolution 24:1586-1591.
Yeh SD, Do T, Chan C, Cordova A, Carranza F, Yamamoto EA, Abbassi M, Gandasetiawan
KA, Librado P, Damia E, et al. 2012. Functional evidence that a recently evolved
Drosophila sperm-specific gene boosts sperm competition. Proceedings of the National
Academy of Sciences 109:2043-2048.
Zdobnov EM, Tegenfeldt F, Kuznetsov D, Waterhouse RM, Simão FA, Ioannidis P, Seppey M,
Loetscher A, Kriventseva EV. 2017. OrthoDB v9.1: cataloging evolutionary and
functional annotations for animal, fungal, plant, archaeal, bacterial and viral orthologs.
Nucleic acids research 45:D744-D749.
Zhang J. 2003. Evolution by gene duplication: an update. Trends in Ecology & Evolution 18:292-
298.
Zhang L, Ren Y, Yang T, Li G, Chen J, Gschwend AR, Yu Y, Hou G, Zi J, Zhou R, et al. 2019.
Rapid evolution of protein diversity by de novo origination in *Oryza*. Nature Ecology &
Evolution 3:679-690.
Zhang W, Landback P, Gschwend AR, Shen B, Long M. 2015. New genes drive the evolution of
gene interaction networks in the human and mouse genomes. Genome biology 16.
Zhang YE, Vibranovski MD, Krinsky BH, Long M. 2010. Age-dependent chromosomal
distribution of male-biased genes in *Drosophila*. Genome Research 20:1526-1533.
Zhang YE, Vibranovski MD, Landback P, Marais GAB, Long M. 2010. Chromosomal
Redistribution of Male-Biased Genes in Mammalian Evolution with Two Bursts of Gene
Gain on the X Chromosome. PLOS Biology 8:e1000494.
Zhao L, Saelao P, Jones CD, Begun DJ. 2014. Origin and spread of *de novo* genes in
Drosophila melanogaster populations. Science 343:769-772.
Zhuang X, Yang C, Murphy KR, Cheng CHC. 2019. Molecular mechanism and history of non-
sense to sense evolution of antifreeze glycoprotein gene in northern gadids.
Proceedings of the National Academy of Sciences 116:4400-4405.
1314