A putative *de novo* evolved gene required for spermatid chromatin condensation in *Drosophila melanogaster*

Emily L. Rivard[1,4,*], Andrew G. Ludwig[1,*], Prajal H. Patel[1,*], Anna Grandchamp[2], Sarah E. Arnold[1,5], Alina Berger[2], Emilie M. Scott[1], Brendan J. Kelly[1,6], Grace C. Mascha[1], Erich Bornberg-Bauer[2,3], Geoffrey D. Findlay[1,**]

1. Department of Biology, College of the Holy Cross, Worcester, MA, USA
2. Institute for Evolution and Biodiversity, University of Münster, Münster, Germany
3. Department of Protein Evolution, Max Planck Institute for Developmental Biology, Tübingen, Germany
4. Current address: Department of Molecular and Cellular Biology, Harvard University, Cambridge, MA, USA
5. Current address: Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA, USA
6. Current address: Department of Entomology, The Ohio State University, Columbus, OH, USA

*equal contribution
**Correspondence to: gfindlay@holycross.edu

<u>Abstract</u>

Comparative genomics has enabled the identification of genes that potentially evolved *de novo*

from non-coding sequences. Many such genes are expressed in male reproductive tissues, but

their functions remain poorly understood. To address this, we conducted a functional genetic

screen of over 40 putative *de novo* genes with testis-enriched expression in *Drosophila*

melanogaster and identified one gene, *atlas,* required for male fertility. Detailed genetic and

cytological analyses showed that *atlas* is required for proper chromatin condensation during the

final stages of spermatogenesis. Atlas protein is expressed in spermatid nuclei and facilitates

the transition from histone- to protamine-based chromatin packaging. Complementary

evolutionary analyses revealed the complex evolutionary history of *atlas*. The protein-coding

portion of the gene likely arose at the base of the *Drosophila* genus on the X chromosome but

was unlikely to be essential, as it was then lost in several independent lineages. Within the last

~15 million years, however, the gene moved to an autosome, where it fused with a conserved

non-coding RNA and evolved a non-redundant role in male fertility. Altogether, this study

provides insight into the integration of novel genes into biological processes, the links between

genomic innovation and functional evolution, and the genetic control of a fundamental

developmental process, gametogenesis.

Author Summary

Genomes are in flux, as genes are constantly added and lost throughout evolution. New genes were once thought to arise almost exclusively via the modification or duplication of existing genes. Recently, however, interest has grown in alternative modes of new gene origination, such as *de novo* evolution from genetic material that previously did not encode proteins. Many *de novo* genes are expressed in male reproductive tissues, but their significance for fertility is not well understood. We screened likely *de novo* genes expressed in the *Drosophila* testis for reproductive roles and found one gene, *atlas*, essential for male fertility. We leveraged genetic and cell biological experiments to investigate roles for Atlas protein in reproduction and found that it is required during sperm development for proper packaging of DNA in the sperm nucleus. Evolutionary analyses of this gene revealed a complicated history, including loss in some lineages, movement between chromosomes, and fusion with a non-protein-coding gene. Studying both the functions and evolutionary histories of new proteins illustrates how they might evolve critical roles in biological processes despite their relative novelty. Furthermore, the study of *atlas* identifies an essential genetic player in the fly testis, an important model system for understanding how gametes are produced.

Introduction

The evolution of new genes is integral to the extensive genotypic and phenotypic diversity observed across species. The best-characterized mechanism of novel gene emergence is gene duplication [1, 2]; however, rapid expansion in high-quality genomic resources has provided mounting evidence of lineage-specific sequences and the existence of alternative modes of new gene origination. One such mechanism is *de novo* evolution, the birth of new genes from previously non-genic or intronic regions, which is now a widely acknowledged source of protein-coding and RNA genes [3-5]. Although *de novo* origination was once considered an unlikely event, catalogs of *de novo* genes have now been published for an expansive range of species [6-13]. Multiple models explain how protein-coding *de novo* genes may acquire both an open reading frame (ORF) and regulatory sequences permitting transcription [14-17]. Interrogation of the biochemical and biophysical properties of the proteins encoded by *de novo* genes has offered initial insight into the mechanisms of emergence and functional potential of these genes [17-20].

The capacity of protein-coding *de novo* genes to evolve important functions is a topic of interest from evolutionary, physiological and molecular perspectives [21]. In the last couple of decades, the products of *de novo* genes have been shown to play diverse roles in a variety of organisms. For example, *de novo* genes function in fundamental molecular processes in yeast, such as *BSC4*, a gene implicated in DNA repair, and *MDF1,* which mediates crosstalk between reproduction and growth [22, 23]. *De novo* genes also evolve roles in organismal responses to disease and changing environmental factors. A putatively *de novo* evolved gene in rice regulates the plant's pathogen resistance response to strains causing bacterial blight [24]. Antifreeze glycoprotein genes, essential for survival in frigid ocean temperatures, evolved *de novo* in the ancestor of Arctic codfishes to coincide with cooling oceans in the Northern Hemisphere [25, 26]. *De novo* genes are additionally implicated in the development and physiology of mammals. In house mice, a *de novo* evolved gene expressed in the oviduct

functions in female fertility by regulating pregnancy cycles [27]. A *de novo* gene found in

humans and chimpanzees regulates the oncogenesis and growth of neuroblastoma, revealing

the relevance of novel genes to human disease [28]. These studies have started to

demonstrate the significance of *de novo* genes, thereby challenging previous assumptions that

only ancient, highly conserved genes can be essential.

Across multicellular animals, male reproductive tissues serve as hubs for new gene

emergence via numerous mechanisms, including *de novo* evolution [12, 29-34]. Proposed

causes of this "out of the testis" phenomenon include the high level of promiscuous transcription

in testis cells [35, 36], the relative simplicity of promoter regions driving expression in the testis

[37], and preferential retention of novel genes with male-biased expression [38]. Sexual

selection also drives rapid evolution of reproductive proteins [39] and could drive the emergence

of new genes as a mechanism of improving male reproductive ability [33]. The testis-biased

expression of novel genes, combined with growing evidence for new genes acting across a

variety of tissue contexts, suggests that many novel genes may function in male reproduction.

For example, a pair of young duplicate genes in *Drosophila, apollo* and *artemis,* are essential for

male and female fertility, respectively [40]. Continued efforts to identify and characterize testis-

expressed novel genes are consequently critical for understanding the genetic basis of male

reproductive phenotypes.

Drosophila serves as an ideal system for interrogating the prevalence, sequence

attributes, expression patterns, and functions of testis-expressed *de novo* genes. The

availability of well-annotated genomes for numerous *Drosophila* species, the tractability of flies

to molecular genetics techniques, and our thorough understanding of *Drosophila* reproductive

processes facilitate comprehensive analyses of novel fly reproductive proteins [41, 42]. As

observed in other biological systems, *Drosophila de novo* genes retained by selection

demonstrate enriched expression in the testis [10, 19, 33, 34]. The expression patterns of

emerging *de novo* genes in the *Drosophila* testis were recently analyzed at single cell resolution

[43], thereby providing insight into the dynamics of novel gene expression throughout

spermatogenesis. In addition to bioinformatic screens that have started to identify *de novo*

genes and large-scale expression analyses of testis-expressed genes, RNAi [44] and

CRISPR/Cas9-based [45] functional screens have identified putative, testis-expressed *de novo*

genes required for fertility. However, a need remains for in-depth experimental and evolutionary

characterization of the genes identified in such screens. Detailed examination of the function of

de novo proteins will enable us to understand how these proteins might integrate into existing

gene networks and become essential.

We previously conducted a pilot functional screen of *de novo* genes with testis-enriched

expression in *D. melanogaster* and identified two novel genes, *goddard* and *saturn*, that are

required for full fertility [46]. *Goddard* knockdown males failed to produce any sperm. *Saturn*

knockdown males produced fewer sperm, which were inefficient at migrating to female sperm

storage organs. Subsequent characterization of Goddard using null deletion alleles and a

biochemically tagged rescue construct showed that the protein localizes to elongating sperm

axonemes and that, in its absence, individualization complexes associate less efficiently with

spermatid nuclei and do not successfully progress along sperm tails [20]. These data

suggested that putative *de novo* genes can evolve essential roles in a rapidly evolving

reproductive process, spermatogenesis.

Here, we expanded this functional screen by evaluating whether any of 42 putative *de*

novo genes that show testis-enriched expression in *D. melanogaster* are required for male

fertility. We identified one gene, which we named *atlas,* whose knockdown or knockout results

in nearly complete male sterility. We found that *atlas* encodes a transition protein that facilitates

spermatid chromatin condensation. The *atlas* gene in *D. melanogaster* arose when a likely *de*

novo evolved protein-coding sequence moved off of the X chromosome and was inserted

upstream of a well-conserved non-coding RNA. While the *atlas* protein-coding sequence has

undergone multiple, independent gene loss events since its apparent origin at the base of the

Drosophila genus, the gene has evolved a critical function in *D. melanogaster*. These results

underscore the importance of detailed functional and evolutionary characterization in

understanding the origins of new protein-coding genes and the selective forces that affect their

subsequent evolution.

Results

An RNAi screen identifies a putative de novo *gene essential for* Drosophila *male fertility*

145   A previous pilot screen of 11 putative *de novo* evolved, testis-expressed genes identified

two genes that are critical for male fertility in *Drosophila melanogaster* [46]. This result, and

other recent work [e.g., 40, 47, 48], suggested that lineage-specific, newly evolved genes can

rapidly become important for fertility, perhaps by gaining interactions with existing protein

networks. To determine more comprehensively the frequency with which potential *de novo*

evolved genes become essential for fertility, we identified *de novo* or putative *de novo* evolved

genes with testis-biased expression. A previous computational analysis identified genes that

are detectable only within the *Drosophila* genus, lack identifiable protein domains, and show no

homology to other known proteins through BLASTP and TBLASTN searches [19]. We filtered

these genes to identify those expressed exclusively or predominantly in the testis, a common

site of *de novo* gene expression in animal species [27, 34, 38, 49]. This resulted in a set of 96

target genes.

Figure 1. An RNAi screen of putative *de novo* genes identifies *CG13541* as a major contributor to
***Drosophila melanogaster* male fertility.** A) All RNAi lines that showed at least partial knockdown of the
target gene were screened in group fertility assays (see Materials and Methods). Relative fertility was
calculated by dividing the average number of progeny produced per female mated to knockdown males
by the average number of progeny produced per female mated to control males in a contemporaneous
experiment. Relative fertility measurements lack error bars because each gene was tested in only 1-2
replicates. Knockdown of *goddard* was used as a positive control. B) A single-mating, single-pair fertility
assay confirms the observed defect when males are knocked down for *CG13541*, as knockdown males

showed significantly reduced fertility (control fertility (mean ± SEM): 109.0 ± 5.3; knockdown fertility: 0.2 ±
0.1; two-sample *t*-test assuming unequal variances, *p* = 5.6 x 10^{-13}).
168
We used testis-specific RNA interference to screen these genes for roles in male fertility.

We obtained RNAi lines from the Vienna *Drosophila* Resource Center (VDRC) and the

Transgenic RNAi Project (TRiP) and constructed additional lines using the TRiP-style

pValium20 vector [50], which induces efficient knockdown in the male germline. We tested an

RNAi line for each of 57 genes by using the *Bam*-GAL4 driver, which is expressed in the male

germline and which we enhanced with a copy of UAS-*Dicer2*. RT-PCR confirmed at least

partial knockdown in lines representing 42 genes (see example in Fig. S1). We then screened

knockdown males for fertility by allowing groups of 7 knockdown males to mate with 5 wild-type

females for 2 days. Progeny counts were standardized to the number of progeny produced by

concurrently mated groups of 7 control males and 5 wild-type females. The results are shown in

Fig. 1A. This initial screen identified *CG13541*, whose knockdown severely reduced male

fertility. We confirmed the result for *CG13541* by performing single-pair mating fertility assays

(Fig. 1B). Consistent with our previous convention of naming testis-expressed genes after

American rocketry [46], we will from here on refer to *CG13541* as *atlas*. While RNAi transgenes

designed to knockdown *CG43072* and *CG33284* caused full and consistently partial sterility,

respectively, we do not further consider these genes because subsequent gene knockout using

CRISPR genome editing indicated that neither gene plays a role in male fertility. In these

cases, the RNAi phenotypes might have been due to off-target knockdown.

CRISPR-mediated gene mutation validates atlas *RNAi results*

We validated the observed fertility defect by using CRISPR/Cas9-based genome editing

to construct putative loss-of-function alleles for *atlas* (Fig. S2). The principal allele we used for

validation and the functional studies described below was a null allele that completely deletes

the *atlas* genomic locus. This allele was generated by targeting each end of the locus with a

gRNA. We made three additional frameshift alleles by inducing double-stranded breaks at a

gRNA target site just downstream of the *atlas* start codon, which induced non-homologous end

joining. Males homozygous for the *atlas* deletion allele have the same fertility defect as

knockdown males (Fig. 2A). Males homozygous for any of three frameshift alleles showed

significantly reduced, but non-zero, fertility (Fig. 2B). It is possible that residual *atlas* function

may be present in these animals, perhaps due to translation initiation at a downstream start

codon, which could generate a shorter protein with partial function. Each frameshift allele

retains the possibility of encoding an N-terminally truncated, but otherwise in-frame, version of

Atlas protein that would lack the first 60 amino acids (out of 172; see Fig. S2). Alternatively, it is

possible that the residual fertility of the frameshift alleles is caused by the gene's intact 3' UTR,

a topic we discuss in more detail below. Finally, we constructed a genomic rescue construct

carrying both the full *atlas* transcribed region and its native regulatory sequences. *atlas* null

males that carried a single copy of the rescue construct had fully restored fertility (Fig. S3).

Overall, these data demonstrate that *atlas* loss, and not an RNAi or CRISPR off-target, causes

nearly complete male sterility.

Figure 2. CRISPR-generated deletion and frameshift alleles of *atlas* confirm the gene's
requirement for male fertility. A) Single-pair fertility assay for males homozygous for the null (Δ*atlas*)
allele or heterozygous controls (Δ*atlas*/SM6). Flies homozygous for the deletion had significantly reduced
fertility (control fertility: 82.9 ± 4.5; null fertility: 0.3 ± 0.6; two-sample t-test assuming unequal variances, *p*
= 5.4 x 10^{-18}). B) Single-pair fertility assays for males homozygous or heterozygous for three frameshift
alleles of *atlas* generated by imprecise non-homologous end joining at a CRISPR/Cas9 target site just

downstream of the start codon: *atlas*[52] (control fertility: 104.7 ± 3.8, mutant fertility: 11.2 ± 4.6; two-sample
t-test assuming unequal variances: $p = 8 \times 10^{-12}$), *atlas*[62] (control fertility: 96.2 ± 5.7; mutant fertility: 8.4 ±
2.9; two-sample *t*-test assuming unequal variances: $p = 6.1 \times 10^{-10}$) and *atlas*[86] (control fertility: 67.5 ± 6.6;
mutant fertility: 17.3 ± 5.8; two-sample *t*-test assuming unequal variances: $p = 9.5 \times 10^{-6}$).
219
220
Atlas *is required for proper spermatid nuclear condensation*

We next examined how *atlas* loss of function impacted male fertility at the cellular level.

Dissection and phase-contrast imaging of *atlas* deletion null or knockdown male reproductive

tracts revealed that while the pre-meiotic and meiotic stages of spermatogenesis appeared

normal, sperm accumulated at the basal end of the testes, rather than in the seminal vesicles

(SVs), over the first week of adulthood (Fig. 3A and Fig. S4). To further characterize the fertility

defects in the absence of *atlas*, we examined the Mst35Bb-GFP ("protamine"-GFP) marker in

null or knockdown backgrounds [51]. Mst35Bb encodes one of two protamine-like proteins

(highly similar paralogs of each other) that bind DNA in mature sperm. Its GFP fusion construct

thus allows visualization of nuclei in late stage spermiogenesis and mature sperm. Consistent

with the observed conglomeration of sperm tails at the basal testes, SVs from either *atlas* null or

knockdown males contained fewer mature sperm (Fig. 3B, Fig. S4C). The nuclei of sperm from

null males also appeared wider and less elongated than those of controls. Together, these data

suggest that *atlas* is required after meiosis, as developing spermatids take on their final

structures.

We next examined two post-meiotic processes: individualization of 64-cell spermatid

cysts into mature sperm, and spermatid nuclear condensation. Individualization initiates when

an actin-rich individualization complex (IC) associates with the bundle of spermatid nuclei. The

IC then proceeds down the sperm tails, expelling cytoplasmic waste and remodeling cell

membranes to form 64 individual sperm. We visualized this process in males 0-1 days old,

when spermatogenesis occurs at high levels, by staining whole mount testes for actin (Fig. 3C-

D). Although ICs associated with nuclear bundles present at the basal end of the testes in both

control and *atlas* null males, we observed significantly fewer nuclear bundle-associated ICs in

nulls (Fig. 3C-D). While control testes typically had several ICs progressing down sperm tails,

we saw a significantly reduced proportion of progressed bundles in nulls (Fig. 3C-D). In some

null testes, we also observed individual investment cones dissociated from progressing ICs (Fig.

3C).

The ability of ICs to assemble at nuclear bundles and progress down sperm tails may be

reduced if nuclear condensation is aberrant [reviewed in ref. 52]. During *Drosophila*

spermiogenesis, round spermatid nuclei undergo a series of stepwise, morphological changes

that are the product of two distinct, but related processes: changes in the chromatin packaging

of DNA, and changes in nuclear shape [53-55]. The end result is thin, condensed nuclei. We

quantified this process in testes dissected from newly eclosed wild-type and *atlas* null males

expressing Mst35Bb-GFP, which marks the final stages of condensation. We shredded the

post-meiotic region of the testes in the presence of a fixative and counted the number of nuclear

bundles that exhibited each of five stages of condensation [53]: round nuclei, early canoe-stage

(unmarked with Mst35Bb-GFP), late canoe-stage (marked with Mst35Bb-GFP), elongated

nuclei, and fully condensed nuclei (Fig. 4). Condensation of the nuclear bundles in *atlas* null

testes progressed at similar rates to controls through the late canoe stage (Table S1).

However, in *atlas* null males, all nuclear bundles that progressed past the canoe stage (which

included ~60% [range: 26-100%] of all observed bundles) showed an aberrant "curled"

phenotype (Fig. 4; Table S1). These data suggest that Atlas protein is required during the later

stages of nuclear condensation and are consistent with the idea that the loss of *atlas* affects

nuclear condensation in a way that reduces IC assembly and sperm individualization (see Fig.

3C).

Figure 3. Cytological investigations of the *atlas* mutant fertility defect. A) Phase contrast microscopy of male reproductive tracts dissected from 7-day-old, unmated control (*w*[1118]) or *atlas* null males. Control males show the expected accumulation of sperm tails in the seminal vesicle (SV), which appear here as a darker brown shading, while null male have an aberrant accumulation of sperm tails at the basal end of the testis (T). B) Visualization of Mst35Bb-GFP in 4-day old control and *atlas* null testes. While Mst35Bb localizes to spermatid nuclei in the absence of *atlas*, the nuclei appear shorter and much less numerous in the outlined SV. C) Representative images from phalloidin staining of *w*[1118] and *atlas* null testes used to assess the association of individualization complexes (ICs) with nuclear bundles and the progression of ICs down the length of sperm tails. D) At top, number of nuclear bundles with ICs associated in control and *atlas* null testes. Significantly more ICs were observed in control testes (control: N = 17, median = 14; mutant: N = 13, median = 7; Wilcoxon rank-sum test W = 34, p = 0.0014). At bottom, proportion of all observed ICs that were intact and that had progressed away from nuclear bundles. Three mutant testes with no observed ICs were excluded from the analysis. A significantly higher proportion of ICs progressed in control testes (control: N = 17, median = 0.27; control: N = 10, median = 0; Wilcoxon rank-sum test W = 28, p = 0.0038).

Figure 4. ***Atlas* null males show aberrant nuclear shaping at and beyond the elongated stage of spermatid nuclear condensation.** Early and late canoe stages were distinguished by the absence or presence of Mst35Bb-GFP, respectively. Late canoe and elongated stages were distinguished by the absence or presence of GFP-positive puncta, respectively. Condensed nuclei were distinguished from elongated nuclei by size. As shown in Table S1, nuclear bundles from *atlas* null testes consistently took on a curved shape after the canoe stage, though the degree of curvature was variable, as exemplified by the two examples of elongated nuclei from *atlas* null testes above.

That condensing spermatid nuclei are misshapen in the absence of *atlas* suggests the

possibility that Atlas protein is critical for nuclear condensation. This idea is further supported

by its predicted biochemical properties. Previously characterized spermatid chromatin binding

proteins are small and highly basic [53, 56, 57], as the excess of positively charged amino acid

side chains facilitates ionic interactions with negatively charged DNA. Many such proteins (i.e.,

Tpl94D, Mst35Ba, Mst35Bb, Prtl99C and Mst77F) also contain a conserved protein domain, the

high-mobility-group box (HMG-box) domain [55, 58-64], suggesting that this type of chromatin

binding protein could have originated through gene duplication and divergence. Consistent with

its putative *de novo* origin, Atlas lacks a detectable HMG-box domain. However, Atlas is

otherwise similar to these other sperm chromatin binding proteins: the ~20 kDa protein has a

highly basic predicted isoelectric point of 10.7, and its primary sequence contains the sequence

KRDK, which matches the canonical consensus sequence for nuclear import, K(K/R)X(K/R)

[65]. To test the hypothesis that Atlas is nuclear localized, and could thus bind DNA, we

generated an *atlas-GFP* transgene under UAS control and expressed it ubiquitously using

tubulin-GAL4 and in the early male germline using *Bam*-GAL4. In both larval salivary glands

and early male germline cells, Atlas-GFP appeared to be nuclear localized (Fig. S5).

While these results were consistent with Atlas protein localizing to the nucleus, they did

not allow us to visualize Atlas in the cells in which it is normally expressed. To do so, we used

CRISPR/Cas9-induced homology directed repair (https://flycrispr.org/scarless-gene-editing/)

[66-68] to create an *atlas*-GFP fusion at the endogenous *atlas* locus (see Fig. S6 and Materials

and Methods). We first confirmed the functionality of the knock-in allele by showing that males

with the *atlas* locus genotype *atlas*-GFP/Δ*atlas* had equivalent fertility to males of genotype

atlas+/Δ*atlas* (Fig. 5A). We then visualized Atlas-GFP fusion protein in whole-mount testes in

conjunction with phalloidin-stained actin (Fig. 5C and Fig. S7A). Atlas-GFP was absent from

seminal vesicles, consistent with its absence from the proteome of mature *D. melanogaster*

sperm [69, 70]. Instead, Atlas-GFP colocalized with condensing nuclear bundles near the basal

end of the testes (Fig. 5C). Actin-based ICs were also observed in the basal testes, but

generally did not co-localize with Atlas-GFP, suggesting that Atlas-GFP is present in

condensing nuclei before IC association (Fig. 5C). This result, taken together with the aberrant

nuclear condensation in the absence of *atlas* (Fig. 4), is consistent with the idea that Atlas is a

transition protein. Transition proteins are chromatin components that act transiently during

spermatid nuclear condensation. A series of transition proteins first replace histones as the

primary DNA binding proteins in the nucleus and then give way to protamines, the proteins that

package chromatin in mature sperm [53, 55, 58].

To further elucidate the role of *atlas* in nuclear condensation, we next examined Atlas-

GFP localization in the presence of either an early spermatid nuclear marker, histone H2Av-

RFP [55, 71], or Mst35Bb-dsRed [51], a marker of nuclei from the late canoe stage through final

condensation. Atlas-GFP showed no co-localization with H2Av-RFP, suggesting that Atlas

functions after histone removal (Fig. 5B). In contrast, some GFP-positive bundles co-localized

with Mst35Bb-dsRed, but others did not (Fig. 5D and Fig. S7B). These data suggest that Atlas

may be one of the final transition proteins used in nuclear condensation before the chromatin

becomes fully condensed with protamines.

Figure 5. An *atlas*-GFP allele generated at the endogenous *atlas* locus is fully functional for male fertility and encodes a protein that localizes to condensing spermatid nuclei. A) A single copy of the *atlas*-GFP allele is sufficient for full fertility when paired with the Δ*atlas* allele, as compared to males heterozygous for the wild-type *atlas* allele (Δ*atlas*/*atlas*-GFP fertility: 86.0 ± 3.2; Δ*atlas*/+ fertility: 84.9 ± 4.1; two-sample t-test assuming unequal variances, *p* = 0.85). B) Atlas-GFP does not co-localize with histone H2Av-RFP, a marker of the initial stages of spermatid nuclear condensation. C) Visualization of Atlas-GFP in the basal portion of whole-mount testes from *atlas-GFP* homozygotes shows that the fusion protein co-localizes with a subset of condensing spermatid nuclear bundles. While actin associates with fully condensed nuclei at the basal testis, Atlas-GFP does not overlap and is also absent from the seminal vesicle. D) Atlas-GFP partially colocalizes with Mst35Bb-dsRed, a marker of the final stage of nuclear condensation, in the basal portion of whole-mount testes. Open arrow: example of co-localization. Filled arrowhead: example of Atlas-GFP that does not co-localize with Mst35Bb-dsRed. Collectively, these data suggest that atlas may serve as a transition protein involved in the final stages of nuclear condensation. The whole testes from which the basal portions are shown in panels C and D are shown in Fig. S7.

To determine the stage(s) of nuclear condensation at which *atlas* functions, we analyzed

the shape of fixed nuclear bundles from shredded testes isolated from *atlas*-GFP males on the

day of eclosion. Based on the stage of the defect in *atlas* null males (Fig. 3-4) and the pattern

of Atlas-GFP-positive bundles in whole-mount testes (Fig. 5), we hypothesized that Atlas-GFP

would localize to the later stages of nuclear condensation. Consistent with this hypothesis, we

did not detect Atlas-GFP in round or early canoe stage bundles (Fig. 6A-B). Atlas-GFP co-

localized with DNA in late canoe stage bundles (Fig. 6C). Interestingly, when nuclei elongated

further, GFP was detected not in the nucleus, but as puncta basal to the nuclei (Fig. 6D; see

also Fig. S7A). Since Atlas-GFP is not observed in mature sperm in the SV (Fig. 5C), these

data suggest that Atlas may function as a transition protein that facilitates the condensation of

spermatid nuclei from histone-based DNA packaging to protamine-like-based DNA packaging

[53] and is then removed from nuclei once protamines bind DNA. Indeed, the appearance of

Atlas in nuclei during the late canoe stage of condensation is similar to the pattern observed for

a previously characterized transition protein, Tpl94D [53]. We hypothesize that the failure of

atlas null sperm to form needle-like nuclei can be explained by the absence of Atlas from the

late canoe nucleus. It is also possible that the apparent removal of Atlas-GFP from nuclei (Fig.

6D) represents a mechanism for removing transition proteins from the nucleus after they exert

their functions. We observed above that some Mst35Bb-GFP also appears to be removed from

the nucleus in puncta during the elongation stage of nuclear condensation (see elongated stage

of control nuclear bundles, Fig. 4), even though other Mst35Bb-GFP molecules ultimately

package DNA in mature, individualized sperm. This could occur if Mst35Bb-GFP is present in

excess of what is needed to package DNA.

Figure 6. Atlas-GFP is present in late canoe-stage spermatid nuclei and then appears to leave the nucleus in puncta. Staging of condensing spermatid nuclei fixed in paraformaldehyde from *atlas*-GFP males stained with TO-PRO-3 DNA stain. Atlas-GFP is not detectable in (A) round stage or (B) early canoe stage nuclei. Atlas-GFP is nuclear localized in the late canoe stage (C). When nuclei become fully elongated (D), puncta of Atlas-GFP appear to be removed from the nucleus.

Evolutionary origins of atlas

To better understand the evolutionary origin of *atlas* and its evolution since emergence,

we used a combination of BLAST- and synteny-based approaches to identify *atlas* orthologs

throughout the genus [46, 72]. One notable feature of this two-exon gene is that the protein-

coding region (519 nucleotides) is contained entirely within the first exon (622 nt); the longer,

second exon (910 nt) appears to be entirely non-coding (Fig. 7). Surprisingly, the second exon

is more widely conserved. BLASTN detected significant matches to this region (range of hit

length: 185-864 nt) in the same genomic location on Muller element C [see ref. 73 for

explanation of Muller elements], as assessed by synteny, in all *Drosophila* species examined,

including distantly related species such as *D. virilis* and *D. grimshawi* (Fig. 7 and S8). The

protein-coding first exon shows a more limited phylogenetic distribution. In most members of

the *melanogaster* group of *Drosophila* (gray box in Fig. 7), this exon is found in a conserved

position, adjacent to the non-coding region on the equivalent of *D. melanogaster* chromosome

2R (Table S2). In *D. ananassae*, however, the protein-coding region is found on the X

chromosome (Muller element A). A putative ortholog for the protein-coding sequence is

detectable by BLASTP in *D. virilis* in a partially syntenic region on the same Muller A element

(Table S2, Fig. S9). These data suggest that the *atlas* protein-coding sequence initially arose

on Muller element A and then moved to Muller element C, giving rise to the gene structure

observed in extant *D. melanogaster* and its sister species.

Figure 7. Molecular evolution and gene expression of *atlas* across the *Drosophila* genus. The gene structure of *atlas* in *D. melanogaster* is shown at top left. The predicted protein-coding sequence is contained entirely within exon 1, while exon 2 encodes the presumed 3' UTR. The gene is located on chromosome 2R, equivalent to Muller element C. The phylogeny shows BLAST- and synteny-based detection of sequences orthologous to the protein-coding sequence and the 3' UTR sequence across *Drosophila* species. Sex-specific adult RNA-seq data were used to assess male expression across species, with RT-PCR verification performed in species marked with asterisks. RNA-seq data for the syntenic region of the 3' UTR in *D. virilis* were ambiguous; see Figs. S8 and S10.

To confirm the lack of *atlas* protein-coding sequences identifiable by BLASTP or

TBLASTN in most non-*melanogaster* group species, we identified the regions syntenic to those

containing *atlas* in *D. ananassae* and *D. virilis* in 11 additional *Drosophila* species and used

more sensitive methods to search for potential orthologs [72]. Specifically, we: a) relaxed the

BLAST cut-offs for detection, since default parameters can cause false-negative results when

searching for potential *de novo* genes in divergent species [74]; b) used adult male RNA-seq

data to detect transcribed areas within each syntenic region that did not match annotated

genes; and, c) predicted the isoelectric point of the potential proteins encoded, under the

hypothesis that Atlas orthologs would have conserved, DNA-binding functions. The results are

summarized in Fig. S9 and Table S2. These searches detected no evidence for *atlas* orthologs

in the following *Drosophila* species: *obscura, miranda, willistoni, hydei, arizonae, mojavensis,*

navojoa, and *grimshawi*. Some of these species have unannotated, male-expressed transcripts

in the regions syntenic to the Muller A location of *atlas* in *D. ananassae* and *D. virilis*, but when

each was compared with BLASTP to the *D. melanogaster* proteome, all matched proteins other

than Atlas, suggesting they may be lineage-specific paralogs of other genes (Fig. S9). In sister

species *D. pseudoobscura* and *D. persimilis*, we detected a male-expressed transcript predicted

to encode a protein with a pI > 10 in the region syntenic to the location of *atlas* in *D. virilis*, but

the predicted protein sequences showed no significant BLASTP similarity to *atlas* orthologs

(Table S2). While this predicted protein may represent a divergent *atlas* ortholog, the abSENSE

method predicts low probabilities of BLASTP detection failure when searching for Atlas protein

in these species (0.02 and 0.04, respectively), so we favor the hypothesis of a lineage-specific,

newly evolved gene in the region. Conversely, in *D. busckii*, we detected, in the region syntenic

to the *D. virilis atlas* locus, a male-expressed transcript predicted to encode a protein with

significant BLASTP identity to *D. melanogaster* Atlas (e = 4e-8), but with a predicted pI of 5.1

and a ~50 percent shorter open-reading frame (Table S2). The ortholog status of this predicted

protein is also unclear, but because of its dramatically altered size and pI, it is unlikely to have a

functional role equivalent to that of *D. melanogaster* Atlas.

To investigate whether the protein-coding region may have reproductive functions in

other species, we used sex-specific RNA-seq data from numerous *Drosophila* species curated

by the Genomics Education Partnership [72; thegep.org] and verified several of these results by

RT-PCR (Fig. 7, Fig. S10). In all species in which *atlas* was detected, the protein-coding region

is expressed specifically in males regardless of its genomic location (Fig. 7). Interestingly, the

non-coding region shows male-specific expression in species lacking an unambiguous,

orthologous coding region, such as *D. pseudoobscura* and *D. mojavensis*. Conversely, while *D.*

yakuba and *D. erecta* express the protein-coding region robustly, we found no RNA-seq

evidence to support expression of the non-coding second exon, in spite of its sequence

conservation (Fig. S8). Based on its high level of sequence conservation, consistent genomic

location and expression in a variety of species, it is possible that what we now consider to be

the 3' untranslated region of *atlas* from *D. melanogaster* was, ancestrally, a non-coding RNA.

The FlyBase database reports two transcript isoforms of *atlas* in *D. melanogaster*: the

atlas-RA isoform is 986 nucleotides, while the *atlas*-RB isoform is 1528 nt. These isoforms

differ in how much of the second, non-coding exon is included in the transcript. We used RT-

PCR of whole male cDNA to assess the presence of these isoforms and their relative

abundances. Primers designed to amplify a region present in both isoforms produced products

that appeared more abundant than primers designed to amplify only the long isoform, even

though both primer pairs appeared to amplify genomic DNA with equal efficiency. Based on

RT-PCR band intensities and controlling for product size and genomic PCR band intensities, we

estimated that the short isoform is about 3-fold more abundant. This difference in abundance is

mirrored in available RNA-seq data, which show approximately 3- to 4-fold higher levels of

expression in the upstream part of exon 2 (Fig. S8), a pattern that also appears in *D. simulans*

and *D. sechellia*. Evaluating the potential significance of this finding awaits functional

characterization of the non-coding region.

As we have observed for other putative *de novo* genes with essential male reproductive

functions [46], the pattern of *atlas* protein-coding sequence presence/absence across the

phylogeny is difficult to explain parsimoniously. If we assume that gene birth events are less

frequent than gene deaths, since the latter can occur through many possible mutational events

and can happen separately along multiple phylogenetic lineages, our data support the

hypothesis of a single origin of the protein-coding sequence at the base of the genus, followed

by independent loss events on the lineages leading to *D. grimshawi*, *D. mojavensis* and *D.*

willistoni, and potentially also *D. pseudoobscura*/*persimilis* (Table S2). We summarize these

findings for 12 representative species of *Drosophila* in Fig. 7. The general patterns of loss do

not change when all species of Table S2 are considered, though an additional loss in the

melanogaster group is likely due to the absence of a detectable ortholog in *D. kikkiwai* and *D.*

serrata. As noted above, the pattern of gene loss can also appear due to orthology detection

failure [74], for which we tried to account with our additional search methods described above.

We also note, however, that the probability of BLASTP-based ortholog detection failure is

relatively low for some *Drosophila* species that lack *atlas*, including *D. pseudoobscura*

(probability of non-detection due to divergence = 0.02), *D. persimilis* (p = 0.04) and *D. willistoni*

(p = 0.06). The probability is higher for other species, *D. mojavensis* (p = 0.33) and *D.*

grimshawi (p = 0.66), underscoring the importance of our additional search strategies. Overall,

our data support the hypothesis of multiple, independent loss events within *Drosophila*.

AbSENSE produces a 1.00 probability of BLASTP-based Atlas ortholog detection failure

outside of *Drosophila*, reflecting the protein's short length and relatively rapid divergence (see

below). Indeed, the protein-coding and non-coding transcriptomes from each species showed

no matches to Atlas protein or cDNA sequences by BLAST. We thus used another synteny-

based approach, summarized in Fig. S11, to look for the protein-coding gene in other Dipterans

with well-resolved genomes: *Musca domestica*, *Glossina morsitans*, *Lucilia cuprina*, *Aedes*

aegypti, *Anopheles darlingi*, *Anopheles gambiae*, *Culex quinquefasciatus* and *Mayetiola*

destructor. In none of these species was a putative homolog found in any potential syntenic

region.

Recognizing the limitation of even this approach, we also used HMMER [75] to search

iteratively either all genomes in ENSEMBL, or all metazoan genomes in ENSEMBL, for

annotated proteins with identity to Atlas from *D. melanogaster* or *D. virilis*. These searches

initially identified significant hits to the Atlas orthologs we identified above from other *Drosophila*

species. When these collections of orthologs were used as queries, no further proteins outside

of *Drosophila* were a significant match. As a control, we performed the same search strategy

with *D. melanogaster* Mst35Bb, a protein whose length, amino acid composition, and function

are similar to Atlas. These searches readily identified orthologs throughout Diptera, consistent

with predictions of its conservation from the OrthoDB database [76]. We also note that a

previous HMMER-based analysis to identify HMG-box-containing spermatid chromatin

condensation proteins in *Drosophila* and other related insects did not detect *atlas* as a member

of the gene family [64], further diminishing the possibility that *atlas* is a divergent paralog of

other transition and protamine-like proteins. Thus, we conclude that *atlas* is a putative *de novo*

evolved gene that is limited phylogenetically to the *Drosophila* genus.

Finally, we used standard tests of molecular evolution to examine the selective

pressures that have shaped Atlas protein within the *melanogaster* group. We aligned the *atlas*

protein-coding sequences from 12 species and used PAML to ask whether a model (M8)

allowing for positive selection, as well as neutral evolution and purifying selection, explained the

data better than models (M7 and M8a) that allowed only neutral evolution and purifying

selection [77, 78]. These data showed that while the *atlas* protein-coding sequence's rate of

evolution was accelerated relative to most *Drosophila* proteins (whole-gene estimated d_N/d_S, ω =

0.41 by PAML model M0), there was no significant evidence for positive selection acting to

recurrently diversify a subset of sites within the protein (Table 2).

Table 2. PAML sites tests for positive selection acting on *atlas* in the *melanogaster*
group.

PAML Model	ω estimate	ln L	Number of parameters	Likelihood ratio test with M8
M0 (uniform ω across all sites)	0.41	-4032.24	23	n/a
M7 (10 site classes, each with $0 \leq \omega \leq 1$)	n/a	-3961.28	24	χ^2 = 1.74, df = 2, *p* = 0.42
M8a (10 site classes as in M7, plus one class with ω = 1)	n/a	-3961.02	25	χ^2 = 1.22, df = 1, *p* = 0.27
M8 (10 site classes as in M7, plus one class with $\omega \geq 1$)	1.39 (9.1% of sites)	-3960.41	26	n/a

<u>Discussion</u>

Across taxa, many *de novo* evolved genes are expressed in the male reproductive

system [12, 31, 33, 34]. Identifying those genes that have evolved essential roles in

reproduction will provide insight into how newly evolved genes integrate with existing cellular

networks [47] and how evolutionary novelties permit adaptation in the face of sexual selection.

Here, we screened 42 putatively *de novo* evolved genes for major effects on male *D.*

melanogaster reproduction. Our primary screen identified three genes whose knockdown

caused an apparent reduction in male fertility. However, subsequent CRISPR-mediated gene

deletion revealed that only one of these genes, *atlas*, was truly essential. This result

underscores the importance of validating genes identified in RNAi screens through traditional

loss-of-function genetics and other approaches.

Loss *of atlas* function reduces fertility by affecting mature sperm production. During

spermiogenesis, *atlas* mutants show aberrant nuclear condensation and an inability to

individualize spermatid bundles successfully. GFP-tagged Atlas protein localizes to condensing

spermatid nuclei in the basal testis and partially co-localizes with Mst35Bb, a protamine around

which DNA is wrapped in mature, individualized sperm. Evolutionary analysis showed that the

atlas protein-coding sequence likely arose at the base of the *Drosophila* genus but was unlikely

to have played an essential role immediately upon birth, as the gene was subsequently lost

along several independent lineages. Within the *melanogaster* group of *Drosophila*, however,

the gene moved from the X chromosome to an autosome, where it formed a single

transcriptional unit with a conserved, non-coding sequence. Since this point, the gene has

encoded a protein with a conserved length, isoelectric point and male-specific expression

pattern, suggesting potential functional conversation over the last ~15 million years.

Comparison of atlas *null and frameshift allele phenotypes*

547   The null deletion and frameshift alleles of *atlas* all caused significantly reduced fertility

(Fig. 2), but the deletion allele resulted in essentially complete sterility, while residual fertility

remained in males homozygous for each frameshift allele. We noted above and depicted in Fig.

S2 how the frameshift alleles have the potential to encode an N-terminally truncated form of

Atlas that would contain amino acids 61-172 of the wild-type protein, if these alleles allow

translation initiation at the methionine-encoding codon 61. Such a truncated protein would be

35 percent shorter than wild-type, lack the predicted nuclear localization sequence, and have a

reduced isoelectric point of 7.1. Each of these factors could contribute to reduced functionality.

555   The broad sequence conservation of the 3' UTR across the *Drosophila* genus, including

in species that lack the *atlas* coding sequence, suggests the alternative possibility that the 3'

UTR also contributes to fertility. In one scenario, the 3' UTR could act as part of the *atlas* locus

by regulating RNA stability and/or the spatiotemporal control of Atlas protein translation. An

alternative possibility is that the 3' UTR functions on its own as a non-coding RNA. Indeed,

since the protein-coding region and the *D. melanogaster* 3' UTR were initially separate genetic

entities (and continue to be in *D. ananassae* and *D. virilis*), it is possible that each contributes to

fertility in a unique way. Future experiments that could distinguish these possibilities are

described below.

564   Based on the current evidence, however, we think that the primary way in which *atlas*

impacts fertility is through its protein-coding sequence. This conclusion is supported by: the

~60-90 percent reduction in fertility in even the frameshift mutants; the stability of the Atlas-GFP

fusion protein and its presence in the spermatid nuclear condensation stages that immediately

precede the timing of the null mutant phenotype; and, the observation that the protein-coding

region shows a more highly conserved expression pattern than the UTR region within the

melanogaster group (Fig. 7 and Fig. S8).

Atlas is an essential transition protein

Several lines of evidence suggest that Atlas is a transition protein that facilitates the

change from histone-based to protamine-based chromatin packaging in spermatid nuclei. Atlas

localizes throughout spermatid nuclei (Fig. 6) and has biochemical properties consistent with

direct DNA interaction. The protein appears specifically at the late canoe stage of nuclear

compaction (Fig. 5B-D). Its lack of overlap with testis-specific histones (Fig. 5B), partial overlap

with Mst35Bb (Fig. 5D), removal from needle-stage nuclei (Fig. 6) and absence from mature

sperm (Figs. 5-6) are all consistent with the expression profile of a transition protein. Several

other transition proteins have been characterized in *D. melanogaster*, including Tpl94D, thmg-1,

thmg-2, and Mst84B [53, 58, 59]. Collectively, the transition proteins vary in the stage of

nuclear condensation at which they first appear and the range of nuclear shapes over which

they are found [79], but otherwise match Atlas in their biochemical properties, transient

expression, and localization throughout the nucleus. Compared to these other transition

proteins, Atlas is present over a fairly narrow range of nuclear condensation stages and reaches

its peak expression just prior to the onset of individualization. *Atlas* is also the only transition

protein gene characterized to date whose removal disrupts fertility, as *Tpl94D*, *thmg-1*, *thmg-2*

and *Mst84B* mutants are all fertile [53, 58, 59]. This may reflect the relatively later timing of

Atlas's expression in spermatid nuclei, reduced functional redundancy between DNA-binding

proteins at the later stages of condensation, a potential interaction between Atlas and an

essential protamine-like protein, and/or a more stringent requirement for DNA binding at these

stages.

Transition proteins give way in spermatid nuclei to protamine-like proteins, which bind

DNA in mature sperm and persist through fertilization. In this way, protamine-like proteins

function analogously to vertebrate protamines, though they are believed to be evolutionarily

independent [57, 64]. In *D. melanogaster*, protamine-like proteins include Mst35Ba, Mst35Bb,

Prtl99C and Mst77F [63, 80, 81]. Interestingly, while all characterized protamine-like proteins

are present in mature sperm, only some are essential for fertility. Knockouts of *Mst35Ba*,

Mst35Bb, or both show occasional nuclear shaping defects, but male fertility is normal [80,

82]. In contrast, mutants of *Prtl99C* or *Mst77F* are sterile. Prtl99C and Mst35Ba/b bind

condensed DNA independently of each other and contribute additively to the shortening of

needle-stage nuclei, but Prtl99C's effect is ~3-fold greater [63, 64]. This difference is apparently

great enough to reduce fertility only in *Prtl99C* mutants. In contrast, *Mst77F* and *Mst35Ba/b*

show a genetic interaction, as *Mst35Ba/b* null flies become nearly sterile in an *Mst77F*

heterozygous background [81]. Furthermore, while Mst35Bb-GFP is expressed in *Mst77F* nulls,

these flies show deformed spermatid nuclei that do not reach a recognizable needle-like stage.

Because *atlas* nulls show considerable phenotypic similarity to *Mst77F* nulls, but not *Prtl99C*

nulls, we hypothesize that Atlas may act in a pathway with Mst77F. Our observation of

inefficient IC movement down sperm tails in *atlas* null testes is reminiscent of a similar

phenotype in *Mst77F* nulls [81], providing further evidence that these proteins may act in a

common pathway. In both cases, ICs can form at misshapen canoe-stage nuclei, but fail at a

subsequent step. While the exact relationship between nuclear abnormalities in the late canoe

stage and individualization is not entirely understood, it is possible that nuclear shape and the

organization of nuclear bundles impact the ability of IC association and IC progression, as is

also observed in mutants of another gene, *dPSMG1*, which controls nuclear shape [59].

Evolution of atlas *and other de novo genes*

618   Because *de novo* genes emerge from non-coding sequences, they typically encode

proteins that are short and lack complex structure [3, 16]. Indeed, expanding the length of the

protein-coding region and evolving higher-level protein structures are hypothesized to be among

the final stages of new gene evolution [4]. In light of these constraints, what kinds of cellular

functions might be available to newly evolved proteins? Vakirlis et al. [83] overexpressed

emerging proto-genes in *S. cerevisiae* and found that those encoding proteins with predicted

transmembrane (TM) domains were more likely to be adaptive, as assessed by the effect of

proto-gene overexpression on growth rate. Such proteins may arise when thymine-rich

intergenic regions undergo mutations that allow protein-coding gene birth and expression, since

many codons with multiple U nucleotides encode amino acids commonly found in TM domains

[83]. Our imaging data, in addition to the prediction tools employed by Vakirlis et al. [83],

suggest that Atlas does not contain a TM domain. However, just as the amino acid

compositional requirements of a TM domain are not overly complex, neither are those of DNA

binding proteins. In essence, these proteins must simply be small, have a high concentration of

positively charged residues, and contain a nuclear localization signal, which itself requires a

small patch of positively charged residues [65]. Thus, DNA binding proteins may be a relatively

easy class of protein to evolve *de novo*.

While many putative *de novo* genes are expressed in the *D. melanogaster* testis [Fig. 1A

and ref. 19], *atlas* was the only verified hit from our screen that was essential for male fertility.

This result raises two related questions. First, why are so many of these other genes expressed

if their knockdown causes no obvious effect? In general, it is common for the knockdown of

protein-coding genes expressed in reproductive tissues in *D. melanogaster* to result in no

detectable fertility defects [46, 84-86]. One set of explanations center on technical issues: a

gene could be expressed outside of the cells targeted by the RNAi driver, knockdown level

could be insufficient to cause a phenotype, and/or a gene may be functionally redundant such

that its knockdown causes no apparent effect. Another hypothesis to explain this pattern is that

while the loss of function of such genes may cause small reductions in fertility that would be

subject to strong negative selection in nature, the conditions used to assay such knockdown

animals in primary screens are rarely tailored to detect differences of this magnitude. A third

possibility, not mutually exclusive with the above, is that while some genes may be expendable

in non-competitive, non-exhaustive mating conditions, their absence may result in lower fitness

in sperm competitive environment, environments in which males mate several times in quick

succession, or environments in which sperm must persist in storage for longer intervals or

during less optimal conditions [87-89]. These latter two possibilities are illustrated by some of

the other spermatid chromatin binding proteins previously characterized as "non-essential" (e.g.,

Mst35Ba/Bb and *Tpl94D*) [58, 80]. While these proteins are not *de novo* evolved, they provide

examples of genes whose mutations cause aberrant cellular phenotypes, such as the

abnormally shaped spermatid nuclei in *Mst35Ba/Bb,* and that may contribute to fertility defects

when placed in a sensitized genetic background [81].

It is also possible that some of the genes we screened are *de novo* genes that have no

functional effects and have thus never been selectively advantageous. For example, 8 of the 42

screened genes have annotated orthologs in either *D. melanogaster* alone or in only *D.*

melanogaster and *D. simulans*. Such recently "born" *de novo* genes could have been

propagated by neutral evolutionary processes but not yet experienced inactivating mutations

that would abolish their coding sequences and/or expression. This pattern would be consistent

with *de novo* genes' high rates of both birth and death observed previously in *Drosophila* [38]

and yeast [83].

A second question raised by our finding that *atlas* encodes an essential transition protein

is: how might *atlas* have evolved to become essential for fertility in *D. melanogaster*, particularly

when other transition proteins appear functionally redundant? Other proteins involved in

spermatid chromatin compaction show variable levels of conservation across *Drosophila*. For

example, the protamines that contribute to DNA packaging in mature sperm [57] are found

across all sequenced *Drosophila* species [61], and orthologs are also reported in FlyBase from

other Dipteran and non-Dipteran insects [90]. However, transition protein Tpl94D is reported to

be restricted to species ranging from *D. melanogaster* to *D. pseudoobscura* [60], as are the

related proteins tHMG1 and tHMG2 with high-mobility group domains [58, 90]. Results like

these suggest that while some protamine-like proteins (i.e., Mst35Ba and Mst35Bb) have

consistently been among the final chromatin-packaging proteins, the specific proteins facilitating

the transition from histones to protamines have likely varied over evolutionary time. Against this

backdrop, and based on our analyses of the protein's presence/absence, biochemical

properties, and expression patterns in extant species, we hypothesize that while the Atlas

protein likely had some DNA-binding ability and male-specific expression upon its origin, it was

only one of several proteins involved in spermatid chromatin compaction. Since *atlas* was lost

independently in several lineages after its birth (Fig. 7), Atlas was likely non-essential at its

outset, but rather evolved an essential function within the *melanogaster* group of species. Such

evolution of essentiality could have occurred because of the loss of a protein with a

complementary function and/or changes in the process of spermiogenesis that thrust Atlas into

a functionally unique role. It is also worth noting that species that have evidently lost *atlas* might

have undergone other compensatory changes in their repertoires of spermatid DNA binding

proteins. For instance, *D. willistoni* lacks *atlas* but appears to have several additional paralogs

of the protamines found only in duplicate in *D. melanogaster*, which could have evolved

transition-protein-like roles.

While our study cannot establish whether the movement of the *atlas* protein-coding

sequence off of the X chromosome onto an autosome early in the evolution of the *melanogaster*

group (Fig. 7) affected the gene's essentiality, such movement remains noteworthy. Prior work

has found a significant dearth of testis-expressed genes on the X chromosome in *Drosophila*

[69, 91-93] and other species [94, 95]. Furthermore, *Drosophila* exhibit suppression of X-linked

testis-expressed genes, and transfer of such genes from the X chromosome to autosomal loci

results in higher expression levels [96-98]. One of several proposed mechanisms for both the

paucity of X-linked testis-expressed genes and the suppression of their expression is meiotic

sex chromosome inactivation (MSCI), in which the X chromosome becomes transcriptionally

silenced earlier than autosomes [99-104]. Thus, genes that affect meiotic or post-meiotic

processes, as *atlas* does, could exert beneficial effects more strongly and/or for a longer period

of time if they become encoded autosomally. While the *atlas* protein-coding sequence appears

to show male-specific expression regardless of its chromosomal location, it is possible that the

movement of *atlas* to chromosome 2 allowed it to evolve a broader or different expression

pattern that expanded or modified its role in spermiogenesis. The complex molecular bases of

both X suppression and "escape" from the X chromosome in *Drosophila* continue to be actively

investigated and debated [102-108], but continued research in this area might inform further

interrogation of the forces driving *atlas* off of the X chromosome.

The movement of the *atlas* protein-coding sequence to chromosome 2 also created the

two-exon gene observed in *D. melanogaster*, in which the longer second exon appears to be

entirely non-coding. This second exon is highly conserved across the genus in both sequence

and genomic location, and it shows male-specific expression in several species that lack the

protein-coding sequence upstream (Fig. 7 and Fig. S8). These patterns of conservation

suggest that the second exon might originally have been a non-coding RNA, a class of molecule

whose importance in *Drosophila* male reproduction has recently become recognized [109, 110].

While previous examples of functional ncRNAs in spermatogenesis have generally acted in

trans to regulate other genes or affect the functions of other proteins, it is also possible that the

long 3' UTR of *atlas* in *D. melanogaster* could affect the translation of *atlas* transcripts. Many

genes functioning in spermatid differentiation are transcribed early in spermatogenesis but

translationally repressed until later in spermiogenesis, a phenomenon that relies on various

forms of post-transcriptional regulation [111, 112]. Future studies of the *atlas* protein-coding

sequence in the absence of its 3' UTR, the expression patterns of Atlas protein in species in

which it is encoded from the X chromosome, or the genetic ablation of the conserved region in

species lacking the protein-coding sequence will provide additional insights.

A final issue raised by our results is the exact timing and mechanism of origin for the

atlas protein-coding sequence. The bioinformatic screen [19] that identified *atlas* and the other

genes tested in Fig. 1 was designed to identify both "*de novo*" genes, defined as protein-coding

regions in *Drosophila* that had recognizable, but non-ORF-maintaining, TBLASTN hits in

outgroup species, and "putative *de novo*" genes, which had no TBLASTN hits in outgroup

species. (Importantly, the screen also eliminated any protein with an identifiable protein

domain, thus reducing the chances of identifying divergent members of gene families.) The vast

majority of the genes we tested with RNAi, including *atlas*, fell into the putative *de novo*

category. The bioinformatic screen's criteria were reasonable for a high-throughput analysis,

but BLAST-based methods have known limitations for detecting orthologous sequences in

diverged species [74, 113, 114]. The lack of identifiable *atlas* protein-coding genes in several

Drosophila species (e.g., *D. pseudoobscura* and *D. willistoni*) is unlikely to be due to BLAST

homology detection failure, and extensive synteny-based searches confirmed the gene's

absence (Fig. S9). BLAST and synteny-based searches for orthologs in non-*Drosophila*

species also did not detect an ortholog, though BLAST searches are not predicted to have

adequate sensitivity for a protein of this size and evolutionary rate, at this level of species

divergence [74]. Hence, in addition to using synteny to search for orthologs, we used HMMER,

which employs hidden Markov models and builds a sequence profile of the target protein using

information from multiple orthologs. Since HMMER also did not detect orthologs outside of

Drosophila, we hypothesize that *atlas* evolved *de novo* at the base of the genus. However,

since we remain unable to identify the non-protein-coding sequence from which *atlas* arose, we

continue to refer to *atlas* as a putative *de novo* gene [5].

Overall, we find that while many putative *de novo* evolved genes are expressed in the *D.*

melanogaster testes, few have major, non-redundant effects on fertility. However, several such

genes have evolved critical roles at distinct stages of spermatogenesis and sperm function. We

showed previously that the putative *de novo* gene *saturn* is required for maximal sperm

production, as well as for the ability of transferred sperm to migrate successfully to sperm

storage organs in females [46]. Another putative *de novo* gene, *goddard*, is required for sperm

production and encodes a cytoplasmic protein that appears to localize to elongating axonemes

[20, 46]. Loss of *goddard* impairs the individualization of spermatid bundles [20], thus exerting

an effect that appears to be upstream of those observed for *saturn* and *atlas*. Here, we report

another novel function for a putative *de novo* gene: encoding an essential transition protein that

is necessary for proper nuclear condensation in spermiogenesis. Taken together, these results

demonstrate that while many *de novo* genes may play subtle roles or share functional

redundancy with other genes, *de novo* genes can also become essential players in complex

cellular processes that mediate successful reproduction.

Materials and Methods

RNA Interference Screen

De novo and putative *de novo* genes inferred to be no older than the *Drosophila* genus

were identified previously [19]. We filtered these genes with publicly available RNA-seq data

[115] to identify those expressed predominantly in the testes [>50% of RPKM sum deriving from

the testes from ModENCODE data; ref. 115], giving a total of 96 genes. To assess each of

these candidates for effects on male fertility, we induced knockdown in the male germline by

crossing UAS-RNAi flies to Bam-GAL4, UAS-Dicer2 flies [46, 116]. Control flies were generated

by crossing the attP-containing genetic background into which UAS-RNAi was inserted to the

same GAL4 line. Flies carrying UAS-RNAi were of two types. Roughly half of the genes had

publicly available lines from the Vienna *Drosophila* Resource Center [117] or the Transgenic

RNAi Project [50]. For the other genes, no publicly available RNAi stock was available, so we

constructed TRiP-style stocks in the pValium20 vector as previously described [85]. These

constructs were integrated into an AttP site in stock BL 25709 (y^1 v^1 P{nos-phiC31\int.NLS}X;

P{CaryP}attP40) from the Bloomington *Drosophila* Stock Center (injections by Genetivision;

Houston, TX, USA) and crossed into a *y v* background to screen for v^+. We attempted at least

two rounds of transgenic production for each gene. In total, we were able to obtain and test

RNAi lines for 57 of the 96 identified genes. Table S3 shows all RNAi lines used and lists the

short hairpin sequences cloned for the TRiP lines we constructed.

We initially screened males knocked down for each candidate gene for major fertility

defects by crossing groups of 7 knockdown or control males to 5 virgin Canton S females,

letting the adults lay eggs for ~48 hours, and then discarding adults and quantifying the resulting

progeny by counting the pupal cases, as previously described [46]. To assess the degree of

knockdown achieved, 10 whole males of each line were homogenized in TRIzol reagent (Life

Technologies, Carlsbad, CA). RNA isolation, DNAse treatment, cDNA synthesis and semi-

quantitative RT-PCR with gene-specific primers were performed as previously described;

amplification of *RpL32* was used as a positive control [46]. We evaluated knockdown efficiency

by agarose gel electrophoresis of RT-PCR products as one of four levels: "complete" if no

product from the knockdown cDNA sample was visible via agarose gel electrophoresis; "near

complete" for a very faint knockdown product that was also much less abundant than the control

product; "partial" for a more robust knockdown product that was still visibly less intense than

control; and "not knocked down" if the product intensity for the knockdown sample equaled or

exceeded that of the control. Any gene that did not show at least partial knockdown was

discarded from further analysis, leaving a total of 42 genes successfully screened. Table S3

shows the degree of knockdown achieved for each line.

CRISPR Genome Editing

To validate RNAi results for *atlas*, *CG43072* and *CG33284*, we used CRISPR/Cas9

genome-editing to generate null alleles that could be used for further analysis, as described

previously [20]. Briefly, our general strategy was to design gRNAs in the pU6.3 vector

(Drosophila Genome Resource Center (DGRC) #1362) that targeted each end of a locus.

These plasmids, along with plasmids encoding gRNAs that targeted the *w+* locus, were co-

injected by Rainbow Transgenics (Camarillo, CA) into embryos laid by *vasa*-Cas9 females in a

w+ background, Bloomington stock #51323 [118]. G_0 animals were crossed to *w-* flies, and

members of G_1 broods with a higher-than-expected fraction of *w-* progeny were individually

crossed to an appropriate balancer line and then PCR-screened for the desired deletion of the

targeted locus.

We also constructed three frameshift, expected loss-of-function alleles for *atlas* by using

CRISPR to induce non-homologous end joining at a single PAM site just downstream of the

atlas start codon. *Vasa*-Cas9 embryos were co-injected and screened for w- progeny as

described above. We then used squish preps to isolate DNA from G1 flies and used a PCR-

RFLP assay to detect mutations. PCR products spanning the gRNA-targeted site were digested

with *Bfa*I (New England Biolabs (NEB), Ipswich, MA); undigested products in which the

expected *Bfa*I site was lost indicated a mutation, which was balanced and then confirmed by

PCR and sequencing of homozygous mutant lines.

We used scarless CRISPR editing and homology-directed repair (HDR) to insert the

GFP protein-coding sequence in-frame at the end of the *atlas* protein sequence (see Fig. S6;

(https://flycrispr.org/scarless-gene-editing/) [66-68]. We first generated an *atlas*-GFP DNA

construct by cloning the *atlas* protein-coding sequence into pENTR and using LR Clonase II

(Thermo Fisher Scientific, Waltham, MA) to recombine the sequence with pTWG (DGRC #1076;

T. Murphy), generating a C-terminally tagged *atlas*:GFP construct. We amplified the *atlas*

fragment from *vasa*-Cas9 strain #51323 genomic DNA. Once *atlas*-GFP was obtained in a

plasmid, we amplified it with primers that contained 5' *Esp*3I sites and overhangs designed for

Golden Gate Assembly (GGA) and that, in the case of the reverse primer, also added on 42

nucleotides downstream of the *atlas* stop codon to reach a PAM site identified by FlyCRISPR

TargetFinder [119] as being optimal for Cas9/gRNA recognition and cleavage. The primer also

introduced a mutation in the PAM site so that insertion of the designed piece of DNA into the

genome *in vivo* would not be subject to re-cutting. We also used the NEB Q5 Site-Directed

Mutagenesis kit to introduce a silent mutation into the *atlas* protein-coding sequence to

eliminate an internal *Esp*3I site. The resulting construct was used as the "left" homology arm for

homology-directed repair (HDR) editing. We constructed a "right" homology arm by using NEB

Q5 PCR to amplify a 982-bp fragment downstream of the PAM site, using primers modified to

contain *Esp*3I sites and overhangs compatible with GGA. We performed GGA by combining

these left and right arms, a plasmid containing a PiggyBac transposase-excisable 3xP3-dsRed

flanked by *Esp*3I sites, and backbone plasmid pXZ13, with *Esp*3I and T4 DNA ligase (NEB). A

combination of colony PCR, restriction digestion and sequencing identified properly assembled

plasmids suitable for HDR.

Vasa-Cas9 embryos were co-injected with the assembled plasmid and a pU6.3 plasmid

encoding a gRNA targeting the region just downstream of the *atlas* stop codon. G0 flies were

crossed to w^{1118} adults, and G1 flies were screened for red fluorescent eyes using the

NIGHTSEA system (NIGHTSEA LLC, Lexington, MA). Six balanced lines from two independent

G1 broods were established. To remove the dsRed from the *atlas* locus, we crossed these lines

to a PiggyBac transposase line (BDSC #8285) and then selected against pBac and dsRed in

the following generation. PCR and sequencing confirmed the expected "scarless" insertion of

GFP at the *atlas* locus.

atlas *Genomic Rescue Line*

We constructed an HA-tagged *atlas* rescue line that contained the *atlas* gene flanked by

1345 bp of sequence upstream of the start codon (but excluding the coding sequence of

upstream gene *CG3124*) and 3000 bp of sequence downstream of the stop codon (including the

full 3' UTR) as follows. Genomic sequences were PCR amplified using Q5 High fidelity

Polymerase (NEB), purified Canton S genomic DNA (Gentra Puregene Tissue Kit, Qiagen,

Germantown, MD), and the atlas rescue F1/R1 and atlas rescue F3/R3 primer sets (see Table

S4). The 3x-HA tag was likewise amplified from pTWH (DGRC 1100; T. Murphy) using atlas

rescue F2/R2 primers. These DNA fragments were subsequently assembled into a XbaI/AscI-

linearized w+attB plasmid (Addgene, Watertown, MA, plasmid 30326, deposited by J.

Sekelsky). The assembled construct was then phiC31 integrated into the PBac{y^+-attP-

3B}VK00037 (Bloomington *Drosophila* Stock Center (BDSC) stock #24872) docking site

(Rainbow Transgenics) and crossed into the *atlas* null background to assess rescue.

Fertility Assays and Sperm Visualization

To validate the finding of reduced fertility for *atlas* knockdown males in the group fertility

assay described above, we performed single-pair fertility assays in which knockdown or mutant

males or their controls were mated individually to Canton S virgin females. Based on previous

experience analyzing genes that resulted in sterility or near-sterility [20, 46], we designed

assays with *N* = 20-30 flies per male genotype. Matings were observed, and males were

discarded after copulation. Females were allowed to lay eggs into the vials for 4 days and then

discarded. Pupal cases were counted as a measure of fertility. Crosses to generate and

mating assays involving RNAi flies were maintained at 25º to optimize knockdown. Before all

assays, flies were reared to sexual maturity (3-7 days) in single-sex groups on cornmeal-

molasses food supplemented with dry yeast grains [46].

To assess the level of fertility conferred by the *atlas*-GFP allele, we crossed *atlas*-GFP

and w^{1118} flies to Δ*atlas*/SM6. Males with genotypes atlas-GFP/Δ*atlas* and +/Δ*atlas* were

compared using the single-pair fertility assay described above.

To observe the production of sperm in knockdown or mutant males, we introduced the

Mst35Bb-GFP marker into these males, which labels mature sperm and late-stage spermatid

nuclei with GFP [51]. Samples were prepared, imaged and analyzed as described previously

[46].

Atlas-GFP Ectopic Expression

We used the Gateway cloning system (Thermo) to construct an *atlas*-GFP transgene

expressed under UAS control (primers in Table S4). The *atlas* protein-coding sequence in

pENTR was recombined with pTWG (*Drosophila* Genomics Resource Center, T. Murphy) as

described above. The resulting plasmid was then inserted into w- flies using P-element-

mediated transposition (Rainbow Transgenics), w+ G1s were selected, and several

independent insertions were balanced. We crossed male UAS-*atlas*:GFP flies to females from

two different driver lines: *tubulin*-GAL4 (to drive ubiquitous expression) and *Bam*-GAL4 (to drive

expression in the early germline). We dissected larval salivary glands of the *tub>atlas*:GFP

males, since these cells are exceptionally large and ideal for visualizing subcellular localization.

We then dissected the testes of *Bam>atlas*:GFP males to evaluate whether the localization

pattern observed in the salivary gland was consistent in testis tissue, albeit not the same cells in

which endogenous atlas appears to be expressed. Protein localization was visualized by

fluorescence confocal microscopy on a Leica SP5 microscope (Leica Microsystems, Wetzlar,

Germany) and images were captured with LASAF as described previously [20].

Imaging Spermatogenesis and Spermatid Nuclear Condensation

897   We used phase-contrast microscopy to examine the stages of spermatogenesis in whole

mount testes [120]. To assess the processes of nuclear condensation and individualization of

64-cell cysts of spermatids in the post-meiotic stages of spermatogenesis, we used

fluorescence and confocal microscopy to visualize actin-based individualization complexes and

nuclei. Samples were processed, and actin and nuclear DNA were visualized with TRITC-

phalloidin (Molecular Probes, Eugene, OR) and TOPRO-3 iodine (Thermo), respectively, as

described previously [20]. The final stages of nuclear condensation were visualized with the

Mst35Bb-GFP marker described above, as well as an equivalent marker, Mst35Bb-dsRed [51].

Earlier nuclear stages were visualized with histone H2AvD-RFP (BDSC stock #23651), which is

present in round spermatid nuclei and the earliest stages of nuclear elongation [53, 121].

Images with H2AvD-RFP were obtained with epifluorescence microscopy, since we lack an

appropriate confocal laser for RFP.

909   To examine spermatid nuclei at various stages of condensation, we visualized nuclear

bundles using TOPRO-3. Testes of newly eclosed (<1 day old) *atlas* null and control males

were dissected in PBS. Testes were then transferred to a droplet of 2% paraformaldehyde on

poly-L-lysine treated glass slides and were gently shredded in the post-meiotic region to release

sperm bundles. Testes were gently squashed beneath coverslips coated in Sigmacote (Sigma

Aldrich, St. Louis, MO). We then froze slides in liquid nitrogen for a few seconds and popped off

of the siliconized coverslip with a razor. Slides were incubated in Coplin jars filled with 95%

ethanol at -20°C for 30 minutes and then mounted in VECTASHIELD (Vector Laboratories,

Burlingame, CA). Nuclear staging was performed by examining the shape of the nuclei. Early

and late canoe stages of condensation were distinguished by the absence or presence of

Mst35Bb-GFP, respectively. Elongated and late canoe stages were distinguished by the

presence or absence, respectively, of vesicles of GFP-tagged nuclear proteins (Atlas-GFP or

Mst35Bb-GFP) located basal to the nuclei. Examples of stages are given in Fig. 4. Confocal

stacks were taken on a Leica SP5 microscope, images were captured by LASAF, and ImageJ

was used to flatten stacks into a single, two-dimensional image. All intact nuclear bundles were

counted for each dissection.

For the experiments measuring nuclear condensation stage (Table S1), a sample size of

N = 10 for each genotype was selected based on the magnitude of the *atlas* null phenotype and

the consistent differences observed in previous dissections of these genotypes with Mst35Bb-

GFP. Likewise, for the IC-nuclear bundle association and IC progression analysis (Fig. 3C-D),

we selected sample sizes of N = ~15 per genotype based on pilot experiments showing that

aberrant actin phenotypes were highly consistent in null testes and previous experience with

such quantification [20].

Evolutionary and Gene Expression Analysis of atlas

We searched for orthologs of the *D. melanogaster* Atlas protein in the original 12

sequenced *Drosophila* species with BLASTP searches in FlyBase [122]. We also used

TBLASTN searches to identify orthologs in species lacking complete protein annotations. We

identified syntenic regions for each species by looking for conserved neighboring genes, such

as *ord* and *CG3124*. In addition to analyzing the *atlas* coding region, we conducted separate

BLASTN searches for the sequence of the *D. melanogaster* 3'UTR across *Drosophila* species

since it has a different conservation pattern than the coding sequence.

To test for sex-specific expression biases for both the ORF and the 3' UTR sequences,

we used adult male- and female-specific RNA-seq data from numerous *Drosophila* species

accessed through the Genomics Education Partnership version of the UCSC Genome Browser

(http://gander.wustl.edu/) and initially collected by Brown et al. [115] and Chen et al. [123] . We

also confirmed these findings experimentally in several species by performing RT-PCR on

cDNA isolated from whole males and whole females, as previously described [46].

To search for *atlas* orthologs in non-*Drosophila* Dipterans, we obtained from ENSEMBL

Metazoa the genomes of *Musca domestica, Glossina morsitans, Lucilia cuprina, Aedes aegypti,*

Anopheles darlingi, Anopheles gambiae, Culex quinquefasciatus and *Mayetiola destructor*. We

performed a synteny search (summarized in Fig. S11) in each species by identifying the nearest

neighbors of *atlas* in the *D. ananassae* and *D. virilis* genomes that had an identifiable homolog

in each species. In all cases, the homologs of the nearest neighbors on each side of *atlas* were

found on different contigs, suggesting synteny breakdown. We obtained up to 1 Mb of

sequence on each side of each identified homolog and queried it with BLASTN, TBLASTN, and

Exonerate [124] for regions with significantly similarity to any portion of the Atlas protein or

cDNA sequences. No significant hits, and no hits better than what could be found in other parts

of the genome, were found. Finally, we used HMMER to search for orthologs in all annotated

proteomes and all metazoan proteomes. We first queried the database with Atlas from either *D.*

melanogaster or *D. virilis* and accepted hits that fell below an e-value cutoff of 0.01 and a

minimum hit length of 3%. These hits were then included iteratively in subsequent searches

until no new significant hits were found.

We analyzed the molecular evolution of the *atlas* protein-coding sequence by obtaining

orthologous protein-coding sequences from *melanogaster* group species. (Analysis out of this

group was not performed due to high sequence divergence and poor alignment quality.) We

used BLASTP to identify these sequences from GenBank and then extracted the coding DNA

sequence for each. Sequences were aligned, checked for recombination, used to construct a

gene tree, and analyzed with the PAML sites test as described previously [78], except that

alignment positions that included gaps were masked from the PAML analysis. We initially

analyzed a set of 13 species (*melanogaster*, *simulans*, *sechellia*, *yakuba*, *erecta*, *suzukii*,

takahashii, *biarmipes*, *rhopaloa*, *ficusphila*, *elegans*, *eugracilis* and *ananassae*); we excluded an

ortholog detected in *D. bipectinata* due to poor alignment. This initial analysis detected a class

of sites with significant evidence of positive selection, but closer inspection of the alignment

revealed that the site with the strongest evidence of selection, corresponding to *D.*

melanogaster residue 31R, may have been driven by a questionable alignment due to an

insertion in that region that was unique to *D. takahashii*. The reported results come from an

analysis that excluded *D. takahashii*, which produced a more reliable alignment and showed no

evidence for any sites under positive selection.

<u>Acknowledgements</u>

We thank Dr. Alexis Hill for assistance with developing the *atlas*-GFP transgene; Gynesis

Vance, Elvis Perez, Ishanpepe Jagusah, Emily Gualdino, and students in the Fall 2016 and Fall

2017 sections of Biology 261 Lab at Holy Cross for assistance with RNAi fertility screens; Dr.

Rob Bellin for assistance with confocal microscopy; Dr. Justin McAlister for use of the

NIGHTSEA system; the Bloomington Stock Center and the Vienna *Drosophila* Resource Center

for fly strains; and, Dr. Mariana Wolfner and members of the Findlay and Bornberg-Bauer labs

for productive discussions about the project and the manuscript. This work was supported by

NSF CAREER Award #1652013 (to GDF) and a Humboldt Fellowship (to AG). GDF and EBB

are also grateful to the Evolution Think Tank at the University of Münster, which funded GDF as

a visiting fellow for in-person collaboration.

References

991
992
1.   Zhang J. Evolution by gene duplication: an update. Trends Ecol Evol. 2003;18(6):292-8.
doi: https://doi.org/10.1016/S0169-5347(03)00033-8.
2.   Lipinski KJ, Farslow JC, Fitzpatrick KA, Lynch M, Katju V, Bergthorsson U. High
spontaneous rate of gene duplication in *Caenorhabditis elegans*. Curr Biol. 2011;21(4):306-10.
doi: 10.1016/j.cub.2011.01.026.
3.   Van Oss SB, Carvunis A-R. *De novo* gene birth. PLoS Genet. 2019;15(5):e1008160. doi:
10.1371/journal.pgen.1008160.
4.   Bornberg-Bauer E, Schmitz JF. Fact or fiction: Updates on how protein-coding genes
might emerge *de novo* from previously non-coding DNA. F1000Res. 2017;6:57. doi:
10.12688/f1000research.10079.1.
5.   McLysaght A, Hurst LD. Open questions in the study of *de novo* genes: what, how and
why. Nat Rev Genet. 2016;17:567.
6.   Li Z-W, Chen X, Wu Q, Hagmann J, Han T-S, Zou Y-P, et al. On the origin of *de novo*
genes in *Arabidopsis thaliana* populations. Genome Biol Evol. 2016;8(7):2190-202. doi:
10.1093/gbe/evw164.
7.   Guerzoni D, McLysaght A. *De novo* genes arise at a slow but steady rate along the
primate lineage and have been subject to incomplete lineage sorting. Genome Biol Evol.
2016;8(4):1222-32. doi: 10.1093/gbe/evw074.
8.   Lu T-C, Leu J-Y, Lin W-C. A comprehensive analysis of transcript-supported de novo
genes in *Saccharomyces sensu stricto* yeasts. Mol Biol Evol. 2017;34(11):2823-38. doi:
10.1093/molbev/msx210.
9.   Zhang L, Ren Y, Yang T, Li G, Chen J, Gschwend AR, et al. Rapid evolution of protein
diversity by de novo origination in *Oryza*. Nat Ecol Evol. 2019;3(4):679-90. doi: 10.1038/s41559-
019-0822-5.
10.   Zhao L, Saelao P, Jones CD, Begun DJ. Origin and spread of *de novo* genes in
Drosophila melanogaster populations. Science. 2014;343(6172):769-72. doi:
10.1126/science.1248286.
11.   Puntambekar S, Newhouse R, Navas JSM, Chauhan R, Vernaz G, Willis T, et al.
Evolutionary divergence of novel open reading frames in cichlids speciation. Sci Rep.
2020;10(1). doi: 10.1038/s41598-020-78555-0.
12.   Ruiz-Orera J, Hernandez-Rodriguez J, Chiva C, Sabidó E, Kondova I, Bontrop R, et al.
Origins of *de novo* genes in human and chimpanzee. PLoS Genet. 2016;11(12):e1005721.
13.   Chamakura KR, Tran JS, O'Leary C, Lisciandro HG, Antillon SF, Garza KD, et al. Rapid
de novo evolution of lysis genes in single-stranded RNA phages. Nat Commun. 2020;11(1). doi:
10.1038/s41467-020-19860-0.
14.   Carvunis A-R, Rolland T, Wapinski I, Calderwood MA, Yildirim MA, Simonis N, et al.
Proto-genes and *de novo* gene birth. Nature. 2012;487:370-4.
15.   Wilson BA, Foy SG, Neme R, Masel J. Young genes are highly disordered as predicted
by the preadaptation hypothesis of *de novo* gene birth. Nat Ecol Evol. 2017;1(6):0146.
16.   Schlötterer C. Genes from scratch – the evolutionary fate of *de novo* genes. Trends
Genet. 2015;31(4):215-9. doi: 10.1016/j.tig.2015.02.007.
17.   Schmitz JF, Ullrich KK, Bornberg-Bauer E. Incipient *de novo* genes can evolve from
frozen accidents that escaped rapid transcript turnover. Nat Ecol Evol. 2018;2(10):1626-32. doi:
10.1038/s41559-018-0639-7.
18.   Vakirlis N, Hebert AS, Opulente DA, Achaz G, Hittinger CT, Fischer G, et al. A molecular
portrait of de novo genes in yeasts. Mol Biol Evol. 2018;35(3):631-45. doi:
10.1093/molbev/msx315.

19.   Heames B, Schmitz J, Bornberg-Bauer E. A continuum of evolving de novo genes drives
protein-coding novelty in *Drosophila*. J Mol Evol. 2020;88(4):382-98. doi: 10.1007/s00239-020-
09939-z.
20.   Lange A, Patel PH, Heames B, Damry AM, Saenger T, Jackson CJ, et al. Structural and
functional characterization of a putative de novo gene in *Drosophila*. Nat Commun.
2021;12:1667. doi: 10.1101/2021.01.18.427054.
21.   Keeling DM, Garza P, Nartey CM, Carvunis A-R. The meanings of 'function' in biology
and the problematic case of de novo gene emergence. eLife. 2019;8:e47014. doi:
10.7554/elife.47014.
22.   Cai J, Zhao R, Jiang H, Wang W. *De novo* origination of a new protein-coding gene in
Saccharomyces cerevisiae. Genetics. 2008;179(1):487-96. doi: 10.1534/genetics.107.084491.
23.   Li D, Yan Z, Lu L, Jiang H, Wang W. Pleiotropy of the *de novo*-originated gene *MDF1*.
Sci Rep. 2014;4(1):7280. doi: 10.1038/srep07280.
24.   Xiao W, Liu H, Li Y, Li X, Xu C, Long M, et al. A rice gene of *de novo* origin negatively
regulates pathogen-induced defense response. PLoS One. 2009;4(2):e4603. doi:
10.1371/journal.pone.0004603.
25.   Baalsrud HT, Tørresen OK, Solbakken MH, Salzburger W, Hanel R, Jakobsen KS, et al.
De novo gene evolution of antifreeze glycoproteins in codfishes revealed by whole genome
sequence data. Mol Biol Evol. 2018;35(3):593-606. doi: 10.1093/molbev/msx311.
26.   Zhuang X, Yang C, Murphy KR, Cheng CHC. Molecular mechanism and history of non-
sense to sense evolution of antifreeze glycoprotein gene in northern gadids. Proc Natl Acad Sci
U S A. 2019;116(10):4400-5. doi: 10.1073/pnas.1817138116.
27.   Xie C, Bekpen C, Künzel S, Keshavarz M, Krebs-Wheaton R, Skrabar N, et al. A de
novo evolved gene in the house mouse regulates female pregnancy cycles. eLife. 2019;8. doi:
10.7554/elife.44392.
28.   Suenaga Y, Islam SMR, Alagu J, Kaneko Y, Kato M, Tanaka Y, et al. NCYM, a *cis*-
antisense gene of MYCN, encodes a *de novo* evolved protein that inhibits GSK3β resulting in
the stabilization of MYCN in human neuroblastomas. PLoS Genet. 2014;10(1):e1003996. doi:
10.1371/journal.pgen.1003996.
29.   Marques AC, Dupanloup I, Vinckenbosch N, Reymond A, Kaessmann H. Emergence of
young human genes after a burst of retroposition in primates. PLoS Biol. 2005;3(11):e357.
30.   Baker RH, Narechania A, Johns PM, Wilkinson GS. Gene duplication, tissue-specific
gene expression and sexual conflict in stalk-eyed flies (Diopsidae). Philos Trans R Soc Lond B
Biol Sci. 2012;367(1600):2357-75. doi: 10.1098/rstb.2011.0287.
31.   Cui X, Lv Y, Chen M, Nikoloski Z, Twell D, Zhang D. Young genes out of the male: an
insight from evolutionary age analysis of the pollen transcriptome. Mol Plant. 2015;8(6):935-45.
doi: 10.1016/j.molp.2014.12.008; 20
10.1016/j.molp.2014.12.008.
32.   Kaessmann H. Origins, evolution, and phenotypic impact of new genes. Genome Res.
2010;20(10):1313-26. doi: 10.1101/gr.101386.109.
33.   Levine MT, Jones CD, Kern AD, Lindfors HA, Begun DJ. Novel genes derived from
noncoding DNA in *Drosophila melanogaster* are frequently X-linked and exhibit testis-biased
expression. Proc Natl Acad Sci U S A. 2006;103(26):9935-9. doi: 10.1073/pnas.0509809103.
34.   Begun DJ, Lindfors HA, Kern AD, Jones CD. Evidence for *de Novo* Evolution of Testis-
Expressed Genes in the *Drosophila yakuba/Drosophila erecta* Clade. Genetics.
2007;176(2):1131-7. doi: 10.1534/genetics.106.069245.
35.   Necsulea A, Kaessmann H. Evolutionary dynamics of coding and non-coding
transcriptomes. Nat Rev Genet. 2014;15(11):734-48. doi: 10.1038/nrg3802.
36.   Soumillon M, Necsulea A, Weier M, Brawand D, Zhang X, Gu H, et al. Cellular source
and mechanisms of high transcriptome complexity in the mammalian testis. Cell Rep.
2013;3(6):2179-90. doi: 10.1016/j.celrep.2013.05.031.

37.   Sorourian M, Kunte MM, Domingues S, Gallach M, Özdil F, Río J, et al. Relocation
facilitates the acquisition of short *cis*-regulatory regions that drive the expression of retrogenes
during spermatogenesis in *Drosophila*. Mol Biol Evol. 2014;31(8):2170-80. doi:
10.1093/molbev/msu168.
38.   Palmieri N, Kosiol C, Schlötterer C. The life cycle of *Drosophila* orphan genes. eLife.
2014;3:e01311. doi: 10.7554/eLife.01311.
39.   Wilburn DB, Swanson WJ. From molecules to mating: Rapid evolution and biochemical
studies of reproductive proteins. J Proteomics. 2016;135:12-25. doi:
10.1016/j.jprot.2015.06.007.
40.   VanKuren NW, Long M. Gene duplicates resolving sexual conflict rapidly evolved
essential gametogenesis functions. Nat Ecol Evol. 2018;2(4):705-12. doi: 10.1038/s41559-018-
0471-0.
41.   Hales KG, Korey CA, Larracuente AM, Roberts DM. Genetics on the fly: a primer on the
Drosophila model system. Genetics. 2015;201(3):815-42. doi: 10.1534/genetics.115.183392.
42.   Demarco RS, Eikenes TH, Haglund K, Jones DL. Investigating spermatogenesis in
Drosophila melanogaster. Methods. 2014;68(1):218-27. doi: 10.1016/j.ymeth.2014.04.020.
43.   Witt E, Benjamin S, Svetec N, Zhao L. Testis single-cell RNA-seq reveals the dynamics
of de novo gene transcription and germline mutational bias in *Drosophila*. eLife. 2019;8. doi:
10.7554/elife.47138.
44.   Reinhardt JA, Wanjiru BM, Brant AT, Saelao P, Begun DJ, Jones CD. *De novo* ORFs in
Drosophila are important to organismal fitness and evolved rapidly from previously non-coding
sequences. PLoS Genet. 2013;9(10):e1003860. doi: 10.1371/journal.pgen.1003860.
45.   Kondo S, Vedanayagam J, Mohammed J, Eizadshenass S, Kan L, Pang N, et al. New
genes often acquire male-specific functions but rarely become essential in *Drosophila*. Genes
Dev. 2017;31(18):1841-6. doi: 10.1101/gad.303131.117.
46.   Gubala AM, Schmitz JF, Kearns MJ, Vinh TT, Bornberg-Bauer E, Wolfner MF, et al. The
goddard and *saturn* genes are essential for *Drosophila* male fertility and may have arisen *de*
novo. Mol Biol Evol. 2017;34(5):1066-82. doi: 10.1093/molbev/msx057.
47.   Abrusán G. Integration of new genes into cellular networks, and their structural
maturation. Genetics. 2013;195(4):1407-17. doi: 10.1534/genetics.113.152256.
48.   Zhang W, Landback P, Gschwend AR, Shen B, Long M. New genes drive the evolution
of gene interaction networks in the human and mouse genomes. Genome Biol. 2015;16(1). doi:
10.1186/s13059-015-0772-4.
49.   Wu DD, Irwin DM, Zhang YP. De novo origin of human protein-coding genes. PLoS
Genet. 2011;7(11). doi: 10.1371/journal.pgen.1002379.
50.   Ni J-Q, Zhou R, Czech B, Liu L-P, Holderbaum L, Yang-Zhou D, et al. A genome-scale
shRNA resource for transgenic RNAi in *Drosophila*. Nat Methods. 2011;8(5):405-7. doi:
10.1038/nmeth.1592.
51.   Manier MK, Belote JM, Berben KS, Novikov D, Stuart WT, Pitnick S. Resolving
mechanisms of competitive fertilization success in *Drosophila melanogaster*. Science.
2010;328(5976):354-7. doi: 10.1126/science.1187096.
52.   Steinhauer J. Separating from the pack: molecular mechanisms of *Drosophila* spermatid
individualization. Spermatogenesis. 2015;5(2):e1041345. doi:
10.1080/21565562.2015.1041345.
53.   Rathke C, Baarends WM, Jayaramaiah-Raja S, Bartkuhn M, Renkawitz R, Renkawitz-
Pohl R. Transition from a nucleosome-based to a protamine-based chromatin configuration
during spermiogenesis in *Drosophila*. J Cell Sci. 2007;120(9):1689-700. doi:
10.1242/jcs.004663.
54.   Fabian L, Brill JA. Drosophila spermiogenesis. Spermatogenesis. 2012;2(3):197-212.
doi: 10.4161/spmg.21798.

55.   Rathke C, Baarends WM, Awe S, Renkawitz-Pohl R. Chromatin dynamics during spermiogenesis. Biochim Biophys Acta. 2014;1839(3):155-68. doi: 10.1016/j.bbagrm.2013.08.004.

56.   Kanippayoor RL, Alpern JHM, Moehring AJ. Protamines and spermatogenesis in *Drosophila* and *Homo sapiens*. Spermatogenesis. 2013;3(2):e24376. doi: 10.4161/spmg.24376.

57.   Jayaramaiah Raja S, Renkawitz-Pohl R. Replacement by *Drosophila melanogaster* protamines and Mst77F of histones during chromatin condensation in late spermatids and role of Sesame in the removal of these proteins from the male pronucleus. Mol Cell Biol. 2005;25(14):6165-77. doi: 10.1128/mcb.25.14.6165-6177.2005.

58.   Gärtner SMK, Rothenbusch S, Buxa MK, Theofel I, Renkawitz R, Rathke C, et al. The HMG-box-containing proteins tHMG-1 and tHMG-2 interact during the histone-to-protamine transition in *Drosophila* spermatogenesis. Eur J Cell Biol. 2015;94(1):46-59. doi: https://doi.org/10.1016/j.ejcb.2014.10.005.

59.   Gärtner SMK, Hundertmark T, Nolte H, Theofel I, Eren-Ghiani Z, Tetzner C, et al. Stage-specific testes proteomics of *Drosophila melanogaster* identifies essential proteins for male fertility. Eur J Cell Biol. 2019;98(2):103-15. doi: https://doi.org/10.1016/j.ejcb.2019.01.001.

60.   Alvi ZA, Chu T-C, Schawaroch V, Klaus AV. Genomic and expression analysis of transition proteins in *Drosophila*. Spermatogenesis. 2016;5(3):e1178518. doi: 10.1080/21565562.2016.1178518.

61.   Alvi ZA, Chu T-C, Schawaroch V, Klaus AV. Protamine-like proteins in 12 sequenced species of *Drosophila*. Protein Pept Lett. 2013;20(1):17-35. doi: 10.2174/092986613804096847.

62.   Dorus S, Freeman ZN, Parker ER, Heath BD, Karr TL. Recent Origins of Sperm Genes in Drosophila. Mol Biol Evol. 2008;25(10):2157-66. doi: 10.1093/molbev/msn162.

63.   Eren-Ghiani Z, Rathke C, Theofel I, Renkawitz-Pohl R. Prtl99C Acts Together with Protamines and Safeguards Male Fertility in *Drosophila*. Cell Rep. 2015;13(11):2327-35. doi: 10.1016/j.celrep.2015.11.023.

64.   Doyen CM, Chalkley GE, Voets O, Bezstarosti K, Demmers JA, Moshkin YM, et al. A Testis-Specific Chaperone and the Chromatin Remodeler ISWI Mediate Repackaging of the Paternal Genome. Cell Rep. 2015;13(7):1310-8. doi: 10.1016/j.celrep.2015.10.010.

65.   Lange A, Mills RE, Lange CJ, Stewart M, Devine SE, Corbett AH. Classical nuclear localization signals: definition, function, and interaction with Importin α. J Biol Chem. 2007;282(8):5101-5. doi: 10.1074/jbc.r600026200.

66.   Hill A, Jain P, Ben-Shahar Y. The *Drosophila* ERG channel seizure plays a role in the neuronal homeostatic stress response. PLoS Genet. 2019;15(8):e1008288. doi: https://doi.org/10.1371/journal.pgen.1008288.

67.   Bier E, Harrison MM, O'Connor-Giles KM, Wildonger J. Advances in Engineering the Fly Genome with the CRISPR-Cas System. Genetics. 2018;208(1):1-18. doi: 10.1534/genetics.117.1113.

68.   Bruckner JJ, Zhan H, Gratz SJ, Rao M, Ukken F, Zilberg G, et al. Fife organizes synaptic vesicles and calcium channels for high-probability neurotransmitter release. J Cell Biol. 2017;216(1):231-46. doi: 10.1083/jcb.201601098.

69.   Dorus S, Busby SA, Gerike U, Shabanowitz J, Hunt DF, Karr TL. Genomic and functional evolution of the *Drosophila melanogaster* sperm proteome. Nat Genet. 2006;38(12):1440-5. doi: 10.1038/ng1915.

70.   Wasbrough ER, Dorus S, Hester S, Howard-Murkin J, Lilley K, Wilkin E, et al. The *Drosophila melanogaster* sperm proteome-II (DmSP-II). J Proteomics. 2010;73(11):2171-85. doi: 10.1016/j.jprot.2010.09.002.

71.   Schuh M, Lehner CF, Heidmann S. Incorporation of *Drosophila* CID/CENP-A and CENP-C into Centromeres during Early Embryonic Anaphase. Curr Biol. 2007;17(3):237-43. doi: 10.1016/j.cub.2006.11.051.

72.   Rele CP, Sandlin KM, Leung W, Reed LK. Manual annotation of genes within *Drosophila*
Species: the Genomics Education Partnership protocol. bioRxiv [Preprint]. 2021 [posted 2020
Dec 11; revised 2020 Dec 12; revised 2020 Dec 14; revised 2020 Dec 15; revised 2021 Jan 27;
revised 2021 May 4; revised 2021 May 21; revised 2021 June 5;
cited 2021 Aug 10]. Available from:
https://www.biorxiv.org/content/10.1101/2020.12.10.420521v8.article-info doi:
https://doi.org/10.1101/2020.12.10.420521
73.   Schaeffer SW, Bhutkar A, McAllister BF, Matsuda M, Matzkin LM, O'Grady PM, et al.
Polytene chromosomal maps of 11 Drosophila species: the order of genomic scaffolds inferred
from genetic and physical maps. Genetics. 2008;179(3):1601-55. doi:
10.1534/genetics.107.086074.
74.   Weisman CM, Murray AW, Eddy SR. Many, but not all, lineage-specific genes can be
explained by homology detection failure. PLoS Biol. 2020;18(11):e3000862. doi:
10.1371/journal.pbio.3000862.
75.   Potter SC, Luciani A, Eddy SR, Park Y, Lopez R, Finn RD. HMMER web server: 2018
update. Nucleic Acids Res. 2018;46(W1):W200-W4. doi: 10.1093/nar/gky448.
76.   Zdobnov EM, Tegenfeldt F, Kuznetsov D, Waterhouse RM, Simão FA, Ioannidis P, et al.
OrthoDB v9.1: cataloging evolutionary and functional annotations for animal, fungal, plant,
archaeal, bacterial and viral orthologs. Nucleic Acids Res. 2017;45(D1):D744-D9. doi:
10.1093/nar/gkw1119.
77.   Yang Z. PAML 4: phylogenetic analysis by maximum likelihood. Mol Biol Evol.
2007;24(8):1586-91. doi: 10.1093/molbev/msm088.
78.   McGeary MK, Findlay GD. Molecular evolution of the sex peptide network in *Drosophila*.
J Evol Biol. 2020;33(5):629-41. doi: 10.1111/jeb.13597.
79.   Hundertmark T, Gärtner SMK, Rathke C, Renkawitz-Pohl R. Nejire/dCBP-mediated
histone H3 acetylation during spermatogenesis is essential for male fertility in *Drosophila*
melanogaster. PLoS One. 2018;13(9):e0203622. doi: 10.1371/journal.pone.0203622.
80.   Rathke C, Barckmann B, Burkhard S, Jayaramaiah-Raja S, Roote J, Renkawitz-Pohl R.
Distinct functions of Mst77F and protamines in nuclear shaping and chromatin condensation
during *Drosophila* spermiogenesis. Eur J Cell Biol. 2010;89(4):326-38. doi:
10.1016/j.ejcb.2009.09.001.
81.   Kimura S, Loppin B. The *Drosophila* chromosomal protein Mst77F is processed to
generate an essential component of mature sperm chromatin. Open Biol. 2016;6(11):160207.
doi: 10.1098/rsob.160207.
82.   Tirmarche S, Kimura S, Sapey-Triomphe L, Sullivan W, Landmann F, Loppin B.
Drosophila protamine-like Mst35Ba and Mst35Bb are required for proper sperm nuclear
morphology but are dispensable for male fertility. G3 (Bethesda). 2014;4(11):2241-5. doi:
10.1534/g3.114.012724.
83.   Vakirlis N, Acar O, Hsu B, Castilho Coelho N, Van Oss SB, Wacholder A, et al. De novo
emergence of adaptive membrane proteins from thymine-rich genomic sequences. Nat
Commun. 2020;11(1). doi: 10.1038/s41467-020-14500-z.
84.   Ravi Ram K, Wolfner MF. Sustained post-mating response in *Drosophila melanogaster*
requires multiple seminal fluid proteins. PLoS Genet. 2007;3(12):e238. doi:
10.1371/journal.pgen.0030238.
85.   Findlay GD, Sitnik JL, Wang W, Aquadro CF, Clark NL, Wolfner MF. Evolutionary rate
covariation identifies new members of a protein network required for *Drosophila melanogaster*
female post-mating responses. PLoS Genet. 2014;10(1):e1004108.
86.   Schnakenberg SL, Matias WR, Siegal ML. Sperm-storage defects and live birth in
Drosophila females lacking spermathecal secretory cells. PLoS Biol. 2011;9(11):e1001192. doi:
10.1371/journal.pbio.1001192.

87.   Yeh SD, Do T, Chan C, Cordova A, Carranza F, Yamamoto EA, et al. Functional
evidence that a recently evolved Drosophila sperm-specific gene boosts sperm competition.
Proc Natl Acad Sci U S A. 2012;109(6):2043-8. doi: 10.1073/pnas.1121327109.
88.   Civetta A, Finn S. Do candidate genes mediating conspecific sperm precedence affect
sperm competitive ability within species? A test case in *Drosophila*. G3 (Bethesda).
2014;4(9):1701-7. doi: 10.1534/g3.114.012476.
89.   Wong A, Albright SN, Giebel JD, Ram KR, Ji S, Fiumera AC, et al. A role for Acp29AB, a
predicted seminal fluid lectin, in female sperm storage in *Drosophila melanogaster*. Genetics.
2008;180(2):921-31. doi: 10.1534/genetics.108.092106.
90.   Larkin A, Marygold SJ, Antonazzo G, Attrill H, Gilberto, Garapati PV, et al. FlyBase:
updates to the *Drosophila melanogaster* knowledge base. Nucleic Acids Res.
2021;49(D1):D899-D907. doi: 10.1093/nar/gkaa1026.
91.   Parisi M, Nuttall R, Naiman D, Bouffard G, Malley J, Andrews J, et al. Paucity of genes
on the *Drosophila* X chromosome showing male-biased expression. Science.
2003;299(5607):697-700. doi: 10.1126/science.1079190.
92.   Parisi M, Nuttall R, Edwards P, Minor J, Naiman D, Lü J, et al. A survey of ovary-, testis-,
and soma-biased gene expression in *Drosophila melanogaster* adults. Genome Biol.
2004;5(6):R40. doi: 10.1186/gb-2004-5-6-r40.
93.   Vibranovski MD, Zhang Y, Long M. General gene movement off the X chromosome in
the *Drosophila* genus. Genome Res. 2009;19(5):897-903. doi: 10.1101/gr.088609.108.
94.   Emerson JJ, Kaessmann H, Betrán E, Long M. Extensive gene traffic on the mammalian
X chromosome. Science. 2004;303(5657):537-40. doi: 10.1126/science.1090042.
95.   Reinke V, San Gil I, Ward S, Kazmer K. Genome-wide germline-enriched and sex-
biased expression profiles in *Caenorhabditis elegans*. Development. 2004;131(2):311-23. doi:
10.1242/dev.00914.
96.   Argyridou E, Huylmans AK, Königer A, Parsch J. X-linkage is not a general inhibitor of
tissue-specific gene expression in *Drosophila melanogaster*. Heredity (Edinb). 2017;119(1):27-
34. doi: 10.1038/hdy.2017.12.
97.   Kemkemer C, Catalán A, Parsch J. 'Escaping' the X chromosome leads to increased
gene expression in the male germline of *Drosophila melanogaster*. Heredity (Edinb).
2014;112(2):149-55. doi: 10.1038/hdy.2013.86.
98.   Argyridou E, Parsch J. Regulation of the X chromosome in the germline and soma of
Drosophila melanogaster males. Genes (Basel). 2018;9(5):242. doi: 10.3390/genes9050242.
99.   Vibranovski MD, Lopes HF, Karr TL, Long M. Stage-specific expression profiling of
Drosophila spermatogenesis suggests that meiotic sex chromosome inactivation drives genomic
relocation of testis-expressed genes. PLoS Genet. 2009;5(11):e1000731. doi:
10.1371/journal.pgen.1000731.
100.   Zhang YE, Vibranovski MD, Krinsky BH, Long M. Age-dependent chromosomal
distribution of male-biased genes in *Drosophila*. Genome Res. 2010;20(11):1526-33. doi:
10.1101/gr.107334.110.
101.   Zhang YE, Vibranovski MD, Landback P, Marais GAB, Long M. Chromosomal
redistribution of male-biased genes in mammalian evolution with two bursts of gene gain on the
X chromosome. PLOS Biol. 2010;8(10):e1000494. doi: 10.1371/journal.pbio.1000494.
102.   Gao G, Vibranovski MD, Zhang L, Li Z, Liu M, Zhang YE, et al. A long-term
demasculinization of X-linked intergenic noncoding RNAs in *Drosophila melanogaster*. Genome
Res. 2014;24(4):629-38. doi: 10.1101/gr.165837.113.
103.   Vibranovski MD, Zhang YE, Kemkemer C, Lopes HF, Karr TL, Long M. Re-analysis of
the larval testis data on meiotic sex chromosome inactivation revealed evidence for tissue-
specific gene expression related to the drosophila X chromosome. BMC Biol. 2012;10(1):49.
doi: 10.1186/1741-7007-10-49.

104.   Mahadevaraju S, Fear JM, Akeju M, Galletta BJ, Pinheiro MMLS, Avelino CC, et al.
Dynamic sex chromosome expression in *Drosophila* male germ cells. Nat Commun. 2021;12(1).
doi: 10.1038/s41467-021-20897-y.
105.   Mikhaylova LM, Nurminsky DI. Lack of global meiotic sex chromosome inactivation, and
paucity of tissue-specific gene expression on the *Drosophila* X chromosome. BMC Biol.
2011;9(1):29. doi: 10.1186/1741-7007-9-29.
106.   Landeen EL, Muirhead CA, Wright L, Meiklejohn CD, Presgraves DC. Sex chromosome-
wide transcriptional suppression and compensatory *cis*-regulatory evolution mediate gene
expression in the *Drosophila* male germline. PLoS Biol. 2016;14(7):e1002499. doi:
10.1371/journal.pbio.1002499.
107.   Meiklejohn CD, Landeen EL, Cook JM, Kingan SB, Presgraves DC. Sex chromosome-
specific regulation in the *Drosophila* male germline but little evidence for chromosomal dosage
compensation or meiotic inactivation. PLoS Biol. 2011;9(8):e1001126. doi:
10.1371/journal.pbio.1001126.
108.   Vibranovski MD. Meiotic Sex Chromosome Inactivation in *Drosophila*. J Genomics.
2014;2:104-17. doi: 10.7150/jgen.8178.
109.   Wen K, Yang L, Xiong T, Di C, Ma D, Wu M, et al. Critical roles of long noncoding RNAs
in *Drosophila* Spermatogenesis. Genome Res. 2016;26(9):1233-44. doi:
10.1101/gr.199547.115.
110.   Bouska MJ, Bai H. Long noncoding RNA regulation of spermatogenesis via the spectrin
cytoskeleton in *Drosophila*. G3 (Bethesda). 2021. doi: 10.1093/g3journal/jkab080.
111.   Lim C, Tarayrah L, Chen X. Transcriptional regulation during Drosophila
spermatogenesis. Spermatogenesis. 2012;2(3):158-66. doi: 10.4161/spmg.21775.
112.   White-Cooper H. Molecular mechanisms of gene regulation during Drosophila
spermatogenesis. Reproduction. 2010;139(1):11-21. doi: 10.1530/rep-09-0083.
113.   Moyers BA, Zhang J. Phylostratigraphic bias creates spurious patterns of genome
evolution. Mol Biol Evol. 2015;32(1):258-67. doi: 10.1093/molbev/msu286.
114.   Moyers BA, Zhang J. Toward reducing phylostratigraphic errors and biases. Genome
Biol Evol. 2018;10(8):2037-48. doi: 10.1093/gbe/evy161.
115.   Brown JB, Boley N, Eisman R, May GE, Stoiber MH, Duff MO, et al. Diversity and
dynamics of the *Drosophila* transcriptome. Nature. 2014;512(7515):393-9. doi:
10.1038/nature12962.
116.   White-Cooper H. Tissue, cell type and stage-specific ectopic gene expression and RNAi
induction in the Drosophila testis. Spermatogenesis. 2012;2(1):11-22. doi: 10.4161/spmg.19088.
117.   Dietzl G, Chen D, Schnorrer F, Su K-C, Barinova Y, Fellner M, et al. A genome-wide
transgenic RNAi library for conditional gene inactivation in *Drosophila*. Nature.
2007;448(7150):151-6. doi: 10.1038/nature05954.
118.   Ge DT, Tipping C, Brodsky MH, Zamore PD. Rapid screening for CRISPR-directed
editing of the *Drosophila* genome using *white* coconversion. G3 (Bethesda). 2016;6(10):3197-
206. doi: 10.1534/g3.116.032557.
119.   Gratz SJ, Ukken FP, Rubinstein CD, Thiede G, Donohue LK, Cummings AM, et al.
Highly specific and efficient CRISPR/Cas9-catalyzed homology-directed repair in *Drosophila*.
Genetics. 2014;196(4):961-71. doi: 10.1534/genetics.113.160713.
120.   White-Cooper H. Spermatogenesis: analysis of meiosis and morphogenesis. In:
Henderson D, editor. Drosophila Cytogenetics Protocols. Methods Mol Biol. 247. Totowa, NJ:
Humana Press; 2004. p. 45-75.
121.   Clarkson M, Saint R. A His2AvDGFP fusion gene complements a lethal His2AvD mutant
allele and provides an in vivo marker for Drosophila chromosome behavior. DNA Cell Biol.
1999;18(6):457-62. doi: 10.1089/104454999315178.

122.   Clark AG, Eisen MB, Smith DR, Bergman CM, Oliver B, Markow TA, et al. Evolution of
genes and genomes on the Drosophila phylogeny. Nature. 2007;450(7167):203-18. doi:
10.1038/nature06341.
123.   Chen Z-X, Sturgill D, Qu J, Jiang H, Park S, Boley N, et al. Comparative validation of the
D. melanogaster modENCODE transcriptome annotation. Genome Res. 2014;24(7):1209-23.
doi: 10.1101/gr.159384.113.
124.   Slater G, Birney E. Automated generation of heuristics for biological sequence
comparison. BMC Bioinformatics. 2005;6(1):31. doi: 10.1186/1471-2105-6-31.
1348