1 **The model diatom Phaeodactylum tricornutum provides insights into the**
2 **diversity and function of microeukaryotic DNA methyltransferases**
3

4 Antoine Hoguin[1][§], Feng Yang[2], Agnès Groisillier[2], Chris Bowler[1], Auguste Genovesio[1],
5 Ouardia Ait-Mohamed[1][*], Fabio Rocha Jimenez Vieira [1][¥][*] and Leila Tirichine[2][*]
6
7 [1]Institut de biologie de l'Ecole normale supérieure (IBENS), Ecole normale supérieure,
8 CNRS, INSERM, PSL Université Paris 75005 Paris, France

9 [2] Nantes Université, CNRS, US2B, UMR 6286, F-44000 Nantes, France
10
11 [§]Current affiliation: Institute of Molecular Plant Sciences, University of Edinburgh,
12 Edinburgh, EH9 3BF, United Kingdom.
13
14 [¥]Current affiliation: Laboratory of Computational and Quantitative Biology - LCQB -
15 UMR 7238 CNRS - Sorbonne Université. Institut de Biologie Paris Seine. 75005 Paris
16
17
18
19 * Authors for correspondence: tirichine-l@univ-nantes.fr,
20 fabio.rocha_jimenez_vieira@sorbonne-universite.fr, ouardia.mohamed@gmail.com

Abstract

Cytosine methylation is an important epigenetic mark involved in the transcriptional
control of transposable elements in mammals, plants and fungi. The Stramenopiles-
Alveolate-Rhizaria (SAR) lineages are a major group of ecologically important marine
microeukaryotes that include the main phytoplankton groups diatoms and
dinoflagellates. However, little is known about their DNA methyltransferase diversity.
Here, we performed an *in-silico* analysis of DNA methyltransferases found in marine
microeukaryotes and showed that they encode divergent DNMT3, DNMT4, DNMT5
and DNMT6 enzymes. Furthermore, we found three classes of enzymes within the
DNMT5 family. Using a CRISPR/Cas9 strategy we demonstrated that the loss of the
DNMT5a gene correlates with a global depletion of DNA methylation and
overexpression of young transposable elements in the model diatom *Phaeodactylum*
tricornutum. The study provides a pioneering view of the structure and function of a
DNMT family in the SAR supergroup using an attractive model species.

Introduction

In eukaryotes the methylation of the fifth carbon of cytosine (5mC) is a well-known epigenetic mark associated with transcriptional repression. It has been implicated in a wide range of cellular processes including the stability of repeat rich centromeric and telomeric regions as well as in repression of transposable element (TEs) expression[1–4]. 5mC is deposited by DNA methyltransferases (DNMTs) capable of *de novo* methylation and is propagated through subsequent cell division by maintenance DNMT enzymes. Eukaryotes have acquired a diverse set of DNMTs by horizontal gene transfer of bacterial DNA cytosine methyltransferase involved in the restriction-methylation system[5]. All DNMTs contain a catalytic protein domain composed of ten conserved motifs (annotated I to X) that provide binding affinity to the DNA substrate and the methyl donor cofactor S-Adenosyl methionine to process the transfer of a methyl group to unmethylated cytosines[6,7]. DNMTs have further diversified over evolutionary time scales in eukaryote lineages and acquired chromatin associated recognition and binding domains giving rise to a wide diversity of DNA methylation patterns[8,9].

The loss and gain of DNMTs have been associated with profound divergence in cell biology and control of gene expression. To date, six main eukaryotic DNMT families have been described and named DNMT1, DNMT2, DNMT3, DNMT4, DNMT5 and DNMT6[10,11]. In Metazoans, the combined activity of the DNMT3 and DNMT1 families allow the deposition and maintenance of DNA methylation patterns during the successive developmental waves of DNA demethylation and remethylation[12]. Zebrafish possesses six DNMT3 family *de novo* methyltransferase genes, *dnmt3–dnmt8*. This group includes both orthologs of mammalian *dnmt3a* and *dnmt3b* as well as fish-specific genes with no mammalian orthologs[13]. In fungi, the DNA methylation machinery consists in a maintenance activity by DNMT1/DIM2, as in *Neurospora crassa*[14], or the activity of ATPase-DNMT5 enzymes as reported in *Cryptococcus neoformans*[11,15]. The DNMT5 enzyme also correlates with a heavy histone linker DNA methylation landscape in *Micromonas pusilla*, the pelagophyte *Aureococcus annophagefferens* and the haptophyte *Emiliania huxleyi*[11]. Fungal DNMT4 relatives are involved in the DNA methylation related process known as Repeat-Induced Point

Mutation and Methylation Induced Premeiotically that leads to TE extinction and/or
stage specific repression as observed in *Aspergillus* and *Neurospora* species[16–19].

Losses and lineage specific duplication of DNMT1 and DNMT3 have occurred
during insect evolution, such as in Diptera lineages[20], leading to secondary loss of
global 5mC methylation. In plants, the acquisition of novel DNMT1 proteins named
Chromomethylases (CMTs) and the divergence of the DNMT3 family led to the
spreading of the asymmetrical non-CG patterns of DNA methylation that is extensively
found in angiosperms[21–23]. DNMT2 is known to methylate tRNAs to yield ribo-5-
methylcytidine in a range of eukaryotic organisms, including humans, mice,
Arabidopsis thaliana, and *Drosophila melanogaster*[24]. It is characterized by its
cytoplasmic localization that contrasts with the exclusively nuclear localization of
Dnmt1 and Dnmt3[25]. Lastly, DNMT6 has been found in *Chlorophyta*, *Haptophyta*,
Ochrophyta, diatoms and dinoflagellates (e.g., *Symbiodinium kawagutii* and
Symbiodinium minutum)[10,11,26,27] but its function remains elusive. Importantly, 5mC is
increasingly reported in eukaryotes of the Stramenopiles-Alveolate-Rhizaria (*SAR)*
lineages as in dinoflagellates[26], diatoms[27] and kelps[28]. However, because of the severe
underrepresentation of marine unicellular eukaryotes in modern sequencing
databases, our understanding of the DNA methylation machinery in these organisms
remains scarce.

Diatoms are a dominant, abundant, and highly diverse group of unicellular brown
microalgae (from 2 to 200 µm) of the stramenopile lineage. It is estimated that diatoms
are responsible for nearly 20% of primary production on earth[29,30]. They are known to
dominate marine polar areas and are major contributors of phytoplankton oceanic
blooms. To date, 5mC has been reported only in four diatoms, *Thalassiosira*
pseudonana[11],*Cyclotella cryptica*[31], *Fragilariopsis cylindrus*[11] and *Phaeodactylum*
tricornutum[11,27]. Diatom methylation patterns strongly contrast with the patterns
observed in animals but also dinoflagellates and plants[32]. Firstly, in *P. tricornutum*, *T.*
pseudonana and *F. cylindrus*, total levels of DNA methylation range from 8% to as low
as 1% of cytosines in the CG context[11] over repeats and TEs usually (but not
exclusively) concentrated in telomeric regions[11,27]. Non-CG methylation is also
detected but is scarce. Diatom genomes are therefore predominantly composed of
isolated highly CG methylated TE islands in an otherwise unmethylated genome and
to that regard are remarkably like fungal methylation profiles. In all diatoms examined
so far, methylated TEs often have low expression[11,27,31]. This is remarkably consistent

with the repressive role of DNA methylation in other eukaryotes and further traces back 5mC-mediated control of TE expression to the last eukaryotic common ancestor. Nonetheless, direct evidence of 5mC repressive role of TEs in diatoms is lacking. Diatom genomes contain predicted proteins similar to members of the DNMT2, DNMT3, DNMT4, DNMT5 and DNMT6 family[11,33]. The conservation of their domain composition across eukaryotic groups as in the yeast *Cryptococcus neoformans* suggests that diatom DNMT5-like C5-MTases play a conserved and specific role in DNA methylation[11,15]. However, the functions of the DNMTs reported in diatoms have not been characterized *in vivo*.

Recent advances in high throughput RNA sequencing technologies led to the development of the Microbial Eukaryote Transcriptome Sequencing Project (MMETSP)[34]. The MMETSP concatenates more than 650 transcriptomes from diverse microeukaryote lineages such as diatoms and dinoflagellates, making it the biggest sequence database for transcriptomes from individual marine microeukaryotes. Here, utilizing the newly defined enhanced Domain Architecture Framework (eDAF) methodology[35], we first explored the structural and phylogenetic diversity of DNMT sequences in marine microeukaryotes from the publicly available MMETSP sequencing databases. Using an integrative approach with available genomes and phylogenetic studies, we provide a DNMT phylogeny focused on the structural and domain diversity found in microeukaryote enzymes and discuss their evolutionary origins. We define, in the DNMT5 family, the sub-families DNMT5a, b and c enzymes, based on structure and phylogenetic assessment. The presence of the predicted DNMT5 family diversity remarkably contrasts with the apparent lack of DNMT1 in most of the MMETSP and microeukaryote databases. Using CRISPR/Cas9 genome editing, we present the functional characterization of the DNMT5a sub-family in the model diatom *P. tricornutum* demonstrating, to our knowledge for the first time in any SAR, the role of this family in the repression of TEs in an early diverging eukaryote lineage[36].

Results

Diversity of DNMT5 methyltransferases in microeukaryotes

To capture the diversity of 5-cytosine DNA methyltransferases encoded in microalgae, we applied a relaxed HMMER search (e-value=1 as the cut-off threshold) for the PFAM DNMT (PF00145) domain on transcriptomes from the MMETSP database. The aim of this approach was to retrieve the maximum of DNMT hits

between distantly related eukaryotes. In this study, we focused on DNMT1, DNMT3,
DNMT4, DNMT5 and DNMT6 gene families that are known or represent putative DNA
modifying enzymes. We retained sequences showing conserved DNMT domains and
depicted their domain structures using eDAF analysis[35]. We built a representative
phylogeny of DNA methyltransferases based on the alignment of conserved DNMT
motifs (Figure 1a, Supplementary Figure 1, Supplementary Data 1). Since DNMT2 is
an aspartic acid transfer RNA methyltransferase[25], published microalgal DNMT2
sequences were used as additional sequences for phylogenetic analysis. The tree
construction exploited the stability of Bayesian approaches to deal with the fast
evolution rates observed in our DNMT sequences. Methods based on posterior
probabilities present more support values than random sampling algorithms when
facing high mutation rates[37–39]. The MMETSP database is composed of transcripts and
is thus deprived from lowly expressed genes or pseudogenes. The relative absence of
a given gene family within a species will therefore be interpreted accordingly.

We found three gene families related to the DNMT5 clade of enzymes that we
named DNMT5a, DNMT5b and DNMT5c (Figure 1a). The sequence alignments show
homology in the functional DNMT motifs (I-IV, VII and X) that contain the S-Adenosyl
methionine binding and catalytic domains within DNMT5s (i.e DNMT5a, b and c)
(Supplementary Figure 2). We noticed that the DNMT5 S-Adenosyl methionine-binding
phenylalanine found in the catalytic motif IV of other DNMTs is replaced by a serine.
Along with other amino acid substitutions within the motifs I to V (Supplementary Figure
2), this serine locks the protein in an autoinhibitory state released by hemi-methylated
DNA binding and the ATPase activity of the SNF domain[40]. The three DNMT5 families
form a supported group of enzymes (posterior probabilities 0.94). The DNMT5a and
DNMT5b clades are well supported (posterior probabilities of 0.98 and 0.97,
respectively). The DNMT5c family is however less supported (posterior probability of
0.88). The relationships between the DNMT5a, b, c sequences are however
unresolved as the DNMT5a, b branch is poorly supported (posterior probability of
0.51). Of note, DNMT5a is found in distantly related eukaryote lineages. We found 76
species with at least one DNMT5 orthologue. We found a DNMT5a in the green alga
Tetraselmis astigmata but also in haptophytes and the marine photosynthetic excavate
euglenozoa *Eutreptiella gymnastica*. The DNMT5a family is also found in
stramenopiles, including diatoms, bolidomonas, pelagophytes and dictiochophytes, as
well in fungi (former *Cryptococcus* DNMT5-related enzymes) (Figure 1a,

Supplementary Data 2). The DNMT5b enzyme is found in diatoms, *Bolidomonas*
pacifica and haptophytes. *Emiliania huxleyi* DNMT5 enzymes are not found in other
haptophytes in the MMETSP database. In addition, the phylogenetic position of E.
huxleyi DNMT5 proteins are not very supportive. First, E. huxleyi DNMT5a protein is
not monophyletic with other ochrophyte DNMT5s and its DNMT5b appears as an
outgroup found in diatoms. (Supplementary Figure 1). Within diatoms, genomes from
both *F. cylindrus* and *Synedra* contain DNMT5a and DNMT5b gene copies
(Supplementary Data 3) but lineage specific loss of DNMT5a is also observed in
Thalassiosirales. Haptophyte DNMT5s could be of lateral gene transfer origin, as in
other microalgae. DNMT5c enzymes are specific to dinoflagellates that are known to
have very fast evolutionary rates and likely divergent base/amino acid compositions.
Dinoflagellate DNMT5c sequences may thus represent a highly divergent DNMT
subgroup that our phylogeny failed to associate with other DNMT5s.
We found that the DNMT5a and b families share a C-terminal SNF2-type
DEXDc/HELICc helicase domain composed of two helicases complemented or not by
a RING finger domain (Figure 1b, Supplementary Data 4). We found that DNMT5b
enzymes display unique features. First, among them, 14 contain an N-terminal laminin
B receptor domain as in *T. pseudonana* (Figure 1b, Supplementary Data 4). Also, other
DNMT5b enzymes contain N-terminal CpG methyl binding domains, as well as HAND
structure domains, methyl-lysine and methyl-arginine TUDOR binding domains
(Supplementary Data 4). Finally, their DNMT domain is longer compared to the
DNMT5a, c due to the presence of spacer sequences between motifs. These
differences in structure may highlight functional diversity between the DNMT5
subfamilies and are consistent with the duplication followed by divergence hypothesis
described above. Accordingly, the DNMT5c family also diverged compared to the
DNMT5a and b enzymes at the protein domain composition. It is indeed characterized
by a long (~1000 amino-acids) N-terminal sequence with no annotated functional
domains (Figure 1b, Supplementary Data 4).
213
214 **The DNMT4 and DNMT1 family methyltransferases in microalgae**
215   In our phylogenetic analysis, the DNMT4 and DNMT1 clades form a poorly
216 supported gene family, as previously described[11,41] (Figure 1a, Supplementary Figure
217 1). DNMT1s are maintenance enzymes in eukaryotes that often associate a DNMT
218 catalytic domain with chromatin binding domains such as Bromo-Adjacent Homology

domains, Plant HomeoDomains, chromodomains and domains required for interaction with accessory proteins. DNMT4 enzymes are related to DIM2 enzymes in fungi [41] and are involved in the Methylation Induced Premeiotically and Repeat-Induced Point Mutation processes. Interestingly, two DNMT4 enzymes were also described in the pennate diatom *F. cylindrus* and the centric diatom *T. pseudonana* based on a previous phylogenetic analysis of DNMT enzymes in microalgae[10]. We first confirmed that orthologues of *T. pseudonana* DNMT4 enzymes are widespread in diatom transcriptomes and genomes. A total of 8 pennate and 23 centric diatoms out of 60 diatom species, express or encode at least one DNMT4 related transcript (Supplementary Data 3). This finding suggests that the family is ancestral in diatoms. In our analysis, no DNMT4 enzymes were found in other species of the MMETSP database. Phylogenetic analysis indicates that RID and diatom DNMT4s may form a moderately supported monophyletic family of enzymes (Figure 1a). At the structural level, both RID and diatom DNMT4 enzymes diverged compared to DNMT1 enzymes, and also between each other. Most diatom DNMT4 enzymes are composed of a single DNMT domain as in *T. pseudonana*, which also contrasts with fungal enzymes (Figure 1b, Supplementary Data 4). Nonetheless, nine diatom DNMT4 proteins possess an additional N-terminal chromodomain as observed in *Thalassiosira miniscula* (Figure 1b, Supplementary Data 3 and Supplementary Data 4). Furthermore, we found two putative DNMT1-like enzymes in the transcriptomic database of two *Raphidophyceae* brown microalgae: *Heterosigma akashiwo* and *Chatonella subsalsa*. They are composed of a conserved DNMT domain and a plant homeodomain (Figure 1b, Supplementary Figure 1, Supplementary Data 4) but poorly defined a monophyletic gene family with either DNMT1s or DNMT4s.

Interestingly, we found a DNMT1-related enzyme in three haptophyte species out of four (*Gephyrocapsa oceanica*, *Isochrysis.sp-CCMP1324* and *Coccolithus pelagicus*) from the MMETSP database that cluster with annotated CMTs found in the coccolithophore *E. huxleyi* (Figure 1a, Supplementary Figure 1). We found that the enzymes of *Gephyrocapsa oceanica* (CAMPEP_0188208858), *Isochrysis-CCMP1324* (CAMPEP_0188844028) and *Emiliania huxleyi* (jgi_215571) have DNMT1-like structures with a Replication Foci Domain followed by a Bromo-Adjacent Homology (in *Emiliana huxleyi* only) and a conserved DNMT domain (Figure 1b, Supplementary Data 4). Haptophyte enzymes seem to distantly relate to the conserved green algal CMT (hCMT2) enzymes (Figure 1a, Supplementary Figure 1).

We detected DNMT1/MET1 transcripts encoding proteins similar to the plant
MET1 enzyme in seven green algae species from MMETSP, mainly from the
Chlamydomonadales order such as in some *Chlamydomonas* species (Figure 1b,
Supplementary Figure 1, Supplementary Data 2).

The DNMT3 and DNMT6 methyltransferases are abundant in diatoms and lack
chromatin associated domains

Our data indicate that the DNMT3 family is not particularly frequent in
microalgae (Figure 2, Supplementary Data 2). DNMT3 is absent in most stramenopiles
except in diatoms; for which genomic and transcriptomic data strongly support its
presence (Supplementary Data 3). DNMT3 seems absent in the studied haptophytes
(Figure 2, Supplementary Data 2). Only one transcript from the cryptomonad
Goniomonas pacifica could be annotated as DNMT3. In addition, we could not identify
DNMT3 enzymes in any green algae in MMETSP, although it is present in red algae
as it is found in the genomes of *Cyanidioschyzon merolae* and *Galdieria sulphuraria*
(Figure 2, Supplementary Data 2). We also report several additional DNMT3
transcripts in dinoflagellates, as previously described[26] (Figure 2, Supplementary Data
2). Upon alignment, dinoflagellate DNMT3 enzymes (including former annotated
enzymes[26]) and *Goniomonas pacifica* DNMT3s are closely related to those from red
algae but diverge from other DNMT3s, while diatoms display their own DNMT3 family
(Supplementary Figure 1). This suggests that the DNMT3 family was iteratively lost
and acquired several times during microalgal evolution. As observed in *P. tricornutum*,
DNMT3 enzymes found in microalgae, all lack chromatin associated domains (Figure
1b, Supplementary Data 4). This contrasts with mammalian DNMT3s[42] that interact
with histone post-translational modifications.

DNMT6 enzymes were found among the most widespread DNMTs in
microeukaryotes. We found a DNMT6 transcript in the MMETSP transcriptomes of
three *Tetraselmis* green algae and seven dinoflagellates (Figure 2, Supplementary
Data 2). In addition, DNMT6 is distributed extensively in stramenopiles, including
Dictyochophyceae, *Crysophyceae* and *Pelagophyceae* (Figure 2, Supplementary Data
2). In diatoms, DNMT6 is very abundant (Supplementary Data 3). DNMT6 is also
present in the non-photosynthetic labyrinthulomycetes *Aplanochytrium stocchinoi* and
probably in *Aplanochytrium keurgelense* (Figure 2, Supplementary Data 2). In addition,
our data strongly support the presence of DNMT6 orthologues in the major

Chromalveolata lineage of *Rhizaria* (Figure 2, Supplementary Data 2), as suggested

in previous reports[26]. DNMT6 enzymes are mostly homogeneous and do not contain

chromatin associated signatures, as in *P. tricornutum* DNMT6 and DNMT3 (Figure 1b,

Supplementary Data 4). Finally, monophyletic relationships within the DNMT6 family

and between microeukaryotes could not be resolved (Supplementary Figure 1).

Single base resolution of DNA methylation in *P. tricornutum* DNMT5:KO lines

The pennate diatom *P. tricornutum* is the model diatom species that we and

others use to study the epigenomic landscape in diatoms, shedding light into the

conservation and divergence of DNA methylation patterns in early diverging

eukaryotes[27,43]. The *P. tricornutum* genome encodes DNMT3 (Phatr3_J47136),

DNMT6 (Phatr3_J47357) and DNMT5a (Phatr3_EG02369) orthologues in single

copies but lacks the DNMT4 and DNMT5b orthologues found in other diatoms

(Supplementary Data 3). We asked whether any of these DNMTs have DNA

methylation function(s) *in vivo*. Using a CRISPR/Cas9-mediated knockout approach,

we screened *P. tricornutum* for DNMT loss of function mutants (see method section).

In this work, we report five independent mutants with homozygous out of frame

deletions generating premature STOP codons in the coding sequence of DNMT5a

named 'M23', 'M25', '7C6', '7C7' and 'M26' DNMT5:KOs. Two mutants, M23 and M25

with frameshift in the coding sequence of DNMT5 were retained for further

investigation (Supplementary Figure 3). Those mutants are also homozygous for other

polymorphisms in the region (Supplementary Figure 3) that could suggest gene

conversion after Cas9 double strand breaks in *P. tricornutum* [44]. No DNMT3 or DNMT6

mutations could be generated using the CRISPR/Cas9 editing strategy.

Using sets of primer pairs targeting the DNMT domain as well as the DEADX

helicase-SNF2 like domain of DNMT5 transcripts, we detected by RT-qPCR a 4- to 5-

fold loss in mRNA levels in both M23 and M25 cell lines (Supplementary Figure 4a,

Supplementary Data 5). 5mC dot blot screening revealed that all DNMT5:KOs had a

4-5 fold loss of DNA methylation compared to the Pt1 8.6 reference ('wild-type')

(Supplementary Figure 4b, Supplementary Figure 5), consistent with the putative role

of DNMT5 in maintaining DNA methylation patterns in diatoms.

To generate a quantitative single base resolution of DNA methylation loss in

DNMT5:KOs, we performed whole genome bisulfite sequencing in M23, M25

(considered as two biological replicates) and the reference, Pt1 8.6 line. We filtered

cytosines by coverage depth considering a 5X coverage in all cell lines as a threshold

and computed CG methylation levels in TEs and genes. We found that CG methylation

is severely impaired in M23 and M25 compared to Pt1 8.6 cell lines (Figure 3a,b,

Supplementary Data 6). This is particularly observed within TEs that are the targets of

DNA methylation in *P. tricornutum* (Figure 3a,b). To get a quantitative view of the loss

of DNA methylation in DNMT5:KOs, we defined differentially methylated regions

(DMRs). We computed DMRs between DNMT5:KOs and WT lines using the bins built-

in DMRcaller[45] tools considering 100 bp bins with a minimal difference of +/- 20% DNA

methylation at CGs (5X coverage) in mutants compared to the Pt1 8.6 line. Those

thresholds were used based on the minimum coverage per cytosine and the

methylation characteristics in our sequencing data (Supplementary Figure 6a,b). We

identified 1715 and 1720 CG DMRs in M23 and M25, respectively (Supplementary

Data 7 and Supplementary Data 8), of which 96% are shared between both mutants

and show a consistent loss of DNA methylation upon knockout of DNMT5a (Figure 3c),

referred in this study as common hypoDMRs. CG common hypoDMRs cover ~0.8% of

the *P. tricornutum* genome. According to the distribution of DNA methylation in the

reference strain, we found that 14.90% (n=454) of annotated TEs are found within

common hypoDMRs (Figure 3d, Supplementary Data 9). In order to take into account

the possible methylation loss occurring in regulatory regions, gene and TE coordinates

were extended by 500 bp and 1 kb, respectively, upstream and downstream of their

start and end sites, considering that intergenic length in *P. tricornutum* varies between

1 kb and 1.5 kb[27]. As a result, respectively 7.76% and 12.23% of TEs are found within

500 bp and 1 kb of common hypoDMR coordinates (Figure 3d, Supplementary Data

9). Consistent with their low level of CG DNA methylation observed in both cell lines,

we found a comparatively low overlap of common hypoDMRs with genes or their

regulatory regions (Figure 3d, Supplementary Data 9). We then asked whether these

common hypoDMRs associate with known regions marked by histone post-

translational modifications. Genomic coordinates of common hypoDMRs overlapped

with previously mapped histone post-translational modification peaks[43]. The number

of common hypoDMRs overlapping with each combination of histone marks is shown

in Figure 3e (Supplementary Data 10). Interestingly, we found that between 80 and

90% of these common hypoDMRs (set size >1500, Figure 3e) overlap with known

regions marked by H3K27me3, H3K9me3 or H3K9me2 defined in the reference Pt1

8.6 line[43]. In addition, 963 (53%) of the common hypoDMRs are found within regions

co-marked by all three repressive histone marks (Figure 3e). This is consistent with

the observation that highly methylated regions described by restriction methylation-

sensitive sequencing (Mcrbc-Chip) also associate with such histone marks[27]. Our data

are consistent with a global loss of DNA methylation in DNMT5:KOs at TE-rich DNA

methylated-H3K27me3, H3K9me2 and H3K9me3 marked regions in the *P. tricornutum*

genome.

361 Gene and TE expression in the absence of DNMT5a in *P. tricornutum*

The control of TEs by the DNA methyltransferase family is a key unifying feature

within eukaryotes[2]. Therefore, we monitored the transcriptional effect of DNMT5a loss

on genes in M23 and M25 backgrounds using whole RNA high throughput sequencing

(Methods section). Given the high level of DNA methylation observed at TEs compared

to genes, we asked whether our RNAseq data captured any TE overexpression that

could be linked to hypoDMRs. We thus analyzed TE-gene transcripts that correspond

to the expression of TE open reading frames (i.e., encoding reverse transcriptase and

integrases) but also genes with TE insertions (Figure 4a), domesticated TEs and mis-

annotated TE loci[27,46]. To identify the most significant changes in mRNA levels, we

focused our analysis on genes and TE-genes showing a significant 2-fold induction or

reduction of expression in mutants compared to the reference line (|LFC| > 1 and an

FDR < 0.01, Supplementary Data 11). In M23 and M25, respectively, a total of 1732

and 806 genes and TE-genes are overexpressed while downregulation was observed

for 1152 and 248 genes and TE-genes (Figure 4b). Stable expression (-1 < LFC < 1

and FDR < 0.01) is observed for 943 genes and TE-genes in M23 and 216 genes and

TE-genes in M25. We found that 557 genes are overexpressed in both cell lines (M23

∩ M25). A total of 225 genes are overexpressed in M25 only (M25-spe) and 1126 are

overexpressed in M23 only (M23-spe). Significantly upregulated genes in both mutants

show consistent overexpression levels (Figure 4c).

In line with the hypothesis that TEs and not genes are directly regulated by DNA

methylation in *P. tricornutum,* we found that 338 TE-genes are upregulated in both

mutants (Figure 4d) which correspond to 56% of overexpressed TE-genes. Gene

ontology (GO) analysis showed that the upregulated TE-genes are enriched in DNA

integration biological function indicating that they mainly correspond to *bona fide* TE

annotations (Figure 4d). In contrast, only 219 (16%) of protein coding genes are

overexpressed in both mutants. They show clear enrichment for GOs associated with

protein folding as well as nucleotide phosphate metabolism and nucleotide binding activity (Figure 4e, Supplementary Data 12). This is typified by the overexpression of chaperone DnaJ domain-containing proteins and Hsp90-like proteins (Supplementary Data 13). The downregulation of genes was not consistent between M23 and M25 as only 35 genes and 16 TE-genes are downregulated in both cell lines (Figure 4f,g, Supplementary Data 14). Expression levels of 12 genes was confirmed by qPCR in the M23 cell line, including DnaJ and HSP90-like protein coding genes mentioned previously (Supplementary Figure 7a,b, Supplementary Data 15). Of note DNMT5a (Phatr3_EG02369) is among the downregulated genes in both mutants (Supplementary Data 14), consistent with qPCR results.

Relationship between CG methylation and expression of TE-genes in *P. tricornutum*

The observed overexpression of TEs in DNMT5:KOs could be directly due to the loss of DNA methylation. To test this, we first determined DNA methylation levels in the 600 upregulated TE-genes in the DNMT5:KO lines (Figure 5a, Supplementary Data 16). For each TE, we also computed the mean-centered normalized LFC (z-score) for each of the M23 and M25 lines (Figure 5a, Supplementary Data 16). We found that the TE-genes with the highest LFC (z-score >2) in the mutants are associated with higher DNA methylation levels in the reference strain. This is the case for each mutant independently, indicating that TEs with the highest upregulation in the DNMT5:KO lines are direct targets of DNA methylation in the reference strain.

We then assessed the relationship between upregulated TE-genes and the common hypoDMRs and found that 62% of upregulated TE-genes are found within these DMRs (Figure 5b). Importantly, this was the case only for TE-genes with overexpression in both cell lines (M23 ∩ M25) and not for M23-spe and M25-spe upregulated TE-genes (Figure 5b). This also means that 40% of upregulated TE-genes cannot be explained by the loss of DNA methylation alone. Similarly, downregulation and stable expression is not associated with common hypoDMRs (Figure 5b). This shows that TE-genes with consistent upregulation are specifically due to the loss of DNA methylation while other TE-gene misregulation is due to cell line specific DNA methylation-independent regulation. Among the 128 upregulated TE-genes in both mutants that are not direct targets of DNA methylation, we found a common hypoDMR in the regulatory region of

42 (in M23) and 15 TE-genes (in M25), respectively, indicating that DNA methylation

loss at these regions was also responsible for their upregulation (Figure 5c).

Next, we assessed TE families as annotated previously[46] (Figure 5d). We find that

overexpressed TE-genes are mostly represented by Copia-like in diatoms (CoDi)

retrotransposons of the CoDi1, CoDi2, CoDi4 and CoDi5 families with a minority of

DNA transposons as the PiggyBack family (Figure 5d). We noticed that the TE families

are found in similar proportions among TEs that overlap the common hypoDMRs and

those that do not. However, when we compared TE lengths, TEs that are upregulated

and overlap with common hypoDMRs are longer than upregulated TEs that are not

overlapping with hypoDMRs (Figure 5e, Supplementary Data 17). This suggests that

younger TEs tend to be direct targets of DNA methylation compared to evolutionary

older TEs family members. Subsequently, loss of DNA methylation causes

upregulation of mainly younger TEs. Filloramo et al.[47] recently described 85 long-LTR-

copia-like (LTR-copia) TEs based on reannotation of the *P. tricornutum* genome by

Oxford Nanopore Technologies long-read sequencing. Such TEs are considered as

potentially still active[47]. They are represented by Copia-like in diatoms (CoDi) of the

CoDi5, CoDi4 and CoDi2 families[47] that corresponds to the TE families found

overexpressed in our datasets (Figure 5d). Accordingly, we found that 75/85 of LTR-

copia are targets of DNA methylation and are associated with common hypoDMRs

(Supplementary Data 18). In addition, by overlapping TE-genes and genomic locations

of LTR-copia, we found that 61/75 of LTR-copia are overexpressed in both mutants

(Supplementary Data 18). Of note, our RNAseq data thus also support the presence

of these TEs in the reference Pt1 8.6 cell line as potentially still active elements. An

example of upregulation at LTR-copia is shown in Figure 5f. Additional shorter TEs

with overexpression also belong to CoDi5, CoDi4 and CoDi2 TE categories suggesting

that an active expression might still remain. Altogether, this strongly suggests that DNA

methylation is involved in the repression of young TEs in the *P. tricornutum* genome.

Discussion

Studies on the evolutionary history of DNMTs have established that the DNA

methylation machinery diverged among eukaryotes along with their respective DNA

methylation patterns[2,11]. However, the diversity of DNMTs found in SAR lineages is

underexplored due to the lack of representative sequences. Based on MMETSP

transcriptomes, we set out to explore the diversity and phylogeny of DNMTs in early

diverging eukaryotes. Besides the absence of genomic sequences, the MMETSP

database only encompasses expressed transcripts from cultured organisms and is

thus deprived of lowly expressed genes and condition-specific expressed genes.

Absence of a given gene family within a species should therefore be interpreted

accordingly. When our analysis found multiple distinct transcripts sharing the same

DNMT subfamily, as in diatoms, we used the most probable open reading frame

translation of the transcripts using eDAF curation to produce our phylogenetic tree.

However, without genomic annotations we cannot rule out that such transcripts result

from alternative transcription originating from a single gene or multi-copy gene families.

Our data are best interpreted when multiple transcripts and annotated genes,

whenever possible, are available, rather than at the species-specific level.

We nonetheless confirm that stramenopiles and dinoflagellates encode a

divergent set of DNMT proteins including DNMT3 and DNMT6 which have no

chromatin associated domains. In addition, our study independently reports the same

DNMT6 enzymes found in the raphidophyceae, *Bigelowella natans* and

Aplanochytrium stochhinoi by earlier work although not specified by the authors[26]. As

reported in trypanosomes[10], we suggest that DNMT6 likely emerged prior to the

Chromalveolata radiation. In trypanosomes, its presence in several lineages does not

predict DNA methylation *per se* and must be further investigated[48].

The DNMT5 enzymes are also very well represented both at the genomic and

transcriptomic levels, even outside the SARs, and are thus likely ancestral to

eukaryotes. We show here that the DNMT domains among the different DNMT5s are

conserved but show a divergence compared to other DNMTs, thus supporting a

common evolutionary origin for all DNMT5 enzymes. DNMT5b enzymes could be

multifunctional enzymes as suggested by the presence of N-terminal HAND domains

found in chromatin remodelers[49], by TUDOR domains found in histone modifying

enzymes, histone post-translational modification readers[50] as well as small RNA

interacting proteins[51,52]. They also contain an SNF2 ATPase domain[11] which plays a

chaperone-like enzyme-remodeling role important for DNA methylation and its

targeting to specific sites[15,53]. DNMT5c enzymes are likely very divergent DNMT5

enzymes that lack ATP-ase SNF domains. The diversity of DNMT5 domains is likely

inherent to its functioning and interaction with other epigenetic processes such as

histone modifications and non-coding RNA. In mammalian cells, TUDOR domain

containing UHRF1 is known to target DNMT1, the functional homologue of DNMT5,

onto newly synthesized DNA substrates during semi conservative DNA replication[54].

Furthermore, TUDOR domain of UHRF1 was reported to play an important role in the

recognition of histone H3K9 methylation[55,56]. While UHRF1, DNMT1 and ATPase

protein containing domains are separate in animals, they form an unusual

multifunctional domain protein in DNMT5 in photosynthetic microeukaryotes. This

domain architecture might be due to the compact genomes of the green microalgae

Micromonas pusilla, Ostreoccocus tauri and Bathycoccus prasinos11.

In our phylogenetic study, the RID/DMTA and diatom DNMT4 enzymes are

related, as shown previously by Huff and Zilberman[11] and Punger and Li[10]. In our case,

because the analysis covers a large evolutionary distance, phylogenetic relationships

between DNMT families should be interpreted accordingly. Therefore, we cannot rule

out the possibility that diatoms and RID families are paraphyletic. The function of

DNMT4 or DNMT4-type enzymes in diatoms is unknown. Among the four diatoms with

a known methylation pattern on TEs, two are lacking DNMT4s (including *P. tricornutum*

presented in this study). The presence of chromodomains known to bind histone post-

translational modifications as in CMT enzymes[57] nonetheless suggests that diatom

DNMT4 might be functional as either a *de novo* or a maintenance enzyme. The lack of

chromatin-associated domains in DNMT3, DNMT6 and other DNMT4 proteins suggest

that the link, if any, between DNA methylation and histone modifications is more

indirect than observed in plants and mammals and might require the activity of

accessory proteins like UHRF1-type[54] or DNMT3-like[58] enzymes that should be further

investigated.

Examining the role of DNMT5a in the pennate diatom *P. tricornutum,* we found

that it is an orthologue of the single DNMT5a protein from *Cryptococcus neoformans,*

which is involved in the maintenance of DNA methylation[11,15]. In that regard, our study

demonstrates that the loss of DNMT5a was sufficient alone to generate a global loss

of CG methylation in *P. tricornutum* similar to *Cryptococcus neoformans*[11]. We further

confirm that TEs are major targets of DNA methylation in diatoms. In addition, DMR

analysis identified regions essentially composed of TEs that show extensive

methylation in the reference strain. HypoDMRs overlap with regions marked by

H3K27me3 but also H3K9me3 which suggest that histone post-translational

modifications and DNA methylation cooperate to maintain TE repression. Genes

appear not to be the primary targets of DNA methylation. Only 51 out of all 9,416

annotated genes are found within DMRs. Among them, 19 were upregulated in both KO mutants. TE methylation is observed in other diatoms such as *F. cylindrus*[11] and *T. pseudonana*[11] where the targeted TEs have low expression[11]. However, those species encode a different set of DNMTs compared to *P. tricornutum. T. pseudonana* appears to lack DNMT5a and has a partial DNMT6 protein while *F. cylindrus* encodes all but DNMT3 (Supplementary Data 3). It is possible that DNMTs show partial functional redundancy in diatoms. In that regard, the DNMT5:KO cell lines presented in this study could be used as a heterologous expression system to decipher the role of other DNMTs in diatoms.

Compared to DNA methylation loss that is observed in different DNMT5:KO cell lines (Supplementary Figure 4, Supplementary Figure 5), gene expression was more inconsistent between cell lines, including when assessed by qPCR validation. We thus make the hypothesis that gene expression is mainly cell line specific in DNMT5:KO cell lines. This divergence in gene expression could be linked to the random insertions of plasmids generated by biolistic transformation. Alternatively, *de novo* and likely random TE insertions upon DNA methylation loss could generate gene expression divergence between cell lines over time.

In our study, we found that 15% of TE-genes are upregulated in the DNMT5:KO cell lines, less than observed in *Arabidopsis thaliana* where the loss of DDM1 (involved in the maintenance of DNA methylation) caused the expression of about 40% of all TE-genes[59]. However, in *P. tricornutum* we found that overexpression and methylation levels are particularly relevant for TEs that have been identified as full length potentially still active LTR-copia elements. Interestingly, in *Arabidopsis thaliana*, the most mobile TEs between different accessions are regulated by the MET2a protein, likely involved in DNA methylation and repression[60]. In addition, such TEs expansion associates with null or loss of function alleles of MET2a[60]. When comparing *P. tricornutum* and *T. pseudonana* genomes, the CoDi2 and CoDi4 families are the main contributors of retrotransposon expansion in *P. tricornutum*[61] although CoDi2 is only found in *P. tricornutum*. We found such TEs to be overexpressed in response to DNA methylation loss. Therefore, DNA methylation seems to be a genome integrity keeper in *P. tricornutum*. Other smaller TEs in the form of TE-genes are also upregulated and may retain some activity in *P. tricornutum*. Upregulation was also observed for TEs that were not targets of DNA methylation in the reference strain but for which a subset was nonetheless found within a 1 kb distance from hypoDMRs, suggesting that initial

repression is likely linked to DNA methylation spreading or proximity which was
reported in a previous work[27].

Highly repetitive TE families are removed in our analysis since only uniquely
mapped reads were aligned. This is true for both transcriptomic and bisulfite
sequencing data. In addition, our transcriptomic analysis is only a snapshot of all TEs
overexpressed at a given time in *P. tricornutum* cell populations. The loss of DNA
methylation could trigger more misregulation of TEs in stress culture conditions, as
previously reported upon nitrogen depletion[27] and exposure to the toxic reactive
aldehyde[61]. DNMT5 mutant cell lines are viable in standard culture conditions used for
P. tricornutum suggesting that co-occurring repressive histone marks reported in
previous studies might be compensating the loss of DNA methylation[35,43]. Additionally,
this suggests that in optimal conditions, loss of DNA methylation is not associated with
drastic biological effects, supporting the lack of a phenotypic response which is
otherwise seen in stress conditions, typically slow growth, smaller cell size and an
atypical morphology. Our study provides direct understanding of DNA methylation
regulation and its role in diatoms offering a robust foundation for future studies in
eukaryotes to further comprehend DNA methylation function and evolution.

Methods
HMMER and Reciprocal Blasy best Hit analysis
We performed an extensive scan of the MMETSP database, enriched with 7 diatom
transcriptomes and genomes from the top 20 most abundant diatoms found in *Tara*
Oceans database[62], using HMMER-search with the model PF00145 to fetch any
DNMT-like, including partial transcripts, sequence within microeukaryotes. We ran
HMMER in a non-stringent fashion to not miss positives DNMT sequences. We used
eDAF approach to filter the expected high number of false positives. It is worth noting
that we initially use HMMER for screening instead of the built-in module of eDAF due
to the time complexity of the latter for extensive searches (tens to hundreds of times
slower than HMMER). Reciprocal BLAST best hit analysis was performed as
previously described[63]. Briefly, the DNMT3 (Phatr3_J47136), DNMT4
(*Thaps3_11011*), DNMT5 (*Phatr3_EG02369*) and DNMT6 (Phatr3_J47357)
orthologues found in *P. tricornutum* or *T. pseudonana* (for DNMT4) were blasted
against a phylogenetically optimized database that include MMETSP transcriptomes.
Putative DNMT sequence hits giving back the corresponding enzyme (DNMT3,

DNMT4, DNMT5 or DNMT6) at the threshold of e-value of 1×10^{-5} in the corresponding

diatom were retained. Candidate enzymes were then analyzed using eDAF.

eDAF-guided domain architecture analysis

enhanced Domain Architecture Framework (eDAF) is a four module computational tool

for gene prediction, gene ontology and functional domain predictions[35]. As previously

described for Polycomb and Trithorax enzymes[35], candidate DNMTs identified by RBH

and HMMER-search were analyzed using the DAMA-CLADE guided built-in functional

domain architecture to predict the occurrences of DNMT domains starting from protein

sequences. The domain architecture of representative enzymes used in this study can

be found in Supplementary Data 4.

Phylogenetic tree analysis

The DNMT domain of candidate enzymes were aligned using ClustalΩ[64] (HHalign

algorithm). The alignment was manually curated and trimmed using trimAL (removing

>25% gap column) to align corresponding DNMT motifs in all gene families. A

convergent phylogenetic tree was then generated using the online CIPRES Science

gateway program[65] using MrBAYES built-in algorithm. Default parameters were used

with the following specifications for calculation of the posterior probability of partition:

sumt.burninfraction=0.5, sump.burningfraction=0.5, 10000000 generations, sampling

each 100. We also used two different models: Estimating the Fixed Rate and GTR.

Cell cultures

Axenic *P. tricornutum* CCMP2561 clone Pt1 8.6 cultures were obtained from the

culture collection of the Provasoli-Guillard National Center for Culture of Marine

Phytoplankton (Bigelow Laboratory for Ocean Sciences, USA.). Cultures were grown

in autoclaved and filtered (0.22 µM) Enhanced Sea Artificial Water (ESAW -

https://biocyclopedia.com/index/algae/algal_culturing/esaw_medium_composition.ph

p) medium supplemented with f/2 nutrients and vitamins without silica under constant

shaking (100rpm). Cultures were maintained in flasks at exponential state in a

controlled growth chamber at 19°C under cool white fluorescent lights at 100 µE m−2

s−1 with a 12h photoperiod. For RNA sequencing and bisulfite experiments, WT and

DNMT5 mutant cultures were seeded in duplicate at 10.000 cells/ml and grown side

by side in 250ml flasks until early-exponential at 1.000.000 cells/ml. Culture growth

was followed using a hematocytometer (Fisher Scientific, Pittsburgh, PA, USA). Pellets

were collected by centrifugation (4000rpm) washed twice with marine PBS

(http://cshprotocols.cshlp.org/content/2006/1/pdb.rec8303) and flash frozen in liquid

nitrogen. Cell pellets were kept at -80°C until use. For bisulfite sequencing, technical

duplicates were pooled to get sufficient materials.

CRISPR/Cas9 mediated gene disruption

The CRISPR/Cas9 knockouts were performed as previously described[44]. Our strategy

consisted in the generation of short deletions and insertions to disrupt the open reading

frame of putative DNMTs of *P. tricornutum*. We introduced by biolistic the guide RNAs

independently of the Cas9 and ShBle plasmids, conferring resistance to Phleomycin,

into the reference strain Pt1 8.6 (referred hereafter as 'reference line' or 'wild-type'-

WT). Briefly, specific target guide RNAs were designed in the first exon of

Phatr3_EG02369 (DNMT5), Phatr3_J47357 (DNMT6) and Phatr3_J36137 (DNMT3)

using the PHYTO/CRISPR-EX[66] software and cloned into the pU6::AOX-sgRNA

plasmid by PCR amplification. For PCR amplification, plasmid sequences were added

in 3' of the guide RNA sequence (minus –NGG) which are used in a PCR reaction with

the template pU6::AOX-sgRNA. Forward primer – sgRNA seq +

GTTTTAGAGCTAGAAATAGC. Reverse primer - sequence to add in 3' reverse

sgRNA seq + CGACTTTGAAGGTGTTTTTTG. This will amplify a new pU6::AOX-

(your_sgRNA). The PCR product is digested by the enzyme DPN1 (NEB) in order to

remove the template plasmid and cloned in TOPO10 *E. coli*. The sgRNA plasmid, the

pDEST-hCas9-HA and the ShBLE Phleomycin resistance gene cloned into the plasmid

pPHAT-eGFP were co-transformed by biolistic in the Pt1 8.6 'Wild Type' strain as

described in[44]. We also generated a line that was transformed with pPHAT-eGFP and

pDEST-hCas9-HA but no guide RNAs. This line is referred as the Cas9:Mock line.

CRISPR/Cas9 – sgRNA transformants were selected by phleomycin resistance

(carried by the plasmid pPHAT-eGFP). 48 hours post-transformation, diatoms were

replated and grown on phleomycin 100µg/ml 50% Enhanced Artificial Sea Water[67]

plates until single colonies appeared (2-3 weeks). Transformants were isolated from

plates and sanger sequenced after PCR using HGS Diamond Taq® as per

manufacturer instruction and the primers DNMT5locus_R and DNMT5locus_F

(Supplementary Data 19). These primers allow the amplification and sequencing of

both alleles of DNMT5. Sequencing using the primer DNMT5locus_R shows that

mutants are homozygous for the deletions. In addition, all mutants show loss of heterozygosity (LOH).

RNA and DNA extraction

Total RNA extraction was performed by classical TRIZOL/Chloroform isolations and precipitation by isopropanol. Frozen cell pellets were extracted at a time in a series of 3 technical extraction/duplicates and pooled. RNA was DNAse treated using DNAse I (ThermoFisher) as per manufacturer's instructions. DNA extraction was performed using the Invitrogen™ Easy-DNA™ gDNA Purification Kit following 'Protocol #3' instructions provided by the manufacturer. Extracted nucleic acids were measured using QUBIT fluorometer and NANODROP spectrometer. RNA and gDNA Integrity were controlled by electrophoresis on 1% agarose gels.

RT-qPCR analysis

 qPCR primers were designed using the online PRIMER3 program v0.4.0 defining 110-150 amplicon size and annealing temperature between 58°C and 62°C. Primer specificity was checked by BLAST on *P. tricornutum* genome at ENSEMBL. For cDNA synthesis, 1µg of total RNA was reverse transcribed using the SuperScript™ III First-Strand (Invitrogen) protocol. For quantitative reverse transcription polymerase chain reaction (RT-qPCR) analysis, cDNA was amplified using SYBR Premix ExTaq (Takara, Madison, WI, USA) according to manufacturer's instructions. CT values for genes of interest were generated on a Roche lightcycler® 480 qpcr system. CT values were normalized on housekeeping genes using the deltaCT method. Normalized CT values for amplifications using multiple couples of primers targeting several cDNA regions of the genes of interest were then averaged and used as RNA levels proxies. Primers used in this study are listed in Supplementary Data 19.

Dot blot

gDNA samples were boiled at 95°C for 10 min for denaturation. Samples were immediately placed on ice for 5 min, and 250-500 ng were loaded on regular nitrocellulose membranes. DNA was then autocrosslinked in a UVC 500 crosslinker – 2 times at 1200uj (*100). The membranes were blocked for 1 hr in 5% PBST-BSA. Membranes were probed for 1 hr at room temperature or overnight at 4°C with 1:1000 dilution of 5mC antibody (OptimAbtm Anti-5-Methylcytosine – BY-MECY 100). 5mC

signals were revealed using 1:5000 dilution of HRP conjugated antirabbit IgG

secondary antibody for 1 hr at room temperature followed by chemo luminescence.

Loading was measured using methylene blue staining.

RNA and Bisulfite sequencing

RNA libraries were prepared by the FASTERIS Company (https://www.fasteris.com).

Total RNA was polyA purified and libraries were prepared for illumina NextSeq

sequencing technologies. For RNAseq analysis, two biological replicates per mutant

were used (M23 and M25). In addition, two biological replicates of a Pt1 8.6 line was

sequenced in the same run as a control (total of 6 samples). Bisulfite libraries and

treatments were performed by the FASTERIS Company and DNA was sequenced on

an Illumina NextSeq instrument. 150bp paired-end reads were generated with 30X

coverage. A 5mC map was also generated in the reference Pt1 8.6 line as a control.

RNAseq analysis

150bp paired-end sequenced reads were subjected to quality control with FastQC

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc). Then, the reads were

aligned on the reference genome of *P. tricornutum* (Phatr3)[46] using STAR (v2.5.3a)[68].

Gene expression levels were quantified using HTseq v0.7.2. Differentially expressed

genes were analyzed using DESeq2 v1.19.37[69] with the following generalized linear

model: ~mutation. FDR values are corrected p.values using the Benjamin and

Hochberg method. Genes are designed significant (DEGs) if the |log2FC| > 1 and the

FDR < 0.05. GO enrichments were calculated using the overrepresentation Fisher's

exact test described in topGO v2.44.0[70]. For each analysis, appropriate DEGs have

been used as input and a GO theme is considered as significant if the p.value < 0.05.

Bisulfite sequencing analysis

Bisulfite   analysis   was   performed   using   Bismark-bowtie   2

(https://www.bioinformatics.babraham.ac.uk/projects/bismark/). We used the default

Bowtie2 implementation of bismark with the specifications that only uniquely mapped

reads should be aligned. All alignments were performed with high stringency allowing

for only one base mismatch ($n = 1$). We also clearly specified that no discordant pairs

of the pair-end reads should be aligned. DNA methylation in the CG, CHG and CHH

contexts was calculated by dividing the total number of aligned methylated reads by the total number of methylated plus un-methylated reads.

DMR calling

Differentially methylated regions were called using the DMRcaller R package v1.22.0[45]. Given the low level of correlation of DNA methylation observed in *P. tricornutum*[11,27] and sequencing coverage in all three cell lines, only cytosines with coverage >=5X in all three lines were kept for further analysis and the bins strategy was favored over other built-in DMRcaller tools. DMRs were defined as 100bp regions with at least an average 20% loss/gain of DNA methylation in either one of the DNMT5:KOs compared to the reference strain. The 'Score test' method was used to calculate statistical significance and threshold was set at p.value <0.01. In addition, to distinguish isolated differentially methylated cytosines from regions with significant loss of DNA methylation, an hypoDMR must contain at least methylated 2 CpG in the reference strain.

Overlap with histone modifications and genomic annotations.

Analysis on bed files were performed using bedtools v2.27.1.[71]. Bedtools intersect with default parameters was used to calculate overlap regions of DMRs with genes and TE-genes. Bedtools window has been used to compute the 500 bp and 1kb upstream and downstream near regions between DMRs, genes and TE-genes.

Percentage overlaps between DMRs as well as the overlap of gene and TE coordinates with histone modifications and DMRs were calculated using the genomation R package v1.22.0[72] and the 'annotateWithFeature' and 'getMembers' functions. For RNAseq analysis, we analyzed the expression of TE-genes as previously described[46]. To define TE-genes in DMRs we crosschecked overlapping TE-genes annotations with *bona fide* TEs in DMRs using 'annotatewithFeature' function. UpSet plots were computed using UpSetR v1.4.0.[73] Heatmaps were produced using the R package ComplexHeatmap (v2.8.0)[74]. Sankey diagram was produced with the R package highcharter (v0.9.4) (https://jkunst.com/highcharter/authors.html). TEs that mapped to less than 3 members of a TE family were discarded. All R plots were obtained using R version 4.0.3.

Data availability

The raw data have been deposited at Gene Expression Omnibus GEO (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE186857). Bisulfite sequencing raw data and bigwig files showing methylation rates (#methylated C/#total number of C) in the context of CHH, CHG and CpG, where H: is A, C or T in the WT, M23 and M25 are under the accession number GSE186855. The raw RNA sequencing data and the TPM counting table are under accession GSE186856.

All the cell lines used in the study, including DNMT5:KOs, are available upon request.

Statistics and Reproducibility

The statistical tests used in this study are indicated in the respective figure legend or in the methods section. We used the function lm() from R to fit the linear model between LFC in the M23 and M25 cell lines. The pairwise Wilcoxon rank-sum test has been used to check for significant differences in TEs lengths between two distributions using the pairwise wilcox.test() function in R (Figure 5e). For multiple-hypotheses testing, p.values were adjusted with the Benjamini-Hochberg procedure. No statistical methods were used to predetermine sample size.

Code availability

All the code that has been used to generate the results in this paper is available from the lead contact upon reasonable request.

Acknowledgements

We thank Catherine Cantrel form IBENS for media preparation. L.T. acknowledges funds from the CNRS, the region of Pays de la Loire (ConnecTalent EPIALG project) and Epicycle ANR project (ANR-19-CE20- 0028-02). C.B. acknowledges funding from the ERC Advanced Awards Diatomite. F.Y. was supported by a PhD fellowship from the Chinese Scholarship Council (CSC-201906310152).

Competing interests

The authors declare no competing interests.

Author Contributions

A.H. and L.T. conceived the study. A.H., F.R.J.V, O.A.M. and L.T. designed the study. L.T. supervised and coordinated the study. A.H. performed the experiments. CB contributed to the discussion of the results. AuGe contributed to the bioinformatic analysis. A.G. grew the mutants and extracted RNA for validation experiments. F.Y. performed QPCR work and gene validation analysis. O.A.M. performed and supervised A.H. for the bioinformatic analysis of RNAseq, gene ontology and bisulfite seq data. A.H. performed the DMR analysis under the supervision of O.A.M. F.R.J.V and A.H. analyzed HMMER, DAMA/CLADE and eDAF data. All authors analyzed and interpreted the data. A.H. and L.T. wrote the manuscript with input from all authors.

References

1.   Kato, M., Miura, A., Bender, J., Jacobsen, S. E. & Kakutani, T. Role of CG and non-CG methylation in immobilization of transposons in Arabidopsis. *Curr. Biol.* (2003) doi:10.1016/S0960-9822(03)00106-4.

2.   Zilberman, D. The evolving functions of DNA methylation. *Current Opinion in Plant Biology* (2008) doi:10.1016/j.pbi.2008.07.004.

3.   Barlow, D. P. & Bartolomei, M. S. Genomic imprinting in mammals. *Cold Spring Harb. Perspect. Biol.* (2014) doi:10.1101/cshperspect.a018382.

4.   Galupa, R. & Heard, E. X-Chromosome Inactivation: A Crossroads Between Chromosome Architecture and Gene Regulation. *Annu. Rev. Genet.* (2018) doi:10.1146/annurev-genet-120116-024611.

5.   Bestor, T. H. DNA methylation: evolution of a bacterial immune function into a regulator of gene expression and genome structure in higher eukaryotes. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* (1990) doi:10.1098/rstb.1990.0002.

6.   Kumar, S. *et al.* The DNA (cytosine-5) methyltransferases. *Nucleic Acids Research* (1994) doi:10.1093/nar/22.1.1.

7.   Cheng, X., Kumar, S., Klimasauskas, S. & Roberts, R. J. Crystal structure of the HhaI DNA methyltransferase. in *Cold Spring Harbor Symposia on Quantitative Biology* (1993). doi:10.1101/SQB.1993.058.01.039.

8.   Zemach, A. & Zilberman, D. Evolution of eukaryotic DNA methylation and the pursuit of safer sex. *Current Biology* (2010) doi:10.1016/j.cub.2010.07.007.

9.   Zemach, A., McDaniel, I. E., Silva, P. & Zilberman, D. Genome-wide
evolutionary analysis of eukaryotic DNA methylation. *Science (80-.).* (2010)
doi:10.1126/science.1186366.

10. Ponger, L. & Li, W. H. Evolutionary diversification of DNA methyltransferases in
eukaryotic genomes. *Mol. Biol. Evol.* (2005) doi:10.1093/molbev/msi098.

11. Huff, J. T. & Zilberman, D. Dnmt1-independent CG methylation contributes to
nucleosome positioning in diverse eukaryotes. *Cell* (2014)
doi:10.1016/j.cell.2014.01.029.

12. Greenberg, M. V. C. & Bourc'his, D. The diverse roles of DNA methylation in
mammalian development and disease. *Nature Reviews Molecular Cell Biology*
(2019) doi:10.1038/s41580-019-0159-6.

13. Seritrakul, P. & Gross, J. M. Expression of the De Novo DNA
Methyltransferases (dnmt3-dnmt8) During Zebrafish Lens Development.
doi:10.1002/dvdy.24077.

14. Kouzminova, E. & Selker, E. U. Dim-2 encodes a DNA methyltransferase
responsible for all known cytosine methylation in Neurospora. *EMBO J.* (2001)
doi:10.1093/emboj/20.15.4309.

15. Dumesic, P. A., Stoddard, C. I., Catania, S., Narlikar, G. J. & Madhani, H. D.
ATP Hydrolysis by the SNF2 Domain of Dnmt5 Is Coupled to Both Specific
Recognition and Modification of Hemimethylated DNA. *Mol. Cell* (2020)
doi:10.1016/j.molcel.2020.04.029.

16. Gladyshev, E. Repeat-Induced Point Mutation and Other Genome Defense
Mechanisms in Fungi. *Microbiol. Spectr.* (2017)
doi:10.1128/microbiolspec.funk-0042-2017.

17. Galagan, J. E. & Selker, E. U. RIP: The evolutionary cost of genome defense.
Trends in Genetics (2004) doi:10.1016/j.tig.2004.07.007.

18. Yang, K. *et al.* The DmtA methyltransferase contributes to Aspergillus flavus
conidiation, sclerotial production, aflatoxin biosynthesis and virulence. *Sci.*
Rep. (2016) doi:10.1038/srep23259.

19. Malagnac, F. *et al.* A gene essential for de novo methylation and development
in ascobolus reveals a novel type of eukaryotic DNA methyltransferase
structure. *Cell* (1997) doi:10.1016/S0092-8674(00)80410-9.

20. Bewick, A. J., Vogel, K. J., Moore, A. J. & Schmitz, R. J. Evolution of DNA
methylation across insects. *Mol. Biol. Evol.* (2017)

doi:10.1093/molbev/msw264.

21. Feng, S. *et al.* Conservation and divergence of methylation patterning in plants
and animals. *Proc. Natl. Acad. Sci. U. S. A.* (2010)
doi:10.1073/pnas.1002720107.

22. Vanyushin, B. F. & Ashapkin, V. V. *DNA methylation in plants. DNA*
Methylation in Plants (2011). doi:10.1146/annurev.arplant.49.1.223.

23. Zhong, X. *et al.* Molecular mechanism of action of plant DRM de novo DNA
methyltransferases. *Cell* (2014) doi:10.1016/j.cell.2014.03.056.

24. Jeltsch, A. *et al.* Mechanism and biological role of Dnmt2 in Nucleic Acid
Methylation. *RNA Biology* (2017) doi:10.1080/15476286.2016.1191737.

25. Goll, M. G. *et al.* Methylation of tRNAAsp by the DNA methyltransferase
homolog Dnmt2. *Science* **311**, 395–398 (2006).

26. De Mendoza, A. *et al.* Recurrent acquisition of cytosine methyltransferases into
eukaryotic retrotransposons. *Nat. Commun.* (2018) doi:10.1038/s41467-018-
03724-9.

27. Veluchamy, A. *et al.* Insights into the role of DNA methylation in diatoms by
genome-wide profiling in Phaeodactylum tricornutum. *Nat. Commun.* (2013)
doi:10.1038/ncomms3091.

28. Fan, X. *et al.* Single-base methylome profiling of the giant kelp Saccharina
japonica reveals significant differences in DNA methylation to microalgae and
plants. *New Phytol.* (2020) doi:10.1111/nph.16125.

29. Armbrust, E. V. The life of diatoms in the world's oceans. *Nature* (2009)
doi:10.1038/nature08057.

30. Malviya, S. *et al.* Insights into global diatom distribution and diversity in the
world's ocean. *Proc. Natl. Acad. Sci. U. S. A.* (2016)
doi:10.1073/pnas.1509523113.

31. Traller, J. C. *et al.* Genome and methylome of the oleaginous diatom Cyclotella
cryptica reveal genetic flexibility toward a high lipid phenotype. *Biotechnol.*
Biofuels (2016) doi:10.1186/s13068-016-0670-3.

32. Lister, R. *et al.* Human DNA methylomes at base resolution show widespread
epigenomic differences. *Nature* (2009) doi:10.1038/nature08514.

33. Maumus, F., Rabinowicz, P., Bowler, C. & Rivarola, M. Stemming Epigenetics
in Marine Stramenopiles. *Curr. Genomics* (2011)
doi:10.2174/138920211796429727.

34.   Keeling, P. J. *et al.* The Marine Microbial Eukaryote Transcriptome Sequencing

Project (MMETSP): Illuminating the Functional Diversity of Eukaryotic Life in

the Oceans through Transcriptome Sequencing. *PLoS Biol.* (2014)

doi:10.1371/journal.pbio.1001889.

35.   Zhao, X. *et al.* Probing the Diversity of Polycomb and Trithorax Proteins in

Cultured and Environmentally Sampled Microalgae. *Front. Mar. Sci.* (2020)

doi:10.3389/fmars.2020.00189.

36.   Strassert, J. F. H., Irisarri, I., Williams, T. A. & Burki, F. A molecular timescale

for eukaryote evolution with implications for the origin of red algal-derived

plastids. doi:10.1038/s41467-021-22044-z.

37.   Simões, T. R., Vernygora, O., Caldwell, M. W. & Pierce, S. E.

Megaevolutionary dynamics and the timing of evolutionary innovation in

reptiles. *Nat. Commun. 2020 111* **11**, 1–14 (2020).

38.   Czech, L., Huerta-Cepas, J. & Stamatakis, A. A Critical Review on the Use of

Support Values in Tree Viewers and Bioinformatics Toolkits. *Mol. Biol. Evol.*

34, 1535–1542 (2017).

39.   Kelk, S. Phylogenetic Networks: Concepts, Algorithms and Applications. *Syst.*

Biol. **61**, 174–175 (2012).

40.   Wang, J. *et al.* Structural insights into DNMT5-mediated ATP-dependent high-

fidelity epigenome maintenance. *Mol. Cell* **82**, 1186-1198.e6 (2022).

41.   Bewick, A. J. *et al.* Diversity of cytosine methylation across the fungal tree of

life. *Nat. Ecol. Evol.* (2019) doi:10.1038/s41559-019-0810-9.

42.   Jurkowska, R. Z., Jurkowski, T. P. & Jeltsch, A. Structure and Function of

Mammalian DNA Methyltransferases. *ChemBioChem* (2011)

doi:10.1002/cbic.201000195.

43.   Veluchamy, A. *et al.* An integrative analysis of post-translational histone

modifications in the marine diatom Phaeodactylum tricornutum. *Genome Biol.*

(2015) doi:10.1186/s13059-015-0671-8.

44.   Nymark, M., Sharma, A. K., Sparstad, T., Bones, A. M. & Winge, P. A

CRISPR/Cas9 system adapted for gene editing in marine algae. *Sci. Rep.*

(2016) doi:10.1038/srep24951.

45.   Catoni, M., Tsang, J. M. F., Greco, A. P. & Zabet, N. R. DMRcaller: A versatile

R/Bioconductor package for detection and visualization of differentially

methylated regions in CpG and non-CpG contexts. *Nucleic Acids Res.* (2018)

doi:10.1093/nar/gky602.

46.   Rastogi, A. *et al.* Integrative analysis of large scale transcriptome data draws a
comprehensive landscape of Phaeodactylum tricornutum genome and
evolutionary origin of diatoms. *Sci. Rep.* (2018) doi:10.1038/s41598-018-
23106-x.

47.   Filloramo, G. V., Curtis, B. A., Blanche, E. & Archibald, J. M. Re-examination of
two diatom reference genomes using long-read sequencing. *BMC Genomics*
22, 1–25 (2021).

48.   Cuypers, B. *et al.* The Absence of C-5 DNA Methylation in *Leishmania*
donovani Allows DNA Enrichment from Complex Samples. *Microorganisms* **8**,
1–18 (2020).

49.   Grüne, T. *et al.* Crystal structure and functional analysis of a nucleosome
recognition module of the remodeling factor ISWI. *Mol. Cell* (2003)
doi:10.1016/S1097-2765(03)00273-9.

50.   Lu, R. & Wang, G. G. Tudor: A versatile family of histone methylation 'readers'.
Trends in Biochemical Sciences (2013) doi:10.1016/j.tibs.2013.08.002.

51.   Pek, J. W., Anand, A. & Kai, T. Tudor domain proteins in development. *Dev.*
(2012) doi:10.1242/dev.073304.

52.   Tóth, K. F., Pezic, D., Stuwe, E. & Webster, A. The pirna pathway guards the
germline genome against transposable elements. in *Advances in Experimental*
Medicine and Biology (2016). doi:10.1007/978-94-017-7417-8_4.

53.   Catania, S. *et al.* Evolutionary Persistence of DNA Methylation for Millions of
Years after Ancient Loss of a De Novo Methyltransferase. *Cell* **180**, 816
(2020).

54.   Bostick, M. *et al.* UHRF1 plays a role in maintaining DNA methylation in
mammalian cells. *Science (80-.).* (2007) doi:10.1126/science.1147939.

55.   Nady, N. *et al.* Recognition of multivalent histone states associated with
heterochromatin by UHRF1 protein. *J. Biol. Chem.* **286**, 24300–24311 (2011).

56.   Rottach, A. *et al.* The multi-domain protein Np95 connects DNA methylation
and histone modification. *Nucleic Acids Res.* **38**, 1796 (2010).

57.   Tajul-Arifin, K. *et al.* Identification and Analysis of Chromodomain-Containing
Proteins Encoded in the Mouse Transcriptome. *Genome Res.* **13**, 1416–1429
(2003).

58.   Ooi, S. K. T. *et al.* DNMT3L connects unmethylated lysine 4 of histone H3 to de

novo methylation of DNA. *Nature* (2007) doi:10.1038/nature05987.

59. Osakabe, A. *et al.* The chromatin remodeler DDM1 prevents transposon

mobility through deposition of histone variant H2A.W. *Nat. Cell Biol.* **23**, 391–

400 (2021).

60. Quadrana, L. *et al.* The Arabidopsis thaliana mobilome and its impact at the

species level. *Elife* **5**, (2016).

61. Maumus, F. *et al.* Potential impact of stress activated retrotransposons on

genome evolution in a marine diatom. *BMC Genomics* (2009)

doi:10.1186/1471-2164-10-624.

62. Dorrell, R. G. *et al.* Phylogenomic fingerprinting of tempo and functions of

horizontal gene transfer within ochrophytes. doi:10.1073/pnas.2009974118/-

/DCSupplemental.

63. Dorrell, R. G. *et al.* Chimeric origins of ochrophytes and haptophytes revealed

through an ancient plastid proteome. *Elife* (2017) doi:10.7554/eLife.23717.

64. Madeira, F. *et al.* The EMBL-EBI search and sequence analysis tools APIs in

2019. *Nucleic Acids Res.* (2019) doi:10.1093/nar/gkz268.

65. Miller, M. A., Pfeiffer, W. & Schwartz, T. Creating the CIPRES Science

Gateway for inference of large phylogenetic trees. in *2010 Gateway Computing*

Environments Workshop, GCE 2010 (2010). doi:10.1109/GCE.2010.5676129.

66. Rastogi, A., Murik, O., Bowler, C. & Tirichine, L. PhytoCRISP-Ex: A web-based

and stand-alone application to find specific target sequences for CRISPR/CAS

editing. *BMC Bioinformatics* (2016) doi:10.1186/s12859-016-1143-1.

67. Vartanian, M., Desclés, J., Quinet, M., Douady, S. & Lopez, P. J. Plasticity and

robustness of pattern formation in the model diatom Phaeodactylum

tricornutum. *J. Compil. © New Phytol.* **182**, 429–442 (2009).

68. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**,

15–21 (2013).

69. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, (2014).

70. Alexa, A. & Maintainer, J. R. Package 'topGO' Type Package Title Enrichment

Analysis for Gene Ontology. (2022).

71. BEDTools: a flexible suite of utilities for comparing genomic features |

Genomics Gateway. http://bioscholar.com/genomics/bedtools-a-flexible-suite-

of-utilities-for-comparing-genomic-features/.

72. Akalin, A., Franke, V., Vlahoviček, K., Mason, C. E. & Schübeler, D.
Genomation: A toolkit to summarize, annotate and visualize genomic intervals.
Bioinformatics (2015) doi:10.1093/bioinformatics/btu775.
73. Lex, A. Sets and intersections. *Nat. Methods* **11**, 779 (2014).
74. Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and
correlations in multidimensional genomic data. *Bioinformatics* **32**, 2847–2849
(2016).

1003

1004

Figure 1

Phylogenetic analysis of DNMTs from MMETSP.

a. Convergent phylogenetic tree of DNMT domains from the MMETSP and reference
genome databases. The sequences selected were from microeukaryotes. Numbers
represent MrBAYES posterior probabilities. Grey branches represent bacterial and
viral DNA cytosine methyltransferase enzymes. We indicate the main lineages found
within each gene family using their corresponding colours next to the tree. **b.**
Schematic representation of the DAMA/CLADE structure of representative DNMT
enzymes. DNMT: DNA methyltransferase; RING: Ring zinc finger domain; DX: Dead
box helicase; Hter: C-terminus-Helicase; LBR: Laminin B receptor; RFD: Replication
Foci Domain; BAH: Bromo-Adjacent Homology; Chromo: Chromodomain; PHD: Plant
HomeoDomain.

1017

Figure 2

Summary of DNMT family lineages found in microeukaryotes.

Full crosses report the presence of a given gene family within lineages. Dashed lines
and crosses indicate the uncertainty in the eukaryotic phylogeny as well as low support
presence of a given DNMT family within lineages. Fungi that share DNMT families with
other eukaryotes presented in this study are shown for comparison purposes. SAR:
Stramenopile Alveolate Rhizaria lineage. Ochrophyte are secondary endosymbiont,
photosynthetic lineages of stramenopiles.

1026

1027 **Figure 3**

1028 **CG methylation profiles in DNMT5:KO lines.**

a. Heatmap of CG methylation levels in Pt18.6 reference (WT), M23 and M25 for TEs (left panel) and genes (right panel). **b.** Violin plot showing the methylation levels in all CGs found in TEs and genes in Pt1.86 and M25, M23. The white dots show the median. **c.** Venn diagram displaying the number of hypoDMRs identified in M23 (M23-spe) (yellow) and M25 (M25-spe) (orange). **d.** Percentages of overlap between common hypoDMRs, genes and TEs. **e.** Association between common hypoDMRs and regions targeted by histone post-translational modifications representative of the epigenetic landscape of *P. tricornutum*. The number of overlapping common hypoDMRs is shown for each histone modification and each combination of histone modifications.

Figure 4

Dynamics of gene and TE-gene expression in DNMT5:KO lines

a. Snapshot of an example TE-gene CG methylation profile. **b.** Differential expression in DNMT5:KOs (M23 and M25 are represented in the upper and lower parts of the volcano plot, respectively) compared with Pt18.6 reference (WT). The upper panel shows a volcano plot that displays the distribution of the fold changes (LFC) in the X-axis and adjusted p.values (-log10FDR) in the Y-axis. The number of genes up and downregulated in each mutant are indicated. The stable genes (1 < LFC > -1 and FDR < 0.01) are shown in grey. The lower panel shows a bar plot that displays the proportion of genes and TE-genes in each expression category (downregulated, stable and upregulated). **c.** Scatter plot comparing fold changes of M23 and M25 upregulated genes. Yellow and orange dots represent specific significantly upregulated genes in M23, M25, respectively (LFC > 1 and FDR < 0.01, M23-sp, M25-sp, respectively). Grey dots represent significantly upregulated genes in both mutants (LFC > 1 and FDR < 0.01, M23 ∩ M25). The solid line represents the linear fit and the grey shading represents 95% confidence interval for the significantly upregulated genes in both mutants. **d.** and **e.** The upper panel represent Venn diagrams showing the numbers of specific (M23-sp, M25-sp) and common (M23 ∩ M25) upregulated TE-genes and genes, respectively, in each mutant compared to the Pt18.6 reference (WT). The lower panel shows the top 10 enriched canonical pathways of upregulated TE-genes and genes, respectively, sorted by p.value in both mutants (M23 ∩ M25) as identified by topGO analysis. The dashed lines show the p.value of 0.05. **f.** Venn diagram displaying

1062 downregulated TE-genes (LFC < -1 and FDR < 0.05) in M23 (M23-spe) (yellow) and

1063 M25 (M25-spe) (orange). **g.** as for **f.** for downregulated genes.

1064

Figure 5

Interplay between CG methylation and TE-gene expression

a. Heatmap of CG methylation levels in Pt18.6 reference (WT), M23 and M25 (left

panel) and LFC normalized levels (Z-scores) (right panel) of the 600 upregulated TE-

genes in M23 and M25 compared to Pt18.6 reference (WT). **b.** Percentages of overlap

between common hypoDMRs and upregulated (red), downregulated (blue) and stable

TE-genes (grey) in M23 only (M23-sp), M25 only (M25-sp) and both mutants (M23 ∩

M25). **c.** Distribution of common (M23 ∩ M25) upregulated TE-genes that overlap with

TE-genes regulatory regions. **d.** Mapping of TEs covered in TE-genes that overlap or

not with common hypoDMRs (queries in the left) onto annotated TEs on Phatr3. Bar

sizes are proportional to the number of TEs in the queries that are assigned to each

TE category. **e.** Violin plot comparing the length (bp) of TEs covered in TE-genes that

overlap or not with common hypoDMRs, Wilcoxon test. The white boxplots show

summary statistics, the bottom of the box displays the lower quartile, the upper part,

the upper quartile and the middle line the median. **f.** IGV snapshot of expression levels

in both replicates of WT and DNMT5:KOs (M23 and M25) (green tracks) and CG

methylation levels in the WT and DNMT5:KOs (M23 and M25) (orange tracks) of an

example LTR-copia (highlighted in yellow). The common hypoDMRs and genes are

also shown in the red and black tracks, respectively.

1084

1085

1086

1087

1088

1089

1090

1091

a.

DNMT4s

diaDNMT4
Diatoms (pennate + centric)

RID/DMTA
Fungi

DNMT1s

△ Raphidophycea D4like enzymes
* Haptophytes CMTs
■ Green algae CMTs
✖ Dinoflagellates/red algal DNMT3s

CMTs

Dim2

Fungal DNMT1

DNMT1/MET1

0.92 0.93
0.92
0.84 0.92

Diatoms (pennate)
Bolidomonas
Pelagophytes
Dictyocophytes
Haptophytes
Green algae
Euglenozoa
Fungi

DNMT5a

0.98 0.51
0.97 0.88

DNMT5b

0.94

DNMT6s

DNMT5s

Diatoms
(pennate + centric)
Bolidomonas
Haptophytes

DNMT5c
Dinoflagellates

0.89
0.84

0.94

Diatoms (pennate + centric)
Labyrinthulomycetes
Crysophytes
Dictiochophytes
Rizharians
Dinoflagellates
Haptophytes
Green algae
Euglenozoa

DNMT2s

✖

Dinoflagellates
Red algae

DNMT3s
Diatoms (pennate + centrics)
Land plants
Vertebrates

b.

DNMT5

DNMT5a
Micromonas pusilla
MicpuC3v2_2783

200 aa

SNF-related

DNMT5b
Thalassiosira pseudonana
Thaps3_3158

SNF-related

DNMT5c
CAMPEP_0197657916

DNMT1-CMT like
Gephyrocapsa oceanica
CAMPEP_0188208858

200 aa

DNMT1/MET1
Chlamydomonas euryale
CAMPEP_0201446650

200 aa

DNMT6
Phaeodactylum tricornutum
Phatr3_J47357

200 aa

DNMT4

RID
Neurosposra crassa
Q8NJW0

200 aa

diaDNMT4
Thalassiosira miniscula
CAMPEP_0201057066

Thalassiosira pseudonana
Thaps3_11011

D1-DNMT4 like
Chatonella subsalsa
CAMPEP_0187160654

200 aa

DNMT3
Phaeodactylum tricornutum
Phatr3_J47136

200 aa

■BAH ; ■chromo ; ■DNMT ; ■DX ; ■Hter ; □LBR ; ■PHD ; ■RFD ; ▮Ring