Whole-cell modeling in yeast predicts compartment-specific proteome constraints that drive metabolic strategies

Ibrahim E. Elsemman[1,7,#], Angelica Rodriguez Prado[2,3,#], Pranas Grigaitis[2,#], Manuel Garcia Albornoz[4], Victoria Harman[5], Stephen W. Holman[5], Johan van Heerden[2], Frank J. Bruggeman[2], Mark M.M. Bisschops[3], Nikolaus Sonnenschein[1], Simon Hubbard[4], Rob Beynon[5], Pascale Daran-Lapujade[3], Jens Nielsen[1,6,*], and Bas Teusink[2,*]

[1] Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, DK2800 Lyngby, Denmark

[2] Systems Biology Lab, Amsterdam Institute of Molecular and Life Sciences, Vrije Universiteit Amsterdam, The Netherlands

[3] Department of Industrial Microbiology, Technical University Delft, Delft, The Netherlands

[4] Division of Evolution & Genomic Sciences, University of Manchester, Manchester, UK

[5] Institute of Systems, Molecular and Integrative Biology, University of Liverpool, Liverpool, UK

[6] Department of Biology and Biological Engineering, Chalmers University of Technology, SE41296 Gothenburg, Sweden

[7] Current address: Department of Information Systems, Faculty of Computers and Information, Assiut University, Assiut, Egypt

[#] These authors contributed equally to this work

[*] To whom correspondence should be addressed: nielsenj@chalmers.se (JN), b.teusink@vu.nl (BT)

When conditions change, unicellular organisms rewire their metabolism to sustain cell

maintenance and cellular growth. Such rewiring may be understood as resource re-allocation

under cellular constraints. Eukaryal cells contain metabolically active organelles such as

mitochondria, competing for cytosolic space and resources, and the nature of the relevant cellular

constraints remain to be determined for such cells. Here we developed a comprehensive metabolic

model of the yeast cell, based on its full metabolic reaction network extended with protein

synthesis and degradation reactions (16304 reactions in total). The model predicts metabolic

fluxes and corresponding protein expression by constraining compartment-specific protein pools

and maximising growth rate. Comparing model predictions with quantitative experimental data

revealed that under glucose limitation, a mitochondrial constraint limits growth at the onset of

ethanol formation - known as the Crabtree effect. Under sugar excess, however, a constraint on

total cytosolic volume dictates overflow metabolism. Our comprehensive model thus identifies

condition-dependent and compartment-specific constraints that can explain metabolic strategies

and protein expression profiles from growth rate optimization, providing a framework to

understand metabolic adaptation in eukaryal cells.

Macromolecular synthesis and energy conservation by metabolism underlies cellular

maintenance, growth and fitness. Unicellular organisms such as yeasts generally display a great

variety of metabolic strategies that lead to competitive fitness across conditions[1]. The associated

reprogramming of metabolism between such metabolic strategies is of key interest in

biotechnology and biomedical research.

One well-known example is "overflow" metabolism in which under aerobic conditions

not all substrate is fully oxidized but secreted as by-products. In cancer cells it is referred to as

the Warburg effect: enhanced glycolytic activity with lactate as byproduct at the expense of

respiration[2]. The same phenomenon is known as the Crabtree effect in *Saccharomyces*

cerevisiae (Baker's yeast)[3]. At sugar limitation yeast respires glucose completely to CO_2; at sugar

excess it displays respirofermentative metabolism, where respiration is combined with ethanol

formation (alcoholic fermentation). The extent to which these two metabolic strategies are used

can be titrated in a glucose-limited chemostat: at a specific "critical" dilution (=growth) rate,

ethanol formation starts and increases linearly with growth rate[4]. Other microorganisms show

similar behaviour[5]: for example, *E. coli* produces acetate at higher growth rates at the expense of

respiration[6].

In the last decade a theoretical framework has been developed that can explain why cells

shift metabolic strategies upon environmental or gene-expression perturbations[5,7-10]. In essence

it is based on the catalytic benefits of proteins and their associated costs[11]. These costs comprise

competition for resources such as building blocks, energy and synthesis machineries, and for

space in cellular compartments. Two key features of this resource allocation paradigm can

explain metabolic adaptations. First, cellular compartments can become "full" when they are

fully occupied with (maximally) active proteins, such that an increase in one protein has to come

at the expense of another. This was postulated as a phenomenological rule based on

experimental observations[12], but also follows naturally from growth-rate maximization[13].

Second, cells allocate their limited resources for protein synthesis according to their

demands[14,15]. Consequently, fractions of needed proteins vary with growth rate within

compartments whose protein content is bounded, and this can lead to "active proteome

constraints" related to full compartments.

Within this framework, the onset of overflow metabolism was explained by the smaller

protein cost of generating ATP through fermentation than respiratory pathways[6,7]; this becomes

important at fast growth when biosynthesis and ribosome demands are high and thus require

large proteome fractions. Earlier work suggests that the proteome-constrained resource

allocation paradigm, which was largely developed for *E. coli*, may also be a powerful perspective

for regulation of eukaryal yeast metabolism, such as ribosome biosynthesis[16], and growth on

different sugars[17]. However, a key feature of the metabolism of a eukaryal cell is the presence of

metabolically active organelles, most prominently mitochondria. Each organelle introduces two

new compartments (intra-organellar space and membrane), and how these compartments

impacts adaptation of metabolism, and which compartments become limiting under different

conditions, is an open question.

Moreover, despite the wealth of experimental data on *Saccharomyces cerevisiae*, a

comprehensive, quantitative, data set in which growth rate is systematically varied and both

fluxes and protein expression levels are measured, which are needed to validate resource

allocation predictions, are still rare (see however, some recent studies[16,18]). Here we generated

such data sets and in parallel developed the most detailed and comprehensive,

compartmentalized and quantitative model of metabolism and protein synthesis of yeast. The

model can compute the costs and benefits of protein expression and translocation; It can be used

to interpret or predict experimentally determined changes in growth rate, (minimal) protein

expression and metabolic fluxes as a result of growth rate optimization through resource

allocation into different, compartmentalized, proteome fractions. Comparison of the model

predictions with the data gives unprecedented insight into our physiological understanding of

this important model organism.

Results

Construction of a comprehensive proteome-constrained yeast model

We extended an existing[19] metabolic genome-scale metabolic model of yeast (GEM) by coupling metabolic fluxes to the synthesis of the catalysing enzyme and added constraints on protein concentrations, expressed as protein fractions of the total proteome (Fig. 1a). We refer to the resulting model as proteome-constrained Yeast (pcYeast). Earlier GEM-based approaches exist that incorporate resource allocation, and for yeast these considered constraints on enzyme activities and total protein content[17,20–23], whereas for *E. coli* constraints and reactions associated with transcription and translation were added[9]. Others considered membrane-area constraints and limitations of protein allocation to specific pathways[8,24]. We combined all these extensions (see Supplementary Notes for detailed information) to make pcYeast: a next-generation yeast GEM and computable knowledge base that incorporates protein expression, translation, folding, translocation and degradation at genome-scale for a compartmentalized, eukaryal, organism. In our current model, we consider the protein compartments most relevant for central metabolism: plasma membrane, cytosol, mitochondrion, and mitochondrial membrane. Other cell compartments such as the nucleus or endoplasmic reticulum are not (yet) specified explicitly - but do occupy volume in the cytosol.

The cellular proteome was divided into metabolically active, ribosomal, and unspecified (UP) proteins. The UP fraction is cytosolic, has an average amino acid composition and is added to always maintain a constant protein density in the cytosol. It has a minimum expression level estimated from the experimental proteomics data (Fig. S1, Supplementary Notes). The minimal UP fraction represents growth-rate independent structural, signalling and "household" proteins. Higher expression of UP than minimal represent both unspecified anticipatory proteins, or metabolic proteins that do not carry flux – including the unsaturated fraction of flux-carrying enzymes, as we will explain.

Metabolic enzymes are assigned to a specific compartment, either cytosol, plasma
membrane, mitochondrial matrix or inner mitochondrial membrane; Mitochondrial proteins
require additional protein transport complexes[25]. For each protein we comprehensively
modelled synthesis and degradation processes, which are responsible for the largest fraction of
cellular energy usage. Our model includes 1,523 proteins whose life cycles are described by
16,304 reactions that include translation initiation, elongation and termination factors,
ribosomal assembly factors, protein-specific folding by chaperones and degradation reactions,
as well as 5'UTR-length dependent energetic costs for translation initiation (Table 1,
Supplementary Notes).

We applied three classes of constraints that couple metabolic fluxes and peptides
synthesis rates (Fig. 1b and Supplementary Notes for details). The *enzyme capacity constraint*
sets the minimal enzyme synthesis rate required to achieve a certain metabolic flux. Thus, all
metabolically-active proteins are modeled to work at their maximal rate and are minimally
expressed in the model; the unsaturated fraction of flux-carrying enzymes is represented by UP,
the unspecified protein that is used to maintain protein density. In this way we prevent choices
about unknown regulatory and kinetic mechanisms that may affect the activity of enzymes;
rather we use the deviation between predicted minimal and measured actual protein expression
levels to indicate the level of saturation of each enzyme. The total enzyme synthesis rate is
constrained by the abundance of ribosomes through a *ribosome capacity constraint*, for both
cytosol and mitochondria. Finally, we added *compartment-specific constraints* on the proteome,
for the cytosol, the plasma membrane, and the mitochondrial matrix and inner membrane, (Fig.
1b). The values for these constraints are based on independent literature data or were fitted to
experimental data (as explicified in Supplementary Notes) and the values are either fixed or
growth-rate dependent, depending on the nature of the constraint.

The steady-state metabolite balances, the enzyme synthesis and degradation balances,
and the compartment-specific proteome constraints together specify a linear program with its
fluxes as optimisation variables, provided the growth rate is treated as a parameter. We use a

binary search algorithm to find the maximum growth rate where the linear program is still

feasible, and a marginal increase in the growth rate would result in an infeasible linear

programming problem. The model returns all the flux values associated with the maximal

feasible growth rate. It should be noted that the structure of the pcYeast model is strain-

independent: this allows subsequent calibration of the model to accommodate and account for

differences in cell physiology and metabolism, inherent to any specific strain of *S. cerevisiae*.

Calibrating the model against experimental data

We performed a series of experiments, using a wild-type *S. cerevisiae* strain CEN.PK 113-

7D, for collection of high-quality datasets of fluxes and protein levels, used either as model input

or for comparison with model predictions. We used glucose-limited continuous cultures

operated at dilution rates close to the critical dilution rate for ethanol formation, to capture

proteome change upon the onset of overflow metabolism. Additionally, we varied the growth

rate in pH-controlled batch experiments, either with different sugar quality or through

translation inhibition. We measured fluxes, including O_2 and CO_2 fluxes (Supplementary Dataset

1), which combined with biomass measurements, allowed to estimate the so-called maintenance

parameters, i.e. ATP usage that is not explicitly accounted for in the model (Supplementary

Notes). Label-free proteome quantification allowed us to reliably estimate proteome fractions of

around 3000 of the 6000 proteins (Supplementary Datasets 2, 3, and 4).

Parameters associated with translation strongly affected our model outcomes, and we

used published quantitative proteomics data[16] to estimate parameters for protein translation,

such as the elongation rate (Supplementary Notes). Following experimental reports we assumed

a constant inactive fraction of ribosomes and a fixed saturation of the actively translating

ribosomes[16,27] and were able to describe the growth-rate dependent ribosome mass fraction

with the model (Fig. 1c). As evidence for correctly capturing the costs of protein synthesis, we

correctly predicted the effect of over-expressing mCherry, an unneeded, "gratuitous" protein, on

the specific growth rate (Fig. 1d).

The model predicts shifts in metabolic strategies

We subsequently used the model to analyse yeast's physiological response to different levels of glucose availability. Traditional Flux Balance Analysis computes continuous chemostat cultures by minimizing glucose uptake rate at fixed growth (=dilution) rate[28]. Here we simulated glucose availability by varying the degree of saturation of the glucose transporter. We needed to constrain the maximal expression level of the glucose transport system based on literature data (Supplementary Notes), as leaving expression free to occupy available membrane space led to unrealistically high expression levels and overestimation of growth rate at low glucose levels. At each saturation level we computed the maximal feasible growth rate and compared model predictions with published data[29], and with data from our glucose-limited chemostat cultures (growth rates between 0.2 – 0.34 h^{-1}). We also included our data from batch cultures on glucose (growth rates 0.37-0.39 h^{-1}) and on trehalose; Trehalose is a disaccharide of two glucose molecules, hydrolyzed extracelullarly[30], thus providing slow release of glucose that supports low growth rates.

As in the chemostat the specific growth rate is equal to the dilution rate, the maximal feasible growth rate that the model predicted can be directly compared to the experimental data (Fig. 2a, 2c-d). The (residual) glucose concentrations were calculated from documented (high) affinity of the transporters, which is close to 1 mM[31]. The resulting relationship between growth rate and residual glucose concentration fit experimental data very well (Fig. 2a), validating our expectation that we could ignore glucose efflux from the cells due to minute levels of intracellular glucose[32] (see Supplementary Notes for details). Increasing glucose transporter saturation increased predicted growth rate, and the effect saturated (Fig. 2b), suggesting that at maximal growth rate further increase in glucose availability has little impact. Predicted biomass yield (Fig. 2c) and fluxes (Fig. 2d) corresponded well with the experimental data, as did the intracellular flux ratios from previously published ^{13}C-labeling flux analysis at three specific growth rates in glucose-limited chemostat cultures (Supplementary Figure 2). In particular, the

model predicted a maximal oxygen consumption rate at dilution rates higher than 0.28 h^{-1}, at the onset of ethanol formation. Above 0.35 h^{-1}, this rate rapidly drops to the low level that is observed under glucose excess (batch) conditions. We conclude that the model can adequately predict the changes in metabolic fluxes when the growth rate is varied through the availability of glucose.

Changes in metabolic strategies are the result of proteome constraints

We used pcYeast to identify the active proteome constraints, *i.e.* the protein pools that limit growth rate, because, according to resource allocation theory, the number of active proteome constraints determines the maximal number of independent metabolic behaviors that are possible in optimal states[5,13]. For this we computed the occupancy of each protein pool: a pool that is fully occupied is indicative of an active constraint. At low growth rates, below 0.28 h^{-1}, the glucose transporter was the only proteome pool that is fully occupied (Fig. 2e). With only glucose uptake as active constraint, pure respiration is the single optimal strategy. At the onset of ethanol formation a second metabolic mode started to carry flux (for formal computation of these modes and the concomitant theory, see Supplementary Notes), and thus a second constraint must have become active. Indeed, at this growth rate the occupancy of the inner-mitochondrial membrane became maximal (Fig. 2e). Thus, the model suggests that under glucose-limited chemostat conditions, the onset of ethanol formation is caused by a limit of the mitochondrial membrane space, and hence the amount of proteins that yeast can maximally express in this compartment.

At a growth rate of 0.35 h^{-1} we found that the unspecified protein level reached its minimal value (Fig. 2e), equivalent to the cytosol being completely filled with maximally active proteins. Further growth rate increase requires higher ribosomes and biosynthetic protein fractions, which now has to come at the expense of the least proteome-efficient pathway. The model confirmed earlier calculations[33] that respiration is less proteome efficient than fermentation (Supplementary Figure 3) and respiration is therefore replaced by fermentation.

The model suggested therefore that at growth rate above 0.35 h^{-1} the second growth-limiting

constraint was shifted from the mitochondrial proteome to the cytosolic proteome. Thus, the

metabolic changes in the model, when growth rate and thus metabolic fluxes increase, are

dictated by the filling up of different cellular compartments with active protein, unique for an

eukaryal cell. The level of detail in our model to suggest the condition-dependent, active,

protein-concentration constraints belonging to different compartments has so far not been

provided by any other model.

Proteomics data validates model predictions

We subsequently measured protein levels with quantitative proteomics and compared

them to the minimal protein levels that the model predicted to be needed to support metabolic

flux. Since we compute minimal levels as if all the enzymes worked at their maximal rate, we

expected to underestimate most proteome fractions. Especially at lower growth rates where

nutrient limitation is most severe, one can expect lower average enzyme saturation, and indeed

we observed larger deviations between predicted minimal protein levels and measured protein

fractions at low growth rates (Fig. 3a). The difference between the predicted minimal level and

the data may be interpreted as a proxy for the average saturation of enzymes. In terms of

protein synthesis costs, the difference between the experimentally measured enzyme expression

and the predicted minimal expression level, however, are covered by the expression of the UP.

We see an overall tendency that the saturation of enzymes increases with growth rate

(Supplementary Figure 4). This is most prominent for the glycolytic pathway; also for amino

acid biosynthesis, the protein expression is higher than expected based on metabolic activity,

indicating also here a substantial undersaturation of the enzymes, as observed before for

bacteria such as *E. coli*[34] and *L. lactis*[35]. We find similar patterns for other biosynthetic pathways,

except for lipids (Supplementary Figure 5).

For mitochondrial proteins involved in the citric acid cycle and respiration, however, we

found that predicted minimal proteome fractions were very close to the measured ones (Fig.

3a). Unless k_{cat} values of mitochondrial enzymes are systematically underestimated, this

indicates that mitochondrial proteins work at higher average saturation than cytosolic proteins -

and seemingly close to their maximal capacity. Regardless of absolute numbers, the saturation of

the mitochondria seems rather constant, suggesting that yeast tunes the total amount of

mitochondria, rather than make excess (hence subsaturated) mitochondria, at least under these

conditions. This may make sense, given the extra costs of mitochondrial components such as

membranes, and for protein translocation of host-derived proteins during mitochondrial

biogenesis, which competes for membrane space with respiratory proteins.

Upon closer inspection, we observed that at the onset of ethanol formation the total

mitochondrial protein fraction started to decrease (Fig. 3b). The observed decay follows the

theoretical dilution-by-growth kinetics if at that point the rate of mitochondrial biosynthesis has

reached a maximum (Fig. 3b). Thus, the data suggests that the rate of mitochondrial biogenesis,

rather than the maximal mitochondrial membrane area currently used by the model, may reach

the host's maximal capacity at the onset of ethanol formation. When we zoom in on the

mitochondrial proteome, we find that the mitochondrial ribosome fraction increased as a

funtion of growth rate, and also other proteins re-allocated (Supplementary Fig. 6). Indeed,

mitochondria are self-replicating entities abiding to the same resource allocation principles as

the host, which even includes selection for fast replication - but obviously severely dictated by

the proteins the host provides. More data related to the mitochondrial biosynthetic processes,

such as mitochondrial ribosomal capacity and protein import machinery would be required to

predict the maximal mitochondrial growth rate from first principles, which is outside the scope

of this study. Nonetheless, the distinct changes of mitochondrial proteins at the critical dilution

rate are consistent with the model prediction that a mitochondrial constraint is responsible for

the onset of ethanol formation under glucose-limited conditions.

Constraints and fluxes under sugar excess conditions

We then varied growth rate (between 0.05 h^{-1} and 0.4 h^{-1}) by providing different sugars, i.e. trehalose, galactose, maltose and glucose during batch cultivation. Ethanol production was already observed on galactose, already at a growth rate of 0.16 h^{-1} so at a much lower growth rate than the critical growth rate of 0.28 h^{-1} under glucose-limited growth (Fig. 4a). Maltose showed intermediate growth rate and fluxes. Initial model simulations with a "naïve" model using the reported catalytic rates of the transporters and catabolic enzymes involved in galactose and maltose metabolism, however, resulted in predicted growth rates and fluxes not far from growth on glucose. This suggests that there are additional cost factors that were not included in the model, and or that *Saccharomyces cerevisiae* is not as well adapted to these sugars.

We therefore used the model as data analysis tool to estimate possible changes in parameters that fit the observed growth rate and corresponding fluxes (see Supplementary Notes for details). Such parameter changes may be interpreted as costs for suboptimal metabolism of carbon sources other than glucose. The onset of ethanol formation at a growth rate of 0.16 h^{-1} required a combination of changes in both sugar uptake and the intracellular proteome (through the minimal UP fraction constraint): a lower sugar uptake capacity alone would be identical to lowering saturation of the transporter as was done for glucose (Fig. 2), and pure respiration would have been found at 0.16 h^{-1}. Conversely, only an increase in minimal UP would have resulted in a proportional flux decrease that we also found with mCherry overexpression (or translation inhibition, Supplementary Figure 7), and more ethanol were to be found.

We had to decrease the maximal galactose uptake rate by a factor of 2.5 compared to glucose. Furthermore, an increase in minimal UP fraction was needed, to 0.49 g/g protein. To fit all fluxes optimally, we also required additional energetic costs (see Supplementary Notes), whose mechanistic underpinning remains to be explored but may be related to the reported toxicity of galactose intermediates[36]. Such a change in energetic costs were not needed to

describe the data for growth on maltose: only a change in the maltose uptake rate and minimal

UP fraction (of 0.34 g/g protein) were required to achieve good fit.

For maltose, a disaccharide of glucose, the reason for the required parameter changes is

not clear. Only a maltose proton-symporter and a maltase protein distinguishes it from growth

on glucose. The transport expression may be tightly regulated as very high maltose uptake rates

can result in substrate-accelerated death[37]. For galactose, the toxicity of its intermediates[36]

results in an evolutionary trade-off with growth on glucose[38]; on galactose yeast cells appear to

be still prepared for growth on glucose, which may prevent them from optimal expression of

proteins on galactose, as shown by expression titration experiments[39]. Indeed, laboratory

evolution experiments on galactose select mutations in Ras/cAMP signalling and adapted strains

show increased growth rates and concomitant increased ethanol fluxes[38]. Interestingly, the

direction of change points to the optimal behaviour predicted by the initial naïve model,

suggesting that the pcYeast model may aid in predicting the direction of evolutionary change

during laboratory evolution experiments (Supplementary Figure 8).

With the updated parameters, we identified for both sugars that the active constraints

limiting growth were the sugar transport expression and the minimal UP fraction constraint

(Figure 4d, Supplementary Notes, Supplementary Figure 9). These active constraints explains

ethanol formation during growth on galactose even though the growth rate is lower than the

critical dilution rate on glucose.

Proteomics data on sugar excess shows re-allocation of metabolic strategies

If growth rate is actively constrained by the cytosolic proteome under galactose, maltose

and glucose excess conditions, it implies that all cytosolic proteins work at their maximum

activity, and changes in flux must be brought about by changes in protein level. We therefore

turned to proteomics again. Comparing the minimal levels of the model with experimental data,

we find again that mitochondrial proteins for the TCA cycle and respiration are very similar to

the predicted minimal levels required to sustain flux (Fig. 4c). Cytosolic proteins were

underestimated - even at sugar excess conditions. (Note however that the expected maximal

attainable activity is not likely at the maximal rate in the forward direction as product inhibition

is inevitable in a chain of enzymes.)

More indicative of "a full cytosol" is that at the onset of ethanol formation (at galactose

growth rate and higher) we find evidence for proportional relationships between protein and

flux for high-flux carrying, pathway-grouped, proteins as a function of growth rate (Fig. 4c). This

is observed even down to the individual protein level (also involving changes in expressed

isozymes), as illustrated for glycolytic and respiratory proteins in Fig. 4d. This implies that

under these conditions, enzyme saturation was constant (and maximal, we expect) and changes

in flux could only be brought about by corresponding changes in enzyme levels. This data

illustrates how mitochondrial proteins are being traded in for glycolytic proteins needed for an

enhanced fermentation and growth rate. It also confirms the model's prediction that the

cytosolic proteome constraint is active during growth on these sugars.

Inhibition of translation highlights the role of environmental signalling in coordination

of metabolism in yeast.

Finally we varied growth rate by translation inhibition by cyclohexamide under

controlled glucose batch conditions, and again measured fluxes, growth rate and proteome

profiles (Fig. 5a). Upon inhibition of translation, we found a decrease in growth rate and close to

proportional decreases in glucose, ethanol and CO_2 fluxes, for both the model and the

experimental data (Fig. 5b). Such behaviour is expected when one dominant constraint is active

and its extent is varied (cf. glucose-limited fully respiratory growth, Fig. 2). In the case of glucose

excess, the model suggested that the cytoplasmic volume was fully occupied with active proteins

(minimal UP constraint was hit), and inhibition of translation required higher expression levels

of ribosomes, taking away limited proteome space for growth-supporting activities.

However, experimental observations compromised this initial explanation. First, for

oxygen the model also predicted a proportional increase with growth rate, but experimentally

the fluxes did not change much as did the expression of enzymes involved in oxygen consumption, such as TCA cycle and oxidative phosphorylation (Fig. 5d). Moreover, the ribosomal proteome fraction increased much less with inhibition than the model predicted (Fig. 5c). Since translation inhibition in the model has the same effect as overexpression of a non-functional protein (Supplementary Figure 7), we followed the earlier observation that the inactive fraction of ribosomes could be recruited for translation, depending on the translational load[16], with only a small improvement (Supplementary Figure 10).

This suggested that either some constraint prevents the ribosomal fraction from increasing to the optimal levels predicted by the model, or the expression of ribosomes in yeast is dominantly regulated by environmental nutrient signalling and less by internal cues. A dominant role of signalling in ribosomal biogenesis has been suggested before[16]. In yeast the TOR pathway appears to be the master regulator of ribosomal biosynthesis and assembly at steady-state growth[40,41]. Following the TOR-specific targets described by Kunkel[41], we find that key target proteins of this signalling pathway, including ribosomal auxiliary factors, had constant expression levels (Supplementary Figure 11 and Supplementary Table 1), supporting the dominant role of external rather than internal cues.

When we constrained ribosomal expression to the measured maximal response, ribosomal expression rapidly became the only active constraint in the model, and the proteome space that became available in the cytosol at the lower growth rates was used for increased respiration (Supplementary Figure 12). This is not observed experimentally, and our data suggest that respiration does not respond to internal cues either. In contrast the fluxes and expression of proteins involved in glycolysis and amino acid metabolism did decrease with growth rate (Fig. 5bd). This suggests that these pathways must be sensitive to internal feedback regulation, as is well known for amino acid metabolism[42]. Thus, the proportional fluxes we found for ethanol and glucose upon translation inhibition, are likely the result of control by demand[43], with lower demand at lower growth rate.

Discussion

In this work, we developed the most comprehensive model of a growing, compartmentalized, eukaryal cell to date. It includes all its known metabolic reactions, and details of the protein synthesis, degradation and transport machinery to express the enzymes. The key of our approach is the application of constraints on protein pools in the different compartments that have direct biochemical meaning and could be independently estimated from literature data. Our approach is unique in level of detail and in dealing with cellular compartmentation, in particular of the mitochondria. We furthermore generated a unique set of high-quality quantitative data on both fluxes and the proteome under different, well-controlled, conditions. Through integration and comparison with the model, we provide deeper insight into the physiology of *Saccharomyces cerevisiae*.

First, we firmly established that metabolic growth strategies of yeast on glucose can be well understood from a proteome-constrained optimisation problem with growth rate as objective. Through our high resolution sampling around the critical dilution rate, we observed distinct changes in proteins exactly at the onset of ethanol formation in the glucose-limited chemostat. We also show that the active constraints that drive these changes can be different under different conditions such as batch growth on galactose - even if ethanol is made in both cases. Our approach to identify the active cellular constraints may resolve some of the discussion in current literature about *the* cause of overflow metabolism, not only in yeast but possibly also in other eukaryotes, including discussion about the Warburg effect in mammalian cells[44].

Second, the proteome constraints of the model are currently based on experimental observations, but further research could drill deeper into their origin. For example, why would the protein density in the cytosol be relatively constant? Does this balance diffusion rates with catalytic capacities[45]? Are the current morphological dimensions of a yeast cell optimal for growth rate? Recent work on selection for cell number showed that smaller cells can be readily selected for[46]. We also identified that the levels of glucose transport and that of mitochondria

need to be constrained to describe the data. Why would yeast not express these components at

higher levels? In the case of mitochondria, the proteomics data suggest that rather than a

maximum mitochondrial membrane area and matrix volume, there is a maximal rate of

mitochondrial biogenesis. Can we calculate this rate from first principles? One could imagine

that an upper limit for mitochondrial "growth rate" exists if all but eight metabolic proteins need

to be transported over the same membrane that must also harbour the full machinery for

oxidative phosphorylation. Moreover, we focused on mitochondrial protein content, and ignored

details on morphology, lipid synthesis, or possible assembly costs. Thus, a next version of the

model will need to address the mitochondrial transport, biosynthesis and morphology in much

more detail.

In the case of glucose transport, the model suggested that further increase in glucose

transporters beyond wild type expression did not increase growth rate substantially and would

likely be invisible for evolution. At maximal saturation of the transporter, glucose transport

expression was (just) no longer an active constraint in our model (Fig. 2e). Thus, it appears as if

yeast expresses just enough glucose transporters to maximise its growth rate under glucose

excess – as found in bacteria[47]. Expressing higher transport levels at lower glucose levels would

enhance growth rate but may not pay off if this state is a transient towards glucose starvation, or

could be outright dangerous if suddenly glucose would become available[37]. The expression level

of the hexose transporters may thus have evolved to be an adaptation to dynamic

environments[48]. Long-term evolution experiments in glucose-limited chemostats indeed show

gene duplications of high-affinity glucose transporters[49], showing that growth limitation, and

hence selection pressure, is on glucose transport under these conditions.

Third, in the case of nutrient uptake limitation, there appears to be "excess" proteome

space that could be filled with anticipatory proteins or heterologous enzymes at no cost in

fitness. Even though the composition of such excess proteome space cannot be predicted with

our model, we were able to predict metabolic fluxes very well: in this nutrient-limited regime

metabolic efficiency (ATP per glucose), not proteome efficiency (ATP per protein), determines

the best growth rate strategy. This explains why Flux Balance Analysis applied to only the

metabolic network has been so successful, but only under nutrient-limited conditions.

Finally, we found linear or even proportional relationships between growth rate and

flux, and between flux and enzyme levels in a sugar excess (batch culture) regime. In terms of

regulation analysis[50], such a regime is characterised by hierarchical regulation with absence of

metabolic regulation, that is, all changes in flux are brought about by changes in enzyme levels,

not their degree of saturation. For glycolysis and amino acid metabolism, the average saturation,

estimated as the ratio of the predicted minimal enzyme level to the expressed enzyme level, at

maximal growth rate is around 0.5, incidently the level predicted as theoretical optimum for

specific reaction rate[51]. In contrast, when growth is limited by glucose availability, the degree of

saturation varies and the model suggests a mixture of hierarchical and metabolic regulation, as

previously observed in chemostats as well[52].

To conclude, we present a mechanistic, compartmentalized, model of an eukaryal

organism in full details, which can act as a valuable, computable, knowledge base. We show how

it can be used to compute protein costs and identify active growth-limiting constraints, and how

it can be combined with quantitative flux and proteomics data to provide unprecedented insight

into cellular physiology. Finally, we show that also in eukaryal cells, metabolic strategies can be

understood on the basis of growth rate optimisation under nutrient and proteome constraints.

What remains to be understood is how the cell's signalling and regulatory networks manage to

implement these (optimal) proteome allocation strategies.

Methods

461   **Model development**

462   The full description of the pcYeast model is provided as Supplementary Notes. The model codes

463   are available per request to the authors and will be published on GitHub upon acceptance of this

464   manuscript.

Strains and shake flask cultivation

The strain used for this study was *Saccharomyces cerevisiae* strain CEN.PK 113-7D[53]. The stocks used for the experiments were grown in 500 mL shake flask containing 100 mL of YPD medium (10 g L^{-1} of Bacto yeast extract, 20 g L^{-1} of peptone and 20 g L^{-1} of D-glucose). The culture was grown up to early stationary phase and 1 mL aliquots were stored in 20% (v/v) of glycerol at -80 °C. For chemostats, pre-cultures were grown in 500 mL shake flasks containing 100 mL of synthetic medium, the pH was set to 6.0 with 2M KOH and the medium was supplemented with 20 g L^{-1} of D-glucose[54]. Shake flasks with medium were inoculated with the 1 mL frozen stocks of the strain and the cultivations were performed in an orbital shaker at 200 rpm at 30 °C. Pre-cultures for batches with translation inhibitors were performed using a similar approach, whereas for batches with different carbon sources the pre-cultures were made with the respective carbon sources instead of D-glucose.

Chemostat cultivations

Chemostat cultivations were performed in 2 L bioreactors (Applikon, Schiedam, The Netherlands) with a working volume of 1.0 L, the dilution rates used in this study were 0.2, 0.23, 0.27, 0.3, 0.32 and 0.34 h^{-1} in two independent replicate cultures. Growth rates were controlled by modifying the inflow rate on each experiment. Synthetic medium according to Verduyn[54] supplemented with 7.5 g L^{-1} of glucose and 0.25 g L^{-1} Pluronic 6100 PE antifoaming agent was supplied to the bioreactor from a 20 L continuously mixed reservoir vessel. Cultures were sparged with dried air at a flow rate of 700 mL min^{-1} and stirred at 800 rpm. The pH of the cultures was maintained at 5.0 by automatic addition of 2 M KOH. If, after at least six volume changes, the cultures dry cell weight concentration and carbon dioxide production rate differed less than 2% over two consecutive volume changes the cultures were considered to be in steady state. For cultures with dilution rates of 0.27, 0.3, 0.32 and 0.34 h^{-1}, cultures were first

maintained at a dilution rate of 0.2 h^{-1} for 15 hours (3 volume changes) prior to increasing the

specific dilution rate to said values.

Batch cultivations with different carbon sources

Batch cultivations (two independent replicate cultures) were performed using synthetic

medium[54], the medium was supplemented with 20 g L^{-1} final concentrations of the carbon

sources, either D-trehalose, D-galactose, D-maltose or D-glucose (Sigma Aldrich). The

bioreactors were inoculated with 100 mL of yeast shake flask cultures, exponentially growing on

the specific carbon source. The final OD_{660} of all pre-cultures was 4. Cultivations were performed

at 30 °C, the pH was kept at 5.0 by automatic addition of 2M KOH. The working volume of the

bioreactors was 1.4 L in 2 L bioreactors (Applikon, Schiedam, The Netherlands). The cultures

were stirred at 8000 rpm and sparged with a flow rate of 700 mL min^{-1} of dried air. Oxygen

levels were kept above 40% of the initial saturation level as measured with Clark electrode

(Mettler Toledo, Greifensee, Switzerland).

Batch cultivations with the translation inhibitor cycloheximide

Batch cultivations (two independent replicate cultures) with the translation inhibitor

cycloheximide were performed as for the batches with different carbon sources, except that all

the batch cultures ran on 20 g L^{-1} of D-glucose and were supplemented with different

concentrations of cycloheximide with the aim of reaching specific growth rates. In total five

growth rates were studied, being 0.06, 0.12, 0.2, 0.32 and 0.41 h^{-1} (adding respective

cycloheximide concentrations of 228.96, 124.51, 52.15, 25.99 and 0 µg L^{-1}).

Analytical methods

Cultures dry weight was measured by filtering 20 mL of culture, the sample was filtered in pre-

dried and pre-weight membrane filters with a pore size of 0.45 µm (Gelman Science), the filter

517 was washed with demineralized water, subsequently it was dried in a microwave (20 min, 350

518 W) and the final weight was measured as described previously.

519 For the measurement of organic acids and residual carbon source concentrations, supernatants

520 of the cultures were used. For carbon-limited chemostat cultures, the samples were directly

521 quenched with cold steel beads and filtered[55], whereas samples from batch cultures were

522 centrifuged (5 min at $16.000\times g$). The supernatants were analysed by high-performance

523 chromatography analysis on an Agilent 1100 HPLC (Agilent Technologies) equipped with an

524 Aminex HPX-87H ion-exchange column (BioRad, Veenendaal, The Netherlands), operated with 5

525 mM H_2SO_4 as the mobile phase at a flow rate of 0.6 mL min^{-1} and at 60 °C. Detection was

526 according to a dual-wavelength absorbance detector (Agilent G1314A) and a refractive-index

527 detector (Agilent G1362A), as described previously.

528 The exhaust gas from batch cultures was cooled down with a condenser (2°C) and dried with a

529 PermaPure Dryer (model MD 110-8P-4; Inacom Instruments, Veenendaal, the Netherlands)

530 before online analysis of carbon dioxide and oxygen with a Rosemount NGA 2000 Analyser

531 (Baar, Switzerland).

533 **Glycogen and trehalose assays**

534 1 mL of culture was taken from the chemostats and directly added to 5 mL of cold methanol (-40

535 °C). The sample was mixed and centrifuged ($4400\times g$, -20 °C for 5 minutes), the supernatant

536 was discarded, and the pellet was washed in 5 mL of cold methanol (-40 °C), and pellets were

537 stored at -80 °C until further processing. Subsequently, the pellets were resuspended in 0.25 M

538 Na_2CO_3 and processed as described previously[56,57]. D-glucose released from trehalose and

539 glycogen were measured with a D-glucose assay kit (K-GLUC Megazyme), two biological

540 replicates and three technical replicates were analysed per condition.

542 **RNA determination**

For RNA determination, 1-2 mL of broth was transferred to a filter (pore size of 0.45 µm, Gelman

Science), after which the filter was washed with cold TCA 5 %. The cells were resuspended in 3

mL of TCA 5% and centrifuged for 15 minutes at 4 °C at 4000 rpm. The supernatant was

removed and the pellet was stored at -20 °C. Finally, samples were processed as described by

Popolo et al., 1982. Two biological replicates and three technical replicates were analysed per

condition.

Protein determination

For the batches with CHX, culture volumes corresponding to 50 mg of DCW were centrifuged,

washed twice with cold demineralized sterile waterand divided into two aliquots of 5 mL. 2 mL

of the aliquot (containing 10 mg DW) was mixed with 1 mL of 3 M NaOH and incubated at 100 °C

for 10 minutes. The final mix was diluted and processed following the copper-sulfate based

method as described previously[58]. The absorbance of the supernatant was measured at 510 nm,

for calibration lyophilized bovine serum albumin (A2153, Sigma Aldrich) was used. Two

biological replicates and 3 technical replicates were analysed per condition.

Proteomics sample preprocessing

Aliquots of 20 mL of culture from chemostats and batches with different carbon sources were

centrifuged (4000 rpm 4 °C, 10 minutes) and washed two times, the final pellet was flash frozen

in liquid nitrogen and stored at -80 °C. Two biological replicates and two technical replicates

were analysed per condition.

Frozen cell pellets were thawed on ice before transfer to Precellys® Lysing Kit 2 ml screw cap

vials with 0.5mm glass beads (Bertin Instruments, France). Lysis was performed in 250 µl lysis

buffer, 50 mM ammonium bicarbonate with cOmplete protease inhibitor cocktail (ROCHE,

Switzerland), using a Minilys Personal Tissue Homogenizer (Bertin Instruments, France), at

maximum speed for 15 cycles of 30 seconds with a one-minute rest on ice between each cycle.

Lysed material was centrifuged for 10 minutes 13,000× *g* at 4°C, the supernatant fraction was

removed and retained. Fresh lysis buffer (250 μl) was added to the insoluble material, which

was resuspended before extraction from the vial via a small hole inserted into the vial base.

Soluble and insoluble fractions were recombined and the total final volume recorded. Protein

concentration was determined using Pierce™ Coomassie Plus Bradford Assay Kit (ThermoFisher

Scientific, UK).

Protein (100 μg) from each sample was treated with 0.05 % (w/v) RapiGest™ SF surfactant

(Waters, UK) at 80 °C for 10 minutes, reduced with 4 mM dithiothreitol (Melford Laboratories

Ltd., UK) at 60 °C for 10 minutes and subsequently alkylated with 14 mM iodoacetamide

(SIGMA, UK) at room temperature for 30 minutes. Proteins were digested with 2 μg Trypsin

Gold, Mass Spectrometry Grade (Promega, US) at 37 °C for 4 hours before a top-up of a further 2

μg trypsin and incubation at 37 °C overnight. Digests were acidified by addition of trifluoroacetic

acid (Greyhound Chromatography and Allied Chemicals, UK) to a final concentration of 0.5 %

(v/v) and incubated at 37 °C for 45 minutes before centrifugation at 13,000× *g* (4°C) to remove

insoluble non-peptidic material.

Proteomics analytics

The sample running order was randomised using a random number generator (Random.org).

Samples were analysed using an UltiMate™ 3000 RSLCnano system (ThermoFisher Scientific)

coupled to a Q Exactive™ HF Hybrid Quadrupole-Orbitrap™ Mass Spectrometer. Protein digests

(1 ug of each) were loaded onto a trapping column (Acclaim PepMap 100 C18, 75 μm x 2 cm, 3

μm packing material, 100 Å) using 0.1 % (v/v) trifluoroacetic acid, 2 % (v/v) acetonitrile in

water at a flow rate of 12 μL min-1 for 7 min.

The peptides were eluted onto the analytical column (EASY-Spray PepMap RSLC C18, 75 μm x

50 cm, 2 μm packing material, 100 Å) at 40°C using a linear gradient of 120 minute shallow

gradient rising from 8 % (v/v) acetonitrile/0.1 % (v/v) formic acid (Fisher Scientific, UK) to 30

% (v/v) acetonitrile/0.1 % (v/v) formic acid at a flow rate of 300 nL min^{-1}. The column was then

596 washed at 1 % A : 99 % B for 8 min, and re-equilibrated to starting conditions. The nano-liquid

597 chromatograph was operated under the control of Dionex Chromatography MS Link 2.14.

598 The nano-electrospray ionisation source was operated in positive polarity under the control of

599 QExactive HF Tune (version 2.5.0.2042), with a spray voltage of 2.1 kV and a capillary

600 temperature of 250°C. The mass spectrometer was operated in data-dependent acquisition

601 mode. Full MS survey scans between m/z 300-2000 were acquired at a mass resolution of

602 60,000 (full width at half maximum at m/z 200). For MS, the automatic gain control target was

603 set to 3e^6, and the maximum injection time was 100 ms. The 16 most intense precursor ions

604 with charge states of 2-5 were selected for MS/MS with an isolation window of 2 m/z units.

605 Product ion spectra were recorded between m/z 200-2000 at a mass resolution of 30,000 (full

606 width at half maximum at m/z 200). For MS/MS, the automatic gain control target was set to

607 1e^5, and the maximum injection time was 45 ms. Higher-energy collisional dissociation was

608 performed to fragment the selected precursor ions using a normalised collision energy of 30 %.

609 Dynamic exclusion was set to 30 s.

611 **Proteomics data analysis**

612 The resulting raw data files generated by XCalibur (version 3.1) were processed using MaxQuant

613 software (version 1.6.0.16)[59]. The search parameters were set as follows: label free experiment

614 with default settings; cleaving enzyme trypsin with 2 missed cleavages; Orbitrap instrument

615 with default parameters; variable modifications: oxidation (M) and Acetyl (protein N-term); first

616 search as default; in global parameters, the software was directed to the FASTA file; for

617 advanced identification "Match between runs" was checked; for protein quantification we only

618 used unique, unmodified peptides. All other MaxQuant settings were kept as default. The false

619 discovery rate (FDR) for accepted peptide spectrum matches and protein matches was set to

620 1%. The CEN.PK113-7D Yeast FASTA file was downloaded from the *Saccharomyces* Genome

621 Database (SGD) (https://downloads.yeastgenome.org/sequence/strains/CEN.PK/CEN.PK113-

622 7D/CEN.PK113-7D_Delft_2012_AEHG00000000/).

624   The resulting MaxQuant output was then analysed using the MSstats package (version 3.5.6)[60]

625   in the R environment (version 3.3.3) to obtain differential expression fold changes with

626   associated p values, along with normalized LFQ and intensity values as described previously[61].

Acknowledgements

629   This work was supported by NWO (NWO ERA-IB-2, project No 053.80.722 to BT and PDL), ERA-

630   IB 4207-00002B DSForsk to JN, and Biotechnology and Biological Sciences Research Council

631   (BB/M025748/1 to S. Hubbard, BB/M025756/1 to RB). PG and BT also acknowledge support by

632   Marie Skłodowska-Curie Actions ITN "SynCrop" (grant agreement No 764591). We thank

633   SURFsara for the HPC resources through access to the Lisa Compute Cluster.

Author Contributions

Conceptualization, funding acquisition and supervision: BT, JN, PDL, RB, S. Hubbard;

experimental data collection: ARP, VH, S. Holman; experimental data analysis: ARP, PG, MGA,

MMMB; computational modeling: IEE, PG; formal analysis: IEE, PG, JvH, FJB, NS, JN, PDL, BT;

writing – original draft: BT; writing – editing: IEE, PG, FJB, JN, BT. All authors have read and

approved the manuscript.

References

1.   Blank, L., Lehmbeck, F. & Sauer, U. Metabolic-flux and network analysis in fourteen

hemiascomycetous yeasts. *FEMS Yeast Res.* **5**, 545–558 (2005).

2.   Pfeiffer, T. & Morley, A. An evolutionary perspective on the Crabtree effect. *Front. Mol.*

Biosci. **1**, 1–6 (2014).

3.   De Deken, R. H. The Crabtree effect: a regulatory system in yeast. *J. Gen. Microbiol.* **44**, 149–

156 (1966).

4. Van Hoek, P., Van Dijken, J. P. & Pronk, J. T. Effect of specific growth rate on fermentative

capacity of baker's yeast. *Appl. Environ. Microbiol.* **64**, 4226–4233 (1998).

5. de Groot, D. H., van Boxtel, C., Planqué, R., Bruggeman, F. J. & Teusink, B. The number of

active metabolic pathways is bounded by the number of cellular constraints at maximal

metabolic rates. *PLOS Comput. Biol.* **15**, e1006858 (2019).

6. Basan, M. *et al.* Overflow metabolism in Escherichia coli results from efficient proteome

allocation. *Nature* **528**, 99–104 (2015).

7. Molenaar, D., van Berlo, R., de Ridder, D. & Teusink, B. Shifts in growth strategies reflect

tradeoffs in cellular economics. *Mol. Syst. Biol.* **5**, 323 (2009).

8. Mori, M., Hwa, T., Martin, O. C., De Martino, A. & Marinari, E. Constrained Allocation Flux

Balance Analysis. *PLoS Comput. Biol.* **12**, 1–24 (2016).

9. O'Brien, E. J., Lerman, J. A., Chang, R. L., Hyduke, D. R. & Palsson, B. Genome-scale models of

metabolism and gene expression extend and refine growth phenotype prediction. *Mol. Syst.*

Biol. **9**, (2013).

10. Scott, M., Gunderson, C. W., Mateescu, E. M., Zhang, Z. & Hwa, T. Interdependence of Cell

Growth and Gene Expression: Origins and Consequences. *Science* **330**, 1099–1102 (2010).

11. Wortel, M. T., Bosdriesz, E., Teusink, B. & Bruggeman, F. J. Evolutionary pressures on

microbial metabolic strategies in the chemostat. *Sci. Rep.* **6**, 29503 (2016).

12. Scott, M. & Hwa, T. Bacterial growth laws and their applications. *Curr. Opin. Biotechnol.* **22**,

559–565 (2011).

13. Wortel, M. T., Peters, H., Hulshof, J., Teusink, B. & Bruggeman, F. J. Metabolic states with

maximal specific rate carry flux through an elementary flux mode. *FEBS J.* **281**, 1547–1555

(2014).

14. Berkhout, J. *et al.* How biochemical constraints of cellular growth shape evolutionary

adaptations in metabolism. *Genetics* **194**, 505–512 (2013).

15. Bosdriesz, E., Molenaar, D., Teusink, B. & Bruggeman, F. J. How fast-growing bacteria

robustly tune their ribosome concentration to approximate growth-rate maximization. *FEBS*

J. **282**, 2029–2044 (2015).

16. Metzl-Raz, E. *et al.* Principles of cellular resource allocation revealed by condition-dependent

proteome profiling. *eLife* **6**, e28034 (2017).

17. Sánchez, B. J. *et al.* Improving the phenotype predictions of a yeast genome-scale metabolic

model by incorporating enzymatic constraints. *Mol. Syst. Biol.* **13**, 935 (2017).

18. Björkeroth, J. *et al.* Proteome reallocation from amino acid biosynthesis to ribosomes

enables yeast to grow faster in rich media. *Proc. Natl. Acad. Sci.* **117**, 21804–21812 (2020).

19. Sánchez, B. J., Feiran Li, Hongzhong Lu, Kerkhoven, E. & Nielsen, J. *SysBioChalmers/yeast-*

GEM: yeast 7.6.0. (Zenodo, 2016). doi:10.5281/ZENODO.1495468.

20. Beg, Q. K. *et al.* Intracellular crowding defines the mode and sequence of substrate uptake by

Escherichia coli and constrains its metabolic activity. *Proc. Natl. Acad. Sci.* **104**, 12663–

12668 (2007).

21. Adadi, R., Volkmer, B., Milo, R., Heinemann, M. & Shlomi, T. Prediction of microbial growth

rate versus biomass yield by a metabolic network with kinetic parameters. *PLoS Comput.*

Biol. **8**, (2012).

22. Labhsetwar, P. *et al.* Population FBA predicts metabolic phenotypes in yeast. *PLOS Comput.*

Biol. **13**, e1005728 (2017).

23. Nilsson, A. & Nielsen, J. Metabolic Trade-offs in Yeast are Caused by F1F0-ATP synthase. *Sci.*

Rep. **6**, 22264 (2016).

24. Zhuang, K., Vemuri, G. N. & Mahadevan, R. Economics of membrane occupancy and respiro-

fermentation. *Mol. Syst. Biol.* **7**, 1–9 (2011).

25. Okamoto, K. The protein import motor of mitochondria: a targeted molecular ratchet driving

unfolding and translocation. *EMBO J.* **21**, 3659–3671 (2002).

26. Kafri, M., Metzl-Raz, E., Jona, G. & Barkai, N. The Cost of Protein Production. *Cell Rep.* **14**, 22–

31 (2016).

27. Bruggeman, F. J., Planqué, R., Molenaar, D. & Teusink, B. Searching for principles of microbial

physiology. *FEMS Microbiol. Rev.* **44**, 821–844 (2020).

28. Famili, I., Forster, J., Nielsen, J. & Palsson, B. O. Saccharomyces cerevisiae phenotypes can be

predicted by using constraint-based analysis of a genome-scale reconstructed metabolic

network. *Proc. Natl. Acad. Sci.* **100**, 13134–13139 (2003).

29. Canelas, A. B., Ras, C., ten Pierick, A., van Gulik, W. M. & Heijnen, J. J. An in vivo data-driven

framework for classification and quantification of enzyme kinetics and determination of

apparent thermodynamic data. *Metab. Eng.* **13**, 294–306 (2011).

30. Jules, M., Guillou, V., François, J. & Parrou, J.-L. Two distinct pathways for trehalose

assimilation in the yeast Saccharomyces cerevisiae. *Appl. Environ. Microbiol.* **70**, 2771–2778

(2004).

31. Diderich, J. A. *et al.* Glucose uptake kinetics and transcription of HXT genes in chemostat

cultures of Saccharomyces cerevisiae. *J. Biol. Chem.* **274**, 15350–15359 (1999).

32. Teusink, B., Diderich, J. A., Westerhoff, H. V., van Dam, K. & Walsh, M. C. Intracellular Glucose

Concentration in Derepressed Yeast Cells Consuming Glucose Is High Enough To Reduce the

Glucose Transport Rate by 50%. *J. Bacteriol.* **180**, 556–562 (1998).

33. Chen, Y. & Nielsen, J. Energy metabolism controls phenotypes by protein efficiency and

allocation. *Proc. Natl. Acad. Sci.* **116**, 17592–17597 (2019).

34. O'Brien, E. J., Utrilla, J. & Palsson, B. O. Quantification and Classification of E. coli Proteome

Utilization and Unused Protein Costs across Environments. *PLOS Comput. Biol.* **12**,

e1004998 (2016).

35. Goel, A. *et al.* Protein costs do not explain evolution of metabolic strategies and regulation of

ribosomal content: does protein investment explain an anaerobic bacterial Crabtree effect?:

Protein costs and evolution of metabolic strategies. *Mol. Microbiol.* **97**, 77–92 (2015).

36. de Jongh, W. A. *et al.* The roles of galactitol, galactose-1-phosphate, and phosphoglucomutase

in galactose-induced toxicity in *Saccharomyces cerevisiae*. *Biotechnol. Bioeng.* **101**, 317–326

(2008).

37. Postma, E., Verduyn, C., Kuiper, A., Scheffers, W. A. & Van Dijken, J. P. Substrate-accelerated

death ofSaccharomyces cerevisiae CBS 8066 under maltose stress. *Yeast* **6**, 149–158 (1990).

38. Hong, K.-K., Vongsangnak, W., Vemuri, G. N. & Nielsen, J. Unravelling evolutionary strategies

of yeast for improving galactose utilization through integrated systems level analysis. *Proc.*

Natl. Acad. Sci. U. S. A. **108**, 12179–12184 (2011).

39. Keren, L. *et al.* Massively Parallel Interrogation of the Effects of Gene Expression Levels on

Fitness. *Cell* **166**, 1282-1294.e18 (2016).

40. Chaker-Margot, M. Assembly of the small ribosomal subunit in yeast: mechanism and

regulation. *RNA* **24**, 881–891 (2018).

41. Kunkel, J., Luo, X. & Capaldi, A. P. Integrated TORC1 and PKA signalling control the temporal

activation of glucose-induced gene expression in yeast. *Nat. Commun.* **10**, 3558 (2019).

42. Chubukov, V., Gerosa, L., Kochanowski, K. & Sauer, U. Coordination of microbial metabolism.

Nat. Rev. Microbiol. **12**, 327–340 (2014).

43. Hofmeyr, J. S. & Cornish-Bowden, A. Regulating the cellular economy of supply and demand.

FEBS Lett. **476**, 47–51 (2000).

44. Vander Heiden, M. G. & DeBerardinis, R. J. Understanding the Intersections between

Metabolism and Cancer Biology. *Cell* **168**, 657–669 (2017).

45. Dill, K. A., Ghosh, K. & Schmit, J. D. Physical limits of cells and proteomes. *Proc. Natl. Acad. Sci.*

108, 17876–17882 (2011).

46. van Tatenhove-Pel, R. J. *et al.* Serial propagation in water-in-oil emulsions selects for

Saccharomyces cerevisiae strains with a reduced cell size or an increased biomass yield on

glucose. *Metab. Eng.* **64**, 1–14 (2021).

47. Jensen, P. R., Michelsen, O. & Westerhoff, H. V. Control analysis of the dependence of

Escherichia coli physiology on the H(+)-ATPase. *Proc. Natl. Acad. Sci.* **90**, 8068–8072

(1993).

48. Buziol, S. *et al.* Dynamic response of the expression of hxt1, hxt5 and hxt7 transport proteins

in Saccharomyces cerevisiae to perturbations in the extracellular glucose concentration. *J.*

Biotechnol. **134**, 203–210 (2008).

49. Gresham, D. *et al.* The Repertoire and Dynamics of Evolutionary Adaptations to Controlled

Nutrient-Limited Environments in Yeast. *PLoS Genet.* **4**, e1000303 (2008).

50. ter Kuile, B. H. & Westerhoff, H. V. Transcriptome meets metabolome: hierarchical and

metabolic regulation of the glycolytic pathway. *FEBS Lett.* **500**, 169–171 (2001).

51. Cornish-Bowden, A. The effect of natural selection on enzymic catalysis. *J. Mol. Biol.* **101**, 1–9

(1976).

52. Daran-Lapujade, P. *et al.* The fluxes through glycolytic enzymes in Saccharomyces cerevisiae

are predominantly regulated at posttranscriptional levels. *Proc. Natl. Acad. Sci.* **104**, 15753–

15758 (2007).

53. Nijkamp, J. F. *et al.* De novo sequencing, assembly and analysis of the genome of the

laboratory strain Saccharomyces cerevisiae CEN.PK113-7D, a model for modern industrial

biotechnology. *Microb. Cell Factories* **11**, 36 (2012).

54. Verduyn, C., Postma, E., Scheffers, W. A. & Van Dijken, J. P. Effect of benzoic acid on metabolic

fluxes in yeasts: A continuous-culture study on the regulation of respiration and alcoholic

fermentation. *Yeast* **8**, 501–517 (1992).

55. Mashego, M. R., van Gulik, W. M., Vinke, J. L. & Heijnen, J. J. Critical evaluation of sampling

techniques for residual glucose determination in carbon-limited chemostat culture

ofSaccharomyces cerevisiae. *Biotechnol. Bioeng.* **83**, 395–399 (2003).

56. Parrou, J. L., Teste, M.-A. & François, J. Effects of various types of stress on the metabolism of

reserve carbohydrates in Saccharomyces cerevisiae: genetic evidence for a stress-induced

recycling of glycogen and trehalose. *Microbiology* **143**, 1891–1900 (1997).

57. Boer, V., Daran, J., Almering, M., Dewinde, J. & Pronk, J. Contribution of the transcriptional

regulator Leu3p to physiology and gene expression in nitrogen- and carbon-limited

chemostat cultures. *FEMS Yeast Res.* **5**, 885–897 (2005).

58. Verduyn, C., Postma, E., Scheffers, W. A. & van Dijken, J. P. Physiology of Saccharomyces

Cerevisiae in Anaerobic Glucose-Limited Chemostat Culturesx. *J. Gen. Microbiol.* **136**, 395–

403 (1990).

59. Tyanova, S., Temu, T. & Cox, J. The MaxQuant computational platform for mass

spectrometry-based shotgun proteomics. *Nat. Protoc.* **11**, 2301–2319 (2016).

60. Choi, M. *et al.* MSstats: an R package for statistical analysis of quantitative mass

spectrometry-based proteomic experiments. *Bioinformatics* **30**, 2524–2526 (2014).

61. Garcia-Albornoz, M. *et al.* A proteome-integrated, carbon source dependent genetic

regulatory network in *Saccharomyces cerevisiae*. *Mol. Omics* **16**, 59–72 (2020).

Fig. 1 pcYeast model formulation and calibration of protein synthesis parameters. **a.** A schematic

overview of reactions in the model, their interdependence and constraints. Metabolic reactions v_i are

proportional to enzyme concentrations e_i that are synthesized at rate $v_{syn,i}$ by the ribosomes R. Each

protein can be degraded with rate $v_{deg,i} = k_{deg} \cdot e_i$ or diluted by growth rate $v_{dil,i} = \mu \cdot e_i$.

Compartment-specific constraints are indicated in the light-blue boxes. **b.** Optimisation problem with

the key constraints, including 1) steady-state mass balances; 2) production of biomass components

such as DNA, lipids, cell wall and polysaccharides. Proteins and tRNA are excluded as their synthesis

rates are optimisation variables 3) enzyme capacity constraints that couple metabolic flux to catalytic

rate $k_{cat,i}$ and the enzyme level, whose value at steady state is determined by its synthesis rate, rates

of enzyme degradation, and dilution by growth. Note we use equalities and hence enzymes work at

their maximal rate and minimal required protein levels are computed; 4) ribosome capacity that

defines an upper bound for protein synthesis rate; 5) compartment-specific proteome constraints that

define the maximal concentration of proteins that can be contained in that compartment, with w_i the

specific volume or area of protein i; 6) a cytosolic protein density constraint that has the same

function as that of proteome constraints, but whose equality forces the cell to fill up any vacant

proteome space with unspecified protein UP with average amino acid composition. **c.** Growth rate was

varied through sugar type (trehalose, galactose, maltose, glucose) or glucose concentration, and

ribosomal protein fraction was determined by proteomics (which was consistent with literature data,

also plotted). The translation rate was calibrated on that data, as detailed in Supplementary Notes. **d.**

Impact of mCherry protein overexpression on growth rate. Symbols show experimental data[26], solid

lines show model predictions based on glucose minimal (SD) medium or rich SC/YPD media. Model

predictions were obtained by varying the proteome mass fraction, occupied by mCherry, and

determining the maximal predicted growth rate at each value of the mass fraction. The relative

growth fitness represents the ratio between the growth rate at certain mCherry expression level vs.

the unperturbed state (no mCherry expression).

Fig. 2 Predicted and measured physiological response of *S. cerevisiae* CEN.PK as a function of glucose availability a. Measured (symbols) and predicted (line) residual glucose concentrations as a function of growth rate. The latter was calculated based on published affinity for glucose and assuming negligible intracellular glucose under these conditions. Note that this resembles a Monod growth curve but with the dependent and independent axis swapped, as we control growth rate in a chemostat. **b.** Maximal feasible growth rates of the model as a function of the glucose transporter saturation. **c.** Measured (symbols) and predicted biomass yield on glucose. **d.** Experimental fluxes from glucose-limited chemostats at different dilution rates (circles) and from two batch experiments (triangles): excess trehalose (which mimicks glucose limitation at low dilution rate[30]) and excess glucose at the highest growth rate. The lines are model predictions; background colors indicate regimes with different active constraints; **e.** Computed proteome occupancy of different constrained protein pools. A fraction of 1 means that the compartment is full with metabolically actively proteins and constrains the growth rate at that condition.

Fig. 3 Proteomics data of selected pathways as a function of glucose availability. Blue symbols are

glucose-limited chemostat data; orange symbols are controlled batch experiments with excess

trehalose (lowest growth rate) or glucose (highest growth rate) **a.** Comparison of predicted minimal

proteome fractions to sustain growth with the experimentally determined proteome fraction for four

pathways. The ratio between the two represents an estimate of the saturation level of the constituent

enzymes. Lines represent the model; experimental data are symbols. **b.** Decay of steady-state

mitochondrial protein fraction with growth rate at onset of ethanol formation suggests a maximal rate

of mitochondrial biosynthesis $v_{syn,max}$.

Fig. 4 Model predictions, fluxes and protein levels plotted as a function of growth rate during hexose

sugar excess conditions (in the order: trehalose, galactose, maltose, glucose) a. Fluxes of sugar

consumption, oxygen consumption and ethanol production. Circles are experimental data, bar plots

indicate model predictions (of both the growth rate and fluxes); **b**. Predicted active constraints under

the different sugar excess conditions as predicted by the mode (see legend of Fig. 2 for details). **c**.

Comparison of predicted minimally needed proteome fractions with experimentally determined ones

suggests differences in saturation level between pathways. Lines represent the model, experimental

data are circles; **d**. Linearity of the expression of individual enzymes in glycolysis (right) and respiration

(left) with growth rate suggests trading in of respiratory protein for fermentative protein. Asterixes

indicate aggregated proteome fractions instead of fractions of individual proteins. The respiratory

proteins converge at 0.474 ± 0.0002 h^{-1}.

Fig. 5 The effect of translation inhibition by cyclohexamide on growth rate, fluxes and proteome

fractions in controlled aerobic batch fermentations on glucose. a. Dependency of culture optical density

(OD) on the time post-inoculation to the medium supplemented with cycloheximide. Lines are values

of consecutive OD measurements, points represent the times when cultures were sampled. **b-d**

Comparison of pcYeast predictions and experimental data: lines are model predictions; symbols are

experimental data points. **b**. Main catabolic fluxes as a function of the growth rate. **c**. Ribosomal

proteome fractions. Data from Fig. 1c are included for comparison. **d.** Proteome fractions measured

for key metabolic pathways, and the minimal proteome fractions predicted by pcYeast.

Table 1. Statistics of the pcYeast model.

Process/Compartment	# of reactions	# of proteins
Total	**24422**	**1520**
Metabolic network	**5774**	**913**
from Yeast7.6	5738	909
manually added metabolic reactions	36	4
Cytoplasm	**2349**	**778**
Plasma membrane	**529**	**114**
Mitochondria	**1089**	**272**
Endomembrane system	**2127**	**133**
Metabolic complex formation, disassembly, dilution	**2787**	**-**
tRNA turnover and modification	**2194**	**56**
Protein synthesis and turnover	**13312**	**403**
Cytoplasmic translation	1512	138
Mitochondrial translation	8	89
Protein folding	1515	31
Protein degradation	1607	42
Protein misfolding, refolding	6061	73
Protein transport	1324	30
Protein dilution by growth	1285	-
Formation of macromolecular complexes	**355**	**196**

867   **Table 2.** Changes to the parameters for simulating sugar excess conditions. NGAM is non-growth

868   related ATP maintenance.

Growth condition	Unit	Glucose (naïve)	Galactose	Maltose
Maximal hexose transporter area	$\mu m^2/cell$	7.5	3.0	3.5
Carbon-related NGAM	$mmol/gDW/h$	0.0	3.0	0.0
Minimal UP fraction	$g\ UP/g\ protein$	0.245	0.49	0.34