A secondary metabolite drives intraspecies antagonism in a gut symbiont that is inhibited

by cell wall acetylation

Mustafa Özçam[1,2], Jee-Hwan Oh[1], Restituto Tocmo[1,3], Deepa Acharya[4], Shenwei Zhang[1], Theresa Astmann[1], Mark

Heggen[1], Silvette Ruiz-Ramírez[1], Fuyong Li[5], Christopher C. Cheng[6], Eugenio Vivas[7], Federico E. Rey[7,] Jan

Claesen[8], Tim Bugni[4], Jens Walter[5,6,9,10], Jan-Peter van Pijkeren[1,11*]

[1] Department of Food Science, University of Wisconsin-Madison, Madison, WI, 53706, USA

[2] Division of Gastroenterology, Department of Medicine, University of California San Francisco, CA, 94141, USA.

[3] Present Address: Department of Pharmacy Practice, University of Illinois at Chicago, Chicago, IL, 60612, USA

[4] Department of Pharmacy, University of Wisconsin-Madison, Madison, WI, 53706, USA

[5] Department of Agriculture, Food and Nutritional Science, University of Alberta, Edmonton, AB T6G 2P5, Canada

[6] Department of Biological Sciences, University of Alberta, Edmonton, AB T6G 2P5, Canada;

[7] Department of Bacteriology, University of Wisconsin-Madison, Madison, WI, 53706, USA

[8] Department of Cardiovascular and Metabolic Sciences and Center for Microbiome and Human Health, Lerner

Research Institute, Cleveland Clinic, Cleveland, OH, 44195, USA

[9] Department of Medicine and APC Microbiome Ireland, University College Cork, Cork T12 K8AF, Ireland;

[10] School of Microbiology, University College Cork, Cork T12 YT20, Ireland.

[11] Lead Contact

*Correspondence: vanpijkeren@wisc.edu

SUMMARY

The mammalian microbiome encodes numerous secondary metabolite biosynthetic gene

clusters, yet their role in microbe-microbe interactions is unclear. Here, we characterized two

polyketide synthase gene clusters (*fun* and *pks*) in the gut symbiont *Limosilactobacillus reuteri.*

The *pks*, but not the *fun* cluster, encodes antimicrobial activity. Forty-one out of 51 *L. reuteri*

strains tested are sensitive to Pks products, which was independent of strains' host origin. The

sensitivity to Pks was also established in intraspecies competition experiments in gnotobiotic

mice. Comparative genome analyses between Pks-resistant and sensitive strains identified an

acyltransferase gene (*act*) that is unique to Pks-resistant strains. Subsequent cell wall analysis of

the wild-type and the *act* mutant strains showed that Act acetylates cell wall components. The

pks mutants lost their competitive advantage and *act* mutants lost their Pks resistance *in vivo*.

Thus, our findings provide insight into how closely related gut symbionts can compete and co-

exist in the gastrointestinal tract.

39 **Keywords:** *Limosilactobacillus reuteri,* gut symbiont, polyketide, polyene, secondary

40 metabolite, biosynthetic gene clusters, acyltransferase, antimicrobial, acetylation, microbial

41 competition

INTRODUCTION

The mammalian gastrointestinal tract is inhabited by trillions of microorganisms that coexist with their host (Sender et al., 2016). While factors like host genetics, immune status, nutritional resources and colonization history affect microbial composition (Bonder et al., 2016; David et al., 2014; Hooper et al., 2015; Martínez et al., 2018; Snijders et al., 2016; Turnbaugh et al., 2009; Zarrinpar et al., 2014), the relationship between microbes is determined by competitive interactions (Boon et al., 2014). Bacteria have developed numerous strategies that mediate survival and competition, including the production of broad and narrow spectrum antimicrobials (Sassone-Corsi et al., 2016). On the other hand, acquisition of resistance genes and/or modification of the cell wall can help bacteria survive the antimicrobial warfare in the gut (Murray and Shaw, 1997).

Some bacterial secondary metabolites have antimicrobial activity against other community members. Their interactions—known as interference competition—are important in the assembly and the maintenance of microbial communities (Jacobson et al., 2018). Most secondary metabolites are produced by biosynthetic gene clusters (BGCs) and polyketide synthase (PKS) gene clusters form a prominent subclass, involved in the biosynthesis of carbon chain backbones from the repeated condensation of acyl-CoA building blocks (Lin et al., 2015; Medema et al., 2014). A survey of 2,430 reference genomes from the Human Microbiome Project uncovered more than 3,000 BGCs (Donia et al., 2014). So far, only seven PKS-like BGCs encoded by gut-associated bacteria have been functionally characterized (Figure S1).

As a class of polyketide synthase products, aryl polyenes are lipids with an aryl head group conjugated to a polyene tail (Lin et al., 2015) and are widely distributed in soil and host-associated bacteria (Cimermancic et al., 2014; Youngblut et al., 2020). Although, many polyene

compounds isolated from terrestrial and marine microbes possess antimicrobial effects *in vitro*

(Herbrík et al., 2020; Lee et al., 2020; Li et al., 2021; Zhao et al., 2021), virtually nothing is

known about their ecological role in microbe-host and microbe-microbe interactions in the

mammalian gastrointestinal tract (Aleti et al., 2019). While metagenome studies are critically

important to identify novel polyene-like BGCs within the microbiome (Hiergeist et al., 2015;

Medema et al., 2011), the assessment of model organisms and their isogenic mutants in the

appropriate ecological context is critical to advance our knowledge on the biological function of

these BGC-derived compounds.

Limosilactobacillus reuteri, until recently known as *Lactobacillus reuteri* (Zheng et al.,

2020), is a gut symbiont species that inhabits the gastrointestinal tract of various vertebrates,

including rodents, birds and primates (Duar et al., 2017). This, combined with the available

genome editing tools (Oh and Van Pijkeren, 2014; Van Pijkeren et al., 2012; Zhang et al., 2018),

make *L. reuteri* an ideal model organism to study microbe-microbe, microbe-phage and microbe-

host interactions (Lin et al., 2018; Oh et al., 2019; Özçam et al., 2019; Walter et al., 2011).

Previously, we screened a *L. reuteri* strain library derived from different hosts and identified that

L. reuteri R2lc activates the aryl hydrocarbon receptor (AhR), a ligand activated transcription

factor that plays an important role in mucosal immunity by inducing the production of

interleukin-22 (Zelante et al., 2013). By whole genome sequencing, comparative genomics, and

targeted gene deletion, we identified that R2lc harbors two genetically distinct PKS clusters (*pks*

and *fun*), each encoded from a multi-copy plasmid. We demonstrated that the *pks* but not the *fun*

cluster is responsible for R2lc mediated AhR activation (Özçam et al., 2019)

In this study, we found that the *pks* cluster in *L. reuteri* R2lc also encodes antimicrobial

activity. Intraspecies competition experiments in gnotobiotic mice with the wild-type and the *pks*

deletion mutant revealed that Pks expression can provide a competitive advantage. Remarkably,

a small number *L. reuteri* strains were resistant to the antimicrobial activity of Pks. We

discovered this resistance was driven by a gene encoding an acyltransferase that increases

acetylation of the cell wall Thus, our findings uncovered mechanisms by which gut symbionts

can compete and co-exist with closely related strains through secondary metabolite production

and cell-wall modification.

RESULTS

L. reuteri **Pks inhibits the competitor strain**

The rodent isolate *L. reuteri* R2lc contains two PKS clusters, *fun* and *pks* (Figure 1A). In

both the human and rodent gut ecosystem, *L. reuteri* R2lc outcompetes most *L. reuteri* strains,

including the rodent isolate *L. reuteri* 100-23 (Duar et al., 2017; Oh et al., 2009). Because select

PKS products have antimicrobial activity (Lin et al., 2015), we tested to what extent Pks and Fun

provide strain R2lc with a competitive advantage. We performed *in vitro* competition

experiments in batch cultures using 1:1 mixtures of the rodent gut isolates *L. reuteri* 100-23 and

L. reuteri R2lc wild type, or our previously generated PKS mutants (R2lcΔ*fun* or R2lcΔ*pks*)

(Özçam et al., 2019). Following 24 hours of incubation, the mixed cultures were plated. On agar,

R2lc and its derivatives form pigmented colonies, while 100-23 colonies are opaque in color. Co-

incubation of R2lc + 100-23 or R2lcΔ*fun* + 100-23 only recovered pigmented colonies; however,

co-incubation of R2lcΔ*pks* + 100-23 yielded a mixture of pigmented and opaque-colored

colonies (% pigmented:opaque colony distribution is 29:71) (Figure 1B). These data suggest that

the *pks* gene cluster, but not the *fun* gene cluster, provides strain R2lc with a competitive

advantage when co-cultured with *L. reuteri* 100-23.

Figure 1. A polyketide synthase cluster in *L. reuteri* R2lc produces polyene-like compounds that lyse competitor strains A) The *fun* cluster (top) spans 13.4 kb containing 12 Open Reading Frames (ORFs), and the *pks* cluster (bottom) spans 11.3 kb containing 15 ORFs. Transport-related, additional biosynthetic, regulatory, and other genes are represented by different colors. **B)** R2lc and R2lcΔ*fun* but not R2lcΔ*pks* inhibits *L. reuteri* 100-23. **C)** R2lc has bactericidal effect against 100-23. Single culture (OD_{600}=0.1) or co-cultures (OD_{600}=0.05 from each strain) were mixed and incubated in MRS broth (pH: 4.0, 37°C) and samples were collected every two hours for up to 12 hours. See also Figure S2. **D)** UPLC-PDA-MS analysis of R2lc and Δ*pks* mutant. R2lc but not Δ*pks* produces unique compounds with a maximum absorption of 380nm (Black: R2lc, blue: Δ*pks*). See also Figure S3. **E)** R2lc-Pks-mediated killing requires cell-cell interaction. Growth (24h) of R2lc and 100-23 separated by a 8 µm filter resulted in killing of 100-23 in a Pks-dependent manner (top left schematic; black bars) while separation by a 0.4 µm filter (top right schematic; blue bars) did not affect 100-23 survival, as determined by viable plate count. ns. no statistical significance ($p > 0.05$); **$p < 0.01$; ***$p < 0.001$. See also Figure S4. **F)** R2lc-Pks lyses competitor strain 100-23. Strain 100-23 was engineered to accumulate recombinant leptin intracellularly. Percentage leptin released (primary y-axis, green) and bacterial counts of 100-23 (secondary y-axis, black) were determined for a monoculture of 100-23 (top panel) and co-cultures of R2lc+100-23 (middle panel) and R2lcΔ*pks*+100-23 (bottom panel). All data shown are means ± SD and are based on three biological replicates.

L. reuteri **Pks has a time-dependent bactericidal effect**

  To understand the dynamics by which R2lc outcompetes 100-23, we performed a time course competition experiment in batch cultures. Mixtures (1:1, $OD_{600} = 0.05$ per strain) of R2lc

+ 100-23, and R2lcΔ*pks* + 100-23 were prepared. As a control, a monoculture of 100-23 was

grown. We harvested samples every two hours and determined the CFU levels by standard plate

count. We observed that R2lc and R2lcΔ*pks* have a similar growth dynamic (μ_{max} of

R2lc= 0.19±0.02, μ_{max} of R2lcΔ*pks*= 0.16±0.003, p = 0.18), and 100-23 grows slightly faster

compared to R2lc and R2lcΔ*pks* in mono-culture (Figure S2A). Up to six hours, mixtures of

R2lc + 100-23 and R2lcΔ*pks* + 100-23 yielded similar levels of 100-23 colonies, which were

comparable to the levels obtained when 100-23 was cultured independently. However, after eight

and ten hours of co-incubation of R2lc + 100-23, 100-23 CFU levels were reduced by three and

five orders of magnitude, respectively, while 100-23 continued to grow in the R2lcΔ*pks* + 100-23

mixture (Figure 1C). Importantly, the sharp decline in 100-23 CFU counts suggests Pks elicits a

strong bactericidal activity against 100-23. Also, the fact that killing of 100-23 is initiated after

6-hours of co-culture suggests that production of Pks may be growth phase dependent.

149   In mice, *L. reuteri*—including strain 100-23—form a biofilm on the stratified squamous

epithelium of the forestomach (Frese et al., 2013; Lin et al., 2018; Savage et al., 1968). Cell

numbers of 100-23, R2lc or R2lcΔ*pks* in gastric biofilms were similar ($4.3\times10^8 \pm 2.6\times10^8$,

$2.0\times10^8 \pm 3.0\times10^7$, $1.7\times10^8 \pm 4.1\times10^7$ CFU/ml, respectively) (Figure S2A). In biofilm co-

cultures, however, R2lc was 750-fold more abundant than 100-23 (p = 0.002). The ability of

R2lc to outcompete 100-23 in biofilm is mediated by Pks, because co-incubation of R2lcΔ*pks*

with 100-23 recovered similar levels of CFUs for both strains (p = 0.27) (Figure S2B).

***L. reuteri pks* cluster produces unique polyene-like compounds**

158   To gain more insight into *L. reuteri* R2lc Pks production *in vitro*, we performed Ultra

Performance Liquid Chromatography coupled with Photodiode Array and Mass Spectrometer

(UPLC-PDA-MS) analysis. By comparing the chromatograms obtained from R2lc and R2lcΔ*pks*

cultures, we found that R2lc produces a family of unique compounds with a maximum

wavelength of 380 nm. These compounds eluted towards the end of the gradient run where the

solvent composition was close to 100% methanol, indicating a hydrophobic characteristic. The

large retention time of the compounds on the C18 column and the maximum absorption

wavelength are consistent with previously identified polyene compounds (Gruber and Steglich,

2007). Therefore, the putative products of the *pks* cluster are predicted to be polyene compounds.

The UV chromatogram extracted from the R2lc culture at different time points shows an

increase in the production of these compounds up to 12 hours of incubation, after which the

intensity reduces as shown in samples at 24 hours (Figure 1D). We subsequently analyzed the

Liquid Chromatography-Mass Spectrometry (LC-MS) data of R2lc and Δ*pks* samples collected

above to identify the polyene products produced by R2lc. The mass spectrum of the peak

between 10.52 min – 10.58 min in the R2lc chromatogram at 12 hours clearly showed an ion

with m/z [M+H]$^+$ value of 257.1172 and corresponding m/z [M+Na]$^+$ value of 279.0092, which

were absent in the 12-hour culture of Δ*pks* (Figure S3). The high-resolution mass for the

compound led to the predicted molecular formula of $C_{16}H_{16}O_3$. The reduction of Pks compounds

after 12 hours of incubation suggests instability of Pks molecules in our experimental setup.

Taken together, these data suggest that the antibacterial compound is produced in a time-

dependent manner.

Pks producing *L. reuteri* R2lc lyses competitor strain through cell-cell interaction

To further enhance our understanding of *L. reuteri* R2lc Pks-mediated killing, we performed an

experiment where strains R2lc and 100-23 were cultured in individual chambers separated by a

0.4 µm or 8µm filter (Figure 1E, top schematic). The 0.4 and 8 µm membranes will allow

flowthrough of small molecules, metabolites and secreted soluble proteins; only the 8 µm

membrane allows flowthrough of bacteria. When the chambers were separated by a 8 µm filter,

we recovered ~ 1,000-fold less 100-23 bacteria in the presence of R2lc wild-type compared to

the 100-23 monoculture ($p < 0.001$). In the presence of R2lcΔpks, levels of 100-23 were 3-fold

less compared to the 100-23 monoculture (Figure 1E, left panels; $p < 0.01$). Viability of 100-23

was not impacted by R2lc when the chambers were separated by a 0.4 µm filter (Figure 1E, right

panels). Since we did not observe antimicrobial activity when cultures were separated by a 0.4

µm membrane, which does allow flowthrough of metabolites, it is plausible that the

antimicrobial compound is not secreted into the extracellular milieu.

To expand on these findings, we performed double-layer agar diffusion assays with a)

filter-sterilized cell-free supernatant of R2lc, b) methanol and acetone extracts of the supernatant,

c) live R2lc cells in MRS and PBS, d) R2lc cells killed following heat- or acetone treatment. Our

findings further validate that the antimicrobial compound is not secreted (Figure S4AB); instead,

only antimicrobial activity is observed when a cell suspension of viable R2lc (Figure S4C) is

used whereas no zone of inhibition was observed with a cell suspension of viable R2lcΔpks

(Figure S4, bottom panel). Also, no zone of inhibition was observed when we inactivated R2lc

by either heat or acetone treatment (Figure S4D). In conclusion, to elicit an antimicrobial effect,

strain R2lc requires to be metabolically active and our data strongly suggest cell-cell interaction

is required.

Next, we designed an experiment to determine if Pks-producing *L. reuteri* R2lc lyses the

target cells using an analogous approach as we previously established (Alexander et al., 2019).

We developed 100-23/leptin, a derivative of strain 100-23 that is engineered to accumulate leptin

intracellularly. Following 12 hours batch culture of the 100-23/leptin monoculture, less than 15%

of the total amount of recombinant leptin was released in the supernatant (Figure 1F, top panel).

We expect that this relatively small amount of leptin is derived from natural lysis, for example,

by biologically active prophages present in 100-23 (Oh et al., 2019). When R2lc Pks is co-

cultured with 100-23/leptin, we found that a significant reduction of 100-23 CFU count directly

correlates to an increase of leptin in the supernatant (Figure 1F, middle panel). However, the co-

culture of R2lcΔ*pks* + 100-23/leptin (Figure 1F, bottom panel) yielded levels of 100-23 and

leptin that were comparable to those observed in the 100-23/leptin monoculture (Figure 1F, top

panel). Collectively, these data show that *L. reuteri* R2lc needs to be alive and likely requires

cell-cell contact to elicit a Pks-mediated bactericidal effect that leads to lysis of the competitor

strain.

L. reuteri* Pks molecules provide a competitive advantage *in vivo

219   To characterize the ecological role of the *pks* antibacterial secondary metabolite gene

cluster *in vivo*, we gavaged germ-free mice (n=4/group) with a 1:1 mixture of R2lc+100-23, or

R2lcΔ*pks*+100-23. To quantify each strain in the fecal material, we engineered R2lc and its

derivatives to encode chloramphenicol resistance (R2lc::*Cm* and R2lcΔ*pks::Cm*) while strain

100-23 was rifampicin-resistant (100-23—Rif[R]). We determined the competition ratio between

R2lc and 100-23 (competition ratio: R2lc CFU count/competitor strain's CFU count) over a

period of six days in feces. We found that *L. reuteri* R2lc wild-type but not the Δ*pks* mutant

gradually outcompetes the 100-23 strain. The R2lc:100-23 competition ratio increased from

107:1 (day 1) up to 980:1 (day 6), while the competition ratio between R2lcΔ*pks*:100-23

declined from 35:1 (day 1) to 4:1 (day 6) (Figure 2A). Similarly, the R2lc:100-23 competition

ratio was higher than the R2lcΔ*pks*:100-23 competition ratio in the forestomach (31,973-fold vs

7.8-fold, *p* = 0.03), cecum (1,515-fold vs. 2.9-fold, *p* = 0.03), colon (1,474-fold vs. 8.9-fold, *p* =

0.03) and jejunum (1,114-fold vs. 5.2-fold, *p* = 0.03) (Figure 2B). Thus, our findings

demonstrate that Pks provides *L. reuteri* R2lc with a competitive advantage throughout the

murine intestinal tract.

Figure 2. *L. reuteri* Pks molecules provide a competitive advantage *in vivo*.

A 1:1 mixture of R2lc+100-23, or R2lcΔ*pks*+100-23 was administered by oral gavage to germ-

free mice. **A)** Competition ratios of the indicated strains in fecal and **B)** intestinal content (day 6)

were determined. In box and whisker plots, the whiskers represent the maximum and minimal

values, and the lower, middle and upper line of the box represent first quartile, median and third

quartile, respectively. Circles represent data from individual mouse. Statistical significance was

determined by Wilcoxon / Kruskal-Wallis Tests; p<0.05 is considered statistically significant.

F.S.: Forestomach.

***L. reuteri* Pks-mediated antimicrobial effect is strain specific**

Now we established that *L. reuteri* Pks provides an *in vivo* and *in vitro* competitive

advantage against strain 100-23, we next investigated to what extent Pks-mediated killing is

strain specific. We performed *in vitro* competition experiments with 51 *L. reuteri* gut isolates

from different host origins: human, rat, mouse, chicken and pig. To quantify strains, we used

derivatives of R2lc and R2lcΔ*pks* that were engineered to be chloramphenicol resistant. For each

competitor strain we isolated a natural rifampicin-resistant mutant. Strain R2lc and the

competitor strain were mixed 1:1 in MRS medium adjusted to pH 4.0, which is pH of the

forestomach, the natural habitat of strain R2lc. The co-cultures were incubated for 24 hours, and

appropriate dilutions were plated on MRS plates containing chloramphenicol (5 µg/ml, for R2lc

and R2lcΔ*pks*) or rifampicin (25 µg/ml, for competitor strain). Competition ratios were

determined after 24 hours of incubation.

Based on non-parametric statistical tests, we identified that R2lc inhibits 41 out of 51

(80.4%) *L. reuteri* strains. Specifically, six out of nine (66.6%) rat-isolates, seven out of ten

(70%) mouse isolates, 16 out of 18 (89%) pig isolates, eight out of nine (88.9%) chicken isolates

and two out of five (40%) human isolates were significantly inhibited by R2lc (R2lc vs

competitor CFU, *P <0.05*). To test if the competitive advantage of R2lc is driven by Pks, we

repeated the competition experiments with R2lcΔ*pks*. We found that deletion of the *pks* cluster

resulted in loss of antimicrobial activity and the Δ*pks* mutant did not show strong inhibition

against competitor strains. However, Δ*pks* mutant still inhibited some competitor strains (16 out

of 51) to a small extent, with observed differences in Colony Forming Units (CFUs) between

Δ*pks* and competitor strains less than two orders of magnitude (Figure 3). These data

demonstrate that R2lc-Pks mediated antimicrobial effect is strain specific and 10 out of 51 tested

L. reuteri strains (19.6%) are not inhibited/outcompeted by the Pks producing strain.

Figure 3. *In vitro* **competition ratio of *L. reuteri* R2lc and R2lc∆*pks* with a panel of 51**

***L. reuteri* strains from different host origin.** Competition ratios between R2lc and the

competitor strain (black) and R2lc∆*pks* and the competitor strain (red) after 24h co-incubation in

MRS (pH 4.0). Data shown are based on at least three biological replicates. The whiskers

represent the maximum and minimal values, and the lower, middle and upper line of the box

represent first quartile, median and third quartile, respectively. Host origin (top) and the *L.*

reuteri phylogenetic lineage (color coding) are indicated. Statistical significance was determined

by Wilcoxon / Kruskal-Wallis Tests. Asterisks (*) represents statistical significance between

R2lc vs competitor strain for the top panel (p<0.05 considered as significant). See also Table S3.

L. reuteri strains resistant to Pks encode an acyltransferase enzyme

Next, we aimed to understand what the underlying mechanism is that allows select strains

to be resistant to *L. reuteri* R2lc Pks. We performed comparative genome analyses to identify

genes unique to strains that are resistant to *L. reuteri* R2lc Pks. We included 24 genome

283   sequences in our analyses, of which four genome sequences were derived from resistant strains

284   (mlc3, Lr4000, 6475 and 20016^T) (Table S1). Our initial genome comparison analyses revealed

285   seven genes unique to three of the four R2lc-resistant strains (mlc3, 6475 and 20016^T) (Table

286   S2).

287   One of the seven unique genes (*act*, Lreu_1368 in strain DSM20016^T, Table S2) is

annotated as an O-acyltransferase. This gene became our focus as several studies demonstrated

that deletion of homologous *act* genes reduces the resistance to enzymes that target the

peptidoglycan cell wall in other Gram-positive bacteria (reviewed in (Ragland and Criss, 2017)).

Moreover, resistance to antimicrobial molecules is typically associated with altered cell-wall

peptidoglycan structures (Vollmer, 2008). In *L. reuteri*, the *act* gene is located in the surface

polysaccharide (SPS) gene cluster. Based on the available genome sequences, we found that

three resistant strains (mlc3, 6475 and 20016^T) all putatively encode a nearly identical Act

protein (≥99% amino acid identity). Three R2lc-Pks-sensitive strains (CR, ATCC 53608 and

one-one) putatively encode a distinct Act protein with 54% amino acid identity to the predicted

Act amino acid sequence of resistant *L. reuteri* strains. Six additional R2lc-Pks-sensitive strains

(I5007, 6799jm-1, lpup, CF48-3A1, Lr4020 and SD2112) putatively encode Act with 26-28%

amino acid identity compared to the predicted Act amino acid sequence of resistant *L. reuteri*

strains. The remainder of the 12 strains that are sensitive to R2lc-Pks do not contain the *act* gene

in the SPS gene cluster (Figure 4).

Figure 4. **Variability in gene content of Surface Polysaccharide gene cluster in *L. reuteri.***

Comparison of surface polysaccharide (SPS) gene clusters of strains resistant and sensitive to

R2lc-Pks. Strains #1-3 (resistant) each putatively encode a nearly identical Act protein with

≥99% amino acid identity; strains #4-6 (sensitive) putatively encode Act that shares 54% amino

acid identity to Act of resistant strains; strains #7-9 (sensitive) putatively encode Act that shares

312   25-28% amino acid identity to Act of resistant strains; strains #10-12 (sensitive) lack the *act*

313   gene. Genes are color-coded with their predicted functions based on the annotation of the SPS

314   gene cluster in *L. reuteri* DSM20016^T. See also Table S1 and Table S2.

315
316   To map which *L. reuteri* strains are resistant or sensitive to R2lc-Pks, we constructed a

317   phylogenetic tree of Act amino acid sequences from 134 *L. reuteri* strains. We found that 43 out

318   of 134 strains (32.3%) contain an *act* gene whose putative product shares 99-100% amino acid

319   identity to the predicted amino acid sequence of resistant *L. reuteri* strains, while the remaining

320   strains putatively encode Act with 23-72% amino acid identity (Figure 5). Also see Table S6.

Figure 5. Phylogenetic tree of *L. reuteri* Act homologs. We constructed a phylogenetic tree

based on Act amino acid sequences derived from 134 Act encoding *L. reuteri* strains to predict

R2lc-resistance and sensitive strains. Forty-three out of 133 strains (32.3%) encode and *act* gene

that shares 99-100% amino acid identify with the *act* gene in ATCC 6475 strain, thus, predicted

to be R2lc-resistant (blue ring). The remaining strains encode an *act* gene that shares 23-54%

amino acid identify with the *act* gene in PTA 6475 (yellow ring). See also Table S6.

The acyltransferase gene in *L. reuteri* confers protection to R2lc-Pks products

To test if the *act* gene confers protection to R2lc-Pks products, we generated *act* mutants

and complemented strains of mlc3 (rat isolate, clade I) and ATCC 6475 (human isolate, clade I).

We observed that strain 6475Δ*act* but not mlc3Δ*act* grows slower *in vitro* compared to the parent

wild-type strains. Growth was partially restored when 6475Δ*act* is complemented (with pNZ-

act), while strain mlc3Δ*act* (with pNZ-*act*) grew faster than wild-type (Figure S5A). In these

strains, we found that inactivation of *act* increased the sensitivity to R2lc-Pks, and *act*

complementation decreased the sensitivity to R2lc-Pks, as is evident from the increased

competition ratios between R2lc *vs* Δ*act* compared to the competition ratios between R2lc *vs*

wild-type and complemented strain (Figure 6A, left panel). In contrast, deletion or

complementation of *act* gene in mlc3 and 6475 did not change the competition ratio of these

strains against R2lcΔ*pks* (Figure 6A, right panel). Overall, our data showed that Act in *L. reuteri*

provides resistance to R2lc-Pks mediated killing.

L. reuteri **Act provides resistance against antimicrobial Pks molecules** *in vivo*

To determine to what extent Act protects against R2lc Pks *in vivo*, we performed

competition experiments in mice. Germ-free mice were administrated with a 1:1 mixture of any

of the following strain combinations: R2lc ± *pks* and mlc3 ± *act* (Figure 6BC). Following

administration, fecal samples were analyzed at day one, three and six, and animals were

sacrificed at day 7 after which we analyzed microbial counts in the stomach, jejunum, cecum and

colon. Fecal analyses revealed that the competition ratios between R2lc and mlc3 or its isogenic

Δ*act* mutant were comparable ($P > 0.05$); however, at day 7, the competition ratio of

R2lc:mlc3Δ*act* was significantly larger compared to R2lc:mlc3 (2,183 ± 3,060-fold *vs* 95 ±

58-fold, respectively; $P < 0.01$) (Figure 6B, left panel). We also observed that in all intestinal

tissues the competition ratio of R2lc∷mlc3Δ*act* was larger compared to R2lc∷mlc3. Specifically,

forestomach (3,777 ± 2,259-fold *vs* 31 ± 48-fold; $P < 0.01$), jejunum (9,510 ± 15,239-fold *vs* 62

± 55-fold; $P < 0.01$), cecum (9,872 ± 17,683-fold *vs* 115 ± 99-fold; $P < 0.01$), colon (959 ±

4,703-fold *vs* 118 ± 104-fold; $P < 0.01$) (Figure 6B, right panel). Thus, the *act* gene provides

protection against R2lc-Pks *in vivo*. To test to what extent R2lc Pks provides R2lc with a

competitive advantage in these experiments, we performed competition experiments with

R2lcΔ*pks*∷mlc3 and R2lcΔ*pks*∷mlc3Δ*act* (Figure 6C). Overall, the competition ratios of

R2lcΔ*pks*∷mlc3 and R2lcΔ*pks*∷mlc3Δ*act* are lower compared to the competition ratios observed

with R2lc wild-type, which demonstrates a clear role for Pks in providing R2lc with a

competitive advantage *in vivo*. Interestingly, the R2lcΔ*pks*∷mlc3Δ*act* competition ratio is higher

than that of R2lcΔ*pks*∷mlc3 in the stomach (1.5-fold vs 0.5-fold, $p = 0.02$), jejunum (2.2-fold vs

0.5-fold, $p = 0.01$) and cecum (2.2-fold vs 1.2-fold, $p = 0.04$) (Figure 6C, right panel), which

could be linked to the role of Act in lysozyme resistance during *in vivo* colonization (Bera et al.,

2005). Indeed, we tested the lysozyme susceptibility of wild-type and *act* mutant strains and

identified that the Δ*act* variants are more susceptible to lysozyme while the complemented

strains survived at levels comparable to that of the wild-type strains (Figure S5B). Therefore,

host lysozyme may also suppress intestinal survival of mlc3Δ*act* thereby providing R2lcΔ*pks* a

competitive advantage. When we tested the susceptibility of the *act* mutants to ampicillin we

determined that the *act* mutants of 6475 (Figure S5C) and mlc3 (Figure S5D) are more sensitive

to ampicillin, while the phenotype of complemented strains was similar to the wild-type strain.

Because Act provides protection to lysozyme and ampicillin, which both attack the

peptidoglycan (Teethaisong et al., 2014; Vanderkelen et al., 2011), we aimed to understand how

Act modifies the peptidoglycan complex.

Figure 6. **The *act* acyltransferase gene protects against R2lc-Pks molecules. A**) Act provides

resistance to R2lc-mediated killing (left panel), which is mediated by the inhibitory activity of an

intact Pks (right panel). **B and C**) Competition ratios of the indicated strains in the previous

germ-free mice in fecal (left panel) and intestinal contents (right panels). When data are

presented as box and whisker plots, the whiskers represent the maximum and minimal values,

and the lower, middle and upper line of the box represent first quartile, median and third quartile,

respectively. Points represent individual data points. Statistical significance was determined by

Wilcoxon / Kruskal-Wallis Tests, p=P value. P<0.05 considered significant. See also Figure S5.

The acyltransferase gene increases acetylation of the cell wall

The bacterial Act enzyme acetylates the C6 hydroxyl of *N*-acylmuramic acid in

peptidoglycan (PG) (Moynihan and Clarke, 2010). To understand the role of *L. reuteri* Act in

acetylation, we performed HPLC analyses to determine acetylation levels of the cell wall in Pks-

resistant and -sensitive strains, relative to total cellular muramic acid (MurN) content. Analysis

of cell wall extracts derived from *L. reuteri* 100-23, a Pks-sensitive strain which does not encode

Act, revealed that 6.65 ± 1.19% was acetylated relative to total MurN. Next, we cloned the *act*

gene from strain mlc3 in the inducible expression vector pSIP411, which was established in

strain 100-23. Induced expression of *act* in 100-23 increased relative acetylation levels to 28.7 ±

9.0%. Inactivation of *act* in *L. reuteri* strains resistant to R2lc-Pks reduced total acetylation

levels (from 70.1 ± 4.9% to 2.3 ± 0.3% for mlc3; and from 61.2 ± 9.1% to 21.3 ± 12.9% for

6475). Overexpression of the *act* gene in mlc3Δ*act* and 6475Δ*act* restored the relative acetylation

levels (from 2.3 ± 0.3% to % to 51.8 ± 15.1% for mlc3Δ*act::act* ; and from 21.3 ± 12.9% to 50.9

± 9.2% for 6475Δ*act::act*) (Table 1). Together these data suggests that *L. reuteri* Act is

responsible for increased acetylation of the cell wall, which we demonstrated is linked to the

resistance to the antimicrobial compound produced by R2lc.

403
404 **Table 1.** O-acetylation levels were determined by base-catalyzed hydrolysis and release of
405 acetate. Percent O-acetylation is reported relative to muramic acid (MurN) content.

Strain	% O-acetylation
100-23	6.65 ± 1.19
100-23 + pSIP control	3.46 ± 0.07
100-23::*act*	28.70 ± 9.01
mlc3	70.12 ± 4.85

mlc3Δ*act* + pSIP control	2.31 ± 0.29
mlc3Δ*act::act*	51.76 ± 15.09
6475	61.21 ± 9.07
6475Δ*act* + pSIP control	21.29 ± 12.92
6475Δ*act::act*	50.90 ± 9.18

406 **ᵃ** Results are from one representative biological replicate measured in triplicate and reported as
407 mean with Standard Deviation.
408

Discussion

In this study, we provided mechanistic insight into intra-species antagonism based on a

secondary metabolite in the gut symbiont *L. reuteri*. We discovered that a biosynthetic gene

cluster (BGC) provides a competitive advantage in the gastrointestinal tract by killing closely

related microbes in a strain-specific manner. Expression of the act gene by select strains of *L.*

reuteri increased resistance against this killing activity.

Closely related strains that occupy the same niche are subject to fierce competition

(Hibbing et al., 2010; Segura Munoz et al., 2020). While the co-existence of two closely related

strains may be stable in the host, long-term evolution may lead to different ecological outcomes

(Edwards et al., 2018). Either evolution leads to changes that form a more stable population—

lineages are overall less competitive—or evolution results in increased fitness of one lineage.

The lineage with increased fitness will replace the competing lineage, or the competing lineage

may evolve in a way that will allow it to persist (Le Gac et al., 2012). Our findings are

interesting in light of these ecological theories. Specifically, four distinct lineages have thus far

been identified among rodent *L. reuteri* isolates (Frese et al., 2011). Our competition analyses

revealed that nearly all strains in lineage I are sensitive to R2lc-Pks products while most strains

in lineage III are resistant. One potential explanation for the latter is the broad distribution of

nearly identical *act* genes in *L. reuteri* strains from different host origin (Figure 5, blue ring),

which suggests *act* exerts an evolutionary pressure that will increase the fitness in the gut

ecosystem. The findings presented in this work, the observation that aryl polyene gene clusters

are widely distributed throughout the host-associated bacteria (Cimermancic et al., 2014), and

that *L. reuteri* 6475Δ*act* has reduced gastrointestinal survival (preliminary data) support the

importance of Act in gut fitness and make it collectively less likely that *act* is acquired through a

neutral process such as genetic drift.

The evolution of a vertebrate symbiont with its host might be reciprocal, resulting in co-

evolution. The beneficial traits that a microbe exerts on its host may have been shaped by natural

selection as they promote host fitness, which is critical for the microbe to thrive in its niche.

R2lc-Pks molecules activates the aryl-hydrocarbon receptor (AhR), a ligand activated

transcription factor (Özçam et al., 2019). Activation of AhR has been shown to induce

production of interleukin-22 (IL-22) (Veldhoen et al., 2008), which enhances the innate immune

response by inducing antimicrobial peptides (i.e. Reg3-lectins) production from the mucosal

layer (Vaishnava et al., 2011; Zheng et al., 2008). It is intriguing to speculate that Pks molecules

provide R2lc with a host-mediated competitive advantage as Pks-mediated AhR activation leads

to antimicrobial peptide production by the host. Collectively, this would support the coevolution

between select microbes and their host.

This work contributes to our understanding of the ecological role of a gut symbiont-

encoded secondary metabolite and provides novel insight into how closely related strains

developed resistance against these antimicrobial molecules by acetylation of the cell-wall

polysaccharide complex. It is plausible bacteria have evolved other mechanisms to survive

exposure to Pks. In this work, we did encounter one strain—the rodent isolate Lr4000—that does

not encode Act but was resistant to R2lc Pks molecules during the batch culture competition

experiments and the mechanism of resistance remains to be elucidated.

With a number of genome editing tools available for *L. reuteri*, we are in a position to use

genetic approaches to study microbe-microbe interactions in a gut symbiont. When these

experiments are placed in the right ecological context, mechanistic insight can be provided on

the formation and stability of microbial communities, which is expected to provide insight how

microbial diversity is maintained within complex ecosystems. This knowledge can be leveraged

to provide rational approaches to select probiotics and next-generation probiotics to ultimately

promote animal and human health.

ACKNOWLEDGEMENTS

We thank Siv Ahrné (Lund University, Sweden) for providing *L. reuteri* R2lc, N2J, N2D,

and N4I, BioGaia AB (Stockholm, Sweden) for providing *L. reuteri* strains ATCC PTA 6475, and

Joseph Skarlupka for technical assistance. We are grateful to the College of Agricultural Life

Sciences (CALS) Statistical Consulting Lab for their assistance in the statistical analysis, and the

University of Wisconsin Biotechnology Center DNA Sequencing Facility for their services. This

work was supported by startup funds from the University of Wisconsin-Madison to J.P.V.P., the

UW-Madison Food Research Institute, and the United States Department of Agriculture, National

Institute of Food and Agriculture (USDA NIFA) Hatch award MSN185615 and grant no. 2018-

6717-27523. M.Ö. received financial support from the Turkish Ministry of National Education and

from the Department of Food Science. M.Ö. and S.Z. were the recipients of the Robert H. and

Carol L. Deibel Distinguished Graduate Fellowship in Probiotic Research, which was awarded by

the Food Research Institute (UW-Madison). J.C. is supported by National Institutes of Health grant

R01 AI153173. Gnotobiotic work was partly supported by the Office of the Vice-Chancellor for

Research and Graduate Education at the University of Wisconsin–Madison, with funding from the

Wisconsin Alumni Research Foundation.

AUTHOR CONTRIBUTIONS

M.Ö.: conceptualization, methodology, formal analysis, investigation, writing–original draft,

writing–review&editing, visualization; J-H.O.: methodology, formal analysis, investigation,

visualization; R.T.: methodology, validation, formal analysis, investigation; D.A.: methodology,

validation, formal analysis; S.Z.: formal analysis, investigation, visualization; T.A.: validation,

formal analysis, investigation; M.H.: validation, investigation; S.R-R.: validation; F.L.:

investigation; C.C.C.: validation; E.V.: methodology; F.E.R.: methodology, resources; J.C.:

investigation; T.B.: resources, supervision; J.W.: resources, writing—original draft, supervision;

J.P.V.P.: conceptualization, methodology, writing–original draft, writing–review&editing,

supervision, project administration, funding acquisition

DECLERATION OF INTEREST

JPVP received unrestricted funds from BioGaia, AB, a probiotic-producing company.

JPVP is the founder of the consulting company Next-Gen Probiotics, LLC. JW has received

grants and honoraria from several food and ingredient companies, including companies that

produce probiotics. JW is a co-owner of Synbiotic Solutions, LLC, and is on the Scientific

Advisory Board of Alimentary Health. M.Ö. was an employee of DuPont Nutrition and

Biosciences. JC is a Scientific Advisor for Seed Health, Inc.

STAR METHODS

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources should be directed to and will be fulfilled by the

Lead Contact, Jan-Peter van Pijkeren (vanpijkeren@wisc.edu).

Material Availability

Requests for the plasmids and strains used in this study should be directed to and will be fulfilled

by the Lead Contact, Jan-Peter van Pijkeren (vanpijkeren@wisc.edu).

Data and Code Availability

506   Any additional information required to reanalyze the data reported in this paper is available from

507   the lead contact upon request.

510   **KEY RESOURCE TABLE**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Bacterial strains		
See Table S3 for bacterial stains used in this study		
Chemicals		
Agar	Alfa	Cat#A10752
Agarose	IBI	Cat# IB70042
LB broth	Acumedia, Neogen	Cat# 7290A
MRS broth	BD	Cat# 288110
M17 broth	BD	Cat# 218561
Potassium Buffered Saline	Gibco	Cat# 14190-144
0.45-µm-pore-size cellulose acetate membrane	Sigma	Product# WHA69012504
0.4-µm-pore-size filter insert	Cell Treat	Part no 230607
8-µm-pore-size filter insert	Cell Treat	Part no 230611
Acetic acid standard solution	Fisher Scientific	Cat# A38S-500
Sulfuric acid	Fisher Scientific	Cat# A300-500
Protease	Sigma	Product# P5147
Magnesium sulfate	Fisher Scientific	Cat# M65-500
RNase A	Thermo Scientific	Cat# EN0531
DNase I	Invitrogen	Cat# 18068015
a-amylase	Sigma	Cat# 10065
Sodium cloride	Fisher Scientific	Cat# S271-500
Tris-HCl	Calbiochem	Product# 648310-M
Sodium dodecyl sulfate	Sigma	Product# L4509
24-well plates	Fisher Scientific	Cat# 12556006
Agarose	DOT Scientific Inc.	Cat# DS170042
Pellet paint	VWR International	Cat# 9049-3
Chow diet	LabDiet	Cat# 5201
DpnI	Fisher Scientific	Cat# FERFD1704
0.22 µm filter, Polyvinylidene fluoride [PVDF]	Millipore	Cat# SLGVM33RS
Sodium hydroxide	Fisher Chemical	Cat# S318100
Methanol	Fisher Chemical	Cat# A456-500
Formic acid	Fisher Chemical	Cat# A11350
Chloramphenicol	Dot Scientific	Cat# DSC61000-25
Erythromycin	Fisher Scientific	Cat# BP920-25

Rifampicin	TCI Chemicals	R0079-25G
Vancomycin	Chem-Impex	Cat# 00315
Ampicillin	DOT Scientific	Cat#DSA40040-25
Ethanol	KOPTEC	Cat# V1016
P_{sak} Induction peptide	Peptide2.0	Sequence: MAGNSSNFIHKIKQIFTHR
Choice Taq DNA polymerase master mix	Denville Scientific	Cat# CB4070-8
Phusion Hot Start polymerase II	Thermo Scientific	Cat# F-549-L
T4 DNA ligase	Thermo Scientific	Cat# EL0011
Polynucleotide kinase	Thermo Scientific	Cat# EK0031
dATP	Promega	Cat# U120A
Ampligase	Lucigen	Cat# A32750
Sodium Hydroxide	Fisher Scientific	Cat# S318-500
HCl	Ricca Chemical Company	Cat# 3440-1
Zirconia/silica beads	Bio Spec	Cat#11079101Z
Critical Commercial Assay Kits		
Qubit dsDNA quantification kit	Invitrogen	Cat# Q32853
Genomic DNA purification kit	Promega	Cat# A1120
GeneJET PCR purification kit	Thermo Scientific	Cat# K0701
Plasmid isolation kit	Promega	Cat# A9340
Leptin ELISA kit	R&D systems	Cat# MOB00B
Experimental Models: Organisms/Strains		
C57BL/6J, Male	Jackson Laboratory	Cat# 000664
Oligonucleotides		
See Table S4 for primers used for this study		
Recombinant DNA		
See Table S5 for plasmids used for this study		
Software and Algorithms		
JMP Pro	SAS	Ver. 11.0.0
Datagraph	Visual Data Tools Inc.	Ver. 4.2.1
Bruker Hystar software	Bruker	Hystar 3.2.
MEGA	(Kumar et al., 2018)	Ver. 10.1.7

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Microbial Strains and Growth Conditions

The strains and plasmids used in this study are listed in Table S3 and S5. *L. reuteri*

strains were cultured in De Man Rogosa Sharpe (MRS) medium (Difco, BD BioSciences). For

growth curve experiments, *L. reuteri* strains were grown for ~16 hours and transferred to pre-

warmed MRS with a starting inoculum of $OD_{600}= 0.1$. Where appropriate, antibiotics were added

to the medium as described below. We determined the OD_{600} at regular intervals as shown in

growth curve figures. For *in vitro* competition and biofilm formation experiments, we used filter-

sterilized (0.22 µm PVDF filter, Millipore) MRS that was adjusted to pH 4.0 with 1 M HCl.

Unless stated otherwise, we prepared bacterial cultures as follows: *L. reuteri* strains were

incubated at 37°C under hypoxic conditions (5% CO_2, 2% O_2). The MRS agar plates were

incubated for 24 hours for colony counting. *Escherichia coli* EC1000 was used as general

cloning host and cultured with aeration at 37°C in lysogeny broth (LB, Teknova). *Lactococcus*

lactis MG1363 harboring pJP042 (VPL2047) was cultured under static condition at 30°C in M17

broth that was supplemented with glucose (0.5% [w/v]). Electrocompetent *E. coli* EC1000 were

prepared as described in (Sambrook and Russell, 2006). Electrocompetent *L. reuteri* cells were

prepared as described in (Van Pijkeren and Britton, 2012). If applicable, MRS for *L. reuteri* was

supplemented with 5 µg/ml erythromycin, 5 µg/ml chloramphenicol or 25 µg/ml rifampicin**.**

Mice

Ethics statement

All mouse experiments were performed in accordance with NIH guidelines, Animal

Welfare Act, and US federal law and were approved by the Application Review for Research

Oversight at Wisconsin (ARROW) committee and overseen by the Institutional Animal Care and

Use Committee (IACUC) under protocol ID M005149-RO1-A01. Conventional pathogen-free and

germ-free mice were housed at Animal Science and Laboratory of Animal Research Facilities

respectively at the University of Wisconsin-Madison.

Mice Strains and Husbandry

Twelve-week-old germ-free male B6 mice (C57BL/6) were maintained in a controlled

environment in plastic flexible film gnotobiotic isolators with a 12 hours light cycle. Sterilized

food (standard chow, LabDiet 5001, St. Louis, MO) and water were provided *ad libitum*.

Reagents and Enzymes

To amplify DNA fragments for cloning and screening, we used Phusion Hot Start DNA

Polymerase II (Thermo Scientific) and Taq DNA polymerase (Denville Scientific), respectively.

We used T4 DNA ligase (Thermo Scientific) for blunt-end ligations. If applicable, we treated

column purified (Thermo Scientific) PCR products with DpnI (Thermo Scientific) to remove

plasmid template DNA. Phosphorylation of double stranded DNA fragments was performed with

T4 Polynucleotide Kinase (Thermo Scientific). Ligase Cycling Reactions (LCR) were performed

as described previously (Kok et al., 2014). Oligonucleotides and double-stranded DNA fragments

were synthesized by Integrated DNA Technologies (IDT; Table S2).

METHOD DETAILS

Construction of *L. reuteri* mutant strains

Construction of suicide shuttle vectors for homologous recombination

To generate mutant strains in lactobacilli we used the counterselection plasmid pVPL3002

(Zhang et al., 2018). To generate *L. reuteri* R2lc*::Cm,* R2lcΔ*pks::Cm,* mlc3Δ*act* and

DSM17938Δ*act*, 500-1000 bp upstream and downstream flanking regions of target genes were

amplified by PCR (see Table S4 for oligonucleotides). We used following oligonucleotides;

oVPL3113-3114 (upstream, R2lc::*Cm*), oVPL3115-3116 (downstream, R2lc::*Cm*), oVPL3194-

3195 (upstream, mlc3Δ*act*), oVPL3196-3197 (downstream, mlc3Δ*act*), oVPL3237-oVPL3238

(upstream, DSM17938Δ*act*) and oVPL3239-oVPL3240 (downstream, DSM17938Δ*act*).

Amplicons were purified (GeneJET PCR Purification Kit, Thermo Scientific). The

pVPL3002 backbone was amplified with oVPL187-oVPL188, purified (GeneJET PCR

Purification Kit, Thermo Scientific) and digested with DpnI. Purified amplicons were quantified

(Qubit, Life Technologies). The amplicons were mixed at equimolar quantities (0.25 pmol),

followed by phosphorylation, ethanol precipitation and LCR (Kok et al., 2014). Following LCR,

DNA was precipitated with Pellet Paint co-precipitant (VWR International), resuspended in 5 µl

sterile water and transformed into electrocompetent *E. coli* EC1000 cells. By PCR, we screened

for insertion of our target sequences using oligonucleotides that flank the multiple cloning site

(oVPL49-oVPL97). Finally, the integrity of deletion and insertion cassettes was determined by

Sanger sequencing. The resultant constructs were named as follow; VPL31130 (contains the

chloramphenicol insertion cassette for R2lc::*Cm* and R2lcΔ*pks::Cm*), and VPL31079 (contains *act*

deletion cassette for mlc3).

Construction of *L. reuteri* mutants by homologous recombination

Three micrograms plasmid DNA was electroporated in electrocompetent *L. reuteri* cells.

For R2lc::*Cm,* bacterial cells were plated on MRS agar containing 5 µg/ml chloramphenicol,

colonies were screened by PCR (oVPLl3117-3118) to confirm double crossover event. For

mlc3Δ*act,* bacterial cells were plated on MRS agar containing 5 µg/ml erythromycin and colonies

were screened for single crossover (SCO) event by PCR with oligonucleotide mixtures oVPL3216-

oVPL3217-oVPL97 (upstream SCO) and oVPL3216-oVPL3217-oVPL49 (downstream SCO).

Following confirmation of SCO, bacterial cells were cultured for one passage in MRS broth

without antibiotic selection, and cells were plated on MRS agar plates containing 400 µg/ml

vancomycin. Vancomycin-resistant colonies represent cells in which a second homologous

recombination event took place (Zhang at al. 2018). To confirm double crossover (DCO), we

performed PCR using oligonucleotides oVPL3216-oVPL3217 (for mlc3Δ*act*) and oVPL3244-

oVPL3245 (for DSM17938Δ*act*). We used Sanger sequencing to verify the integrity of the

recombinant strains.

Inactivation of *L. reuteri* 6475 acyltransferase by recombineering

To inactivate the gene putatively encoding acyltransferase (LAR_1287) in *L. reuteri* 6475,

we applied single-stranded DNA recombineering using previously established procedure (Van

Pijkeren et al., 2012). An 80-mer oligonucleotide identical to the lagging strand with five

consecutive mismatches yielded, when incorporated, two in-frame stop codons (Van Pijkeren and

Britton, 2012). Electrocompetent *L. reuteri* 6475 cells harboring pJP042 (VPL2047) were

transformed with 100 µg oVPL3166. To identify recombinants, we plated cells on MRS plates and

recombinant genotypes were identified by mismatch amplification mutation assay (MAMA) PCR

(Cha et al., 1992) using oligonucleotide combination oVPL3163-3164-3165. After colony

purification, the pure genotype recombinants were confirmed by MAMA-PCR and Sanger

sequencing. The resulted mutant strain was named as *L. reuteri* 6475Δ*act.* To cure pJP042, cells

were grown in plain MRS for two passages and plated on MRS agar. Then cells were inoculated

in MRS containing 5 µg/ml erythromycin to identify cells from which pJP042 was cured.

Construction of Δ*act* mutants, pSIP controls, and complemented strains

To complement *act* into *L. reuteri* 6475Δ*act* (VPL4241) and mlc3Δ*act* (VPL4308), we

amplified by PCR the *act* gene from *L. reuteri* 6475 and mlc3 (oVPL3188-3189), the EFtu

promoter from *L. reuteri* 6475 (oVPL 1447-1448) and the pNZ8048 backbone (oVPL309-310).

PCR products were quantified, purified (Fisher Scientific, GeneJET), and DpnI treated (Fisher

Scientific, FastDigest). Subsequenlty, amplicons were purified again and mixed at equimolar

concentrations (0.25 pmol), phosphorylated (T4 polynucleotide kinase, Thermo Scientific), and

ethanol precipitated. DNA fragments were assembled by Ligase Cycling Reactions (LCR)

(oVPL 4262-4264), followed by precipitation with Pellet Paint co-precipitant (VWR

International), resuspension in 10 μL sterile water and transformation into electrocompetent

EC1000 cells. By colony PCR inserts were identified (oVPL1447 and oVPL3188), and the

integrity of the constructs was verified by Sanger sequencing. The resulting strains were labeled

E. coli VPL31603 and VPL31601 which contain pNZ8048-*act$_{mlc3}$* and pNZ8048-*act$_{6475}$*,

respectively. Hereafter, pNZ8048-*act$_{mlc3}$* and pNZ8048-*act$_{6475}$* are referred to as pVPL31603 and

pVPL31601, respectively.

One microgram of pVPL31603 and pVPL31601 were transformed into electrocompetent

rifampicin-resistant *L. reuteri* mlc3Δ*act* (VPL4248) and *L. reuteri* 6475Δ*act* (VPL4241),

respectively. Cells were plated on MRS containing 5 μg/mL erythromycin and screened via

colony PCR (oVPL1447 and oVPL3188). The mutant strains were labeled VPL31666 (mlc3Δ*act*

+ pVPL31603) and VPL31664 (6475Δ*act* + pVPL31601). Using the above method, control

strains were generated by transforming pNZ8048 empty vector into rifampicin-resistant *L.*

reuteri mlc3, mlc3Δ*act*, 6475, and 6475Δ*act*, and were labeled VPL 31668, 31675, 31613,

31673, respectively.

Generating rifampicin resistant *L. reuteri* strains

To enumerate the bacteria co-cultured with R2lc∷*cm*, we generated mutants that naturally

acquired mutations to yield resistance to rifampicin. Briefly, *L. reuteri* strains were grown in MRS

for 16 hours and plated on MRS agar containing 25 µg/ml rifampicin followed by 24 hours of

incubation at 37°C under hypoxic condition. Rifampicin resistant colonies were purified by a

conventional streak plate method and the corresponding cultures were stored at -80°C.

Whole Genome Sequencing

Genomic DNA was isolated using the genomic DNA purification kit (Wizard, Promega),

and DNA concentrations were determined using the Qubit® (dsDNA High Sensitivity Assay Kit,

Life Technologies). Whole genome sequencing was performed at the University of Wisconsin-

Madison Biotechnology Center. Samples were prepared according to the Celero PCR Workflow

with Enzymatic Fragmentation (NuGEN). Quality and quantity of the finished libraries were

assessed using an Agilent bioanalyzer and Qubit® dsDNA HS Assay Kit, respectively. Libraries

were standardized to 2 nM. Paired end, 250 bp sequencing was performed using the Illumina

NovaSeq6000. Images were analyzed using the standard Illumina Pipeline, version 1.8.2.

Comparative Genome and Bioinformatic Analyses

Reads were assembled *de novo* using SPAdes (version 3.11.1) software. Assembled draft

genomes were uploaded to National Center of Biotechnology Information (NCBI) ((Özçam and

van Pijkeren, 2019; Zhang et al., 2020)) and the Joint Genome Institute Integrated Microbial

Genomes and Microbes (JGI-IMG) database to perform comparative genome analyses. The web-

based comparative genome database Phylogenetic Profiler from the JGI-IMG was used to identify

genes present only in *L. reuteri* strains that are resistant to R2lc (ATCC PTA6475 [named as MM4-

1A in JGI-IMG], mlc3, DSM20016 and Lr4000).

We used *L. reuteri* JCM112 acyltransferase protein sequence (LAR_1287) as a query

sequence to search the National Center for Biotechnology Information (NCBI) Nonredundant

protein sequence (nr) and JGI-IMG database to identify homologous among *L. reuteri* strains.

Partial protein sequences were excluded from the data set. Amino acid sequences of *L. reuteri act*

genes were aligned by using MUSCLE (Edgar, 2004), and we constructed the phylogenetic tree

with MEGA 7.0 software (Kumar et al. 2016).

In vitro **competition assay with *L. reuteri* strains**

L. reuteri strains were grown in MRS for 16 hours. The rifampicin resistant competition

strains were co-cultured with either R2lc::*Cm* or R2lcΔ*pks::Cm* at optical density OD_{600}: 0.05

(from each strain in co-culture) in a pre-warmed MRS broth (pH 4.0). The co-cultures were then

incubated for 24 hours at 37°C. After incubation, a serial dilution was performed and 100 µl from

appropriate dilution was plated to MRS plates containing chloramphenicol (5 µg/ml) or rifampicin

(25 µg/ml). After 24 hours of incubation, we determined the competition ratios.

In vitro **biofilm assay**

L. reuteri R2lc, R2lcΔ*pks* and 100-23 cultures were inoculated into 2 ml MRS broth at

$OD_{600}= 0.1$ (0.05 from each strain if co-culture) in 24-well plates (Fisher Scientific) and incubated

for 24 hours. The culture supernatant was removed, and bacterial biofilm was washed three

times with 1 ml PBS. The adherent cells were scraped from the well and re-suspended in 2

ml PBS. For colony enumeration, the suspended cultures were plated on MRS plates containing

chloramphenicol (5 µg/ml) for R2lc and R2lcΔ*pks* or rifampicin (25 µg/ml) for 100-23, and

incubated for 24 hours.

Split-well (filter separated) competition experiment

Two cultures (R2lc:Cm^R or R2lc Δ*pks*:Cm^R vs. 100-23, Rif^R) were separated by two

inserts with different membrane pore sizes in 6-well plates as shown in Figure 1E. Filter inserts

(0.4 or 8 µm; CellTreat) were placed in a 6-well plate to create upper and lower layers.

Overnight cultures (16 hours at 37°C in MRS) of 100-23 and either R2lc:Cm^R or R2lcΔ*pks*:Cm^R

were washed once in MRS (pH= 4.0) by centrifugation (5 min at $5,000 \times g$), resuspended in

MRS to make initial OD_{600}= 0.05. Three mL of R2lc:Cm^R (or R2lc Δ*pks*:Cm^R) and 100-23 (Rif^R)

suspensions were added to upper (R2lc:Cm^R or R2lcΔ*pks*:Cm^R) and lower wells (100-23 [Rif^R]),

respectively. Plates were incubated for 24 hours at 37°C under hypoxic conditions. The viability

of the cells (CFU/mL) was determined by the standard plate count method as described above.

Sample Preparation for Double-Layer Agar Antimicrobial Assay

Live cells from R2lc:Cm^R or R2lc Δ*pks*:Cm^R overnight cultures (16 hours) were prepared

in MRS or PBS following a wash with MRS and PBS, respectively. Bacterial culture supernatant

was collected following centrifugation (1 min at $15,000 \times g$). Heat-treated (dead) cells were

prepared following heat treatment at 70°C for 30 min. For secondary metabolite extraction from

cells, acetone (1 vol) or methanol (1/50 vol) was added to the washed cell pellets obtained from

fresh culture followed by shaking at 500 rpm at room temperature for 3 hours (Eppendorf,

Thermomixer 5382). Before drop plating, all supernatant and cell extracts were filter-sterilized

(0.22 µm, EMD Millipore).

Double-Layer Agar Antimicrobial Assay

Bacterial culture, supernatant, and cell extracts of R2lc:Cm^R or R2lc Δpks:Cm^R were

subjected to the antimicrobial assay using the double-layer agar containing 100-23 cells. Two

hundred μL of 100-23 overnight culture diluted in MRS (OD$_{600}$= 0.5) was mixed with 3 mL of

pre-warmed (50°C) 0.2% agarose in dH$_2$O followed by pouring the mixture onto the MRS-agar

plate. Solidified top-agar was dried for 15 min at room temperature. Ten μL of each sample was

dropped onto the top agar and dried until completely absorbed. All double-layer agar plates were

incubated at 37°C under a hypoxic condition (2% O$_2$, 5% CO$_2$) for 24 hours.

Construction of leptin producing 100-23 strain and leptin release experiment

Rifampicin-resistant 100-23 strain (VPL4251) was transformed with pVPL31131, the

plasmid that encodes murine leptin (Alexander et al., 2019), to yield strain VPL31598 (100-

23/leptin). Cm^R-derivatives of *L. reuteri* R2lc (VPL4231) and R2lcΔpks (VPL4209), and 100-

23/leptin were grown in MRS, the latter supplemented with 5 μg/ml erythromycin. One ml of

each bacterial culture was centrifuged (2 min 16,000 × g) and washed in an equal volume of

MRS, which was subsequently added to 40 mL pre-warmed MRS. Mono-culture experiments for

100-23/leptin and co-culture experiments for R2lc + 100-23/leptin, R2lcΔpks +100-23/leptin

were initiated with a starting inoculum of OD$_{600}$= 0.1. At 0, 2, 4, 6, 8, and 10 hour time points,

we collected samples of each culture and plated on MRS-agar supplemented with 25 μg/ml

rifampicin to determine 100-23/leptin CFU counts. In addition, we collected samples of each

culture to harvest cell-free supernatant and total cell lysate for leptin quantification. Nine

hundred microliters culture were centrifuged for 1 minute at $16,000 \times g$, filter-sterilized, and

stored at -20°C. To obtain total cell lysate, we used a bead beater (BioSpec) to lyse 1 mL culture

samples in tubes containing ~100 µL 0.1 mm zirconia/silica beads with 3 minutes total bead

beating time, performed in two 90 seconds stages with 1 min cooling on ice in between. After

bead beating, samples were spun down (1 minute at 10,000 rpm) and the supernatant was filter-

sterilized and stored at -20°C. Supernatant and cell lysate samples were thawed on ice and leptin

levels were quantified by ELISA.

***In vivo* competition experiment**

Germ-free B6 (C57BL/6J, male 12-week-old, n=4-5mice/group) mice were maintained in

sterile biocontainment cages in the gnotobiotic animal facility at the University of Wisconsin-

Madison. Mice were colonized following a single oral gavage of 200 µl *L. reuteri* cocktail in

PBS (1:1 ratio, ~2×10^8 CFU). Each group was gavaged with a mixture of either *L. reuteri*

R2lc::*Cm* + competition strain or *L. reuteri* R2lcΔ*pks::Cm* + competition strain. Twenty-four

hours following colonization, fecal samples were collected daily from individual mice to

determine the fecal CFUs. Fecal samples were homogenized and diluted in PBS followed by

plating on MRS-Cm and MRS-Rif. After 24 hours of incubation colonies were counted. At day

7, mice were sacrificed by CO_2 asphyxiation and digesta from forestomach, jejunum, cecum and

colon were collected, weighed and re-suspended with PBS (100 mg/ml) to determine the CFUs

per 100 mg content.

Ampicillin Resistance Experiment

L. reuteri 6475 + pCtl, 6475Δ*act* + pCtl, 6475Δ*act* + pNZ8048-*act$_{6475}$*, mlc3 + pCtl,

mlc3Δ*act* +pCtl, and mlc3Δ*act* + pNZ8048-*act$_{mlc3}$*, were grown in MRS supplemented with 5

µg/ml erythromycin for 16 hours. The cultures were centrifuged, washed twice in 1 mL MRS,

and sub-cultured into 10 mL pre-warmed MRS with no antibiotic. Sub-cultures grew to

OD_{600}=0.6 and subsequently sub-cultured again into 1 mL aliquots containing a range of

concentrations of ampicillin with a starting inoculum of OD_{600}=0.01. Samples were transferred to

a 96-well flat-bottom plate for 24 hour growth curve measurements using a MultiSkan Sky Plate

Reader (ThermoFisher, Waltman, MA, USA) with 37°C, kinetic loop with 5 second shaking

intervals every 30 min.

Determination of peptidoglycan O'-acylation by HPLC

Peptidoglycan (PG) was extracted following a previously reported method (Ha et al., 2016)

with some modifications. After 16 hours of incubation, the cells were harvested by centrifugation

at $5,000 \times g$ and 4 °C for 5 min, washed twice with 10 mM sodium phosphate buffer (pH 6.5) and

then resuspended in 50 ml of water (pH 5.5 to 6.0). The cell suspension was boiled in an equal

volume of 8% (w/v) sodium dodecyl sulfate (SDS, 4% w/v final concentration) in 25 mM SP

buffer (pH 6.5) for 1 hour under reflux with stirring. Samples were centrifuged ($30,000 \times g$) and

the SDS-insoluble PG was washed (5 times) with sterile double distilled H_2O to completely

remove SDS and lyophilized. Lyophilized PG was dissolved in 4 ml of 1:1 mixture of 10 mM Tris-

HCl (pH 6.5) and 10 mM NaCl and sonicated for 2 min. The PG suspension was treated with 100

µg/ml α-amylase (from *Bacillus sp.*, Sigma), 10 µg/ml DNase I (Invitrogen), 50 µg/ml RNase A

(Thermo Scientific), and 20 mM $MgSO_4$ solution and incubated overnight at 37 °C with shaking.

The PG suspension was further treated with 200 µg/ml protease (from *Streptomyces griseus*,

767 Sigma) and incubated overnight at 37 °C with shaking. Samples were then re-extracted by boiling

768 in 1% SDS for 40 min, washed, lyophilized and stored at -20 °C until use.

769  For the analysis of acetate release from PG, lyophilized PG (20 mg) was dissolved in 150

770 µl ddH$_2$O and mixed with an equal volume of either 160 mM NaOH or 160 mM sodium

771 phosphate buffer (pH 6.5) and incubated overnight at 37°C with shaking. The peptidoglycan was

772 collected by centrifugation at 15,000 × g for 20 min. The supernatant was filtered through a 0.45-

773 µm-pore-size cellulose acetate membrane and quantitation of released acetate was carried out in

774 a Dionex UltiMate 3000 HPLC equipped with an LPG-3400 quaternary pump, a WPS-3000

775 analytical autosampler, and a DAD-3000 diode array detector. The filtered supernatants or 100

776 µM of acetic acid standard solution were injected onto a 300 × 7.8 mm Rezex^{TM} ROA-Organic

777 Acid H$^+$ (8%) column and eluted isocratically with 20 mM H$_2$SO$_4$ at 0.5 ml min^{-1}. Total MurNAc

778 was liberated by acid-catalyzed hydrolysis (6 N HCl, 100°C for 1.5 hour). MurNac was then

779 determined by HPLC and quantified against a calibration curve established using standard

780 MurNac. The column was maintained and 40 °C and absorbance was monitored at 210 nm.

782 **LC/MS Sample preparation**

783  An aliquot of 1.5 ml from each of the cultures were collected in Eppendorf tubes and

784 centrifuged at 2152 × g for 5 min. The supernatants were collected and transferred to dram vials

785 and the cell pellets were incubated for 1 hour in 100 µl methanol. Next, the samples were

786 centrifuged again and the methanol extracts were added to 1 dram vials. A Gilson GX-271 liquid

787 handling system was used to subject 900 µL of the samples to automated solid phase extraction

788 (SPE). Extracts were loaded onto pre-conditioned (1 ml MeOH followed by 1 ml H$_2$O) EVOLUTE

789 ABN SPE cartridges (25 mg absorbent mass, 1 ml reservoir volume; Biotage, S4 Charlotte, NC).

Samples were subsequently washed using H_2O (1 ml) to remove media components, and eluted

with MeOH (500 μL) directly into an LC/MS-certified vial.

UHPLC/HRESI-qTOF-MS Analysis of Extracts

LC/MS data were acquired using a Bruker MaXis ESI-qTOF mass spectrometer (Bruker,

Billerica, MA) coupled with a Waters Acquity UPLC system (Waters, Milford, MA) with a PDA

detector operated by Bruker Hystar software, as previously described (Hou et al., 2012). Briefly,

a solvent system of MeOH and H_2O (containing 0.1% formic acid) was used on an RP C-18 column

(Phenomenex Kinetex 2.6μm, 2.1 × 100 mm; Phenomenex, Torrance, CA) at a flow rate of 0.3

ml/min. The chromatogram method started with a linear gradient from MeOH/ H_2O (10%/90%)

to MeOH/ H_2O (97%/3%) in 12 min, then held for 2 min at MeOH/ H_2O (97%/3%). Full scan mass

spectra (m/z-150-1550) were measured in positive Electrospray Ionization (ESI) mode. The mass

spectrometer was operated using the following parameters: capillary, 4.5 kV; nebulizer pressure,

1.2 bar; dry gas flow, 8.0 L/min; dry gas temperature, 205 °C; scan rate, 2 Hz. Tune mix (ESI-L

low concentration; Agilent, Santa Clara, CA) was introduced through a divert valve at the end of

each chromatographic run for automated internal calibration. Bruker Data Analysis 4.2 software

was used for analysis of chromatograms.

808
809

SUPPLEMENTARY MATERIALS

A Streptococcus mutans mutanobactin A
(PKS/NRPS)

B Escherichia coli colibactin
(PKS/NRPS)

C Bacteroides vulgatus commendamide
(N-acyl-amide)

D Bacteroides fragilis α-galactosylceramide
(sphingolipid)

E Limosilactobaillus reuteri reutericyclin
(tetramic acid)

F Escherichia coli APE_{Ec}
(aryl polyene)

G Streptococcus agalactiae granadaene
(ornithine (rhamno)polyene)

Figure S1. Structures of bioactive fatty acid-like, polyketide and NRPS/PKS hybrid

compounds in bacteria from the gastrointestinal tract. So far, there are only seven microbial

metabolites from GI tract have been chemically identified. These are an antifungal PKS/NRPS

hybrid metabolite, Mutanobactin A produced by *Steptococcus mutans,* a genotoxin PKS/NRPS

hybrid metabolite, Colibactin produced by *Escherichia coli*; a G-Protein Coupled Receptors

(GPCR) agonist N-acyl-amide metabolite, Mutanobactin A, produced by *Bacteroides vulgatus*;

an invariant Natural Killer Cell (iNTK) activator sphingolipid metabolite, α-galactosylceramide,

819 produced by *Bacteroides fragilis*; an antimicrobial tetramic acid metabolite, Reutericyclin,

820 produced by *Limosilactobacillus reuteri*; an antioxidant aryl polyene molecule, APEEC,

821 produced by *Escherichia coli*; a polyene ornithine compound, Granadaene, produced by

822 *Streptococcus agalactiae*. Related to Introduction.

825 **Figure S2. A polyketide synthase cluster in *L. reuteri* R2lc inhibits biofim formation of *L.**

826 *reuteri* **100-23. A)** R2lc and R2lcΔ*pks* have a similar growth dynamic, and 100-23 grows slightly

827 faster compared to R2lc and R2lcΔ*pks* in mono-culture. **B)** Quantification of the number of CFU

828 in single and **C)** co-culture biofilm. For box and whisker plots, the whiskers represent the

829 maximum and minimal values, and the lower, middle and upper line of the box represent first

830 quartile, median and third quartile, respectively. Circles represent individual data points. All data

831 represent an average of at least three independent experiments. Statistical significance was

832 determined by Wilcoxon / Kruskal-Wallis Tests p<0.05 considered as significant. p=P value.

833 Related to Figure 1.

Figure S3. R2lc but not *pks* mutant produces unique compounds. The mass spectrum of the

peak between 10.52 min –10.58 min in the R2lc chromatogram at 12 hours clearly showed an ion

with m/zvalue of 257.1172 as the [M+H]+and 279.0092 as the corresponding [M+Na]+ which

were absent in the corresponding 12-hour culture of Δpks. Bruker Smart Formula algorithm,

which uses both the exact mass of the molecular ion and the isotopic pattern allowed accurate

determination of the molecular formula as $C_{16}H_{16}O_3$. Related to Figure 1.

Figure S4. *Pks*-mediated killing mechanism requires Pks-producing strain to be alive.

Double-layer agar diffusion assay with R2lc (top panel) or R2lcΔ*pks* (bottom panel) a) cell-free

supernatants; b) concentrated methanol (which concentrates the pigment (Özçam et al., 2019)

and acetone extracts; c) live cells in MRS and PBS; d) cells killed by treatment of heat or

acetone. Related to Figure 1.

Figure S5. Growth dynamics of wild-type, *act* mutant and complemented strains of *L. reuteri*

6475 and mlc3 in MRS and MRS supplanted with lysozyme and ampicillin. **A)** Deletion and

complementation of the *act* gene reduces the growth of *L. reuteri* 6475, while complementation

but not deletion of the *act* gene increases the growth of *L. reuteri* mlc3. **B)** Deletion of the *act*

gene in *L. reuteri* 6475 and mlc3 increases the lysozyme sensitivity, while the complemented

855   strains survived at levels comparable to that of the wild-type strains. **C)** the *act* mutants of 6475

856   and **D)** mlc3 are more sensitive to lysozyme, and complemented strains had a phenotype similar

857   to that of wild-type strains. Related to Figure 6.

861   **Table S1**. Strains used for comparative genome analyses.

Strain Number	Strain	Sensitive/Resistant*
1	CR	S
2	100-23	S
3	L1604-1	S
4	ATCC53608	S
5	3c6	S
6	L1600-1	S
7	one-one	S
8	rat19	S
9	JCM1081	S
10	I5007	S
11	6799jm-1	S
12	lpuph	S
13	CF48-3A1	S
14	1366	S
15	N2J	S
16	LR4020	S
17	SD2112	S
18	AD23	S
19	N4I	S
20	100-93	S
21	PTA6475	R
22	DSM20016^T	R
23	mlc3	R
24	Lr4000	R

862   *Sensitive/resistant phenotypes were determined based on statistical analyses shown in
863   Figure 4. S: sensitive, R: Resistant. Related to Figure 4.
864
865
866   **Table S2.** Unique genes in R2lc-resistat 6475, DSM20016^T and mlc3.

Locus Tag	Gene Name	Length (aa)
Lreu_0435	hypothetical protein	69
Lreu_0848	holin, Cph1 family	149
Lreu_1143	transcription regulator, XRE family	67
Lreu_1353	hypothetical protein	522
Lreu_1362	glycosyltransferase	324
Lreu_1368	acyltransferase	355
Lreu_1869	hypothetical protein	175

867   aa: amino acid. Related to Figure 4.

869 **Table S3.** Bacterial strains used in this study.

Genus and Species	Strain	VPL identifier	Description/ Origin	Source
Escherichia coli	EC1000	VPL1009	Cloning host	(Leenhouts et al., 1996)
Lactococcus lactis ssp. cremoris	MG1363	VPL2005	*L. lactis* MG1363 harboring pSIP411	Laboratory stock
Limosilactobaillus reuteri	R2lc	VPL1053	Rat	Siv Ahrné—JGI 2716884882
Limosilactobaillus reuteri	R2lc::*Cm*	VPL4231	Insertion of *Cm* gene in R2lc chromosome	This study
Limosilactobaillus reuteri	R2lcΔ*fun::Cm*	VPL4234	Insertion of *Cm* gene in R2lcΔ*fun* chromosome	This study
Limosilactobaillus reuteri	R2lcΔ*pks::Cm*	VPL4209	Insertion of *Cm* gene in R2lcΔ*pks* chromosome	(Özçam et al., 2019)
Limosilactobaillus reuteri	6475 (pJP042)	VPL2047	6475 harboring pJP042 plasmid	(Van Pijkeren and Britton, 2012)
Limosilactobaillus reuteri	6475 + pNZ8048	VPL31613	6475—Rif^R + pNZ8048	This study
Limosilactobaillus reuteri	6475Δ*act*	VPL4241	Deletion of act gene in 6475	This study
Limosilactobaillus reuteri	6475Δ*act* + pNZ8048	VPL31673	6475Δ*act*—Rif^R + pNZ8048	This study
Limosilactobaillus reuteri	6475Δ*act*—Rif^R (pSIP411)	VPL31153	6475Δ*act*—Rif^R harboring pSIP411 plasmid	This study
Limosilactobaillus reuteri	6475Δ*act::act*—Rif^R	VPL4250	6475Δ*act*—Rif^R harboring pSIP-*act* plasmid	This study
Limosilactobaillus reuteri	6475Δ*act::act*—Rif^R	VPL31664	6475 Δ*act* + pJP31601	This study
Limosilactobaillus reuteri	6475/leptin	VPL31313	6475 producing leptin	(Alexander et al., 2019)
Limosilactobaillus reuteri	100-23	VPL1049	Rat	JGI: 2500069000
Limosilactobaillus reuteri	100-23—Rif^R	VPL4251	100-23 Rif resistant	This study
Limosilactobaillus reuteri	100-23/leptin	VPL315998	100-23—Rif^R + pVPL31313	This study
Limosilactobaillus reuteri	100-23—Rif^R (pSIP411)	VPL31155	100-23—Rif^R harboring pSIP411 plasmid	This study
Limosilactobaillus reuteri	100-23::*act*—Rif^R	VPL4249	Deletion of act gene in 100-23, Rif resistant	This study
Limosilactobaillus reuteri	6475	VPL1014	Human	BioGaia AB
Limosilactobaillus reuteri	6475—Rif^R	VPL4296	6475, Rif resistant	This study
Limosilactobaillus reuteri	SD2112	VPL1013	Human	BioGaia AB
Limosilactobaillus reuteri	SD2112—Rif^R	VPL4297	SD2112, Rif resistant	This study
Limosilactobaillus reuteri	CSB7	VPL1168	Chicken	Jens Walter
Limosilactobaillus reuteri	CSB7—Rif^R	VPL4298	CSB7, Rif resistant	This study
Limosilactobaillus reuteri	KYE26	VPL1162	Chicken	Jens Walter
Limosilactobaillus reuteri	KYE26—Rif^R	VPL4299	KYE26, Rif resistant	This study
Limosilactobaillus reuteri	LK159	VPL1156	Chicken	Jens Walter
Limosilactobaillus reuteri	LK159—Rif^R	VPL4301	LK159, Rif resistant	This study
Limosilactobaillus reuteri	L4	VPL1173	Chicken	Jens Walter

Limosilactobaillus reuteri	L4—Rif^R	VPL4302	L4, Rif resistant	This study
Limosilactobaillus reuteri	mouse2	VPL1070	Mouse	(Frese et al., 2011)
Limosilactobaillus reuteri	Mouse2—Rif^R	VPL4252	Mouse2, Rif resistant	This study
Limosilactobaillus reuteri	N2J	VPL1052	Rat	(Frese *et al*. 2011)
Limosilactobaillus reuteri	N2J—Rif^R	VPL4253	N2J, Rif resistant	This study
Limosilactobaillus reuteri	one-one	VPL1060	Mouse	(Frese *et al*. 2011)
Limosilactobaillus reuteri	one-one—Rif^R	VPL4254	one-one, Rif resistant	This study
Limosilactobaillus reuteri	ATCC 53608	VPL1090	Pig	BioGaia AB EMBL LN906634
Limosilactobaillus reuteri	ATCC 53608—Rif^R	VPL4255	ATCC 53608, Rif resistant	This study
Limosilactobaillus reuteri	L1600-1	VPL1064	Mouse	(Frese *et al*. 2011)
Limosilactobaillus reuteri	L1600-1—Rif^R	VPL4256	L1600-1, Rif resistant	This study
Limosilactobaillus reuteri	N2D	VPL1067	Rat	Siv Ahrné
Limosilactobaillus reuteri	N2D—Rif^R	VPL4257	N2D, Rif resistant	This study
Limosilactobaillus reuteri	DSM17938	VPL4171	Human	BioGaia AB
Limosilactobaillus reuteri	DSM17938—Rif^R	VPL4258	DSM17938, Rif resistant	This study
Limosilactobaillus reuteri	DSM17938Δ*act*	VPL4313	Deletion of *act* gene in DSM17938	This study
Limosilactobaillus reuteri	DSM17938Δ*act*—Rif^R	VPL4315	DSM17938Δ*act*, Rif resistant	This study
Limosilactobaillus reuteri	Lr4000	VPL1071	Mouse	BioGaia AB
Limosilactobaillus reuteri	Lr4000—Rif^R	VPL4259	Lr4000, Rif resistant	This study
Limosilactobaillus reuteri	CF48-3A1	VPL1086	Human	BioGaia AB—JGI2502171173
Limosilactobaillus reuteri	CF48-3A1—Rif^R	VPL4260	CF48-3A1, Rif resistant	This study
Limosilactobaillus reuteri	L1604-1	VPL1066	Mouse	(Frese *et al*. 2011)
Limosilactobaillus reuteri	L1604-1—Rif^R	VPL4261	L1604-1, Rif resistant	This study
Limosilactobaillus reuteri	CR	VPL1059	Rat	(Frese *et al*. 2011)
Limosilactobaillus reuteri	CR—Rif^R	VPL4262	CR, Rif resistant	This study
Limosilactobaillus reuteri	mlc3	VPL1050	Mouse	JGI: 2506381016
Limosilactobaillus reuteri	mlc3—Rif^R	VPL4263	mlc3, Rif resistant	This study
Limosilactobaillus reuteri	mlc3 + pNZ8048	Vpl31668	VPL4263 + pNZ8048	This study
Limosilactobaillus reuteri	mlc3Δ*act*	VPL4246	Deletion of *act* gene in mlc3	This study
Limosilactobaillus reuteri	mlc3Δ*act*—Rif^R	VPL4308	mlc3Δ*act*, Rif resistant	This study
Limosilactobaillus reuteri	mlc3Δ*act* + pNZ8048	VPL31675	VPL4308+ pNZ8048	This study
Limosilactobaillus reuteri	mlc3Δ*act*—Rif^R (pSIP411)	VPL31177	mlc3Δ*act*—Rif^R, harboring pSIP411 plasmid	This study
Limosilactobaillus reuteri	mlc3Δ*act::act*—Rif^R	VPL4248	mlc3Δ*act*—Rif^R harboring pSIP-*act* plasmid	This study
Limosilactobaillus reuteri	mlc3Δ*act::act*—Rif^R	VPL31666	mlc3Δ*act* + pVPL31603	This study
Limosilactobaillus reuteri	I5007	VPL1082	Pig	JGI: 2554235423
Limosilactobaillus reuteri	I5007—Rif^R	VPL4264	I5007, Rif resistant	This study
Limosilactobaillus reuteri	Lr4020	VPL1072	Mouse	(Frese *et al*. 2011)
Limosilactobaillus reuteri	Lr4020—Rif^R	VPL4265	Lr4020, Rif resistant	This study
Limosilactobaillus reuteri	Lp167-67	VPL1085	Pig	BioGaia AB JGI 2599185361
Limosilactobaillus reuteri	Lp167-67—Rif^R	VPL4266	Lp167-67, Rif resistant	This study
Limosilactobaillus reuteri	N4I	VPL1063	Rat	(Frese *et al*. 2011)

Limosilactobaillus reuteri	N4I—Rif[R]	VPL4267	N4I, Rif resistant	This study
Limosilactobaillus reuteri	3c6	VPL1083	Pig	JGI: 2599185333
Limosilactobaillus reuteri	3c6—Rif[R]	VPL4268	3c6, Rif resistant	This study
Limosilactobaillus reuteri	FUA3043	VPL1062	Rat	(Frese *et al*. 2011)
Limosilactobaillus reuteri	FUA3043—Rif[R]	VPL4269	FUA3043, Rif resistant	This study
Limosilactobaillus reuteri	Rat 19	VPL1069	Rat	(Frese *et al*. 2011)
Limosilactobaillus reuteri	Rat 19—Rif[R]	VPL4270	Rat 19, Rif resistant	This study
Limosilactobaillus reuteri	6799jm-1	VPL1051	Mouse	(Frese *et al*. 2011)
Limosilactobaillus reuteri	6799jm-1—Rif[R]	VPL4271	6799jm-1, Rif resistant	This study
Limosilactobaillus reuteri	100-93	VPL1047	Mouse	(Frese *et al*. 2011)
Limosilactobaillus reuteri	100-93—Rif[R]	VPL4272	100-93, Rif resistant	This study
Limosilactobaillus reuteri	Lpuph-1	VPL1056	Mouse	JGI: 2506381017
Limosilactobaillus reuteri	Lpuph-1—Rif[R]	VPL4273	Lpuph-1, Rif resistant	This study
Limosilactobaillus reuteri	DSM 20016	VPL1046	Human	JGI: 640427118
Limosilactobaillus reuteri	DSM 20016—Rif[R]	VPL4274	DSM 20016, Rif resistant	This study
Limosilactobaillus reuteri	AD 23	VPL1048	Rat	(Frese *et al*. 2011)
Limosilactobaillus reuteri	AD 23—Rif[R]	VPL4275	AD 23, Rif resistant	This study
Limosilactobaillus reuteri	173/4	VPL1135	Pig	Jens Walter
Limosilactobaillus reuteri	173/4—Rif[R]	VPL4276	173/4, Rif resistant	This study
Limosilactobaillus reuteri	4S17	VPL1122	Pig	(Frese *et al*. 2011)
Limosilactobaillus reuteri	4S17—Rif[R]	VPL4277	4S17, Rif resistant	This study
Limosilactobaillus reuteri	1366	VPL1115	Chicken	Jens Walter
Limosilactobaillus reuteri	1366—Rif[R]	VPL4278	1366, Rif resistant	This study
Limosilactobaillus reuteri	JW2015	VPL1126	Pig	Jens Walter
Limosilactobaillus reuteri	JW2015—Rif[R]	VPL4279	JW2015, Rif resistant	This study
Limosilactobaillus reuteri	32	VPL1137	Pig	Jens Walter
Limosilactobaillus reuteri	32—Rif[R]	VPL4280	32, Rif resistant	This study
Limosilactobaillus reuteri	3S3	VPL1120	Pig	Jens Walter
Limosilactobaillus reuteri	3S3—Rif[R]	VPL4281	3S3, Rif resistant	This study
Limosilactobaillus reuteri	13S14	VPL1118	Pig	Jens Walter
Limosilactobaillus reuteri	13S14—Rif[R]	VPL4282	13S14, Rif resistant	This study
Limosilactobaillus reuteri	LK150	VPL1112	Chicken	Jens Walter
Limosilactobaillus reuteri	LK150—Rif[R]	VPL4283	LK150, Rif resistant	This study
Limosilactobaillus reuteri	JCM 1081	VPL1110	Chicken	Jens Walter
Limosilactobaillus reuteri	JCM 1081—Rif[R]	VPL4284	JCM 1081, Rif resistant	This study
Limosilactobaillus reuteri	HW8	VPL1166	Chicken	Jens Walter
Limosilactobaillus reuteri	HW8—Rif[R]	VPL4300	HW8, Rif resistant	This study
Limosilactobaillus reuteri	JW2019	VPL1129	Pig	Jens Walter
Limosilactobaillus reuteri	JW2019—Rif[R]	VPL4286	JW2019, Rif resistant	This study
Limosilactobaillus reuteri	1133 (146/2)	VPL1133	Pig	Jens Walter
Limosilactobaillus reuteri	1133 (146/2)—Rif[R]	VPL4287	1133, Rif resistant	This study
Limosilactobaillus reuteri	CP415	VPL1146	Pig	Jens Walter
Limosilactobaillus reuteri	CP415—Rif[R]	VPL4288	CP415, Rif resistant	This study
Limosilactobaillus reuteri	1063	VPL1143	Pig	Jens Walter
Limosilactobaillus reuteri	1063—Rif[R]	VPL4289	1063, Rif resistant	This study
Limosilactobaillus reuteri	1068	VPL1142	Pig	Jens Walter
Limosilactobaillus reuteri	1068—Rif[R]	VPL4291	1068, Rif resistant	This study
Limosilactobaillus reuteri	1048	VPL1141	Pig	Jens Walter
Limosilactobaillus reuteri	1048—Rif[R]	VPL4292	1048, Rif resistant	This study
Limosilactobaillus reuteri	1013	VPL1140	Pig	Jens Walter
Limosilactobaillus reuteri	1013—Rif[R]	VPL4293	1013, Rif resistant	This study

Limosilactobaillus reuteri	1704	VPL1139	Pig	Jens Walter
Limosilactobaillus reuteri	1704—Rif[R]	VPL4294	1704, Rif resistant	This study
Limosilactobaillus reuteri	FUA3048	VPL1063	Rat	(Frese *et al.* 2011)
Limosilactobaillus reuteri	FUA3048 —Rif[R]	VPL4295	FUA3048, Rif resistant	This study

871 VPL: Van Pijkeren Laboratory strain identification number. pVPL: Van Pijkeren Laboratory

872 plasmid identification number. Rif[R]: rifampicin-resistant; *act*: acyltransferase; JGI: Joint Genome

873 Institute; Cm: Chloramphenicol; Rif: Rifampicin.

877 **Table S4.** Oligonucleotides used in this study.

Oligonucleotide	Sequence (5'-3')	Target/Comment
oVPL49	acaatttcacacaggaaacagc	F. Insert screening of pVPL3002.
oVPL97	cccccattaagtgccgagtgc	R. Insert screening of pVPL3002.
oVPL187	taccgagctcgaattcactgg	R. amplifies pVPL3002 backbone.
oVPL188	atcctctagagtcgacctgc	F. amplifies pVPL3002 backbone.
oVPL3113	acgcacgacaggaagaatttg	F. amplifies u/s flanking region of CmR[+] insertion cassette.
oVPL3114	agactcgagccttgtggc	R. amplifies u/s flanking region of CmR[+] insertion cassette
oVPL3115	aagtacgaacgataatcagccc	F. amplifies d/s flanking region of CmR[+] insertion cassette.
oVPL3116	agccagtattatgacgggtc	R. amplifies d/s flanking region of CmR[+] insertion cassette
oVPL2856	agtgtcatggcgcattaacg	F. amplifies CmR[+] gene.
oVPL2857	ttataaaagccagtcattaggcc	R. amplifies CmR[+] gene
oVPL3166	tattaatgttattactagcaaacgaattaataaaatcctattaacta ggattaggatgcaacgatggaccaggataagca	*L. reuteri* 6475 *act* gene recombineering oligo.
oVPL3163	tgaacaaattcaggccaattatctgg	F. screening *act* gene
oVPL3164	tgtagttgtttggcgtcaggc	R. screening *act* gene
oVPL3165	cgttgcatcctaatcctagttaatag	MAMA oligonucleotide for *act* gene
oVPL659	tgccccgttagttgaagaag	F. amplifies pSIP411 backbone.
oVPL660	attctgctcccgcccttatg	R. amplifies pSIP411 backbone.
oVPL3188	ttaaggaattatcatcctaacaaatttattc	F. amplifies *act* gene in *L. reuteri* 6475.
oVPL3189	atgcttaatagtaagagactccac	R. amplifies *act* gene in *L. reuteri* 6475.
oVPL3194	acgatattgaatctttgcgactcc	F. amplifies u/s flanking region of *act* gene in *L. reuteri* mlc3.
oVPL3195	tagtggagtctcttactattaag	R. amplifies u/s flanking region of *act* gene in *L. reuteri* mlc3.
oVPL3196	aggaatattgtctggagtaagc	F. amplifies d/s flanking region of *act* gene in *L. reuteri* mlc3.
oVPL3197	ccattgtgactgtacatagtt	R. amplifies d/s flanking region of *act* gene in *L. reuteri* mlc3.
oVPL3198	aaacgacggccagtgaattcgagctcggtaacgatattgaatc tttgcgactccattatg	LCR bridging oligonucleotide to ligate plasmid backbone + u/s flanking region of mlc3Δ*act* deletion cassette.
oVPL3199	aattatgcttaatagtaagagactccactaaggaatattgtctgg agtaagcttaggaat	LCR bridging oligonucleotide to ligate u/s + d/s flanking region of mlc3Δ*act* deletion cassette.
oVPL3200	gtagataataactatgtacagtcacaatggatcctctagagtcg acctgcaggcatgcaa	LCR bridging oligonucleotide to ligate d/s + plasmid backbone of mlc3Δ*act* deletion cassette.
oVPL3216	atcaccaccgaagagatacc	F. *L. reuteri* mlc3Δ*act* deletion cassette screening oligonucleotide.
oVPL3217	agttgggtggataagtatgac	R. *L. reuteri* mlc3Δ*act* deletion cassette screening oligonucleotide.

oVPL3220	ataaaatactattacaaggagattttagccatgcttaatagtaag agactccactacata	Bridging oligonucleotide to ligate plasmid backbone + *act* gene for *act* gene overexpression cassettes.
oVPL3221	aataaatttgttaggatgataattccttaagaattcggtaccccg ggttcgaaggcgcca	Bridging oligonucleotide to ligate plasmid backbone + *act* gene for *act* gene overexpression cassettes.
oVPL3237	acgtgtctcctatgttaataccg	F. amplifies u/s flanking region of *act* gene in *L. reuteri* DSM17938.
oVPL3238	tataccaaattgtgctggat	R. amplifies u/s flanking region of *act* gene in *L. reuteri* DSM17938.
oVPL3239	aattattcgtataatagttgatcc	F. amplifies d/s flanking region of *act* gene in *L. reuteri* DSM17938.
oVPL3240	ccaactcacttgacccggta	R. amplifies d/s flanking region of *act* gene in *L. reuteri* DSM17938.
oVPL3241	aaacgacggccagtgaattcgagctcggtaacgtgtctcctat gttaataccggggaaaa	LCR bridging oligonucleotide to ligate plasmid backbone + u/s flanking region of DSM17938Δ*act* deletion cassette.
oVPL3242	actgtaactaatccagcacaatttggtataaattattcgtataata gttgatcctcttgt	LCR bridging oligonucleotide to ligate u/s + d/s flanking region of DSM17938Δ*act* deletion cassette.
oVPL3243	gttttcttttaccgggtcaagtgagttggatcctctagagtcgac ctgcaggcatgcaa	LCR bridging oligonucleotide to ligate d/s + plasmid backbone of DSM17938Δ*act* deletion cassette.
oVPL3244	gaaattgttcgtcgctatgga	F. *L. reuteri* DSM17938Δ*act* deletion cassette screening oligonucleotide.
oVPL3245	tcccagatttctggccaac	R. *L. reuteri* DSM17938Δ*act* deletion cassette screening oligonucleotide.
oVPL309	tctcgctttgattgttctatcgaaag	F. amplifies backbone pNZ, oligos starting d/s of IR region.
oVPL310	ataaggaagataaatcccataagggc	R. amplifies backbone pNZ, oligos starting d/s of IR region.
oVPL1447	cgaattaatagaaaaacattagtcaaatac	F. amplifies EF-tu promoter from plasmid pJG001
oVPL1448	taatgaaaacctcctgataaatttacaag	R. amplifies EF-tu promoter from plasmid pJG001
oVPL4262	ttcgctttcgatagaacaatcaaagcgagacgaattaatagaaa aacattagtcaaatac	bridging oligonucleotide from eftu promoter to act gene of 6475 or mlc3
oVPL4264	aataaatttgttaggatgataattccttaaataaggaagataaatc ccataagggcggga	bridging oligonucleotide from act gene of 6475 or mlc3 to pnz8048 backbone

878 oVPL: van Pijkeren Laboratory oligonucleotide identification number. F: Forward, R: Reverse,
879 u/s: upstream, d/s: downstream; LCR: Ligase Cycling Reaction.
880
881 **Table S5**. Plasmids used in this study.

Plasmid	Characteristic	Source
pSIP411	EmR, Sakacin-P based expression vector	(Sørvig et al., 2005)
pVPL3002	pORI19 harboring *L. reuteri* derived ddlF258Y	(Zhang et al., 2018)
pVPL31150	pSIP411 harboring *L. reuteri* 6475 *act* gene	This Study
pVPL2042	EmR, pNZ8048 derivative. Cm marker was replaced by Em marker	(Zhang et al., 2018)
pVPL31130	EmR, derivative of vector pVPL3002 in which the *L. reuteri* R2lc*::Cm* and R2lcΔ*pks::Cm* insertion cassette was cloned in the MCS site.	This Study
pVPL31079	EmR, derivative of vector pVPL3002 in which the *L. reuteri* mlc3Δ*act* deletion cassette was cloned in the MCS site.	This Study
pVPL31139	EmR, derivative of vector pVPL3002 in which the *L. reuteri* DSM17938Δ*act* deletion cassette was cloned in the MCS site.	This Study
pVPL31601	EmR, derivative of vector pVPL3002 in which the *L. reuteri* mlc3 *act* gene was cloned in the MCS site.	This Study
pVPL31603	EmR, derivative of vector pVPL3002 in which the *L. reuteri* ATCC6475 *act* gene was cloned in the MCS site.	This Study

882 pVPL: Van Pijkeren Laboratory plasmid identification number. EmR: Erythromycin-resistant;
883 CmR: Chloramphenicol-resistant. Related to STAR Methods.

884 **Table S6.** Acyltransferase gene in *L. reuteri* strains

Host	Host	Strain	Lineage	ACT_genes
Rodent	Mouse (Mus musculus)	480_44	I	No hits found
Rodent	Mouse (Mus musculus)	482_46	I	No hits found
Rodent	Mouse (Mus musculus)	484_39	I	No hits found
Rodent	Mouse	I49	I	No hits found
Rodent	Rat	L106	I	Identities = 102/377 (27%)
Rodent	Rat	L107	I	Identities = 102/377 (27%)
Rodent	Rattus norvegicus	L109	I	Identities = 191/354 (54%)
Rodent	Rattus norvegicus	L110	I	Identities = 191/354 (54%)
Rodent	Rattus norvegicus	L111	I	Identities = 191/354 (54%)
Avian	Cape teal	L5	I	No hits found
Avian	Cape teal	L6	I	No hits found
Rodent	Mouse	lpuph1	I	Identities = 63/246 (26%)
Rodent	Mouse (Mus musculus)	LR0	I	No hits found
Avian	Bird (Gallus gallus)	P43	I	No hits found
Rodent	Rat	TD1	I	Identities = 102/377 (27%)
Human	Human	DSM 20016	II	Identities = 352/355 (99%)
Human	Human (Homo sapiens)	IRT	II	Identities = 352/355 (99%)
Human	Human	JCM1112	II	Identities = 352/355 (99%)
Human	Amaru indians	L26	II	Identities = 193/354 (55%)
Human	Amaru indians	L27	II	Identities = 193/354 (55%)
Human	Human	L28	II	Identities = 352/355 (99%)
Human	Human	L29	II	Identities = 352/355 (99%)
Human	Human	L30	II	Identities = 351/355 (99%)
Human	Human	L31	II	Identities = 351/355 (99%)
Human	Human	L32	II	Identities = 352/355 (99%)
Human	Human	L33	II	Identities = 352/355 (99%)
Human	Human	L37	II	Identities = 352/355 (99%)
Human	Human (Homo sapiens)	MM2-3	II	Identities = 352/355 (99%)
Human	Human (Homo sapiens)	MM4-1a	II	Identities = 352/355 (99%)
Rodent	Rat	100-23	III	No hits found
Rodent	Mouse	103b	III	Identities = 355/355 (100%)
Rodent	Mouse	107k	III	No hits found
Rodent	Mouse	111b	III	Identities = 355/355 (100%)
Rodent	Mouse	609d	III	No hits found
Human	Human (Homo sapiens)	DS12_10	III	No hits found
Rodent	Mouse	L118	III	No hits found
Rodent	Mouse	L119	III	No hits found
Rodent	Mouse	L120	III	Identities = 106/379 (28%)
Human	Amaru indians	L25	III	No hits found
Rodent	Marmota vancouveriensis	L92	III	Identities = 355/355 (100%)
Rodent	Sourdough	LTH2584	III	No hits found
Rodent	Sourdough	LTH5448	III	Identities = 193/354 (55%)
Rodent	Mouse	mlc3	III	Identities = 355/355 (100%)
pig	Pig	ATCC 53608	IV	Identities = 192/354 (54%)
pig	Pig	I5007	IV	Identities = 107/384 (28%)

pig	Pig	KLR1001	IV	Identities = 77/250 (31%)
pig	Pig	KLR2001	IV	Identities = 192/354 (54%)
pig	Pig	KLR3002	IV	No hits found
Primate	Agile gibbon	L123	IV	Identities = 85/321 (26%)
Primate	Lion-tailed Macaque	L125	IV	Identities = 58/152 (38%)
Primate	Orangutan	L126	IV	No hits found
Primate	Japanese macaque	L45	IV	No hits found
Primate	Japanese macaque	L46	IV	No hits found
Primate	Japanese macaque	L47	IV	No hits found
Primate	Agile gibbon	L48	IV	No hits found
Primate	Agile gibbon	L49	IV	No hits found
Primate	Agile gibbon	L50	IV	No hits found
Primate	Gorilla	L55	IV	No hits found
Primate	Sulawesi macaque	L59	IV	No hits found
Primate	Mandrill	L60	IV	No hits found
Primate	Mandrill	L61	IV	No hits found
Primate	Mandrill	L62	IV	No hits found
Primate	Monkey	L68	IV	No hits found
Primate	Monkey	L69	IV	No hits found
Primate	Orangutan	L71	IV	No hits found
Primate	Orangutan	L72	IV	No hits found
Primate	Orangutan	L73	IV	No hits found
pig	Pig	lp167-67	IV	No hits found
pig	Pig	pg-3b	IV	No hits found
pig	Pig	ZLR003	IV	No hits found
pig	Pig	20-2	V	Identities = 97/388 (25%)
pig	Pig	3c6	V	No hits found
Avian	Bird (Gallus gallus)	1366	VI	No hits found
Avian	Gallus gallus	An166	VI	Identities = 97/388 (25%)
Avian	Gallus gallus	An71	VI	Identities = 97/388 (25%)
Human	Human (Homo sapiens)	CF48-3A	VI	Identities = 111/391 (28%)
Avian	Bird (Gallus gallus)	CSF8	VI	No hits found
Human	Homo sapiens	DS17_10	VI	No hits found
Avian	Gallus gallus	JCM1081	VI	No hits found
Avian	African spoonbiil	L1	VI	Identities = 111/391 (28%)
Avian	Golden pheasant	L10	VI	Identities = 82/272 (30%)
Avian	Golden pheasant	L11	VI	Identities = 82/272 (30%)
Avian	Golden pheasant	L12	VI	Identities = 82/272 (30%)
Avian	Poultry	L17	VI	Identities = 97/388 (25%)
Avian	Quail	L19	VI	Identities = 111/391 (28%)
Avian	Argus pheasent	L2	VI	Identities = 197/354 (56%)
Avian	Quail	L20	VI	Identities = 111/391 (28%)
Avian	Quail	L21	VI	Identities = 111/391 (28%)
Avian	Turkey	L22	VI	No hits found
Avian	Turkey	L23	VI	Identities = 352/355 (99%)
Avian	Turkey	L24	VI	Identities = 111/391 (28%)
Avian	Argus pheasent	L3	VI	Identities = 352/355 (99%)
Human	Human	L34	VI	Identities = 111/391 (28%)

Human	Human	L35	VI	Identities = 111/391 (28%)
Human	Human	L36	VI	Identities = 111/391 (28%)
Avian	Argus pheasent	L4	VI	Identities = 352/355 (99%)
Human	Png	L43	VI	Identities = 153/353 (43%)
Avian	Francolin	L7	VI	Identities = 111/391 (28%)
Avian	Francolin	L8	VI	Identities = 111/391 (28%)
Avian	Francolin	L9	VI	Identities = 111/391 (28%)
Human	Human (Homo sapiens)	M27U15	VI	Identities = 111/391 (28%)
Human	Human (Homo sapiens)	MD-IIE-43	VI	No hits found
Human	Human (Homo sapiens)	MM34-4A	VI	Identities = 111/391 (28%)
Human	Human (Homo sapiens)	RC-14	VI	No hits found
Human	Human (Homo sapiens)	SD2112	VI	Identities = 111/391 (28%)
Rodent	Rat	L108	VII	No hits found
Rodent	Springhaas	L112	VII	Identities = 193/354 (55%)
Rodent	Springhaas	L113	VII	Identities = 193/354 (55%)
Rodent	Springhaas	L114	VII	Identities = 193/354 (55%)
Rodent	Porqupine	L121	VII	Identities = 252/355 (71%)
Human	Png	L38	VII	No hits found
Human	Png	L39	VII	No hits found
Human	Png	L40	VII	No hits found
Human	Png	L41	VII	No hits found
Human	Png	L42	VII	No hits found
Human	Png	L44	VII	No hits found
Primate	Black howler monkey	L51	VII	Identities = 193/354 (55%)
Primate	Lion-tailed Macaque	L56	VII	Identities = 109/384 (28%)
Primate	Lion-tailed Macaque	L57	VII	No hits found
Primate	Lion-tailed Macaque	L58	VII	No hits found
Primate	Patas monkey	L65	VII	No hits found
Primate	Monkey	L70	VII	Identities = 254/355 (72%)
Rodent	Capy C	L85	VII	No hits found
Rodent	Capy C	L86	VII	No hits found
Rodent	Coendou prehensilis	L87	VII	No hits found
Rodent	Dasyprocta leporina	L88	VII	No hits found
Rodent	Dasyprocta leporina	L89	VII	No hits found
Primate	Black howler monkey	L124	IX	Identities = 83/329 (25%)
Primate	Black howler monkey	L52	IX	Identities = 83/329 (25%)
Primate	Black howler monkey	L53	IX	Identities = 83/329 (25%)
Primate	Patas monkey	L66	IX	No hits found
Primate	Patas monkey	L67	IX	No hits found
Primate	Mandrill	L63	Close to IX	No hits found
Primate	Mandrill	L64	Close to IX	No hits found
Rodent	Apodemus agrarius	L77	Close to *L. reuteri*	Identities = 106/384 (28%)
Rodent	Apodemus agrarius	L78	Close to *L. reuteri*	Identities = 106/384 (28%)
Rodent	Apodemus agrarius	L79	Close to *L. reuteri*	Identities = 106/384 (28%)
Rodent	Apodemus agrarius	L80	Close to *L. reuteri*	Identities = 106/384 (28%)

Related to Figure 5.

References

Aleti, G., Baker, J.L., Tang, X., Alvarez, R., Dinis, M., Tran, N.C., Melnik, A. V., Zhong, C.,

Ernst, M., Dorrestein, P.C., et al. (2019). Identification of the Bacterial Biosynthetic Gene

Clusters of the Oral Microbiome Illuminates the Unexplored Social Language of Bacteria during

Health and Disease. MBio *10*.

Alexander, L.M., Oh, J.-H., Stapleton, D.S., Schueler, K.L., Keller, M.P., Attie, A.D., and

Pijkeren, J.-P. van (2019). Exploiting Prophage-Mediated Lysis for Biotherapeutic Release by

Lactobacillus reuteri. Appl. Environ. Microbiol. *85*.

Bera, A., Herbert, S., Jakob, A., Vollmer, W., and Götz, F. (2005). Why are pathogenic

staphylococci so lysozyme resistant? The peptidoglycan O-acetyltransferase OatA is the major

determinant for lysozyme resistance of *Staphylococcus aureus*. Mol. Microbiol. *55*, 778–787.

Bonder, M.J., Kurilshikov, A., Tigchelaar, E.F., Mujagic, Z., Imhann, F., Vila, A.V., Deelen, P.,

Vatanen, T., Schirmer, M., Smeekens, S.P., et al. (2016). The effect of host genetics on the gut

microbiome. Nat. Genet. *48*, 1407–1412.

Boon, E., Meehan, C.J., Whidden, C., Wong, D.H.-J., Langille, M.G.I., and Beiko, R.G. (2014).

Interactions in the microbiome: communities of organisms and communities of genes. FEMS

Microbiol. Rev. *38*, 90–118.

Cha, R.S., Zarbl, H., Keohavong, P., and Thilly, W.G. (1992). Mismatch amplification mutation

assay (MAMA): Application to the c-H-ras gene. Genome Res. *2*, 14–20.

Cimermancic, P., Medema, M.H., Claesen, J., Kurita, K., Wieland Brown, L.C., Mavrommatis,

K., Pati, A., Godfrey, P.A., Koehrsen, M., Clardy, J., et al. (2014). Insights into secondary

metabolism from a global analysis of prokaryotic biosynthetic gene clusters. Cell *158*, 412–421.

David, L.A., Maurice, C.F., Carmody, R.N., Gootenberg, D.B., Button, J.E., Wolfe, B.E., Ling,

A. V, Devlin, A.S., Varma, Y., Fischbach, M.A., et al. (2014). Diet rapidly and reproducibly

alters the human gut microbiome. Nature *505*, 559–563.

Donia, M.S., Cimermancic, P., Schulze, C.J., Wieland Brown, L.C., Martin, J., Mitreva, M.,

Clardy, J., Linington, R.G., and Fischbach, M.A. (2014). A Systematic Analysis of Biosynthetic

Gene Clusters in the Human Microbiome Reveals a Common Family of Antibiotics. Cell *158*,

1402–1414.

Duar, R.M., Frese, S.A., Lin, X.B., Fernando, S.C., Burkey, T.E., Tasseva, G., Peterson, D.A.,

Blom, J., Wenzel, C.Q., Szymanski, C.M., et al. (2017). Experimental Evaluation of Host

Adaptation of *Lactobacillus reuteri* to Different Vertebrate Species. Appl. Environ. Microbiol.

83, e00132-17.

Edgar, R.C. (2004). MUSCLE: Multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res. *32*, 1792–1797.

Edwards, K.F., Kremer, C.T., Miller, E.T., Osmond, M.M., Litchman, E., and Klausmeier, C.A.

(2018). Evolutionarily stable communities: a framework for understanding the role of trait

evolution in the maintenance of diversity. Ecol. Lett. *21*, 1853–1868.

Frese, S.A., Benson, A.K., Tannock, G.W., Loach, D.M., Kim, J., Zhang, M., Oh, P.L., Heng,

N.C.K., Patil, P.B., Juge, N., et al. (2011). The evolution of host specialization in the vertebrate

gut symbiont *Lactobacillus reuteri*. PLoS Genet. *7*.

Frese, S.A., MacKenzie, D.A., Peterson, D.A., Schmaltz, R., Fangman, T., Zhou, Y., Zhang, C.,

Benson, A.K., Cody, L.A., Mulholland, F., et al. (2013). Molecular Characterization of Host-

Specific Biofilm Formation in a Vertebrate Gut Symbiont. PLoS Genet. *9*.

Le Gac, M., Plucain, J., Hindré, T., Lenski, R.E., and Schneider, D. (2012). Ecological and

evolutionary dynamics of coexisting lineages during a long-term experiment with *Escherichia*

coli. Proc. Natl. Acad. Sci. U. S. A. *109*, 9487–9492.

Gruber, G., and Steglich, W. (2007). Calostomal, a polyene pigment from the gasteromycete

Colostoma cinnabarinum (Boletales). Zeitschrift Fur Naturforsch. - Sect. B J. Chem. Sci. *62*,

129–131.

Ha, R., Frirdich, E., Sychantha, D., Biboy, J., Taveirne, M.E., Johnson, J.G., DiRita, V.J.,

Vollmer, W., Clarke, A.J., and Gaynor, E.C. (2016). Accumulation of Peptidoglycan *O* -

Acetylation Leads to Altered Cell Wall Biochemistry and Negatively Impacts Pathogenesis

Factors of *Campylobacter jejuni*. J. Biol. Chem. *291*, 22686–22702.

Herbrík, A., Corretto, E., Chroňáková, A., Langhansová, H., Petrásková, P., Hrdý, J., Čihák, M.,

Krištůfek, V., Bobek, J., Petříček, M., et al. (2020). A Human Lung-Associated *Streptomyces* sp.

TR1341 Produces Various Secondary Metabolites Responsible for Virulence, Cytotoxicity and

Modulation of Immune Response. Front. Microbiol. *10*, 3028.

Hibbing, M.E., Fuqua, C., Parsek, M.R., and Peterson, S.B. (2010). Bacterial competition:

surviving and thriving in the microbial jungle. Nat. Rev. Microbiol. *8*, 15–25.

Hiergeist, A., Gläsner, J., Reischl, U., and Gessner, A. (2015). Analyses of Intestinal Microbiota:

Culture versus Sequencing. ILAR J. *56*, 228–240.

Hooper, L. V, Littman, D.R., Macpherson, A.J., and Program, M.P. (2015). Interactions between

the microbiota and the immune system. Science (80-.). *336*, 1268–1273.

Hou, Y., Braun, D.R., Michel, C.R., Klassen, J.L., Adnani, N., Wyche, T.P., and Bugni, T.S.

(2012). Microbial Strain Prioritization Using Metabolomics Tools for the Discovery of Natural

Products. Anal. Chem. *84*, 4277–4283.

Jacobson, A., Lam, L., Rajendram, M., Tamburini, F., Honeycutt, J., Pham, T., Van Treuren, W.,

Pruss, K., Stabler, S.R., Lugo, K., et al. (2018). A Gut Commensal-Produced Metabolite

Mediates Colonization Resistance to *Salmonella* Infection. Cell Host Microbe *24*, 296-307.e7.

Kok, S. De, Stanton, L.H., Slaby, T., Durot, M., Holmes, V.F., Patel, K.G., Platt, D., Shapland,

E.B., Serber, Z., Dean, J., et al. (2014). Rapid and reliable DNA assembly via ligase cycling

reaction. ACS Synth. Biol. *3*, 97–106.

Kumar, S., Stecher, G., Li, M., Knyaz, C., and Tamura, K. (2018). MEGA X: Molecular

Evolutionary Genetics Analysis across Computing Platforms. Mol. Biol. Evol. *35*, 1547–1549.

Lee, A.J., Cadelis, M.M., Kim, S.H., Swift, S., Copp, B.R., and Villas-Boas, S.G. (2020).

Epipyrone A, a Broad-Spectrum Antifungal Compound Produced by *Epicoccum nigrum* ICMP

19927. Molecules *25*, 5997.

Leenhouts, K., Buist, G., Bolhuis, A., Berge, A. ten, Kiel, J., Mierau, I., Dabrowska, M.,

Venema, G., and Kok, J. (1996). A general system for generating unlabelled gene replacements

in bacterial chromosomes. Mol. Gen. Genet. MGG *253*, 217–224.

Li, J., Sang, M., Jiang, Y., Wei, J., Shen, Y., Huang, Q., Li, Y., and Ni, J. (2021). Polyene-

Producing Streptomyces spp. From the Fungus-Growing Termite *Macrotermes barneyi* Exhibit

High Inhibitory Activity Against the Antagonistic Fungus *Xylaria*. Front. Microbiol. *12*, 649962.

Lin, X.B., Lohans, C.T., Duar, R., Zheng, J., Vederas, J.C., Walter, J., and Gänzle, M. (2015).

Genetic determinants of reutericyclin biosynthesis in *Lactobacillus reuteri*. Appl. Environ.

Microbiol. *81*, 2032–2041.

Lin, X.B., Wang, T., Stothard, P., Corander, J., Wang, J., Baines, J.F., Knowles, S.C.L.,

Baltrūnaitė, L., Tasseva, G., Schmaltz, R., et al. (2018). The evolution of ecological facilitation

within mixed-species biofilms in the mouse gastrointestinal tract. ISME J. 1.

Martínez, I., Maldonado-Gomez, M.X., Gomes-Neto, J.C., Kittana, H., Ding, H., Schmaltz, R.,

Joglekar, P., Cardona, R.J., Marsteller, N.L., Kembel, S.W., et al. (2018). Experimental

evaluation of the importance of colonization history in early-life gut microbiota assembly. Elife

7.

Medema, M.H., Blin, K., Cimermancic, P., de Jager, V., Zakrzewski, P., Fischbach, M.A.,

Weber, T., Takano, E., and Breitling, R. (2011). antiSMASH: rapid identification, annotation and

analysis of secondary metabolite biosynthesis gene clusters in bacterial and fungal genome

sequences. Nucleic Acids Res. *39*, W339-46.

Medema, M.H., Cimermancic, P., Sali, A., Takano, E., and Fischbach, M.A. (2014). A

Systematic Computational Analysis of Biosynthetic Gene Cluster Evolution: Lessons for

Engineering Biosynthesis. PLoS Comput. Biol. *10.*

Moynihan, P.J., and Clarke, A.J. (2010). O-acetylation of peptidoglycan in gram-negative

bacteria: identification and characterization of peptidoglycan O-acetyltransferase in *Neisseria*

gonorrhoeae. J. Biol. Chem. *285*, 13264–13273.

Murray, I.A., and Shaw, W. V. (1997). O-Acetyltransferases for chloramphenicol and other

natural products. Antimicrob. Agents Chemother. *41*, 1–6.

Oh, J.H., and Van Pijkeren, J.P. (2014). CRISPR-Cas9-assisted recombineering in *Lactobacillus*

reuteri. Nucleic Acids Res. *42.*

Oh, J.-H., Lin, X.B., Zhang, S., Tollenaar, S.L., Özçam, M., Dunphy, C., Walter, J.E., and

Pijkeren, J.-P. van (2019). Prophages in *Lactobacillus reuteri* are associated with fitness trade-

offs but can increase competitiveness in the gut ecosystem. Appl. Environ. Microbiol.

Oh, P.L., Benson, A.K., Peterson, D.A., Patil, P.B., Moriyama, E.N., Roos, S., and Walter, J.

(2009). Diversification of the gut symbiont *Lactobacillus reuteri* as a result of host-driven

evolution. ISME J. *4*, 377–387.

Özçam, M., and van Pijkeren, J.-P. (2019). Draft Genome Sequence of Aryl Hydrocarbon

Receptor Activator Strains *Lactobacillus reuteri* R2lc and 2010 . Microbiol. Resour. Announc. *8*.

Özçam, M., Tocmo, R., Oh, J.H., Afrazi, A., Mezrich, J.D., Roos, S., Claesen, J., and van

Pijkeren, J.P. (2019). Gut symbionts *Lactobacillus reuteri* R2lc and 2010 encode a polyketide

synthase cluster that activates the mammalian aryl hydrocarbon receptor. Appl. Environ.

Microbiol. *85*.

Van Pijkeren, J.P., and Britton, R.A. (2012). High efficiency recombineering in lactic acid

bacteria. Nucleic Acids Res. *40*, 1–13.

Van Pijkeren, J.-P., Neoh, K.M., Sirias, D., Findley, A.S., and Britton, R.A. (2012). Exploring

optimization parameters to increase ssDNA recombineering in *Lactococcus lactis* and

Lactobacillus reuteri. Bioengineered *3*, 209–217.

Ragland, S.A., and Criss, A.K. (2017). From bacterial killing to immune modulation: Recent

insights into the functions of lysozyme. PLoS Pathog. *13*, e1006512.

Sambrook, J., and Russell, D.W. (2006). Transformation of *E. coli* by Electroporation. Cold

Spring Harb. Protoc. *2006*, pdb.prot3933.

Sassone-Corsi, M., Nuccio, S.-P., Liu, H., Hernandez, D., Vu, C.T., Takahashi, A.A., Edwards,

R.A., and Raffatellu, M. (2016). Microcins mediate competition among Enterobacteriaceae in the

inflamed gut. Nature *540*, 280–283.

Savage, D.C., Dubos, R., and Schaedler, R.W. (1968). The gastrointestinal epithelium and its

autochthonous bacterial flora. J. Exp. Med. *127*, 67–76.

Segura Munoz, R.R., Mantz, S., Schmaltz, R.J., Walter, J., and Ramer-Tait, A.E. (2020).

Experimental evaluation of ecological principles to understand and modulate the outcome of

bacterial strain competition in gut microbiomes.

Sender, R., Fuchs, S., and Milo, R. (2016). Revised Estimates for the Number of Human and

Bacteria Cells in the Body. PLoS Biol. *14*, e1002533.

Snijders, A.M., Langley, S.A., Kim, Y.-M., Brislawn, C.J., Noecker, C., Zink, E.M., Fansler,

S.J., Casey, C.P., Miller, D.R., Huang, Y., et al. (2016). Influence of early life exposure, host

genetics and diet on the mouse gut microbiome and metabolome. Nat. Microbiol. *2*, 16221.

Sørvig, E., Mathiesen, G., Naterstad, K., Eijsink, V.G.H., and Axelsson, L. (2005). High-level,

inducible gene expression in *Lactobacillus sakei* and *Lactobacillus plantarum* using versatile

expression vectors. Microbiology *151*, 2439–2449.

Teethaisong, Y., Autarkool, N., Sirichaiwetchakoon, K., Krubphachaya, P., Kupittayanant, S.,

and Eumkeb, G. (2014). Synergistic activity and mechanism of action of *Stephania suberosa*

Forman extract and ampicillin combination against ampicillin-resistant *Staphylococcus aureus*. J.

Biomed. Sci. *21*, 1–11.

Turnbaugh, P.J., Ridaura, V.K., Faith, J.J., Rey, F.E., Knight, R., and Gordon, J.I. (2009). The

effect of diet on the human gut microbiome: a metagenomic analysis in humanized gnotobiotic

mice. Sci. Transl. Med. *1*, 6ra14.

Vaishnava, S., Yamamoto, M., Severson, K.M., Ruhn, K.A., Yu, X., Koren, O., Ley, R.,

Wakeland, E.K., and Hooper, L. V. (2011). The antibacterial lectin RegIIIγ promotes the spatial

segregation of microbiota and host in the intestine. Science (80-.). *334*, 255–258.

Vanderkelen, L., Van Herreweghe, J.M., Vanoirbeek, K.G.A., Baggerman, G., Myrnes, B.,

Declerck, P.J., Nilsen, I.W., Michiels, C.W., and Callewaert, L. (2011). Identification of a

bacterial inhibitor against g-type lysozyme. Cell. Mol. Life Sci. *68*, 1053–1064.

Veldhoen, M., Hirota, K., Westendorf, A.M., Buer, J., Dumoutier, L., Renauld, J.-C., and

Stockinger, B. (2008). The aryl hydrocarbon receptor links TH17-cell-mediated autoimmunity to

environmental toxins. Nature *453*, 106–109.

Vollmer, W. (2008). Structural variation in the glycan strands of bacterial peptidoglycan. FEMS

Microbiol. Rev. *32*, 287–306.

Walter, J., Britton, R.A., and Roos, S. (2011). Host-microbial symbiosis in the vertebrate

gastrointestinal tract and the *Lactobacillus reuteri* paradigm. Proc. Natl. Acad. Sci. *108*, 4645–

4652.

Youngblut, N.D., De La Cuesta-Zuluaga, J., Reischer, G.H., Dauser, S., Schuster, N., Walzer, C.,

Stalder, G., Farnleitner, A.H., and Ley, R.E. (2020). Large-Scale Metagenome Assembly

Reveals Novel Animal-Associated Microbial Genomes, Biosynthetic Gene Clusters, and Other

Genetic Diversity.

Zarrinpar, A., Chaix, A., Yooseph, S., and Panda, S. (2014). Diet and feeding pattern affect the

diurnal dynamics of the gut microbiome. Cell Metab. *20*, 1006–1017.

Zelante, T., Iannitti, R., Cunha, C., DeLuca, A., Giovannini, G., Pieraccini, G., Zecchi, R.,

D'Angelo, C., Massi-Benedetti, C., Fallarino, F., et al. (2013). Tryptophan catabolites from

microbiota engage aryl hydrocarbon receptor and balance mucosal reactivity via interleukin-22.

Immunity *39*, 372–385.

Zhang, S., Oh, J.-H., Alexander, L.M., Özçam, M., and Van Pijkeren, J.-P. (2018). D-Ala-D-Ala

ligase as a broad host-range counterselection marker in vancomycin-resistant lactic acid bacteria.

J. Bacteriol. JB.00607-17.

Zhang, S., Özçam, M., and van Pijkeren, J.-P. (2020). Draft Genome Sequences of 12

Lactobacillus reuteri Strains of Rodent Origin. Microbiol. Resour. Announc. *9*.

Zhao, W.T., Shi, X., Xian, P.J., Feng, Z., Yang, J., and Yang, X.L. (2021). A new fusicoccane

diterpene and a new polyene from the plant endophytic fungus *Talaromyces pinophilus* and their

antimicrobial activities. Nat. Prod. Res. *35*, 124–130.

Zheng, J., Wittouck, S., Salvetti, E., Franz, C.M.A.P., Harris, H.M.B., Mattarelli, P., O'toole,

P.W., Pot, B., Vandamme, P., Walter, J., et al. (2020). A taxonomic note on the genus

Lactobacillus: Description of 23 novel genera, emended description of the genus Lactobacillus

beijerinck 1901, and union of *Lactobacillaceae* and *Leuconostocaceae*. Int. J. Syst. Evol.

Microbiol. *70*, 2782–2858.

Zheng, Y., Valdez, P.A., Danilenko, D.M., Hu, Y., Sa, S.M., Gong, Q., Abbas, A.R., Modrusan,

Z., Ghilardi, N., de Sauvage, F.J., et al. (2008). Interleukin-22 mediates early host defense

against attaching and effacing bacterial pathogens. Nat. Med. *14*, 282–289.

1079