A quantitative gibberellin signalling biosensor reveals a role for gibberellins in internode specification at the shoot apical meristem

Bihai Shi[1*], Amelia Felipo-Benavent[2*], Guillaume Cerutti[1], Carlos Galvan-Ampudia[1], Lucas Jilli[2], Geraldine Brunoud[1], Jérome Mutterer[2], Lali Sakvarelidze-Achard[2], Jean-Michel Davière[2], Alejandro Navarro-Galiano[3], Ankit Walia[4], Shani Lazary[5], Jonathan Legrand[1], Roy Weinstein[5], Alexander M. Jones[4], Salomé Prat[3], Patrick Achard[2†], Teva Vernoux[1†]

[1] Laboratoire Reproduction et Développement des Plantes, Univ Lyon, ENS de Lyon, CNRS, INRAE, INRIA, F-69342, Lyon, France

[2] Institut de biologie moléculaire des plantes, CNRS, Université de Strasbourg, 67084 Strasbourg, France

[3] Centre for Research in Agricultural Genomics, 08193 Cerdanyola, Barcelona, Spain

[4] Sainsbury Laboratory, Cambridge University, Cambridge CB2 1LR, United Kingdom

[5] Department of Molecular Biology and Ecology of Plants, Tel Aviv University, Tel Aviv 69978, Israel

[*]These authors contributed equally

[†]To whom correspondence should be addressed: T.V. (teva.vernoux@ens-lyon.fr) or P.A. (patrick.achard@ibmp-cnrs.unistra.fr)

Abstract

Growth at the shoot apical meristem (SAM) is essential for shoot architecture construction. The phytohormones gibberellins (GA) play a pivotal role in coordinating plant growth, but their role in the SAM remains mostly unknown. Here, we developed a ratiometric GA signalling biosensor by engineering one of the DELLA repressors, to suppress its master regulatory function in GA transcriptional responses while preserving its degradation upon GA sensing. We demonstrate that this novel degradation-based biosensor accurately reports on cellular changes in GA levels and perception during development. We used this biosensor to map GA signalling activity in the SAM. We show that high GA signalling is found primarily in cells located between organ primordia that are the precursors of internodes. By gain- and loss-of-function approaches, we further demonstrate that GAs regulate cell division plane orientation to establish the typical cellular organisation of internodes, thus contributing to internode specification in the SAM.

Introduction

The shoot apical meristem (SAM) at the tip of shoot axes comprises a stem cell niche whose activity produces lateral organs and stem segments in a modular iterative fashion during the whole plant life. Each of these repetitive units or phytomere includes an internode and lateral organs at a node and an axillary meristem at the leaf axil[1]. The growth and organization of phytomeres change during development. In *Arabidopsis thaliana*, internode growth is inhibited during the vegetative phase and axillary meristems rest dormant at the axils of rosette leaves. Upon floral transition, the SAM turns into an inflorescence meristem, producing elongated internodes and axillary buds that form branches at the axils of cauline leaves, and later flowers without leaves[2]. While substantial progress has been made on our understanding of the mechanisms controlling the initiation of leaves, flowers and branches, much less is known on how internodes are initiated.

Growth via cell division and expansion is essential for reiterative organogenesis at the SAM. The tetracyclic diterpenoid hormones gibberellins (GA) are key growth regulators[3–5], with a crucial role in many embryonic and post-embryonic developmental processes[6]. Central to the GA signalling pathway are the five DELLA proteins, GIBBERELLIC ACID INSENSITIVE (GAI), REPRESSOR OF GA1-3 (RGA) and (RGL) RGA-Like 1-3[7]. These nuclear proteins are composed of an N-terminal DELLA/TVHYNP domain and of a GRAS domain. The GRAS domain allows DELLAs to interact with diverse target transcription factors and transcriptional regulators and to suppress growth by modulating their activity[7]. Binding of GA to the GIBBERELLIN INSENSITIVE DWARF1 (GID1) GA receptor promotes GID1 interaction with the N-terminal domain of DELLAs, triggering DELLA degradation by the ubiquitin-dependent proteasome pathway[7–10], and in turn, de-repressing GA responses. Despite the relatively specialised role of the GRAS and DELLA/TVHYNP domains, residues

required for DELLA degradation and partner protein interaction are widely distributed within

the protein sequence.

The identification of genes encoding GA catabolic enzymes as direct targets of the class I

KNOX meristem identity regulators has led to propose that GA levels are low in the SAM

cells, while high GA concentrations trigger growth of lateral organs[3–5]. Low GA levels could

then contribute to SAM maintenance[3]. However, more recent analysis indicate also that GAs

promote the increase in SAM size during floral transition by regulating the division and

expansion of inner SAM cells[11,12], in a similar manner as in the root[13,14]. DELLAs were

likewise found to limit meristem size by directly regulating the expression of the cell-cycle

inhibitor KRP2 in the internal part of the SAM (the rib zone)[12]. Together, these findings

support a role for GA in positively regulating cell division in the inner tissues of the SAM,

and thus SAM size. At the same time, several genes encoding GA biosynthetic and

degradation enzymes are expressed specifically in lateral organs[4,11], illustrating a complex

spatio-temporal GA distribution in the SAM to likely fulfil different functions.

Accessing spatio-temporal GA distribution has been instrumental to better understand

function of these hormones in different tissues and at various developmental stages.

Accumulation of GA in the root endodermis and regulation of their cellular level via GA

transport were discovered by using bioactive fluorescein (Fl)-tagged GA[15,16]. More recently,

the nlsGPS1 GA FRET sensor revealed that GA levels are correlated with cell elongation in

roots, stamen filaments and dark-grown hypocotyls[17]. Here, building on the knowledge on the

GA signalling pathway, we report on the engineering and characterization of a quantitative

degradation-based GA signalling biosensor. We used this novel biosensor, in conjunction

with GA-Fls and nlsGPS1, to map the spatio-temporal distribution of GA signalling activity

and to quantitatively analyse how GAs regulate cell behaviour in the SAM epidermis. We

demonstrate that GAs control the orientation of division planes of SAM cells located between

organ primordia, therefore specifying the typical cellular organization of internodes.

Results

Modifying the RGA protein for sensor construction (Fig. 1; Extended Data Fig. 1, 2)

To generate a degradation-based GA signalling biosensor, we engineered a DELLA protein to

meet two criteria, (i) a specific degradation upon GA perception; and (ii) a minimal

interference with GA signalling. To do so, we modified DELLAs to preserve the interaction

with GID1 while abolishing interactions with partner transcription factors, by introducing

mutations in the GRAS domain. Among the 5 *Arabidopsis* DELLAs, RGA displays one of the

highest GA-dependent degradation rate[18], and RGA-GFP fusions are widely used as GA

signalling reporter[8]. Leveraging on the results of GRAS domain mutant analyses in rice[19] and

Arabidopsis[20,21](Fig. 1a, Extended Data Fig. 1a), we generated four modified RGA versions

(RGA^{m1} to RGA^{m4}) and tested their ability to meet the above defined criteria (Extended Data

Fig. 1). We first used a yeast-two hybrid (Y2H) assay to test the binding capacity of these

modified candidates with three well known DELLA interacting partners, JAZ1, TCP14 and

IDD2[22]. While RGA^{m1} had a minor effect on interactions, RGA^{m2}, RGA^{m3} and RGA^{m4} lost

their capacity to interact with these partners (Fig. 1b, Extended Data Fig. 1b). In addition,

RGA^{m2}, RGA^{m3} and RGA^{m4} were able to bind GID1 in the presence of GA, thus suggesting

that these DELLA candidates are still degraded in response to GA (Fig. 1c, Extended Data

Fig. 1c). To assess this possibility, we explored GA-dependent degradation of RGA^{m2},

RGA^{m3} and RGA^{m4} fused to GFP in transient expression assays. These three candidates were

degraded after GA treatment (Fig. 1d, Extended Data Fig. 1d), RGA^{m2}-GFP having the fastest

degradation kinetics although it was slightly more stable than RGA-GFP. Noteworthy,

RGA^{m2} harbours three amino acid substitutions in the GRAS PFYRE motif (Fig. 1a), that is

highly conserved in all plant DELLAs. Therefore, this variant is likely to have similar

properties in plant species other than *Arabidopsis*.

Based on the above results, we selected RGA^{m2} for further analysis. We next showed that

RGA^{m2} is unable to bind with the BZR1 DELLA interacting partner in yeast and a larger

screen confirmed that these mutations abolishes interactions with practically all known

DELLA partners (Fig. 1b, Extended Data Fig. 2). Moreover, co-immunoprecipitation studies

demonstrated that binding of RGA^{m2} to IDD2 is also strongly reduced *in planta* compared to

RGA (Fig. 1e). Finally, transient expression assays confirmed that, while RGA represses

BZR1 and TCP14 transcriptional activities, RGA^{m2} did not (Fig. 1f-g). Taken together, these

data indicate that the expression of RGA^{m2} (hereafter named RGA^{mPFYR}) in plants has a

negligible impact on GA signalling. Thus, RGA^{mPFYR} constitutes a suitable DELLA variant

form for engineering a degradation-based biosensor that monitors GA signalling activity and

more specifically the combinatorial effect of GA and of its complex perception machinery.

Engineering a GA signalling sensor (Fig. 2; Extended Data Fig. 3-5)

To create a ratiometric GA signalling sensor, we fused the RGA^{mPFYR} protein to the fast

maturing yellow fluorescent protein VENUS[23] and co-expressed this fusion protein together

with a nuclear-localized non-degradable reference protein, TagBFP-NLS, under the

pUBQ10[24] or *pRPS5a*[25,26] constitutive promoter. We used the 2A self-cleaving peptide to

allow for a stoichiometric production of both fluorescent proteins, enabling quantification of

GA signalling activity using fluorescence intensity ratio between RGA^{mPFYR}-VENUS and

TagBFP[27,28] (Fig. 2a). We named the sensor lines qRGA^{mPFYR} (quantitative RGA^{mPFYR}) and

first analysed their TagBFP fluorescence pattern in vegetative and reproductive tissues (Fig.

2b and Extended Data Fig. 3). TagBFP fluorescence of *pUBQ10::qRGA^{mPFYR}* lines was

homogeneously distributed in hypocotyl, the vegetative shoot meristem, cotyledons and roots

(except in the root tip) but the TagBFP signal was unevenly distributed in the inflorescence

SAM, with a stronger signal in organ boundaries (Extended Data Fig. 3b). Conversely, the

TagBFP signal was strong in the root tip and the vegetative SAM in *pRPS5a::RGA^mPFYR* lines

(Fig. 2b), and showed a homogenous distribution in the inflorescence SAM (Extended Data

Fig. 3e). Hence, the choice of one of these constructs depends on the analysed tissue: we used

pUBQ10::RGA^mPFYR in subsequent experiments in seedlings, and *pRPS5a::RGA^mPFYR* for

experiments performed in inflorescence SAM .

To validate that qRGA^mPFYR can detect changes in GA levels or signalling activity, we next

investigated the effects of exogenous GA and paclobutrazol (PAC; an inhibitor of GA

biosynthesis) treatments. Noteworthy, the growth of qRGA^mPFYR seedling hypocotyls was

similar to wild-type under mock conditions and upon GA or PAC treatment, although

hypocotyls were slightly longer after GA treatment (Fig. 2c). Furthermore, shoot development

and plant fertility were not significantly affected in qRGA^mPFYR plants (Fig. 2d and Extended

Data Fig. 3g), confirming that qRGA^mPFYR negligibly interferes with plant growth and GA

responses. Also, while the TagBFP signal was unaffected in hypocotyls upon GA and PAC

application, RGA^mPFYR-VENUS fluorescence was respectively reduced and increased after

these treatments, alike the *pRGA::GFP-RGA* positive control (Fig. 2e). Similar results were

observed in qRGA^mPFYR roots. In contrast, VENUS fluorescence did not change in any of the

treatments in RGA^m3-VENUS and RGA^m4-VENUS lines, consistently with the lower GA-

dependent degradation rate of these variants (Extended Data Fig. 4). In further image analyses,

in order to fully cover the range of values in VENUS/TagBFP fluorescence ratio, we used 3 -

RGA^mPFYR-Venus/TagBFP as a quantitative measure of GA signalling activity (hereafter

named "GA signalling"; Fig. 2a; see also Supplementary Method 1). This quantitative

approach confirmed statistically significant changes in GA signalling activity in hypocotyls,

with an increase and decrease respectively upon GA and PAC treatments compared to

untreated seedlings (Fig. 2e,f). Exogenous GA and PAC treatments induced similar responses

in the SAM, although the effect of PAC was less pronounced than in hypocotyl (Extended

data Fig. 3h-y). Furthermore, GA signalling activity increased in the SAM with both

exogenous GA concentration and treatment duration (Extended Data Fig. 5), showing that

$qRGA^{mPFYR}$ is also suitable to be used as a GA signalling sensor in the SAM.

Next, we tested if $qRGA^{mPFYR}$ is able to report for changes in endogenous GA levels in

growing hypocotyls. Hypocotyl elongation is widely used to study the effect of light and dark

on cell expansion, a growth process that requires GA signalling[17,29]. While in dark-grown

hypocotyls, cells elongate along an oriented spatio-temporal gradient (from the base to the tip),

in the light, hypocotyl cells elongate rather homogenously and continuously during the entire

growth period[30]. Thus, we investigated GA signalling distribution in $qRGA^{mPFYR}$ hypocotyls

grown in light and dark conditions, and concomitantly measured hypocotyl lengths from 2

days after sowing (DAS). Whereas at 3 DAS, hypocotyl lengths of light and dark-grown

seedlings were similar, dark-grown hypocotyls were substantially longer than that of light-

grown hypocotyls from 4 DAS and reached a plateau at 7 DAS (Fig. 2g). This growth

response was identical to wild type (Col-0) (Fig. 2g). We then compared GA signalling

activity in the basal region of light- and dark-grown $qRGA^{mPFYR}$ hypocotyls at 3, 5, and 7

DAS (Fig. 2h). Consistent with the growth dynamics, GA signalling was not statistically

different between dark- and light-grown seedlings at 3 DAS. However, at 5 DAS, GA

signalling activity was significantly higher in the dark than in the light, mirroring the

activation of cell expansion in darkness. At 7 DAS, a shift in the GA response occurred, with

GA signalling activity being lower in the dark than in the light, which correlates with the

completion of hypocotyl elongation in dark condition. Thus, $qRGA^{mPFYR}$ allows following

dynamic changes in GA signalling activity in growing hypocotyls.

qRGA^{mPFYR} fluorescence depends on GA receptor activity (Fig. 3 and Extended Data Fig.

6)

As GA signalling activity depends on both GA concentration and GA perception, we analysed

the expression of the three *GID1* receptors in vegetative and reproductive tissues. In seedlings,

GID1-GUS reporter lines showed that *GID1a* and *c* are highly expressed in cotyledons (Fig.

3a-c). Moreover, all three receptors are expressed in leaves, lateral root primordia, root tip

(excluding root cap for *GID1b*) and vasculature (Fig. 3a-c). In the SAM, we only detected a

GUS signal for GID1b and 1c (Extended data Fig. 6a-c). *In situ* hybridization confirmed these

expression patterns and further demonstrated that *GID1c* is homogenously expressed at a low

level in the SAM while *GID1b* shows higher expression at the SAM periphery (Extended data

Fig. 6d-l). A *pGID1b::2xmTQ2-GID1b* translational fusion further revealed a graded *GID1b*

expression ranging from low or no expression in the SAM centre to high expression in organ

boundaries (Extended data Fig. 6m). Thus, GID1 receptors are unevenly distributed across

and within tissues. Moreover, we observed that overexpressing *GID1* (*pUBQ10::GID1a-*

mCherry) enhanced qRGA^{mPFYR} sensitivity to external GA application in hypocotyls (Fig. 3d).

Taken together, these results suggest that differential expression of GID1 receptors can

modulate qRGA^{mPFYR} emission ratio, confirming that this biosensor reports for the

combinatorial action of GA and GA receptors.

A GA signalling map in the shoot apical meristem (Fig. 4; Extended Data Fig. 7-9)

Distribution of GA signalling within the SAM has remained elusive so far. Thus, we used

plants expressing qRGA^{mPFYR} together with the *pCLV3::mCherry-NLS* stem cell reporter to

compute a high-resolution quantitative map of GA signalling activity, focusing on the L1

layer (epidermis; Fig. 4a-b, see Methods and Supplementary Method 1) given the key role of

L1 in controlling growth at the SAM[31]. Here, expression of *pCLV3::mCherry-NLS* provided a

fixed geometric reference for analysing the spatio-temporal distribution of GA signalling

activity[32]. Although GAs were proposed to be required for lateral organ development[4], we

observed that GA signalling was lower in flower primordia (P) from the P_3 stage onward (Fig.

4a-b), while young P_1 and P_2 primordia had intermediate activity similar to the one found in

the central zone (Fig. 4a-b). Higher GA signalling activity was found in the boundaries of

organ primordia, starting from P_1/P_2 (on the lateral sides of the boundary) and culminating

from P_4, and in all peripheral zone cells located between primordia (Fig. 4a-b; Extended data

Fig. 7a-b). This higher GA signalling activity was not only monitored in the epidermis, but

also in the L2 and upper L3 layers (Extended data Fig. 7b). This reveals a GA signalling

pattern mostly mirroring primordia distribution. The inter-primordia region (IPR) distribution

results from the progressive establishment of a high GA signalling activity between

developing primordia and the central zone, while in parallel GA signalling activity decreases

in primordia (Fig. 4c-d).

The distribution of GID1b and GID1c receptors (Extended Data Fig. 6b-c, g-m) suggests that

differential expression of GA receptors contributes to shape the GA signalling activity pattern

in the SAM. We wondered if differential accumulation of GA could also be involved. To

investigate this possibility, we used the nlsGPS1 GA FRET sensor[17]. An increased emission

ratio was detected in nlsGPS1 SAMs treated with 10 μM GA_{4+7} for 100 min (Extended Data

Fig. 8), indicating that nlsGPS1 responds to changes in GA concentration in the SAM as it

does in roots[17]. The spatial distribution of the nlsGPS1 emission ratio indicates that GA levels

are relatively low in the SAM external layers, but it shows they are elevated in the centre of

the SAM and in the inter-primordia region (Fig. 4e, Extended Data Fig. 8a, c). We also

treated SAMs with fluorescent GAs (GA_3-, GA_4-, GA_7-Fl) or with Fl alone as a negative

control. The Fl signal was distributed in the whole SAM, including the central zone and

primordia, although with lower fluorescence (Fig. 4f, Extended Data Fig. 9d). In contrast, all

the three GA-Fls specifically accumulated in primordia boundaries and also, to different

degrees, in the rest of the IPR with GA_7-Fl accumulating in most IPR cells (Fig. 4g, Extended

Data Fig. 9). Fluorescence intensity quantification demonstrated higher IPR cells to non-IPR

cells intensity ratios in the GA-Fl treated SAMs, compared to Fl-treated SAMs (Fig. 4h,

Extended Data Fig. 9c). Taken together, these results suggest that GAs are present at higher

levels in IPR cells. This indicates that the SAM GA signalling activity pattern results from

both differential expression of the GA receptors and from the differential accumulation of GA

in IPR cells. Our analysis thus identifies an unexpected spatio-temporal GA signalling pattern,

with a lower activity in the centre of the SAM and in primordia, while activity is elevated in

the IPR of the peripheral zone.

Correlation analyses suggest a role for GA signalling in cell division plane orientation in

the shoot apical meristem (Fig. 5)

To understand the role of differential GA signalling activity at the SAM, we analysed the

correlation between GA signalling activity, cell expansion and cell division using time-lapse

live imaging of qRGA^{mPFYR} *pCLV3::mCherry-NLS* SAMs. Given the role of GA in growth

regulation, a positive correlation was expected with cell expansion parameters. We thus first

compared maps of GA signalling activity to those of cell surfacic growth rate (as a proxy for

cell expansion intensity for a given cell and daughter cells if it divides) and growth anisotropy,

which measures the directionality of cell expansion (here also for a given cell and daughter

cells if it divides; Fig. 5a-b, see Methods and Supplementary Method 1). Our cell surfacic

growth intensity map of the SAM was consistent with previous observations[33,34], with a

minimal growth rate in boundaries and maximal rates in developing flowers (Fig. 5a). A

principal component analysis (PCA) showed an anti-correlation between GA signalling

activity and cell surfacic growth intensity (Fig. 5c). It further showed that the main axis of

variability, encompassing GA signalling input and growth intensity, is orthogonal to the

direction defined by high expression of *CLV3*, which argues in favour of excluding cells from

the centre of the SAM in the rest of the analysis. Spearman correlation analyses confirmed the

PCA results (Fig. 5d), suggesting that higher GA signalling in the IPR does not lead to cell

expansion. However, the correlation analysis demonstrated a mild positive association

between GA signalling activity and growth anisotropy (Fig. 5c,d), suggesting that higher GA

signalling in the IPR act on cell growth orientation and possibly cell division plane

positioning.

Thus, we next studied the correlation between GA signalling and cell division activity, by

identifying newly formed cell walls during a time course analysis (Fig. 5e). This methodology

allows us to measure both the frequency and orientation of cell divisions. Strikingly, we found

that cell division frequency was similar in the IPR and the rest of the SAM (non-IPR, Fig. 5f),

showing that differences in GA signalling between IPR and non-IPR cells do not have a major

effect on cell division. This, together with the positive correlation between GA signalling and

growth anisotropy, lead us to ask whether GA signalling activity could act on cell division

plane orientation. We measured the orientation of new cell walls as the angle relative to the

radial axis connecting the centre of the meristem to the centre of the new cell walls (Fig. 5g),

and observed that cells had a clear tendency to divide at angles closer to 90° relative to the

radial axis, with the highest frequency observed at 70-80° (23.28%) and 80-90° (22.62%) (Fig.

5h) i.e. corresponding to cell divisions oriented in the circumferential/transverse direction. To

explore the contribution of GA signalling to this cell division behaviour, we analysed

separately the cell division parameters in the IPR and in the rest of the SAM (Fig. 5h). We

observed that the angle of cell division was different in the IPR compared to the rest of SAM

cells (non-IPR) or the entire SAM (Fig. 5h); IPR cells showing a higher rate of transverse cell

divisions, i.e. 70-80° and 80-90° (33.86% and 30.71% respectively) (Fig. 5h). Thus, our

observations reveal a link between high GA signalling and cell division plane orientation, that

parallels the correlation between GA signalling activity and growth anisotropy (Fig. 5c,d). To

further establish spatial conservation of this link, we measured the orientation of division

planes in IPR cells above primordia starting at stage P_3, given that the highest GA signalling

activity is detected in this region from stage P_4 (Fig. 4). The division angles of the IPR above

P_3 and P_4 did not show statistically significant differences, although an increase in the

frequency of transverse cell divisions was observed in the IPR above P_4 (Fig. 5i). Differences

in cell division plane orientation were however statistically significant in IPR cells above P_5,

in which the frequency of transverse cell divisions was drastically increased (Fig. 5i). Taken

together, these results suggest that GA signalling controls orientation of cell division in the

SAM coherently with previous reports[35,36], high GA signalling likely inducing a transverse

orientation of cell divisions in the IPR.

GA signalling activity positively regulates transverse cell divisions in the shoot apical

meristem (Fig. 6; Extended Data Fig. 10-12)

Cells in the IPR are expected not to be incorporated into primordia but rather in the

internodes[2,37,38]. A transverse orientation of cell divisions in the IPR could generate the typical

organization in parallel longitudinal cell files of the internode epidermis. Our observations

above indicate that GA signalling is likely to act in this process by regulating cell division

orientation.

Loss-of-function of multiple *DELLA* genes leads to constitutive GA response and thus *della*

mutants could be used to test this hypothesis[39]. We first analysed the expression patterns of

the five *DELLA* genes in the SAM. Transcriptional fusion GUS lines[40] showed that *GAI, RGA,*

and *RGL1* are expressed in the SAM and to a much lesser extent, *RGL2* (Extended Data Fig.

10a-d). *In situ* hybridization further showed that *GAI* mRNA specifically accumulates in

primordia and developing flowers (Extended Data Fig. 10e). *RGL1* and *RGL3* mRNA were

detected throughout the SAM dome and in older flowers, while *RGL2* mRNA was more

abundant in the boundary regions (Extended Data Fig. 10f-h). Confocal imaging of

pRGL3::RGL3-GFP SAM confirmed the expression observed with *in situ* hybridization and

showed that the RGL3 protein accumulates in the central part of the SAM (Extended Data Fig.

10i). Using a *pRGA::GFP-RGA* line, we also found that the RGA protein accumulates in the

SAM, but its abundance is reduced in boundaries starting from P_4 (Extended Data Fig. 10j).

Notably, the expression patterns of *RGL3* and *RGA* are compatible with a higher GA

signalling activity in the IPR, as detected with qRGA$^{\text{mPFYR}}$ (Fig. 4). In addition, these data

indicate that all *DELLAs* are expressed in the SAM and that collectively, their expression

covers the entire SAM.

We next analysed the cell division parameters in wild-type (L*er*, control) and quintuple

(global) *gai-t6 rga-t2 rgl1-1 rgl2-1 rgl3-4 della* mutants (Fig. 6a-b). Interestingly, we

observed a statistically significant change in frequency distribution of cell division angles in

the SAM of global *della* mutant compared to wild-type (Fig. 6c). This change in the global

della mutant resulted from an increase in frequency of 80-90° angles (34.71 % vs 24.55 %)

and to a lesser extent of 70-80° angles (23.78 % vs 20.18 %), i.e. corresponding to transverse

cell divisions (Fig. 6c). The frequency of non-transverse divisions (0-60°) was also lower in

global *della* mutant (Fig. 6c). The increased occurrence of transverse cell divisions was easily

visible in the SAM of global *della* mutant (Fig. 6b). The frequency of transverse cell divisions

in the IPR was also higher in global *della* mutant compared to wild-type (Fig. 6d). Outside of

the IPR region, distribution of the angle of cell division was more homogeneous in wild-type,

while in the global *della* mutant it was skewed toward tangential divisions, as in the IPR (Fig.

6e). Collectively, these observations indicate that constitutive activation of GA signalling in

the SAM induces transverse cell division both in the IPR and the rest of the SAM.

The orientation of cell division has been proposed to be influenced by the geometry of the

SAM and notably by tensile stresses prescribed by the curvature of the tissue[41]. We thus

analysed the SAM shape of the global *della* mutant. As previously shown[12], the size of the

global *della* mutant SAM was bigger than wild-type (Extended Data Fig. 11a,b,d). However,

the SAM curvature was identical in the two genotypes (Extended Data Fig. 11b,e,f,g,h,j). We

observed a comparable increase in size in the quadruple *gai-t6 rga-t2 rgl1-1 rgl2-1 della*

mutant, again without modification of the curvature compared to wild-type (Extended Data

Fig. 11c,d,f,i,j). Frequency of cell division orientation was also affected in the quadruple *della*

mutant, but to a lesser extent than in the global *della* mutant (Extended Data Fig. 12). This

dosage effect, along with the absence of effects on curvature, suggests that changes in the

occurrence of transverse cell division depends on changes in GA signalling activity rather

than in SAM geometry, and that the remaining RGL3 activity in quadruple *della* mutants

limits the change in cell division orientation caused by the loss of DELLA activity. Taken

together, our results indicate that GA signalling positively regulates transverse orientation of

cell division.

We then tested the effect of inhibiting GA signalling specifically in the IPR. To do so, we

used the *CUP-SHAPED COTYLEDON 2* (*CUC2*) promoter to drive expression of the

dominant negative gai-1 protein fused to VENUS (in a *pCUC2::gai-1-VENUS* line). *CUC2*

promoter drives expression in a large part of the IPR (including the boundary cells) in the

SAM starting from P_4 (Extended Data Fig. 11k). The distribution of cell division angles in the

entire SAM or in the IPR of *pCUC2::gai-1-VENUS* plants did not show statistically

significant differences with respect to the wild-type (Fig. 6f-i), although unexpectedly, we

found in these plants a higher frequency of 80-90° divisions in non-IPR cells. We thus

analysed the geometry of *pCUC2::gai-1-VENUS* SAMs and observed that size of the SAM

was reduced, whereas it had a significantly higher curvature (Extended Data Fig. 11l-q). This

change in the *pCUC2::gai-1-VENUS* SAM curvature could generate a mechanical stress

distribution whereby high circumferential stress starts at a short distance from the SAM

centre[42]. This could counteract in part the effect of GA signalling changes by increasing the

probability of cell division with circumferential/transverse orientation, thus explaining our

observations.

Taken together, these data support a direct role for GA signalling in transverse orientation of

the cell division plane in the IPR.

High GA signalling initiates internode specification in the shoot apical meristem (Fig. 7;

Extended Data Fig. 13)

Transverse orientation of division planes in the IPR as a result of higher GA signalling

activity opens the possibility that GAs pre-organize radial cell files in the epidermis within the

SAM to specify the cellular organization later found in the internode epidermis. Indeed, such

cell files are often visible in the SAM images of the global *della* mutant (Fig. 6b). Therefore,

to further understand the developmental function of the GA signalling spatial pattern in the

SAM, we used time-lapse imaging to analyse the cell spatial organization in the IPR in wild

type (L*er* and Col-0), global *della* mutant and *pCUC2::gai-1-VENUS* transgenic plants.

We marked L*er* cells above and on the side of P_4 according to their developmental fate

(analysed 34 hrs after the first observation, i.e. more than two plastochrones) using three

different colours: yellow for those incorporated into the primordia in the vicinity of P_4, green

for those located in the IPR and orange for those contributing to both (Fig. 7a-c). At t0 (0 h),

1-2 layers of IPR cells were visible in front of P_4 (Fig. 7a). As expected, when these cells

divide, they mostly do it with a transverse division plane (Fig. 7a-c). Similar results were

obtained with Col-0 SAMs (focusing on P_3 that had a comparable folding at the boundary

than P_4 in L*er*), although in this genotype the formation of the crease at the flower boundary

hides the IPR cells more rapidly (Fig. 7g-i). Thus, their division pattern and localization in the

IPR support the idea that these cells correspond to internode precursors.

Compared to Ler, 1-2 extra layers of IPR cells were observed in front of P$_4$ at t0 (0 h) in the

SAM of global $della$ mutants. These cells divided several times in 34 hrs (Fig. 7d-f, compared

to 7a-c), and in consequence, the mostly transverse divisions of IPR cells led to a higher

population of cells organized in radial cell files (Fig. 7d-f, compared to 7a-c). This indicates

that the higher GA signalling activity in global $della$ mutant SAMs promotes internode

specification. We conducted a similar analysis in $pCUC2::gai-1-VENUS$ plants. Since

expression of this transgene causes changes in SAM geometry (Extended Data Fig.

11m,n,p,q), we analysed IPR cells above the first primordium showing a comparable folding

at the boundary as Col-0 P$_3$. In these plants, opposite to global $della$ mutants, much less cell

divisions occurred in the IPR and there was no clear sign of an organization in radial cell files

(Fig. 7j-l), thus showing that inhibition of GA signalling in the IPR perturbs the specification

of the cellular organisation of internodes in the SAM. In lines with these results, we were able

to detect the appearance of internodes in the global $della$ mutant just below the SAM using

electron microscopy, while flowers remained compacted in Ler (Fig. 7m-n). By contrast,

organs were much more compacted in the SAM of $pCUC2::gai-1-VENUS$ than in Col-0 (Fig.

7o-p), consistent with the taller and shorter inflorescence stem of the global $della$ mutant[39,43]

and $pCUC2::gai-1-VENUS$ plants, respectively (Extended Data Fig. 13). Our results thus

support the hypothesis that higher GA signalling activity in the IPR specifies the cellular

organization of internodes in the SAM, through a regulation of the orientation of cell division

planes (Extended Data Fig. 14).

Discussion

Here, we developed a new ratiometric GA signalling biosensor, qRGA^{mPFYR}, that provides

information on GA function at the cellular level by allowing quantitative mapping of GA

signalling activity that results from the combinatorial action of GA and GA receptors

concentrations, with minimal interference with the endogenous signalling pathway. To this

end, we have engineered a modified DELLA protein, RGA^{mPFYR}, that has lost its capacity to

bind DELLA interacting partners but that remains sensitive to GA-induced proteolysis.

qRGA^{mPFYR} responds to both exogenous and endogenous changes in GA levels and its

dynamic sensing properties allow the assessment of spatio-temporal changes in GA signalling

activity during developmental processes. qRGA^{mPFYR} is also a highly flexible tool as it can be

adapted to a variety of tissues simply by changing, if needed, the promoter used for its

expression and is very likely transferable to other species given the conserved nature of the

GA signalling pathway and of the PFYRE motif in angiosperms.

Internode development is a key trait for plant architecture and crop improvement. qRGA^{mPFYR}

revealed a higher GA signalling activity specifically in cells of the IPR, that are the precursors

of internodes. By combining quantitative image analysis and genetics, we show that the GA

signalling pattern imposes circumferential/transverse cell division planes in the SAM

epidermis, shaping the cell division organization required for internode development. Few

developmental regulators of the orientation of the cell division plane during development

have been identified[44,45]. Our work provides a striking example of how GAs act in controlling

this cellular parameter. Mechanical stress has also been suggested to regulate cell division

plane orientation by acting on cortical microtubule[34,41,46] and our results are compatible with a

combined effect of mechanical stress and GAs in this regulation. GA signalling could affect

the orientation of the cell division plane also through an effect on cortical microtubule

orientation[35,36,47]. Likewise, previous works have highlighted the importance of the DELLA

interacting proteins TCP14 and 15 in the control of internode patterning[48,49] and these factors

could convey GA action, together with *BREVIPEDICELLUS* (*BP*) and *PENNYWISE* (*PNY*), which regulate internode development and have been shown to affect GA signalling[2,50]. Early cytological studies showed that both the inner tissues and the peripheral zone of the SAM are required for internode development in *Arabidopsis*[2,37]. The fact that GAs positively regulate cell division in the inner tissues supports a dual function of GAs in regulating meristem size and internode at the SAM. Patterns of oriented cell division are also highly regulated in the inner SAM tissues, and this regulation is essential to stem growth[44]. It will be interesting to explore whether GAs also play a role in orienting cell division planes in the inner tissues of the SAM and thus synchronize internode specification and development within the SAM.

Reference

1. Hay, A. & Tsiantis, M. *KNOX* genes: versatile regulators of plant development and diversity. *Development* **137**, 3153–3165 (2010).

2. Smith, H. M. S. & Hake, S. The interaction of two homeobox genes, *BREVIPEDICELLUS* and *PENNYWISE*, regulates internode patterning in the *Arabidopsis* inflorescence. *Plant Cell* **15**, 1717–1727 (2003).

3. Hay, A. *et al.* The Gibberellin pathway mediates KNOTTED1-type homeobox function in plants with different body plans. *Curr. Biol.* **12**, 1557–1565 (2002).

4. Jasinski, S. *et al.* KNOX action in *Arabidopsis* is mediated by coordinate regulation of cytokinin and gibberellin activities. *Curr. Biol.* **15**, 1560–1565 (2005).

5. Shani, E., Yanai, O. & Ori, N. The role of hormones in shoot apical meristem function. *Curr. Opin. Plant Biol.* **9**, 484–489 (2006).

6. Hedden, P. & Sponsel, V. A century of gibberellin research. *J. Plant Growth Regul.* **34**, 740–760 (2015).

7. Davière, J.-M. & Achard, P. Gibberellin signaling in plants. *Development* **140**, 1147–1151

(2013).

8. Silverstone, A. L. *et al.* Repressing a repressor: gibberellin-induced rapid reduction of the

RGA protein in *Arabidopsis. Plant Cell* **13**, 1555–1566 (2001).

9. Fu, X. *et al.* Gibberellin-mediated proteasome-dependent degradation of the barley DELLA

protein SLN1 repressor. *Plant Cell* **14**, 3191–3200 (2002).

10. Ueguchi-Tanaka, M. *et al. GIBBERELLIN INSENSITIVE DWARF1* encodes a soluble

receptor for gibberellin. *Nature* **437**, 693–698 (2005).

11. Kinoshita, A. *et al.* Regulation of shoot meristem shape by photoperiodic signaling and

phytohormones during floral induction of *Arabidopsis. eLife* **9**, e60661 (2020).

12. Serrano-Mislata, A. *et al. DELLA* genes restrict inflorescence meristem function

independently of plant height. *Nat. Plants* **3**, 749–754 (2017).

13. Achard, P. *et al.* Gibberellin signaling controls cell proliferation rate in *Arabidopsis. Curr.*

Biol. **19**, 1188–1193 (2009).

14. Ubeda-Tomás, S. *et al.* Gibberellin signaling in the endodermis controls *Arabidopsis* root

meristem size. *Curr. Biol.* **19**, 1194–1199 (2009).

15. Shani, E. *et al.* Gibberellins accumulate in the elongating endodermal cells of *Arabidopsis*

root. *Proc. Natl. Acad. Sci.* **110**, 4834–4839 (2013).

16. Tal, I. *et al.* The *Arabidopsis* NPF3 protein is a GA transporter. *Nat. Commun.* **7**, 11486

(2016).

17. Rizza, A., Walia, A., Lanquar, V., Frommer, W. B. & Jones, A. M. *In vivo* gibberellin

gradients visualized in rapidly elongating tissues. *Nat. Plants* **3**, 803–813 (2017).

18. Wang, F. *et al.* Biochemical Insights on degradation of *Arabidopsis* DELLA proteins

gained from a cell-free assay system. *Plant Cell* **21**, 2378–2390 (2009).

19. Hirano, K. *et al.* Characterization of the molecular mechanism underlying gibberellin

perception complex formation in rice. *Plant Cell* **22**, 2680–2696 (2010).

20. Hussain, A., Cao, D. & Peng, J. Identification of conserved tyrosine residues important for

gibberellin sensitivity of *Arabidopsis* RGL2 protein. *Planta* **226**, 475–483 (2007).

21. Hussain, A., Cao, D., Cheng, H., Wen, Z. & Peng, J. Identification of the conserved

serine/threonine residues important for gibberellin-sensitivity of *Arabidopsis* RGL2 protein.

Plant J. **44**, 88–99 (2005).

22. Velde, K. V. D., Ruelens, P., Geuten, K., Rohde, A. & Straeten, D. V. D. Exploiting

DELLA signaling in cereals. *Trends Plant Sci.* **22**, 880–893 (2017).

23. Shaner, N. C., Steinbach, P. A. & Tsien, R. Y. A guide to choosing fluorescent proteins.

Nat. Methods **2**, 905–909 (2005).

24. Grefen, C. *et al.* A ubiquitin-10 promoter-based vector set for fluorescent protein tagging

facilitates temporal stability and native protein distribution in transient and stable expression

studies. *Plant J.* **64**, 355–365 (2010).

25. Weijers, D. *et al.* An *Arabidopsis* minute-like phenotype caused by a semi-dominant

mutation in a *RIBOSOMAL PROTEIN S5* gene. *Dev. Camb. Engl.* **128**, 4289–4299 (2001).

26. Tol, N. van *et al.* Enhancement of *Arabidopsis* growth characteristics using genome

interrogation with artificial transcription factors. *PLOS ONE* **12**, e0174236 (2017).

27. Wend, S. *et al.* A quantitative ratiometric sensor for time-resolved analysis of auxin

dynamics. *Sci. Rep.* **3**, 2052 (2013).

28. Goedhart, J. *et al.* Quantitative co-expression of proteins at the single cell level –

application to a multimeric FRET sensor. *PLOS ONE* **6**, e27321 (2011).

29. Achard, P. *et al.* DELLAs contribute to plant photomorphogenesis. *Plant Physiol.* **143**,

1163–1172 (2007).

30. Gendreau, E. *et al.* Cellular basis of hypocotyl growth in *Arabidopsis thaliana. Plant*

Physiol. **114**, 295–305 (1997).

31. Savaldi-Goldstein, S., Peto, C. & Chory, J. The epidermis both drives and restricts plant

shoot growth. *Nature* **446**, 199–202 (2007).

32. Galvan-Ampudia, C. S. *et al.* Temporal integration of auxin information for the regulation

of patterning. *eLife* **9**, e55832 (2020).

33. Uyttewaal, M. *et al.* Mechanical stress acts via katanin to amplify differences in growth

rate between adjacent cells in *Arabidopsis. Cell* **149**, 439–451 (2012).

34. Burian, A. *et al.* A correlative microscopy approach relates microtubule behaviour, local

organ geometry, and cell growth at the *Arabidopsis* shoot apical meristem. *J. Exp. Bot.* **64**,

5753–5767 (2013).

35. Marc, J. & Hackett, W. P. Changes in the pattern of cell arrangement at the surface of the

shoot apical meristem in *Hedera helix* L. following gibberellin treatment. *Planta* **186**, 503–

510 (1992).

36. Locascio, A., Blázquez, M. A. & Alabadí, D. Dynamic regulation of cortical microtubule

organization through prefoldin-DELLA interaction. *Curr. Biol.* **23**, 804–809 (2013).

37. Vaughan, J. G. The morphology and growth of the vegetative and reproductive apices of

Arabidopsis thaliana (L.) Heynh., *Capsella bursa-pastoris* (L.) Medic. and *Anagallis arvensis*

L. *J. Linn. Soc. Lond. Bot.* **55**, 279–301 (1955).

38. Jackson, D., Veit, B. & Hake, S. Expression of maize *KNOTTED1* related homeobox

genes in the shoot apical meristem predicts patterns of morphogenesis in the vegetative shoot.

Development **120**, 405–413 (1994).

39. Cheng, H. *et al.* Gibberellin regulates *Arabidopsis* floral development via suppression of

DELLA protein function. *Development* **131**, 1055–1064 (2004).

40. Gallego Giraldo, C. *et al.* Role of the gibberellin receptors GID1 during fruit-set in

Arabidopsis. Plant J. **79**, 1020–1032 (2014).

41. Louveaux, M., Julien, J.-D., Mirabet, V., Boudaoud, A. & Hamant, O. Cell division plane

orientation based on tensile stress in *Arabidopsis thaliana. Proc. Natl. Acad. Sci.* **113**, E4294–

E4303 (2016).

42. Hamant, O. *et al.* Developmental patterning by mechanical signals in *Arabidopsis.*

Science **322**, 1650–1655 (2008).

43. Kumar, S. V. *et al.* Transcription factor PIF4 controls the thermosensory activation of

flowering. *Nature* **484**, 242–245 (2012).

44. Bencivenga, S., Serrano-Mislata, A., Bush, M., Fox, S. & Sablowski, R. Control of

oriented tissue growth through repression of organ boundary genes promotes stem

morphogenesis. *Dev. Cell* **39**, 198–208 (2016).

45. Yoshida, S. *et al.* Genetic control of plant development by overriding a geometric division

rule. *Dev. Cell* **29**, 75–87 (2014).

46. Rasmussen, C. G. & Bellinger, M. An overview of plant division-plane orientation. *New*

Phytol. **219**, 505–512 (2018).

47. Wenzel, C. L., Williamson, R. E. & Wasteneys, G. O. Gibberellin-induced changes in

growth anisotropy precede gibberellin-dependent changes in cortical microtubule orientation

in developing epidermal cells of barley leaves. kinematic and cytological studies on a

gibberellin-responsive dwarf mutant, M489. *Plant Physiol.* **124**, 813–822 (2000).

48. Davière, J.-M. *et al.* Class I TCP-DELLA interactions in inflorescence shoot apex

determine plant heigh. *Curr. Biol.* **24**, 1923–1928 (2014).

49. Kieffer, M., Master, V., Waites, R. & Davies, B. TCP14 and TCP15 affect internode

length and leaf shape in *Arabidopsis. Plant J.* **68**, 147–158 (2011).

50. Khan, M. *et al.* Repression of lateral organ boundary genes by *PENNYWISE* and *POUND-*

FOOLISH is essential for meristem maintenance and flowering in *Arabidopsis*. *Plant Physiol.*

169, 2166–2186 (2015).

51. Koini, M. A. *et al.* High temperature-mediated adaptations in plant architecture require the

bHLH transcription factor PIF4. *Curr. Biol.* **19**, 408–413 (2009).

52. Lee, S. *et al.* Gibberellin regulates *Arabidopsis* seed germination via *RGL2*, a *GAI/RGA-*

like gene whose expression is up-regulated following imbibition. *Genes Dev.* **16**, 646–658

(2002).

53. Pfeiffer, A. *et al.* Integration of light and metabolic signals for stem cell activation at the

shoot apical meristem. *eLife* **5**, e17023 (2016).

54. Karimi, M., Inzé, D. & Depicker, A. GATEWAY^{TM} vectors for Agrobacterium-mediated

plant transformation. *Trends Plant Sci.* **7**, 193–195 (2002).

55. Galinha, C. *et al.* PLETHORA proteins as dose-dependent master regulators of

Arabidopsis root development. *Nature* **449**, 1053–1057 (2007).

56. Simon, M. L. A. *et al.* A PtdIns(4)P-driven electrostatic field controls cell membrane

identity and signalling in plants. *Nat. Plants* **2**, 1–10 (2016).

57. Shcherbakova, D. M., Hink, M. A., Joosen, L., Gadella, T. W. J. & Verkhusha, V. V. An

orange fluorescent protein with a large stokes shift for single-excitation multicolor FCCS and

FRET imaging. *J. Am. Chem. Soc.* **134**, 7913–7923 (2012).

58. Resentini, F. *et al.* TCP14 and TCP15 mediate the promotion of seed germination by

gibberellins in *Arabidopsis thaliana*. *Mol. Plant* **8**, 482–485 (2015).

59. Gallego-Bartolomé, J. *et al.* Molecular mechanism for the interaction between gibberellin

and brassinosteroid signaling pathways in *Arabidopsis*. *Proc. Natl. Acad. Sci.* **109**, 13446–

13451 (2012).

60. de Lucas, M. *et al.* A molecular framework for light and gibberellin control of cell

elongation. *Nature* **451**, 480–484 (2008).

61. Castrillo, G. *et al.* Speeding cis-trans regulation discovery by phylogenomic analyses

coupled with screenings of an arrayed library of *Arabidopsis* transcription factors. *PLOS ONE*

6, e21524 (2011).

62. Vernoux, T., Kronenberger, J., Grandjean, O., Laufs, P. & Traas, J. PIN-FORMED 1

regulates cell fate at the periphery of the shoot apical meristem. *Dev. Camb. Engl.* **127**, 5157–

5165 (2000).

63. Rozier, F., Mirabet, V., Vernoux, T. & Das, P. Analysis of 3D gene expression patterns in

plants using whole-mount RNA in situ hybridization. *Nat. Protoc.* **9**, 2464–2475 (2014).

64. Schroeder, W., Martin, K. & Lorensen, B. *The visualization toolkit: an object-oriented*

approach to 3D graphics. (Kitware, 2006).

65. Landrein, B. *et al.* Mechanical stress contributes to the expression of the *STM* homeobox

gene in *Arabidopsis* shoot meristems. *eLife* **4**, e07811 (2015).

Methods

Growth conditions and Plant material

Plants were grown on soil or *in vitro* on 1x Murashige-Skoog (MS) medium (Duchefa) supplemented with 1% sucrose and 1% agar (Sigma) under standard conditions (16h photoperiod at 22°C), except for hypocotyl and root growth experiments, for which the seedlings were grown on vertical plates under continuous light and 22°C.

We used the following *Arabidopsis* mutants and transgenic lines in the L*er* background: *gai-t6 rga-t2 rgl1-1 rgl2-1* quadruple *della*[39], *gai-t6 rga-t2 rgl1-1 rgl2-1 rgl3-4* global *della*[51] or Col-0: *pGID1a::GID1a-GUS, pGID1b::GID1b-GUS, pGID1c::GID1c-GUS, pGAI::GUS, pRGA::GUS, pRGL1::GUS, rgl2-5* (a promoter trap GUS line)[52], *pRGA::GFP-RGA*[8], *pRGL3::RGL3-GFP*, *pCLV3::mCherry-NLS*[53], *nlsGPS1*[17].

All other transgenic lines were generated in the Col-0 background. To generate these lines and to perform transient assays in *Nicotiana benthamiana* leaves, plasmids were constructed in the following way. RGA cDNA (Agi code AT2G01570) and RGA^{m1}, RGA^{m2,} RGA^{m3} and RGA^{m4} mutant variants were obtained by PCR using specific primers numbered 1 to 8 in Supplementary Table 1, and inserted into pDONR221 (Thermo Fisher Scientific) by Gateway cloning and recombined with pB7FWG2[54] to generate *p35S::RGA-GFP* and *p35S::RGA$^{m1/m2/m3/m4}$-GFP*. To generate *pRPS5a::qRGA^{mPFYR}*, *pUBQ10:qRGA^{mPFYR}*, *pRPS5a::RGA$^{m1/m3/m4}$-VENUS-2A-BFP*, *pUBQ10::RGA$^{m1/m3/m4}$-VENUS-2A-BFP* and *pCUC2::gai-1-VENUS*, the promoter region of *pRPS5a* (1.7 kb fragment), *pUBQ10* (2.5 kb fragment) or *pCUC2* (3.2 kb fragment) inserted into pDONR P4-P1R (Thermo Fisher Scientific), *RGA^{mPFYR}-VENUS*, *RGA$^{m1/m3/m4}$-VENUS* or *gai-1* cDNA inserted into pDONR221, and *2A-TagBFP* or *VENUS* (for *pCUC2::gai-1-VENUS)* inserted into pDONR P2R-P3 (Thermo Fisher Scientific), were recombined into pB7m34GW[54]. To obtain *pUBQ10::GID1a-mCherry*, *GID1a* cDNA inserted into pDONR221 was recombined with

pDONR P4-P1R-*pUBQ10* and pDONR P2R-P3-*mCherry* into pB7m34GW. *p35S:IDD2-RFP*

was obtained by recombining *IDD2* cDNA inserted in pDONR221 into pB7RWG2[54]. To get

pGID1b::2xmTQ2-GID1b, a 3.9-kb fragment upstream of the coding region of *GID1b* and a

4.7-kb fragment including *GID1b* cDNA (1.3 kb) and terminator (3.4 kb) were first amplified

using primers in Supplementary Table 1, then inserted into pDONR P4-P1R (Thermo Fisher

Scientific) and pDONR P2R-P3 (Thermo Fisher Scientific), respectively, and finally

recombined into pGreen 0125[55] destination vector with pDONR221 2xmTQ2[56] by Gateway

cloning. To make *pCUC2::LSSmOrange*, the promoter sequence of *CUC2* (3229 bp upstream

the ATG), followed by the coding sequence for the large stokes shift mOrange

(LSSmOrange)[57] with a N7 nuclear localization signal, and the NOS transcription terminator,

was assembled into the pGreen Kanamycin destination vector using 3 fragment Gateway

recombination system (Invitrogen). For transactivation assays, *pTCP-LUC* and *pBRE-LUC*

reporter constructs (containing synthetic promoters) and *p35S::3xHA-VP16-TCP14* and

p35S::3xHA-VP16-BZR1 encoding effector proteins were used as previously described[58,59] .

Plant binary vectors were inserted into *Agrobacterium tumefaciens* GV3101 strain and

respectively introduced into *Nicotiana benthamiana* leaves by agro-infiltration and

Arabidopsis Col-0 by floral dip. *pUBQ10::qRGA^{mPFYR} pUBQ10::GID1a-mCherry* and

pCLV3::mCherry-NLS qRGA^{mPFYR} were isolated from F3 and F1 progeny of the appropriate

crosses, respectively.

Pharmacological treatments

Chemical treatments with GA (GA$_3$, Sigma or Duchefa) and paclobutrazol (PAC, Duchefa)

were performed at the concentrations and time indicated in figures. For Y2H assays, 100 μM

GA$_3$ were added in selective media to promote interaction between RGA and GID1. For RGA

and RGA$^{m1/m2/m3/m4}$ degradation kinetics, *N. benthamiana* agro-infiltrated leaf discs were

incubated with 100 μM GA_3 and 100 mM cycloheximide (Sigma) over 300 min. For

hypocotyl length measurements and fluorescence analyses of $qRGA^{mPFYR}$ hypocotyls and

roots, seedlings were grown for 5 days on MS agar medium and then transferred for 4 days on

MS agar plates supplemented with 5 μM PAC or 10 μM GA_3. For GA_3 treatment on

$qRGA^{mPFYR}$, GA_{4+7} treatment on nlsGPS1, and GA-Fl treatment on L*er*, dissected shoot

apices were inserted into Apex Culture Medium (ACM, 1/2x MS medium (Duchefa), 1%

sucrose, 1% agarose, 2 mM MES (Sigma), 1x vitamin solution (myo-Inositol 100 mg/L,

nicotinic acid 1 mg/L, pyridoxine hydrochloride 1 mg/L, thiamine hydrochloride 10 mg/L,

glycine 2 mg/L), 200 nM N6-Benzyladenine) with indicated concentrations of GA/GA-Fl,

and also immersed under 200 μl of GA/GA-Fl solution of indicated concentrations for a

indicated period of time. For PAC treatment on $qRGA^{mPFYR}$, 50 μM PAC were sprayed on the

whole inflorescence every two days for a period of 5 days before observation.

Yeast two-hybrid assays

For Y2H assays, both the full-length and the C-terminal part of RGA and $RGA^{m1/m2/m3/m4}$

(named M5 version, amino acids 199 to 587; the N-terminal part is subject to self-activation

in yeast[60]) inserted into pDONR221 were recombined into pGBKT7 (Clontech) to obtain BD-

RGA, $BD-RGA^{m1/m2/m3/m4}$, BD-M5RGA and $BD-M5RGA^{m1/m2/m3/m4}$. On the other hand, JAZ1,

TCP14, IDD2, BZR1, GID1a, GID1b and GID1c cDNAs inserted into pDONR221 were

fused to the activation domain GAL4 (AD) after recombination into pGADT7 (Clontech).

Direct interaction assays were carried out following the Clontech procedures. Briefly, BD-

M5DELLA and AD-JAZ1, AD-TCP14, AD-IDD2, AD-BZR1 and, on the other hand, BD-

DELLA (full length) and AD-GID1 constructs were co-transformed in the yeast strain AH109

and interactions tests were surveyed on selective medium lacking tryptophan, leucine and

histidine. In some cases, the medium was supplemented with 3-amino-1, 2, 4 triazole (3AT,

Sigma) or with GA to promote interaction between DELLA and GID1.

To confirm that the RGA^{m2} mutation impaired interaction with all RGA protein partners, we

used the C-terminal part of RGA and RGA^{m2} (amino acids 199 to 587) and of GAI (amino

acids 148 to 533) for positive control, to probe the Arabidopsis transcription factors REGIA +

REGULATORS (RR) arrayed library, following the protocol described in Castrillo G. et al.[61].

TFs in the RR library were fused to GAL4 activation domain of the pDEST22 vector and

independently transformed into the yeast strain YM4271 in 96-well plates. The RGA^{m2}

protein fused to the GAL4 BD in the pDEST32 vector was transformed into the pJ694 yeast

strain and used as bait. Replicates of the library were grown overnight on SD-Trp solid media

and inoculated together with 100 μL of an overnight RGA^{m2} culture grown on SD-Leu on

microtiter plates containing 100 μL of YPDA per well. Plates were incubated for 2 days at

30°C for mating, and diploid colonies were selected in new 96-well plates containing 200 μL

of SD-Leu/Trp. 5 μL of the diploid cell cultures were lastly tested for protein interaction by

placing them on solid SD medium lacking both Leu and Trp (positive growth control), and on

SD medium lacking Leu, Trp and His, in the presence of different concentrations of 3-

aminotriazol (3-AT) (Sigma-Aldrich). Results were expressed in the form of a heat map for

the strength of interaction according to the colony growth after five days of incubation at

30°C.

Co-immunoprecipitation assays

CoIP assays of IDD2-RFP with RGA-GFP or RGA^{m2}-GFP were performed on *N.*

benthamiana agro-infiltrated leaves with *p35S::IDD2-RFP*, *p35S::RGA-GFP* or

p35S::RGA^{m2}-GFP. Three days after infiltration, total proteins were extracted with the native

extraction buffer [Tris-HCl (pH 7.5) 50 mM, glycerol 10%, Nonidet P-40 0.1% supplemented

with Complete Protease Inhibitors 1X (Roche)], and then incubated for 2h at 4°C with 50 µl

of anti-GFP antibody conjugated with paramagnetic beads (Miltenyi Biotec). After incubation,

samples were loaded onto a magnetic column system (µ columns ; Miltenyi Biotec) to recover

the immunoprotein complexes according to manufacturer's protocol. The immunoprecipitated

(RGA-GFP and RGA^{m2}-GFP) and co-immunoprecipitated (IDD2-RFP) proteins were

detected by western-blot with anti-GFP (JL8; Clontech) and anti-RFP (6G6; Chromotek),

respectively.

Immunodetection analyses

Plant materials were ground in 2x SDS-PAGE buffer followed by heating at 95°C for 5 min.

After centrifugation at 13000 *g* for 5 min, total proteins were separated on 8.5% SDS-PAGE

gel and transferred to an immobilon-P (PVDF) membrane (Millipore). Membranes are then

saturated with blocking buffer (TBS 1x, Tween-20 0.1%; milk 5%) and incubated with a

2000-fold dilution of anti-GFP (JL8; Clontech), anti-RGA (6G6; Chromotek), anti-HA

(Sigma) or anti-actin (Agrisera) antibodies, and a 5000-fold dilution of peroxidase-conjugated

goat anti-rabbit or mouse IgG (Invitrogen). Signals were detected using the Luminata Forte

Western HRP Substrate (Millipore).

Transactivation assays

Transactivation assays were performed using the Dual-Glo Luciferase Assay System

(Promega) according to manufacturer's instructions. Briefly, *N. benthamiana* leaves were

agro-infiltrated with *pTCP-LUC* and *pBRE-LUC* reporter constructs, *p35S::3xHA-VP16-*

TCP14 or *p35S::3xHA-VP16-BZR1* encoding effector proteins, and *p35S::RGA-GFP* or

p35S::RGA^{m2}-GFP. Three days after infiltration, total proteins were extracted in lysis buffer

(Promega), then Firefly and control Renilla LUC activities were quantified with FLUOstar

Omega luminometer (BMG Labtech) using OMEGA2 software version 5.50 R4. For loading

control, protein levels were analysed by immunodetection.

GUS staining

For GUS expression detection in inflorescence apex, 28-d-old plants grown under LD

conditions were used. After fixation in 90% cold acetone at room temperature for 20 min,

shoot apexes were transferred into GUS staining solution containing 1 mM potassium

ferrocyanide, 1 mM potassium ferricyanide and 1 mg/ml 5-bromo-4-chloro-3-indolyl-*ß*-D-

glucuronide (X-Gluc), vacuum-infiltrated on ice for 15 min, and incubated overnight at 37 °C

in dark before washing with ethanol series and microscope detection. For the experiment with

seedlings, 7-d-old plantlets grown in vitro under LD conditions were vacuum-infiltrated for

15 min in GUS staining solution containing 2 mM potassium ferrocyanide, 2 mM potassium

ferricyanide and 0.25 mg/ml X-Gluc, and incubated at 37°C for 24 h. Then GUS solution was

replaced with ethanol 70% and seedlings were observed using optical microscope.

In situ **hybridizations**

RNA in situ hybridizations were performed according to Vernoux et al.[62] with some minor

modifications on fixation. Briefly, shoot apices with ~1cm stem were collected and

immediately fixed in FAA solution (3.7% Formaldehyde, 5% Acetic Acid, 50% Ethanol)

precooled at 4 ºC. After vacuum treatments for 2x 15 min, the fixative was changed and the

samples were incubated overnight. Antisense probes of the cDNA and 3'-UTR of *GID1a*,

GID1b, *GID1c*, *GAI*, *RGL1*, *RGL2* and *RGL3* were synthesized as described in Rozier et al.[63]

using primers indicated in Supplementary Table 1. Immunodetection of the digoxigenin-

labelled probes was performed using an anti-digoxigenin antibody, and sections were stained

with 5-Bromo-4-chloro-3-indoryl Phosphate (BCIP)/Nitroblue Tetrazolium (NBT) solutions.

Microscopy

For confocal microscopy observations of hypocotyl, complete seedlings were put on slides and images were obtained with a Zeiss LSM 780 confocal microscope. SAM live imaging was performed as described[32]. Briefly, dissected SAMs were let to recover overnight after dissection. Confocal images were taken with a Zeiss LSM 710 or 700 confocal laser scanning microscope equipped with a water-dipping lens (W Plan-Apochromat 40x/1.0 DIC). The confocal settings were as previously described[32] and detailed below. To observe GFP or GFP with propidium iodide (PI), a laser of 488 nm was used to excite, and the emission for GFP was 500-520 nm (in some cases it was 500-540 nm), for PI was 610-650 nm. For imaging of qRGA^{mPFYR} or qRGA^{mPFYR} with mCherry and PI, lasers of 514 nm, 405 nm, 561 nm and 488 nm were used for the excitation of VENUS, TagBFP, mCherry and PI, respectively, and the corresponding emission wavelength was 520-560 nm, 430-460 nm, 580-615 nm and 620-660 nm. mTURQUOISE2 (mTQ2) was excited by 445 nm laser and the emission range was 470-510 nm. For FRET detection of nlsGPS1, an acquisition mode of spectral imaging (λ-scan) with emission wavelength from 461 nm to 597 nm was used with excitation at 458 nm.

For optical microscopy, photographs of plants were taken with a LEICA MZ12 stereoscopic microscope equipped with a ZEISS AxioCam ICc5 camera head or Canon camera.

For scanning electronic microscopy of fresh shoot apex, a Hirox SH-3000 table-top microscope equipped with Coolstage (-20 °C to -30 °C) was used.

Image processing

Confocal stacks were processed in Fiji (fiji.sc) to get max projection or orthogonal views. Older primordia were also removed in Fiji. For 3D visualization of confocal stacks, we used

the Zeiss ZEN2 software (Fig. 7) and a rendering using the VTK library[64] (Fig. 5e, Fig. 6). In

both cases, parameters were adjusted accordingly to show mainly the L1 cells.

Quantification and statistical analysis

GA sensor quantification

For qRGA^{mPFYR} quantification and visualization, images of hypocotyls were analyzed using a

Python script that performs nuclei detection and signal quantification in 3D. The details of the

algorithms used in the analysis are provided in Supplementary Method 1 (Section 2 – Nuclei

detection and signal quantification).

For nlsGPS1 quantification and visualization, a series of Fiji macros were used (macros are

deposited in Protocol Exchange). Briefly, the "seg-auto.ijm" macro was run for segmentation

of nuclei based on the sum of signals from all the 14 channels of the spectral imaging. After

removing little objects corresponding to signal noise using the "object-screening.ijm" macro,

the ratio of DxAm (channel 8) to DxDm (channel 3) of each segmented nucleus was

calculated by running the "3D-ratio.ijm", and from here, the 3D construction of the SAM,

with every nucleus showing its ratio value, was also achieved and printed. Statistical analysis

was done in R.

SAM image sequence analysis

To analyze time-lapse confocal acquisitions of SAMs and obtain quantitative measures of GA

signalling as well as cellular growth parameters, we developed a computational pipeline. It

consists of 3D watershed segmentation for cell identification from the PI staining, nuclei

detection and qRGA quantification, temporal registration using the PI image channel and

surfacic growth estimation based on expert cell lineages. Individual SAM sequences were

then aligned in order to perform population-scale statistics. Extensive details on the

algorithms used in this pipeline are provided in Supplementary Method 1.

Hypocotyl length measurement

To measure hypocotyl length, seedlings were grown in vertical MS medium (Duchefa) with 0.8% (w/v) phytoagar and 1% sucrose for 5 days and transferred to MS, MS supplemented with 10 μM GA3 or 5 μM Paclobutrazol for 4 days in continuous light at 22ºC. Plates were scanned and hypocotyl length was measured from the images using Fiji.

Cell division orientation quantification

New cell walls were identified by comparing images obtained at 0 h and 10 h, and masked with a manually drawn line in Fiji. Then, a macro was used to skeletonize and then to measure the angles of the drawn cell walls with an expert-defined centre of the SAM. Further angle frequency distribution was done in Excel using Pivot table. Statistical analysis of Kolmogorov-Smirnov test was performed using an online tool: http://www.physics.csbsju.edu/stats/KS-test.n.plot_form.html.

Curvature quantification

The MorphoGraphX software was used to quantify cellular curvature of L1 cells of SAMs of different genetic backgrounds. Statistical analysis was done in R.

Meristem size measurement

To measure meristem size, a method described before was used[65]. Briefly, the SAM radius was determined by drawing a circle that covers I_1 and I_2 and that the center of which is roughly overlapping with the geometrical center of the SAM surface using Fiji software. Statistical analysis was done in R.

Synthesis and characterization of GA-Fluorescein (GA-Fl)

GA-Fluorescein (GA-Fl) were synthesized using a previously described protocol[15]. Extensive details on synthesis and characterization are provided in Supplementary Method 2.

 Figure legends

 Figure 1. The modified RGA^{m2} protein is an inactive DELLA that remains sensitive to

 GA.

a, Schematic representation of the domain structure of a typical DELLA protein. The nucleic

acids and amino acids mutated in RGA^{m2} are indicated in red. **b**, Yeast two hybrid (Y2H)

assays in which RGA and RGA^{m2} were tested pairwise with four known DELLA-interacting

partners: JAZ1, TCP14, IDD2 and BZR1. Empty pGBKT7 and pGADT7 vectors were

included as negative controls. Photos show the growth of the yeast on control media (GM)

and on selective media (SM). **c**, Pairwise Y2H interaction assays between RGA or RGA^{m2}

and the three GA receptors GID1a, GID1b and GID1c. Photos show the growth of the yeast

on control media (GM), selective media (SM) and SM media supplemented with 100 μM GA$_3$.

d, Time-course analysis of GA-induced degradation of RGA (upper panel) and RGA^{m2}

protein (lower panel). Immunodetection of RGA-GFP and RGA^{m2}-GFP protein in *35S::RGA-*

GFP and *35S::RGA^{m2}-GFP N. benthamiana* agro-infiltrated leaves treated with 100 mM

cycloheximide (CHX) and 100 μM GA$_3$ for the indicated times. Numbers indicate RGA-GFP

and RGA^{m2}-GFP levels relative to Actin levels, used as loading control. **e**,

Coimmunoprecipitation assays between RGA or RGA^{m2} and IDD2. Protein extracts from

different combinations of *N. benthamiana* agro-infiltrated leaves with *35S::RGA-GFP*,

35S::RGA^{m2}-GFP and *35S::IDD2-RFP* were immunoprecipitated with anti-GFP antibodies.

The co-immunoprecipitated protein (IDD2-RFP) was detected by anti-RFP antibodies. **f-g,**

Effect of RGA and RGA^{m2} on BZR1 (**f**) and TCP14 (**g**) transcriptional activities in *N.*

benthamiana agro-infiltrated leaves with a combinations of BZR1, TCP14, RGA and RGA^{m2}

effector constructs and corresponding Luciferase/Renilla reporter constructs, as indicated (top

panels). Transcriptional activities are represented as the ratio of Luciferase and Renilla (used

as internal control) activities, relative to the value obtained for the reporter construct alone

that was set to 1. Different letters denote significant differences ($p < 0.05$) using ANOVA

Tukey's tests for multiple comparisons. Bottom panels: immunodetection of RGA-GFP,

RGA^{m2}-GFP, HA-VP16-BZR1 and HA-VP16-TCP14 from *N. benthamiana* agro-infiltrated

leaves used for transcriptional activity assays.

Extended Data Figure 1. Characterization of mutant DELLA proteins.

a, Position of the mutated nucleic acids and amino acids of the four candidate modified

DELLA proteins used in the study: RGA^{m1}, RGA^{m2}, RGA^{m3}, RGA^{m4}. *: stop codon. **b**,

Pairwise Y2H interaction assays between RGA, RGA^{m1}, RGA^{m3}, RGA^{m4} and three DELLA-

interacting proteins, JAZ1, TCP14 and IDD2. Empty pGBKT7 and pGADT7 vectors were

included as negative controls. Photos show the growth of the yeast on control media (GM)

and on selective media (SM). **c**, Pairwise Y2H interaction assays between RGA, RGA^{m3},

RGA^{m4} and GID1a, GID1b and GID1c. Photos show the growth of the yeast on control media

(GM), selective media (SM) and SM media supplemented with 100 μM GA$_3$. **d**, Time-course

analysis of GA-induced degradation of RGA, RGA^{m2}, RGA^{m3} and RGA^{m4}. Left panel:

Immunodetection of RGA-GFP, RGA^{m2}-GFP, RGA^{m3}-GFP and RGA^{m4}-GFP proteins

respectively in *35S::RGA-GFP*, *35S::RGA^{m2}-GFP*, *35S::RGA^{m3}-GFP* and *35S::RGA^{m4}-GFP*

N. benthamiana agro-infiltrated leaves treated with 100 mM cycloheximide (CHX) and 100

μM GA$_3$ for the indicated times. Actin is used as sample loading control. Right panel:

Quantification of the immunoblot signals depicted as a graph. Values indicate RGA-GFP,

RGA^{m2}-GFP, RGA^{m3}-GFP and RGA^{m4}-GFP signals relative to Actin signals.

Extended Data Figure 2. RGA^{m2} loses its ability to bind with RGA-interacting partners.

Extended pairwise Y2H interaction assays between GAI, RGA and RGA^{m2} and with 91

DELLA-interacting transcription factors. The colour code and numbers (0 to 5) indicate the

strength of the interactions based on the size of the yeast colonies on selective media (without

leucine, tryptophan and adenine: -LWH). Empty pGBKT7 and pGADT7 vectors were

included as negative controls.

Figure 2. The qRGA^{mPFYR} sensors monitors changes in GA levels.

a, Schematic representation of the qRGA^{mPFYR} construct composed of two elements: the

sensor module (RGA^{mPFYR}-VENUS), and the normalization element (TagBFP fused with a

nuclear localization signal (NLS)). The two elements are linked by a 2A self-cleaving peptide,

and driven by the same promoter allowing stoichiometric expression. GA signalling activity is

measured as 3 minus the ratio between VENUS and TagBFP signal intensities. **b**, TagBFP

expression pattern of *pUBQ10::qRGA^{mPFYR}* and *pRPS5a::qRGA^{mPFYR}* sensors monitored in

cotyledon, vegetative SAM, hypocotyl and root of 7-day-old seedlings. **c**, Hypocotyl length of

wild-type (Col-0) and *pUBQ10::qRGA^{mPFYR}* seedlings. Upper panel: Hypocotyl length at 5

and 9 days after sowing (DAS). Lower panel: Hypocotyl length of 9-d-old seedlings grown

for 4 days on MS media and then transferred to MS media supplemented with 5 µM PAC or

10 µM GA$_3$, and controls (Mock). Different letters denote significant differences ($p < 0.05$)

using ANOVA Tukey's tests for multiple comparisons. **d**, Boxplot representations of the

number of rosette leaves (upper panel) and final plant height (lower panel) of wild-type (Col-

0) and *pRPS5a::qRGA^{mPFYR}* adult plants. *P* values are from Wilcoxon rank-sum tests. **e**,

Representative confocal images of RGA^{mPFYR}-VENUS and TagBFP signals in hypocotyls of

9-d-old *pUBQ10::qRGA^{mPFYR}* seedlings that have been treated with PAC or GA (and Mock

control), as in **c**. The *pRGA::GFP-RGA* transgenic line was used as a positive control of the

treatments. Intensity heatmap representation of GA signalling activity is shown with the same

colour code in the last vertical panel. **f**, Boxplot representation of GA signalling activity

(means of at least 8 seedlings) measured in seedling hypocotyls grown in the same conditions

as in **c** and shown in **e**. For statistical analysis, a Kruskal-Wallis test between MS/PAC and

MS/GA conditions was applied. The experiment was repeated twice with similar results. **g**,

Hypocotyl length measurement of wild-type (Col-0) and *pUBQ10::qRGA*^{mPFYR} seedling

hypocotyls grown either in dark or light conditions from day 2 to day 9 post sowing. **h**,

Boxplot representation of GA signalling activity measured in basal region of seedling

hypocotyls (means of 15 to 31 seedlings) grown in light or dark conditions, at 3, 5 and 7 DAS.

A Kruskal-Wallis test between light and dark-grown seedlings was applied. The experiment

was repeated twice with similar results.

Extended data Figure 3. Expression patterns of *pUBQ10::qRGA*^{mPFYR} and

***pRPS5a::qRGA*^{mPFYR} in the SAM and responses of *pRPS5a::qRGA*^{mPFYR} to changes in**

GA levels.

a-c, Fluorescence signals of RGA^{mPFYR}-VENUS (a), TagBFP-NLS (b) and merge channels (c)

in *pUBQ10::qRGA*^{mPFYR} inflorescence SAM. **d-f**, Fluorescence signals of RGA^{mFPYR}-VENUS

(d), TagBFP-NLS (e) and overlay (f) in *pRPS5a::qRGA*^{mPFYR} SAM. **g**, Representative

growing wild-type (Col-0) and *pRPS5a::qRGA*^{mPFYR} plants with flowers and elongated

siliques indicative of normal fertility (arrows). **h-p**, Fluorescence signals of RGA^{mPFYR}-

VENUS (h, m), TagBFP-NLS (i, n) and overlay (j, o) in *pRPS5a::qRGA*^{mPFYR} SAM, 2h after

a treatment with 100 μM GA$_3$ (h-j) and mock controls (m-o). (k, p) Heatmap representations

of GA signalling activity from (j, o). The older floral primordia were removed digitally. (l)

Boxplot representation of GA signalling activity in SAM treated as in (j, o). **q-u**,

Fluorescence signals of RGA^{mFPYR}-VENUS (q, v), TagBFP-NLS (r, w) and overlay (s, x) in

pRPS5a::qRGA^{mPFYR} SAM, 5 days after a treatment with 50 μM PAC (v-x) and mock (q-s). (t,

y) Heatmap representations of GA signalling activities from (s, x). Older floral primordia

were removed digitally. (u) Boxplot representation of GA signalling activity is SAM treated

as in (s, x). *P* values are from Wilcoxon rank-sum tests. Experiments were repeated twice

with similar results. Scale bars = 20 µm (a-f, h-j, m-o, q-s, v-x) and 1 mm (g).

Extended Data Figure 4. Behaviour of the different GA signalling sensor lines in

response to changes in GA levels in seedling root.

Representative VENUS and TagBFP signals obtained in seedling roots of qRGA^{mPFYR},

qRGA^{m3} and qRGA^{m4} lines driven with *pUBQ10* and *pRPS5a* promoters, treated with 5 µM

PAC or 10 µM GA$_3$ (and Mock controls), as in Figure 2. *pRGA::GFP-RGA* is used as a

positive control of GA and PAC treatments.

Extended data Figure 5. Dose and temporal responses of qRGA^{mPFYR} to exogenous GA

treatment in the SAM.

a-i, Overlay of VENUS and TagBFP maximum intensity projection (a-d) and corresponding

heatmap representation of GA signalling activity (e-h) of qRGA^{mPFYR} SAMs treated for 3 h

with 1 µM (b, f), 10 µM (c, g) and 100 µM (d, h) GA$_3$, and Mock controls (a, e). Boxplot

representation of GA signalling activity is shown in (i). **j-z**, Overlay of VENUS and TagBFP

maximum intensity projection (j-m, r-u) and corresponding heatmap representation of GA

signalling activity (n-q, v-y) of qRGA^{mPFYR} SAM treated with 100 µM GA$_3$ for 1 h (s, w), 2 h

(t, x) and 3 h (u, y), and mock controls (j-q). Boxplot representation of GA signalling activity

is shown in (z). *P* values are from Wilcoxon rank-sum tests. Experiments were repeated twice

with similar results. Scale bars = 20 µm.

Figure 3. The qRGA^{mPFYR} sensor signal depends on GA receptor activity.

a-c, Expression pattern of *GID1a*, *GID1b* and *GID1c* in shoots (upper panel) and root tips

(lower panel) of 7-d-old *pGID1a::GID1a-GUS* (a), *pGID1b::GID1b-GUS* (b) and *pGID1c::*

GID1c-GUS (c) seedlings. **d**, Boxplot representation of GA signalling activity in hypocotyls

of 5-d-old *pUBQ10::RGA^{mPFYR}* (pink) and *pUBQ10::RGA^{mPFYR} pUBQ10::GID1a-mCherry*

(cyan) seedlings treated with 100 μM GA_3 for 30, 90 and 180 min, and untreated controls. *P-*

values are from Kruskal-Wallis tests.

Extended data Figure 6. Expression patterns of GA receptors in the SAM.

a-c, Expression patterns of *GID1a*, *GID1b* and *GID1c* in the SAM of *pGID1a::GID1a-GUS*

(a), *pGID1b::GID1b-GUS* (b) and *pGID1c::GID1c-GUS* (c) adult plants. **d-l**, *In situ*

localization of *GID1a* (d-f), *GID1b* (g-i) and *GID1c* (j-l) mRNA in inflorescence apex. 3

consecutive sections are shown for each. **m**, Maximum intensity projection showing the

expression of *pGID1b::2xmTQ2-GID1b*. Scale bars = 200 μm (a-c) and 20 μm (d-m).

Figure 4. A GA signalling activity map reveals high levels of GA signalling activity in

inter-primordia cells in the SAM.

a, Maximum intensity projection showing the expression of *pRPS5a::qRGA^{mPFYR}* and

pCLV3::mCherry-NLS in the SAM (overlay). CZ, central zone; P, primordium. **b**, Heatmap

representation of L1 GA signalling activity averaged from seven SAMs aligned using the

CLV3 domain as a reference. **c-d**, Quantification of GA signalling activity in central zone

(CZ), inter-primordia region (IPR) and primordia (P) indicated by different colours as shown

in (**c**). *P* values are from one way ANOVA. **e,** 3D visualization of nlsGPS1 emission ratio in

the SAM. Primordia stages are estimated according to morphology. Arrows highlight higher

nlsGPS emission ratio in the boundaries and IPR. **f-g,** Fluorescence distribution in wild-type

(L*er*) SAMs treated with fluorescein (Fl, **f**) and GA_4-Fl (**g**). The inserts show the same images

merged with the PI channel that visualizes cell membranes (magenta). **h**, Comparison of the

ratio of average fluorescence intensity in the IPR to that in the non-IPR (excluding primordia)

between Fl and GA$_4$-Fl treatment in the SAM. *P* value is from a Wilcoxon rank-sum test. The

experiment was repeated twice with similar results. Scale bars = 20 µm.

Extended data Figure 7. Digital longitudinal sections through the centre of the SAM and

primordia of qRGA$^{\text{mPFYR}}$ plants.

a, Maximum intensity projection showing the expression of *pRPS5a::qRGA$^{\text{mPFYR}}$*

pCLV3::mCherry-NLS plants in the SAM as in Fig 4a. CZ, central zone. **b**, Signals for

qRGA$^{\text{mPFYR}}$-VENUS, TagBFP-NLS and mCherry-NLS in digital longitudinal sections

through the centre of CZ and primordia P$_1$ to P$_8$ along the white dotted lines indicated in (**a**).

Scale bars = 20 µm.

Extended data Figure 8. Response of nlsGPS1 to GA treatment in the SAM.

a-d, 3D visualization of nlsGPS1 expression in the SAM before (**a**, **c**) and after (**b**, **d**) mock (**a**,

b) and 10 µM GA$_{4+7}$ treatment (**c**, **d**). **e**, Boxplot representation of nlsGPS1 emission ratios

from nuclei analysed in three SAMs. The number of nuclei N is indicated. *P* value is from a

Wilcoxon rank-sum test. The experiment was repeated twice with similar results.

Extended data Figure 9. Fluorescence distribution in SAMs treated with Fl and GA$_3$-,

GA$_4$- and GA$_7$-Fl.

a-b, Fluorescence distribution in wild-type (L*er*) SAMs treated with GA$_3$-Fl (**a**) and GA$_7$-Fl

(**b**). The inserts show the same images overlaid with the PI channel that visualizes cell

membrane (magenta). **c**, Ratio of average fluorescence intensity in the IPR to that in the non-

IPR (excluding primordia) after GA$_3$-Fl and GA$_7$-Fl treatment in the SAM, compared to Fl

treatment (Fig 4f, h). *P* values are from Wilcoxon rank-sum tests. **d-g,** Digital longitudinal

sections through the centre of CZ and primordia P$_3$ to P$_7$ of L*er* SAMs treated with 5 µM

fluorescein (**d**), GA$_3$-Fl (**e**), GA$_4$-Fl (**f**), or GA$_7$-Fl (**g**) (green) and stained with PI (red). Scale

bars = 20 µm.

Figure 5. High GA signalling activity correlates positively with growth anisotropy and

transverse cell division orientation.

a-b, Averaged surfacic growth (**a**) and growth anisotropy (**b**) heat maps (used as proxies for

cell expansion intensity and direction, respectively) in the SAM. **c**, PCA analysis including

the following variables: GA signalling, surfacic growth intensity, surfacic growth anisotropy

and CLV3 expression. **d**, Spearman correlation study between GA signalling, surfacic growth

intensity and surfacic growth anisotropy in tissue scale but excluding CZ. **e**, 3D visualization

of Col-0 SAM L1 cells using confocal microscopy. New cell walls formed in the SAM (but

not primordia) in 10 h are coloured according to their angle values. The insert shows the

corresponding 3D image at 0 h. **f**, Boxplot representation of cell division frequency in the IPR

and non-IPR of Col-0 SAM. *P* value is from Wilcoxon rank-sum test. **g**, A schematic diagram

showing how the angle of a new cell wall (magenta) relative to the radial direction from the

centre of the SAM (white dotted line) was measured. Only acute angle values (i.e: 0-90°) are

considered. **h**, Frequency histograms of division plane orientation of cell from the entire SAM

(dark blue), the IPR (medium blue) and non-IPR (light blue), respectively. **i**, Frequency

histograms of division plane orientation of cells located in front and sides of P$_3$ (light green),

P$_4$ (mediate green) and P$_5$ (dark green), respectively. *P* values are from Kolmogorov-Smirnov

tests. The experiment was repeated twice with similar results.

1012

1013   **Extended Data Figure 10. Expression patterns of DELLA-encoding genes and DELLA**

1014   **proteins in the SAM.**

a-d, GUS signal in the shoot apex of *pGAI::GUS* (**a**), *pRGA::GUS* (**b**), *pRGL1::GUS* (**c**) and

rgl2-5 (a promoter trap GUS line) mutant (**d**). **e-h**, *In situ* hybridization serial sections through

a shoot apex using probes of GAI (**e**), RGL1 (**f**), RGL2 (**g**) and RGL3 (**h**). **i-j,** Intensity

heatmaps of maximum intensity-projection of *pRGL3::RGL3-GFP* (**i**) and *pRGA::GFP-RGA*

(**j**) in the SAM. Arrowheads mark the regions with lower GFP-RGA expression. Scale bars =

20 µm.

Figure 6. Cell division orientation distribution in the SAM are modified in global *della*

mutant and *pCUC2::gai-1-VENUS* transgenic plant.

a-b, 3D visualisation of the L1 layer of PI-stained L*er* (**a**) and global *della* mutant (**b**) SAM

using confocal microscopy. New cell walls formed in the SAM but not primordia in 10 h are

shown and coloured according to their angle values. Inserts show the SAM at 0 h. Arrows in

(**b**) highlight examples of aligned cell files in global *della* mutant. **c-e**, Comparison of

frequency distribution of division plane orientation of cells in the entire SAM (**d**), IPR (**e**) and

non-IPR (**f**) between L*er* and global *della*. *P* values are from Kolmogorov-Smirnov tests. The

experiment was repeated three times with similar results. **f-g**, 3D visualisation of confocal

image stacks of PI-stained SAMs of Col-0 (**i**) and *pCUC2::gai-1-VENUS* (**j**) transgenic plants.

New cell walls formed in the SAM but not primordia in 10 h are shown as in (**a,b**). **h-j**,

Comparison of frequency distribution of division plane orientation of cells located in the

entire SAM (h), IPR (i) and non-IPR (j) between Col-0 and *pCUC2::gai-1-VENUS*

transgenics. *P* values are from Kolmogorov-Smirnov tests. The experiment was repeated three

times with similar results.

Extended Data Figure 11. Analysis of SAM shape of global and quadruple *della* mutants

and *pCUC2::gai-1-VENUS* transgenic plants.

a-c, Maximum projection (**a-c**) and orthogonal views (XZ dimension, lower panels; YZ

dimension, left panels) of PI-stained SAMs of L*er* (**a**), global *della* (**b**) and quadruple *della*

mutant (**c**) using confocal microscopy. **d**, Comparison of SAM size between L*er*, global and

quadruple *della* mutants. *P* values are from Kruskal-Wallis tests. **e**,**f**, XZ point plots of L1

nuclei of L*er* (blue), global *della* (orange) and quadruple *della* (green) showing the tissue-

level curvature of the SAM. **g-j**, Heatmaps of cellular curvature of L1 cells in the SAM

(including P_4) of L*er* (**g**), global *della* (**h**) and quadruple *della* (**i**). Statistics are shown in (**j**). *P*

values are from Kruskal-Wallis tests. **k**, Maximum projection view of the SAM of

pCUC2::LSSmOrange-NLS line. The insert shows the same image superimposed with the

expression of *pRPS5a::TagBFP-NLS*. **l-m**, Maximum projection (**l-m**) and orthogonal views

(XZ dimension, lower panels; YZ dimension, left panels) of PI-stained SAM of *pCUC2::gai-*

1-VENUS (**m**) transgenic plant and its wild-type control (**l**). **n**, XZ point plots of L1 nuclei of

wild-type (blue) and *pCUC2::gai-1-VENUS* transgenic plants (yellow) showing the tissue-

level curvature of the SAM. **o-q**, Heatmaps of cellular curvature of L1 cells in the SAM of

wild-type (**o**) and *pCUC2::gai-1-VENUS* (**p**) transgenic plants. Statistics are shown in (**q**). *P*

value is from Wilcoxon rank-sum test. Scale bars = 20 μm.

1056

Extended Data Figure 12. Cell division plane orientation distribution in the SAM of

quadruple *della* mutants.

a-b, 3D visualization of the L1 of PI-stained SAM of quadruple *della* mutant using confocal

microscopy. New cell walls formed in the SAM (but not in primordia) in 10 h are shown and

coloured according to their angle values. (**a**) is representative of 6 out of 13 SAMs while (**b**)

represents the remaining 7 SAMs where a higher frequency of transverse cell divisions could

be detected. Inserts are images at 0 h. (**c**) Comparison of frequency distributions of division

plane orientation between cells located in the entire SAM of L*er* and quadruple *della* mutants.

P value is from Kolmogorov-Smirnov test.

1066

Figure 7. Emergence of the internode cellular organisation in the inter-primordia region

in wild-type and in plants with modified GA signalling activity.

a-l, Time-lapse (0 h, 24 h, 34 h) visualization of the L1 of the SAM from confocal

mciroscopy of L*er* (**a-c**), global *della* (**d-f**), Col-0 (**g-i**) and *pCUC2::gai-1-VENUS* transgenic

(**k-n**). The imaging time is indicated at the right up corner of each panel. Cells outlined with

yellow dotted lines are those incorporated into a primordium at 34 h, and cells in green are

found in the IPR at 34 h. Those in orange produce both primordium and IPR cells. White

dotted lines mark the edge of the elder primordium (that have been removed at 34h except in

(**c**)). **m-p,** Scanning electron microscopy images the shoot apex of L*er* (**m**), global *della* (**n**),

Col (**o**), and *pCUC2::gai-1-VENUS* (**p**) allowing to analyse the early establishment of

internode just below the SAM. The different size of arrows indicates different range of size of

the early internode in the different genetic background.

1079

1080   **Extended Data Figure 13. Altered plant height in *pCUC2::gai-1-VENUS* transgenic**

1081   **plants.**

1082   Representative 37-d-old wild-type (**a**) and 56-d-old *pCUC2::gai-1-VENUS* transgenic plants

1083   (**b**) grown under long-day conditions. Scale bars = 1 cm.

1084

1085   **Extended Data Figure 14. Hypothesis of the role of GA signalling in internode**

1086   **specification.**

1087   Higher levels of GA signalling activity in the IPR specify the cellular organization of

1088   internode in the SAM via a regulation on cell division orientation.

1089

1090

1091

Acknowledgments

We thank the colleagues of the Laboratoire Reproduction et Développement des Plantes for fruitful discussions on this work and for sharing material and advice. We also thank Miguel Perez-Amador, Jan Lohmann, Yvon Jaillais, Tai-ping Sun, Miguel Blazquez, David Alabadi and the NASC for seeds and plasmids. We acknowledge the contribution of SFR Biosciences (UMS3444/CNRS, US8/Inserm, ENS de Lyon, UCBL) facility PLATIM for assistance with microscopy and image analysis. This work was supported by the ANR-16-CE13-0014 (GrowthDynamics) grant to T.V. and P.A; a European Research Council grant (GAtransport) to R.W..

Author contribution

P.A. and T.V. designed the study and supervised the work; B.S., A.F.-B, P.A. and T.V. designed the experiments. B.S., A.W. and A.M. performed the FRET analysis; A.N.-G. and S.P. conducted the large-scale two-hybrid analysis; S.L. and R.W. designed the synthesis strategy and synthesized the GA-fl fluorescent probes. G.C. developed the pipeline for quantitative image analysis with the help of J.L.; B.S., A.F.-B., and G.C. performed the image analysis. B.S., A.F.-B., C.G.-A., L.J., G.B., J.M., L.S.-A. and J.-M.D. performed all the other experiments; all authors were involved in data analysis; B.S., A.F.-B., P.A. and T.V. wrote the manuscript with inputs from all authors.

Financial competing interests

The authors declare no competing interests.

1115   **Corresponding authors**

1116   Correspondence and request for materials should be addressed to T.V. (teva.vernoux@ens-

1117   lyon.fr) or P.A. (patrick.achard@ibmp-cnrs.unistra.fr).

1118

Data and software availability

All experimental data and quantified data that support the findings of this study are available

from the corresponding authors upon request. Quantitative image and geometry analysis

algorithms are provided in Python libraries timagetk, cellcomplex, tissue_nukem_3d and

sam_atlas (https://gitlab.inria.fr/mosaic/) made publicly available under the CECILL-C

license. The script used to process hypocotyl images is publicly available in the

qrga_nuclei_quantification project (https://gitlab.inria.fr/gcerutti/). The pipelines used to

analyze SAM image sequences and to produce map visualizations are provided in a separate

project (https://gitlab.inria.fr/mosaic/publications/sam_spaghetti) as Python scripts.

Fig. 1

a

c

b

d

Extended Data Fig. 1

Intensity of the interaction — 0 1 2 3 4

Plate-Position in MINILIBRARY	AGI	TF Name	Interaction in - LWH			
			GAI	RGA	RGAm2	ø
002-A01	At2g31380	STH	4	0	0	0
002-A02	At5g25810	ERF040/TINY	4	0	0	0
002-A03	At1g27050	HB54	4	0	0	0
002-A04	At5g66700	HB-8/HB-53	4	0	1	0
002-A05	At2g31380	STH	4	0	0	0
002-A06	At2g28910	CXIP4	4	0	0	0
002-A07	At4g21040	DOF4.3	4	0	0	0
002-A08	At4g01350	DC1	4	0	0	0
002-A09	At4g01350	DC1	4	0	0	0
002-A10	At2g02080	ATIDD4	4	4	0	0
002-A11	At4g03250	HDZIP	4	0	0	0
002-A12	At4g03250	HDZIP	4	0	0	0
002-B01	At4g17460	HAT1	4	0	0	0
002-B02	At3g04420	ANAC048	4	1	0	0
002-B03	At1g73100	SUVH3	3	0	0	0
002-B04	At3g02310	AGL4/SEP2	4	0	0	0
002-B05	At1g17590	NFYA8	4	0	0	0
002-B06	At1g77920	bZIP50/TGA7	3	0	0	0
002-B07	At5g06839	bZIP65/TGA10	4	0	0	0
002-B08	At1g25340	MYB116	3	0	0	0
002-B09	At5g35550	TT2/MYB123	4	0	0	0
002-B10	At4g22680	MYB85	4	0	1	0
002-B11	At5g62320	MYBCU15/MYB99	4	0	0	0
002-B12	At5g62320	MYBCU15/MYB99	4	0	0	0
002-C01	At1g17770	SUVH7/SDG17/SET17	4	0	0	0
002-C02	At1g09540	MYB61	4	4	0	0
002-C03	At2g36890	BIT1/MYB38/RAX2	4	0	0	0
002-C04	At5g12870	MYB46	4	1	0	0
002-C05	At3g09600	RVE8/LCL5	4	0	0	0
002-C06	At3g09600	RVE8/LCL5	3	1	0	0
002-C07	At4g24240	WRKY7	4	4	0	0
002-C08	At4g22070	WRKY31	4	3	0	0
002-C09	At1g69810	WRKY36	4	1	0	0
002-C10	At5g24520	TTG1	4	1	0	0
002-C11	At3g47620	TCP14	4	1	0	0
002-C12	At1g69690	TCP15	4	0	0	0
002-D01	At1g31630	AGL86	4	0	0	0
002-D02	At5g26650	AGL36	4	1	0	0
002-D03	At1g22590	AGL87	4	0	0	0
002-D04	At2g24430	ANAC038	4	1	0	0
002-D05	At2g45050	GATA2	3	0	0	0
002-D06	At3g21175	GATA24/ZML1	4	0	0	0
002-D07	At1g24625	ZFP7	4	2	0	0
002-D08	At2g41940	ZFP8	4	2	0	0
002-D09	At4g16610	C2H2	3	0	0	0
002-D10	At5g43700	AUX2-11/IAA4	3	0	0	0
002-D11	At3g23050	IAA7	3	0	0	0
002-D12	At3g02150	TCP13	4	3	0	0
002-E01	At2g24430	ANAC038	4	0	0	0
002-E02	At2g45050	GATA2	4	0	0	0
002-E03	At1g70000	myb-type HTH	4	0	0	0
002-E04	At3g20310	ERF083/ERF7	4	3	1	1
002-E05	At5g18560	PUCHI/ERF086	4	0	0	0
002-E06	At1g49480	REM19/RTV1	4	1	1	0
002-E07	At1g14600	Homeodomain/G2-like?	4	0	1	0
002-E08	At5g16560	KAN1	4	1	0	0
002-E09	At5g05090	Homeodomain/G2-like?	4	2	0	0
002-E10	At1g51140	bHLH122/EN70/ Flowering bHLH3/FBH3	4	4	0	0
002-E11	At2g42280	Flowering bHLH4/ FBH4	4	0	0	0
002-E12	At1g61660	bHLH112/EN64	4	3	0	0
002-F01	At5g64750	ABR1/ERF111	4	1	0	0
002-F02	At5g54070	HSFA9	4	0	0	0
002-F03	At5g09460	bHLH143/EN129/AJAX3	4	0	0	0
002-F04	At1g19210	ERF017	4	0	0	0
002-F05	At5g17690	LHP1/TFL2	3	1	0	0
002-F06	At4g13980	HSF12/HSFA5	4	1	0	0
002-F07	At2g01760	ARR14	3	0	0	0
002-F08	At2g31220	bHLH10/EN23	4	0	0	0
002-F09	At4g02590	bHLH59/UNE12	4	0	0	0
002-F10	At4g30935	WRKY32	4	4	1	0
002-F11	At1g68920	bHLH49/EN82	4	1	0	0
002-F12	At2g40340	ERF48/DREB2C	4	0	0	0
002-G01	At3g09600	RVE8/LCL5	4	0	0	0
002-G02	At1g51600	GATA28/ZML2	4	0	0	0
002-G03	At3g29035	NAC3/ANAC059/ORS1	4	3	0	0
002-G04	At5g66700	HB-8/HB-53	3	0	0	0
002-G05	At4g37750	ANT	4	3	0	0
002-G06	At5g60120	TOE2	4	1	0	0
002-G07	At1g43160	RAP2-6/ERF108	4	0	0	0
002-G08	At1g77200	ERF037	4	1	0	0
002-G09	At3g57600	DREB2F	4	0	0	0
002-G10	At2g28910	CXIP4	4	0	0	0
002-G11	At2g28910	CXIP4	4	0	0	0
002-G12	At1g31630	AGL86	4	0	0	1
002-H01	At5g26650	AGL36	4	0	0	0
002-H02	At1g69690	TCP15	4	1	0	0
002-H03	At3g47620	TCP14	4	0	0	0
002-H04	At5g29000	PHL1	4	0	0	0
002-H05	No gene	Control	4	0	0	0
002-H06	At1g09530	bHLH8/EN100/PIF3	4	4	0	0
002-H07	At2g43010	SRL2/PIF4	4	0	0	0
002-H08	At2g33880	HB-3/STIP/WOX9A	4	3	0	0

Extended Data Fig. 2

Fig. 2

Extended data Fig. 3

Extended Data Fig. 4

Extended data Fig. 5

Fig. 3

Extended data Fig. 6

Fig. 4

a
P₆
P₃
P₁
P₈
CZ
P₄
P₅
P₂
P₇
pRPS5a::RGA^mPFYR-VENUS
pCLV3::mCherry-NLS
pRPS5a::TagBFP-NLS

b
P₁
P₂
P₃
P₄
P₅
P₆
P₇
P₈

Extended data Fig. 7

Extended data Fig. 8

Extended data Fig. 9

a Surfacic growth intensity

b Surfacic growth anisotropy

c PCA Component 2 (38.82%)
Surfacic growth intensity
Surfacic growth anisotropy
GA signalling
CLV3 expression
PCA Component 1 (38.82%)

d Map correlations (Spearman)
All zones except CZ

GA signalling
Surfacic growth intensity — *** -0.86
Surfacic growth anisotropy — *** -0.08 / *** 0.19

e P₅ P₂ P₃ P₄ P₁ P₆ Col 10 h 0 h 0° 90°

f Percentage of cell division — IPR, Non-IPR, $P = 0.7959$

g SAM centre

h Frequency (%)
SAM (N = 408, 10 SAMs)
IPR (N = 163, 10 SAMs)
Non-IPR (N = 245, 10 SAMs)
$P < 0.01$
$P < 0.01$
Angle (°)

i Frequency (%)
P3 (N = 36, 10 SAMs)
P4 (N = 53, 10 SAMs)
P5 (N = 40, 10 SAMs)
$P = 0.155$
$P = 0.043$
Angle (°)

Fig. 5

a   *pGAI::GUS*

b   *pRGA::GUS*

c   *pRGL1::GUS*

d   *rgl2*

e   0 μm   8 μm   16 μm   24 μm

P$_5$ P$_5$ P$_1$ I$_1$ I$_1$ P$_5$ I$_1$

GAI *in situ*

f   0 μm   8 μm   16 μm

RGL1 *in situ*

i   255   0   *pRGL3::RGL3-GFP*

g   0 μm   8 μm   16 μm

RGL2 *in situ*

j   255   0   P$_4$ P$_3$ P$_2$ P$_5$ *pRGA::GFP-RGA*

h   0 μm   8 μm   16 μm

RGL3 *in situ*

Extended Data Fig. 10

Fig. 6

Extended Data Fig. 12

Fig. 7

a Wild-type

b *pCUC2::gai-1-VENUS*

Extended Data Fig. 13

Higher GA signalling
activity in the IPR

Preferential circumferential
cell division orientation

Specification of the internode
cellular organization

Extended Data Fig. 14