The activity and functions of subarctic soil microbial communities vary across vegetation types

Viitamäki Sirja[1*], Pessi Igor S.[1,2], Virkkala Anna-Maria[3,4], Niittynen Pekka[3], Julia Kemppinen[5], Eronen-Rasimus Eeva[1, 6], Luoto Miska[2,3] & Hultman Jenni[1,2*]

[1] Department of Microbiology, University of Helsinki, Finland

[2] Helsinki Institute of Sustainability Science (HELSUS), Helsinki, Finland.

[3] Department of Geosciences and Geography, University of Helsinki, Finland

4 Woodwell Climate Research Center, Falmouth, Massachusetts, USA

[5] Geography Research Unit, University of Oulu, Finland

[6] Finnish Environment Institute, Marine Research Centre, Helsinki, Finland

* Correspondence:

Corresponding Authors

sirja.viitamaki@helsinki.fi

jenni.hultman@helsinki.fi

Keywords: microbial communities, transcriptomics, microbial ecology, tundra, climate change

Abstract

Global warming changes the activity of soil microbial communities in high latitudes, which might result in higher greenhouse gas emissions. However, these microbial processes involved in GHG production and consumption are not thoroughly understood. We analyzed 116 soil metatranscriptomes from 73 tundra sites and investigated how bacterial and archaeal communities and their functions vary horizontally (i.e. vegetation type) and vertically (i.e. topsoil organic and mineral layers) during the summer season, in soil types that differed in pH, moisture, soil organic matter (SOM), and carbon and nitrogen content. Active microbial communities were significantly different in the organic and mineral soil layers. Additionally, the communities differed significantly between the different vegetation types both in the organic and mineral layers. Various plant polymer degraders were particularly active in shrub-dominated ecosystems with high SOM and low pH, less known mixotrophic groups (such as Chloroflexi) were active lower SOM and higher pH. Additionally, we detected transcripts of alphaproteobacterial methanothrophs, which potentially moderate methane release from tundra soils in deeper soil layer. Our results provide new insights into the diversity and activity of microbial communities of high latitudes under climate change.

1   Introduction

The Arctic is experiencing some of the most rapid and severe climate change on Earth (IPCC 2018). Over the next 100 years, climate change is expected to accelerate, contributing to major ecological changes (Post et al. 2019). Tundra soils harbor large carbon (C) stocks that are susceptible to microbial degradation with rising global temperatures (Johnston et al. 2019, Mackelprang *et al*. 2011). Soil microorganisms are critical in C and nutrient cycling in terrestrial ecosystems (Falkowski et al. 2008). Therefore, understanding soil microbial communities and how they correspond to local environmental variation is essential to predict the impacts of environmental change on ecosystem functioning (Buckeridge et al. 2013, Mod et al. 2021, Virkkala et al. 2018).

Warming and elevated atmospheric carbon dioxide (CO_2) concentrations stimulate plant productivity and can thereby increase the net carbon sink of arctic ecosystems (Friend *et al.*, 2014; Berner *et al.*, 2020). However, warmer temperatures also increase soil decomposition (Bond-Lamberty *et al.*, 2018) and plant productivity, and this provides substrates for microbes, which subsequently stimulate decomposition and the release of CO_2 (priming effect, reviewed in Jansson & Hofmockel, 2020). Although GHG emissions from soil are measurable (Bond-Lamberty & Thomson, 2010), soil microbial communities and their functions have not been explored in detail

due to technical limitations and vast diversity. This in turn hinders a comprehensive understanding

of the net contribution of microbes and their feedback to climate change.

In high-latitude soils, microbial communities are exposed to extreme environmental stressors,

such as fluctuating temperatures, long periods of darkness, and intensive UV radiation and drought.

In addition, vegetation is typically heterogenous in areas, where the local growing conditions

(microclimate, soil moisture, soil nutrients) can vary greatly at small spatial scales (le Roux *et al.*,

2013; Kemppinen *et al.*, 2021). Different vegetation types affect soil biotic and abiotic factors,

through which vegetation can influence the local soil microbial community. For example, tundra

soils are relatively acidic (pH 4-6; Hobbie & Gough, 2002; Männistö *et al.*, 2007) and soil acidity is

generally one of the most predominant drivers of the composition and activity of microbial

communities (Chu *et al.*, 2010). In addition, the quantity and quality of soil organic matter (SOM)

and nitrogen (N) availability and the C/N ratio affect microbial processes and primary production in

Arctic tundra soils (Koyama *et al.*, 2014, Zhang *et al.*, 2014).

Soil microbial communities are taxonomically and functionally diverse. Unlike higher

organisms, microbial communities in Arctic soils can be as diverse as in other soil biomes across

the globe (Neufeld & Mohn 2005; Chu *et al.* 2010). Recent studies have revealed Acidobacterial

predominance in acidic soils and stability of microbial community composition over seasons in

subarctic tundra (Männistö *et al.*, 2007; Männistö *et al.*, 2009), suggesting that Acidobacteria may

also have a significant role in community functions. In addition, recent research has shown

methanogenesis and methanotroph activity in peatlands experiencing permafrost thaw (McCalley *et

al.*, 2014; Singleton *et al.*, 2018; Woodcroft *et al.*, 2018). Previous studies on high-Arctic and

permafrost soil microbial communities have also shown activity of diverse organic matter

decomposition, methanogenesis, and methanotrophy (eg. Hultman *et al.*, 2015; Schostag *et al.*,

2015; Tveit *et al.*, 2015). However, soil microbial studies rarely compare multiple vegetation types

or investigate drier upland tundra, which is a noteworthy ecosystem that covers ca. 90% of the

Arctic (Raynolds *et al.*, 2019). In general, these studies were performed with small sample sizes

(Hultman *et al.*, 2015; Schostag *et al.*, 2015; Singleton *et al.*, 2018; Woodcroft *et al.*, 2018),

limiting our understanding of the spatial variability in microbial communities. Further, several

molecular methods have been used, including phospholipid fatty acid (PLFA), terminal restriction

fragment analysis (T-RFLP) (Männistö *et al.*, 2007) and metagenomics (Pessi *et al.* 2020). In

metatranscriptomics the target molecule is RNA, allowing analysis of both function and activity of

organisms at the time of sampling. However, transcriptomics analyses have not yet been fully

explored. Therefore, transcriptomics provides a new tool to thoroughly examine the tundra

microbial community composition and functions.

Using a metatransctriptomics approach targeting total RNA, we investigated soil bacterial and

archaeal diversity and activity. We analyzed 73 sites during the active growing season in a

mountain tundra ecosystem. The area covers a wide range of microclimatic and soil nutrient

conditions and different vegetation types. The samples represent four different tundra vegetation

types (namely barren tundra, deciduous shrub, evergreen shrub and meadow; Figure 1), and both

organic and mineral layers when available.

91   2   Materials and methods

92   2.1 Study setting and sampling

The study area is located in Kilpisjärvi, northwestern Finland, and extends to the Scandinavian

Mountains. The 3 km^2 area covers parts of two mountains, Mount Saana (69°02′N, 20°51′E) and

Mount Korkea-Jehkas (69°04′N, 20°79′E) and the valley between them. The elevation range in the

study area is 320 m, with the highest point on Mount Saana at 903 m a.s.l. The study area is

topographically heterogeneous and part of the arctic-alpine tundra biome. Consequently, the area

has relatively broad environmental gradients of microclimate, soil moisture, and soil pH, for

example (Kemppinen *et al.*, 2021). The vegetation type is mainly mountain heath dominated by

dwarf shrubs, such as *Empetrum nigrum* and *Betula nana*, and to a lesser extent by *Juniperus*

communis, Vaccinium vitis-idaea, *V. uliginosum,* and *V. myrtillus* (Kemppinen *et al.*, 2019).

However, due to the fine-scale environmental variation and broad gradients, the landscape forms a

mosaic of different vegetation types, as both vegetation cover and plant species composition can

vary over very short distances (le Roux *et al.*, 2013). The soils in the area are mostly poorly

developed leptosols with shallow organic layers and occasional podzolization; however, the

meadows have soils with thicker organic layers. Permafrost is absent from soils but can be found in

the bedrock above 800 m a.s.l. (King and Seppälä 1987). The average air temperature and

precipitation sum in July measured at the Kilpisjärvi meteorological station (69°05′N 20°79′E,

480 m a.s.l.) for the period 1981-2010 is 11.2°C and 73 mm, respectively (Pirinen et al. 2012).

Samples were collected in July 2017 from 73 sites (Figure 1). Sampling sites consisted of

four vegetation types, which were classified using a similar system as in the Circumpolar Arctic

Vegetation Map, namely barren soil, deciduous shrub, evergreen shrub, and meadow (Walker *et al.*

2005). All sampling equipment was disinfected with 70% ethanol before and between samples to

avoid contamination. The soil was bored with a 50-mm diameter stainless-steel soil corer with a

plastic inner casing. When available, both organic and mineral layer sub-samples were collected.

The samples were collected below the rhizosphere, with a 5-cm target depth for the organic layer

sample, whereas the mineral layer sample was taken from the lowest part of the core from 10-15

cm. The samples were placed in a Whirl-Pak sampling bag (Nasco, Fort Atkinson, WI, USA) with a

metal spoon and immediately frozen on dry ice and kept frozen at -80°C until nucleic acid

extraction. Samples were collected from 73 sites, from which 116 metatranscriptomes were

sequenced.

2.2 Soil physicochemical data

For analysis of soil physicochemical properties, approximately 0.2 dm^3 of soil was collected with a

steel cylinder (diameter 5.5 cm, height 7.5) and stored at 4 °C. The soils were lyophilized according

to the Finnish standard SFS300 and pH analysis was performed according to international standard

ISO 10390. SOM content was analyzed by loss on ignition analysis according to standard SFS

3008. CNS analysis (carbon, nitrogen, and sulfur) was performed with a Vario Micro Cube

(Elementar, Langenselbold, Germany). Mineral samples were sieved through a 2-mm plastic sieve

and the organic samples for the CNS analysis were homogenized by hammering the material into

smaller pieces.

2.3 Nucleic acid extraction

Three replicate DNA/RNA extractions were performed for each sample. Samples were kept on ice

during weighing and extraction and all steps were performed promptly with nuclease-free labware

to avoid RNA degradation. Solutions and water were treated with 0.1% diethylpyrocarbonate.

Nucleic acids were extracted using a modified hexadecyltrimethylammonium bromide (CTAB),

phenol-chloroform, and bead-beating protocol (Griffiths *et al.*, 2000; DeAngelis *et al.*, 2009). On

dry ice, approximately 0.5 g of frozen soil was transferred to a 2-mL Lysing Matrix E tube (MP

Biomedicals, Heidelberg, Germany) and 0.5 mL of CTAB buffer (consisting of equal amounts of

10% CTAB in 1 M sodium chloride and 0.5 M phosphate buffer in 1 M NaCl), 50 μl 0.1 M

ammonium aluminum sulfate ($NH_4(SO_4)_2 \cdot 12 H_2O$), and 0.5 mL phenol:chloroform:isoamyl

alcohol (25:24:1) was added. After bead-beating with FastPrep (MP Biomedicals, Heidelberg,

Germany) at 5.5 m s^{-1} for 30 s, 0.5 mL chloroform was added. Nucleic acids were precipitated with

polyethylene glycol 6000 (PEG6000, 30% in 1.6 M NaCl) and washed with ethanol. Nucleic acids

were once again extracted from the leftover soil pellet to maximize RNA. Nucleic acids were

resuspended in 25 μl of Buffer EB and 250 μl Buffer RLT with β-mercaptoethanol added. Buffers

EB and RLT were from an AllPrep DNA/RNA Mini Kit (Qiagen, Hilden, Germany). All

centrifugations were performed at 4°C. Finally, RNA and DNA were purified with AllPrep

DNA/RNA Mini Kit, where RNA was treated with DNAse I (Qiagen, Hilden, Germany). The

amount and integrity of RNA were assessed on a Bioanalyzer RNA 2100 Nano/Pico Chip with

Total RNA Assay (Agilent, Santa Clara, CA, USA). To ensure that RNA was DNA-free, a check-

up PCR with universal primers and gel electrophoresis was performed. Triplicates were pooled by

combining equal amounts of RNA from each triplicate.

2.4 Sequencing

Complementary DNA (cDNA) libraries were constructed from the pooled RNAs with Ultra II RNA

Library Prep Kit for Illumina (New England Biolabs, Ipswich, MA, USA). cDNA concentrations

were measured using a Qubit fluorometer with a dsDNA BR/HS kit (Invitrogen, Carlsbad, CA,

USA). Before sequencing, the libraries were analyzed with a Fragment analyzer (Advanced

Analytical, Ames, IA) and small cDNA fragments were removed to avoid primer binding to the

flowcell and to reduce cluster density. Single-end sequencing was performed on an Illumina

NextSeq 500 (Illumina, San Diego, CA, USA) with 150 cycles at the Institute of Biotechnology,

Helsinki, Finland.

2.5 Data processing and analysis

As the total RNA approach was used (Urich *et al.*, 2008), ribosomal RNA was not depleted after

extraction and RNA transcripts were sequenced. One sample (site 11202, organic layer) yielded

80.4 million reads and it was reduced randomly with seqtk (Li 2016) with random seed 100 to a 4-

million read subset. The quality of the sequences was assessed using default settings with FastQC

(v. 0.11.5, Andrews 2010) and MultiQC (v. 1.3, (Ewels *et al.*, 2016)) before and after trimming.

Trimming and quality filtering was performed with Cutadapt (v. 1.10, (Martin, 2011)) with quality

cutoff 25 and minimum adapter overlap 10. A mineral layer sample from site 11210 was discarded

due to a low number of reads. Metaxa2 (v. 2.1.3, (Bengtsson-Palme *et al.*, 2018)) was used to

identify reads mapping to the small subunit (SSU) rRNA. These were then classified using the

mothur (v. 1.40.5, (Schloss *et al.*, 2009)) classify.seqs command with a confidence cutoff of 60%

and SILVA release 132 (Quast *et al.*, 2013) as the reference database. The taxonomy of abundant

taxa was manually updated according to the Genome Taxonomy database (Parks *et al.*, 2018; Parks

et al., 2020). Reads were mapped onto the Kyoto Encyclopedia of Genes and Genomes (KEGG)

Prokaryote database release 86 (Kanehisa & Goto, 2000) using DIAMOND blastx (v. 2.1.3,

Buchfink *et al.* 2015) with an E-value of 0.001. Blast results were parsed with KEGG-tools (v. 1.3,

Pessi 2020). Briefly, the KEGG orthology (KO) identifier of the best BLAST hit was assigned to

each read and mapped to the KEGG module hierarchy, and spurious pathways were removed using MinPath (v. 1.4, (Ye & Doak, 2009). Since genes for methane and ammonia oxidation are not distinguished in KEGG classification we performed BLAST (Altschul *et al.*, 1990) for these genes against a manually curated database of eight *pmoA* and eight *amoA* genes from bacteria and archaea.

Due to differences in read numbers per sample, the metatranscriptome data were subsampled for statistical analyses of protein-encoding genes to avoid skewing caused by the low number of observations in low-read samples. After subsampling, the number of samples represented in each vegetation type was 6 (barren), 7 deciduous shrub, 18 (evergreen shrub) in the mineral layer and 5 (barren), 6 (deciduous shrub), 19 (evergreen shrub), and 8 (meadow) in the organic layer. For the mineral layer, meadow samples were discarded from statistical analyses due to a low sample number (n=4) after subsampling. Statistical analyses and visualization were performed with R (v. 3.6.2, R Core Team 2020). Differences between microbial communities and functional genes among vegetation types (Table 1) were tested with permutational multivariate analysis of variance using distance matrices (PERMANOVA, function adonis, R-package vegan v. 2.5-6, Oksanen et al. 2019) and pairwise permutation MANOVA (functions pairwise.perm.manova R-package RVAideMemoire v.0.9-78 (Hervé, 2021) and vegdist, R-package vegan v. 2.5-6). For the statistical analyses of functional genes, samples were subsampled from 116 to 73 samples to avoid skewing caused by the low number of observations, using a limit of 9654 KEGG hits (function rrarefy, R-package vegan v. 2.5-6). Differences between vegetation types for bacterial and archaeal genera and between soil types for selected genes were tested with one-way analysis of variance (ANOVA, functions lm and aov, R-core package) and pairwise t-test (function pairwise.t.test, R-core package). Soil physicochemical properties were tested with Kruskal-Wallis test and pairwise Wilcoxon test with Bonferroni correction (functions kruskal.test and pairwise.wilcox.test, R-core package). Differences in microbial communities in metagenomic and metatranscriptomic datasets were visualized in principal coordinates analysis plot, using Bray-Curtis dissimilarities (function ordinate, Phyloseq v. 1.30.0 (McMurdie & Holmes, 2013).

Table 1. Number of samples sequenced from each of the vegetation type in the study area.

Soil layer	Barren	Deciduous shrub	Evergreen shrub	Meadow	**Total**
Organic	6	14	37	12	**69**
Mineral	7	10	23	7	**47**

209   **2.6 Data availability**

210   Sequences have been deposited to the European Nucleotide Archive under accession number
211   PRJEB45463.

3 Results

3.1 Soil conditions across vegetation types

We observed high variability in soil conditions between organic and mineral layers and across
vegetation types. Among all samples collected, SOM content varied from 2% to 92%, gravimetric
soil water concentration from 9% to 432%, and pH from 3.9 to 5.8 (Figure 1). Soil properties
differed between soil layers (Kruskal-Wallis test; $P < 0.01$, Figure 1). The organic layer in the
meadow sites was less acidic (Pairwise Wilcoxon test; $P < 0.05$) and contained less SOM than the
organic layers in sites with deciduous or evergreen shrubs ($P < 0.05$). Sites with meadow vegetation
also had a lower C/N ratio than sites with deciduous or evergreen shrubs or barren soil ($P < 0.01$).
The physicochemical properties of the mineral layer were not significantly different between
vegetation types. In mineral layer pH was overall higher than in the organic layer whereas SOM,
N%, C/N and gravimetric soil water concentration were lower.

Figure 1. a) Map of sampling sites. Saana fell is located in the southern part of the study area. Typical vegetation in meadow (b), evergreen shrub (c), deciduous shrub (d), and barren (e) sites. f) Soil physicochemical properties (pH, soil organic matter (SOM), nitrogen (N), carbon-nitrogen ratio (C/N), and gravimetric water content) in each vegetation type. Categories with the same letter are not statistically different. One outlier from mineral C/N plot was removed.

3.2 Microbial community composition across sites and vegetation types

We analyzed 281.1 million sequence reads obtained with a total RNA approach (Urich *et al.*, 2008) from 116 samples (Supplementary table S1). First, we analyzed the small subunit ribosomal RNA (SSU rRNAs) from the metatranscriptomes. SSU rRNA represented $35 \pm 2\%$ (mean $\pm$ standard deviation) of the reads. On the domain level, over 80% of the assigned SSU rRNAs in the total metatranscriptomes were bacterial, 0.1 % Archaeal, and 19% were of eukaryotic origin, with fungal SSU rRNA representing 12% of the sequences. Out of the identified fungal phyla, Ascomycota represented 8.2% and Basidiomycota 2.8%. Furthermore, 0.2% of the sequences in both soils were recognized as SSU rRNA but could not be assigned to the domain level.

Figure 2. Order-level bacterial abundances in metatranscriptomes (RNA) and metagenomes (DNA) in organic (upper panel) and mineral (lower panel) layers. Samples of the same vegetation type were pooled and unclassified taxa were removed.

According to the analysis of bacterial 16S rRNA, the predominant active phyla in the microbial communities were Actinobacteria ($27 \pm 9\%$) with order Acidothermales and Solirubrobacterales; Acidobacteria ($17 \pm 3\%$) with orders Acidobacteriales and Solibacterales; Alphaproteobacteria ($16 \pm 3\%$) with orders Rhizobiales and Acetobacterales, and Planctomycetes

(14 ± 3%) with order Gemmatales (Figure 2). Moreover, Deltaproteobacteria, Chloroflexi,

Gammaproteobacteria, and Verrucomicrobia were also active in these soils. Out of the Archaeal

reads Thaumarchaea (0.12 ± 0.15%) were most active, with Group 1.1c (0.05 ± 0.05%) and

Nitrososphaerales (0.05 ± 0.11%) being the most active taxa. The relative abundances at the genus

level in each vegetation type and soil layer are listed in Supplementary table S2.

Figure 3. a) Genus-level classified square root transformed relative abundances of the 100 most

abundant bacterial genera for each vegetation type and each site, and b) Square root transformed

Archaeal taxa with mean relative abundance > 0.001, lowest rank classification level shown. A

column of pixels represents a site.

257   The most active genera were *Acidothermus* (Class Actinobacteria, 13 ± 5.7%) and

Candidatus Solibacter (Acidobacteriia, 3.11 ± 0.83%), followed by *Bryobacter* (Acidobacteriia,

2.06 ± 0.38%) *Pajaroellobacter* (Deltaproteobacteria, 1.64 ± 0.55%), and *Roseiarcus*

(Alphaproteobacteria, 1.48 ± 1.07%, Figure 3a). Archaea were not particularly abundant in the

metatranscriptomes of these soils, but the most active archaeal groups belonging to Thaumarchaea

were more active in the mineral than organic layer (Figure 3b).

When comparing the 16S rRNA from metatranscriptomes (i.e., RNA; active microbes) to

the 16S rRNA genes from metagenomes (i.e., DNA, Pessi *et al.*, 2020), the most abundant and

active microbial groups were the same but notable differences appeared among the datasets (Figures

2 and 4a). Orders with greater relative abundance in metatranscriptomes included Gemmatales

(Planctomycetes)**,** Acidothermales (Actinobacteria), Solibacterales (Acidobacteria), and

Myxococcales (Deltaproteobacteria), whereas Subgroup 2 (Acidobacteria), Rickettsiales

(Alphaproteobacteria), Dormibacterota, Chitinophagales (Verrucomicrobia), and

Gemmatimonadales (Gemmatimonadetes) showed higher relative abundance in metagenomes. In

metatranscriptomes, the orders with relative abundance more than 1% covered $78 \pm 0.02\%$ of active

orders, but only $64 \pm 0.02\%$ in metagenomes. Additionally, more than 1500 bacterial and archaeal

genera were identified from metatranscriptomes and less than 500 bacterial and archaeal genera

from metagenomes.

Figure 4. a. Principal coordinates analysis plot showing differences between 16S rRNA genes from

metagenomes and 16S rRNA from metatranscriptomes among vegetation types and soil layers,

based on square root-transformed Bray-Curtis dissimilarities. b. Ordination plot showing the

differences in active microbial communities through 16S rRNA genes in organic and mineral layer.

Active microbial communities were different in organic and mineral layers (PERMANOVA; $R^2 =$

0.06642; $P < 0.001$, Figure 4b) and therefore further analyses were conducted separately for each

layer. Interestingly, the communities also differed significantly between the four different

vegetation types both in organic (PERMANOVA; $R^2 = 0.15866$; $P < 0.001$) and in mineral layer

(PERMANOVA; $R^2 = 0.13057$; $P < 0.01$). The communities in the organic layer were significantly

different at the meadow site compared to the deciduous or evergreen shrub sites (pairwise

PERMANOVA, *P* <0.01) or barren sites (pairwise PERMANOVA, *P* <0.05), whereas communities

in the mineral layer were significantly different only at the meadow and evergreen shrub sites

(pairwise PERMANOVA, *P* <0.01).

Genus-level comparisons were conducted for the metatranscriptomes among vegetation

types and between different soil layers. Considering genera with mean abundance twofold larger

than the ´mean of all genera, in the organic layer alphaproteobacterial genera *Bradyrhizobium*,

Nitrobacter, *Rhodoplanes* and *Rhodomicrobium* were significantly (ANOVA/pairwise T-test;

P<0.01) more active in meadow sites than sites with other vegetation types (Figure 5a). In addition,

Acidothermus (Actinobacteria) was significantly more active in deciduous and evergreen shrub sites

than in meadow sites, whereas *Roseiarcus* (Rhizobiales), *Acidipila* (Acidobacteriales), *Granulicella*

(Acidobacteriales), *Edaphobacter* (Acidobacteriales), and *Gemmata* (Gemmatales) were more

active in deciduous and evergreen shrub and barren sites than in meadow sites. The following five

genera were significantly more active in the mineral layer across all vegetation types:

Anaeromyxobacter (Myxococcales) and uncultured lineages from Ktedonobacterales,

Pedospherales, and Pyrinomonadales. However, these were more active in meadow site compared

to others (Figure 5b) and *Bauldia* (Rhizobiales) was more active in meadow and barren sites than in

evergreen shrub sites (Figure 5b).

Figure 5. Boxplot showing genera with mean abundance larger than the twofold mean of all genera
that are significantly more active in certain vegetation types in a) organic layer (*P* < 0.01) and b)
mineral layer (*P* < 0.01).

3.3 Active microbial community functions across sites and vegetation types

The differences between the organic and mineral layer for protein-encoding genes were small but

significant (PERMANOVA, R^2 = 0.03044, P < 0.001). Protein-encoding genes were significantly

different (PERMANOVA, R^2 = 0.09852, P < 0.001) between vegetation types in the organic layer,

and pairwise comparisons showed that meadow sites were different from shrub and barren sites

(pairwise PERMANOVA, P <0.01) and evergreen shrub sites differed from barren sites (pairwise

PERMANOVA, P <0.05). No significant differences for functional genes were observed between

vegetation types in mineral layer.

The most abundantly transcribed genes that were mapped to KEGG (Kanehisa & Goto,

2000) pathways are involved in genetic information processing, including i) folding, sorting, and

degradation, ii) transcription, and iii) metabolism (Figure 6a). ABC transporter genes were widely

transcribed, including transport system substrate-binding proteins for ribose (*rbsB*), D-xylose

(*xylF*), and sorbitol/mannitol (*smoE*, *mtlE*), and multiple sugar transport system ATP-binding

proteins (*msmX*, *msmK*, *malK*, *sugC*, *ggtA*, *msiK*), indicating the decomposition of plant polymers

composed of these sugars. Other widely transcribed ABC transporters included branched-chain

amino acid transport system protein genes (*livKGFHM*) and urea transport system substrate-binding

protein genes (*urtA*). In addition to transporters, chaperon genes *groeL*, *dnaK,* and cold shock

protein gene *cspA*, involved in survival in cold temperatures, were among the most transcribed

genes in all samples. Gene *coxL/cutL* encoding the large subunit of aerobic carbon-monoxide

dehydrogenase was widely expressed. Two genes involved in nitrogen uptake, glutamine synthetase

glnA and ammonium transporter *amt*, were among the most abundant transcripts.

Interestingly, *pmoCAB-amoCAB* genes involved in methane and ammonia oxidation were

transcribed significantly more (*pmoC*: $R^2 = 0.2539$, *pmoB*: $R^2 = 0.1541$, *pmoA* $R^2 = 0.2372$; $P <$

0.001; ANOVA) in the mineral layer than in the organic layer (Figure 6b). In KEGG classification,

genes for methane and ammonia oxidation are not distinguished and therefore we performed

BLAST (Altschul *et al.*, 1990) for these genes against a manually curated database of eight *pmoA*

and eight *amoA* genes from different organisms. Results suggested that 96 % of the *pmoA-amoA*

hits were *pmoA*, indicating activity of methane oxidation by particulate methane monooxygenase

(pMMO) in mineral layers. There was virtually no transcription of soluble methane monooxygenase

(sMMO) genes *mmoXYZBCD* or genes *mxaF* or *xoxF*, which encode the methanol dehydrogenases

for methanol oxidation to formaldehyde (Figure 6b). Genes for formaldehyde assimilation in serine

pathway, including *glyA* encoding glycine hydroxymethyltransferase, were transcribed. This

indicates utilization of the serine cycle instead of the ribulosemonophosphate (*RuMP*) cycle for

formaldehyde assimilation in these microbial communities.

Although some individual genes showed significantly different transcription in vegetation

types (data not shown), transcriptome profiles based on KEGG hits across the vegetation types were

mostly homogenous. Genes with a KEGG classification represented only a small fraction (1.39 ±

0.27%) of the reads that were recognized as protein-encoding genes. While genes with no KEGG

classification were not analyzed in this study, they were mostly proteins with unknown function

(Supplementary Figure S3).

Figure 6. a) KEGG classified square-root transformed gene transcripts with mean relative abundance > 0.002. b) KEGG classified square-root transformed gene transcripts for methane oxidation and serine pathway formaldehyde assimilation. A column of pixels represents a site.

4 Discussion

We investigated over 100 soil samples across a subarctic landscape to gain new insights into the microbial community composition together with their previously uncharacterized functions in the tundra. This study suggests that microbial community composition and activity are strongly linked to vegetation types via soil moisture as well as organic matter quality and quantity.

The most active microbial phyla were Actinobacteria, Acidobacteria, Alphaproteobacteria, and Planctomycetes, consistent with previous studies from Arctic regions (Hultman *et al.*, 2015; Tas *et al.*, 2018; Tripathi *et al.*, 2019). These phyla, excluding Planctomycetes, were also among the most abundant in the metagenomics dataset from these samples (Pessi et al., 2020). Although the microbial community composition was similar across vegetation types, there were significant

differences at the genus level in the active microbes. Bacterial and archaeal communities and their activities in the organic layers differed across vegetation types. In particular, meadow sites were distinct from deciduous and evergreen shrub sites. Actinobacteria (Acidothermus), Acidobacteria (*Acidipila, Granulicella,* and *Edaphobacter),* and Planctomycetes (*Gemmata)* were significantly more active in the organic layer of the deciduous and evergreen shrub sites or barren sites than in meadow sites, whereas Alphaproteobacteria (*Bradyrhizobium*, *Nitrobacter*, *Rhodoplanes,* and *Rhodomicrobium)* were significantly more active in meadow sites.

In the organic layer, soil physicochemical factors also varied according to vegetation type, but not in the mineral layer. These linkages are likely related to the vegetation being the primary source of material for the organic layer, whereas the mineral layer properties are more affected by bedrock and soil texture, for example (Jenny 1941). Soil samples from shrub sites had significantly higher SOM concentrations and lower pH, whereas meadow sites had lower SOM concentrations and higher pH. The difference between shrub and meadow sites is presumably linked to the differences in organic matter quality; evergreen shrubs, such as *Empetrum nigrum*, (the dominant plant species of the study area) produce recalcitrant, acidic, and slowly decomposing litter, whereas meadows are dominated by forbs, grasses, and sedges, which produce litter that decomposes faster than the litter of *E. nigrum* (Hobbie *et al.*, 2000) and have higher nutrient concentrations. However, the two shrubs are difficult to separate as they usually form mixed communities within the study area. In the mineral layer, the environmental conditions can be more homogeneous and plant roots may not have a direct effect on the microbial community, which leads to more uniform microbial communities between different vegetation types. Although mineral samples were collected below the true rhizosphere, the indirect impact of plant roots is considerable in the proximate soil (Haichar *et al.*, 2008).

Acidobacteria were the most active phylum in all vegetation types, consistent with a previous PLFA study on the same area (Männistö *et al.*, 2007). The genera *Acidipila*, *Granulicella*, and *Edaphobacter* from order Acidobacteriales (also known as Subgroup 1), were particularly active in the organic layer with shrub vegetation. Of these, *Granulicella* are chemoorganotrophic bacteria capable of degrading various heteropolysaccharides, and several described species have been found in acidic upland soils (Pankratov & Dedysh, 2010; Männistö *et al.*, 2013). Acidobacteriales were also predominant in Northern Russian raised bogs (Ivanova *et al.*, 2020).

In this study, *Candidatus* Solibacter Ellin6076 and *Bryobacter* from Acidobacterial order Solibacterales were the second and third most active genera, with no significant variation according to vegetation type. Both have been found in acidic boreal soils (Kulichevskaya *et al.*, 2009; Rawat *et al.*, 2012). Genome analysis of *Candidatus* Solibacter revealed the ability not only to utilize

complex plant cell-wall polysaccharides and simple sugars but might also be able to use carbon

monoxide (CO), a toxic gas, as a complementary energy source in a mixotrophic lifestyle (Ward *et*

al., 2009). Moreover, the carbon monoxide dehydrogenase gene *coxL/cutL* was widely expressed

here, which may indicate the usage of CO in these soils. Thus, these bacteria are likely to have a

role in carbon cycling in these tundra soils. Additionally, *Candidatus* Solibacter can participate in

nitrogen cycling as they harbor candidate nitrate-reduction genes. *Bryobacter* are moss-associated,

acidotolerant, strictly aerobic, slow-growing chemoorganotrophic bacteria, which inhabit acidic

wetlands and soils and are also capable of hydrolyzing various heteropolysaccharides

(Kulichevskaya *et al.*, 2009). These acidophilic bacteria likely have a role as diverse degraders of

organic matter in tundra soils, as they have been shown to survive temperature and moisture

fluctuation and low-nutrient conditions (Ward *et al.*, 2009; Rawat *et al.*, 2012).

Actinobacteria were widely represented in all vegetation types, with a large number of

unclassified taxa. The most abundant active genus, *Acidothermus*, is the only described genus in the

family *Acidothermaceae* and has one described species (*Acidothermus cellulolyticus,* Mohagheghi

et al., 1986). This thermophilic, acidophilic, and cellulolytic species was first isolated from acidic

hot springs and has been studied mostly because of its endoglucanase for biotechnological

applications (Mohagheghi *et al.*, 1986). In tundra soils, *Acidothermus* likely has a role as a degrader

of cellulose from plant litter. Another active actinobacterial group abundant in these soils, order

Solirubrobacterales, is not well characterized. These are mesophilic heterotrophs and based on 16S

rRNA gene analysis, Solirubrobacterales have been encountered in diverse environments, such as

weathered serpentine rock (Khilyas *et al.*, 2019), Antarctic soil (Chong *et al.*, 2012), agroecosystem

soils (Shange *et al.*, 2012), *Sphagnum* peat bog (Ivanova *et al.*, 2016), and shrub-associated desert

soil (Saul-Tcherkas & Steinberger, 2011). This wide range of habitats suggests that this order

includes mixotrophic bacteria with the ability to survive in environments with fluctuating

conditions, including temperature, nutrients, and UV radiation.

In addition to Actinobacteria and Acidobacteria, the following three orders from phylum

Planctomycetes were among the most active bacterial groups in these soils: Gemmatales,

Isosphaerales, and Tepidisphaerales. These orders are abundant in acidic raised bogs with high

organic matter but low nutrient content (Ivanova *et al.*, 2020) and they have chitinolytic potential in

Sphagnum-dominated wetlands (Ivanova *et al.*, 2018). *Aquisphera* and *Singulisphaera* from order

Isosphaerales were among the most active genera. These acidophilic bacteria have been isolated

from northern Russian peatlands (Kulichevskaya *et al.*, 2009; Kulichevskaya *et al.*, 2012) and can

degrade several biopolymers under acidic, microaerobic, and low-temperature conditions.

Intriguingly, metatranscriptomes revealed that Planctomycetes were more active than what could be

expected based on the metagenomes, indicating that interpretation based solely on metagenomic data would underestimate their importance. As Planctomycetes were widely active in all vegetation types and genus *Gemmata* was significantly more active in the shrub sites, the results suggest that Planctomycetes are potentially active and diverse OM degraders in tundra soils.

Diverse Alphaproteobacteria were active in all sites, and the order Rhizobiales was particularly predominant in all vegetation types. *Roseiarcus* was among the most active genera in shrub sites with low pH and high OM. The only described species from this genus, *Roseiarcus fermentas*, was isolated from a methanotrophic consortium enriched in acidic *Sphagnum* peat (Kulichevskaya *et al.*, 2014). These bacteria belong to a cluster of environmental sequences, which have been obtained from wetlands and forest soils by molecular methods and are probably adapted to live in acidic soils in alternating high UV radiation and dark periods and varying levels of oxygen (Kulichevskaya *et al.*, 2014).

Potential CH_4 oxidation was observed particularly in the mineral layer, as particulate methane monooxygenase (pMMO) was transcribed together with the activity of methanotrophic bacteria, such as *Methylocapsa* (Rhizobiales). This genus has only a few described species, including *M. acidiphila* isolated from Siberian acidic *Sphagnum* bog (Dedysh *et al.*, 2002), *M. palsarum* isolated in subarctic palsa soil (Dedysh *et al.*, 2015), and *M. gorgona* from subarctic soil (Tveit *et al.*, 2015). These bacteria possess pMMO and use the serine pathway for one-carbon compound utilization, which was also transcribed here. Additionally, *Methylocystis*, *Methylovirgula,* and *Methylocella* showed activity in these soils. Contrastingly, Singleton et al. (2018) found gammaproteobacterial but no alphaproteobacterial methanotrophs active in permafrost thaw fens. Based on the results shown here and by others (Belova *et al.*, 2020), *Methylocapsa* and other methanotrophs from order the Rhizobiales could have a significant role as CH_4 oxidizers in subarctic tundra soils.

Several other Rhizobiales with diverse metabolism were particularly active in meadows. Although many known members of *Bradyrhizobium* are chemoorganotrophs and symbiotic N-fixers, some grow chemolithotrophically in the presence of H_2, CO_2, and low levels of O_2. Genes involved in N fixation were not observed in these soils. Nitrite-oxidizing *Nitrobacter* and anaerobic photoheterotrophic *Rhodoplanes* with capacity for complete denitrification were active in meadow sites. Overall, Alphaproteobacteria are an active and metabolically diverse group in these tundra soils.

We observed the activity of several candidate divisions of phylum Chloroflexi, especially in the mineral layer. One of the most diverse and active divisions were Ktedonobacterales, which are abundant in extreme environments with fluctuating temperature, intense solar radiation, and low

OM content. Ktedonobacterales are also potential atmospheric CO and H_2 oxidizers (King & King, 2014; Islam *et al.*, 2019). In addition to Chloroflexi, members of the uncultivated phyla Dormibacterota and Eremiobacterota were active in the mineral layer. As Ktedonobacteria, they can exploit atmospheric H_2, CO_2, and CO for energy and carbon sources (Ji *et al.*, 2017). Transcription of carbon monoxide dehydrogenase might indicate Ktedonobacterales, Dormibacterota and Eremiobacterota and the previously mentioned *Candidatus* Solibacter oxidize CO in these soils. *Candidatus* Udaeobacter and uncultured lineage ADurb.Bin063-1 from phylum Verrucomicrobia were particularly active in meadows. *Candidatus* Udaeobacter potentially oxidizes trace gas H_2 to generate energy and benefits antibiotics release in soil (Willms *et al.*, 2020). These are poorly characterized bacteria, yet they were active and potentially have traits to survive in harsh, competitive conditions with the ability to scavenge atmospheric trace gases.

Archaeal 16S rRNA represented only 0.1% in the studied tundra samples. Thaumarchaeal genera *Nitrosophaera* and *Nitrosotalea* and candidate division Group 1.1c were the most active Archaea in the soils and were significantly more active in the mineral layer. *Nitrosophaera* and *Nitrosotalea* are capable of NH_3 oxidation but are suspected of a mixotrophic or even heterotrophic lifestyle (Tourna *et al.*, 2011; Lehtovirta-Morley *et al.*, 2014). The cultured strains were isolated from temperate soils and their temperature optima were at mesophilic scale. Group 1.1c is the dominating archaeal taxa in temperate acidic forest and forest peat soils (Kemnitz *et al.*, 2007; Stopnišek Nejc *et al.*, 2010). These archaea grow at low pH (Lehtovirta *et al.*, 2009) at an optimal temperature of 30°C (Weber *et al.*, 2015). Group 1.1c does not oxidize NH_3 or carry the gene *amoA*, in contrast to cultured relatives from groups 1.1a and 1.1b do, which is why they are suspected to be heterotrophic (Weber *et al.*, 2015). Temperature may be the growth-limiting factor of these Archaea, and their activity in carbon-rich northern soils may be higher with warming climate. Indeed, they do have a role in the tundra soil ecosystem that is yet to be discovered.

5 Conclusions

Understanding the microbial communities and their functions is key to acquiring a process-level understanding about tundra biogeochemistry and accurate models of greenhouse gas release. It is of global relevance to investigate this particularly in the Arctic, where soil organic carbon stocks are large and vulnerable to climate change and have drastic impacts on the global carbon cycle. Our results revealed the large number of active bacteria and archaea across tundra soils. The activity of these communities varied according to vegetation types; in particular, communities in the organic layer at the meadow sites were distinct from deciduous and evergreen shrub sites. Communities at

the deciduous and evergreen shrub sites had a similar microbial community composition and activity. Vegetation type-driven differences in microbial communities in the mineral layer were less evident. Potential organic matter degraders belonging to Actinobacteria, Acidobacteria, Alphaproteobacteria, and Planctomycetes were active in shrub soils high SOM content, whereas poorly known Chloroflexi and Dormibacterota were active in the deeper, low SOM mineral layer. Chloroflexi and Dormibacterota use potentially atmospheric trace gases for growth. In addition, alphaproteobacterial methanotrophs are potential moderators of methane release, particularly in the mineral layers of subarctic tundra soils. Genes involved in nitrogen fixation were not observed in these soils, however nitrite-oxidizing *Nitrobacter* sp. and *Rhodoplanes* sp. with capacity for complete denitrification were active in meadow sites. Microbes in these soils thus impact the tundra ecosystem functioning in various ways and play an important role in climate change feedbacks.

6   Acknowledgments

The study was funded by the Academy of Finland (Grant 1314114) and the University of Helsinki three-year grant to JH. SV was funded by the Doctorate Program in Microbiology and Biotechnology. PN was funded by Kone Foundation and Nessling Foundation. JK was funded by the Arctic Interactions at the University of Oulu and Academy of Finland (grant no. 318930, Profi 4). AMV acknowledges Gordon and Betty Moore Foundation (Grant 8414). Permission to perform fieldwork was granted by Metsähallitus. We acknowledge the CSC IT center for science LTD for the computational resources and Mr. Kimmo Mattila for assistance. We acknowledge the Kilpisjärvi Biological Station and staff and members of the BioGeoClimate Modelling Lab for use of the premises and assistance with fieldwork.

7   References

Altschul SF, Gish W, Miller W, Myers EW & Lipman DJ (1990) Basic local alignment search tool J Mol Biol **215:** 403-410.

Andrews S. FastQC: A Quality Control Tool for High Throughput Sequence Data. 2019.

Belova SE, Danilova OV, Ivanova AA, Merkel AY & Dedysh SN (2020) Methane-Oxidizing Communities in Lichen-Dominated Forested Tundra Are Composed Exclusively of High-Affinity USCalpha Methanotrophs. Microorganisms **8:** 10.3390/microorganisms8122047.

Bengtsson-Palme J, Richardson RT, Meola M *et al.* (2018) Metaxa2 Database Builder: enabling taxonomic identification from metagenomic or metabarcoding data using any genetic marker. Bioinformatics **34:** 4027-4033.

Berner LT, Massey R, Jantz P *et al.* (2020) Summer warming explains widespread but not uniform
greening in the Arctic tundra biome. Nat Commun **11:** 4621-020-18479-5.

Bond-Lamberty B & Thomson A (2010) Temperature-associated increases in the global soil
respiration record. Nature **464:** 579-582.

Bond-Lamberty B, Bailey VL, Chen M, Gough CM & Vargas R (2018) Globally rising soil
heterotrophic respiration over recent decades. Nature **560:** 80-83.

Buchfink B, Xie C & Huson DH (2015) Fast and Sensitive Protein Alignment using DIAMOND.
Nature Methods **12:** 59–60.

Buckeridge, K. M., Banerjee, S., Siciliano, S. D., & Grogan, P. (2013). The seasonal pattern of soil
microbial community structure in mesic low arctic tundra. Soil Biology and Biochemistry, **65**: 338-
347.

Chong CW, Convey P, Pearce DA & Tan IKP (2012) Assessment of soil bacterial communities on
Alexander Island (in the maritime and continental Antarctic transitional zone). Polar Biol **35:** 387-
399.

Chu H, Fierer N, Lauber CL, Caporaso JG, Knight R & Grogan P (2010) Soil bacterial diversity in
the Arctic is not fundamentally different from that found in other biomes. Environ Microbiol.
12(11):2998-3006.

DeAngelis KM, Brodie EL, DeSantis TZ, Andersen GL, Lindow SE & Firestone MK (2009)
Selective progressive response of soil microbial community to wild oat roots. ISME J **3:** 168-178.

Dedysh SN, Didriksen A, Danilova OV, Belova SE, Liebner S & Svenning MM (2015)
Methylocapsa palsarum sp. nov., a methanotroph isolated from a subArctic discontinuous
permafrost ecosystem. Int J Syst Evol Microbiol **65:** 3618-3624.

Dedysh SN, Khmelenina VN, Suzina NE, Trotsenko YA, Semrau JD, Liesack W & Tiedje JM
(2002) Methylocapsa acidiphila gen. nov., sp. nov., a novel methane-oxidizing and dinitrogen-
fixing acidophilic bacterium from Sphagnum bog. Int J Syst Evol Microbiol **52:** 251-261.

Ewels P, Magnusson M, Lundin S & Käller M (2016) MultiQC: summarize analysis results for
multiple tools and samples in a single report. Bioinformatics **32:** 3047-3048.

Falkowski PG, Fenchel T & Delong EF (2008) The microbial engines that drive Earth's
biogeochemical cycles. Science 320: 1034-1039.

Friend AD, Lucht W, Rademacher TT *et al.* (2014) Carbon residence time dominates uncertainty in
terrestrial vegetation responses to future climate and atmospheric CO2. Proc Natl Acad Sci U S A
111: 3280-3285.

Griffiths RI, Whiteley AS, O'Donnell AG & Bailey MJ (2000) Rapid method for coextraction of
DNA and RNA from natural environments for analysis of ribosomal DNA- and rRNA-based
microbial community composition. Appl Environ Microbiol **66:** 5488-5491.

Haichar FZ, Marol C, Berge O, Rangel-Castro JI, Prosser JI, Balesdent J, Heulin T & Achouak W (2008) Plant host habitat and root exudates shape soil bacterial community structure. ISME J **2:** 1221-1230.

Hervé M (2021) RVAideMemoire: Testing and Plotting Procedures for Biostatistics.

Hobbie SE & Gough L (2002) Foliar and soil nutrients in tundra on glacial landscapes of contrasting ages in northern Alaska. Oecologia **131:** 453-462.

Hobbie SE, Schimel JP, Trumbore SE & Randerson JR (2000) Controls over carbon storage and turnover in high-latitude soils. Global Change Biol **6:** 196-210.

Hultman J, Waldrop MP, Mackelprang R *et al.* (2015) Multi-omics of permafrost, active layer and thermokarst bog soil microbiomes. Nature **521:** 208-212.

IPCC, 2018: Global warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty [V. Masson-Delmotte, P. Zhai, H. O. Pörtner, D. Roberts, J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J. B. R. Matthews, Y. Chen, X. Zhou, M. I. Gomis, E. Lonnoy, T. Maycock, M. Tignor, T. Waterfield (eds.)]. In Press.

Islam ZF, Cordero PRF, Feng J *et al.* (2019) Two Chloroflexi classes independently evolved the ability to persist on atmospheric hydrogen and carbon monoxide. The ISME Journal **13:** 1801-1813.

Ivanova AA, Zhelezova AD, Chernov TI & Dedysh SN (2020) Linking ecology and systematics of acidobacteria: Distinct habitat preferences of the Acidobacteriia and Blastocatellia in tundra soils. PLOS ONE **15:** e0230157.

Ivanova AA, Wegner C, Kim Y, Liesack W & Dedysh SN (2018) Metatranscriptomics reveals the hydrolytic potential of peat-inhabiting Planctomycetes. Antonie Van Leeuwenhoek **111:** 801-809.

Ivanova AA, Kulichevskaya IS, Merkel AY, Toshchakov SV & Dedysh SN (2016) High Diversity of Planctomycetes in Soils of Two Lichen-Dominated Sub-Arctic Ecosystems of Northwestern Siberia. Frontiers in Microbiology **7:** 2065.

Jansson JK & Hofmockel KS (2020) Soil microbiomes and climate change. Nat Rev Microbiol **18:** 35-46.

Jenny H (1941) Factors Of Soil Formation A System of Quantitative Pedology. Dover Publications, New York, 281 p.

Ji M, Greening C, Vanwonterghem I *et al.* (2017) Atmospheric trace gases support primary production in Antarctic desert surface soil. Nature **552:** 400-403.

Johnston, E. R., Hatt, J. K., He, Z., Wu, L., Guo, X., Luo, Y., ... & Konstantinidis, K. T. (2019). Responses of tundra soil microbial communities to half a decade of experimental warming at two critical depths. Proceedings of the National Academy of Sciences, **116**(30), 15096-15105.

Kanehisa M & Goto S (2000) KEGG: kyoto encyclopedia of genes and genomes. Nucleic Acids
Res **28:** 27-30.

Kemnitz D, Kolb S & Conrad R (2007) High abundance of Crenarchaeota in a temperate acidic
forest soil. FEMS Microbiol Ecol **60:** 442-448.

Kemppinen J, Niittynen P, Aalto J, le Roux PC & Luoto M (2019) Water as a resource, stress and
disturbance shaping tundra vegetation. Oikos 128: 811-822.

Kemppinen J, Niittynen P, le Roux PC *et al.* (2021) Consistent trait-environment relationships
within and across tundra plant communities. Nat Ecol Evol **5:** 458-467.

Khilyas IV, Sorokina AV, Elistratova AA, Markelova MI, Siniagina MN, Sharipova MR,
Shcherbakova TA, D'Errico ME & Cohen MF (2019) Microbial diversity and mineral composition
of weathered serpentine rock of the Khalilovsky massif. PLoS One **14:** e0225929.

King CE & King GM (2014) Description of Thermogemmatispora carboxidivorans sp. nov., a
carbon-monoxide-oxidizing member of the class Ktedonobacteria isolated from a geothermally
heated biofilm, and analysis of carbon monoxide oxidation by members of the class
Ktedonobacteria. Int J Syst Evol Microbiol **64:** 1244-1251.

King L and M Seppälä (1987) Permafrost Thickness and Distribution in Finnish Lapland - Results
of Geoelectrical Soundings. Polarforschung 57:127-147.

Koyama A, Wallenstein MD, Simpson RT & Moore JC (2014) Soil bacterial community
composition altered by increased nutrient availability in Arctic tundra soils. Front Microbiol **5:** 516.

Kulichevskaya IS, Danilova OV, Tereshina VM, Kevbrin VV & Dedysh SN (2014) Descriptions of
Roseiarcus fermentans gen. nov., sp. nov., a bacteriochlorophyll a-containing fermentative
bacterium related phylogenetically to alphaproteobacterial methanotrophs, and of the family
Roseiarcaceae fam. nov. Int J Syst Evol Microbiol **64:** 2558-2565.

Kulichevskaya IS, Baulina OI, Bodelier PL, Rijpstra WI, Damsté JS & Dedysh SN (2009)
Zavarzinella formosa gen. nov., sp. nov., a novel stalked, Gemmata-like planctomycete from a
Siberian peat bog. Int J Syst Evol Microbiol **59:** 357-364.

Kulichevskaya IS, Serkebaeva YM, Kim Y, Rijpstra WI, Damsté J,S.Sinninghe, Liesack W &
Dedysh SN (2012) Telmatocola sphagniphila gen. nov., sp. nov., a novel dendriform planctomycete
from northern wetlands. Frontiers in microbiology **3:** 146-146.

le Roux PC, Aalto J & Luoto M (2013) Soil moisture's underestimated role in climate change
impact modelling in low-energy systems. Glob Chang Biol **19:** 2965-2975.

Lehtovirta LE, Prosser JI & Nicol GW (2009) Soil pH regulates the abundance and diversity of
Group 1.1c Crenarchaeota. FEMS Microbiol Ecol **70:** 367-376.

Lehtovirta-Morley L, Ge C, Ross J, Yao H, Nicol GW & Prosser JI (2014) Characterisation of
terrestrial acidophilic archaeal ammonia oxidisers and their inhibition and stimulation by organic
compounds. FEMS Microbiol Ecol **89:** 542-552.

Li H., Seqtk: a fast and lightweight tool for processing FASTA or FASTQ sequences. 2016

Mackelprang R, Waldrop M, DeAngelis K *et al.* (2011) Metagenomic analysis of a permafrost
microbial community reveals a rapid response to thaw. Nature **480,** 368–371.
637
638   Mod, H.K., Buri, A., Yashiro, E. et al. Predicting spatial patterns of soil bacteria under current and
639   future environmental conditions. ISME J (2021).
640
Männistö MK, Tiirola M & Haggblom MM (2009) Effect of freeze-thaw cycles on bacterial
communities of arctic tundra soil. Microb Ecol **58:** 621-631.

Männistö MK, Tiirola M & Haggblom MM (2007) Bacterial communities in Arctic fjelds of
Finnish Lapland are stable but highly pH-dependent. FEMS Microbiol Ecol **59:** 452-465.

Männistö MK, Kurhela E, Tiirola M & Haggblom MM (2013) Acidobacteria dominate the active
bacterial communities of Arctic tundra with widely divergent winter-time snow accumulation and
soil temperatures. FEMS Microbiol Ecol **84:** 47-59.

Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal **17**

McCalley CK, Woodcroft BJ, Hodgkins SB *et al.* (2014) Methane dynamics regulated by microbial
community response to permafrost thaw. Nature **514:** 478-481.

McMurdie PJ & Holmes S (2013) phyloseq: an R package for reproducible interactive analysis and
graphics of microbiome census data. PLoS One **8:** e61217.

Mohagheghi A, Grohmann K, Himmel M, Leighton L & Updegraff (986) Dmy1: Isolation and
Characterization of Acidothermus cellulolyticus gen. nov., sp. nov., a New Genus of Thermophilic,
Acidophilic, Cellulolytic Bacteria. International Journal of Systematic and Evolutionary
Microbiology, **36:** 435-443.

Neufeld JD, Mohn WW(2005) Unexpectedly high bacterial diversity in arctic tundra relative to
boreal forest soils, revealed by serial analysis of ribosomal sequence tags. Appl Environ Microbiol.
71(10):5710-5718.

Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al. vegan: Community
Ecology Package. 2019.

Pankratov TA & Dedysh SN (2010) Granulicella paludicola gen. nov., sp. nov., Granulicella
pectinivorans sp. nov., Granulicella aggregans sp. nov. and Granulicella rosea sp. nov., acidophilic,
polymer-degrading acidobacteria from Sphagnum peat bogs. Int J Syst Evol Microbiol **60:** 2951-
2959.

Parks DH, Chuvochina M, Chaumeil PA, Rinke C, Mussig AJ & Hugenholtz P (2020) A complete
domain-to-species taxonomy for Bacteria and Archaea. Nat Biotechnol **38:** 1079-1086.

Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil PA & Hugenholtz P
(2018) A standardized bacterial taxonomy based on genome phylogeny substantially revises the tree
of life. Nat Biotechnol **36:** 996-1004.

Pessi IS, Viitamäki S, Eronen-Rasimus E, Delmont TO, Luoto M & Hultman J (2020) Truncated
denitrifiers dominate the denitrification pathway in tundra soil metagenomes. bioRxiv
2020.12.21.419267.

Pessi IS. keggR: a tool to parse the results of BLAST/DIAMOND similarity searches made against
the KEGG GENES prokaryotes database. github.com/igorspp/keggR 2020.

Pirinen P, Simola H, Aalto J, Kaukoranta JP, Karlsson P, Ruuhela R (2012) Tilastoja Suomen
ilmastosta 1981–2010 (Climatological statistics of Finland 1981–2010). Finnish Meteorological
Institute Reports 2012.

Post E, Alley RB, Christensen TR *et al.* (2019) The polar regions in a 2 degrees C warmer world.
Sci Adv **5:** eaaw9883.

Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J & Glockner FO (2013) The
SILVA ribosomal RNA gene database project: improved data processing and web-based tools.
Nucleic Acids Res **41:** D590-6.

R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Rawat SR, Männistö MK, Bromberg Y & Haggblom MM (2012) Comparative genomic and
physiological analysis provides insights into the role of Acidobacteria in organic carbon utilization
in Arctic tundra soils. FEMS Microbiol Ecol **82:** 341-355.

Raynolds MK, Walker DA, Balser A *et al.* (2019) A raster version of the Circumpolar Arctic
Vegetation Map (CAVM). Remote Sens Environ **232:** 111297.

Saul-Tcherkas V & Steinberger Y (2011) Soil microbial diversity in the vicinity of a Negev Desert
shrub--Reaumuria negevensis. Microb Ecol **61:** 64-81.

Schloss PD, Westcott SL, Ryabin T *et al.* (2009) Introducing mothur: open-source, platform-
independent, community-supported software for describing and comparing microbial communities.
Appl Environ Microbiol **75:** 7537-7541.

Schostag M, Stibal M, Jacobsen CS, Baelum J, Tas N, Elberling B, Jansson JK, Semenchuk P &
Prieme A (2015) Distinct summer and winter bacterial communities in the active layer of Svalbard
permafrost revealed by DNA- and RNA-based analyses. Front Microbiol **6:** 399.

Shange RS, Ankumah RO, Ibekwe AM, Zabawa R & Dowd SE (2012) Distinct Soil Bacterial
Communities Revealed under a Diversely Managed Agroecosystem. PLOS ONE **7:** e40338.

Singleton CM, McCalley CK, Woodcroft BJ *et al.* (2018) Methanotrophy across a natural
permafrost thaw environment. ISME J **12:** 2544-2558.

Stopnišek Nejc, Gubry-Rangin Cécile, Höfferle Špela, Nicol Graeme W., Mandič-Mulec Ines &
Prosser James I. (2010) Thaumarchaeal Ammonia Oxidation in an Acidic Forest Peat Soil Is Not
Influenced by Ammonium Amendment. Appl Environ Microbiol **76:** 7626-7634.

Tas N, Prestat E, Wang S, Wu Y, Ulrich C, Kneafsey T, Tringe SG, Torn MS, Hubbard SS &
Jansson JK (2018) Landscape topography structures the soil microbiome in arctic polygonal tundra.
Nat Commun **9:** 777-018-03089-z.

Tourna M, Stieglmeier M, Spang A *et al.* (2011) *Nitrososphaera viennensis*, an ammonia oxidizing
archaeon from soil. Proc Natl Acad Sci USA **108:** 8420-8425.

Tripathi BM, Kim HM, Jung JY, Nam S, Ju HT, Kim M & Lee YK (2019) Distinct Taxonomic and
Functional Profiles of the Microbiome Associated With Different Soil Horizons of a Moist Tussock
Tundra in Alaska. Front Microbiol **10:** 1442.

Tveit AT, Urich T, Frenzel P & Svenning MM (2015) Metabolic and trophic interactions modulate
methane production by Arctic peat microbiota in response to warming. Proc Natl Acad Sci USA
112: E2507-E2516.

Urich T, Lanzen A, Qi J, Huson DH, Schleper C & Schuster SC (2008) Simultaneous assessment of
soil microbial community structure and function through analysis of the meta-transcriptome. PLoS
One **3:** e2527.

Virkkala A-M, Virtanen T, Lehtonen A, Rinne J, and Luoto M (2018) The current state of CO2 flux
chamber studies in the Arctic tundra: A review. Prog. Phys. Geogr., **42:**162–184.

Walker DA, Raynolds MK, Daniëls FJ, Einarsson E, Elvebakk A, Gould WA, Katenin AE, Kholod
SS, Markon CJ, Melnikov ES, Moskalenko NG, Talbot SS, Yurtsev BA and The other members of
the CAVM Team (2005) The Circumpolar Arctic vegetation map. Journal of Vegetation Science,
16: 267-282.

Ward NL, Challacombe JF, Janssen PH *et al.* (2009) Three genomes from the phylum
Acidobacteria provide insight into the lifestyles of these microorganisms in soils. Appl Environ
Microbiol **75:** 2046-2056.

Weber EB, Lehtovirta-Morley LE, Prosser JI & Gubry-Rangin C (2015) Ammonia oxidation is not
required for growth of Group 1.1c soil Thaumarchaeota. FEMS Microbiol Ecol **91:** fiv001.

Willms IM, Rudolph AY, Göschel I, Bolz SH, Schneider D, Penone C, Poehlein A, Schöning I &
Nacke H (2020) Globally Abundant "Candidatus Udaeobacter" Benefits from Release of
Antibiotics in Soil and Potentially Performs Trace Gas Scavenging. mSphere **5:** e00186-20.

Woodcroft BJ, Singleton CM, Boyd JA *et al.* (2018) Genome-centric view of carbon processing in
thawing permafrost. Nature **560:** 49-54.

Ye Y & Doak TG (2009) A parsimony approach to biological pathway reconstruction/inference for
genomes and metagenomes. PLoS Comput Biol **5:** e1000465.

Zhang X, Xu S, Li C, Zhao L, Feng H, Yue G, Ren Z & Cheng G (2014) The soil carbon/nitrogen
ratio and moisture affect microbial community structures in alkaline permafrost-affected soils with
different vegetation types on the Tibetan plateau. Res Microbiol **165:** 128-139.