Mathematical modeling of the effects of Wnt-10b on bone metabolism

Carley V. Cook[a,b], Mohammad Aminul Islam[a,b], Brenda J. Smith[c], Ashlee N. Ford Versypt[a,b,d,*]

[a]*School of Chemical Engineering, Oklahoma State University, Stillwater, OK, USA*
[b]*Department of Chemical and Biological Engineering, University at Buffalo, The State University of New York, Buffalo, NY, USA*
[c]*Department of Nutritional Sciences, Oklahoma State University, Stillwater, OK, USA*
[d]*Institute for Computational and Data Sciences, University at Buffalo, The State University of New York, Buffalo, NY, USA*

Abstract

Bone health is determined by many factors including bone metabolism or remodeling. Wnt-10b has been shown to alter osteoblastogenesis through pre-osteoblast proliferation and differentiation as well as the osteoblast apoptosis rate, which collectively lead to the increase of bone density. To model this change, we adapted a previously published model of bone remodeling. The resulting model is a single compartment system that includes ordinary differential equations for active osteoclasts, pre-osteoblasts, osteoblasts, and osteocytes and a differential equation that tracks the amount of bone present at the remodeling site. Our alterations to the original model consist of extending it past a single remodeling cycle and implementing a direct relationship to Wnt-10b. Four new parameters were estimated and validated using normalized data from mice. The model connects Wnt-10b to bone metabolism and predicts the change in trabecular bone volume caused by a change in Wnt-10b. We find that this model predicts the expected increase in pre-osteoblasts and osteoblasts while also pointing to a decrease in osteoclasts when Wnt-10b is increased.

Keywords: Bone remodeling, Wnt signaling, Mathematical modeling, Bone cells

1. Introduction

Osteoporosis, characterized by decreased bone mass and structural deterioration, results from an imbalance in the tissue's metabolic processes. In the adult skeleton, bone is remodeled within millions of basic multicellular units (BMU) and involves the activity of osteoclasts, osteoblasts, osteocytes, and their precursors [1–4]. An important regulator in bone remodeling is the canonical Wnt/β-catenin pathway [5, 6]. This pathway is important for growth and development [6], maintenance of teeth [6], and immune responses [7–9]. This pathway is activated by glycoproteins in the wingless-related integration site family (Wnt). Of the nineteen different types of Wnt, only seven are known to activate the canonical Wnt/β-catenin pathway. Activation with a fizzled receptor and a low-density lipoprotein receptor-related protein coreceptor bound

to Wnt prevents β-catenin deconstruction by the proteasome. [6, 8]. Avoiding deconstruction, β-catenin activates gene expressions related to Wnt expression in the nucleus.

Expression of Wnt-10b, a canonical Wnt/β-catenin pathway activator, occurs in various systems in the body including both the bone microenvironment and the immune system [10]. While over activation of the canonical Wnt/β-catenin pathway has been associated with a tumor environment [7, 9, 10], normal levels of Wnt-10b are required for T cells to develop and function properly [10]. When targets of the β-catenin pathway are disrupted, T cells do not properly develop. In one study, T cell factor-1 is disrupted in a mouse model causing the T cells to not develop properly [11], and in another study complete deletion of lymphoid enhancing factor-1 led to nonviable mice [12]. T cells can also be a source of Wnt-10b [10, 13].

T-cell produced Wnt-10b has been implicated in altering bone volume in mice [13, 14] and has been simulated by us [15]. This suggests another link between the bone and the immune system as Wnt-10b is necessary for bone remodeling. Wnt-10b can also be produced by cells of osteoblastic lineage. A change in Wnt-10b levels has been shown to alter bone volume significantly [13, 16–18]. Wnt-10b regulates bone volume via the canonical pathway by activating genes related to osteoblastogenesis, pre-osteoblast proliferation, and cell survival [10, 19, 20]. The extent to which changes in Wnt-10b levels alter these factors and how these alterations affect the cell populations and bone volume as a whole is currently unknown.

Currently, there are a few published mathematical/computational models on bone metabolism. One model describes the autocrine and paracrine interactions of osteoblasts and osteoclasts using ordinary differential equations (ODEs) that track the cell populations as well as changes in trabecular bone mass [21]. This model has been updated to include a relationship with the parathyroid hormone (PTH) [22]. These publications along with a model that tracks spatial trabecular bone volume changes [23] have been used to develop a human based ODE model that tracks how different signals cause the development of active osteoclasts and osteoblasts and a change in trabecular bone mass [24]. None of these models include Wnt-10b; therefore, each lacks a vital part of the bone remodeling system. The authors of [25] have pointed out the importance of including Wnt-10b in a bone remodeling model using a network model that identifies the Wnt and PTH pathways as the most important systems in regulating bone remodeling.

A few of the existing bone metabolism models do explore the role Wnt-10b has in bone remodeling [26–30]. However, the models in one subset [26–28] assume a constant basal concentration of Wnt-10b and focus on how an increase of sclerostin competes for the Wnt binding sites. The models in the other subset [29, 30] allow the Wnt-10b concentration to vary but only in the presence of a tumor. These models connect Wnt-10b through Hill functions that alter an assortment of the terms for pre-osteoblast proliferation, differentiation from pre-osteoblasts to osteoblasts, and osteoblast apoptosis (Table 1). None of these models address potential changes in Wnt-10b levels caused by T cells or BMU cells.

The present work aims to improve understanding of bone metabolism by developing a computational model that directly connects the remodeling cycle to altered Wnt-10b concentrations that can be constant or

Table 1: Wnt-10b relationships presented in existing models for bone remodeling

Model	Wnt-10b type	Wnt-10b interactions
[26]	Constant	Not explicitly modeled
[27]	Constant	Pre-osteoblast proliferation
[28]	Constant	Pre-osteoblast proliferation Pre-osteoblast differentiation Osteoblast apoptosis
[29]	Dynamic	Pre-osteoblast proliferation
[30]	Dynamic	Pre-osteoblast proliferation Pre-osteoblast differentiation

dynamically induced by other physiologically processes such as an immune response. The model development is covered in Section 2, which includes an overview of the model, the data used for the model, and the techniques used to determine parameter values. After validating the model, we used it to explore how Wnt-10b directly effects important cell populations in a BMU and how these population changes alter bone volume (Section 3). The model will allow us in the future to explore potential ways to shift bone metabolism away from an imbalance that favors bone resorption to prevent bone loss or restore bone, but not without some limitations discussed in Section 4.

2. Methods

We alter a published model, which is referred to as Graham 2013 [24], to include a mathematical relationship that represents how Wnt-10b interacts with bone formation (Figure 1). This is a single compartment model that tracks the important cells involved in the bone remodeling cycle as well as bone volume. We extend past a single remodeling cycle and add a direct relationship to Wnt-10b stimuli. We add a relationship to Wnt-10b by utilizing published data to fit model parameters.

Figure 1: Interactions between bone cell populations considered in the Graham 2013 model [24] and the added impact of Wnt-10b (1-3). Pre-osteoblasts (P), osteoblasts (B), osteocytes (S), and osteoclasts (C) are tracked by differential equations. Pre-osteoclasts (PC) are assumed constant and are not explicitly modeled. The interactions include an assortment of autocrine (not depicted here) and paracrine (single dashed lines) signaling factors. Single dashed lines with arrows represent activating relationships, and single dashed lines with a block end represent inhibitory relationships. Solid lines represent transformations: self-proliferation, differentiation, and apoptosis. Wnt-10b (Wnt) stimulates osteoblastogenesis through pre-osteoblast proliferation (1) and differentiation to osteoblasts (2). Wnt-10b inhibits osteoblast apoptosis (3). Impacts of bone cells on bone volume from a combination of bone resorption and formation are represented by double dashed lines with block and arrow ends, respectively.

2.1. Graham 2013 model

The Graham 2013 model includes five ODEs that track the changes in bone volume (z) through the active populations of osteocytes (S), pre-osteoblasts (P), osteoblasts (B), and osteoclasts (C) [24]. The model does not include an equation for the population of pre-osteoclasts. The model includes important autocrine and paracrine signaling factors that are represented by power law relationships, g_{ij} and f_{ij}, where i and j are the two cell types involved in the signaling. This section summarizes the parts of the Graham 2013 model and parameter values that we adopt directly from [24] without alteration (Table 2).

2.1.1. Osteocytes

Eq. 1 describes the dynamics of the osteocyte cell population, S. Mature osteoblasts convert into an osteocyte phenotype at a rate of α_1. The term $\left(1 - \frac{S}{K_s}\right)_+$ represents the effectiveness of sclerostin regulation by osteocytes, and the subscript $+$ means that the value must remain greater than or equal to

zero. If the quantity inside the parentheses were to be evaluated as less than zero, then the term would be set to zero. Note that although sclerostin regulation does include a Wnt pathway, we are focusing on Wnt-10b excreted from T cells or from a genetic perturbation, not Wnt-10b produced within a normal balanced remodeling cycle. It is assumed in the model that over the duration of remodeling, osteocytes will not die. Instead of a death term, the osteocyte population is reduced from the steady state value of 200 cells to 180 cells at the start of each remodeling cycle. This reduction leads to a decreased level of sclerostin and allows for the RANKL activation of osteoclasts (Eq. 2). This process represents the initial biomechanical action that triggers a remodeling cycle.

$$\frac{dS}{dt} = \alpha_1 B^{g_{31}} \left(1 - \frac{S}{K_s} \right)_+ \tag{1}$$

2.1.2. Osteoclasts

Eq. 2 describes the dynamics of the activated osteoclast cell population, C. A large amount of pre-osteoclasts are assumed to be available without a significant change in the population, so the pre-osteoclast population is not modeled. The production of osteoclasts depends on a differentiation rate of α_4 and an interaction of the receptor activator of NF-κB (RANK)/RANK ligand (RANKL)/osteoprotegerin (OPG) system that is described by the term $S^{g_{41}} P^{g_{42}} (\epsilon + B)^{g_{43}} \left(1 - \frac{S}{K_s} \right)_+^{g_{44}}$. OPG from osteoblasts can act as a decoy receptor for RANKL. This interaction is represented as $(\epsilon + B)^{g_{43}}$. This term includes a very small number, ϵ, to prevent dividing by zero when the osteoblast population is zero, since g_{43} is a negative integer. The second part of the equation represents osteoclast apoptosis with a first order rate constant β_3.

$$\frac{dC}{dt} = \alpha_4 S^{g_{41}} P^{g_{42}} (\epsilon + B)^{g_{43}} \left(1 - \frac{S}{K_s} \right)_+^{g_{44}} - \beta_3 C^{f_{34}} \tag{2}$$

2.2. Model enhancements for Wnt-10b dependence

This section covers the parts of the model that we alter to explicitly account for Wnt-10b changes from stimuli outside the normal bone remodeling cycle. Several parts of the equations remain the same as the Graham 2013 model (Table 2), but with new terms added or parameters adjusted in each of the following equations (Table 5). The adjusted parameters are denoted with the subscript *adj*. As these additions are related to Wnt-10b, we introduce a variable Wnt into the equations that represents the normalized fold change of Wnt-10b present compared to the normal levels of Wnt-10b in the system. Note that if the Wnt-10b levels are normal, the remodeling cycle is normal as well, and Wnt takes on a value of zero. Figure 1 visually describes how Wnt-10b alters the remodeling cycle.

We connect the concentration of Wnt-10b to the remodeling cycle through a Hill function in the form of

$$\pi_{Wnt} = \frac{Wnt}{K_M + Wnt} \tag{3}$$

Table 2: Parameter values and definitions directly from the Graham 2013 model [24]

Parameter	Definition	Value	Units
α_1	Osteoblast embedding rate	0.5	day^{-1}
α_2	Differentiation rate of pre-osteoblast precursors	0.1	day^{-1}
α_3	Pre-osteoblast proliferation rate	0.1	day^{-1}
α_4	Differentiation rate of osteoclast precursors	0.1	day^{-1}
β_1	Differentiation rate of pre-osteoblasts	0.1	day^{-1}
β_2	Osteoblast apoptosis rate	0.1	day^{-1}
δ	Pre-osteoblast apoptosis rate	0.1	day^{-1}
ϵ	Avoid 0 denominator	1	cells
K_s	Critical value of osteocyte population	200	cells
g_{31}	Osteoblast autocrine signaling	1	dimensionless
g_{21}	Osteocyte paracrine signaling of pre-osteoblasts	2	dimensionless
g_{22}	Sclerostin regulation of osteoblastogenesis	1	dimensionless
g_{32}	Pre-osteoblast autocrine signaling	1	dimensionless
g_{41}	Osteocyte paracrine signaling of osteoclasts	1	dimensionless
g_{42}	Pre-osteoblast paracrine signaling of osteoclasts	1	dimensionless
g_{43}	Osteoblast paracrine signaling of osteoclasts	-1	dimensionless
g_{44}	Sclerostin regulation of osteoclastogenesis	1	dimensionless
f_{12}	Pre-osteoblast paracrine signaling of osteoblasts	1	dimensionless
f_{14}	Osteoclast paracrine signaling of osteoblasts	1	dimensionless
f_{23}	Osteoblast autocrine signaling for apoptosis	1	dimensionless
f_{34}	Osteoclast autocrine signaling for apoptosis	1	dimensionless
k_1	Bone resorption rate	0.698331	%volume cells^{-1} day^{-1}
k_2	Balanced bone formation rate	0.015445	%volume cells^{-1} day^{-1}

Note: k_1 was manually adjusted from the rounded value of 0.7 reported in the Graham 2013 model [24] to six significant figures as for k_2 to maintain no steady state change in bone volume at the end of a remodeling cycle (in the absence of Wnt-10b perturbations) when the populations of all cell types return to their initial conditions.

We assume that K_M remains the same for both the pre-osteoblast and osteoblast interactions because they are both from osteoblast lineage. A Hill function relationship is chosen based on the previous models related to Wnt-10b (Table 1). This particular form is chosen so that when the Wnt-10b concentration is not perturbed, π_{Wnt} will take the value of zero, and the model will return to a steady state of no net bone volume change at the end of a remodeling cycle.

2.2.1. Pre-osteoblasts

Eq. 4 describes the dynamics of the pre-osteoblast cell population, P. The first four terms are unchanged from the Graham 2013 model. Pre-osteoblasts differentiate at a rate of α_2 from a large population of stem cells. This differentiation is triggered by the decrease in sclerostin signaling of osteocytes at the beginning of a remodeling cycle . The pre-osteoblast cell population can also be increased by proliferation of existing cells at a rate of α_3. The pre-osteoblast population is decreased by differentiation into osteoblasts regulated by paracrine signals from the osteoclast population and by apoptosis with a first order rate constant of δ. The new terms we include here account for Wnt-10b effects on the pre-osteoblast proliferation and pre-osteoblast differentiation to osteoblasts, both via the factor defined in Eq. 3. The fifth term in Eq. 4 represents the increase in pre-osteoblast proliferation rate due to excess Wnt-10b. The sixth term quantifies Wnt-10b direct effects on pre-osteoblast differentiation independent from osteoclast regulation.

$$\frac{dP}{dt} = \alpha_2 S^{g_{21}} \left(1 - \frac{S}{K_s}\right)_+^{g_{22}} + \alpha_3 P^{g_{32}} \left(1 - \frac{S}{K_s}\right)_+ - \beta_1 P^{f_{12}} C^{f_{14}} - \delta P + \alpha_{3adj} \pi_{Wnt} P - \beta_{1adj} \pi_{Wnt} P \quad (4)$$

2.2.2. Osteoblasts

Eq. 5 describes the dynamics of the osteoblast cell population, B. Pre-osteoblasts mature into osteoblasts at a rate of β_1 with osteoclast regulation and normal Wnt-10b ($Wnt = 0$), but when the amount of Wnt-10b is altered, the maturation rate changes by $\beta_{1adj} \pi_{Wnt}$ as discussed above for Eq. 4. Wnt-10b also changes how fast osteoblasts die. We reduce the osteoblast apoptosis rate constant β_2 by $\beta_{2adj} \pi_{Wnt}$. The final term in Eq. 5 represents the conversion of osteoblasts into osteocytes.

$$\frac{dB}{dt} = \beta_1 P^{f_{12}} C^{f_{14}} + \beta_{1adj} \pi_{Wnt} P - (\beta_2 - \beta_{2adj} \pi_{Wnt}) B^{f_{23}} - \alpha_1 B^{g_{31}} \left(1 - \frac{S}{K_s}\right)_+ \quad (5)$$

2.2.3. Trabecular bone volume

Eq. 6 shows the dynamics of trabecular bone volume, z, at a single remodeling site. Bone volume is reduced by osteoclasts at a rate of k_1. This rate is slightly different than the rate in Graham 2013 due to rounding. In the Graham model, the reported k_1 value results in a slightly unbalanced remodeling cycle (a new steady state results at the end of each cycle and amplified after repeated cycles). We manually adjust k_1 to match the same number of significant figures as k_2 until we obtain a completely balanced remodeling cycle–that is, no steady state change in bone volume at the end of one remodeling cycle (in the absence of Wnt-10b perturbations) when the populations of all cell types return to their initial conditions. Note that

this alteration is not related to a change in Wnt-10b concentration. Osteoblasts build back the bone at a rate of k_2.

$$\frac{dz}{dt} = -k_1 C + k_2 B \tag{6}$$

2.3. Mouse data

This section covers the data used and the assumptions made concerning the data during the development of the model. Data originally presented in graphs was extracted using Plot Digitizer (version 2.6.8) [31], a tool that digitizes the axes and returns the data points based on the pixel locations. We utilized three published data sources from *in vivo* experiments with C57BL/6 mice (Table 3). Two sources from the same lab [16, 17] were used to estimate the parameters, and the third source from another lab [18] was used to validate the model.

2.3.1. Trabecular bone volume

The trabecular $\frac{\text{bone volume}}{\text{total volume}}$ (BV/TV) data was normalized against the control groups using

$$\frac{\text{altered} - \text{baseline}}{\text{baseline}} * 100 = \text{normalized BV/TV} \tag{7}$$

turning BV/TV data into relative change in trabecular bone volume. We refer to "altered BV/TV" to denote mice that were genetically altered to over- or under-produce Wnt-10b, while "baseline BV/TV" denotes unaltered litter mates. We normalized the mouse data for use in the model previously parameterized for humans. We assumed that the normalized BV/TV data is consistent from mice to human. Note that since our model starts at a baseline of 100 percent of normal bone volume, we expect our simulation to produce z that is 100 plus the expected normalized bone volume. We also took into account the different remodeling cycle lengths of our model and mice. Mice have a twelve to fifteen day remodeling cycle, but our model has a 100-day remodeling cycle for humans [32]. We held constant the number of remodeling cycles between mice and our model when comparing them. We assumed that the three month old mice could be modeled as having undergone six cycles and that the six month old mice could be modeled as having undergone twelve cycles. Thus, we ran the mathematical model for either six or twelve 100-day cycles.

2.3.2. Wnt-10b fold change

For our model, we take fold change of Wnt-10b to be

$$\frac{\text{altered levels} - \text{baseline levels}}{\text{baseline levels}} = \text{Normalized Wnt10b fold change} \tag{8}$$

Note that the notation for "altered" and "baseline" remains consistent with the previous section except Wnt-10b is in place of BV/TV. This is done so that the model will produce a balanced remodeling cycle (returning to initial steady state) when $Wnt = 0$, representing a normal baseline level of Wnt-10b.

Table 3: Bone volume data from mice with genetically altered Wnt-10b expression

Normalized Wnt-10b fold change	Age (months)	Altered BV/TV (%)	Baseline BV/TV (%)	Normalized BV/TV	Source
-1	3	4.3	7.4	-29.7	[16]
+1.8	3	8.04	6.35	26.6	[18]
+1.8	6	4.42	3.24	36.6	[18]
+5	3	18.1	10.7	69.2	[16]
+50	6	15.8	3.6	339	[17]

Note: To produce the 50-fold change of Wnt-10b in mice, Bennett et al. [17] followed the procedures in Longo et al. [33].

2.4. Numerical methods

The system of ODEs for the populations of the bone cells and the bone volume in Eqs. 1–2 and 4–6 are solved simultaneously using the `ode15s` function in MATLAB, which is a variable-step, variable-order solver based on numerical differentiation formulas of orders one to five for stiff problems. The relative tolerance is set to 10^{-7}. The time step is set to 551 steps per 100-day cycle. The NonNegative option is turned on, because all of the dependent variables (cell populations and bone volume) must have strictly non-negative values physiologically. The initial conditions are provided in Table 4.

Table 4: Initial conditions for the species tracked in Eqs. 1–2 and 4–6

Symbol	Initial condition	Definition	Units
S	180	Osteocyte population at time t	cells
P	0	Pre-osteoblast population at time t	cells
B	0	Osteoblast population at time t	cells
C	0	Osteoclast population at time t	cells
z	100	Relative bone volume at time t	%

2.5. Parameter estimation

We fit the parameters α_{3adj}, β_{1adj}, β_{2adj}, and K_M using the MATLAB nonlinear least-squares function `lsqcurvefit`. The multicycle effect of Wnt-10b is on bone volume. Thus, we solve the system of ODEs using the numerical methods described in the previous section for multiple cycles. The ODE solver is called

for one remodeling cycle at a time. The solutions from the final time at the end of one cycle are used as the initial conditions for the subsequent cycle, except cell populations that have a value less than one are set to zero to not allow fractional values less than one discrete cell to persist. The process is repeated for an integer number of cycles. The bone volume calculated at the end of the series of remodeling cycles is taken as the model output for the parameter estimation algorithm and is compared to the normalized BV/TV data. The input is the magnitude of normalized Wnt-10b fold change. We assume that the saturation parameter K_M could take on a value between 1 and 100, and we bound the parameter fitting accordingly. We bound β_{1adj} between zero and one based on the magnitudes of the other parameters. We set the value of α_{3adj} to be β_{1adj} plus a value between zero and one that is found by the solver. This makes α_{3adj} greater than β_{1adj} to ensure that the total effect of increasing Wnt-10b on pre-osteoblasts yields a net increase in formation. Since β_{2adj} and β_2 collectively impact osteoblast apoptosis, we check that the net apoptosis rate satisfies

$$\beta_2 - \beta_{2adj}\pi_{Wnt} \geq 0. \tag{9}$$

For `lsqcurvefit` the optimization options are set to have step tolerance of 10^{-14}, function tolerance of 10^{-14}, optimality tolerance of 10^{-6}, and 10,000 maximum function evaluations.

2.6. Code sharing

To enable code reuse, we share our MATLAB code including parameter values, scripts for plotting, and documentation in an open-source software repository [34]. We also include the systems biology markup language (SBML) version of our model for others to reuse readily. This version can be easily converted to Python [35].

3. Results and Discussion

Table 5: Estimated parameter values and definitions

Parameter	Definition	Value	Units
α_{3adj}	Wnt-10b dependent pre-osteoblast proliferation rate	2.45×10^{-1}	day^{-1}
β_{1adj}	Wnt-10b dependent differentiation rate of pre-osteoblasts	2.15×10^{-1}	day^{-1}
β_{2adj}	Wnt-10b dependent osteoblast apoptosis rate	1.28×10^{-3}	day^{-1}
K_M	Half saturation for Wnt-10b	8.24	dimensionless

Parameter estimation resulted in parameters (Table 5) that fit the data [16, 17] with a residual norm of 26.81. Figure 2A shows the `lsqcurvefit` fitting results at the end of six or twelve remodeling cycles. Figure 2B shows the cumulative changes in bone volume after these remodeling cycles are plotted against

varying fold changes in Wnt-10b for six or twelve remodeling cycles. Note that the data points are the same for A and B but are represented on different axes. The the parameter fitting results visually agree with the data well. With fitted parameters, the model gives the bone volume progression over multiple remodeling cycles for the full range of Wnt-10b perturbations explored. The initial conditions for these simulations are provided in Table 4. As Wnt-10b increases, the bone volume also increases.

Figure 2: Fitting and simulation results of Bone volume and Wnt-10b. A) Fitting results for three normalized Wnt-10b fold changes shown in the legend. These specific fold changes were used to parameterize the model by comparison of the final bone volume with the corresponding data points for the Bennett 2005 data [17] and the Bennett 2007 data [16]. B) Simulated normalized BV/TV relationship with normalized Wnt-10b fold change. The parameter fitting simulation results produce an acceptable residual norm of 26.81. Data points were compared with the cumulative final simulated BV/TV after six remodeling cycles (red solid curve) or after twelve remodeling cycles (blue dotted curve). Fold changes of Wnt-10b range from -1 to 50 in this simulation. Note that the data used in A and B is the same but represented on different axes.

To determine if the model is a good predictor of the change in bone volume that occurs when Wnt-10b levels are perturbed, we utilized a separate set of data [18]. The simulation results fall within the error of the data at six remodeling cycles, but overestimate the results at twelve remodeling cycles (Figure 3). This model validation is particularly good for six or fewer remodeling cycles. The shaded area and the error bar of the data at twelve cycles do overlap to a reasonable extent. The offset at later time points could be due to the fact that the mice in the study used for validation had intermittent doses of Wnt-10b via weekly injections, which possibly allowed for fluctuating Wnt-10b levels. On the contrary, the mice were genetically altered in the studies we used for modeling fitting. Thus, the Wnt-10b levels were presumably constant for the genetically altered mice, and the reported levels of Wnt-10b in the injection experiments may not have captured the dynamics. The shaded area in Figure 3 shows the simulation results within the standard deviation of the Wnt-10b reported fold change, assuming that the fold change is constant throughout the duration of the twelve cycles. The bone volume would shift lower if the Wnt-10b levels would be allowed to diminish between injections. Alternatively, the offset could show a decreased sensitivity to Wnt-10b over long term treatment.

Figure 3: Validation of model with data from Roser-Page et al. [18]. The simulation shown is the result of running the model with a 1.8 normalized fold change of Wnt-10b for twelve remodeling cycles. The grey shaded area shows the simulation results for the standard deviation (± 0.6) of the reported Wnt-10b fold change. The data [18] corresponds to six and twelve remodeling cycles. The simulation falls within the error of the data for six remodeling cycles, but overestimates for twelve remodeling cycles.

For each fold change in Wnt-10b shown in Figure 2A, we also determined the corresponding numbers of the activated populations of osteocytes, pre-osteoblasts, osteoblasts, and osteoclasts (Figure 4). The results show a positive correlation with pre-osteoblast and osteoblast populations and a slight negative correlation with osteoclasts (Figure 5). Even though we did not alter the osteoclast equation (Eq. 2), the population is related to the ratio of pre-osteoblasts to osteoblasts. The model predicts that the ratio of pre-osteoblasts to osteoblasts and the ratio of osteoclasts to osteoblasts both decrease with increasing Wnt-10b (Figure 6). The cumulative numbers of cells of each type were determined by integrating the areas under the curves (AUC) shown in Figure 5 using the `trapz` function in MATLAB. The AUC were used for comparison instead of the dynamic number of cells given that the initial conditions for numbers of activated cells of all three types are zero.

Figure 4: Activated cell population dynamic results for (A) osteocytes, (B) pre-osteoblasts, (C) osteoblasts, and (D) osteoclasts at a normalized 50-fold increase in Wnt-10b. The activated cell populations follow identical dynamics for each remodeling cycle while Wnt-10b concentration is constant.

Figure 5: Activated cell population results for a single remodeling cycle with varying Wnt-10b concentrations shown in the legends. (A) The osteocyte dynamics show little change across Wnt-10b concentration. (B) The pre-osteoblast dynamics show a very slight positive correlation with Wnt-10b. (C) The osteoblast dynamics show a positive correlation with Wnt-10b. (D) Osteoclasts decrease with increasing Wnt-10b.

Figure 6: Ratios of the cumulative numbers (area under the curve, AUC) of (A) pre-osteoblasts and (B) osteoclasts to osteoblasts. As Wnt-10b increases, both of the ratios decrease.

The reduction of osteoclasts population has been experimentally tested for mice with a normalized 5-fold increase in Wnt-10b; they found that the change in the number of osteoclasts on the perimeter of the bone was not statistically significant [16]. However, based on the physiology of bone remodeling, it is expected that a change in the ratio of pre-osteoblast and osteoblasts would alter osteoclast populations through the RANK/RANKL/OPG pathway [27, 36, 37]. In this pathway, pre-osteoblasts, osteoblasts, and osteocytes secrete RANKL that binds to RANK receptors on osteoclasts thus increasing osteoclastogenesis. Osteoblasts and osteocytes also secrete OPG, which is a competitive inhibitor of RANKL (Figure 7). As as the osteocyte dynamics are shown to be relatively consistent, OPG levels change due to a change in osteoblast dynamics. It is also important to note that the predicted change in osteoclast population is still small enough that it would be hard to quantify experimentally. Determining the indirect relationship between activated osteoclast population and Wnt-10b would help to further validate the osteoclast population aspect of the model.

Figure 7: RANK/RANKL/OPG pathway where osteoblastic cell populations regulate the osteoclasts (C) cell populations. Pre-osteoblasts (P), osteoblasts (B), and osteocytes(S) secrete RANKL causing an increase in osteoclasts. Osteoblasts and osteocytes also inhibit osteoclastogenesis by secreting OPG, a competitive inhibitor of RANKL.

This model does provide insight into bone metabolism, but it is not without its limitations. The data utilized was normalized mice data, and the remodeling cycle was set to 100 days instead of a typical 200 day human remodeling time. These issues can be resolved if more data becomes available. In the future this model could be expanded to include other direct relationships with cell signaling molecules of interest such as TNF-α (experimentally shown to alter osteoclast formation and activity) and IL-6 (experimentally shown to alter osteoclast formation) [38].

4. Conclusions

Understanding how Wnt-10b alters the dynamics of bone metabolism could lead to new therapeutic targets for osteoporosis. Though there are a few published models focused on bone metabolism, the model developed here provides new insight on how chronic changes in Wnt-10b affect a single bone remodeling cycle or a sequence of cycles. Experimentally Wnt-10b has been shown to increase osteoblastogenesis and decrease osteoblast apoptosis rates. Our model is able to reproduce the expected changes in pre-osteoblasts and osteoblasts and is able to predict changes in bone volume caused by Wnt-10b. Our model also points

to a slight decrease in osteoclasts that is hard to determine experimentally but is mechanistically sound. These results could be used to design future experiments that could further enhance understanding of how Wnt-10b regulates bone metabolism.

Acknowledgment

Research reported in this publication was supported by the National Institute of General Medical Sciences of the National Institutes of Health under award number R35GM133763. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

We would like to thank Dr. Joshua D. Ramsey at Oklahoma State University for the guidance provided as a master's committee member for C.V. Cook. We would also like to thank Dr. Stelios T. Andreadis, Dr. Viviana Monje-Galvan, and Dr. Rudiyanto Gunawan at the University at Buffalo for their manuscript suggestions. Lastly, we would like to thank the other members of the Systems Biomedicine & Pharmaceutics Lab at the University at Buffalo for providing feedback on this project.

References

[1] A. Parfitt, Targeted and nontargeted bone remodeling: relationship to basic multicellular unit origination and progression, Bone 30 (1) (2002) 5–7.

[2] A. M. Parfitt, Osteonal and hemi-osteonal remodeling: the spatial and temporal framework for signal traffic in adult human bone, J. Cell. Biochem. 55 (3) (1994) 273–286.

[3] E. F. Eriksen, Cellular mechanisms of bone remodeling, Rev. Endocr. Metab. Disord. 11 (4) (2010) 219–227.

[4] L. J. Raggatt, N. C. Partridge, Cellular and molecular mechanisms of bone remodeling, J. Biol. Chem. 285 (33) (2010) 25103–25108.

[5] K. S. Houschyar, C. Tapking, M. R. Borrelli, D. Popp, D. Duscher, Z. N. Maan, M. P. Chelliah, J. Li, K. Harati, C. Wallner, S. Rein, D. Pforringer, G. Reumuth, G. Grieb, S. Mouraret, M. Dadras, J. M. Wagner, J. Y. Cha, F. Siemers, M. Lehnhardt, B. Behr, Wnt pathway in bone repair and regeneration - what do we know so far, Front. Cell. Dev. Biol. 6 (2018) 170.

[6] P. Duan, L. F. Bonewald, The role of the wnt/β-catenin signaling pathway in formation and maintenance of bone and teeth, Int. J. Biochem. Cell. Biol. 77 (Pt A) (2016) 23–29.

[7] X. Li, Y. Xiang, F. Li, C. Yin, B. Li, X. Ke, Wnt/β-catenin signaling pathway regulating t cell-inflammation in the tumor microenvironment, Front. Immunol. 10 (2019) 2293.

[8] O. Silva-Garcia, J. J. Valdez-Alarcon, V. M. Baizabal-Aguirre, Wnt/β-catenin signaling as a molecular target by pathogenic bacteria, Front. Immunol. 10 (2019) 2135.

[9] S. G. Pai, B. A. Carneiro, J. M. Mota, R. Costa, C. A. Leite, R. Barroso-Sousa, J. B. Kaplan, Y. K. Chae, F. J. Giles, Wnt/β-catenin pathway: modulating anticancer immune response, J. Hematol. Oncol. 10 (1) (2017) 101.

[10] P. Wend, K. Wend, S. Krum, G. Miranda-Carboni, The role of Wnt-10b in physiology and disease, Acta Physiol. 204 (1) (2012) 34–51.

[11] J. Castrop, S. Verbeek, F. Hofhuis, H. Clevers, Circumvention of tolerance for the nuclear t-cell protein TCF-1 by immunization of TCF-1 knock-out mice, Immunobiology 193 (2-4) (1995) 281–287.

[12] C. Vangenderen, R. M. Okamura, I. Farinas, R. G. Quo, T. G. Parslow, L. Bruhn, R. Grosschedl, Development of several organs that require inductive epithelial-mesenchymal interactions is impaired in LEF-1-deficient mice, Genes Dev. 8 (22) (1994) 2691–2703.

[13] A. M. Tyagi, M. Yu, T. M. Darby, C. Vaccaro, J. Y. Li, J. A. Owens, E. Hsu, J. Adams, M. N. Weitzmann, R. M. Jones, R. Pacifici, The microbial metabolite butyrate stimulates bone formation via T regulatory cell-mediated regulation of WNT10B expression, Immunity 49 (6) (2018) 1116–1131 e7.

[14] M. Terauchi, J. Y. Li, B. Bedi, K. H. Baek, H. Tawfeek, S. Galley, L. Gilbert, M. S. Nanes, M. Zayzafoon, R. Guldberg, D. L. Lamar, M. A. Singer, T. F. Lane, H. M. Kronenberg, M. N. Weitzmann, R. Pacifici, T lymphocytes amplify the anabolic activity of parathyroid hormone through wnt10b signaling, Cell Metab. 10 (3) (2009) 229–40.

[15] M. A. Islam, C. V. Cook, B. J. Smith, A. N. Ford Versypt, Mathematical modeling of the gut-bone axis and implications of butyrate treatment on osteoimmunology, Ind. Engr. Chem. Res. 60 (49) (2021) 17814–17825.

[16] C. N. Bennett, H. Ouyang, Y. L. Ma, Q. Zeng, I. Gerin, K. M. Sousa, T. F. Lane, V. Krishnan, K. D. Hankenson, O. A. MacDougald, Wnt-10b increases postnatal bone formation by enhancing osteoblast differentiation, J. Bone Miner. Res. 22 (12) (2007) 1924–1932.

[17] C. N. Bennett, K. A. Longo, W. S. Wright, L. J. Suva, T. F. Lane, K. D. Hankenson, O. A. MacDougald, Regulation of osteoblastogenesis and bone mass by Wnt-10b, Proc. Natl. Acad. Sci. U. S. A. 102 (9) (2005) 3324–3329.

[18] S. Roser-Page, T. Vikulina, M. Zayzafoon, M. N. Weitzmann, CTLA-4Ig-induced T cell anergy promotes Wnt-10b production and bone formation in a mouse model, Arthritis Rheumatol. 66 (4) (2014) 990–999.

[19] M. M. Zaiss, R. M. Jones, G. Schett, R. Pacifici, The gut-bone axis: how bacterial metabolites bridge the distance, J. Clin. Invest. 129 (8) (2019) 3018–3028.

[20] H. Jing, X. Su, B. Gao, Y. Shuai, J. Chen, Z. Deng, L. Liao, Y. Jin, Epigenetic inhibition of Wnt pathway suppresses osteogenic differentiation of BMSCs during osteoporosis, Cell Death Dis. 9 (2) (2018) 1–12.

[21] S. V. Komarova, R. J. Smith, S. J. Dixon, S. M. Sims, L. M. Wahl, Mathematical model predicts a critical role for osteoclast autocrine regulation in the control of bone remodeling, Bone 33 (2) (2003) 206–215.

[22] S. V. Komarova, Mathematical model of paracrine interactions between osteoclasts and osteoblasts predicts anabolic action of parathyroid hormone on bone, Endocrinology 146 (8) (2005) 3589–3595.

[23] B. P. Ayati, C. M. Edwards, G. F. Webb, J. P. Wikswo, A mathematical model of bone remodeling dynamics for normal bone cell populations and myeloma bone disease, Biol. Direct 5 (1) (2010) 28.

[24] J. Graham, B. Ayati, S. Holstein, J. Martin, The role of osteocytes in targeted bone remodeling: a mathematical model, PLoS ONE 8 (5) (2013) e63884.

[25] C. J. Proctor, A. Gartland, Simulated interventions to ameliorate age-related bone loss indicate the importance of timing. front endocrinol (lausanne). 2016; 7: 61, Front. Endocrinol. 7 (2016) 61.

[26] R. J. Eudy, M. R. Gastonguay, K. T. Baron, M. M. Riggs, Connecting the dots: Linking osteocyte activity and therapeutic modulation of sclerostin by extending a multiscale systems model, CPT: Pharmacometrics Syst. Pharmacol. 4 (9) (2015) 527–536.

[27] M. Martin, V. Sansalone, D. M. L. Cooper, M. R. Forwood, P. Pivonka, Mechanobiological osteocyte feedback drives mechanostat regulation of bone in a multiscale computational model, Biomech. Model. Mechanobiol. 18 (5) (2019) 1475–1496.

[28] V. Lemaire, D. R. Cox, Dynamics of bone cell interactions and differential responses to PTH and antibody-based therapies, Bull. Math. Biol. 81 (9) (2019) 3575–3622.

[29] P. R. Buenzli, P. Pivonka, B. S. Gardiner, D. W. Smith, Modelling the anabolic response of bone using a cell population model, J. Theor. Biol. 307 (2012) 42–52.

[30] A. Farhat, D. Jiang, D. Cui, E. T. Keller, T. L. Jackson, An integrative model of prostate cancer interaction with the

bone microenvironment, Math. Biosci. 294 (2017) 1–14.

[31] Huwaldt, J. A., Plot Digitizer, `https://sourceforge.net/projects/plotdigitizer/files/plotdigitizer/2.6.8/` (2015

(accessed June 12, 2021)).

[32] R. L. Jilka, The relevance of mouse models for investigating age-related bone loss in humans, J. Gerontol. A Biol. Sci.

Med. Sci. 68 (10) (2013) 1209–1217.

[33] K. A. Longo, W. S. Wright, S. Kang, I. Gerin, S.-H. Chiang, P. C. Lucas, M. R. Opp, O. A. MacDougald, Wnt-10b inhibits

development of white and brown adipose tissues, J. Biol. Chem. 279 (34) (2004) 35503–35509.

[34] C. V. Cook, A. N. Ford Versypt, Wnt10bBoneCompartment, http://github.com/ashleefv/Wnt10bBoneCompartment (ac-

cessed 22.02.13). (2022). `doi:10.5281/zenodo.6066256`.

[35] S. M. Ruggiero, A. N. Ford Versypt, SBMLtoODEpy: A software program for converting SBML models into ODE models

in Python, J. Open Source Software 4 (41) (2019) 1643.

[36] J. M. Kim, C. J. Lin, Z. Stavre, M. B. Greenblatt, J. H. Shim, Osteoblast-osteoclast communication and bone homeostasis,

Cells 9 (9).

[37] T. J. Martin, G. A. Rodan, Intercellular Communication during Bone Remodeling, Elsevier, 2008, pp. 547–560.

[38] M. Ponzetti, N. Rucci, Updates on osteoimmunology: what's new on the cross-talk between bone and immune system,

Front. Endocrinol. 10 (2019) 236.