# Mechanosensory trichome cells evoke a mechanical stimuli–induced immune response in plants

Mamoru Matsumura[1,10], Mika Nomoto[1,2,10]*, Tomotaka Itaya[1,2], Yuri Aratani[3], Mizuki Iwamoto[4], Takakazu Matsuura[5], Yuki Hayashi[1], Toshinori Kinoshita[1,6], Izumi C Mori[5], Takamasa Suzuki[7], Shigeyuki Betsuyaku[4,8], Steven H Spoel[9], Masatsugu Toyota[3], Yasuomi Tada[1,2]*

[1]Division of Biological Science, Graduate School of Science, Nagoya University, Aichi, Japan.

[2]Center for Gene Research, Nagoya University, Aichi, Japan.

[3]Department of Biochemistry and Molecular Biology, Saitama University, Saitama, Japan.

[4]Graduate School of Life and Environmental Sciences, University of Tsukuba, Ibaraki, Japan.

[5]Institute of Plant Science and Resources (IPSR), Okayama University, Okayama, Japan.

[6]Institute of Transformative Bio-Molecules (WPI-ITbM), Nagoya University, Aichi, Japan.

[7]Department of Biological Chemistry, College of Bioscience and Biotechnology, Chubu University, Aich, Japan.

[8]Faculty of Life and Environmental Sciences Microbiology Research Center for Sustainability, University of Tsukuba, Ibaraki, Japan.

[9]Institute of Molecular Plant Sciences, School of Biological Sciences, University of Edinburgh, Edinburgh, UK.

[10]These authors contributed equally.

Lead contact

*Correspondence: nomoto@gene.nagoya-u.ac.jp (M.N.); ytada@gene.nagoya-u.ac.jp (Y.T.).

## Abstract

Perception of pathogen-derived ligands by corresponding host receptors is a pivotal strategy in eukaryotic innate immunity. In plants, this is complemented by circadian anticipation of infection timing, promoting basal resistance even in the absence of pathogen threat. Here, we report that trichomes, hair-like structures on the epidermis, directly sense external mechanical forces caused by raindrops to anticipate waterborne infections in *Arabidopsis thaliana*. Exposure of leaf surfaces to mechanical stimuli initiates the concentric propagation of intercellular calcium waves away from trichomes to induce defence-related genes. Propagating calcium waves enable effective immunity against pathogenic microbes through the calmodulin-binding transcription activator 3 (CAMTA3) and mitogen-activated protein kinases. We propose a novel layer of plant immunity in which trichomes function as mechanosensory cells to detect potential risks.

## Introduction

Innate immunity is an evolutionarily conserved front line of defence across the plant and animal kingdoms. In plants, pattern-recognition receptors (PRRs), such as leucine-rich repeat receptor-like kinases (LRR-RLKs) and LRR receptor proteins (LRR-RPs), specifically recognize microbe-associated molecular patterns (MAMPs) as non-self molecules, leading to the activation of pattern-triggered immunity (PTI) to limit pathogen proliferation[1,2]. While adapted pathogens have evolved virulence effectors that can circumvent PTI, plants also deploy disease resistance (*R*) genes, primarily encoding nucleotide-binding LRR proteins, which mount effector-triggered immunity (ETI)[3-5]. ETI often culminates in a hypersensitive response as well as acute and localized cell death induced at the site of infection and accompanied by profound transcriptional changes of defence-related genes to retard pathogen growth[4,5]. These ligand-receptor systems are largely dependent on a transient increase in intracellular calcium concentration ($[Ca^{2+}]_i$), followed by the initiation of phosphorylation-dependent signalling cascades, including mitogen-activated protein kinases (MAPKs) and calcium-dependent protein kinases, that orchestrate a complex transcriptional network and the activity of immune mediators[6,7].

In addition to PTI and ETI, plant immunity can be induced periodically in the absence of pathogen threat, a process that is under the control of the circadian clock and driven by daily oscillations in humidity as well as light-dark cycles[8-10]. Such responses enable plants to prepare for the potential increased risk of infection at the time when microbes are anticipated to be most infectious. Therefore, the anticipation of potentially pathogenic microorganisms through sensing of climatological changes on the one hand and their specific detection on the other constitute two distinct layers of the plant immune system.

Among the climatological factors that affect the outcome of plant-microbe

interactions, rain is a major cause of devastating plant diseases, as fungal spores and bacteria

are spread through rain-dispersed aerosols or ballistic particles splashed from neighbouring

infected plants. In addition, raindrops contain plant pathogens, including *Pseudomonas*

*syringae* and *Xanthomonas campestris*, and negatively regulate stomatal closure, which

facilitates pathogen entry into leaf tissues[11-13]. High humidity, which is usually associated

with rain, enhances the effects of bacterial pathogen effectors, such as HopM1, and

establishes an aqueous apoplast for aggressive host colonization[14]. These findings suggest

that it would be beneficial for plants to recognize rain as an early risk factor for infectious

diseases.

How do plants respond to rain? Rain induces the expression of mechanosensitive

*TOUCH* (*TCH*) genes in plants[15]. Mechanostimulation may affect a variety of plant

physiological processes mediated by hormones such as auxin, ethylene, and gibberellin[16-19].

*Arabidopsis thaliana* seedlings exposed to rain-simulating water spray accumulate the

immune phytohormone jasmonic acid (JA) to promote the expression of JA-responsive

genes[20]. Thus, rain modulates both mechanotransduction and hormone-signalling pathways

that could affect the growth and development of plants as well as environmental responses.

However, the regulatory mechanisms underpinning the rain-activated signalling pathway

have not been fully elucidated.

Here, we report a novel layer of the plant immune system evoked by sensing

mechanostimulation of falling raindrops: trichomes, hair-like cells on the leaf surface,

function as mechanosensory cells that mount an effective immune response against both

biotrophic and necrotrophic pathogens. When trichomes are mechanically stimulated,

intercellular calcium waves are concentrically propagated away from them, and this is

followed by the activation of MAPKs and initiation of the calcium- and calmodulin-binding

transcription activator (CAMTA)-regulated immune response. We propose that plants

directly recognize rain as a risk factor and evoke a rapid immune response that substantially

contributes to early detection of, and protection from, potential pathogens.

## Results and discussion

### Rain and mechanical stimuli induce mechanosensitive genes involved in plant immunity

To investigate the effect of rain on transcriptional changes in *Arabidopsis* leaves, we

performed transcriptome deep sequencing (RNA-seq) of wild-type Columbia (Col-0)

*Arabidopsis* treated with artificial raindrops (Supplementary Fig. 1a; Methods). After

applying only 10 falling droplets, we detected the marked induction of 1,050 genes 15 min

after treatment (Supplementary Table 1). Gene Ontology (GO) analysis of these genes

revealed a striking enrichment in categories associated with plant immunity, as evidenced by

the expression of major immune regulators including *WRKY DNA-BINDING PROTEIN*

(*WRKY*) genes, *CALMODULIN BINDING PROTEIN 60-LIKE* g (*CBP60g*), *MYB DOMAIN*

*PROTEIN* (*MYB*) genes, *ETHYLENE RESPONSE FACTOR* (*ERF*) genes, and *MAP KINASE*

(*MPK*) genes[21,22] (Fig. 1a, Supplementary Table 1, Supplementary Table 2). The touch-

induced genes *TCH2* and *TCH4* were also highly upregulated in response to one falling

raindrop (falling) compared to a water droplet placed directly on the leaf surface (static)

(Supplementary Fig. 1b). These results suggested that mechanosensation is involved in

altering transcriptional activity.

To validate this hypothesis, we mechanically stimulated rosette leaves by gently

brushing them one to ten times along the main veins with a small paint brush (Supplementary

Fig. 1c; Methods) and analysed the expression profile of the immune regulator *WRKY33*,

which was responsive to raindrops. *WRKY33* expression was maximally induced 15 min after

brushing the leaves one to four times (Supplementary Fig. 1d). Next, we compared gene

expression patterns between leaves that were brushed once and those that received 10 falling

raindrops. Both raindrops and brushing strongly upregulated *TCH2*, *TCH4*, *WRKY33*,

*WRKY40*, *WRKY53*, *CBP60g*, *MYB51*, *ERF1*, and *JASMONATE-ZIM-DOMAIN PROTEIN*

*1* (*JAZ1*) expression [15,21,22] (Fig. 1b, c), suggesting that raindrops are likely recognized as a

mechanical stimulus.

Then, to comprehensively identify mechanosensitive genes, we performed an RNA-

seq analysis of leaves brushed once. We identified 1,241 genes that were significantly

induced 15 min after this treatment relative to control plants (Supplementary Table 3). These

mechanical stimuli (MS)-induced genes were primarily categorized as plant immune

responses, such as response to chitin, defence response, and immune system response (Fig.

1d, Supplementary Table 4). We found that 87.3% of raindrop-induced genes and 73.9% of

MS-induced genes overlapped (Fig. 1e): this set of 917 genes expressed upon both treatments

was enriched in GO categories associated with stress responses (Supplementary Fig. 1e).

Furthermore, the expression levels of these 917 genes, including major immune regulators,

were strongly positively correlated between the two treatments (Pearson correlation

coefficient r = 0.917) (Fig. 1f). These transcriptome analyses indicated that falling raindrops

stimulate the expression of mechanosensitive genes involved in environmental stress

responses, including plant immunity.

**Rain and MS rapidly activate plant immune responses**

To further characterize raindrop-induced genes, we conducted a comparative analysis with

published transcriptome datasets. Many raindrop- and MS-induced genes were also

expressed during major plant immune responses, such as those triggered by the immune

phytohormones salicylic acid (SA), which is effective against biotrophic pathogens (21%;

193/917 genes), and JA, which mounts immune responses to necrotrophic pathogens (11.8%;

108/917 genes); the bacterial-derived peptide flg22, which activates PTI (37%; 339/917

genes); and the bacterial pathogen *Pseudomonas syringae* pathovar *maculicola* ES4326 (*Psm*

ES4326) (25.8%; 237/917 genes)[1,2,23-25] (Fig. 2a, b). In total, 58.6% (537/917 genes) of

raindrop- and MS-induced genes overlapped with those induced in response to different

immune elicitors, suggesting that raindrops activate mechanosensitive immune responses.

Since stress-responsive gene expression is either positively or negatively regulated

by phytohormones, we determined the changes in the accumulation levels of six

phytohormones [SA, JA, JA-isoleucine (JA-Ile), abscisic acid (ABA), gibberellic acid 4

(GA$_4$), and indole-3-acetic acid (IAA)] in leaves treated with 10 falling droplets and in those

brushed once. No significant changes in the levels of the phytohormones, except JA and JA-

Ile, were observed 5 min after treatment (Fig. 2c), consistent with the previous report that

water spray induces JA-mediated transcriptional changes[20]. The slight increase in JA and JA-

Ile could explain the observation that only 11.8% of raindrop- and MS-induced genes are JA-

responsive (Fig. 2a). Although 21% of raindrop- and MS-induced genes overlap with SA-

responsive genes (Fig. 2a), SA levels were not significantly increased in response to

raindrops and MS (Fig. 2c). Therefore, most mechanosensitive genes, whose expression is

induced 5 min after treatment with raindrops, are presumably regulated independently of

phytohormonal responses. A previous report demonstrated that GA accumulation is reduced

by "bending" leaves twice per day for 2 weeks[19]. Here, significant changes in GA levels were

not detected upon transient application of raindrops or MS (data not shown). These results

indicated that plants differentially respond to MS depending on their intensity and duration.

Activation of MAPKs is one of the earliest cellular events and a hallmark of plant

immune responses. In particular, PRRs promptly activate a phosphorylation cascade

involving MPK3 and MPK6 in response to MAMPs, whereby the downstream immune

components of PTI are phosphorylated to promote transcriptional reprogramming[1,6,7,26].

Because 37% of raindrop- and MS-induced genes were also upregulated by flg22 treatment

(Fig. 2a), we examined whether a MAPK cascade is activated in responses to raindrops and

MS by immunoblot analysis with the anti-p44/p42 antibody, which detects phosphorylated

MPK3/MPK6[27,28]. Upon treatment of rosette leaves with 4 falling raindrops or MS (1

brushing), phosphorylation of MPK3/MPK6 was induced within 3 min and remained high

for 10 min after each treatment (Fig. 2d, e), indicating that MPK3/MPK6 activation precedes

the expression of mechanosensitive genes detected 10 min after MS application

(Supplementary Fig. 1d). The kinetics of MS-activated MPK3/MPK6 were reminiscent of

those observed upon activation of the PRR protein FLS2 and its coreceptor BRI1-

ASSOCIATED RECEPTOR KINASE 1 (BAK1), which are responsible for recognition of

the bacterial flg22 epitope[1,26]. Wild-type, *fls2*, and *bak1* mutant plants displayed comparable

levels of phosphorylated MPK3/MPK6 in response to MS (Fig. 2f), however, suggesting that

FLS2 and BAK1 are not positively involved in raindrop-elicited mechanotransduction.

We then performed a comparative analysis of raindrop- and MS-induced genes

against published transcriptome datasets describing the specific and conditional activation of

MPK3/MPK6 in transgenic *Arabidopsis* plants carrying a constitutively active variant of

tobacco (*Nicotiana tabacum*) *MAP KINASE Cab 2* (*NtMEK2*) under the control of the

dexamethasone-inducible promoter[28]. Approximately 27.5% (252/917 genes) of both

raindrop- and MS-induced genes were upregulated by MPK3/MPK6[28] (Supplementary Fig.

2a, Supplementary Table 5), and these upregulated genes were highly enriched in categories

associated with plant immunity (Supplementary Fig. 2b, Supplementary Table 6), suggesting

that MAPKs play a critical role in mechanotransduction.

**Rain and MS confer resistance to both biotrophic and necrotrophic pathogens**

We then investigated whether raindrops and MS confer resistance to pathogenic microbes.

Raindrops containing the spores of the necrotrophic pathogen *Alternaria brassicicola* Ryo-1

were placed on fully expanded leaves after pretreatment with raindrops or MS for 3 h at an

interval of 15 min. Both stimuli significantly suppressed lesion development compared to

control plants without pretreatment (Fig. 2g, h). Pretreatment of leaves with MS for 3 h also

efficiently protected plants from infection with the biotrophic pathogen *Psm* ES4326 (Fig.

2i). These results confirmed that mechanostimulation induces a PTI-like response to confer

a broad spectrum of resistance to both biotrophic and necrotrophic pathogens, as MS activate

immune MAPKs and upregulate a large subset of flg22-induced genes. In support of this

argument, exposure to the fungal cell wall, chitin, also upregulated 42.1% (386/917 genes)

of raindrop-induced genes (Supplementary Fig. 2c).

**Mechanosensitive genes are regulated by calmodulin-binding transcription activator 3**

To dissect rain-induced mechanotransduction, we searched for a conserved *cis*-regulatory

element in the promoter sequences of mechanosensitive genes. From an unbiased promoter

analysis of the top 300 genes among 917 differentially expressed genes, we obtained the

highest enrichment for the CGCG box (CGCGT or CGTGT), which is recognized by

calmodulin (CaM)-binding transcription activators (CAMTAs) that are conserved from

plants to mammals[29-33] (Fig. 3a). The *Arabidopsis* transcription factor CAMTA3 (also named

SIGNAL RESPONSIVE1 [SR1]) is a negative regulator of plant immunity because *camta3*

null mutants exhibit constitutive expression of defence-related genes and enhanced resistance

to virulent *P. syringae* infection[34,35]. CAMTA transcription factors possess a CaM-binding

(CaMB) domain and an IQ domain to which CaM binds in a calcium-dependent manner to

negate their function (Supplementary Fig. 3a). Mutation of the IQ domain, such as in

CAMTA3[A855V], suppresses the constitutive expression of defence-related genes seen in the

*camta2 camta3* double mutant when expressed in this background but is no longer regulated

by calcium-mediated responses[36,37]. In agreement with our promoter analysis, 28.7% of

constitutively upregulated genes (309/1,075 genes) in the *camta1 camta2 camta3* triple

mutant overlapped with raindrop- and MS-induced genes detected in wild-type plants[38]

(Supplementary Fig. 3b, Supplementary Table 7). Upon application of raindrops and MS,

*WRKY33* and *CBP60g* transcript levels were significantly reduced in plants expressing the

*CAMTA3[A855V]* variant compared to a *CAMTA3-GFP* transgenic line expressing a transgene

that complemented the phenotype of the *camta2 camta3* mutant (Fig. 3b, c), suggesting that

CAMTA3 is involved in mechanotransduction.

To confirm whether CAMTA3 directly targets mechanosensitive genes, we

investigated the genome-wide distribution of CAMTA3-binding sites by chromatin

immunoprecipitation followed by deep sequencing (ChIP-seq) using *CAMTA3[A855V]-GFP*

plants, as the mutant protein stably represses the transcription of CAMTA3-regulated genes.

With the aid of model-based analysis of ChIP-seq (MACS2) software, we identified 2,641

and 2,728 CAMTA3-binding genes, respectively, in two replicates ($P < 0.05$); about 40% of

these peaks are located in the promoter regions and another 30% in gene bodies

(Supplementary Fig. 3c, Supplementary Table 8). The overlap between the two replicates

highlighted 2,011 CAMTA3-targeted genes that included 272 raindrop- and 297 MS-induced

genes such as *TCH2*, *TCH4*, and *CBP60g* (Fig. 3d), consistent with our hypothesis that

CAMTA3 regulates the transcription of mechanosensitive genes.

To validate the results from the promoter analysis of mechanosensitive genes, we

next investigated specific DNA sequences to which CAMTA3 selectively binds by analysing

CAMTA3-binding peaks by Multiple EM for Motif Elicitation (MEME)-ChIP (Methods).

We again identified the CGCG box (CGCGT or CGTGT) as the motif with the highest

enrichment score ($3.4 \times 10^{-44}$) (Fig. 3e). The subsequent visualization of ChIP-seq profiles

via the Integrative Genomics Viewer (IGV)[39] demonstrated that CAMTA3 is primarily

enriched at the CGCG boxes of mechanosensitive genes, including *TCH2*, *TCH4*, *CAM2*,

*CBP60g*, *CALMODULIN LIKE 23* (*CML23*), and *WRKY40* (Fig. 3f). GO analysis on 314

CAMTA-targeted genes (Fig. 3d, shown in red) to define the biological functions of these

genes showed a significant enrichment in categories related to immune and environmental

responses (Supplementary Fig. 3d, e). We thus investigated whether CAMTA3 is required

for the immune responses. *camta2 camta3 CAMTA3-GFP* transgenic plants effectively

mounted an enhanced disease resistance against *P. syringae* in response to MS, while *camta2*

*camta3 CAMTA3^{A855V}-GFP* plants were significantly compromised in resistance (Fig. 3g).

These results demonstrate that CAMTA3 negatively regulates the plant immune responses

by binding to the CGCG box in raindrop- and MS-induced gene promoters and represses the

expression of these genes.

Since mechanostimulation rapidly activates MPK3/MPK6 (Fig. 2d, e), we

investigated whether CAMTA3 mediates the activation of these MPKs. Using *camta2*

*camta3 CAMTA3-GFP* and *camta2 camta3 CAMTA3^{A855V}-GFP*, we detected the

phosphorylation of MPK3/MPK6 independently of CAMTA3 activity (Supplementary Fig.

3f). In addition, the calcium ionophore A23187 clearly induced the phosphorylation of MPK3

and MPK6 (Supplementary Fig. 3g). These results suggested that the mechanotransduction

initiated by raindrops and MS may cause a $Ca^{2+}$ influx that negates the repressive effect of

CAMTA3 and concomitantly activates the MAPK cascade, as previously proposed[26].

**MS initiates intercellular calcium waves concentrically away from trichomes**

To visualize how mechanostimulation induces the expression of immune genes *in planta*, we

generated *Arabidopsis* transgenic lines with the promoter sequences of *WRKY33* and

*CBP60g* driving the expression of nucleus-targeted enhanced *YELLOW FLUORESCENT*

*PROTEIN* (*YFP-NLS*) (*WRKY33pro:EYFP-NLS* and *CBP60gpro:EYFP-NLS*). *WRKY33*

expression is regulated by both MPK3/MPK6 and CAMTA3 (Supplementary Table 5,

Supplementary Table 8), while *CBP60g* is not mediated by MPK3/MPK6 (Supplementary

Table 5). When half leaves were gently brushed (Supplementary Fig. 1c), we detected YFP

fluorescence in the *WRKY33pro:EYFP-NLS* and *CBP60gpro:EYFP-NLS* transgenic lines as

localized, clustered groups of cells only in the brushed half (Fig. 4a, Supplementary Fig. 4a).

Closer inspection of the stimulated regions revealed that both genes were induced in cells

surrounding trichomes, hair-like structures projecting outward from the epidermal surface

(Fig. 4b, c, Supplementary Fig. 4b, c).

Trichomes function as chemical and physical barriers against insect feeding and are

likely involved in drought tolerance and protection against ultraviolet irradiation[40,41].

Mechanostimulation of a single trichome induces $Ca^{2+}$ oscillations within the proximal skirt

cells that surround the base of trichomes[42], suggesting that the mechanical force could be

focused on only skirt cells (Supplementary Fig. 5). However, since mechanostimulation by

raindrops and MS confers resistance to pathogens in whole leaves, we hypothesized that

trichomes activate a $Ca^{2+}$ signal in a large area of leaves, as shown in Figure 4a.

To visualize changes in cytosolic $Ca^{2+}$ concentrations ($[Ca^{2+}]_{cyt}$) induced by MS on

the leaf surface, we used transgenic *Arabidopsis* expressing the GFP-based $[Ca^{2+}]_{cyt}$ indicator

GCaMP3[43,44]. Leaf brushing induced a marked increase of $[Ca^{2+}]_{cyt}$ in the surrounding leaf

area of trichomes 1 min after stimulation (Fig. 4d, Supplementary Video 1). Flicking a single

trichome with a silver chloride wire triggered an intercellular calcium wave that propagated

concentrically away from the trichome and surrounding skirt cells at a speed of 1.0 μm/s (Fig.

4e, f, Supplementary Video 2). This pattern showed striking consistency with the area of

induction observed with the *WRKY33pro:EYFP-NLS* and *CBP60gpro:EYFP-NLS* reporters

(Fig. 4a, b; Supplementary Fig. 4a, b). The base of trichomes exhibited a rapid and transient

increase in $[Ca^{2+}]_{cyt}$ before the concentric propagation of calcium waves was initiated (Fig.

4g, Supplementary Video 3).

**Trichomes are mechanosensory cells activating plant immunity**

To investigate the possible involvement of trichomes in mechanosensation in *Arabidopsis*

leaves and activation of the immune response, we observed calcium waves using the

knockout mutant of *GLABROUS1* (*GL1*)[45], which lacks trichomes. The *gl1* mutant exhibits

effective basal resistance comparable to that of wild-type Col-0 plants[46], and its local

resistance to *Psm* ES4326 and *A. brassicicola* Ryo-1 is similar to the levels of Col-0 plants

(Supplementary Fig. 6). The mechanostimulation-induced propagation of concentric calcium

waves was compromised in the *gl1* mutant (Fig. 5a, b), confirming that trichomes are true

MS sensors and initiate calcium waves (Supplementary Videos 4, 5). Furthermore,

approximately 70.5% of mechanosensitive genes were expressed in a trichome-dependent

manner (Fig. 5c, Supplementary Fig. 7a, Supplementary Table 9), and transcript levels of 18

representative mechanosensitive immune genes were markedly lower at all time points in the

*gl1* mutant than they were in the wild type in RNA-seq analysis of leaves brushed 4 times

(Fig. 5d), suggesting that trichomes serve as the main sensor of MS. Similarly, compared to

wild-type plants, the transcription of raindrop-induced *WRKY33*, *WRKY40*, and *CBP60g*, as

well as the activation of MPK3/MPK6 by MS, was also significantly reduced in the *gl1*

mutant (Fig. 5e, f, Supplementary Fig. 7b). Moreover, MS-induced resistance against *Psm*

ES4326 infection was abrogated in the *gl1* mutant (Fig. 5g). As with *Psm* ES4326, the

application of MS to wild-type plants prior to inoculation with *A. brassicicola* significantly

limited lesion development, whereas the *gl1* mutant did not show enhanced disease resistance

in response to MS (Fig. 5h).

309           Our work highlights a novel layer of plant immunity that is triggered by an

unexpected function of trichomes as mechanosensory cells. When trichomes are

mechanically stimulated, intercellular calcium waves are concentrically propagated away

from the trichomes, followed by the activation of CAMTA3-dependent immune responses

(Fig. 6). Rapid phosphorylation of MAPKs also is a prerequisite for mechanosensitive gene

expression, as MPK3/MPK6 mediate the phosphorylation of mechanosensitive WRKY33 for

its activation[47,48]. This notion is supported by the finding that the expression of 252 genes

among 917 raindrop- and MS-induced genes is regulated by MPK3/MPK6 (Supplementary

Fig. 2a), and their promoter sequences possess the W-box (TTGACC) for WRKYs as the

most enriched *cis*-regulatory elements (Supplementary Fig. 8). The molecular mechanism by

which calcium mediates the activation of MPK3/MPK6 has yet to be elucidated.

320           Mechanostimulation by repeatedly bending leaves confers resistance to the

necrotrophic pathogen *Botrytis cinerea* via JA accumulation[49]. In addition, a subset of JA-

responsive genes upregulated by water spray is mediated by MYC2/MYC3/MYC4

transcription factors[20]. These observations strongly indicate that mechanosensation causes

profound JA-dependent changes in gene transcription, promoting plant immune responses to

necrotrophic pathogens. The JA- and MYC-dependent pathway does not play a major role in

the expression of mechanosensitive *TCH* genes, however, indicating that

mechanotransduction is regulated by other signaling pathways. Our work demonstrated that

raindrops and MS only partially activate the JA signal but rather strongly induce a PTI-like

response via the $Ca^{2+}$- and CAMTA3-dependent pathway, which is highly effective against

both necrotrophs and biotrophs (Fig. 2g-i). Because rain disseminates diverse pathogens with

different parasitic strategies, including fungi, bacteria, and virus[50,51], it is highly reasonable

that plants perceive raindrops as a risk factor and activate broad-spectrum resistance.

Plants possess mechanosensory cells with a variety of functions, such as flower

antennas of *Catasetum* species for pollination, tentacles of *Drosera rotundifolia* for insect

trapping, root hairs of *Arabidopsis* for water tracking, and red cells of *Mimosa pudica* for

evading herbivores[52]. The carnivorous Venus flytrap (*Dionaea muscipula*) captures insects

by sensing mechanostimulation via sensory hairs on leaf lobes[53]. To monitor diverse MS

applied to plants, several sensing mechanisms have been proposed that include the detection

of cell wall components, distortion of the plasma membrane, and the displacement of the

plasma membrane against the cell wall[54]. In all these systems, a transient increase in $[Ca^{2+}]_{cyt}$

is thought to play a pivotal role in short- and long-term responses. Indeed, two successive

stimulations of sensory hairs of the flytrap are required to meet the threshold of $[Ca^{2+}]_{cyt}$ for

rapid closure of the leaf blade[53,55]. As the trichome on the leaf surface is widely conserved

among many land plants, there may be a common and novel intercellular network of cell-cell

communication that initiates calcium waves for activating immune responses.

**Contact for reagents and subject details**

Further information and requests for reagents may be directed to and will be fulfilled by the corresponding authors Yasuomi Tada (ytada@gene.nagoya-u.ac.jp) and Mika Nomoto (nomoto@gene.nagoya-u.ac.jp).

**Experimental model and subject details**

**Plants**

*Arabidopsis thaliana* accession Columbia-0 (Col-0) was the background for all plants used in this study. *WRKY33pro:EYFP-NLS* (Col-0) and *CBP60gpro:EYFP-NLS* (Col-0) were generated as previously described[56]. *35Spro:GCaMP3* (Col-0), *gl1* [Col(*gl1*)], *camta2 camta3 CAMTA3pro:CAMTA3-GFP*, and *camta2 camta3 CAMTA3pro:CAMTA3^{A855V}-GFP* were previously reported[36,38]. The *Arabidopsis* mutants *fls2* (SALK_093905) and *bak1-3* (SALK_034523) were obtained from the Arabidopsis Biological Resource Center (ABRC). *35Spro:GCaMP3* was introduced into the *gl1* mutant background by crossing. The selection of homozygous lines was performed by genotyping using primers listed in Supplementary Table 10. Plants were grown on soil (peat moss; Super Mix A and vermiculite mixed 1:1) at 22°C under diurnal conditions (16-h-light/8-h-dark cycles) with 50-70% relative humidity. *WRKY33pro:EYFP-NLS* (Col-0) and *CBP60gpro:EYFP-NLS* (Col-0) were sown on soil and grown in a growth room at 23°C in constant light as previously described[56]. *35Spro*:*GCaMP3* (Col-0) and *35Spro*:*GCaMP3* (*gl1*) were grown on Murashige and Skoog (MS) plates [1% (w/v) sucrose, 0.01% (w/v) myoinositol, 0.05% (w/v) MES, and 0.5% (w/v) gellan gum pH 5.8] as previously described[44,57].

**Artificial raindrop treatment**

Reverse osmosis (RO) water was kept in a 500 mL beaker until the water temperature reached

room temperature (22°C). A transfusion set (NIPRO Infusion Set TI-U250P, Nipro, Osaka,

Japan) was installed on a steel stand with the beaker at a height of 1.2 m (H-type Stand I3,

As One, Osaka, Japan) and was adjusted to release 13 µL water droplets (Supplementary Fig.

1a). In this setting, the applied mechanical energy to the leaf surface is equivalent to one in

which 5.8 µL of raindrops reach a terminal velocity of 6.96 m/s[58]. This size raindrop is

frequently observed in nature; thus, the impact of simulated rain is comparable with that of

true rain[58]. The adaxial side of leaves from 4-week-old plants was treated with 10 droplets

for RNA-seq, and 1, 4 or 10 droplets for quantitative RT-PCR (RT-qPCR). The adaxial side

of leaves from 4-week-old plants was treated with one falling or static droplet

(Supplementary Fig. 1b). Sample leaves were collected 15 min after treatment and stored at

-80 °C until use.

**Brush treatment**

The adaxial side of leaves from 4-week-old plants was brushed once for RNA-seq and 4 for

RT-qPCR along the main veins at an angle of 30-40° (KOWA nero nylon drawing pen flat

12, Kowa, Aichi, Japan) (Supplementary Fig. 1c). Sample leaves were collected 15-, 30- and

60 min after treatment for RNA-seq and 15 min for RT-qPCR, and stored at -80°C until use.

**RNA-seq library construction**

Total RNA was extracted from 80-100 mg frozen samples using Sepasol-RNA I Super G

(Nacalai Tesque, Kyoto, Japan) and the TURBO DNase free kit (Thermo Fisher Scientific,

IL, USA) according to the manufacturers' protocols. Total RNA was further purified with

the RNeasy RNA Isolation Kit (QIAGEN, Hilden, Germany) and assessed for quality and

quantity with a Nanodrop spectrophotometer (Thermo Fisher Scientific). We used 1 µg total

RNA for mRNA purification with NEBNext Oligo d(T)$_{25}$ (NEBNext poly(A) mRNA

Magnetic Isolation Module; New England Biolabs, MA, USA), followed by first-strand

cDNA synthesis with the NEBNext Ultra DNA Library Prep Kit for Illumina (New England

Biolabs) and NEBNext Multiplex Oligo for Illumina (New England Biolabs) according to

the manufacturer's protocols. For the analysis of raindrop- and MS-induced gene expression,

the amount of cDNA was determined on an Agilent 4150 TapeStation System (Agilent, CA,

USA). cDNA libraries were sequenced as single-end reads for 81 nucleotides on an Illumina

Nextseq 550 (Illumina, CA, USA). The reads were mapped to the *Arabidopsis thaliana*

reference genome (TAIR10, http://www.arabidopsis.org/) online (BaseSpace, Illumina,

https://basespace.illumina.com/). Pairwise comparisons between samples were performed

with the EdgeR[59] package on the web (Degust, https://degust.erc.monash.edu/). For the

comparative analysis of differentially expressed genes between leaves in the *gl1* mutant and

Col-0, the amount of cDNA was determined by the QuantiFluor dsDNA System (Promega,

WI, USA). cDNA libraries were sequenced as single-end reads for 36 nucleotides on an

Illumina Nextseq 500 (Illumina). The reads were mapped to the *Arabidopsis thaliana*

reference genome (TAIR10) via Bowtie[60] with the options "--all --best --strata". Pairwise

comparisons between samples were performed with the EdgeR package in the R program[59].

Enrichment of GO categories for biological processes was determined using BiNGO

(http://www.psb.ugent.be/cbd/papers/BiNGO/Home.html) ($P < 0.05$)[61].

**Re-analysis of immune-related transcriptome datasets**

We used the following public transcriptome datasets for the comparative analysis with the

RNA-seq data obtained in this study: 10-day-old *Arabidopsis* seedlings treated with 1 μM

flg22 (Array Express; E-NASC-76)[62], 8-day-old *Arabidopsis* seedlings treated with 40 μM

chitin (Gene Expression Omnibus GSE74955), leaves from 4-week-old *Arabidopsis* plants

inoculated with *Pseudomonas syringae* pv. *maculicola* (*Psm*) ES4326 (24 h post inoculation)

(GSE18978), 2-week-old *Arabidopsis* seedlings treated with 0.5 mM SA or 50 µM JA (DNA

Data Bank of Japan DRA003119), 12-day-old *Arabidopsis GVG-NtMEK2$^{DD}$* seedlings

treated with 2 µM DEX for 0 and 6 h (NCBI Sequence Read Archive SRP111959), and 4-

week-old *Arabidopsis camta1 camta2 camta3* triple mutant (GSE43818). The overlaps

between differentially expressed genes in each transcriptome dataset were evaluated as Venn

diagrams (http://bioinformatics.psb.ugent.be/webtools/Venn/).

**RT-qPCR**

Total RNA was extracted from 30-40 mg leaf tissues with Sepasol-RNA I Super G and the

TURBO DNase free kit (Thermo Fisher Scientific) according to the manufacturer's protocols,

followed by reverse transcription with the PrimeScript RT reagent kit (Takara Bio, Shiga,

Japan) using oligo dT primers. RT-qPCR was performed on the first-strand cDNAs diluted

20-fold in water using KAPA SYBR FAST qPCR Master Mix (2x) kit (Roche, Basel,

Switzerland) and gene-specific primers in a LightCycler 96 (Roche). Primer sequences are

listed in Supplementary Table 10.

**Quantification of plant hormones**

The adaxial side of leaves from 4-week-old plants was treated with 10 raindrops (raindrop),

brushed once (MS), and cut (wounding). Sample leaves (0.07-0.1 g) were collected 5 min

after treatment and stored at -80°C until use. SA, JA, JA-Ile, ABA, IAA, and GA$_4$ were

extracted and purified by solid-phase extraction. The contents of these hormones were

quantified using liquid chromatography-electrospray tandem mass spectrometry (LC-ESI-

MS/MS) (triple quadrupole mass spectrometer with 1260 high-performance LC, G6410B;

Agilent Technologies Inc., CA, USA), as previously reported[63].

## ChIP assay

Approximately 0.7 g of 2-week-old *camta2 camta3 CAMTA3pro:CAMTA3^{A855V}-GFP* seedlings was fixed in 25 mL 1% formaldehyde under vacuum for three cycles of 2 min ON/2 min OFF using an aspirator (SIBATA, Tokyo, Japan). Subsequently, 1.5 mL of 2 M glycine was added to quench the cross-linking reaction under vacuum for 2 min. The samples were then washed with 50 mL double-distilled water and stored at -80°C until use. Frozen samples were ground to a fine powder with a mortar and pestle in liquid nitrogen and dissolved in 2.5 mL nuclei extraction buffer (10 mM Tris-HCl pH 8.0, 0.25 M sucrose, 10 mM $MgCl_2$, 40 mM β-mercaptoethanol, protease inhibitor cocktail)[64]. Samples were filtered through two layers of Miracloth (Calbiochem, CA, USA) and centrifuged at 17,700 g at 4°C for 5 min. The pellets were resuspended in 75 μL nuclei lysis buffer [50 mM Tris-HCl pH 8.0, 10 mM EDTA, 1% (w/v) SDS]. After incubation first at room temperature for 20 min and then on ice for 10 min, the samples were mixed with 225 μL ChIP dilution buffer without Triton [16.7 mM Tris-HCl pH 8.0, 167 mM NaCl, 1.2 mM EDTA, 0.01% (w/v) SDS]. Chromatin samples were sonicated for 35 cycles of 30 sec ON/30 sec OFF using a Bioruptor UCW-201 (Cosmo Bio, Tokyo, Japan) to produce DNA fragments, followed by the addition of 375 μL ChIP dilution buffer without Triton, 200 μL ChIP dilution buffer with Triton [16.7 mM Tris-HCl pH 8.0, 167 mM NaCl, 1.2 mM EDTA, 0.01% (w/v) SDS, 1.1% (w/v) Triton X-100], and 35 μL 20% (w/v) Triton X-100. After centrifugation at 17,700 g at 4°C for 5 min, 900 μL solubilized sample was split into two 2.0 mL PROKEEP low-protein-binding tubes (Watson Bio Lab USA, CA, USA) and incubated with 0.75 μL anti-GFP antibody (for immunoprecipitation [IP]) (ab290; Abcam, Cambridge, UK) or Rabbit IgG-Isotype Control (Input) (ab37415; Abcam) for 4.5 h with gentle rocking, and an 18 μL aliquot was used as the input control. Then, samples from *camta2 camta3 CAMTA3pro:CAMTA3^{A855V}-GFP* were mixed with 50 μL of a slurry of Protein A agarose beads (Upstate, Darmstadt, Germany) and

incubated at 4°C for 1 h with gentle rocking. Beads were washed twice with 1 mL low-salt

wash buffer [20 mM Tris-HCl pH 8.0, 150 mM NaCl, 2 mM EDTA, 0.1% (w/v) SDS, 1%

(w/v) Triton X-100], twice with 1 mL high-salt wash buffer [20 mM Tris-HCl pH 8.0, 500

mM NaCl, 2 mM EDTA, 0.1% (w/v) SDS, 1% (w/v) Triton X-100], twice with 1 mL LiCl

wash buffer [10 mM Tris-HCl pH 8.0, 0.25 M LiCl, 1 mM EDTA, 1% (w/v) sodium

deoxycholate, 1% (w/v) Nonidet P-40], and twice with 1 mL TE buffer [10 mM Tris-HCl pH

8.0, 1 mM EDTA]. After washing, beads were resuspended in 100 µL elution buffer [1%

(w/v) SDS, 0.1 M NaHCO$_3$] and incubated at 65°C for 30 min. For the input controls, 41.1

µL TE buffer, 8.7 µL 10% (w/v) SDS, and 21 µL elution buffer were added to 18 µL of each

solubilized sample. Both supernatant and input samples were mixed with 4 µL of 5 M NaCl

and incubated at 65°C overnight to reverse the cross-linking, followed by digestion with 1

µL Proteinase K (20 mg/ml) (Invitrogen, CA, USA) at 37°C for 1 h. ChIP samples were

mixed with 500 µL Buffer NTB and purified using the PCR clean-up gel extraction kit

following the manufacturer's instructions (MACHEREY-NAGEL, Düren, Germany).

**ChIP-seq library construction**

ChIP-seq libraries for the input and two biological replicates were constructed from 2 ng

purified DNA samples with the NEB Ultra II DNA Library Prep Kit for Illumina (New

England Biolabs) according to the manufacturer's instructions. The amount of DNA was

determined on an Agilent 4150 TapeStation System (Agilent). All ChIP-seq libraries were

sequenced as 81-nucleotide single-end reads using an Illumina NextSeq 550 system.

**Analysis of ChIP-seq**

Reads were mapped to the *Arabidopsis thaliana* reference genome (TAIR10,

http://www.arabidopsis.org/) using Bowtie2 with default parameters[60]. The Sequence

Alignment/Map (SAM) file generated by Bowtie2 was converted to a Binary Alignment/Map

(BAM) format file by SAMtools[65]. To visualize mapped reads, Tiled Data Files (TDF) file

were generated from each BAM file using the igvtools package in the Integrative Genome

Browser (IGV)[39]. ChIP-seq peaks were called by comparing the IP with the Input using

Model-based Analysis of ChIP-Seq (MACS2) with the "-p 0.05 -g 1.19e8" option ($P <$

$0.05$)[66]. The peaks were annotated using the nearest gene using the Bioconductor and the

ChIPpeakAnno packages in the R program, from which we identified 2,011 genes detected

in both biological replicates. Enrichment of GO categories of the set of 314 genes overlapping

between raindrop- and MS-induced genes for biological processes was determined using

BiNGO (http://www.psb.ugent.be/cbd/papers/BiNGO/Home.html)[61]. Sequences of the

peaks were extracted from the *Arabidopsis thaliana* genome as FASTA files with Bedtools[67].

To identify the candidates of CAMTA3-binding motifs, the FASTA files were subjected to

MEME (Multiple EM for Motif Elicitation)-ChIP with the default parameter (-meme-minw

6-meme-maxw 10)[68], and a density plot of the distribution of the motifs was generated.

**Immunoblot analysis for detection of MPK3 and MPK6 phosphorylation**

The adaxial side of leaves from 4-week-old plants was brushed four times or treated with

four raindrops, and samples (0.1-0.15 g) were snap-frozen in liquid nitrogen. Total proteins

were extracted in protein extraction buffer [50 mM Tris-HCl pH 7.5, 150 mM NaCl, 2 mM

DTT, 2.5 mM NaF, 1.5 mM $Na_3VO_4$, 0.5% (w/v) Nonidet P-40, 50 mM β-glycerophosphate,

and proteinase inhibitor cocktail] and centrifuged once at 6,000 g, 4°C, for 20 min and twice

at 17,000 g, 4°C for 10 min. The supernatant was mixed with SDS sample buffer [50 mM

Tris-HCl pH 6.8, 2% (w/v) SDS, 5% (w/v) glycerol, 0.02% (w/v) bromophenol blue, and

200 mM DTT] and heated at 70°C for 20 min. The protein samples were subjected to SDS-

PAGE electrophoresis and transferred onto a nitrocellulose membrane (GE Healthcare, IL,

USA). The membrane was incubated with an anti-phospho-p44/42 MAPK polyclonal

antibody (Cell Signalling Technology, MA, USA) (1:1,000 dilution) and goat anti-rabbit

IgG(H+L)-HRP secondary antibody (BIO-RAD, CA, USA) (1:2,000 dilution). The bands for

MPK3/6 were visualized using chemiluminescence solution mixed 5:1 with ImmunoStar

Zeta (FUJIFILM Wako Chemicals, Osaka, Japan) and SuperSignal West Dura Extended

Duration Substrate (Thermo Fisher Scientific). The Rubisco bands were stained with

Ponceau S (Merck Sharp & Dohme Corp., NJ, USA) as a loading control. The

phosphorylation levels of MPK3 and MPK6 were quantified with the blot analysis plug-in in

ImageJ (https://imagej.nih.gov/ij/).

**Treatment with the calcium ionophore A23187**

Twelve-day-old Col-0 seedling was treated with 50 µM calcium ionophore A23187 for 15-,

30- and 60 min. Samples were processed for the phosphorylation of MPK3 and MPK6 as

described in the "Immunoblot analysis for detection of MPK3 and MPK6 phosphorylation"

section. The leaf tissue was stored at -80°C until use.

**Promoter-reporter imaging**

The 3.0-kbp promoters for *WRKY33* and *CBP60g*, both of which covered the previously

analyzed respective regulatory sequences, were amplified from Col-0 genomic DNA by PCR

and cloned into the pENTR/D-TOPO vector (Invitrogen). The promoter regions were

recombined using Gateway technology into the binary vector pBGYN. The resulting

pBGYN-pWRKY33-YFP-NLS and pBGYN-pCBP60g-YFP-NLS vectors were introduced

into *Agrobacterium tumefaciens* GV3101 (pMP90) and then into *Arabidopsis* Col-0 plants

using the floral dip method. A representative homozygous line was selected for each

construct for further detailed analyses.

Promoter-reporter imaging was performed using an MA205FA automated

stereomicroscope (Leica Microsystems, Wetzlar, Germany) and DFC365FX CCD camera

(Leica Microsystems) in 12-bit mode. Chlorophyll autofluorescence and YFP fluorescence

were detected through Texas Red (TXR) (excitation 560/40 nm, extinction 610 nm) and YFP

(excitation 510/20 nm, extinction 560/40 nm) filters (Leica Microsystems). To image

fluorescence emanating from the *WRKY33pro:EYFP-NLS* (Col-0) and *CBP60gpro:EYFP-*

*NLS* (Col-0) plants[56], the leaves of 3-week-old *Arabidopsis* plants were brushed 10 times at

an interval of 15 min for 2 h or left untreated.

**Promoter analysis**

The statistical analysis for overrepresented transcriptional regulatory elements across

transcriptome datasets described above was calculated using a prediction program as

previously reported[32]. The *P* values were calculated using Statistical Motif Analysis in

Promoter                    or                    Upstream                    Sequences

(https://www.arabidopsis.org/tools/bulk/motiffinder/index.jsp). Figures of promoter motif

sequences are generated with WebLogo (https://weblogo.berkeley.edu/logo.cgi).

**Real-time $[Ca^{2+}]_{cyt}$ imaging**

We used 4-week-old and 3-week-old plants expressing the GFP-based cytosolic $Ca^{2+}$

concentration ($[Ca^{2+}]_{cyt}$) indicator GCaMP3[43,44]. To image the fluorescence from the

GCaMP3 reporter (in Col-0 and *gl1*) in whole leaves, the adaxial sides of leaves from 4-

week-old plants were brushed. To monitor the calcium waves propagating from trichomes, a

single trichome from a 2-week-old seedling was flicked with a silver chloride wire. Samples

were imaged with a motorized fluorescence stereomicroscope (SMZ-25; Nikon, Tokyo,

Japan) equipped with a 1× objective lens (NA = 0.156, P2-SHR PLAN APO; Nikon) and an

sCMOS camera (ORCA-Flash 4.0 V2; Hamamatsu Photonics, Shizuoka, Japan) as

described[44].

**Propidium iodide staining**

A stock solution of 10 mM propidium iodide (PI) was prepared with phosphate-buffered

saline (PBS). Rosette leaves of 4-week-old Col-0 plants were cut into 5 mm squares, floated

in a glass petri dish with 20 µM PI solution, and incubated for 1 h at room temperature.

Stained tissues were observed under the all-in-one fluorescence microscope (BZ-X800;

KEYENCE CORPORATION, Osaka, Japan) equipped with a 20x objective lens (CFI S Plan

Fluor LWD ADM 20XC, Nikon) and TRITC dichroic mirror (excitation 545/25 nm,

extinction 605/70 nm) (KEYENCE).

**Bacterial infection**

568    MS were applied to the adaxial leaf surface of 4-week-old plants by brushing 4 times at an

569    interval of 15 min for 3 h. Sample leaves were then inoculated by infiltration, using a plastic

570    syringe (Terumo Tuberculin Syringe 1 mL; TERUMO), with *Psm* ES4326 (OD$_{600}$ = 0.001)

571    resuspended in 10 mM MgCl$_2$. Bacterial growth was measured 2 days after inoculation as

572    described previously[69].

**Fungal infection**

*Alternaria brassicicola* strain Ryo-1 was cultured on 3.9% (w/v) potato dextrose agar plates

(PDA; Becton, Dickinson and Company, NJ, USA) for 4-20 days at 28°C in the dark. After

incubation of the agar plates for 3-7 days under ultraviolet C light, a conidial suspension of

*A. brassicicola* was obtained by mixing with RO water[70]. The adaxial side of leaves from 4-

week-old plants was treated with 10 droplets or MS by brushing at an interval of 15 min for

3 h, followed by spotting with 5 µL conidia suspension (2 x 10$^5$ per mL) of *A. brassicicola*

on the adaxial side of leaves. Inoculated plants were placed at 22°C under diurnal conditions

(16-h-light/8-h-dark cycles) with 100% relative humidity. The lesion size of fungal infection

was measured with ImageJ 3 days after inoculation.

**Statistics and reproducibility**

GraphPad Prism 9 (GraphPad software, CA, USA) was used for all statistical analyses. Two-

sided one-way analysis of variance (one-way ANOVA) or two-way analysis of variance

(two-way ANOVA) was used for multiple comparisons. Unless stated otherwise, sample size

n represents technical replicates. In RT-qPCR, n ≥ 3; in bacterial growth assays, n = 8; in

real-time $[Ca^{2+}]_{cyt}$ imaging assays of *35Spro:GCaMP3* (Col-0) and *35Spro:GCaMP3* (*gl1*),

n = 14 and 9, respectively; and in fungal disease propagation assays, n = 29 (Fig. 2g) and n

= 15 (the other figures). All experiments were performed at least three times with similar

results (biological replicates). In all graphs, asterisks indicate statistical significance tested

by Student's *t* test (two groups) or one/two-way ANOVA (multiple groups).

**Reporting summary**

Further information on research design is available in the Nature Research Reporting

Summary linked to this article.

**Date availability**

The authors declare that all data supporting the findings of this study are available within this

article and its Supplementary Information files. RNA-seq and ChIP-seq data have been

deposited in the DDBJ Sequence Read Archive (DRA) at the DNA Data Bank (DDBJ;

http://www.ddbj.nig.ac.jp/) through the accession numbers DRA011970, DRA009248 and

DRA011123.

## References

1. Couto, D. & Zipfel, C. Regulation of pattern recognition receptor signalling in plants. *Nat. Rev. Immunol.* **16**, 537-552 (2016).

2. Zipfel, C. et al. Bacterial disease resistance in *Arabidopsis* through flagellin perception. *Nature* **428**, 764-767 (2004).

3. Chisholm, S. T., Coaker, G., Day, B. & Staskawicz, B. J. Host-microbe interactions: shaping the evolution of the plant immune response. *Cell* **124**, 803-814 (2006).

4. Cui, H., Tsuda, K. & Parker, J. E. Effector-triggered immunity: from pathogen perception to robust defense. *Annu. Rev. Plant Biol.* **66**, 487-511 (2015).

5. Jones, J. D. & Dangl, J. L. The plant immune system. *Nature* **444**, 323-329 (2006).

6. Boudsocq, M. et al. Differential innate immune signalling via Ca(2+) sensor protein kinases. *Nature* **464**, 418-422 (2010).

7. Macho, A. P. & Zipfel, C. Plant PRRs and the activation of innate immune signaling. *Mol. Cell* **54**, 263-272 (2014).

8. Mwimba, M. et al. Daily humidity oscillation regulates the circadian clock to influence plant physiology. *Nat. Commun.* **9**, 4290 (2018).

9. Wang, W. et al. Timing of plant immune responses by a central circadian regulator. *Nature* **470**, 110-114 (2011).

10. Zhou, M. et al. Redox rhythm reinforces the circadian clock to gate immune response. *Nature* **523**, 472-476 (2015).

11. Madden, L. V. Effects of rain on splash dispersal of fungal pathogens. *CANADIAN J. Plant Pathol.* **19**, 225-230 (1997).

12. Schwartz, H. F., Otto, K. L. & Gent, D. H. Relation of temperature and rainfall to development of Xanthomonas and Pantoea leaf blights of onion in Colorado. *Plant Dis.* **87**, 11-14 (2003).

13. Melotto, M., Underwood, W., Koczan, J., Nomura, K. & He, S. Y. Plant stomata function in innate immunity against bacterial invasion. *Cell* **126**, 969-980 (2006).

14. Xin, X. F. et al. Bacteria establish an aqueous living space in plants crucial for virulence. *Nature* **539**, 524-529 (2016).

15. Braam, J. & Davis, R. W. Rain-, wind-, and touch-induced expression of calmodulin and calmodulin-related genes in Arabidopsis. *Cell* **60**, 357-364 (1990).

16.    Jaffe, M. J. Thigmomorphogenesis: The response of plant growth and development
to mechanical stimulation : With special reference to *Bryonia dioica*. *Planta* **114**, 143-
157 (1973).

17.    Sanyal, D. & Bangerth, F. Stress induced ethylene evolution and its possible
relationship to auxin-transport, cytokinin levels, and flower bud induction in shoots
of apple seedlings and bearing apple trees. *Plant Growth Regul.* **24**, 124-134 (1998).

18.    Johnson, K. A., Sistrunk, M. L., Polisensky, D. H. & Braam, J. *Arabidopsis thaliana*
responses to mechanical stimulation do not require ETR1 or EIN2. *Plant Physiol.* **116**,
643-649 (1998).

19.    Lange, M. J. & Lange, T. Touch-induced changes in *Arabidopsis* morphology
dependent on gibberellin breakdown. *Nat. Plants* **1**, 14025 (2015).

20.    Van Moerkercke, A. et al. A MYC2/MYC3/MYC4-dependent transcription factor
network regulates water spray-responsive gene expression and jasmonate levels. *Proc.*
*Natl. Acad. Sci. U S A* **116**, 23345-23356 (2019).

21.    Li, B., Meng, X., Shan, L. & He, P. Transcriptional regulation of pattern-triggered
immunity in plants. *Cell Host Microbe* **19**, 641-650 (2016).

22.    Pandey, S. P. & Somssich, I. E. The role of WRKY transcription factors in plant
immunity. *Plant Physiol.* **150**, 1648-1655 (2009).

23.    Dong, X., Mindrinos, M., Davis, K. R. & Ausubel, F. M. Induction of Arabidopsis
defense genes by virulent and avirulent Pseudomonas syringae strains and by a cloned
avirulence gene. *Plant Cell* **3**, 61-72 (1991).

24.    Fu, Z. Q. & Dong, X. Systemic acquired resistance: turning local infection into global
defense. *Annu. Rev. Plant Biol.* **64**, 839-863 (2013).

25.    Howe, G. A. & Jander, G. Plant immunity to insect herbivores. *Annu. Rev. Plant Biol.*
**59**, 41-66 (2008).

26.    Tena, G., Boudsocq, M. & Sheen, J. Protein kinase signaling networks in plant innate
immunity. *Curr. Opin. Plant Biol..* **14**, 519-529 (2011).

27.    Galletti, R., Ferrari, S. & De Lorenzo, G. Arabidopsis MPK3 and MPK6 play
different roles in basal and oligogalacturonide- or flagellin-induced resistance against
*Botrytis cinerea*. *Plant Physiol.* **157**, 804-814 (2011).

28.    Su, J. et al. Active photosynthetic inhibition mediated by MPK3/MPK6 is critical to
effector-triggered immunity. *PLoS Biol.* **16**, e2004122 (2018).

29.   Bouche, N., Scharlat, A., Snedden, W., Bouchez, D. & Fromm, H. A novel family of
calmodulin-binding transcription activators in multicellular organisms. *J. Biol. Chem.*
**277**, 21851-21861 (2002).

30.   Doherty, C. J., Van Buskirk, H. A., Myers, S. J. & Thomashow, M. F. Roles for
*Arabidopsis* CAMTA transcription factors in cold-regulated gene expression and
freezing tolerance. *Plant Cell* **21**, 972-984 (2009).

31.   Finkler, A., Ashery-Padan, R. & Fromm, H. CAMTAs: calmodulin-binding
transcription activators from plants to human. *FEBS Lett.* **581**, 3893-3898 (2007).

32.   Yamamoto, Y. Y. et al. Identification of plant promoter constituents by analysis of
local distribution of short sequences. *BMC Genomics* **8**, 67 (2007).

33.   Yang, Y. et al. Genome-wide identification of *CAMTA* gene family members in
*Medicago truncatula* and their expression during root nodule symbiosis and hormone
treatments. *Front. Plant Sci.* **6**, 459 (2015).

34.   Du, L. et al. Ca(2+)/calmodulin regulates salicylic-acid-mediated plant immunity.
*Nature* **457**, 1154-1158 (2009).

35.   Galon, Y. et al. Calmodulin-binding transcription activator (CAMTA) 3 mediates
biotic defense responses in *Arabidopsis*. *FEBS Lett.* **582**, 943-948 (2008).

36.   Kim, Y. S. et al. CAMTA-mediated regulation of salicylic acid immunity pathway
genes in Arabidopsis exposed to low temperature and pathogen infection. *Plant Cell*
**29**, 2465-2477 (2017).

37.   Nie, H. et al. SR1, a calmodulin-binding transcription factor, modulates plant defense
and ethylene-induced senescence by directly regulating NDR1 and EIN3. *Plant*
*Physiol.* **158**, 1847-1859 (2012).

38.   Kim, Y., Park, S., Gilmour, S. J. & Thomashow, M. F. Roles of CAMTA transcription
factors and salicylic acid in configuring the low-temperature transcriptome and
freezing tolerance of Arabidopsis. *Plant J.* **75**, 364-376 (2013).

39.   Thorvaldsdottir, H., Robinson, J. T. & Mesirov, J. P. Integrative Genomics Viewer
(IGV): high-performance genomics data visualization and exploration. *Brief*
*Bioinformatics* **14**, 178-192 (2013).

40.   Gutschick, V. P. Biotic and abiotic consequences of differences in leaf structure. *New*
*Phytol.* **143**, 3-18 (2002).

41.   Wagner, G. J., Wang, E. & Shepherd, R. W. New approaches for studying and
exploiting an old protuberance, the plant trichome. *Ann. Bot.* **93**, 3-11 (2004).

42.  Zhou, L. H. et al. The Arabidopsis trichome is an active mechanosensory switch. *Plant Cell Environ.* **40**, 611-621 (2017).

43.  Tian, L. et al. Imaging neural activity in worms, flies and mice with improved GCaMP calcium indicators. *Nat. Methods* **6**, 875-881 (2009).

44.  Toyota, M. et al. Glutamate triggers long-distance, calcium-based plant defense signaling. *Science* **361**, 1112-1115 (2018).

45.  Larkin, J. C., Oppenheimer, D. G., Lloyd, A. M., Paparozzi, E. T. & Marks, M. D. Roles of the GLABROUS1 and TRANSPARENT TESTA GLABRA genes in Arabidopsis trichome development. *Plant Cell* **6**, 1065-1076 (1994).

46.  Xia, Y. et al. The *glabra1* mutation affects cuticle formation and plant responses to microbes. *Plant Physiol.* **154**, 833-846 (2010).

47.  Mao, G. et al. Phosphorylation of a WRKY transcription factor by two pathogen-responsive MAPKs drives phytoalexin biosynthesis in *Arabidopsis*. *Plant Cell* **23**, 1639-1653 (2011).

48.  Tsuda, K. & Somssich, I. E. Transcriptional networks in plant immunity. *New Phytol.* **206**, 932-947 (2015).

49.  Chehab, E. W., Yao, C., Henderson, Z., Kim, S. & Braam, J. *Arabidopsis* touch-induced morphogenesis is jasmonate mediated and protects against pests. *Curr. Biol.* **22**, 701-706 (2012).

50.  Bauer, H. et al. The contribution of bacteria and fungal spores to the organic carbon content of cloud water, precipitation and aerosols. *Atmospheric Research* **64**, 109-119 (2002).

51.  Prussin, A. J. 2nd, Marr, L. C. & Bibby, K. J. Challenges of studying viral aerosol metagenomics and communities in comparison with bacterial and fungal aerosols. *FEMS Microbiol. Lett.* **357**, 1-9 (2014).

52.  Huynh, T.-P. & Haick, H. Learning from an intelligent mechanosensing system of plants. *Adv. Mater. Technol.* **4**, 1800464 (2019).

53.  Scherzer, S., Federle, W., Al-Rasheid, K. A. S. & Hedrich, R. Venus flytrap trigger hairs are micronewton mechano-sensors that can detect small insect prey. *Nat. Plants* **5**, 670-675 (2019).

54.  Bacete, L. & Hamann, T. The Role of mechanoperception in plant cell wall integrity maintenance. *Plants (Basel, Switzerland)* **9**,   (2020).

55.  Suda, H. et al. Calcium dynamics during trap closure visualized in transgenic Venus
flytrap. *Nat. Plants* **6**, 1219-1224 (2020).

56.  Betsuyaku, S. et al. Salicylic acid and jasmonic acid pathways are activated in
spatially different domains around the infection site during effector-triggered
immunity in *Arabidopsis thaliana. Plant Cell Physiol.* **59**, 8-16 (2018).

57.  Murashige, T. & Skoog, F. A revised medium for rapid growth and bio assays with
Tobacco tissue cultures. *Physiol. Plant.* **15**, 473-497 (1962).

58.  Gunn, R. & Kinzer, G. D. The terminal velocity of fall for water droplet in stagnant
air. *J. Atmospheric Sci.* **6**, 243-248 (1949).

59.  Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package
for differential expression analysis of digital gene expression data. *Bioinformatics* **26**,
139-140 (2010).

60.  Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat.*
*Methods* **9**, 357-359 (2012).

61.  Maere, S., Heymans, K. & Kuiper, M. BiNGO: a Cytoscape plugin to assess
overrepresentation of gene ontology categories in biological networks.
*Bioinformatics* **21**, 3448-3449 (2005).

62.  Denoux, C. et al. Activation of defense response pathways by OGs and flg22 elicitors
in *Arabidopsis* seedlings. *Mol. Plant* **1**, 423-445 (2008).

63.  Matsuura, T., Mori, I. C., Himi, E. & Hirayama, T. Plant hormone profiling in
developing seeds of common wheat (*Triticum aestivum L.*). *Breeding Science* **69**, 601-
610 (2019).

64.  Yamaguchi, N. et al. PROTOCOLS: Chromatin immunoprecipitation from
Arabidopsis tissues. *The arabidopsis book* **12**, e0170 (2014).

65.  Li, H. et al. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078-2079 (2009).

66.  Zhang, Y. et al. Model-based Analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, R137
(2008).

67.  Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics* **26**, 841-842 (2010).

68.  Machanick, P. & Bailey, T. L. MEME-ChIP: motif analysis of large DNA datasets.
*Bioinformatics* **27**, 1696-1697 (2011).

69. Cao, H., Bowling, S. A., Gordon, A. S. & Dong, X. Characterization of an Arabidopsis mutant that is nonresponsive to inducers of systemic acquired resistance. *Plant Cell* **6**, 1583-1592 (1994).

70. Hiruma, K. et al. *Arabidopsis ENHANCED DISEASE RESISTANCE 1 is required for* pathogen-induced expression of plant defensins in nonhost resistance, and acts through interference of *MYC2*-mediated repressor function. *Plant J.* **67**, 980-992 (2011).

**Acknowledgements**

We thank M. F. Thomashow for the seeds of CAMTA3 variants. This work was supported by JSPS KAKENHI Grant Numbers JP23120520, JP25120718, and JP18K19334 (to Y.T.), JP15H05955 (to Y.T. and T.K.), JP19H05363, and JP21H00366 (to M.N.).

**Author contributions**

M.M., M.N., S.H.S., and Y.T. designed the research. M.M. and T.I. established the artificial rain device. M.M. optimized the protocols for artificial raindrop and brush treatment. M.M. and M.N. constructed the Illumina sequencing libraries for RNA-seq and ChIP-seq. T.S. performed RNA-seq and analysis. M.N. performed the ChIP-seq and analysis of CAMTA3. T.M. and I.M. performed the quantification of phytohormones. M.M., Y.H., and T.K. performed the detection of MPK3 and MPK6 phosphorylation. Y.A. and M.T. generated *35Spro*:*GCaMP3* (*gl1*) plants, and M.N., M.T., and Y.A. visualized real-time $[Ca^{2+}]_{cyt}$. M.I. and S.B. generated the transgenic lines *WRKY33pro:EYFP-NLS* (Col-0) and *CBP60gpro:EYFP-NLS* (Col-0). M.M., M.I., and S.B. performed promoter-reporter imaging. M.N. and M.M. performed the rest of the experiments. M.M., M.N., S.H.S., and Y.T. wrote the manuscript with input from all authors.

## Competing interests

The authors declare no competing interests.

**Fig. 1 | Raindrop-induced gene expression highly correlates with mechanical stimuli (MS)-induced gene expression in *Arabidopsis*. a** Enriched Gene Ontology (GO) categories of 1,050 raindrop (10 droplets)-induced genes in the wild type (Col-0). The top 10 categories are shown in ascending order of *P* values. **b**, **c** Transcript levels of MS-induced and defence-related genes in 4-week-old Col-0 plants 15 min after being treated with 10 falling droplets (raindrop, **b**) or 1 brushing (MS, **c**), determined by RT-qPCR and normalized to *UBIQUITIN 5* (*UBQ5*). Data are presented as mean ± SD. **d** Enriched GO categories of 1,241 MS (1 brushing)-induced genes in Col-0. The top 10 categories are shown in ascending order of *P* values. **e** Venn diagram of the overlap between transcriptome datasets from raindrop- and MS-induced genes (*P* < 0.05). **f** Radar chart of intensity compared with mock (log₂FC) and

803    Pearson correlation coefficient (r = 0.917288423) of 917 raindrop- and MS-induced genes

804    (left). Intensities of major immune regulator genes induced by raindrops and MS in RNA-

805    seq analysis ($log_2FC$) (right).

**Fig. 2 | Raindrop-induced mechanosensation triggers defence responses in *Arabidopsis.***
**a**, **b** Venn diagram (**a**) and the upset plot (**b**) between 917 raindrop- and MS-induced genes and transcriptome datasets obtained from salicylic acid (SA), jasmonic acid (JA), and flg22 (PAMP) treatment and *Psm* ES4326 infection ($P < 0.05$). Overlap with raindrop- and MS-induced genes: SA, 21%, 193/917 genes; JA, 11.8%, 108/917 genes; flg22, 37%, 339/917 genes; *Psm* ES4326, 25.8%, 237/917 genes; any of the four factors, 58.6%, 537/917 genes. **c** Fresh weight (ng/g) of plant hormones SA, JA, JA-isoleucine (JA-Ile), abscisic acid (ABA), and indole-3-acetic acid (IAA) 5 min after 10 falling droplets (raindrop), 1 brushing (MS), and cutting (wounding). **d**, **e** Raindrop (4 droplets)- (**d**) and MS (4 brushing)-induced (**e**) MAPK activation in Col-0. Total proteins were extracted from 4-week-old plants treated with raindrops and detected by immunoblot analysis with anti-p44/42 MAPK antibodies. Relative phosphorylation levels are shown below each blot. **f** MS-induced MAPK activation in Col-0, *fls2*, and *bak1-3*. Total proteins were extracted from 4-week-old plants after 5 min of MS treatment (1 brushing) and detected by immunoblot analysis with anti-p44/42 MAPK antibodies. Relative phosphorylation levels are shown below each blot. **g, h** Disease progression of *Alternaria brassicicola* in Col-0 leaves 3 days after inoculation with (+) or without (–) raindrop (10 droplets) pretreatment (**g**) or with (+) or without (–) MS (4 brushing) pretreatment (**h**). Error bars represent SE. Asterisks indicate significant difference (one-way ANOVA; ****$P < 0.0001$). **i** Growth of *Psm* ES4326 in Col-0 leaves 2 days after inoculation with (+) or without (–) MS (4 brushing) pretreatment. An outline of the experiment is provided at left. Error bars represent SE. Asterisks indicates significant difference (one-way ANOVA; ****$P < 0.0001$). Cfu, colony-forming units.

**Fig. 3 | MS-induced genes are regulated by CAMTA3. a** Promoter analysis of the top 300
(among 917 genes) raindrop- and MS-induced genes in terms of expression levels revealed
that the CAMTA-binding CGCG box [CGC(/T)GT] were overrepresented among these genes.
**b**, **c** Transcript levels of *WRKY33* and *CBP60g* in 4-week-old *camta2 camta3*
*CAMTA3pro:CAMTA3-GFP* (*CAMTA3*) and *camta2 camta3 CAMTA3pro:CAMTA3^{A855V}*-
*GFP* [*CAMTA3*(*A855V*)] plants 15 min after the plants were treated with 1 falling droplet (**b**)

or brushed 4 times (**c**), determined by RT-qPCR and normalized to *UBQ5* transcript levels.

Data are presented as mean ± SD. **d** Venn diagram depicting the overlap between genes with

CAMTA3-binding sites in their promoters, as determined by ChIP-seq, and raindrop- and

MS-induced genes as determined by RNA-seq. A total of 314 genes, shown in red, were

identified as CAMTA3-target genes. **e** The CGCG box was identified as an overrepresented

motif among the sequence peaks of 314 genes by MEME-ChIP. **f** Localization of CAMTA3

on the promoters of the MS-induced genes *TCH2*, *TCH4*, *CAM2*, *CBP60g*, *CML23*, and

*WRKY40*, as representative of the 314 genes shown in (**d**). Blue and red lines indicate

CGCGT and CGTGT, respectively. **g** Growth of *Psm* ES4326 in *camta2 camta3*

*CAMTA3pro:CAMTA3-GFP* (*CAMTA3*) and *camta2 camta3 CAMTA3pro:CAMTA3$^{A855V}$*-

*GFP* [*CAMTA3*(*A855V*)] plants 2 days after inoculation with (+) or without (–) MS (4

brushing) pretreatment. Error bars represent SE. Asterisks indicate significant difference

(one- and two-way ANOVA; ****$P < 0.0001$). Cfu, colony-forming units. NS, not significant.

**Fig. 4 | Trichomes initiate intercellular calcium waves. a–c** YFP fluorescence in a whole
leaf from *WRKY33pro:EYFP-NLS* (Col-0) with (MS, bottom half) or without brushing
(untreated, top half) (**a**), along with zoomed-in views of brushed (**b**) and untreated (**c**) areas.
Arrowheads indicate trichomes (**b, c**). Scale bars, 0.5 mm (**a**), 0.3 mm (**b, c**). **d** $Ca^{2+}$ imaging
using *35Spro:GCaMP3* (Col-0). The leaf surface of a 4-week-old plant was treated with MS
by brushing. MS-induced intercellular calcium waves propagated concentrically from

trichomes. Scale bar, 1.0 mm. See also Supplementary Video 1. **e** $Ca^{2+}$ imaging using

*35Spro:GCaMP3* (Col-0). A single trichome from a 2-week-old seedling was flicked with a

silver chloride wire. MS-induced intercellular calcium waves propagated concentrically from

the trichome (dashed outline). Scale bar, 0.2 mm. See also Supplementary Video 2. **f** $[Ca^{2+}]_{cyt}$

changes at sites indicated by numbers in (**e**). **g** Side view of a trichome whose neck was

flicked with a silver chloride wire. MS-induced intercellular $Ca^{2+}$ influx was transiently

observed in the trichome base (arrowheads) followed by the formation of circular waves.

Scale bar, 0.1 mm. See also Supplementary Video 3.

**Fig. 5 | Trichomes are mechanosensory cells. a** $Ca^{2+}$ imaging using *35Spro:GCaMP3* (Col-0) and *35Spro:GCaMP3* (*gl1*). Leaf surfaces were exposed to MS by brushing. MS-induced calcium waves were compromised in the *gl1* mutant. See also Supplementary Videos 4 and 5. Scale bars, 0.5 mm. **b** $[Ca^{2+}]_{cyt}$ signature of (**a**). **c** Venn diagram of transcriptome datasets

for MS-induced genes in Col-0 and *gl1* ($P < 0.05$). NS, not significant. Lower, fold change

(FC) (*gl1*)/FC (Col-0) < 0.5. High, MS (*gl1*)/Mock (*gl1*), $\log_2$FC $\geq 1$ in *gl1*. $P < 0.05$. **d**

Heatmap of differentially expressed defence-related genes obtained from transcriptome

datasets from Col-0 and *gl1* plants treated with MS (4 brushing). **e** Transcript levels of

*WRKY33*, *WRKY40*, and *CBP60g* in 4-week-old Col-0 and *gl1* plants 15 min after treatment

with 4 falling droplets, determined using RT-qPCR and normalized to *UBQ5*. Data are

presented as mean ± SD. Asterisks indicate significant difference (one- and two-way

ANOVA; *$P < 0.05$). **f** MS-induced MAPK activation in Col-0 and *gl1*. Total proteins were

extracted from 4-week-old leaves 5 min after MS treatment and detected by immunoblot

analysis with anti-p44/42 MAPK antibodies. Relative phosphorylation levels are shown

below each blot. **g** Growth of *Psm* ES4326 in Col-0 and *gl1* leaves 2 days after inoculation

with (+) or without (–) MS (4 brushing) pretreatment. Error bars represent SE. Asterisks

indicate significant difference (one- and two-way ANOVA; ***$P < 0.001$). Cfu, colony-

forming units. NS, not significant. **h** Disease progression of *Alternaria brassicicola* in Col-0

and *gl1* leaves 3 days after inoculation with (+) or without (–) MS (4 brushing) pretreatment.

Error bars represent SE. Asterisks indicate significant difference (one- and two-way ANOVA;

****$P < 0.0001$). NS, not significant.

**Fig. 6 | Trichomes activate broad-spectrum disease resistance.** Model showing how trichomes directly sense the mechanical impact of raindrops as an emergency signal in anticipation of possible infections. Mechanosensory trichome cells initiate intercellular calcium waves in response to MS. $[Ca^{2+}]_{cyt}$ initiates the de-repression of $Ca^{2+}$/CaM-dependent CAMTA3 and activates the phosphorylation of MPK3 and MPK6, thereby inducing WRKY-dependent transcription.