Eye-hand coordination during online reach corrections is task-dependent

Anouk J. de Brouwer[1] & Miriam Spering[1,2,3]

[1] Department of Ophthalmology & Visual Sciences, University of British Columbia, Vancouver, BC, Canada
[2] Institute for Computing, Information and Cognitive Systems, University of British Columbia, Vancouver, BC, Canada
[3] Djavad Mowafaghian Centre for Brain Health, University of British Columbia, Vancouver, BC, Canada

Running title: Eye-hand coordination during online reach corrections

Corresponding author: Miriam Spering, miriam.spering@ubc.ca

Abstract

To maintain accurate movements, the motor system needs to deal with errors that can occur due to inherent noise, changes in the body, or disturbances in the environment. Here, we investigated the temporal coordination of rapid corrections of the eye and hand in response to a change in visual target location during the movement. In addition to a 'classic' double-step task in which the target stepped to a new position, participants performed a set of modified double-step tasks in which the change in movement goal was indicated by the appearance of an additional target, or by a spatial or symbolic cue. We found that both the absolute correction latencies of the eye and hand and the relative eye-hand correction latencies were dependent on the visual characteristics of the target change, with increasingly longer latencies in tasks that required more visual and cognitive processing. Typically, the hand started correcting slightly earlier than the eye, especially when the target change was indicated by a symbolic cue, and in conditions where visual feedback of the hand position was provided during the reach. Our results indicate that the oculomotor and limb-motor system can be differentially influenced by processing requirements of the task and emphasize that temporal eye-hand coordination is flexible rather than rigid.

Key words: online control; visuomotor; perturbation; reaction time; reaching; eye movements

1   Introduction

2   How humans adjust and optimize movements to correct for errors that are due to sensory and 3   motor noise, changes in the body, or external disturbances is a major focus of current 4   neuroscience research. In the laboratory, tasks that artificially produce movement errors have 5   revealed the ability to rapidly correct for errors during the movement (i.e., movement corrections), 6   as well as the ability to adjust the movement to consistent errors over the course of several 7   repetitions (i.e., motor adaptation). Here, we investigate the temporal coordination of movement 8   corrections in eye and hand movements in a reaching task.

9   The double-step task has been widely used to study both movement corrections and motor 10   adaptation in eye and hand movements. In this classic paradigm, the visual target is displaced at 11   the time of movement onset to simulate a spatial error. In the case of saccadic eye movements, 12   the endpoint error after the saccade toward the initial target location triggers a second, corrective 13   saccade to the new target location (Becker and Jürgens 1979; Hallett and Lightstone 1976; Joiner 14   et al. 2010; Tian et al. 2013). If the target is repeatedly displaced to the same location, motor 15   adaptation will result in an adjustment of the initial saccade (Herman et al. 2013; McLaughlin 16   1967; Tian et al. 2009). The double-step paradigm has also been used extensively to study 17   corrections of reach movements, as well as to study adaptation of reach movements (Magescas 18   and Prablanc 2006). Because the duration of reach movements is much longer than that of 19   saccades, movement corrections in response to a displacement of the reach target typically occur 20   online (i.e., 'in flight') (Georgopoulos et al. 1981; Goodale et al. 1986; Megaw 1974; Soechting 21   and Lacquaniti 1983) with reaction times as short as ~110 ms (Brenner and Smeets 1997; Day 22   and Lyon 2000; Prablanc and Martin 1992). Another popular paradigm to study reach corrections 23   is the cursor displacement paradigm, in which hand movements correct within short latencies to 24   a perceived deviation of the reach trajectory (Brenner and Smeets 2003; Franklin and Wolpert 25   2008; Sarlegna et al. 2003; Saunders and Knill 2003). Together, these findings indicate that visual 26   information is continuously used to control and correct movements.

27   The significant progress in understanding how the sensorimotor system corrects for errors 28   in eye and hand movements is largely based on studies that have investigated these two motor 29   systems separately. The few studies that measured both eye and hand movements in response 30   to a displacement of a visual target have made several interesting observations. First, hand 31   movement corrections start earlier when a corrective saccade accompanies the reach, as 32   opposed to when the eyes are instructed to fixate (Abekawa et al. 2014; Diedrichsen et al. 2004),

showing a facilitative effect of eye movements on hand movements. Second, whereas the eye
usually leads the hand by (several) hundred milliseconds when initiating a goal-directed reach
(e.g., Land and Hayhoe 2001; Prablanc et al. 1979), the hand might start correcting for spatial
errors before the eye (Abekawa et al. 2014; Gritsenko et al. 2009), although the opposite finding
(i.e., the eye corrects before the hand) has also been reported (Neggers and Bekkering 2002).
Here, we investigated the detailed temporal coordination of eye and hand movement corrections
by assessing the timing of corrections in various stimulus and task conditions. For hand
movements, it has been shown that the visual characteristics of the target (Kozak et al. 2019;
Veerman et al. 2008), as well as the presence of visual distractors (Reichenbach et al. 2014) can
influence the duration within which corrections are initiated, but eye movements were either not
measured (Veerman et al. 2008) or participants were instructed to fixate (Reichenbach et al.
2014) in these studies.

On one hand, we might expect a tight temporal coupling of corrective eye and hand
movements, independent of stimulus and task conditions, based on the close behavioral and
neurophysiological connection of both movements (de Brouwer et al. 2021; Land and Hayhoe
2001; Prablanc et al. 1979). On the other hand, the temporal coupling might differ across
conditions. For example, recent work has shown that errors in the reach trajectory (i.e., due to a
cursor jump) evoke the earliest and most vigorous corrections when gaze is directed at the reach
target, as compared to surrounding locations (de Brouwer et al. 2018). As such, looking at the
target as early as possible might be particularly important when peripheral visual feedback of the
hand can be used to improve reach accuracy (i.e., when the hand is visible). In other conditions,
it might be optimal to maintain fixation at the original target to prevent a temporary disturbance of
the retinal image by eye movements (Mrotek and Soechting 2007; Neggers and Bekkering 2000;
Ross et al. 2001), and perform a reach correction in peripheral vision.

Participants performed a set of tasks in which they were asked to reach, as rapidly and
accurately as possible, toward a visual target using a robotic manipulandum, while their hand and
eye movements were recorded. In a subset of trials, an unpredictable change in the movement
goal occurred. We included a classic double-step task (Hallett and Lightstone 1976), in which the
target 'stepped' to a new location, triggering an immediate, reactive saccade to that location. In
addition, we designed a set of modified double-step tasks with the intention to manipulate the
latency of the saccade to the new movement goal. In these tasks, the change in movement goal
was indicated by the appearance of an additional target, or by a spatial or symbolic cue, placing
different demands on visual and cognitive processing. Whereas the visual target displacement
could be expected to trigger an immediate, reactive saccade, the additional target and cue

conditions were designed to prevent triggering an immediate saccade and instead produce a later,
voluntary saccade. Thus, we hypothesized that our modified double-step tasks would delay the
corrective saccade to the new goal location, but the question remained whether this would also
delay the reach correction (Abekawa et al. 2014; Diedrichsen et al. 2004), or whether the reach
correction would be prioritized to be completed as rapidly as possible. In addition to varying the
visual characteristics of the target change, we manipulated the presence of visual feedback of the
hand. We hypothesized that the presence of hand feedback would speed up the corrective
saccade to allow optimal monitoring of the reach trajectory (de Brouwer et al. 2018), and
potentially speed up the reach correction (Reichenbach et al. 2009). Overall, we aim to contribute
to a better understanding of the temporal coordination of eye and hand movements during online
reach corrections, the role of visual and cognitive demands in different task contexts, and the role
of visual feedback.

Methods

Participants

Nineteen participants (mean age 26 years, range 19-36 years, 5 female) completed the
experiment and were compensated $12/hour for their participation. Three other participants did
not complete the experiment because the experimenter could not achieve sufficiently accurate
calibration of the eye tracker (mean error <1.5° and maximum error <3.0° visual angle). All
participants were self-reported right-handed and had normal or corrected-to-normal visual acuity.
The study was approved by the University of British Columbia Behavioural Research Ethics
Board. Participants provided written informed consent before the start of the experiment.
Of the participants who completed the experiment, three participants were excluded from
the analysis because they had less than 4 (out of 10) valid trials for any combination of task ×
target change time × target change direction in two or more tasks. The most common reason for
an insufficient number of valid trials was that an initial saccade to the target could not be detected,
likely due to eye tracking difficulties. This exclusion resulted in 16 complete and analyzed data
sets for the experiment.

Experimental setup

Participants performed reaching movements to visual targets with their right hand. They were seated in a chair with their forehead resting against a pad and their hand holding on to the handle of a robotic manipulandum that moves in the horizontal plane (KINARM End-Point Robot; BKIN Technologies Ltd., Kingston, ON, Canada). A mirror was mounted horizontally above the handle, and an LCD monitor was mounted horizontally above the mirror, such that stimuli were projected in the plane of the handle when looking onto the mirror while the view of the arm was blocked (Fig. 1A). Kinematics of the handle were recorded with a temporal resolution of 1129 Hz, resampled to 1000 Hz. Eye movements of the left eye were recorded using a built-in monocular video-based eye tracker (Eyelink 1000, SR Research Ltd., Kanata, ON, Canada) with a temporal resolution of 500 Hz, resampled to 1000 Hz. The pupil was detected with a proprietary algorithm that accounted for small head movements, which were measured using a target sticker placed on the participant's forehead or cheek. The eye tracker was calibrated for the 2D horizontal workspace using proprietary algorithms (BKIN Technologies).

Task and visual stimuli

Figure 1B provides a schematic illustration of the tasks. The hand position was represented on the screen as a cursor (1 cm diameter white circle) aligned with the handle. All other stimuli were presented in red (5.9 cd/m^2) on a black background (0.9 cd/m^2) for clear, high-contrast visibility. Each trial began with the presentation of a start position (2 cm diameter circle) at the horizontal center of the screen near the participant. Participants were instructed to move the cursor and their gaze to the start position to initiate the trial. The cursor had to be at the start position for 250 ms, and the recorded eye position on the screen had to be within a 5 cm radius of the start position at the end of the 250-ms period. After a random delay of 250 to 500 ms, the start position disappeared and the reach target (2 cm diameter circle) appeared 20 cm in front of the start position and in line with the start position or 5 cm to the left or right of the start position. Participants were instructed to reach toward the target as quickly and accurately as possible. In 40% of trials, the movement goal remained unchanged during the trial. In the remaining 60% of trials, a sudden change in movement goal from the central to either the left or right target location was presented. Participants were instructed to move their hand to the new target as soon as possible. The target disappeared 250 ms after detection of the reach offset, defined by the velocity of the handle falling below a threshold of 2 cm/s for 250 ms. To encourage participants to perform fast reaching

movements throughout the experiment, a message 'Too slow' was displayed on the screen if the movement time was longer than 800 ms in unperturbed trials, or longer than 1100 ms in trials with a change in movement goal. A new trial started after a 500 ms inter-trial-interval.

Participants performed five versions of the main task, in separate blocks (Fig. 1B). (1) In the classic double-step task, a single reach target stepped from the central to the left or right target position. (2) In the placeholder task, placeholders (2 cm diameter open circles) were presented in addition to the target at the two non-target locations. When the target stepped, it would 'swap' position with one of the placeholders. (3) The overlap task was similar to the placeholder task, except that when the target stepped, the central circle would remain filled (i.e., the old target would stay in its position when the new target was presented). (4,5) In the line and arrow cue tasks, three filled targets were presented at the three target locations. (4) In the line (i.e., spatial) cue task, a reach to the left or right target, or a change in movement goal from the central to the left or right target, was indicated by the appearance of a vertical line (10 cm × 0.2 cm) presented 10 cm to the left or right of the midline, and vertically centered between the start and target position. (5) In the arrow (i.e., symbolic) cue task, a reach or change to the left or right target was indicated by the central circle changing into a leftward or rightward pointing triangle (2.5 × 1.5 cm).

To ensure that the time of the change was unpredictable and that our results were not determined by the time of the target change, the target jump or appearance of the cue was triggered by one of three events: the onset of the saccade (y gaze position reached a third of the distance between the start and target position), the onset of the reach movement (the cursor had fully moved out of the start position), or the time where the handle passed the midpoint between the start position and the target position. The execution of the saccade and the onset of the reach movement are commonly used target change triggers in target displacement tasks (e.g., Diedrichsen et al. 2004; Gritsenko et al. 2009; Prablanc and Martin 1992; Oostwoud Wijdenes et al. 2013). We included the late target change time to make sure that the timing of the corrective saccade would not be influenced by the saccade refractory period, which is commonly observed to be around 150 ms (Carpenter 1977).

Figure 1. Experimental setup and tasks. A) Participants performed reaching movements in the horizontal plane using a KINARM robotic manipulandum. Vision of the hand was blocked by a mirror onto which the stimuli were projected such that they appeared in the plane of the handle. The mirror is depicted as transparent to illustrate the location of the hand and the manipulandum. B) In each task, participants moved their hand from a start position to one of three target positions 20 cm in front of the start position (both shown as red circles). In a subset of trials, the movement goal changed from the central position to a position 5 cm to the left or right of the central position (shown by the grey arrow pointing to the left that was not visible to the participant). In the double-step task (1), the target stepped from the central to the left/right position (the dotted circle indicates the original, invisible target position). The placeholder task (2) was identical to the double-step task, except that the two alternative target positions were indicated with open circles together with the target position. The overlap task (3) was identical to the placeholder task, except that the original target position remained 'filled in' when the target jumped to its new position. In the line cue task (4), all three possible target locations were indicated, and the (new) target was indicated by a spatial cue: a line on the left for the left target, a line on the right for the right target, no line for the central target). In the arrow cue task (5), the (new) target was indicated by a symbolic cue location at the central target: a leftward pointing arrow for the left target, a rightward pointing arrow for the right target, a circle for the central target.

Procedure

Each participant performed one practice block and ten experimental blocks, with short breaks in between blocks. Half of the experimental blocks were performed with online visual feedback of the hand position during the reach, and the other half was performed with endpoint feedback only. In the latter case, the cursor disappeared when the reach target was presented and reappeared after movement offset. The order of tasks and feedback was counterbalanced. Twenty different orders for the five tasks were created by starting with a 5×5 Latin Square in which every task occurs once in each row and once in each column, and then duplicating this square three times while randomly permuting the columns. Half of the participants performed every task first with online cursor feedback and then with endpoint cursor feedback, while the other half performed

every task first with endpoint cursor feedback and then with online cursor feedback. The eye
tracker was typically recalibrated at the start of each block, however, in some cases where
calibration proved difficult, the eye tracker was recalibrated every two blocks. No instructions on
eye movements were given except for directing the eyes to the start position at the beginning of
the trial. The practice block consisted of 20 trials without a target change. Each experimental
block consisted of 20 repetitions of unperturbed trials to the central target, 10 repetitions of
unperturbed trials to the left/right target, 10 repetitions of trials in which the movement goal
changed to the left/right during the saccade to the target, 10 repetitions of trials in which the
movement goal changed to the left/right target following reach onset, and 10 repetitions of trials
in which the movement goal changed to the left/right following the handle passing the point
midway between the start and target position, resulting in a total of 100 trials per block.
Participants took about 90 minutes to complete the experiment.

194 Data analysis

195 The timing of the appearance of all visual stimuli and events (start position, target, target
196 displacement, cue) was corrected offline by the delay in the system (57±5 ms), which was
197 measured using a photodiode after the completion of data collection. Trials were excluded if the
198 offline analysis showed that the target change had occurred before the event that triggered it
199 (<0.5% of trials).

200 Hand and eye movement data preprocessing

The x and y positions of the center of the handle in the horizontal plane were used for the analysis
of hand movements. For each trial, the onset of the reach was defined as the first moment in time
where the resultant velocity of the handle was greater than 5 cm/s. In the offline analysis, the
reach offset was defined as the first moment in time where the resultant velocity of the handle
was smaller than 5 cm/s for 250 ms after the handle had passed the point midway between the
start and target position. Trials were discarded if the onset of the reach occurred before the target
appeared (1% of trials), or more than 1000 ms after the target appeared (<0.5%). Trials were also
discarded if the y amplitude of the reach was less than half the target distance, or the x amplitude
was less than half of the target displacement in change trials (1%).
The x and y eye positions on the screen, as obtained from the calibrated eye tracker, were
used for analysis of the eye movements. Eye blinks, intervals in which the pupil signal was
missing, and intervals in which the eye was detected outside the screen were removed from the

x and *y* eye positions. For intervals with missing data for less than 100 ms (i.e., shorter than a
typical blink duration), *x* and *y* eye position were linearly interpolated. Eye positions were filtered
with a 2nd order lowpass Butterworth filter with a cut-off frequency of 30 Hz. *X* and *y* eye velocity
$(\dot{x}, \dot{y})$ were computed by numerically differentiating *x* and *y* eye position signals.

Next, 2D eye positions on the screen were converted to an eye-based 3D coordinate
system (Singh, Perry & Herter, 2016), assuming that the height of the stimulus plane relative to
the eye is fixed (i.e., *z* is constant). The 3D eye positions on the screen were then transformed to
an eye-based spherical coordinate system according to

$$\rho = \sqrt{x^2 + y^2 + z^2}$$

$$\theta = tan^{-1}\left(\frac{y}{x}\right)$$

$$\phi = cos^{-1}\left(\frac{z}{\rho}\right)$$

224   with ρ the radial distance from the eye to the point of gaze on the screen, θ the azimuthal angle
225   in the *xy*-plane, and ϕ the elevation angle. Next, ρ, θ, and ϕ were differentiated to obtain $\dot{\rho}$, $\dot{\theta}$ and
226   $\dot{\phi}$:

$$\dot{\rho} = \frac{x \cdot \dot{x} + y \cdot \dot{y} + z \cdot \dot{z}}{\rho}$$

$$\dot{\theta} = \frac{\dot{x} \cdot y - x \cdot \dot{y}}{x^2 + y^2}$$

$$\dot{\phi} = \frac{z(x \cdot \dot{x} + y \cdot \dot{y}) - (x^2 + y^2)\dot{z}}{(x^2 + y^2 + z^2)\sqrt{x^2 + y^2}}$$

230   Note that $\dot{z}$ is assumed to be zero, reducing some parts of the equations to zero. Finally, the eye
231   angular velocity ω was calculated according to

$$\omega = \sqrt{(\dot{\theta} \cdot sin(\phi))^2 + \dot{\phi}^2}$$

Saccades were detected by searching for intervals in which angular eye velocity was larger than
100°/s for at least 4 ms. Next, saccade onset was defined as the last sample in which eye velocity
was below a threshold of 30°/s for 2 ms. Saccade offset was defined as the first sample in which
eye velocity fell below a threshold of 30°/s for 2 ms.

Target saccades were defined as saccades that started between 50 ms before (predictive
saccades) and 1000 ms after target appearance, with a *y* onset position before the midpoint
between start and target position, and a minimum *y* amplitude of half the target distance. Trials
were discarded if a target saccade was not detected (4% of trials). Corrective saccades from the
initial to the final target were defined as saccades that started at least 50 ms after the target

change, with a *y* onset and offset position past the midpoint between start and target position,
and a minimum *x* amplitude of half the target displacement. Change trials were discarded if a
corrective saccade was not detected (1% of trials). 93% of trials were included in the analysis.

Hand and eye movement analysis

We computed the latencies of the initial saccade and reach movements to the target and, for
target change trials, the latencies of the corrective responses of the eye and hand. The latency of
the corrective saccade was defined as the latency of the first saccade directed towards the new
target position following the target change. The latency of the corrective hand movement was
defined using a variant of the extrapolation method applied to the velocity of the hand in the *x*
direction (i.e., the direction of the perturbation) (Oostwoud Wijdenes et al. 2014). We first
corrected the *x* velocity in individual change trials by subtracting the average *x* velocity in no-
change trials to the central target in the same block. Next, for each trial we obtained the two points
at which the additional *x* velocity reached 30% and 70% of the first peak in velocity, and fitted a
line through these points and all data points in between. The onset of the reach correction was
defined as the time where this line crossed zero. In some trials, the increase in *x* velocity was not
approximately linear in the interval between the 30% and 70% peak velocity points, or there was
no clear, single peak in *x* velocity, resulting in a poor fit. We therefore discarded trials in which the
R^2 of the fit was below 0.95 or in which the extrapolated onset occurred before the target change
(2% of trials).

Hypotheses and statistical analysis

We designed a set of double-step tasks to determine the effect of target and cue characteristics
on the temporal coordination of corrections of the eye and hand. We hypothesized that, with
respect to the classic double-step task, corrective saccade latency would increase in the
placeholder task, because of the necessity to identify which of the items is the displaced target
(Oostwoud Wijdenes et al. 2014; Smeets et al. 2016), as well as in the overlap task, as latency
increases are a well-known effect of stimulus overlap (Saslow 1967). We predicted a greater
increase in corrective saccade latency in the line and arrow cue tasks, potentially even until after
the cursor hit the target, due to the lack of visual stimuli triggering a reactive saccade to the new
target. We also hypothesized that corrective saccade latencies would be shorter when online
cursor feedback was provided compared to when endpoint feedback was provided, to allow the
earliest possible use of visual feedback to help guide the incoming cursor (de Brouwer et al.,

2018). If the reach and saccadic system show strong temporal coupling, we further expect a
constant delay between reach and saccade correction latencies across tasks, with high
correlations within individuals. Weak coupling would result in low correlations within individuals,
and potentially in varying delays between corrective responses across tasks.

For each of the latencies, we computed the median value per subject, task and feedback
condition for no change trials to the central target (20 repetitions), and per subject, task, feedback
condition, target change time, and target change direction for change trials (10 repetitions per trial
type). These median values were averaged across left and right target change directions,
resulting in four values for each combination of task and feedback. To rule out any delays in the
corrective saccade due to a saccade refractory period when the target changes during or
immediately after the saccade, we assessed the effect of change time on the latency of
corrections in the classic double-step task. On average, the target change was visible on the
screen 86 ms after saccade onset, 54 ms after reach onset, and 57 ms after the hand passed the
midway point, or 242, 343, and 506 ms after target appearance, respectively. Although there was
a significant effect of change time on corrective saccade latency, with the shortest latencies when
the change occurred after reach onset, there was no significant difference between latencies
when the change occurred after saccade onset and when the change occurred midway during
the reach (i.e., with a long delay after the initial saccade). This suggests that factors other than
the saccade refractory period determined the corrective saccade latency. We therefore averaged
the data across change times, calculating the mean and within-subject standard error of the mean
(Cousineau 2005) for each combination of task and feedback.

Statistical analyses were performed in R using the rstatix package (v 0.7.0). We performed
5 (task) $\times$ 2 (feedback) repeated measures ANOVAs, applying a Greenhouse-Geisser correction
when the assumption of sphericity was violated, and we calculated generalized eta squared effect
sizes (η_G^2). Significant main effects were followed up by pairwise comparisons with Bonferroni
correction for multiple comparisons.

Results

300 The main goal of this study was to determine the influence of the characteristics of the visual
301 target or cue that indicates a sudden change in movement goal, on the timing of rapid corrective
302 responses of the eye and hand. An additional goal was to determine the effect of feedback of the
303 hand on the timing of these corrections. In the following sections, we will first show task and

feedback effects on correction latencies of the eye and hand. We then report how the temporal

relation between eye and hand corrections was affected by task and feedback.

Initial movement latencies

Regardless of task and feedback condition, each double-step trial started with a saccade toward

the initial (central) target that was followed by an initial reach toward the central target. To provide

insight into the timeline of events in our task, we report the mean latencies of these initial

movements, calculated per task and feedback condition, in Table S1. Briefly, the mean latencies

ranged between 143 and 162 ms for initial saccades, and between 197 and 260 ms for initial

reach movements. Since all correction latencies in our main analyses were assessed relative to

the target change, which was triggered by the execution of the saccade or reach, we did not test

the effects of task or feedback on these initial latencies. The short initial latencies suggest that

participants were anticipating the appearance of the target. The Pearson correlation coefficients

between saccade and reach latencies, calculated separately for each participant and then

averaged across participants, ranged between 0.39 and 0.59 (Table S1), indicating moderate

temporal coupling during the initial movement phase.

Effects of task and feedback on movement corrections

Latencies of corrective saccades and reach corrections

In response to the change in movement goal, both the eye and the hand rapidly initiated a

correction in all conditions. Examples of these corrections and their timing are shown in Figure 2

for a representative participant, in blocks with online cursor feedback. Figure 2A shows that the

initial straight-ahead reach trajectory was followed by a correction to the left or right, with later

target changes (brighter colours) eliciting later trajectory adjustments requiring a greater change

in reach direction. The eye and hand started correcting at around the same time (Fig. 2B), and

the corrective saccade was of much shorter duration than the reach correction.

Averaging across participants revealed a clear task-dependency of correction latencies

(Fig. 3A, B). Figure 3A shows that corrective saccade latencies (relative to the time of target

change) were modulated by the characteristics of the visual target or cue that indicated the target

change. Latencies were shortest in the classic double-step task (*M*=191 ms), longer in the

placeholder task (*M*=244 ms), longer again in the overlap and line cue tasks (*M*=271 and 271

ms), and longest in the arrow cue task (*M*=305 ms). This observation was reflected in a significant

main effect of task in a 5 (task) × 2 (feedback) ANOVA ($F(4,60)=67.5$, $p<0.001$, $\eta_G^2=0.63$). Post-

hoc pairwise comparisons confirmed significant differences between all tasks ($p<0.01$), except

between the overlap task and the line cue task ($p=1.000$; all p-values for pairwise comparisons

here and below were adjusted with a Bonferroni correction). Corrective saccade latencies did not

differ between blocks with online cursor feedback and blocks with endpoint cursor feedback (main

effect $F(1,15)=0.2$, $p=0.647$, $\eta_G^2=0.001$). There was no significant interaction between task and

feedback ($F(4,60)=1.5$, $p=0.220$, $\eta_G^2=0.007$).

Reach correction latencies were also dependent on the characteristics of the visual target

or cue that indicated the target change (Fig. 3B). Reach correction latencies were shortest in the

double-step task ($M=194$ ms), longer in the placeholder task ($M=238$ ms), and longest in the

overlap ($M=265$ ms), line cue ($M=266$ ms) and arrow cue ($M=280$ ms) tasks. This was reflected

in a significant main effect of task ($F(4,60)=57.7$, $p<0.001$, $\eta_G^2=0.52$), with significant pairwise

differences between all tasks (all $p<0.001$), except between the overlap, line cue, and arrow cue

tasks ($p>0.05$). In addition, reach correction latencies were shorter with online cursor feedback

($M=240$ ms) than with endpoint cursor feedback ($M=258$ ms; main effect of feedback $F(1,15)=9.2$,

$p=0.008$, $\eta_G^2=0.08$). The effect of feedback was significant for all tasks ($p<0.05$) except the arrow

cue task ($p=0.375$), as revealed by pairwise comparisons following up on the significant

interaction between task and feedback ($F(4,60)=3.2$, $p=0.018$, $\eta_G^2=0.01$).

In summary, corrective saccade latencies were longer in the modified double-step tasks

than in the classic double-step task, in line with our expectations. Reach correction latencies

followed a similar pattern to saccade latencies, with the exception that there was no further

increase in latency for the arrow cue task. We also found that while corrective saccade latencies

were not influenced by the presence of online cursor feedback, reach corrections started earlier

when online cursor feedback was provided. Knowing that both corrective saccade latencies and

reach correction latencies were affected by the task, the next question is how the relative timing

and the correlation between saccade and reach corrections were influenced by the task.

Figure 2. Eye and hand corrections in response to a change in movement goal in individual trials of an example participant, in tasks with online cursor feedback. A) Hand trajectories. Each trajectory is a single trial, with the brightness indicating the target change time as indicated in the legend in B. Note that the results were averaged across target change times (see Methods). B) Timing of corrections relative to the target change. Reach corrections are shown by the longer horizontal bars (one for each trial), with the brightness indicating the target change time. Corrective saccades are shown by the short dark-colored bars. For each target change time, trials are sorted according to the latency of the reach correction.

Figure 3. Timing of eye and hand corrections in response to a change in movement goal, as a function of task and cursor feedback. A) Latencies of corrective saccades relative to the target change. Each small dot represents a participant and each square represents the average across participants (*n*=16). B) As A, but showing reach correction latencies. C) Differences between corrective saccade latencies and reach correction latencies, with positive values indicating that the eye started correcting first, and negative values indicating that the hand started correcting first. D) Correlations between corrective saccade and reach correction latencies. For all panels, colors represent different tasks, filled symbols represent online cursor feedback and open symbols represent endpoint feedback only. Error bars represent ± one within-subjects standard error of the mean [note that these are often hidden behind the symbol showing the average value].

Temporal coordination of movement corrections

To determine the influence of task on the temporal coordination of corrections of the eye and hand, we calculated the relative latencies of corrective saccades and reach trajectory corrections, as well as the correlations between latencies. Figure 3C shows that the hand often started correcting before the eye. The relative latencies of corrections of the hand and eye showed a significant effect of task ($F(4,60)$=9.8, p<0.001, η_G^2=0.13) and feedback ($F(1,15)$=15.3, p=0.001, η_G^2=0.13), as well as a significant interaction ($F(4,60)$=3.4, p=0.014, η_G^2=0.01). First, the lead of the hand was greater in the arrow cue task (*M*=25 ms) than in the other four tasks (all *p*<0.001). The relative latencies in the double-step, placeholder, overlap, and line cue tasks (*M*=-3, *M*=7, *M*=6 and *M*=6 ms, respectively) did not differ significantly between each other (all *p*>0.05). Second, the lead of the hand was greater when the hand cursor was visible during the reach (*M*=18 ms) than when the cursor was only shown at the end of the reach (*M*=-1 ms). Pairwise comparisons to unpack the interaction showed that this difference was significant for all tasks (all *p*<0.01) except the arrow cue task (*p*=0.080).

As an additional measure of the temporal coordination of corrections, we calculated

Pearson correlation coefficients between corrective saccade and reach correction latencies (Fig.

3D). These correlations showed a significant effect of task ($F(4,60)=18.0$, $p<0.001$, $\eta_G^2=0.36$) but

not feedback ($F(1,15)=0.3$, $p=0.623$, $\eta_G^2=0.002$), and no significant interaction ($F(4,60)=1.1$,

$p=0.343$, $\eta_G^2=0.02$). Correlations were lower in the double-step task ($M=0.18$) than in the overlap

($M=0.52$, $p<0.001$), line cue ($M=0.57$, $p<0.001$), and arrow cue tasks ($M=0.44$, $p<0.001$).

Correlations were also lower in the placeholder task ($M=0.29$) than in the overlap ($p=0.003$), line

cue (p<0.001), and arrow cue tasks ($p=0.026$), and correlations were higher in the line cue task

than in the arrow cue task ($p=0.020$).

In summary, the hand generally started correcting its trajectory in response to a change

in movement goal before a corrective saccade occurred. Although in some cases the relative

delays between corrections of the hand and eye were around zero, the longer neuromuscular

delays and higher inertia of the arm (Gribble et al. 2002) imply that even in these cases, the neural

signal to execute a correction occurred earlier for the hand than for the eye. This is in contrast to

the temporal coordination of the initiation of goal-directed eye-hand movements, where the eye

typically leads the hand. Taken together with the results on absolute latencies above, the greater

hand-eye delay in the arrow cue task compared to the other tasks was the result of the corrective

saccade being delayed. Further, the greater lead of the hand when online cursor feedback was

provided resulted from an earlier response of the hand. Finally, correlations between correction

latencies differed between tasks, with lower correlations in the classic double-step task and

placeholder task than in the overlap, line cue, and arrow cue tasks.

428 Discussion

This study aimed at determining the effects of visual target and cue characteristics on temporal

eye-hand coordination when hand movements had to rapidly correct for a change in movement

goal location, while the eyes were free to move. To date, few studies have investigated online

corrections of goal-directed reach movements while also measuring eye movements. In addition,

the task demands in these studies were limited to simple target step paradigms. We found that

across stimulus and task conditions, participants initiated a corrective saccade in addition to

correcting the hand trajectory. We report the following key findings: First, the correction latency in

eye and hand depended on visual and cognitive processing demands of the task. For example,

task versions that contained a cue indicating the change in movement goal triggered longer-

latency corrections as compared to task versions in which participants simply followed the target

when it stepped to a new location. Second, eye and hand corrections were not initiated
simultaneously: the hand started correcting before the eye. The hand-eye correction delay was
greatest when the target change was signaled by a symbolic cue. Third, correction latencies of
the eye and hand were weakly correlated, within individuals and tasks, in target step tasks, and
moderately correlated in tasks where the target change was indicated by an additional target or
cue, requiring an intentional correction. Fourth, online visual feedback of the hand cursor,
compared to endpoint feedback only, sped up the initiation of online corrections of the hand
trajectory but did not affect the latency of corrective saccades. Together, our findings show that
the timing and coordination of rapid movement corrections of the eye and hand depend on the
visual and cognitive demands of the change in movement goal. These findings provide important
insights into the role of stimulus and task demands during rapid movement corrections, and will
be discussed in detail below.

Our first key finding is that the latencies of both corrective saccades and reach corrections
depend on the processing demands of the task. Previous studies already showed that visual
target characteristics can affect reach correction latencies in response to a target change (Kozak
et al. 2019; Veerman et al. 2008). Here, we show that corrective saccade latencies are also
affected by the visual as well as cognitive demands of the task. Specifically, consistent with our
hypothesis, corrective saccade latencies increased, relative to those in the classic double-step
task, in a placeholder task where all three possible targets were shown and the target swapped
position with one of the placeholders, and further increased in an overlap task where the new
target was shown in addition to the initial target. Corrective saccade latencies also increased
relative to the classic double-step task when the target change was indicated by a spatial (line)
cue, and further increased when the target was indicated by a symbolic (arrow) cue. The latencies
of corrections of the reach trajectory showed a similar pattern, but without a further increase in
latency in the arrow cue task. Note that in all of our tasks, reach correction latencies were slower
than commonly reported. This is most likely a result of the extra mass and inertia of the robotic
manipulandum, but could also have been caused by relatively low contrast of stimuli on the screen
(Veerman et al. 2008). Nevertheless, the task-dependency of latencies emphasizes the
importance of visual characteristics of the target and the requirements of the task for the timing
of movement corrections.

Our second key finding is that corrections in eye and hand are not initiated simultaneously,
and–as the absolute latency–the relative latency of corrections depends on the visual and
cognitive processing demands of the task. Whereas many previous studies reported that the eye
typically leads the hand when initiating a movement (Bowman et al. 2009; Land and Hayhoe 2001;

Neggers and Bekkering 2000; Prablanc et al. 1979), here we show that the opposite can be true
when performing a correction, as Abekawa and colleagues (2014) have previously shown for
corrective movements in response to a target displacement (see also Gritsenko et al. 2009).
Although the relative latencies between corrections of the hand and the eye were around zero in
most of the tasks in the current study, it is important to note that even in cases where corrections
of the eye and hand were detected at around the same time from their velocities, the longer
neuromuscular delay and greater inertia of the arm and hand imply that the neural signal to
execute a correction was sent earlier to the arm muscles than to the eye muscles. The finding
that the reach correction is initiated before the corrective saccade indicates that the initial
correction of the reach must be based on an approximate computation of the visual movement
goal location (Franklin et al. 2016), since high-resolution retinal information as well extraretinal
information of the new target position does not become available until the eye lands on the new
target. By contrast, the later part of the correction could be more refined, as shown for trajectory
corrections in response to cursor displacements (Cross et al. 2019). Further, the delay of the
corrective saccade was greater in the arrow cue task compared to the other tasks, as a result of
a further increase in corrective saccade but not reach correction latencies. In this particular task,
the eye appeared to linger, likely reflecting the time it took to process the shape information of the
visual cue, requiring the involvement of the ventral visual stream (Goodale and David Milner
1992). Moreover, in the arrow cue task, the cue was presented centrally, akin to tasks involving
endogenous visual spatial attention. Such tasks typically involve slower and more sustained shifts
of attention. By contrast, the line cue task involved an exogenous cue, which is known to trigger
faster, transient shifts of visual spatial attention (Busse et al. 2008; Carrasco 2011). The time
course of events in our tasks reflects the differences between endogenous and exogenous visual
spatial attention. However, it is somewhat surprising that the extra processing time in the symbolic
cue task affected the eye, but did not cause a further increase in reach correction latencies.

As another aspect of temporal eye-hand coordination, we showed that, within individuals
and tasks, correction latencies of the eye and hand are weakly to moderately correlated. Our third
key finding is that these correlations were higher for tasks that involved a slower, intentional or
voluntary correction (overlap and cue tasks) than in tasks that involved a faster, more automatic
correction (classic double-step and placeholder tasks). This is similar to the findings reported by
Sailer and colleagues (2000) for direct goal-directed actions. Generally, correlations in movement
latencies imply a link in the mechanisms driving these movements (de Brouwer et al. 2021).
Higher correlations in more demanding tasks might indicate an overlap in the latency distribution
of eye and hand movements for voluntary but not for reactive saccades, or might indicate the

presence of a 'bottleneck' in information processing that affects both the oculomotor and the limb-
motor system (though this would not explain the greater hand-eye delay in the arrow cue task).
Even though correction latencies were overall correlated, the initiation of eye and hand corrections
is likely driven by separate movement commands subserved by different neural circuitry (de
Brouwer et al. 2021).

Our fourth key finding is that online visual feedback on the hand (cursor) location, as
compared to endpoint feedback only, sped up the initiation of hand movement corrections
(Reichenbach et al. 2009), resulting in a greater lead of the hand. Corrective saccade latencies
did not differ between blocks with online and endpoint cursor feedback This was in contrast to our
hypothesis that online cursor feedback would (also) reduce the latency of corrective saccades,
given the fact that directing the eyes to the reach target improves monitoring of the reach trajectory
in peripheral vision (de Brouwer et al. 2018), providing an important reason to move the eyes to
the new target location as soon as possible. Presumably, cursor feedback increases certainty
about the location of the hand (Acerbi et al. 2017; Izawa and Shadmehr 2008), and therefore
accelerates the correction, but this facilitation does not transfer to the eye.

With respect to the neural mechanisms underlying movement corrections, previous
studies on online corrections of reaching movements have emphasized the role of the posterior
parietal cortex (Desmurget et al. 1999; Pisella et al. 2000), as well as the possibility of a
subcortical pathway guiding both the eye and the limb when performing corrections (Day and
Brown 2001; Reynolds and Day 2012) (see also Cross et al. 2019; Pruszynski et al. 2010). The
exact contributions of these areas are still unknown (for reviews see Archambault et al. 2015;
Gaveau et al. 2014). Reynolds and Day proposed that a subcortical circuit allows for very fast
responses, while a cortical pathway drives slightly slower but more flexible responses. Although
our results cannot provide conclusive evidence about the pathways involved in online corrections,
the short latencies in the classic double-step task and longer latencies in the other tasks are
consistent with the idea of a subcortical and cortical circuit. Further, the involvement of the ventral
visual stream can explain the additional increase in latencies in the arrow cue task.

In summary, we designed a set of double-step tasks and showed that the correction of the
reach trajectory in response to a target change is accompanied by a corrective saccade. Both the
absolute and relative latencies of the corrective saccade and reach correction depended on the
visual characteristics of the target change. Corrections were initiated later when the task required
more visual and cognitive processing, and the hand typically started correcting before the eye,
especially when the change was initiated by a symbolic cue. Our results highlight that the hand
and eye are coordinated in a flexible manner that is suited for the task at hand. Our findings also

provide a framework for discussing latency differences obtained across tasks and studies and emphasize the importance of taking stimulus and task conditions into account when assessing eye-hand coordination. This is especially important when aiming to generalize laboratory results to the more visually and cognitively demanding conditions of real-world tasks.

Acknowledgements

We would like to thank Hannah Brown for help with data collection. This work was funded by an NSERC Discovery Grant and Accelerator Supplement (RGPIN 418493) awarded to Miriam Spering.

References

Abekawa N, **Inui T**, **Gomi H**. Eye-hand coordination in on-line visuomotor adjustments. *Neuroreport* 25: 441–445, 2014.

Acerbi L, **Vijayakumar S**, **Wolpert DM**. Target Uncertainty Mediates Sensorimotor Error Correction. *PLoS One* 12: e0170466, 2017.

Archambault PS, **Ferrari-Toniolo S**, **Caminiti R**, **Battaglia-Mayer A**. Visually-guided correction of hand reaching movements: The neurophysiological bases in the cerebral cortex. *Vision Res* 110: 244–256, 2015.

Becker W, **Jürgens R**. An analysis of the saccadic system by means of double step stimuli. *Vision Res* 19: 967–983, 1979.

Bowman MC, **Johansson RS**, **Flanagan JR**. Eye-hand coordination in a sequential target contact task. *Exp Brain Res* 195: 273–283, 2009.

Brenner E, **Smeets JB**. Fast Responses of the Human Hand to Changes in Target Position. *J Mot Behav* 29: 297–310, 1997.

Brenner E, **Smeets JBJ**. Fast corrections of movements with a computer mouse. *Spat Vis* 16: 365–376, 2003.

de Brouwer AJ, **Flanagan JR**, **Spering M**. Functional Use of Eye Movements for an Acting System. *Trends Cogn Sci* 25: 252–263, 2021.

de Brouwer AJ, **Gallivan JP**, **Flanagan JR**. Visuomotor feedback gains are modulated by gaze position. *J Neurophysiol* 120: 2522–2531, 2018.

Busse L, **Katzner S**, **Treue S**. Temporal dynamics of neuronal modulation during exogenous and endogenous shifts of visual attention in macaque area MT. *Proc Natl Acad Sci U S A* 105: 16380–16385, 2008.

Carpenter RHS. *Movements of the Eyes*. 1977.

Carrasco M. Visual attention: the past 25 years. *Vision Res* 51: 1484–1525, 2011.

Cousineau D. Confidence intervals in within-subject designs: A simpler solution to Loftus and Masson's method. *Tutorials in Quantitative Methods for Psychology* 1: 42–45, 2005.

Cross KP, **Cluff T**, **Takei T**, **Scott SH**. Visual Feedback Processing of the Limb Involves Two Distinct Phases. *The Journal of Neuroscience* 39: 6751–6765, 2019.

Day BL, Brown P. Evidence for subcortical involvement in the visual control of human reaching. *Brain* 124: 1832–1840, 2001.

Day BL, **Lyon IN**. Voluntary modification of automatic arm movements evoked by motion of a visual target. *Experimental Brain Research* 130: 159–168, 2000.

Desmurget M, **Epstein CM**, **Turner RS**, **Prablanc C**, **Alexander GE**, **Grafton ST**. Role of the posterior parietal cortex in updating reaching movements to a visual target. *Nat Neurosci* 2: 563–567, 1999.

Diedrichsen J, **Nambisan R**, **Kennerley SW**, **Ivry RB**. Independent on-line control of the two hands during bimanual reaching. *Eur J Neurosci* 19: 1643–1652, 2004.

Franklin DW, **Reichenbach A**, **Franklin S**, **Diedrichsen J**. Temporal Evolution of Spatial Computations for Visuomotor Control. *J Neurosci* 36: 2329–2341, 2016.

Franklin DW, **Wolpert DM**. Specificity of reflex adaptation for task-relevant variability. *J Neurosci* 28: 14165–14175, 2008.

Gaveau V, **Pisella L**, **Priot A-E**, **Fukui T**, **Rossetti Y**, **Pélisson D**, **Prablanc C**. Automatic online control of motor adjustments in reaching and grasping. *Neuropsychologia* 55: 25–40, 2014.

Georgopoulos AP, **Kalaska JF**, **Massey JT**. Spatial trajectories and reaction times of aimed movements: effects of practice, uncertainty, and change in target location. *J Neurophysiol* 46: 725–743, 1981.

Goodale MA, **David Milner A**. Separate visual pathways for perception and action. *Trends in Neurosciences* 15: 20–25, 1992.

Goodale MA, **Pelisson D**, **Prablanc C**. Large adjustments in visually guided reaching do not depend on vision of the hand or perception of target displacement. *Nature* 320: 748–750, 1986.

Gribble PL, **Everling S**, **Ford K**, **Mattar A**. Hand-eye coordination for rapid pointing movements. Arm movement direction and distance are specified prior to saccade onset. *Exp Brain Res* 145: 372–382, 2002.

Gritsenko V, **Yakovenko S**, **Kalaska JF**. Integration of predictive feedforward and sensory feedback signals for online control of visually guided movement. *J Neurophysiol* 102: 914–930, 2009.

Hallett PE, **Lightstone AD**. Saccadic eye movements towards stimuli triggered by prior saccades. *Vision Res* 16: 99–106, 1976.

Herman JP, **Blangero A**, **Madelain L**, **Khan A**, **Harwood MR**. Saccade adaptation as a model
of flexible and general motor learning. *Exp Eye Res* 114: 6–15, 2013.

Izawa J, **Shadmehr R**. On-line processing of uncertain information in visuomotor control. *J*
Neurosci 28: 11360–11368, 2008.

Joiner WM, **Fitzgibbon EJ**, **Wurtz RH**. Amplitudes and directions of individual saccades can
be adjusted by corollary discharge. *J Vis* 10: 22.1–12, 2010.

Kozak RA, **Kreyenmeier P**, **Gu C**, **Johnston K**, **Corneil BD**. Stimulus-Locked Responses on
Human Upper Limb Muscles and Corrective Reaches Are Preferentially Evoked by Low Spatial
Frequencies. *eNeuro* 6, 2019.

Land MF, **Hayhoe M**. In what ways do eye movements contribute to everyday activities? *Vision*
Res 41: 3559–3565, 2001.

Magescas F, **Prablanc C**. Automatic drive of limb motor plasticity. *J Cogn Neurosci* 18: 75–83,
2006.

McLaughlin SC. Parametric adjustment in saccadic eye movements. *Perception &*
Psychophysics 2: 359–362, 1967.

Megaw ED. Possible modification to a rapid on-going programmed manual response. *Brain Res*
71: 425–441, 1974.

Mrotek LA, **Soechting JF**. Target interception: hand-eye coordination and strategies. *J*
Neurosci 27: 7297–7309, 2007.

Neggers SF, **Bekkering H**. Ocular gaze is anchored to the target of an ongoing pointing
movement. *J Neurophysiol* 83: 639–651, 2000.

Neggers SFW, **Bekkering H**. Coordinated control of eye and hand movements in dynamic
reaching. *Hum Mov Sci* 21: 349–376, 2002.

Oostwoud Wijdenes L, **Brenner E**, **Smeets JBJ**. Analysis of methods to determine the latency
of online movement adjustments. *Behav Res Methods* 46: 131–139, 2014.

Pisella L, **Gréa H**, **Tilikete C**, **Vighetto A**, **Desmurget M**, **Rode G**, **Boisson D**, **Rossetti Y**. An
"automatic pilot" for the hand in human posterior parietal cortex: toward reinterpreting optic
ataxia. *Nat Neurosci* 3: 729–736, 2000.

Prablanc C, **Echallier JF**, **Komilis E**, **Jeannerod M**. Optimal response of eye and hand motor
systems in pointing at a visual target. I. Spatio-temporal characteristics of eye and hand
movements and their relationships when varying the amount of visual information. *Biol Cybern*
35: 113–124, 1979.

Prablanc C, **Martin O**. Automatic control during hand reaching at undetected two-dimensional
target displacements. *J Neurophysiol* 67: 455–469, 1992.

Pruszynski JA, **Andrew Pruszynski J**, **King GL**, **Boisse L**, **Scott SH**, **Randall Flanagan J**,
Munoz DP. Stimulus-locked responses on human arm muscles reveal a rapid neural pathway
linking visual input to arm motor output. *European Journal of Neuroscience* 32: 1049–1057,
2010.

Oostwoud Wijdenes L, **Brenner E**, **Smeets JBJ**. Comparing Online Adjustments to Distance
and Direction in Fast Pointing Movements. *Journal of Motor Behavior* 45: 395–404, 2013.

Reichenbach A, **Franklin DW**, **Zatka-Haas P**, **Diedrichsen J**. A dedicated binding mechanism
for the visual control of movement. *Curr Biol* 24: 780–785, 2014.

Reichenbach A, **Thielscher A**, **Peer A**, **Bülthoff HH**, **Bresciani J-P**. Seeing the hand while
reaching speeds up on-line responses to a sudden change in target position. *J Physiol* 587:
4605–4616, 2009.

Reynolds RF, **Day BL**. Direct visuomotor mapping for fast visually-evoked arm movements.
Neuropsychologia 50: 3169–3173, 2012.

Ross J, **Morrone MC**, **Goldberg ME**, **Burr DC**. Changes in visual perception at the time of
saccades. *Trends Neurosci* 24: 113–121, 2001.

Sailer U, **Eggert T**, **Ditterich J**, **Straube A**. Spatial and temporal aspects of eye-hand
coordination across different tasks. *Exp Brain Res* 134: 163–173, 2000.

Sarlegna F, **Blouin J**, **Bresciani J-P**, **Bourdin C**, **Vercher J-L**, **Gauthier GM**. Target and hand
position information in the online control of goal-directed arm movements. *Exp Brain Res* 151:
524–535, 2003.

Saslow MG. Effects of Components of Displacement-Step Stimuli Upon Latency for Saccadic
Eye Movement. *Journal of the Optical Society of America* 57: 1024, 1967.

Saunders JA, **Knill DC**. Humans use continuous visual feedback from the hand to control fast
reaching movements. *Exp Brain Res* 152: 341–352, 2003.

Singh T, **Perry CM**, **Herter TM**. A geometric method for computing ocular kinematics and
classifying gaze events using monocular remote eye tracking in a robotic environment. *J
Neuroeng Rehabil* 13: 10, 2016.

Smeets JB, **Oostwoud Wijdenes L**, **Brenner E**. Movement Adjustments Have Short Latencies
Because There is No Need to Detect Anything. *Motor Control* 20: 137–148, 2016.

Soechting JF, **Lacquaniti F**. Modification of trajectory of a pointing movement in response to a
change in target location. *J Neurophysiol* 49: 548–564, 1983.

Tian J, **Ethier V**, **Shadmehr R**, **Fujita M**, **Zee DS**. Some perspectives on saccade adaptation.
Ann N Y Acad Sci 1164: 166–172, 2009.

Tian J, **Ying HS**, **Zee DS**. Revisiting corrective saccades: Role of visual feedback. *Vision
Research* 89: 54–64, 2013.

Veerman MM, **Brenner E**, **Smeets JBJ**. The latency for correcting a movement depends on the
visual attribute that defines the target. *Exp Brain Res* 187: 219–228, 2008.