Negative modulation of macroautophagy by HERPUD1 is counteracted by an increased ER-lysosomal network with impact in drug-induced stress cell survival

Vargas G.[1❖], Cortés O.[1❖], Arias-Muñoz E.[1,2], Hernández S.[1], Cerda-Troncoso C.[1], Hernández L.[1], González A. E.[5]&, Tatham M. H.[3], Bustamante H.[4], Retamal C.[1], Cancino J.[1], Varas-Godoy M.[1], Hay R. T.[3], Rojas-Fernandez A.[3,6], Cavieres V. A.[1,2*] and Burgos P. V.[1,2*]

[1] Centro de Biología Celular y Biomedicina (CEBICEM), Facultad de Medicina y Ciencia, Universidad San Sebastián, Lota 2465, Santiago 7510157, Chile.

[2] Centro de Envejecimiento y Regeneración (CARE-UC), Facultad de Ciencias Biológicas, Pontificia Universidad Católica, Santiago 8330023, Chile

[3] Center for Gene Regulation and Expression, College of Life Sciences, University of Dundee, DD1 4HN, Dundee 4HN UK

[4] Instituto de Microbiología Clínica, Facultad de Medicina, Universidad Austral de Chile, Valdivia 5110566, Chile.

[5] Instituto de Fisiología, Facultad de Medicina, Universidad Austral de Chile, Valdivia 5110566, Chile.

[6] Instituto de Medicina & Centro Interdisciplinario de Estudios del Sistema Nervioso (CISNe), Universidad Austral de Chile, Valdivia 5110566, Chile.

* Correspondence: vcavieresr@docente.uss.cl (V.A.C.); patricia.burgos@uss.cl (P.V.B.); Tel.: +56-9-65887097 (V.A.C.); +56-2-22606309 (P.V.B.)

❖ These authors contributed equally to this work

& Current address: Institute of Biochemistry II, School of Medicine, Goethe University Frankfurt, Theoder-Stern-Kai 7, 60590 Frankfurt am Main, Germany.

ABSTRACT

Macroautophagy and the ubiquitin proteasome system work as an interconnected network in the maintenance of cellular homeostasis. Indeed, efficient activation of macroautophagy upon nutritional deprivation is sustained by degradation of preexisting proteins by the proteasome. However, the specific substrates that are degraded by the proteasome in order to activate macroautophagy are currently unknown. By quantitative proteomic analysis we identified several proteins downregulated in response to starvation but independently of ATG5 expression. Among them, the most significant was HERPUD1, an ER protein of short-half life and a well-known substrate of the proteasome. We found that increased HERPUD1 stability by deletion of its ubiquitin-like domain (UBL) plays a negative role on basal and induced macroautophagy. Moreover, we found it triggers ER expansion by reordering the ER in crystalloid structures, but in the absence of unfolded protein response activation. Surprisingly, we found ER expansion led to an increase in the number and function of lysosomes establishing a tight network with the presence of membrane-contact sites.

44 Importantly, a phosphomimetic S59D mutation within the UBL mimics UBL deletion on its
45 stability and the ER-lysosomal network expansion revealing an increase of cell survival
46 under stress conditions. Altogether, we propose stabilized HERPUD1 downregulates
47 macroautophagy favoring instead a closed interplay between the ER and lysosomes with
48 consequences in drug-cell stress survival.
49
50 **INTRODUCTION**
51
Macroautophagy (from here referred to as autophagy) is a catabolic pathway that mediates
the engulfment of aberrant or damaged cytoplasmic constituents into double-membrane
autophagosomes that subsequently fuse with lysosomes to form a hybrid organelle called
the autolysosome that mediates the degradation of the cargo by acid hydrolases (Mizushima
et al. 2008; Khaminets, Behl, and Dikic 2016). Autophagy is also implicated in the
degradation of cellular constituents under basal conditions, playing an essential role in the
maintenance of cellular homeostasis upon a variety of environmental conditions such as
nutrient restriction or other stressors (Murrow and Debnath 2013). Autophagy is highly
inducible by environmental changes being a highly dynamic process that resolves a variety
of cellular demands (Murrow and Debnath 2013). In fact, increased autophagy is protective
in different cells and organisms, playing a crucial role in cell maintenance and survival under
different insults (Moreau, Luo, and Rubinsztein 2010). On the other hand, defects in
autophagy enhance cell vulnerability under harmful conditions such as those present in the
tumor microenvironment (Camuzard et al. 2020).
66
Although initially autophagy was thought to work independently of the ubiquitin proteasome
system (UPS), increasing evidence shows many layers of both negative and positive
regulation (Bustamante et al. 2018), revealing an interconnected network with important
roles in cellular homeostasis and maintenance (Korolchuk, Menzies, and Rubinsztein 2010).
Inhibitors of the proteasome 20S catalytic core with the use of β-subunits blockers triggers
an enhancement in the biogenesis of autophagosomes (Zhu, Dunner, and McConkey 2010).
In contrast, impairment of the proteasome 19S regulatory particle with an inhibitor of
PSMD14, a proteasomal deubiquitinating enzyme, blocks the biogenesis of
autophagosomes (Demishtein et al. 2017; Bustamante et al. 2020). To date a limited number
of substrates of the UPS system are known to play a regulatory role in autophagy (Jia and
Bonifacino 2019; Thayer et al. 2020) and many aspects about the functional role of this
interconnected network between autophagy and UPS remain elusive.
79
To identify UPS substrates that could act as negative regulators of autophagy, we conducted
a quantitative SILAC proteomic analysis in cells stably depleted of ATG5 by shRNA-
mediated knockdown. ATG5 protein is part of a complex with ATG12 and ATG16L that
controls an essential step in the autophagosome formation (Walczak and Martens 2013). We
focused on proteins downregulated in response to nutrient deprivation, but not because of
autophagy activation. The protein with the most significant downregulation, in wild type and
ATG5 depleted cells was the Homocysteine-responsive endoplasmic reticulum-resident
ubiquitin-like domain (UBL) member 1 protein named as HERPUD1. This protein is a
transmembrane ER-resident protein with low levels of expression due to its short half-life by
rapid proteasomal degradation (K. Kokame et al. 2000; Sai et al. 2003).
90

Here, we found that increased HERPUD1 stability through the deletion of its UBL domain
causes a decrease in basal and induced autophagy. Additionally, it promotes an ER
expansion which is organized in stacked tubules forming crystalloid ER-like structures but
independent of ER stress. Furthermore, we uncovered that higher HERPUD1 stability has a
positive impact in lysosomal function, establishing an ER-lysosomal network with the
presence of membrane contact-sites (MCS). Further, combining bioinformatics and site-
directed mutagenesis we found the phosphomimetic S59D mutant within the UBL domain of
HERPUD1 mimics the effect of the UBL deletion increasing its levels and inducing the
expansion of the ER-lysosomal network, a phenotype that promotes stress cell survival.
These findings thus identify HERPUD1 as a hotspot platform to promote stress cell survival
by inducing an expansion of the ER-lysosomal network when autophagy slows down.
102
103 **MATERIALS AND METHODS**
104
Reagents
Bafilomycin A1 (BafA1, cat#B1793), tunicamycin (Tun, cat#T7765), thapsigargin (Tg,
cat#T9033), cisplatin (CDDP, cat#479306), Sulforhodamine B (SRB, cat#230162), Earle's
balanced salt solution (EBSS, Cat#E2888), puromycin dihydro-chloride (cat#P8833), and
protease inhibitors cocktail (cat#P8340) were purchased from Sigma-Aldrich (St. Louis, MO,
USA). MG132 (cat#474790) was purchased from Merck Millipore (Burlington, MA, USA).
LysoTracker^{TM} Red DND-99 (cat#L7528), 4′,6-diamidino-2-phenylindole (DAPI) (cat#D-1306)
and TRIzol^{TM} (cat#15596018) were purchased from ThermoFisher Scientific (Waltham, MA,
USA), Magic Red® (cat#6133) was purchased from Immunochemistry Technologies, LLC
(Bloomington, IN, USA). The QuikChange II XL direct-mutagenesis kit was obtained from
Stratagene (cat#200522, La Jolla, CA, USA) and the Vybrant Apoptosis Pacific Blue-annexin
V kit and 7AAD from Invitrogen (cat#A35122).
Antibodies
118
The following monoclonal antibodies were used: mouse anti-β-ACTIN clone BA3R (cat#
MA5-15739, Thermo Fisher Scientific), mouse anti-XBP-1S clone E7M5C (cat#47134S, Cell
Signaling Technology, Danvers, MA, USA), mouse anti-FLAG clone M2 (cat#F1804, Sigma
Aldrich), mouse anti-GRP78/BiP clone 40/BiP (cat# 610978, BD Biosciences, San Jose, CA,
USA), mouse anti-LAMP1 clone H4A3 (cat# 610978, Developmental Studies Hybridoma
Bank, Iowa City, IA, USA), rabbit monoclonal anti-CALNEXIN clone C5C9 (cat#2679S, Cell
Signaling Technologies), rabbit monoclonal anti-ATF4 clone D4B8 (cat#11815S, Cell
Signaling Technologies), rabbit monoclonal anti-HERPUD1 clone EPR9649 (cat#ab150424,
Abcam), rat monoclonal anti-GRP94 clone SPM249 (cat#ab233979, Abcam, Cambridge,
UK). We used the following polyclonal antibodies: rabbit anti-LC3 (cat#2775S, Cell Signaling
Technology), rabbit anti-PERK (cat#P0074, Sigma-Aldrich), goat anti-CATHEPSIN-D
(cat#AF1014, R&D Systems, Minneapolis, MN, USA), rabbit anti-HERPUD1 (cat#BML-
PW9705, ENZO Life Sciences, Farmingdale, NY, USA). Horseradish peroxidase-conjugated
secondary antibodies were purchased from Jackson ImmunoResearch Laboratories (West
Grove, PA, USA), Alexa fluorophore-conjugated secondary antibodies were purchased from
Thermo Fisher Scientific.
135
Mass Spectrometry based proteomics and statistical analysis

Human H4 neuroglioma cells stably expressing shLuc and shATG5 were grown in
Dulbecco's modified Eagle's medium lacking all amino acids except L-lysine and L-arginine,
which were replaced with either unlabelled amino-acids (Lys0 and Arg0) or stable isotopic
forms 13C6 15N2-lysine (Lys8) and 13C6 15N4-arginine (Arg10) (Cambridge Isotope
Laboratories). The medium was supplemented with 10% dialyzed FBS using previous
published methods (Golebiowski et al. 2009; Yin et al. 2012). Two separate cultures of each
shLuc and shATG5 cells were grown in light amino acids (Lys0 + Arg0) or heavy amino
acids (Lys8 + Arg10) over 9 replication cycles to achieve over 96% incorporation and similar
cell counts per culture. Both cell cultures were divided in two, and cells were grown under
control conditions in DMEM + 10% FBS or starvation in EBSS media for 4 h so that each cell
type grown with each label were treated with both conditions (8 cultures in total). Cells were
washed and lysed in 1.2 x LDS sample buffer with reducing agent, obtaining crude cell
extracts at approximately 1mg/ml. To allow all experimental conditions to be compared with
one another a light reference mix was obtained by combining all four light amino acid
conditions in a 1:1:1:1 ratio by volume. This was then mixed 1:1 ratio (v:v) with each heavy
amino acid extract. The same comparisons were made in reverse by combining all heavy
amino acid samples into a reference mix and combining this 1:1 (v:v) with each of the four
individual light amino acid conditions. See Figure 1A for experimental design. All 8 mixes
were fractionated by SDS-PAGE and stained gels were cut into 3 slices per lane before
tryptic peptides were extracted. The resultant 24 samples of dried down peptides were
resuspended in 35 μl 0.1% TFA 0.5% acetic acid. Peptide samples were analyzed by LC-
MS/MS twice; the first using 9 μl peptide sample run over a 90 min peptide fractionation
gradient, and the second using 18 μl peptide sample run over a 240 min fractionation
gradient. Peptides were analyzed using a Q exactive mass spectrometer (Thermo Scientific)
coupled to an EASY-nLC 1000 liquid chromatography system (Thermo Scientific), using an
EASY-Spray ion source (Thermo Scientific), running a 75 μm x 500 mm EASY-Spray column
at 45°C. A top 10 data-dependent method was applied. Full scan spectra (m/z 300-1800)
were acquired with resolution R= 70,000 at m/z 200 (after accumulation to a target value of
1,000,000 with maximum injection time of 20ms). The most intense ions were fragmented by
HCD and measured with a resolution of R= 17,500 at m/z 200 (target value of 500,000
maximum injection time of 60 ms) and intensity threshold of 2.1×10^4. Peptide match was set
to "preferred", a 40 second dynamic exclusion list was applied, and ions were ignored if they
had an assigned charge state of 1, 8 or >8. All 48 data files were analyzed simultaneously in
MaxQuant (v1.5.2.8) using default parameters excepting the selection of SILAC labels,
activation of 'requantify' and 'match between runs', using a uniport 'HUMAN' proteome
database (downloaded 24/02/2015- 73920 entries) as search space. Raw files derived from
the same mix were grouped under the 'Experiment' heading as Mix01-Mix08. Raw files were
given MaxQuant experimental design 'fraction' numbers such that spectra derived from the
same HPLC gradient and from the equivalent gel slices across different lanes would be
matched, as well as one sliced either side. 4395 protein groups were listed as informed in
the Supp. File 1, proteinGroups.txt sheet. This list was edited to leave 3911 protein groups
by removing decoy proteins, protein listed as potential contaminants, proteins identified only
by modified peptides, and proteins without a H/L ratio reported for any comparison (see
Supp. File 1, accepted sheet). The average of normalized forward and reverse ratios of
starvation/control were carried forward for statistical analysis, but only for those with H/L
ratios reported for both. These values along with average $\log_{10}$ protein intensities were used
in Perseus to calculate Significance B values to identify statistical outliers (SigB<0.001) for
the three ratios. Data for these shortlisted proteins are summarized in Supp. File 1,

Plasmids and site-directed mutagenesis

For all HERPUD1 constructs generated in this study, previously described cDNAs encoding
either the full-length or the ΔUBL deletion mutant human HERPUD1, both with a C-terminal
FLAG-tag, were used as templates (Sai et al. 2003). pCI-HERPUD1-FLAG and pCI-ΔUBL-
FLAG were digested with EcoRI and NotI restriction enzymes to obtain an insert, which was
sub-cloned to a lentiviral pLVX-IRES-Puro vector (Takara Bio Inc, CA, USA) containing a
puromycin resistance gene. The substitution S59D and S59A were introduced into the pLVX-
IRES-FLAG-tagged HERPUD1 vector using the QuikChange II XL direct-mutagenesis kit
(Stratagene, cat#200522) and the mutagenesis service of GenScript (Hong Kong, China).

Cell culture and generation of stable cell lines
198
Maintenance of H4 human neuroglioma cells stably expressing either shRNAs against ATG5
or luciferase genes was performed as previously described (González et al. 2017). HeLa
cells were obtained from the American Type Culture Collection (Manassas, VA, USA). HeLa-
derived cell lines were cultured in Dulbecco's modified Eagle's medium (DMEM; Thermo
Fisher Scientific) supplemented with 10% (vol/vol) heat-inactivated fetal bovine serum (FBS;
Thermo Fisher Scientific), and 100 U/ml penicillin/100 mg/ml streptomycin (Thermo Fisher
Scientific), in a 5% CO_2 atmosphere at 37ºC. The generation of HeLa stable cell lines
expressing all different variants of FLAG-tagged HERPUD1 cloned in the pLVX-IRES-Puro
vector were generated by transfection with Lipofectamine 2000 (Invitrogen) according to
manufacturer's instructions. After 24 h the cells were selected and maintained with 2 µg/ml
of puromycin.
210

Preparation of protein extracts, Electrophoresis, SDS-PAGE and Western blot analysis
213
Cells were washed with ice-cold phosphate buffered saline (PBS) and lysed in
Radioimmunoprecipitation assay buffer (RIPA lysis buffer) [50 mM Tris-HCl , 150 mM NaCl,
5 mM EDTA, 1% NP-40, 1% sodium deoxycholate, 0.1% SDS, pH 7.4], supplemented with a
cocktail of protease inhibitors [416 µM 4-(2-Aminoethyl)benzenesulfonyl fluoride, 0.32 µM
Aprotinin, 16 µM Bestatin, 5.6 µM E-64, 8 µM Leupeptin and 6 µM Pepstatin A; Sigma-
Aldrich] and phosphatase inhibitors (1 mM NaF, 0,3 mM $Na_2P_2O_7$ and 1 mM Na_3VO_4; Sigma-
Aldrich). Cell lysates were collected and lysed for 30 min at 4°C in rotation. Then, extracts
were sonicated with ultrasonic power three times on ice with pulses of 2-3 sec at 40 mA
using a tip sonicator system. Extracts were further centrifuged for 20 min at 13.000xg at 4°C.
The supernatants were collected, and protein concentration was quantified using the BCA
assay (ThermoFisher Scientific). The protein extracts were denatured at 65°C for 5 min and
analyzed using our previous described methods (González et al. 2017; Bustamante et al.
2020).
227

Transmission electron microscopy
229

HeLa cells were fixed for 16 h by immersion in 2.5% glutaraldehyde in 0.1 M cacodylate
buffer (pH 7.0) at room temperature, and then washed three times with a cacodylate buffer
for 2 h. Cells were post-fixed with 1% osmium tetroxide (OsO_4) for 2 h and washed 3 times
with double distilled water. Then, the cells were treated with 1% aqueous uranyl for 90 min,
and sequentially dehydrated through an acetone battery 50%, 70%, 95%, and 100% for 20
min each. Cells were pre-embedded in epon/acetone 1:1 overnight and then in pure epon for
4 h. Finally, cells were embedded in fresh resin and polymerized in an oven at 60°C for 48 h.
Ultrafine sections (80 nm) were obtained using an ultramicrotome Leica Ultracut R. The
sections were incubated with 4% uranyl acetate in methanol for 2 min and lead citrate for 5
min. The grids were visualized using a Philips Tecnai 12 electron microscope (Eindhoven,
The Netherlands) at 80 kV. This work was performed in the Advanced Microscopy Facility of
the Faculty of Biological Sciences at Pontificia Universidad Católica de Chile.

Fluorescence microscopy, data acquisition and quantification

Cells grown on glass coverslips were washed with PBS-Ca^{2+}/Mg^{2+} and then fixed,
permeabilized and stained using our published protocols (Bustamante et al. 2020; Cavieres
et al. 2020). The images were acquired using a TCS SP8 laser-scanning confocal
microscope (Leica Microsystems, Wetzlar, Germany) equipped with a 63× oil immersion
objective (1.4 NA), photomultipliers (PMT), hybrid detectors (HyD) system using 405 nm,
488 nm, 561 nm and 643 nm laser lines for excitation running the LASX Leica software. For
image quantification, 16-bit (1024x1024) images were acquired under identical settings
avoiding signal saturation. Cell and nucleus area measurements were performed by using
ICY   software   (Quantitative   Image   Analysis   Unit,   Institut   Pasteur,
http://icy.bioimageanalysis.org/). A pipeline was created to completely automate image
analysis by using the following sequential plugins: active contours (cell segmentation), hk-
means (threshold detection), wavelet spot detector (spot detection) and studio
(colocalization). Total Fluorescence Integrated intensity measurement was performed by
using ImageJ (FIJI) (Schindelin et al. 2012). The number of dots (lysosomes) was performed
by using the LOG detector algorithm available on the TrackMate plugin (FIJI). For live cell
imaging assays, HeLa cells were grown in glass bottom culture dishes (MatTek Corporation,
Ashland, MA, USA) and labeled with the following Invitrogen probes: ER-Tracker™ Blue-
White DPX (E12353), LysoTracker™ Red DND-99 (L7528, Invitrogen) and Magic-red
(Immunochemistry Technologies, LLC) according to the manufacturer's protocol. Before
image acquisition, culture medium was replaced with phenol red-free DMEM supplemented
with HEPES (10 mM, pH 7.4), and images were acquired with the 63x oil immersion
objective (1.4 NA) of the TCS SP8 laser-scanning confocal microscope, running the Leica
Application Suite LAS X software, coupled to a controlled temperature chamber (UNO-temp
controller, OKOLAB) acquiring 16-bit images at 37ºC (488 laser for excitation; HyD: 510–550
nm; 1024 × 1024 pixels; frame average 2). For volume analysis, z-stack (0.3 m z-interval,
1024x1024, 180 m pixel size) images were quantified using 3D analysis plugin by using
ICY software. Total Fluorescence Integrated intensity and dots number were measured
using ImageJ.
271
272 **RNA isolation and RT-PCR analysis**
273
274 Total RNA extraction from HeLa cells, oligo-dT and MMLV reverse transcriptase and
275 quantitative reverse transcription PCR of the cDNA template (RT-PCR) was carried out as

previously described (Bustamante et al. 2020). The specific primer pairs used for CYCLOPHILIN-A, CYCA (NM_001300981.2) and XBP1 (NM_001079539.2) were the following: **CYCA-F** TCGAGTTGTCCACAGTCAGC, **CYCA-R** TTCATCTGCACTGCCAAGAC, **XBP1-F** CGCTTGGGGATGGATGCCCTG and **XBP1-R** CCTGCACCTGCTGCGGACT.

Cell viability assays

Exponentially growing cells stably expressing either FLAG-tagged HERPUD1-WT or HERPUD1-S59D were trypsinized and seeded at 20,000 cells per well in 96-well microplates and allowed to attach for 6 h at 37°C and 5% CO_2. Then, cells were incubated with CDDP in serial dilutions from 1 mM to 1 µM in 2% FBS medium and were incubated for 24 h at 37°C and 5% CO_2. After drug incubation, the IC50 was obtained using the Sulforhodamine B (SRB) cell cytotoxicity assay (Blois, Smith, and Josephson 2011). Briefly, cells were fixed (10% trichloroacetic acid, 4°C, 1 h), water washed, dried and stained (0.4% SRB v/v in 0.1% acetic acid, 1 h at RT), and then washed four times (1% acetic acid). Dissolved SRB (10 mM Tris-base, pH 10) was quantified (564 nm, Synergy HT BioTek reader). CDDP treatments were done in quadruplicate in at least three independent experiments. The IC50 values associated with the cytotoxic effects of CDDP were calculated using GraphPad Prism software (version 8.2; GraphPad Software, San Diego, CA, USA) using non-linear regression model and dose-response equations (log(inhibitor) vs. normalized response). After drug incubation, apoptotic cells were analyzed with the commercial kit Vybrant Apoptosis Pacific Blue-annexin V (cat#A35122, Invitrogen) using the protocol provided by the manufacturer. Briefly, cells stably expressing either FLAG-tagged HERPUD1-WT or HERPUD1-S59D were treated with 10 µM CDDP for 24 h. Cells were harvested and centrifuged at 800 x g for 5 min at room temperature and washed with cold PBS 1X. Cells were re-centrifuged, and the pellet resuspended in 100 µL annexin-binding buffer. Then, 5 µL of the annexin V conjugate was added to the cell suspension and incubated for 15 min at room temperature. After, 400 µL of annexin-binding buffer was added, gently mixed, and maintained on ice for later analysis in a BD FACSCanto II flow cytometer (Flow Cytometer Facility of Cell 4 Cell, Santiago, Chile), with a previous incubation with 3 µL of 7-Amino-Actinomycin D (7AAD) to exclude non-viable cells (included in the kit).

Densitometric Quantification and Statistical Analysis

The amount of immunoblot signal was estimated using Image J software version 1.48v (Wayne Rasband, NIH, http://imagej.nih.gov). For each condition, protein bands were quantified from at least three independent experiments in order to ensure adequate statistical power. Data analysis was performed using Microsoft Excel 2013 for Windows (Redmond, WA, USA) or GraphPad Prism 6. Results are represented in graphs depicting the mean ± standard deviation. Statistical significance of data comparisons from two groups comparisons was determined with two-tailed unpaired Student's T-test for parametric data. Values of $P < 0.05$ (*), $P < 0.01$ (**), $P < 0.001$ (***) were regarded as statistically significant and are indicated in the figures.

Acknowledgments: We thank the generous donation of pCI-HERPUD1-Flag and pCI-HERPUD1-ΔUBL-Flag by Doctor Koichi Kokame from National Cerebral and Cardiovascular

321 Center, Japan. Additionally, we thank Dr. Lavandero and Dr. Quiroga for the critical
322 discussion of the manuscript.
323

Funding: This research was funded
by Fondo Nacional de Desarrollo Científico y Tecnológico of Chile
(FONDECYT; http://www.conicyt.cl/fondecyt) No. 1190928 to MV-G, No. 1171649 to P.V.B.
& No. 11150532 to A.R.F.; Associative Investigation Program
(PIA; https://www.conicyt.cl/pia) including No. ACT-172066 to P.V.B. & No. AFB-170005 to
P.V.B.; Academy Insertion Program (PAI; http://www.conicyt.cl/pai) No. 79150075 to
A.R.F.; Fondo de Equipamiento Científico y Tecnológico of Chile
(FONDEQUIP; http://www.conicyt.cl/fondequip) No. EQM150118 to P.V.B.; Cooperation
International Programme (CONICYT-RCUK; https://www.conicyt.cl/pci) No. DPI20140068 to
P.V.B.

334
335 **Statement of interest:** The authors declare that there is no conflict of interest.
336
337 **Competing interest:** The authors declare that they have no competing interests.
338
339 **Data availability statement:** All the original data supporting the results is available.
340
341 **RESULTS**
342
343 ***HERPUD1 is a regulator of autophagy under the control of its UBL domain.***
344
Previous reports have demonstrated a close interplay between autophagy and the Ubiquitin-
Proteasome System (Bustamante et al. 2018), however, to date, few proteasomal substrates
are known as modulators of autophagy (Jia and Bonifacino 2019; Guarascio et al. 2020;
Thayer et al. 2020). To search for potential novel candidates that could be downregulated by
the proteasome in order to activate autophagy, we performed a SILAC-based proteomic
study to quantitatively determine the proteome of H4 neuroglioma cells under basal and
induced autophagy by EBSS starvation conditions. To eliminate all proteins downregulated
because of autophagy activation, we compared the proteome of H4 cells where autophagy is
inhibited by stable depletion of ATG5 by shRNA-mediated knockdown (shATG5) respect to
control H4 cells expressing an shRNA against the luciferase gene (shLuc), both cell lines
previously characterized (González et al. 2017; Tapia et al. 2019) (Fig. 1A). ATG5 protein is
part of a complex with ATG12 and ATG16L that controls an essential step in the
autophagosome formation (Walczak and Martens 2013). Silencing of ATG5 causes a strong
inhibition in LC3B-positive autophagosomes, a phenotype also previously confirmed in our
H4 cell lines (González et al. 2017). Among all the proteins downregulated by EBSS
starvation, we found that in both H4 cell lines, shLuc (Fig. 1B) and shATG5 (Fig. 1C) the
most significantly downregulated protein was HERPUD1, a protein originally identified as a
homocysteine-inducible gene, that is also upregulated by endoplasmic reticulum (ER) stress
(K. Kokame et al. 2000; Koichi Kokame, Kato, and Miyata 2001). Importantly, HERPUD1 is
an ER-stress membrane protein whose levels under non-stressful conditions are low due to
proteasome degradation (K. Kokame et al. 2000; Sai et al. 2003). Indeed, pharmacological
inhibition of the proteasome leads to a rapid increase of HERPUD1 levels (Sai et al. 2003;
Miura et al. 2010). In addition to HERPUD1, we found several other proteins significantly
down- or up-regulated by EBSS treatment (Fig. 1D). However, while some proteins were
down- or up-regulated by EBSS treatment in both, shLuc and shATG5 stable expressing cell
lines (Fig. 1D, red dots), many other hits were only down or up-regulated dependent on

ATG5 protein expression (Fig. 1D, purple dots). The complete list of proteins that responded
significantly to EBSS treatment in both cell lines is shown in Suppl. Fig 1. Here, we focus on
the characterization of HERPUD1, the hit with the highest score of downregulation in both
cell lines, working with the hypothesis that its reduction by EBSS treatment could be
indicative of its role as negative regulator of autophagy activity. Previous studies had shown
that silencing HERPUD1 triggers the autophagic flux (Quiroga et al. 2013). Therefore, in this
study, and considering that HERPUD1 levels are increased under ER stress, we study the
cellular consequences of HERPUD1 stability on autophagy function. HERPUD1 is an
integral membrane protein with both termini facing the cytoplasm, with the ubiquitin-like
domain (UBL) located in its N-terminus (Fig. 2A; UBL: purple color, left side). Here, we
cloned full-length HERPUD1 into the pLVX-IRES-Puro vector with a FLAG-tagged in its C-
terminal end (Fig. 2A; FLAG: green color) designed as HERPUD1-WT. In addition, we
cloned a FLAG-tagged HERPUD1 version lacking the residues between Val^{10}-Cys^{86}
designed as HERPUD1-ΔUBL (Fig. 2A; right side), as previously reported (Sai et al. 2003).
Further, we generated puromycin-resistant HeLa cells stably expressing either HERPUD1-
WT or HERPUD1-ΔUBL, considering that previous characterization of HERPUD1 in HeLa
cells was done only by transient transfection (Sai et al. 2003). Western blot analysis using
anti-HERPUD1, and anti-FLAG antibodies showed that the levels of HERPUD1-ΔUBL were
higher than the levels induced by the overexpression of HERPUD1-WT (Fig. 2B, lane 3 and
lane 5). In addition, western blot analysis with an anti-HERPUD1 antibody confirmed
HERPUD1 is expressed in low levels under non-stressful conditions (Fig. 2B, lane 1). The
UBL is a domain that resembles ubiquitin in terms of their primary sequence and three-
dimensional structure, considered a general interaction motif with the proteasome
particularly with the 19S regulatory particle of the 26S proteasome (R. Hartmann-Petersen
and Gordon 2004). Interestingly, it has been previously shown that deletion of UBL in
HERPUD1 abolishes its proteasomal degradation (Sai et al. 2003). To confirm this, we
treated the cells during 4 h with 20 µM MG132, a potent blocker of the proteasome activity
(Tsubuki et al. 1996; D. H. Lee 1998). As expected, we observed that this treatment caused
an increase in the levels of overexpressed HERPUD1-WT (Fig. 2B, lane 3 and 4). Similarly,
we observed a slight increase in the levels of endogenous HERPUD1 (Fig. 2B, lane 1 and
2). In contrast, HERPUD1-ΔUBL did not respond to this treatment, observing similar levels in
the absence or presence of MG132 (Fig. 2B, lane 5 and 6). Together, these results
confirmed HERPUD1-ΔUBL is a tool to explore the effect of HERPUD1 stability on
autophagy.
405
Therefore, we investigated the effect of HERPUD1-ΔUBL stable expression by analyzing the
subcellular distribution of the microtubule-associated protein 1 light chain 3B (LC3B), a
classical marker of autophagy, from here referred for simplicity as LC3 (Tanida, Ueno, and
Kominami 2008). Immunofluorescence analysis of LC3 in cells expressing HERPUD1-WT
showed basal autophagy represented by LC3-positive membrane dots that correspond to
autophagosomes decorated with lipidated LC3 (LC3-II) (Fig. 3A; left, control). In comparison,
we noticed that HERPUD1-ΔUBL cells showed a slight decrease in the number of
autophagosomes which was accompanied with a higher LC3 cytosolic distribution (LC3-I)
(Fig. 3A; right, control). Because autophagosomes are constantly forming autolysosomes
through the fusion with acidic lysosomes for degradation, we tested the effect of BafA1, a
drug that raises the lysosomal pH resulting in the perturbation of the autophagic flux
(González et al. 2017). As expected, treatment of HERPUD1-WT cells with BafA1 showed a
higher number of autophagosomes (Fig. 3A, left, BafA1). In contrast, this treatment in

HERPUD1-ΔUBL cells showed fewer numbers of autophagosomes compared to HERPUD1-
WT cells (Fig. 3A, right, BafA1). Furthermore, we biochemically validated these results
performing western blot analysis of endogenous LC3 (Fig. 3B). We found that under basal
conditions overexpression of HERPUD1-ΔUBL caused an increase in LC3-I levels, in
comparison to HERPUD1-WT (Fig. 3B, lane 1 and 2). In agreement with this observation,
HERPUD1-ΔUBL cells showed a significant reduction in the LC3-II/LC3-I ratio (0.47 ± 0.22),
respect to control cells (1.00 ± 0.05) (Fig. 3C). A similar finding was observed in the
presence of BafA1 treatment, observing in HERPUD1-ΔUBL cells a significant decrease in
the ratio LC3-II/LC3-I (0.94 ± 0.37), respect control cells (3.01 ± 0.78) (Fig. 3B, lane 3 and
4). Importantly, similar findings were obtained under induced autophagy by EBSS starvation
medium (Suppl. Fig. 2). We observed that the LC3-II/LC3-I ratio in HERPUD1-ΔUBL cells in
the presence of EBSS plus BafA1 was diminished with respect HERPUD1-WT cells (Suppl.
Fig. 2A, line 5 and 6 and Suppl. Fig. 2B). Altogether, these findings strongly indicate that
increased stability of HERPUD1 plays a negative effect on autophagy.
433
434   ***Increased stability of HERPUD1 by the deletion of its UBL domain triggers expansion***
435   ***of the ER organized in stacked tubules and crystalloid ER-like structures in the***
436   ***absence of ER stress.***
437
Because previous reports have shown that defective autophagy leads to ER expansion (Ou
et al. 1995; Jung et al. 2008) we investigated whether HERPUD1 stability could be linked
with ER remodeling. In contrast to previous studies where only transient expression of
HERPUD1-ΔUBL was characterized (Sai et al. 2003), we investigated the subcellular
distribution of FLAG-tagged HERPUD1-ΔUBL but in stable HeLa cell lines, in comparison
with its wild type version. Immunofluorescence analysis with an anti-FLAG antibody showed
a stronger signal for HERPUD1 in the absence of its UBL domain (Fig. 4A, FLAG, lower left
panel) in comparison to HERPUD1-WT (Fig. 4A, upper left panel). A similar result was also
observed with an anti-HERPUD1 antibody (Suppl. Fig. 3A). To unveil if this phenotype was
affecting the ER in general, we analyzed by immunofluorescence the distribution of
endogenous ER proteins markers. First, we tested endogenous CALNEXIN, an ER
membrane resident protein that acts as a molecular chaperone of glycoproteins (Ou et al.
1995) and GRP94, an ER resident membrane protein of the Heat-Shock Protein (HSP) 90
family (Marzec, Eletto, and Argon 2012). We found profound changes in the distribution of
both proteins showing a similar pattern to anti-FLAG in HERPUD1-ΔUBL cells (Fig. 4A,
CALNEXIN and GRP94, lower panel), compared to HERPUD1-WT where a characteristic
ER pattern is observed. Indeed, HERPUD1-ΔUBL shows a nice colocalization with
CALNEXIN and GRP94 (Fig. 4A, lower merge). In addition, and to exclude the possibility
that this phenotype was specific to CALNEXIN and GRP94, we analyzed the ER pattern
using an ER-tracker probe in live cells, observing clear changes in the morphology of the ER
respect to HERPUD1-WT (Suppl. Fig. 3B). We noticed that in comparison to HERPUD1-
ΔUBL, HERPUD1-WT did not show a high colocalization with CALNEXIN and GRP94 (Fig.
4A, upper merge), which can be explained by the low levels of HERPUD1-WT expression
due to its constant degradation by the proteasome under basal conditions (Sai et al. 2003).
In this regard, localization of HERPUD1 at the ER has been well documented (K. Kokame et
al. 2000). Whereas HERPUD1-ΔUBL did not cause an increase in the levels of CALNEXIN
and GRP94 measured by Western blot analysis (Suppl. Fig. 3C y 3D), we concluded that
HERPUD1 stability by the deletion of its UBL domain causes an ER-like expansion
phenomena that does not involve the increase in the levels of CALNEXIN and GRP94.

HERPUD1 is known to be upregulated under ER stress, a condition reported to cause ER
membrane expansion to alleviate this condition (Schuck et al. 2009). Therefore, we studied
whether HERPUD1-ΔUBL could be causing ER proliferation. To assess this, we performed
immunofluorescence analysis with an anti-GRP94 antibody. Interestingly, we observed that
HERPUD1-ΔUBL expressing cells present a large and dense ER network extending
throughout the entire cytoplasm including the periphery of the cell, compared to the less
extended ER network observed in HERPUD1-WT cells (Fig. 4B, left and right panel). To gain
a more comprehensive understanding of the ER differences between HERPUD1-WT and
HERPUD1-ΔUBL expressing cells, we measured the ER volume using 3D images at high
magnification taken with a z-interval of 0.3 µm followed by a z-stack maximum intensity
projection. For visualization, we analyzed the distribution of GRP94 in the whole cell using a
heat map gradient ranging from non-intensity (black color) to the higher intensity (red color)
(Fig. 4B, left and right panel). This analysis confirmed HERPUD1-ΔUBL increases the
volume of the ER and the expansion of this organelle, compared to the expression of
HERPUD1-WT. Indeed, quantitative analysis showed a significant increase in the ER
volume with the expression of HERPUD1-ΔUBL (2361 ± 967), compared to HERPUD1-WT
(1700 ± 689) (Fig. 4C). The ER morphological changes observed by confocal microscopy
prompted us to perform ultra-structural analysis by transmission electron microscopy (TEM).
While cells expressing HERPUD1-WT showed the common ER cytoplasmic structures (Fig.
4D left panel [ER]), unexpectedly HERPUD1-ΔUBL showed a high number of staked tubular
ER structures oriented in a hexagonal spatial distribution (Fig. 4D center panel [ER]). Others
have referred to these structures as energetically stable structures formed by a remarkable
proliferation of smooth ER, resembling crystalloid structures of ER (Chin et al. 1982;
Anderson, Orci, and Brown 1983; Borgese, Francolini, and Snapp 2006), which in
appearance can be compared with "honeycomb like-structures" (Fig. 4D center panel [ER]).
Additionally, a higher magnification (Fig. 4D, right panel), confirmed the ER contained in the
crystalloid structure corresponds to smooth ER evidenced by the absence of attached
ribosome structures.
495
To gain insights whether the ER proliferation triggered by the expression of HERPUD1-
ΔUBL was the result of an ER stress response, we studied X-box binding protein 1 (XBP1)
as a reporter of ER stress and the unfolded protein response (UPR). First, we analyzed by
RT-PCR the XBP1 mRNA processing, from the unspliced inactive XBP1 mRNA (u*XBP1*)
form to the spliced active XBP1 mRNA (s*XBP1*) form (Fig. 5A), which is considered a
hallmark of the UPR response (Yoshida et al. 2001). RT-PCR analysis of mRNA from HeLa
WT untreated cells showed a single band as expected (Fig. 5A, lane 1). The same analysis
including tunicamycin (Tun) (inhibitor of N-linked glycosylation (Heifetz, Keenan, and Elbein
1979) and thapsigargin (Tg) (blocker of ER Ca2+ import (Thastrup et al. 1990)) as positive
inducers of ER stress, produced the appearance of two bands in HeLa-WT cells, indicating
the s*XBP1* form in response to ER stress (Fig. 5A, lane 2 and 3). In contrast, a stable
expression of either HERPUD1-WT or HERPUD1-ΔUBL showed no detection of s*XBP1,*
observing only the expression of the u*XBP1* form (Fig. 5A, lane 4 and 5), similar to control
HeLa WT untreated cells (Fig. 5A, lane 1). Analysis of PCR bands relative to *CYCLOPHILIN*
A (CYC-A) as a housekeeping control showed no difference between cells expressing either
HERPUD1-WT or HERPUD1-ΔUBL proteins (Fig. 5A, lane 4 and 5). Similar findings were
observed by western blot analysis (Fig. 5B). s*XBP1* is rapidly translated to a highly active
transcription factor, known as XBP1, responsible for the upregulation of a variety of UPR
genes (Yoshida et al. 2001). To examine the significance of the UBL domain of HERPUD1 in

the response to ER stress induction, HERPUD1-WT cells and HERPUD1-ΔUBL cells were
treated with 2 μM Tg for 2, 4 and 6 h and we performed western blot analysis for XBP1
detection. We observed XBP1 protein was almost undetectable in both cell lines in the
absence of a stressor (Fig. 5B, lane 1 and 5). In contrast, we observed a robust induction of
XBP1 in both cell lines upon treatment for 2, 4 and 6 h with 2 μM Tg, observing similar XBP1
expression levels after 6 h of treatment in both cell lines (Fig. 5B, lane 4 and 8). Quantitative
analysis confirmed this conclusion, observing no significant differences between HERPUD1
WT (58.99 ± 9.18) and HERPUD1-ΔUBL (59.59 ± 17.49) expressing cells (Fig. 5C). In
addition to XBP1, we tested by western blot analysis the ER resident transmembrane protein
PERK, a kinase that undergoes hyperphosphorylation in response to ER stress shown as a
shift in its mobility during SDS-polyacrylamide gel electrophoresis (Bertolotti et al. 2000).
This kinase mediates the attenuation of the global translation by the phosphorylation of
eIF2α (Bertolotti et al. 2000; Harding, Zhang, et al. 2000). In contrast to its differential
electrophoretic migration with Tun and Tg (Suppl. Fig. 4, lane 2 and 3), no changes in PERK
migration were observed in HERPUD1-WT or HERPUD1-ΔUBL cells (Suppl Fig. 4, lane 4
and 5). Analysis of the activating transcription factor 4 (ATF4), a protein highly expressed
upon UPR response (Harding, Novoa, et al. 2000) showed again no changes in ATF4 levels
in either cell line (Suppl. Fig. 4, lane 4 and 5). In contrast, a robust detection of ATF4 was
found upon the addition of Tun and Tg (Suppl. Fig. 4, lane 2 and 3). Altogether, our findings
strongly indicate that the maintenance of the ER expansion triggered by HERPUD1 stability
is not the consequence of ER stress.
536

537 *Increased HERPUD1 stability by the deletion of its UBL domain leads to a remarkable*
538 *increase in lysosomal number and function*

539
ER proliferation that excludes ribosomes is able to regulate ER contacts with lysosomes
(Friedman et al. 2013; C. A. Lee and Blackstone 2020). Lysosomes are an important
catabolic organelle of eukaryotic cells, containing a diverse repertoire of acidic hydrolases
(luminal pH of 4.5–5.0) that can digest macromolecules such as sugars, lipids and proteins
and even entire organelles (H. Xu and Ren 2015). We asked whether expanded ER by
increased HERPUD1 stability could have an impact on endolysosomal organelles. To
visualize these compartments, HERPUD1-WT and HERPUD1-ΔUBL expressing HeLa cells
were stained with an antibody against the endogenous lysosomal-associated membrane
protein 1 (LAMP1), a membrane protein found on late endosomes and lysosomes (Griffiths
et al. 1988). Unexpectedly, we observed a robust increase in LAMP1 positive structures
located around the entire cytoplasm (Fig. 6A). In agreement with this finding, quantification
analysis showed a significant increase in the number of LAMP1 positive structures in
HERPUD1-ΔUBL (88.72 ± 16.77) respect to control cells (69.03 ± 13.79) (Fig.6B). To
determine if the increase in LAMP1 structures corresponded to lysosome organelles, we
performed co-staining of LAMP1 with CATHEPSIN-D (CAT-D), a luminal hydrolytic
lysosomal enzyme (Benes, Vetvicka, and Fusek 2008), observing several LAMP1 positive
structures positive to CAT-D (Suppl. Fig 5A, lower merge). In addition, we noticed a
significant increase in the integrated fluorescence intensity of CAT-D in HERPUD1-ΔUBL
(1.47 ± 0.88), compared to HERPUD1-WT expressing cells (1.00 ± 0.20) (Suppl. Fig 5B).
Considering this increase in CAT-D fluorescence we evaluated if HERPUD1-ΔUBL also
impacts lysosomal activity, as assessed by measuring lysosomal acidity and activity in live
cells. We first measured if HERPUD1 stability affects the range of lysosomal pH, using the

pH-sensitive lysosomal dye LysoTracker Red, which accumulates and emits red
fluorescence in acidic compartments with pH <6.5 (De Duve et al. 1974; Chou, Paul
Krapcho, and Hacker 2001). Then, we measured CATHEPSIN-B activity using the Magic
Red assay (Boonacker et al. 2003). As shown in Figure 6C, expression of HERPUD1-ΔUBL
causes a significant increase in the number of LysoTracker positive structures (211.95 ±
78.90), compared to control cells (117.85 ± 48.97) (Fig. 6C and 6D). Similarly, we found an
increase in the number of structures positive to Magic Red fluorescence (Figure 6E). This
was confirmed by measuring the integrated fluorescence intensity, which revealed a
significant increase of this parameter in HERPUD1-ΔUBL cells (1.56 ± 0.43), compared to
control cells (1.00 ± 0.19) (Fig. 6F). Altogether, these findings show that ER expansion
triggered by HERPUD1 increased stability is a potent strategy to promote lysosomal
function.
574
575 ***The phosphomimetic mutant S59D in the UBL domain promotes HERPUD1 stability***
576 ***mimicking the effects of UBL deletion on the ER and autophagy.***
577
Further, we searched for a mechanism that could mimic the phenotype observed by the
deletion of UBL on HERPUD1. As shown by the analysis with PyMOL molecular graphics
system (Suppl. Fig 6A), the UBL domain of HERPUD1 (PDB 1WGD; green color) resembles
UBIQUITIN (PDB 2MSG; purple color) in terms of their three-dimensional structure.
UBIQUITIN as well as UBL domains are known targets of phosphorylation under cellular
stress (Kondapalli et al. 2012; Swaney, Rodríguez-Mias, and Villén 2015; Sauvé et al. 2018).
In fact, oxidative stress promotes phosphorylation of UBIQUITIN at Ser65 causing the
accumulation of ubiquitylated proteins due to a reduction in global protein turnover rates
(Koyano et al. 2014; Swaney, Rodríguez-Mias, and Villén 2015). In the same line, it is known
that the UBL domain on PARKIN is phosphorylated by PINK1, and is responsible for the
activation of its E3 ubiquitin-ligase activity (Sauvé et al. 2018), both essential players of
mitochondria quality control by mitophagy (Jin and Youle 2012). Thereby, we searched for
Ser residues as possible candidates of phosphorylation in the UBL domain of HERPUD1
using the KinasePhos2.0 web tool. Five residues were predicted in position Ser16, Ser27,
Ser33, Ser59 and Ser90 (Suppl. Fig 6B, Ser residues in red color). For all these five Ser
residues, we generated phosphoinert (substitutions to alanine) and phosphomimetic
(substitutions to aspartic acid) mutant versions. Stably transfected HeLa cells were
generated for each mutant. Among all mutations tested, the substitution S59D, but not the
S59A, showed the strongest increase in HERPUD1 signal, as is shown by
immunofluorescence with an anti-FLAG antibody (Fig. 7A and 7B). Importantly, we found
that the phosphomimetic S59D mutant presents an ER pattern very similar to HERPUD1-
ΔUBL cells, confirmed by the co-staining with GRP94 (Fig. 7A, right panel merge). In
contrast, no changes were observed with the phosphoinert S59A mutant, showing a similar
phenotype than HERPUD1-WT (Fig. 7A and 7B). Thus, further characterization included the
comparison between HERPUD1-WT and HERPUD1-S59D. In addition, under basal
conditions western blot analysis using anti-HERPUD1 and anti-FLAG antibodies showed
higher levels of HERPUD1-S59D (40.09 ± 5.31) compared to either HERPUD1-WT (1.00 ±
0.53) or HERPUD1-S59A (2.88 ±1.76) (Fig. 7C, lane 5 compared to lane 1 and 3 and Fig.
7D). We also measured the levels of these versions upon treatment for 4 h with 20 mM
MG132, which is a proteasomal inhibitor, known to abolish HERPUD1 proteasomal
degradation (Yan et al. 2014). As expected, MG132 treatment caused a significant increase
in HERPUD1-WT (16.55 ± 7.13) (Fig. 7C, lane 1 compared to lane 2 and Fig. 7D). A similar

result was obtained with HERPUD1-S59A (20.36 ± 9.65) (Fig. 7C, lane 3 compared to lane 4
and Fig. 7D). In contrast, we observed non-significant differences in the levels of HERPUD1-
S59D in the absence (40.09 ± 5.31) or presence of MG132 (47.17 ± 11.74) (Fig. 7C, lane 5
and 6 and Fig. 7D). Further, and based on the negative impact of HERPUD1 stability on
autophagy by the deletion of its UBL domain, we next investigated if the stability of
HERPUD1-S59D could have a similar outcome. Biochemically, we found that HERPUD1-
S59D cells have a significant reduction in the LC3-II/LC3-I ratio (0.66 ± 0.02), in comparison
to HERPUD1-WT (1.00 ± 0.18) (Fig. 7E and 7F). This result was corroborated by
immunofluorescence analysis of LC3 in the absence or presence of BafA1 showing that
HERPUD1-S59D cells present a lesser number of autophagosomes, compared to
HERPUD1-WT (Suppl. Fig. 7). Moreover, in agreement with previous reports indicating that
a reduction in autophagy enhances ER expansion and cell size (Khaminets et al. 2015;
Miettinen and Björklund 2015), we noticed that HERPUD1-S59D cells were significantly
larger in size (2.52 ± 0.98), compared to HERPUD1-WT (1.00 ± 0.4) and HERPUD1-S59A
cells (0.88 ± 0.29) (Fig. 7A and Suppl. Fig. 8A). Importantly, cell size quantification showed
HERPUD1-ΔUBL were also larger in size (1.36 ± 0.64) than HERPUD1-WT (1.00 ± 0.4), but
smaller than HERPUD1-S59D (2.52 ± 0.98) (Suppl. Fig. 8A). In addition, because nuclei size
remains proportional to the cell size in a wide range of genetic backgrounds and growth
conditions (Huber and Gerace 2007), we studied the nuclei size in all cell lines. Surprisingly,
nuclei size in HERPUD1-S59D cells were significantly larger (1.74 ± 0.72) in comparison to
all tested cell lines; HERPUD1-WT (1.00 ± 0.49), HERPUD1-S59A (1.12 ± 0.47) and
HERPUD1-ΔUBL (1.11 ± 0.52) (Suppl. Fig. 8B). Altogether, our findings support the idea
that HERPUD1 increased stability might play an important function in cellular plasticity by
commanding a program that controls ER expansion with impact in cell and nuclei size. The
results with the phosphomimetic version of HERPUD1 opens the alternative of a program
activated under the control of phosphorylation.

The phosphomimetic HERPUD1-S59D mutant promotes ER-lysosomal network with
impact in stress cell survival

Because of the increase in the number of functional lysosomes under the expression of
HERPUD1-ΔUBL, we investigated if HERPUD1-S59D expressing cells could have a similar
output. First, we performed immunofluorescence of LAMP1, observing a significant increase
in the number of LAMP1 positive structures in HERPUD1-S59D cells (135.28 ± 21.32), in
comparison to HERPUD1-WT (69.03 ± 13.79) and HERPUD1-S59A (58.42 ± 14.55) (Supp.
Fig. 9A and 9B). Next, analysis with LysoTracker in HERPUD1-S59D cells (183.35 ± 60.05),
in comparison to HERPUD1-WT (118.18 ± 51.00) and HERPUD1-S59A (86.13 ± 41.46)
confirmed HERPUD1-S59D cells have a significant increase in the number of lysosomes per
cell (Suppl. Fig. 9C and 9D). Similar results were obtained with Magic Red where the
quantification showed an increase of almost twice the intensity of Magic Red in HERPUD1-
S59D cells (2.25 ± 0.32) in comparison with HERPUD1-WT (1.00 ± 0.15) and HERPUD1-
S59A (1.02 ± 0.24) (Suppl. Fig 9E and 9F).

One important aspect of cellular plasticity is the establishment of a network between the ER
and the lysosomes that helps in the constant adaptation to particular cellular needs. In this
regard, it is known that the ER forms membrane-contact sites (MCSs) with lysosomes (Valm
et al. 2017), acting as a potent spatiotemporal organizer of endolysosomal biology (Neefjes,
Jongsma, and Berlin 2017). We hypothesized that the expansion of the ER-lysosomal

network should be accompanied by the appearance of MCSs between these two organelles.
Therefore, we studied the spatial distribution of the ER and lysosomes upon stability of
HERPUD1 with the expression of HERPUD1-ΔUBL and HERPUD1-S59D. For this, multiple
confocal images were obtained with CALNEXIN and LAMP1 co-staining by
immunofluorescence and analyzed in a super-resolution scale. As expected, we observed
that similar to HERPUD1-WT expression, expression of HERPUD1-ΔUBL and HERPUD1-
S59D form ER-lysosomes MCSs (Fig. 8 middle panel). MCSs are also shown in a three-
dimensional (3D) reconstruction with high magnification (Fig. 8 inset lower panel).
Altogether, our findings strongly suggest that expansion of ER by HERPUD1 stability not
only promotes the remodeling of the lysosomal network but also the formation of MCSs
between these two organelles.

ER expansion has been known for decades as a trigger for crystalloid ER (Chin et al. 1982)
and ER whorls (Feldman, Swarm, and Becker 1981; Schuck et al. 2009), however only
recent studies have proposed this phenomenon could be part of a program necessary to
overcome ER stress with impact in stress cell survival (Schuck et al. 2009; F. Xu et al.
2021). Moreover, taking in consideration that HERPUD1 stability triggers ER expansion, ER-
lysosomal network and MCSs between these two organelles, a protein known to be
upregulated under ER stress, we studied the effect of HERPUD1 stability in the cell viability
of HeLa cells in response to cisplatin (CDDP). CDDP is the most frequently used
chemotherapeutic agent for the treatment of some types of cancers, including cervical
cancer in accordance with the model of HeLa cells (Dasari and Bernard Tchounwou 2014).
In addition, it is known that alleviation of ER stress attenuates CDDP-induced apoptosis (Wu
et al. 2018). Thus, we investigated the resistance to drug-induced cytotoxicity of HERPUD-
WT and HERPUD1-S59D cells in response to varying doses of CDDP for 24 h with the
Sulforhodamine B (SRB) assay. We found a significant increase in the resistance of
HERPUD1-S59D cells in comparison with HERPUD1-WT at 2, 4, 8 and 16 μM of CDDP
(Fig. 9A). We corroborated this result measuring apoptosis of these cells in the absence or
presence of 10 μM CDDP for 24 h by flow cytometry using Annexin-V staining method. In
agreement with the SRB assay, we observed HERPUD1-S59D cells showed less apoptosis
compared to HERPUD1-WT cells in response to CDDP treatment (Fig. 9B). These results
strongly indicate HERPUD1 stability helps to overcome cisplatin-induced cytotoxicity. In this
regard, we propose that this differential response could be mediated by the expansion of the
ER/lysosomal network, an aspect that should be investigated in other cancer cell models.

DISCUSSION

Post-translational responses allow cells to respond swiftly to stress conditions with
consequences in their proteome composition. Autophagy and the UPS are two closely
related pathways that adjust their functions in response to cellular demands in order to
maintain cellular homeostasis (Korolchuk, Menzies, and Rubinsztein 2010). During
starvation overall protein degradation rises by activation of both autophagy and the UPS
(Mizushima, Yoshimori, and Ohsumi 2011; VerPlank et al. 2019). Indeed, before up-
regulation of autophagy, efficient synthesis of new proteins is sustained by degradation of
preexisting proteins by the proteasome, which shows that the proteasome also plays a
crucial role in cell survival after nutritional stress (Vabulas and Hartl 2005). In agreement
with this, it is known that the proteasome activity transits from a latent to an activated state
(Asano et al. 2015; Collins and Goldberg 2020). However, if UPS activation upon starvation

mediates the degradation of specific proteins that could slow-down autophagy is unknown.
Downregulation of HERPUD1 upon nutrient starvation in ATG5-depleted cells opens that
alternative. In this regard, it has been previously proposed that HERPUD1 depletion up-
regulates autophagy (Quiroga et al. 2013) and the degradation of cytosolic protein
aggregates (Miura et al. 2010). These findings support the idea that basal levels of
HERPUD1 could act as a break in the activation of basal autophagy.

Under ER stress, HERPUD1 is upregulated even faster than ER chaperones (K. Kokame et
al. 2000; Bergmann et al. 2018), where it participates in Endoplasmic Reticulum-Associated
Degradation (ERAD) (Schulze et al. 2005). HERPUD1 facilitates the assembly of the HRD1
complex, also known as the retrotranslocon, key in the retrotranslocation of unfolded
proteins from the ER to the cytosol for proteasomal degradation upon demand (Leitman et
al. 2014; Schulz et al. 2017). However, the function of HERPUD1 in the absence of ER
stress is less understood. Here, we propose HERPUD1 acts as a negative regulator of
autophagy controlled by its proteasome dependent stability, a mechanism that could operate
in the absence of ER stress under the control of phosphorylation of its UBL domain.

One interesting aspect of HERPUD1 is the UBL domain present in its N-terminus region. In
general, UBL-containing proteins share the ability to interact with the 19S regulatory particle
of the 26S proteasome promoting its activation (Rasmus Hartmann-Petersen and Gordon
2004; Yu and Matouschek 2017; Collins and Goldberg 2020). The UBL domain itself can
stimulate multiple proteasome activities in a similar fashion to ubiquitin chains (H. T. Kim and
Goldberg 2018; Collins and Goldberg 2020). However, unlike ubiquitin and its homologous
(e.g., SUMO and Nedd8), UBL-containing proteins cannot be conjugated to other proteins.
The human genome encodes over 60 UBL-containing proteins, where 15 of them have been
studied in their ability to bind and regulate proteasome activity (Collins and Goldberg 2020).
HERPUD1 is one member of this group, however its role as a positive modulator of the
proteasomal activity remains uncharacterized. Moreover, because UBL-containing proteins
can stimulate proteasome activity (H. T. Kim and Goldberg 2018; Collins and Goldberg
2020), it would be interesting to explore the effect of HERPUD1-ΔUBL and HERPUD1-S59D
on the activity of the proteasome under normal and starvation conditions.

In addition, recent findings show UBL-containing proteins can also play a regulatory role in
autophagy, such as USP14 (E. Kim et al. 2018), NUB1 (Guarascio et al. 2020), Elongin B
(Antonioli et al. 2016), UHRF1 (Shi et al. 2020), OASL (Batista-Silva et al. 2018), BAT3
(Sebti et al. 2014) and UBQLN (Rothenberg et al. 2010; Şentürk et al. 2019; Yang and
Klionsky 2020). Our findings show that the stabilization of HERPUD1 by lacking its UBL
domain causes a reduction in LC3-II/LC3-I ratio, which positions HERPUD1 as a member of
this growing list of UBL-containing proteins that function as regulators of autophagy.
However, how HERPUD1 stability mediates the reduction in LC3-II/LC3-I ratio needs further
studies. One aspect to be explored is the finding that HERPUD1 interacts with UBQLN (T. Y.
Kim et al. 2008). UBQLN is a cytosolic protein, that in addition to HERPUD1, interacts with
LC3 and ubiquitinated cargos in autophagosomes (Rothenberg et al. 2010). Furthermore, it
is known that silencing UBQLN leads to a reduction in the lipidation of LC3-I to LC3-II that
correlates with a diminished number of autophagosomes (Rothenberg et al. 2010). Because
UBQLN binds HERPUD1 independently of its UBL domain (T. Y. Kim et al. 2008), it is
possible that an increase in HERPUD1 stability at the ER could mediate the sequestration of
UBQLN in this compartment affecting its function in other membranes such as

autophagosomes (Rothenberg et al. 2010). In this regard, it has been previously proposed
that recruitment of UBQLN to the ER by HERPUD1 could bring the proteasome and the
ubiquitinated substrates to specific microdomains of the ER, which could hypothetically be
the step that promotes the ERAD pathway (T. Y. Kim et al. 2008). Because our results show
that HERPUD1 stability maintain the ER-lysosomal network, additional work is needed to
determine if HERPUD1:UBQLN interaction could play a role in the delivery of substrates to
the ER-to-lysosomes-associated degradation (ERLAD) (Fregno et al. 2018; Fregno and
Molinari 2019), considering the increase in the lysosomal degradation function. As
HERPUD1-S59D mimics the effect on HERPUD1 stability we propose that recruitment of
UBQLN at the ER is controlled by phosphorylation of HERPUD1.

Ubiquitin and UBL domains are targets of phosphorylation (Kondapalli et al. 2012; Koyano et
al. 2014; Wauer et al. 2015). The best-known example is the phosphorylation of the UBL
domain of the ubiquitin ligase PARKIN by the Ser/Thr kinase PINK1 on Ser65 (Kondapalli et
al. 2012; Koyano et al. 2014). This post-translational modification orchestrates its enzymatic
E3 ligase activity (Aguirre et al. 2017), participating in the ubiquitination of mitochondrial
proteins during mitophagy. Here, we propose that phosphorylation of the UBL domain in
HERPUD1 must affect its noncovalent binding to the proteasome explaining its stability
increase, as occurs with HERPUD1-ΔUBL. Further studies are needed to define the
physiological triggers of HERPUD1 phosphorylation, and the kinase involved.

This study is the first report indicating HERPUD1 stability mediates ER expansion but not as
a consequence of ER stress. In this regard, it is well known that the ER expands to alleviate
ER stress (Schuck et al. 2009). However, the ER expansion phenomenon is not always
homeostatic. ER expansion requires an adequate supply of membrane lipids although the
mechanisms that govern ER biogenesis are yet unclear. For example, the ER expands
several folds when B lymphocytes differentiate into antibody-secreting plasma cells (Wiest et
al. 1990), when hepatocytes increase its P450 detoxification system (Feldman, Swarm, and
Becker 1981), in response to Epidermal Growth Factor (EGF) (Caldieri et al. 2017) and
statins (Chin et al. 1982). Interestingly, lipid synthesis activation causes expansion of the ER
and resistance to ER stress even in cells lacking the UPR, highlighting the physiological
importance of ER membrane biogenesis in homeostasis. A future challenge will be to sort
out the connection between HERPUD1 and lipid synthesis, a link recently suggested by
genomics (Van Der Laan et al. 2018).

In this regard, an interesting observation to be considered is that HERPUD1 increased
stability mimics the effect of statins in reference to the appearance of a crystalloid ER (Chin
et al. 1982). Statins are cholesterol reducing agents acting as blockers of the cholesterol
biosynthesis by the inhibition of 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA)
reductase. Interestingly, statins possess beneficial effects in a variety of human diseases.
Importantly, a growing number of studies refer to statins as ER stress reducing agents
(Mollazadeh et al. 2018; T. Zhang et al. 2018), modulators of autophagy (Ashrafizadeh et al.
2020) and inducers of lysosomal biogenesis (Y. Zhang et al. 2020). However, if statins
promote those effects by a mechanism related with HERPUD1 stability is unknown.

Expansion of ER by HERPUD1 stability is correlated with a slow-down in autophagy. In
agreement with this, it is known that the knockdown of the two major autophagy regulators,
ATG5 and BECN1, likewise trigger ER expansion (Khaminets et al. 2015). Moreover, a

similar ER expansion is also observed by ER-phagy inhibition, a selective form of autophagy
responsible for the degradation of excess ER (Khaminets et al. 2015; Grumati, Dikic, and
Stolz 2018). However, studies investigating if the UPR is activated or not in response to
inhibition of autophagosomal biogenesis or ER-Phagy dysfunction are still lacking. While the
opposite effect has been reported, observing that excessive ER-Phagy results in activation
of the UPR response (Liao et al. 2019).

Our findings also indicate that ER expansion by HERPUD1 stability is accompanied by an
increase in the number of functional lysosomes. Because several studies have indicated that
UBL-containing proteins cause a positive modulation of the UPS system we can speculate
that HERPUD1-ΔUBL and HERPUD1-S59D could have a negative impact in the proteasome
activity. Disturbances in the UPS and autophagy function are known to be compensated by
lysosome biogenesis (Jackson and Hewitt 2016). In addition, we have already discussed the
possibility that HERPUD1 stability could trigger the recruitment of UBQLN to the ER.
Interestingly, it has been recently discovered that UBQLN plays a crucial role in the
maintenance of the acidic pH of lysosomes and in a closed interplay with the ER (Şentürk et
al. 2019; Yang and Klionsky 2020). Further work is needed to determine if
HERPUD1:UBQLN interaction could play a role at this level. Furthermore, based on recent
findings (Toulmay and Prinz 2011; Henne 2017), it is necessary to explore if MCSs between
ER and lysosomes could play a role in the transfer of lipids to lysosomes to compensate for
the massive expansion of the ER under HERPUD1 stability by degradation.

Together, our findings highlight novel insights into the possible role of HERPUD1 as a
regulator of autophagy and its participation in the maintenance of the ER structure.
Moreover, our results suggest that HERPUD1 stabilization promotes the lysosomal function
which could promote ER-lysosome intercommunication even in conditions where the UPR is
not activated. The mechanism behind this regulation remains to be elucidated, specially the
UBL phosphorylation-dependent stabilization of HERPUD1 and its effect on the pathways
discussed above. Furthermore, if this regulation has a role in cell pathological conditions it
should be analyzed in future studies.

Figure legends

Figure 1. SILAC-based proteomic study reveals HERPUD1 as a possible modulator of
autophagy. (A). Design of the SILAC experiment to monitor changes of the cellular
proteome of H4 cells during starvation and shRNA-mediated knockdown of ATG5 (shATG5)
or shRNA against the luciferase gene (shLuc). 'Reference' mixes of all cell extracts were
prepared for light and heavy conditions for 'forward' and 'reverse' SILAC experiments and
were mixed 1:1 with individual cell extracts, giving a total of 8 mixes. **(B-C)** Scatter plot
comparing 'Forward' and 'Reverse' Log2 Starvation/Normal ratio data for cells shLuc (B) and
shATG5 (C). **(D)** Comparison of average Log_2 Starvation/Normal ratios in shLuc (x-axis) and
shATG5 (y-axis). Protein outliers under either or both knockdown conditions are colored as
indicated.
829

Figure 2. Deletion of UBL domain increases HERPUD1 protein levels. (A) Schematic
representation HERPUD1-WT (left image) and ΔUBL (right image) at the ER membrane,
both FLAG-tagged at the carboxyl-terminus (shown in green). The Ubiquitin-Like domain

(UBL, amino acids 10-86) are represented in purple. **(B)** HeLa cells untransfected (UT) or
stably expressing the WT or ΔUBL versions of HERPUD1 were not treated (lanes 1, 3 and 5)
or treated with 20 μM of MG132 for 4 h (lanes 2, 4 and 6). Detergent-soluble protein extracts
were analyzed by western blot with anti-HERPUD1 and anti-FLAG antibodies. β-ACTIN was
used as loading control. Position of molecular mass markers is indicated on the left.

Figure 3. The stabilization of HERPUD1 by its UBL deletion negatively regulates
autophagy. (A) HeLa cells stably expressing HERPUD1-WT-FLAG or HERPUD1-ΔUBL-
FLAG were grown in glass coverslips and were treated or not with 100 nM BafA1 for 4 h.
Cells were fixed and incubated with an antibody to LC3 followed by incubation with Alexa-
488-conjugated donkey anti-rabbit IgG. Stained cells were examined by confocal
microscopy. Scale bar 10 μm. **(B)** HeLa cells stably expressing HERPUD1-WT-FLAG or
HERPUD1-ΔUBL-FLAG untreated (lanes 1 and 2) or treated with 100 nM BafA1 for 4 h
(lanes 3 and 4). Detergent-soluble protein extracts were analyzed by western blot with a
rabbit polyclonal antibody against LC3. Monoclonal antibody against β-ACTIN (clone BA3R)
was used as loading control. Position of molecular mass markers is indicated on the left. **(C)**
Densitometry quantification of LC3-I and LC3-II protein levels from images as those shown
in B. Relative levels are expressed as the ratio of LC3-II to LC3-I. Bars represent the mean ±
standard deviation. Statistical analysis was performed using two-tailed unpaired Student's t-
test (n = 3 ** p < 0.01, *** p < 0.001)

Figure 4. HERPUD1-ΔUBL stabilization alters the ER morphology. (A) HeLa cells stably
expressing HERPUD1-WT-FLAG or HERPUD1-ΔUBL-FLAG grown in glass coverslips were
fixed, permeabilized and triple-labeled with a mouse monoclonal antibody against FLAG,
with a rabbit monoclonal antibody against CALNEXIN and with a rat monoclonal antibody
against GRP94 followed by incubation with Alexa-488-conjugated donkey anti-rabbit IgG
(green channel), Alexa-594-conjugated donkey anti-rat IgG (red channel) and Alexa-647-
conjugated donkey anti-mouse IgG (blue channel). Images were acquired using a TCS SP8
laser-scanning confocal microscope. The fourth image on each row is the merge of blue,
green and red channels; yellow indicates colocalization of the red and green channels, cyan
indicates colocalization of the green and blue channels, magenta indicates colocalization of
the red and blue channels, and white indicates colocalization of all three channels. Scale
bar, 10 μm. **(B)** HeLa cells stably expressing HERPUD1-WT-FLAG or HERPUD1-ΔUBL-
FLAG grown in glass coverslips were fixed, permeabilized and labeled with rat monoclonal
antibody against GRP94 followed by incubation with Alexa-594-conjugated donkey anti-rat
IgG. Stained cells were examined by fluorescence microscopy. Images were acquired using
a TCS SP8 laser-scanning confocal microscope. Pseudocolor image was created using all
the serial confocal sections of HeLa HERPUD1-WT-FLAG (left image) and HERPUD1-
ΔUBL-FLAG (right image). The scale on the right represents the maximum (red) to minimum
(black) intensity measured for GPR94. **(C)** Quantification of ER-volume obtained from serial
image reconstruction (z-stack 0.3 μm z-interval, 1024x1024, 180 μm pixel size). Volume is
depicted in a scatter plot, open circles represent HeLa HERPUD1-WT-FLAG (n=79) and
open squares HERPUD1-ΔUBL-FLAG (n=72). Statistical analysis was performed using two-
tailed unpaired Student's t-test (*** p < 0.001). **(D)** TEM micrograph shows HeLa HERPUD1-
WT-FLAG or HERPUD1-ΔUBL-FLAG (left and center images, respectively) at a lower
magnification. Crystalloid ER structure is visible as a honeycomb in HERPUD1-ΔUBL
(central image). An ER crystalloid structure from HERPUD1-ΔUBL-FLAG at a higher

magnification is shown (right image from dashed square in the center image, ER:
endoplasmic reticulum; M: mitochondria). Scale bar 500 nm.

Figure 5. The stability of HERPUD1 by the deletion of its UBL domain does not trigger
endoplasmic reticulum stress. (A) Splicing of *XBP1* was analyzed by RT-PCR. Total
RNAs were obtained from HeLa cells stably expressing HERPUD1-WT-FLAG or HERPUD1-
ΔUBL-FLAG, additionally, total RNA was obtained from HeLa WT cells untreated or treated
with 2 μM Thapsigargin (Tg) or 5 μg/mL Tunicamycin (Tun) for 4 h as a control of XBP1
splicing. Then, cDNA was synthesized and mRNA expression of XBP1 was analyzed using
specific primers. (*uXBP1*: unspliced form, *sXBP1*: spliced form). CYCLOPHILIN-A (CYC-A)
was used as an internal control. **(B)** HeLa cells stably expressing HERPUD1-WT-FLAG or
HERPUD1-ΔUBL-FLAG were treated with 2 μM Thapsigargin (Tg) for different timepoints (0
h, 2 h, 4 h and 6 h). Detergent-soluble protein extracts were analyzed by western blot with a
monoclonal antibody against XBP1. Monoclonal antibody against β-ACTIN was used as
loading control. Position of molecular mass markers is indicated on the left. **(C)** Densitometry
quantification of XBP1 protein levels from images as those shown in B. Bars represent the
mean ± standard deviation. Statistical analysis was performed using two-tailed unpaired
Student's t-test (n = 3 n.s not statistically significant, ** $p < 0.01$, *** $p < 0.001$).

Figure 6. Expression of HERPUD1-ΔUBL increases lysosomal number and function.
(A) HeLa cells stably expressing HERPUD1-WT-FLAG or HERPUD1-ΔUBL-FLAG grown in
glass coverslips were fixed, permeabilized and labeled with a mouse monoclonal antibody
against LAMP1 followed by incubation with Alexa-594-conjugated donkey anti-mouse IgG,
and DAPI for nuclei staining. Stained cells were examined by fluorescence microscopy.
Images were acquired using a TCS SP8 laser-scanning confocal microscope. Scale bar 10
μm. **(B)** Quantification of LAMP1 positive structures per cell. In the scatter plot each circle
and square represents the average of positive LAMP1 dots per cell from a frame of HeLa
HERPUD1-WT (n=559) or HERPUD1-ΔUBL (n=700) respectively. The quantified cells were
from 3 independent experiments. **(C and E)** HeLa cells stably expressing HERPUD1-WT-
FLAG or HERPUD1-ΔUBL-FLAG were grown in glass bottom culture dishes and labeled
with **(C)** LysoTracker™ Red DND-99 or **(E)** Magic Red®. For live cell imaging analysis,
culture medium was replaced with phenol red-free DMEM supplemented with HEPES (10
mM, pH 7.4) and images were acquired with TCS SP8 laser-scanning confocal microscope
at 37ºC. Scale bar 10 μm. **(D)** Quantification of LysoTracker™ dots per cell. Open circles
and open squares represent the average of fluorescence signals for each cell of HeLa
HERPUD1-WT-FLAG (n=139) or HERPUD1-ΔUBL-FLAG (n=138) respectively. **(F)**
Quantification of Magic Red® positive structures. In the scatter plot each circle and square
represents the average of positive dots for Magic Red per cell from a frame of HeLa
HERPUD1-WT-FLAG (n=1279) or HERPUD1-ΔUBL-FLAG (n=984) respectively. The
quantified cells were from 3 independent experiments. Statistical analysis was performed
using two-tailed unpaired Student's t-test (*** $p < 0.001$)

Figure 7. Stabilization of HERPUD1 by S59D mutation alters the ER morphology and
decreases autophagy. (A) HeLa cells stably expressing HERPUD1-WT-FLAG, HERPUD1-
S59A-FLAG or HERPUD1-S59D-FLAG were grown in glass coverslips, fixed, permeabilized,
and double-labeled with mouse monoclonal antibody against FLAG and with a rat
monoclonal antibody against GRP94 followed by incubation with Alexa-488-conjugated
donkey anti-rabbit IgG (green channel) and Alexa-594-conjugated donkey anti-rat IgG (red

channel). Images were acquired using a TCS SP8 laser-scanning confocal microscope. The third image on each column is the merge of green and red channels; yellow indicates colocalization of red and green channels. Scale bar: 10 μm. **(B)** Quantification of fluorescence FLAG signal from indicated cells. **(C)** HeLa cells stably expressing HERPUD1-WT-FLAG, HERPUD1-S59A-FLAG or HERPUD1-S59D-FLAG were treated or not with 10 μM of MG132 for 4 h. Detergent-soluble protein extracts were analyzed by western blot with anti-HERPUD1 and anti-FLAG antibodies. β-ACTIN was used as a loading control. Position of molecular mass markers is indicated on the left. **(D)** Densitometry quantification of HERPUD1 western blot signal from images as those shown in C. Bars represent the mean ± standard deviation. **(E)** HeLa cells stably expressing HERPUD1-WT-FLAG or HERPUD1-S59D-FLAG untreated (lanes 1 and 3) or treated with 100 nM BafA1 for 4 h (lanes 2 and 4). Detergent-soluble protein extracts were analyzed by western blot with a rabbit polyclonal antibody to LC3B. Monoclonal antibody against β-ACTIN was used as a loading control. Position of molecular mass markers is indicated on the left. **(F)** Densitometry quantification of LC3-I and LC3-II protein levels from images as those shown in E. Relative levels are expressed as the ratio of LC3-II to LC3-I from images as those shown in E. Bars represent the mean ± standard deviation. Statistical analysis was performed using two-tailed unpaired Student's t-test (n = 3 n.s. not statistically significant, *p < 0.05, ** p < 0.01, *** p < 0.001)

Figure 8. Formation of lysosomal membrane contact sites with the extended ER network in HERPUD1-ΔUBL and HERPUD1-S59D cells. HeLa cells stably expressing HERPUD1-WT-FLAG, HERPUD1-ΔUBL-FLAG or HERPUD1-S59D-FLAG were grown in glass coverslips and then fixed, permeabilized and double-labeled with a rabbit monoclonal antibody against CALNEXIN and a mouse monoclonal antibody against LAMP1 followed by incubation with Alexa-488-conjugated donkey anti-rabbit IgG (green channel), Alexa-594-conjugated donkey anti-mouse IgG (red channel) and DAPI for nuclei staining. All images were acquired using a TCS SP8 laser-scanning confocal microscope in z-series. Upper pannel correspond to inverted z-projection of maximum intensity of CALNEXIN images showing the different ER patterns between cell lines. Middle pannel show 3D reconstructions of CALNEXIN (green channel) and LAMP1 (red channel) images. Scale bar, 10 μm. Lower pannel show a higher magnification of the 3D reconstructions. Scale bar, 1 μm (Note: HERPUD1-WT-FLAG cells are smaller than HERPUD1-S59A-FLAG and HERPUD1-S59D-FLAG).

Figure 9. Stabilization of HERPUD1 by S59D mutation decreases cell death mediated by CDDP. (A) HeLa cells stably expressing HERPUD1-WT-FLAG or HERPUD1-S59D-FLAG were treated with different concentrations of cisplatin (CDDP) and then a SRB assay was performed. The absorbance values of each point were normalized to control cells (without treatment) and transformed to a percentage. Experiments were performed at least three times, and the results are expressed as mean ± standard deviation. Statistical analysis was performed using two-tailed unpaired Student's t-test (n = 4 n.s. not statistically significant and *p < 0.05) **(B)** HeLa cells stably expressing HERPUD1-WT-FLAG or HERPUD1-S59D-FLAG were treated or not with CDDP 10 μM for 24 h. Afterwards, cells were collected by trypsinization and stained with Annexin V-Pacific Blue and 7AAD. Scatter plots representing the different quadrants are labeled as Q1 (necrotic cells), Q2 (viable cells), Q3 (dead cells) and Q4 (apoptotic cells). The graph shows the percentage of cells present in each quadrant.

Supplementary Figure 1. Table summarizing all proteins identified as significantly differing in abundance (Significance B <0.001) between normal and serum starved conditions for either shLuc or shATG5 treated cells, or both. Number cells are color coded by response to starvation; Blue = reduced abundance, red = increase, white = no change. Number Cells with a black border met significance criteria. Blank number cells correspond to missing data from one or both forward and reverse experiments.

Supplementary Figure 2. The stabilization of HERPUD1 by its UBL deletion negatively regulates autophagy under starvation. **(A)** HeLa cells stably expressing HERPUD1-WT-FLAG or HERPUD1-ΔUBL-FLAG untreated (lanes 1 and 2) or treated with EBSS for 4 h in the absence or presence of 100 nM BafA1 (lanes 3 to 6). Detergent-soluble protein extracts were analyzed by western blot with a rabbit polyclonal antibody against LC3. Monoclonal antibody against β-ACTIN (clone BA3R) was used as a loading control. Position of molecular mass markers is indicated on the left. **(B)** Densitometry quantification of LC3-I and LC3-II protein levels from images as those shown in A. Relative levels are expressed as the ratio of LC3-II to LC3-I. Bars represent the mean ± standard deviation. Statistical analysis was performed using two-tailed unpaired Student's t-test ($n = 3$ * $p < 0.05$, ** $p < 0.01$).

Supplementary Figure 3. HERPUD1-ΔUBL stabilization alters the ER morphology without changing ER resident protein levels. (A) HeLa cells stably expressing HERPUD1-WT-FLAG or HERPUD1-ΔUBL-FLAG grown in glass coverslips were fixed, permeabilized and incubated with an antibody against HERPUD1 followed by incubation with Alexa-488-conjugated donkey anti-rabbit IgG. Stained cells were examined by confocal microscopy. Scale bar 10 μm. **(B)** HeLa cells stably expressing HERPUD1-WT-FLAG or HERPUD1-ΔUBL-FLAG were grown in glass bottom culture dishes and labeled with ER-Tracker™ Blue-White DPX. Stained cells were examined by fluorescence microscopy. Images were acquired using a TCS SP8 laser-scanning confocal microscope. Scale bar 10 μm. **(C)** HeLa cells HERPUD1-WT-FLAG or HERPUD1-ΔUBL-FLAG. Detergent-soluble protein extracts were analyzed by western blot with a rabbit monoclonal antibody against CALNEXIN, rat monoclonal antibody against GRP94 and rabbit monoclonal antibody against HERPUD1. Monoclonal antibody against β-ACTIN (clone BA3R) was used as a loading control. Position of molecular mass markers is indicated on the left. **(D)** Densitometry quantification of CALNEXIN and GRP94 protein levels from images as those shown in C. Bars represent the mean ± standard deviation. Statistical analysis was performed using two-tailed unpaired Student's t-test ($n = 3$, n.s. not statistically significant).

Supplementary Figure 4. **The stability of HERPUD1 by the deletion of its UBL domain does not trigger endoplasmic reticulum stress.** HeLa WT cells untreated or treated with 2 μM Thapsigargin (Tg) or 5 μg/mL Tunicamycin (Tun) for 4 h (lane 1 to 3) and untreated HeLa cells stably expressing HERPUD1-WT-FLAG or HERPUD1-ΔUBL-FLAG (lane 4 and 5) were lysed. Detergent-soluble protein extracts were analyzed by western blot with a rabbit polyclonal antibody against P-PERK/PERK, rabbit monoclonal antibody against ATF4 and rabbit monoclonal antibody against HERPUD1. Monoclonal antibody against β-ACTIN (clone

1018 BA3R) was used as a loading control. Position of molecular mass markers is indicated on
1019 the left.
1020

Supplementary Figure 5. **Expression of HERPUD1-ΔUBL increases lysosomal**
organelle number. (A) HeLa cells stably expressing HERPUD1-WT-FLAG or HERPUD1-
ΔUBL-FLAG grown in glass coverslips were fixed, permeabilized, and double-labeled with
mouse monoclonal antibody against LAMP1 and with a goat polyclonal antibody against
CATHEPSIN-D (CAT-D) followed by incubation with Alexa-488-conjugated donkey anti-goat
IgG (green channel), Alexa-594-conjugated donkey anti-mouse IgG (red channel) and DAPI
for nuclei staining. Images were acquired using a TCS SP8 laser-scanning confocal
microscope. The third image on each row is the merge of red and green channels; yellow
indicates overlapping localization of the red and green channels. Scale bar, 10 μm. **(B)**
Quantification of CATHEPSIN-D positive structures per cell. In the scatter plot open circles
and open squares represent the average of fluorescence signal for each cell in a frame of
HeLa HERPUD1-WT-FLAG (n=559) or HERPUD1-ΔUBL-FLAG (n=700) respectively.
Quantified cells were from 3 independent experiments. Statistical analysis was performed
using two-tailed unpaired Student's t-test (*** $p < 0.001$).
1035

Supplementary Figure 6. Predicted phosphorylation sites of the UBL domain in
HERPUD1. **(A)** The Ubiquitin-like Domain of HERPUD1 (PDB 1WGD, green color) and
ubiquitin (PDB 2MSG, purple color) resolved structures were aligned by PyMOL molecular
graphics system. **(B)** Predicted phosphorylated serines are shown in red in the resolved
structure of the Ubiquitin-like Domain of HERPUD1 (PDB 1WGD, green color). Each arrow
points to the specific residue shown in the figure.
1042

Supplementary Figure 7. Stabilization of HERPUD1 by S59D mutation decreases
autophagosomes. HeLa cells stably expressing HERPUD1-WT-FLAG or HERPUD1-S59D-
FLAG grown in glass coverslips were treated or not with 100 nM BafA1 for 4 h. Cells were
fixed, permeabilized and incubated with a polyclonal rabbit antibody against LC3 followed by
incubation with Alexa-488-conjugated donkey anti-rabbit IgG. Stained cells were examined
by confocal microscopy. Images were acquired using a TCS SP8 laser-scanning confocal
microscope. Scale bar 10 μm.
1050

Supplementary Figure 8. **Expression of HERPUD1-S59D increases nuclear and cell**
size. (A) Cell size analysis of HeLa cells stably expressing HERPUD1-WT-FLAG,
HERPUD1-ΔUBL-FLAG, HERPUD1-S59A-FLAG and HERPUD1-S59D-FLAG. Cells grown
in glass coverslips were marked with Wheat Germ Agglutinin (WGA), fixed and stained with
DAPI for nuclei visualization. Stained cells were examined by fluorescence microscopy. Cell
area was measured by selection of cell contour. **(B)** Analysis of the nuclei size of HeLa
HERPUD1-WT-FLAG, HERPUD1-ΔUBL-FLAG, HERPUD1-S59A-FLAG and HERPUD1-
S59D-FLAG. Cells grown in glass coverslips were stained with Hoechst for nuclei
visualization. Stained cells were examined by fluorescence microscopy. Nuclei area was
measured by selection of nuclei contour.
1061

1062 **Supplementary Figure 9. Expression of HERPUD1-S59D increases lysosomal number**
1063 **and function. (A)** HeLa cells stably expressing HERPUD1-WT-FLAG, HERPUD1-S59A-
1064 FLAG or HERPUD1-S59D-FLAG grown in glass coverslips were fixed, permeabilized and
1065 labeled with a mouse monoclonal antibody against LAMP1 followed by incubation with

Alexa-594-conjugated donkey anti-mouse IgG and DAPI for nuclei staining. Stained cells were examined by fluorescence microscopy. Images were acquired using a TCS SP8 laser-scanning confocal microscope. Scale bar 10 μm. **(B)** Quantification of LAMP1 positive structures per cell. In the scatter plot each circle, triangle and rhombus represent the average of positive LAMP1 dots per cell from a frame of HeLa HERPUD1-WT-FLAG (n=559), HERPUD1-S59A-FLAG (n=557) and HERPUD1-S59D-FLAG (n=575) respectively. The quantified cells were from 3 independent experiments. **(C and E)** HeLa cells stably expressing HERPUD1-WT-FLAG, HERPUD1-S59A-FLAG or HERPUD1-S59D-FLAG were grown in glass bottom culture dishes and labeled with **(C)** LysoTracker™ Red DND-99 or **(E)** Magic Red®. For live cell imaging, cell culture medium was replaced with phenol red-free DMEM supplemented with HEPES (10 mM, pH 7.4), and images were acquired with TCS SP8 laser-scanning confocal microscope at 37ºC. Scale bar 10 μm. **(D)** Quantification of LysoTracker™ dots per cell. Open circles, triangles and rhombuses represent the average of fluorescence signals for each cell of HeLa HERPUD1-WT-FLAG (n=223), HERPUD1-S59A-FLAG (n=233) or HERPUD1-S59D-FLAG (n=124) respectively. **(F)** Quantification of Magic Red® positive structures. In the scatter plot each open circle, triangle and rhombus represent the average of positive dots for Magic Red per cell from a frame of HeLa HERPUD1-WT-FLAG (n=718), HERPUD1-S59A-FLAG (n=1183) or HERPUD1-S59D-FLAG (n=670) respectively. The quantified cells were from 3 independent experiments. Statistical analysis was performed using two-tailed unpaired Student's t-test (*** $p < 0.001$).

REFERENCES

Aguirre, Jacob D., Karen M. Dunkerley, Pascal Mercier, and Gary S. Shaw. 2017. "Structure of Phosphorylated UBL Domain and Insights into PINK1-Orchestrated Parkin Activation." *Proceedings of the National Academy of Sciences of the United States of America* 114(2): 298–303. https://pubmed.ncbi.nlm.nih.gov/28007983/ (March 30, 2021).

Anderson, R. G.W., L. Orci, and M. S. Brown. 1983. "Ultrastructural Analysis of Crystalloid Endoplasmic Reticulum in UT-1 Cells and Its Disappearance in Response to Cholesterol." *Journal of Cell Science*.

Antonioli, Manuela, Federica Albiero, Mauro Piacentini, and Gian Maria Fimia. 2016. "Temporal Regulation of Autophagy Response by the CULLIN 4-AMBRA1-CULLIN 5 Axis." *Molecular and Cellular Oncology* 3(5). https://pubmed.ncbi.nlm.nih.gov/27857967/ (March 30, 2021).

Asano, Shoh et al. 2015. "A Molecular Census of 26S Proteasomes in Intact Neurons." *Science* 347(6220): 439–42. https://pubmed.ncbi.nlm.nih.gov/25613890/ (March 30, 2021).

Ashrafizadeh, Milad, Zahra Ahmadi, Tahereh Farkhondeh, and Saeed Samarghandian. 2020. "Modulatory Effects of Statins on the Autophagy: A Therapeutic Perspective." *Journal of Cellular Physiology* 235(4): 3157–68. https://pubmed.ncbi.nlm.nih.gov/31578730/ (March 30, 2021).

Batista-Silva, Leonardo Ribeiro, Rychelle Clayde Affonso Medeiros, Flávio Alves Lara, and Milton Ozório Moraes. 2018. "Type I Interferons, Autophagy and Host Metabolism in Leprosy." *Frontiers in Immunology* 9(APR). https://pubmed.ncbi.nlm.nih.gov/29755459/ (March 30, 2021).

Benes, Petr, Vaclav Vetvicka, and Martin Fusek. 2008. "Cathepsin D-Many Functions of One Aspartic Protease." *Critical Reviews in Oncology/Hematology* 68(1): 12–28.

Bergmann, Timothy J. et al. 2018. "Chemical Stresses Fail to Mimic the Unfolded Protein Response Resulting from Luminal Load with Unfolded Polypeptides." *Journal of*

Biological Chemistry 293(15): 5600–5612. https://pubmed.ncbi.nlm.nih.gov/29453283/ (June 21, 2020).

Bertolotti, Anne et al. 2000. "Dynamic Interaction of BiP and ER Stress Transducers in the Unfolded-Protein Response." Nature Cell Biology 2(6): 326–32. https://pubmed.ncbi.nlm.nih.gov/10854322/ (March 31, 2021).

Blois, Joseph, Adam Smith, and Lee Josephson. 2011. "The Slow Cell Death Response When Screening Chemotherapeutic Agents." Cancer Chemotherapy and Pharmacology 68(3): 795–803. https://pubmed.ncbi.nlm.nih.gov/21193989/ (March 30, 2021).

Boonacker, Emil et al. 2003. "Rapid Assay to Detect Possible Natural Substrates of Proteases in Living Cells." BioTechniques 35(4): 766–74. https://pubmed.ncbi.nlm.nih.gov/14579742/ (March 31, 2021).

Borgese, Nica, Maura Francolini, and Erik Snapp. 2006. "Endoplasmic Reticulum Architecture : Structures in Flux." Current Opinion in Cell Biology 18(4): 358–64. https://pubmed.ncbi.nlm.nih.gov/16806883/ (March 30, 2021).

Bustamante, Hianara A. et al. 2018. "Interplay between the Autophagy-Lysosomal Pathway and the Ubiquitin-Proteasome System: A Target for Therapeutic Development in Alzheimer's Disease." Frontiers in Cellular Neuroscience 12. https://pubmed.ncbi.nlm.nih.gov/29867359/ (March 30, 2021).

Bustamante, Hianara A et al. 2020. "The Proteasomal Deubiquitinating Enzyme PSMD14 Regulates Macroautophagy by Controlling Golgi-to-ER Retrograde Transport." Cells 9(3): 777. https://pubmed.ncbi.nlm.nih.gov/32210007/ (June 23, 2020).

Caldieri, Giusi et al. 2017. "Reticulon 3-Dependent ER-PM Contact Sites Control EGFR Nonclathrin Endocytosis." Science 356(6338): 617–24. https://pubmed.ncbi.nlm.nih.gov/28495747/ (March 30, 2021).

Camuzard, Olivier, Sabine Santucci-Darmanin, Georges F. Carle, and Valérie Pierrefite-Carle. 2020. "Autophagy in the Crosstalk between Tumor and Microenvironment." Cancer Letters 490: 143–53. https://pubmed.ncbi.nlm.nih.gov/32634449/ (March 30, 2021).

Cavieres, Viviana A. et al. 2020. "Human Golgi Phosphoprotein 3 Is an Effector of RAB1A and RAB1B." PLoS ONE 15(8 August). https://pubmed.ncbi.nlm.nih.gov/32790781/ (March 30, 2021).

Chin, D. J. et al. 1982. "Appearance of Crystalloid Endoplasmic Reticulum in Compactin-Resistant Chinese Hamster Cells with a 500-Fold Increase in 3-Hydroxy-3-Methylglutaryl-Coenzyme A Reductase." Proceedings of the National Academy of Sciences of the United States of America 79(4 I): 1185–89. https://pubmed.ncbi.nlm.nih.gov/6951166/ (March 30, 2021).

Chou, Kai Ming, A. Paul Krapcho, and Miles P. Hacker. 2001. "Impact of the Basic Amine on the Biological Activity and Intracellular Distribution of an Aza-Anthrapyrazole: BBR 3422." Biochemical Pharmacology 62(10): 1337–43. https://pubmed.ncbi.nlm.nih.gov/11709193/ (March 30, 2021).

Collins, Galen A., and Alfred L. Goldberg. 2020. "Proteins Containing Ubiquitin-like (Ubl) Domains Not Only Bind to 26S Proteasomes but Also Induce Their Activation." Proceedings of the National Academy of Sciences of the United States of America 117(9): 4664–74. https://pubmed.ncbi.nlm.nih.gov/32071216/ (March 30, 2021).

Dasari, Shaloam, and Paul Bernard Tchounwou. 2014. "Cisplatin in Cancer Therapy: Molecular Mechanisms of Action." European Journal of Pharmacology 740: 364–78. https://pubmed.ncbi.nlm.nih.gov/25058905/ (March 30, 2021).

Demishtein, Alik et al. 2017. "SQSTM1/P62-Mediated Autophagy Compensates for Loss of Proteasome Polyubiquitin Recruiting Capacity." Autophagy 13(10): 1697–1708. https://pubmed.ncbi.nlm.nih.gov/28792301/ (March 30, 2021).

De Duve, Christian et al. 1974. "Lysosomotropic Agents." Biochemical Pharmacology 23(18). https://pubmed.ncbi.nlm.nih.gov/4606365/ (March 30, 2021).

Feldman, Dorothy, Richard L. Swarm, and Janet Becker. 1981. "Ultrastructural Study of Rat Liver and Liver Neoplasms after Long-Term Treatment with Phenobarbital." Cancer Research.

Fregno, Ilaria et al. 2018. " ER -to-lysosome-associated Degradation of Proteasome-
resistant ATZ Polymers Occurs via Receptor-mediated Vesicular Transport ." *The*
EMBO Journal 37(17). https://pubmed.ncbi.nlm.nih.gov/30076131/ (March 30, 2021).
Fregno, Ilaria, and Maurizio Molinari. 2019. "Proteasomal and Lysosomal Clearance of
Faulty Secretory Proteins: ER-Associated Degradation (ERAD) and ER-to-Lysosome-
Associated Degradation (ERLAD) Pathways." *Critical Reviews in Biochemistry and*
Molecular Biology 54(2): 153–63. https://pubmed.ncbi.nlm.nih.gov/31084437/ (March
30, 2021).
Friedman, Jonathan R. et al. 2013. "Endoplasmic Reticulum-Endosome Contact Increases
as Endosomes Traffic and Mature." *Molecular Biology of the Cell* 24(7): 1030–40.
https://pubmed.ncbi.nlm.nih.gov/23389631/ (March 30, 2021).
Golebiowski, Filip et al. 2009. "System-Wide Changes to Sumo Modifications in Response to
Heat Shock." *Science Signaling* 2(72). https://pubmed.ncbi.nlm.nih.gov/19471022/
(March 30, 2021).
González, Alexis E. et al. 2017. "Autophagosomes Cooperate in the Degradation of
Intracellular C-Terminal Fragments of the Amyloid Precursor Protein via the
MVB/Lysosomal Pathway." *FASEB Journal* 31(6): 2446–59.
https://pubmed.ncbi.nlm.nih.gov/28254759/ (June 23, 2020).
Griffiths, Gareth et al. 1988. "The Mannose 6-Phosphate Receptor and the Biogenesis of
Lysosomes." *Cell* 52(3): 329–41. https://pubmed.ncbi.nlm.nih.gov/2964276/ (March 30,
2021).
Grumati, Paolo, Ivan Dikic, and Alexandra Stolz. 2018. "ER-Phagy at a Glance." *Journal of*
Cell Science 131(17). https://pubmed.ncbi.nlm.nih.gov/30177506/ (March 30, 2021).
Guarascio, Rosellina et al. 2020. "Negative Regulator of Ubiquitin-Like Protein 1 Modulates
the Autophagy-Lysosomal Pathway via P62 to Facilitate the Extracellular Release of
Tau Following Proteasome Impairment." *Human Molecular Genetics* 29(1): 80–96.
https://pubmed.ncbi.nlm.nih.gov/31691796/ (March 30, 2021).
Harding, Heather P., Yuhong Zhang, et al. 2000. "Perk Is Essential for Translational
Regulation and Cell Survival during the Unfolded Protein Response." *Molecular Cell*
5(5): 897–904. https://pubmed.ncbi.nlm.nih.gov/10882126/ (March 30, 2021).
Harding, Heather P., Isabel Novoa, et al. 2000. "Regulated Translation Initiation Controls
Stress-Induced Gene Expression in Mammalian Cells." *Molecular Cell* 6(5): 1099–1108.
https://pubmed.ncbi.nlm.nih.gov/11106749/ (March 30, 2021).
Hartmann-Petersen, R., and C. Gordon. 2004. "Ubiquitin-Proteasome System." *Cellular and*
Molecular Life Sciences 61(13). https://pubmed.ncbi.nlm.nih.gov/15224183/ (March 30,
2021).
Hartmann-Petersen, Rasmus, and Colin Gordon. 2004. "Integral UBL Domain Proteins: A
Family of Proteasome Interacting Proteins." *Seminars in Cell and Developmental*
Biology 15(2): 247–59. https://pubmed.ncbi.nlm.nih.gov/15209385/ (March 30, 2021).
Heifetz, Aaron, Roy W. Keenan, and Alan D. Elbein. 1979. "Mechanism of Action of
Tunicamycin on the UDP-GlcNAc:Dolichyl-Phosphate GlcNAc-1 -Phosphate
Transferase." *Biochemistry* 18(11): 2186–92.
https://pubs.acs.org/doi/abs/10.1021/bi00578a008 (April 27, 2021).
Henne, William Mike. 2017. "Discovery and Roles of ER-Endolysosomal Contact Sites in
Disease." *Advances in experimental medicine and biology* 997: 135–47.
https://pubmed.ncbi.nlm.nih.gov/28815527/ (March 30, 2021).
Huber, Michael D., and Larry Gerace. 2007. "The Size-Wise Nucleus: Nuclear Volume
Control in Eukaryotes." *Journal of Cell Biology* 179(4): 583–84.
https://pubmed.ncbi.nlm.nih.gov/17998404/ (March 30, 2021).
Jackson, Matthew P., and Eric W. Hewitt. 2016. "Cellular Proteostasis: Degradation of
Misfolded Proteins by Lysosomes." *Essays in Biochemistry* 60(2): 173–80.
https://pubmed.ncbi.nlm.nih.gov/27744333/ (March 30, 2021).
Jia, Rui, and Juan S. Bonifacino. 2019. "Negative Regulation of Autophagy by Uba6-Birc6–
Mediated Ubiquitination of Lc3." *eLife* 8. https://pubmed.ncbi.nlm.nih.gov/31692446/
(March 30, 2021).

Jin, Seok Min, and Richard J. Youle. 2012. "PINK1-and Parkin-Mediated Mitophagy at a
Glance." *Journal of Cell Science* 125(4): 795–99.
https://pubmed.ncbi.nlm.nih.gov/22448035/ (March 30, 2021).
Jung, Hye Seung et al. 2008. "Loss of Autophagy Diminishes Pancreatic β Cell Mass and
Function with Resultant Hyperglycemia." *Cell Metabolism* 8(4): 318–24.
https://pubmed.ncbi.nlm.nih.gov/18840362/ (April 26, 2021).
Khaminets, Aliaksandr et al. 2015. "Regulation of Endoplasmic Reticulum Turnover by
Selective Autophagy." *Nature* 522(7556): 354–58.
https://pubmed.ncbi.nlm.nih.gov/26040720/ (March 30, 2021).
Khaminets, Aliaksandr, Christian Behl, and Ivan Dikic. 2016. "Ubiquitin-Dependent And
Independent Signals In Selective Autophagy." *Trends in Cell Biology* 26(1): 6–16.
https://pubmed.ncbi.nlm.nih.gov/26437584/ (March 30, 2021).
Kim, Eunkyoung et al. 2018. "Dual Function of USP14 Deubiquitinase in Cellular
Proteasomal Activity and Autophagic Flux." *Cell Reports* 24(3): 732–43.
https://pubmed.ncbi.nlm.nih.gov/30021169/ (March 30, 2021).
Kim, Hyoung Tae, and Alfred L. Goldberg. 2018. "UBL Domain of Usp14 and Other Proteins
Stimulates Proteasome Activities and Protein Degradation in Cells." *Proceedings of the
National Academy of Sciences of the United States of America* 115(50): E11642–50.
https://pubmed.ncbi.nlm.nih.gov/30487212/ (March 30, 2021).
Kim, Tae Yeon, Eunmin Kim, Sungjoo Kim Yoon, and Jong Bok Yoon. 2008. "Herp
Enhances ER-Associated Protein Degradation by Recruiting Ubiquilins." *Biochemical
and Biophysical Research Communications* 369(2): 741–46.
https://pubmed.ncbi.nlm.nih.gov/18307982/ (March 30, 2021).
Kokame, K., K. L. Agarwal, H. Kato, and T. Miyata. 2000. "Herp, a New Ubiquitin-like
Membrane Protein Induced by Endoplasmic Reticulum Stress." *Journal of Biological
Chemistry* 275(42): 32846–53. https://pubmed.ncbi.nlm.nih.gov/10922362/ (June 21,
2020).
Kokame, Koichi, Hisao Kato, and Toshiyuki Miyata. 2001. "Identification of ERSE-II, a New
Cis-Acting Element Responsible for the ATF6-Dependent Mammalian Unfolded Protein
Response." *Journal of Biological Chemistry* 276(12): 9199–9205.
https://pubmed.ncbi.nlm.nih.gov/11112790/ (March 30, 2021).
Kondapalli, Chandana et al. 2012. "PINK1 Is Activated by Mitochondrial Membrane Potential
Depolarization and Stimulates Parkin E3 Ligase Activity by Phosphorylating Serine 65."
Open Biology 2(MAY). https://pubmed.ncbi.nlm.nih.gov/22724072/ (March 30, 2021).
Korolchuk, Viktor I., Fiona M. Menzies, and David C. Rubinsztein. 2010. "Mechanisms of
Cross-Talk between the Ubiquitin-Proteasome and Autophagy-Lysosome Systems."
FEBS Letters 584(7): 1393–98. https://pubmed.ncbi.nlm.nih.gov/20040365/ (March 30,
2021).
Koyano, Fumika et al. 2014. "Ubiquitin Is Phosphorylated by PINK1 to Activate Parkin."
Nature 510(7503): 162–66. https://pubmed.ncbi.nlm.nih.gov/24784582/ (June 24,
2020).
Van Der Laan, Sander W. et al. 2018. "From Lipid Locus to Drug Target through Human
Genomics." *Cardiovascular Research* 114(9): 1258–70.
https://pubmed.ncbi.nlm.nih.gov/29800275/ (March 30, 2021).
Lee, Crystal A., and Craig Blackstone. 2020. "ER Morphology and Endo-Lysosomal
Crosstalk: Functions and Disease Implications." *Biochimica et Biophysica Acta -
Molecular and Cell Biology of Lipids* 1865(1).
https://pubmed.ncbi.nlm.nih.gov/31678515/ (March 30, 2021).
Lee, Do Hee. 1998. "Proteasome Inhibitors: Valuable New Tools for Cell Biologists." *Trends
in Cell Biology* 8(10): 397–403. https://pubmed.ncbi.nlm.nih.gov/9789328/ (March 30,
2021).
Leitman, Julia et al. 2014. "Herp Coordinates Compartmentalization and Recruitment of
HRD1 and Misfolded Proteins for ERAD." *Molecular Biology of the Cell* 25(7): 1050–60.
https://pubmed.ncbi.nlm.nih.gov/24478453/ (June 23, 2020).
Liao, Yangjie et al. 2019. "Excessive ER-Phagy Mediated by the Autophagy Receptor

FAM134B Results in ER Stress, the Unfolded Protein Response, and Cell Death in HeLa Cells." *Journal of Biological Chemistry* 294(52): 20009–23. https://pubmed.ncbi.nlm.nih.gov/31748416/ (March 30, 2021).

Marzec, Michal, Davide Eletto, and Yair Argon. 2012. "GRP94: An HSP90-like Protein Specialized for Protein Folding and Quality Control in the Endoplasmic Reticulum." *Biochimica et Biophysica Acta - Molecular Cell Research* 1823(3): 774–87. https://pubmed.ncbi.nlm.nih.gov/22079671/ (March 31, 2021).

Miettinen, Teemu P., and Mikael Björklund. 2015. "Mevalonate Pathway Regulates Cell Size Homeostasis and Proteostasis through Autophagy." *Cell Reports* 13(11): 2610–20. https://pubmed.ncbi.nlm.nih.gov/26686643/ (March 30, 2021).

Miura, Hikari et al. 2010. "Deletion of Herp Facilitates Degradation of Cytosolic Proteins." *Genes to Cells* 15(8): 843–53. https://pubmed.ncbi.nlm.nih.gov/20604806/ (March 30, 2021).

Mizushima, Noboru, Beth Levine, Ana Maria Cuervo, and Daniel J. Klionsky. 2008. "Autophagy Fights Disease through Cellular Self-Digestion." *Nature* 451(7182): 1069–75. https://pubmed.ncbi.nlm.nih.gov/18305538/ (March 30, 2021).

Mizushima, Noboru, Tamotsu Yoshimori, and Yoshinori Ohsumi. 2011. "The Role of Atg Proteins in Autophagosome Formation." *Annual Review of Cell and Developmental Biology* 27: 107–32. https://pubmed.ncbi.nlm.nih.gov/21801009/ (March 30, 2021).

Mollazadeh, Hamid et al. 2018. "The Effect of Statin Therapy on Endoplasmic Reticulum Stress." *Pharmacological Research* 137: 150–58. https://pubmed.ncbi.nlm.nih.gov/30312664/ (March 30, 2021).

Moreau, Kevin, Shouqing Luo, and David C. Rubinsztein. 2010. "Cytoprotective Roles for Autophagy." *Current Opinion in Cell Biology* 22(2): 206–11. https://pubmed.ncbi.nlm.nih.gov/20045304/ (March 30, 2021).

Murrow, Lyndsay, and Jayanta Debnath. 2013. "Autophagy as a Stress-Response and Quality-Control Mechanism: Implications for Cell Injury and Human Disease." *Annual Review of Pathology: Mechanisms of Disease* 8: 105–37. https://pubmed.ncbi.nlm.nih.gov/23072311/ (March 30, 2021).

Neefjes, Jacques, Marlieke M.L. Jongsma, and Ilana Berlin. 2017. "Stop or Go? Endosome Positioning in the Establishment of Compartment Architecture, Dynamics, and Function." *Trends in Cell Biology* 27(8): 580–94. https://pubmed.ncbi.nlm.nih.gov/28363667/ (March 30, 2021).

Ou, W. J. et al. 1995. "Conformational Changes Induced in the Endoplasmic Reticulum Luminal Domain of Calnexin by Mg-ATP and Ca2+." *Journal of Biological Chemistry* 270(30): 18051–59. https://pubmed.ncbi.nlm.nih.gov/7629114/ (March 30, 2021).

Quiroga, Clara et al. 2013. "Herp Depletion Protects from Protein Aggregation by Up-Regulating Autophagy." *Biochimica et Biophysica Acta - Molecular Cell Research* 1833(12): 3295–3305. https://pubmed.ncbi.nlm.nih.gov/24120520/ (March 30, 2021).

Rothenberg, Cara et al. 2010. "Ubiquilin Functions in Autophagy and Is Degraded by Chaperone-Mediated Autophagy." *Human Molecular Genetics* 19(16): 3219–32. https://pubmed.ncbi.nlm.nih.gov/20529957/ (March 30, 2021).

Sai, Xiaorei et al. 2003. "The Ubiquitin-like Domain of Herp Is Involved in Herp Degradation, but Not Necessary for Its Enhancement of Amyloid β-Protein Generation." *FEBS Letters* 553(1–2): 151–56. https://pubmed.ncbi.nlm.nih.gov/14550564/ (March 30, 2021).

Sauvé, Véronique et al. 2018. "Mechanism of Parkin Activation by Phosphorylation." *Nature Structural and Molecular Biology* 25(7): 623–30. https://pubmed.ncbi.nlm.nih.gov/29967542/ (March 30, 2021).

Schindelin, Johannes et al. 2012. "Fiji: An Open-Source Platform for Biological-Image Analysis." *Nature Methods* 9(7): 676–82. https://pubmed.ncbi.nlm.nih.gov/22743772/ (March 30, 2021).

Schuck, Sebastian et al. 2009. "Membrane Expansion Alleviates Endoplasmic Reticulum Stress Independently of the Unfolded Protein Response." *Journal of Cell Biology* 187(4): 525–36. https://pubmed.ncbi.nlm.nih.gov/19948500/ (March 30, 2021).

Schulz, Jasmin et al. 2017. "Conserved Cytoplasmic Domains Promote Hrd1 Ubiquitin

Ligase Complex Formation for ER-Associated Degradation (ERAD)." *Journal of Cell Science* 130(19): 3322–35. https://pubmed.ncbi.nlm.nih.gov/28827405/ (March 30, 2021).

Schulze, Andrea et al. 2005. "The Ubiquitin-Domain Protein HERP Forms a Complex with Components of the Endoplasmic Reticulum Associated Degradation Pathway." *Journal of Molecular Biology* 354(5): 1021–27. https://pubmed.ncbi.nlm.nih.gov/16289116/ (March 30, 2021).

Sebti, Salwa et al. 2014. "BAT3 Modulates P300-Dependent Acetylation of P53 and Autophagy-Related Protein 7 (ATG7) during Autophagy." *Proceedings of the National Academy of Sciences of the United States of America* 111(11): 4115–20. https://pubmed.ncbi.nlm.nih.gov/24591579/ (March 30, 2021).

Şentürk, Mümine et al. 2019. "Ubiquilins Regulate Autophagic Flux through MTOR Signalling and Lysosomal Acidification." *Nature Cell Biology* 21(3): 384–96. https://pubmed.ncbi.nlm.nih.gov/30804504/ (March 30, 2021).

Shi, Xiaojuan et al. 2020. "Silencing UHRF1 Enhances Cell Autophagy to Prevent Articular Chondrocytes from Apoptosis in Osteoarthritis through PI3K/AKT/MTOR Signaling Pathway." *Biochemical and Biophysical Research Communications* 529(4): 1018–24. https://pubmed.ncbi.nlm.nih.gov/32819559/ (March 30, 2021).

Swaney, Danielle L, Ricard A Rodríguez-Mias, and Judit Villén. 2015. "Phosphorylation of Ubiquitin at Ser65 Affects Its Polymerization, Targets, and Proteome-wide Turnover." *EMBO reports* 16(9): 1131–44. https://pubmed.ncbi.nlm.nih.gov/26142280/ (June 24, 2020).

Tanida, Isei, Takashi Ueno, and Eiki Kominami. 2008. "LC3 and Autophagy." *Methods in Molecular Biology* 445: 77–88. https://pubmed.ncbi.nlm.nih.gov/18425443/ (March 30, 2021).

Tapia, Diego et al. 2019. "KDEL Receptor Regulates Secretion by Lysosome Relocation- and Autophagy-Dependent Modulation of Lipid-Droplet Turnover." *Nature Communications* 10(1). https://pubmed.ncbi.nlm.nih.gov/30760704/ (March 30, 2021).

Thastrup, O et al. 1990. "Thapsigargin, a Tumor Promoter, Discharges Intracellular Ca2+ Stores by Specific Inhibition of the Endoplasmic Reticulum Ca2(+)-ATPase." *Proceedings of the National Academy of Sciences* 87(7).

Thayer, Julia A. et al. 2020. "The PARK10 Gene USP24 Is a Negative Regulator of Autophagy and ULK1 Protein Stability." *Autophagy* 16(1): 140–53. https://pubmed.ncbi.nlm.nih.gov/30957634/ (March 30, 2021).

Toulmay, Alexandre, and William A. Prinz. 2011. "Lipid Transfer and Signaling at Organelle Contact Sites: The Tip of the Iceberg." *Current Opinion in Cell Biology* 23(4): 458–63. https://pubmed.ncbi.nlm.nih.gov/21555211/ (March 30, 2021).

Tsubuki, Satoshi et al. 1996. "Differential Inhibition of Calpain and Proteasome Activities by Peptidyl Aldehydes of Di-Leucine and Tri-Leucine." *Journal of Biochemistry* 119(3): 572–76. https://pubmed.ncbi.nlm.nih.gov/8830056/ (March 30, 2021).

Vabulas, Ramunas M., and F. Ulrich Hartl. 2005. "Cell Biology: Protein Synthesis upon Acute Nutrient Restriction Relies on Proteasome Function." *Science* 310(5756): 1960–63. https://pubmed.ncbi.nlm.nih.gov/16373576/ (March 30, 2021).

Valm, Alex M. et al. 2017. "Applying Systems-Level Spectral Imaging and Analysis to Reveal the Organelle Interactome." *Nature* 546(7656): 162–67. https://pubmed.ncbi.nlm.nih.gov/28538724/ (March 30, 2021).

VerPlank, Jordan J.S., Sudarsanareddy Lokireddy, Jinghui Zhao, and Alfred L. Goldberg. 2019. "26S Proteasomes Are Rapidly Activated by Diverse Hormones and Physiological States That Raise CAMP and Cause Rpn6 Phosphorylation." *Proceedings of the National Academy of Sciences of the United States of America* 116(10): 4228–37. https://pubmed.ncbi.nlm.nih.gov/30782827/ (March 30, 2021).

Walczak, Marta, and Sascha Martens. 2013. "Dissecting the Role of the Atg12-Atg5-Atg16 Complex during Autophagosome Formation." *Autophagy* 9(3): 424–25. https://pubmed.ncbi.nlm.nih.gov/23321721/ (March 30, 2021).

Wauer, Tobias et al. 2015. "Ubiquitin Ser65 Phosphorylation Affects Ubiquitin Structure,

Chain Assembly and Hydrolysis." *The EMBO Journal* 34(3): 307–25. https://pubmed.ncbi.nlm.nih.gov/25527291/ (March 30, 2021).

Wiest, D. L. et al. 1990. "Membrane Biogenesis during B Cell Differentiation: Most Endoplasmic Reticulum Proteins Are Expressed Coordinately." *Journal of Cell Biology* 110(5): 1501–11. https://pubmed.ncbi.nlm.nih.gov/2335560/ (March 30, 2021).

Wu, Yuping et al. 2018. "Alleviation of Endoplasmic Reticulum Stress Protects against Cisplatin-Induced Ovarian Damage." *Reproductive Biology and Endocrinology* 16(1). https://pubmed.ncbi.nlm.nih.gov/30176887/ (March 30, 2021).

Xu, Fang et al. 2021. "COPII Mitigates ER Stress by Promoting Formation of ER Whorls." *Cell Research* 31(2): 141–56. https://pubmed.ncbi.nlm.nih.gov/32989223/ (March 30, 2021).

Xu, Haoxing, and Dejian Ren. 2015. "Lysosomal Physiology." *Annual Review of Physiology* 77: 57–80. https://pubmed.ncbi.nlm.nih.gov/25668017/ (March 30, 2021).

Yan, Long et al. 2014. "Ube2g2-Gp78-Mediated HERP Polyubiquitylation Is Involved in ER Stress Recovery." Journal of Cell Science 127(7): 1417–27. https://pubmed.ncbi.nlm.nih.gov/24496447/ (June 21, 2020).

Yang, Ying, and Daniel J. Klionsky. 2020. "A Novel Role of UBQLNs (Ubiquilins) in Regulating Autophagy, MTOR Signaling and v-ATPase Function." *Autophagy* 16(1): 1–2. https://pubmed.ncbi.nlm.nih.gov/31516068/ (March 30, 2021).

Yin, Yili et al. 2012. "SUMO-Targeted Ubiquitin E3 Ligase RNF4 Is Required for the Response of Human Cells to DNA Damage." *Genes and Development* 26(11): 1196–1208. https://pubmed.ncbi.nlm.nih.gov/22661230/ (March 30, 2021).

Yoshida, Hiderou et al. 2001. "XBP1 MRNA Is Induced by ATF6 and Spliced by IRE1 in Response to ER Stress to Produce a Highly Active Transcription Factor." *Cell* 107(7): 881–91. https://pubmed.ncbi.nlm.nih.gov/11779464/ (March 30, 2021).

Yu, Houqing, and Andreas Matouschek. 2017. "Recognition of Client Proteins by the Proteasome." *Annual Review of Biophysics* 46: 149–73. https://pubmed.ncbi.nlm.nih.gov/28301771/ (March 30, 2021).

Zhang, Tao et al. 2018. "HMG-CoA Reductase Inhibitors Relieve Endoplasmic Reticulum Stress by Autophagy Inhibition in Rats with Permanent Brain Ischemia." *Frontiers in Neuroscience* 12(JUN). https://pubmed.ncbi.nlm.nih.gov/29970982/ (March 30, 2021).

Zhang, Youzhi et al. 2020. "Simvastatin Improves Lysosome Function via Enhancing Lysosome Biogenesis in Endothelial Cells." *Frontiers in Bioscience - Landmark*.

Zhu, K., K. Dunner, and D. J. McConkey. 2010. "Proteasome Inhibitors Activate Autophagy as a Cytoprotective Response in Human Prostate Cancer Cells." *Oncogene* 29(3): 451–62. https://pubmed.ncbi.nlm.nih.gov/19881538/ (March 30, 2021).

Figure 1

A

B

Figure 2

A

B

C

Figure 3

A

	FLAG	CALNEXIN	GRP94	Merge
HERPUD1-WT				
HERPUD1-ΔUBL				

B

HERPUD1-WT   HERPUD1-ΔUBL

GRP94 fluorescence intensity

Max

Min

C

ER Volume (μm³)

***

WT   ΔUBL
HeLa HERPUD1

D

HERPUD1-WT   HERPUD1-ΔUBL   HERPUD1-ΔUBL

Bar: 500 nm

Figure 4

A

	HeLa WT			HeLa HERPUD1	
pb	C	Tun	Tg	WT	ΔUBL
300 —						*uXBP1* / *sXBP1*
100 —						*CYC-A*
	1	2	3	4	5

B

HERPUD1 WT   HERPUD1 ΔUBL

kDa   0   2   4   6   0   2   4   6   h Tg 2 μM

70 —

50 —   XBP1

40 —   β-ACTIN

1   2   3   4   5   6   7   8

C

HeLa HERPUD1

XBP1 levels (Normalized units)

n.s.

***

**

0   6   0   6   h Tg 2μM

WT   ΔUBL

Figure 5

Figure 6

Figure 7

Figure 8

A

% Survival (y-axis, 0 to 150)
CDDP Concentration μM (x-axis: 1, 2, 4, 8, 16)

Legend:
● HERPUD1-WT
■ HERPUD1-S59D

n.s. * * * *

B

HERPUD1-WT - CDDP

0,36 %   4,90 %

Q1 Q2
Q3 Q4

93,8 %   0,91 %

HERPUD1-WT + CDDP

0,33 %   6,42 %

Q1 Q2
Q3 Q4

80,3 %   12,9 %

HERPUD1-S59D - CDDP

1,21 %   5,37 %

Q1 Q2
Q3 Q4

91,5 %   1,92 %

HERPUD1-S59D + CDDP

1,08 %   3,50 %

Q1 Q2
Q3 Q4

86,5 %   8,90 %

7AAD (y-axis)

Annexin V (x-axis)

Figure 9