DEVELOPMENT OF NOVEL MULTIPLEX PCR FOR RAPID DIAGNOSIS OF COINFECTED HEMO-PARASITES IN CATTLE

Pankaj Kumar[#], Abhay Kumar, Kamal Sarma, Paresh Sharma[1], Rashmi Rekha Kumari[2] and Manish Kumar[3]

ICAR Research Complex for Eastern Region, Patna

P.O. Bihar Veterinary College, Patna -800 011, INDIA

1. National Institute of Animal Biotechnology, Hyderabad -500 032, Telangana, India.

2. Department of Veterinary Pharmacology and Toxicology, Bihar Veterinary College, BASU, Patna-800 014, Bihar, India

3. Department of Biosciences and Bioengineering, Indian Institute of Technology Guwahati, Guwahati-781039, Assam, India

#Corresponding author

Pankaj Kumar, Division of Livestock and Fisheries Management, ICAR Research Complex for Eastern Region,

Patna-800 014, Bihar, India

 Email: Pankaj.Kumar1@icar.gov.in

Keywords: Novel, Multiplex PCR, Theileria, Babesia, Anaplasma, Cattle, Coinfection

ABSTRACT

A novel, rapid and specific multiplex polymerase chain reaction has been developed for the diagnosis of hemo-parasitic infection in bovine blood by three of the most common hemo-parasites. The reported method relied on the detection of the three different bovine hemoparasites isolated from red blood cells (RBCs) of cattle by conventional Giemsa stained blood smear (GSBS) and confirmed by multiplex PCR. The designed multiplex primer sets can amplify 205, 313 and 422 bp fragments of *apocytochrome b*, *sporozoite and macroschizont 2 (spm2)* and *16S rRNA* gene for *Babesia bigemina*, *Theileria annulata* and *Anaplasma marginale*, respectively. This multiplex PCR was sensitive with the ability to detect the presence of 150 ng of genomic DNA. The primers used in this multiplex PCR also showed highly specific amplification of specific gene fragments of each respective parasite DNA without the

presence of non-specific and non-target PCR products. This multiplex PCR system was used to diagnose GSBS

confirmed blood samples (N=12) found infected or co-infected with hemoparasites. A comparison of the two

detection methods revealed that 58.33% of specimens showed concordant diagnoses with both techniques. The

specificity, positive predictive value and kappa coefficient of agreement was highest for diagnosis of *B. bigemina*

and lowest for *A. marginale*. The overall Kappa coefficient for diagnosis based on GSBS for multiple pathogen

compared to multiplex PCR was 0.56 slightly behind the threshold of 0.6 of agreement. Therefore, confirmation

should always be made based on PCR to rule out false positive due to differences in subjective observations, stain

particles and false negative due to low level of parasitaemia. The simplicity and rapidity of this specific multiplex

PCR method make it suitable for large-scale epidemiological studies and for follow-up of drug treatments.

Introduction

Recent studies suggest that vector borne diseases has emerged due to climate change in tropical countries

(Balakrishnan 2017; El-Sayed and Kamel 2020; Kumar et al. 2021; Kumari et al. 2019). Tick population has

increased considerably and so is the tick borne infections in susceptible host. It was observed that cattle population

often suffer from coinfection of more than one pathogens transmitted by the ticks. These infections cannot be

diagnosed symptomatically as the symptoms are overlapping and elusive. Similarly microscopic examination based

on conventional blood smear examination often requires expertise observation and may sometimes be misleading in

co-infections or due to subjective observations. PCR based diagnosis is very specific and sensitive in detecting these

pathogens, however may require multiple PCR for diagnosis of pathogen load in samples suspected to have co-

infections. Considering these limitations of diagnosis of multiple tick transmitted diseases, an attempt has been made

to design a multiplex PCR to detect co-infections of these pathogens in a single run with the objective to provide

prompt and complete diagnosis to the farmers.

Materials and Methods

Ethics. The study involved drawing of 3 ml blood from jugular vein aseptically from clinically affected animal with

the consent of the owners. These were either brought for treatment to veterinary hospitals or as a call for treatment at

farmer's door by the local veterinarian. There are no specific ethical guidelines for a blood sample collection from

clinical cases, and hence no prior approval was mandatory. Treatment was provided to all animals sampled.

Animal. Blood samples were selectively collected from crossbred cattle (*Bos taurus* × *Bos indicus* breeds) (n=30) in

an EDTA vacutainer tube and a clot activator tube (BD, Franklin, USA) suspected to have infection of tick

transmitted diseases. Cattle were chosen from the farmer's field, dairy farms and veterinary hospitals around

endemic pockets of Manner block of the peri-urban Patna located in the Gangetic plains of Bihar, India. The

selection of cattle was made on the available clinical history and symptoms associated with tick infestation.

Clinical examination. Selected animals from the endemic area were clinically examined during sampling. Rectal

temperature for understanding the severity, mucosal membranes for status of anemia and pre-scapular lymph nodes

for any enlargement, respiration pattern was also observed. Eye was examined for any opacity or abnormal

discharge. Any other abnormality observed during clinical examination was also recorded.

Blood examination. The collected blood samples were processed for microscopic examination, hematological

examination, and whole blood genomic DNA isolation. For microscopic examination of blood and lymph-node

aspirate, a duplicate thin smear was prepared on a glass slide from each sample and fixed with methanol for 5 min.

The fixed smear on the glass slide was stained with Giemsa stain (Himedia) for 20 min, washed, air-dried, and

examined under oil (100x) immersion. The presence of piroplasms in blood smear through microscopic examination

was considered positive for *Theileria* spp. The hemoglobin (Hb) concentration (g/dL) in blood was measured using

HemoCue® cuvettes and its analyzer. PCV was calculated based on correlation between hemoglobin and PCV

expressed as Hb concentration (g/dL) = 0.3 PCV + 3 (Turkson and Ganyo 2015). RBC count and WBC counts were

measured manually under microscope by routine procedures (Jain 1999).

DNA Isolation. Infected blood samples (300 µL) found positive based on GSBS were used to extract whole genomic

DNA using a commercial kit (GCC) as per the manufacturer's instruction and stored at -20°C. The quality and

quantity of genomic DNA were checked by running the extracted DNA on agarose gel (1%), and the purity was

measured using nanodrop readings.

Primer Designing. The primer was designed for identification of three most common pathogens (*Theileria annulata,*

Anaplasma marginale and *Babesia bigemina*) in bovine based on our experience and infection prevailing in the

region (Table 1). The primers of all the three pathogens was designed to get nearly similar melting temperature

(Tm~57-59°C). Primer for *T. annulata* (313 bp) was designed using NCBI reference sequence XM_947453.1 for

2763 bp sequence of *T. annulata* Spm2 protein partial mRNA (Pain et al., 2005). The *sporozoite and macroschizont*

2 (*spm2*) gene sequence is reported to be specific to *T. annulata* parasitic infection (Prabhakaran et al. 2021; Tian et

al. 2018). Primer for simultaneous detection of *A. marginale* (422 bp) was designed using NCBI reference sequence

for 16S ribosomal RNA sequence of *A. marginale* str. Florida as Gene ID-7398331. The primer for the third

common pathogen *B. bigemina* (205 bp) was designed using NCBI reference nucleotide sequence AF109354.1 for

720 bp of *apocytochrome b* gene (Bilgiç et al. 2013).

Multiplex PCR. The thermal PCR program was standardized for multiplex PCR to detect the genomic DNA of three

haemo-parasite (*T. annulata, B. bigemina, and A. marginale*) in a single run using DNA extracted from cattle blood

samples. All isolated DNA samples from cattle blood that was found infected with single or multiple haemo-parasite

by GSBS was used to run the multiplex PCR.

Briefly, PCR assay was performed in a total volume of 25 μl containing 3 μl (50 ng/μl) of the genomic DNA and 2

μl primers each of forward and reverse primer prepared after mixing of equal volume of 10 micro-molar working

solution of the three forward and reverse primers and 12.5 μl of OneTaq Hot Start master mix with buffer (Imperial

Life Sciences (P) Limited), and ultra-pure nuclease free water to make the final volume. Amplification involved a

hot start of 5 min at 95˚C, followed by 35 cycles of 1 min at 95˚C, 1 min at 58˚C, 1 min at 72˚C, and a final

extension step of 10 min at 72˚C.

The amplified PCR products were subjected to electrophoresis on 2% agarose gel stained with ethidium bromide,

visualized under UV light, and photographed in trans-illuminator apparatus. The amplicons molecular sizes were

estimated by including a Quick-Load 100 bp DNA Ladder (Imperial Life Sciences Private Limited) and visualised

and documented on Gel-Doc (Azure Biosystems c200). DNA positive controls of *T. annulata* were kindly provided

by NAIB, Hyderabad, while positive control of *A. marginale* (MK834271) and *B. bigemina* (MH936010) was used

from DNA found positive after specific amplification and sequencing for species specific amplified products.

Distilled water served as negative control.

Statistics: Data were analysed for mean and standard error. Analysis of variance (ANOVA) using post hoc LSD test

of significance, if any, was observed at P<0.05. The observation of GSBS and Multiplex PCR was compared for

sensitivity (S_e), specificity (S_p), positive predictive values (PPV) and negative predictive value (NPV) by the method

elaborated earlier (Trevethan 2017) . Cohen's kappa coefficient was calculated by the method described (Landis and

Koch 1977).

Results

The study indicated that the endemic region had high (76.67%) infection of tick transmitted haemo-parasite

in selected crossbred cattle based on conventional Giemsa blood smear examination (Fig. 1). The infected animals

had clinical signs such as high fever, drop in milk yield, inappetance to anorexia, high percentage (30.44) of infected

cattle were co-infected with more than one haemo-parasite. The haematological changes observed in these cattle are

indicative of significant RBCs destruction, anaemia (reduced Hb level) and leukopenia in infected cattle compared

to non-infected cattle (Table 2). The study corroborated with earlier reports of these changes associated with tick

transmitted haemo-parasite in cattle (Ganguly et al. 2017; Larcombe et al. 2019; Prabhakaran et al. 2021) and

buffaloes (Kumar et al. 2021). Molecular confirmation based on PCR amplification requires multiple attempts to

detect co-infected samples. Multiplex PCR based on species specific primers was observed to be useful and 100 %

specific in detecting the co-infection of tick transmitted haemo-parasite in a single run (Table 3). The visualization

was clear and specific on gel doc examination (Fig. 2). Negative samples did not result in any amplification. Further,

the results of multiplex PCR considered as confirmatory was compared with results of GSBS. The unmatched results

of GSBS was considered as false positive and negative. The sensitivity (S_e), specificity (S_p), positive predictive

values (PPV), negative predictive values (NPV) and Cohen's kappa coefficient of the GSBS observation with

multiplex PCR are depicted in table 4. The S_p, PPV and kappa coefficient of agreement was highest for diagnosis of

B. bigemina and lowest for *A. marginale*. The S_e was highest for detection of *T. annulata* and lowest for *B.*

bigemina. The overall Kappa coefficient for diagnosis of these parasites by the both the test are in moderate (0.56)

agreement, though has high S_e (78.9), S_p (76.5), and PPV (78.9) percentage. This the multiplex PCR is able to detect

false negative samples based on GSBS due to low level of parasitaemia and also able to rectify false positive results

due to differences in subjective observations.

Discussion

Tick transmitted diseases with high infection in bovine population of Bihar with *Theileria* spp., *A*

marginale, and *B. bigemina* and their co-infection has been frequently reported (Kala et al. 2018; Kumar et al. 2019;

Kumar et al. 2021; Prabhakaran et al. 2021; Roy 2021). Our finding also suggest high rate of infection of these

hemo-parasite and association of more than one type parasite in the animals screened by GSBS. These results have

few limitation and dependent on expert examination and staining procedures (Prabhakaran et al. 2021). It was

observed that cattle infected with these parasites adversely affected haematological parameters. These changes are

associated with capability of infecting parasite to reach the circulation and invade the RBCs and corroborates with

earlier findings from mixed infected cattle of southern India (Jayalakshmi et al. 2019). Severity of changes varied

depending upon the type and degree of RBCs infected by these parasites. The confirmatory molecular diagnostic

method was developed to detect co-infected state in animal in single run by multiplex PCR using specific gene

primers which was found useful and corroborates with the findings of GSBS. *Spm2 gene* used for detection of *T.*

annulata has not been attempted earlier for multiplex PCR development and has been reported to be expressed both

in sporozoites and in later stages (macroschizont-infected leucocytes and piroplasms) of life-cycle of *T. annulata*

(Knight et al. 1998). *Apocytochrome b* gene has been reported to be very effective for detection of *B. bigemina* and

B. bovis by targeting a species-specific region of this gene (Ganzinelli et al. 2020). Similar methods of multiplex

PCR are also reported based on different target genes. Kundave et al., (2018) developed multiplex PCR using

Tams1, *18S rRNA* and *16S rRNA* genes of *T. annulata*, *B. bigemina* and *A. marginale*, respectively. Bilgiç et al.,

(2013) developed a multiplex PCR assay for simultaneous detection of *T. annulata*, *A. marginale* and *B. bovis* in

cattle using *cytochrome b* gene, *major surface protein–1β* encoding gene, and *VESA–1α* gene, respectively. The

results of S_e, S_p, PPV, NPV and Cohen's kappa coefficient of the GSBS observation with multiplex PCR indicates

that diagnosis of *A. marginale* is difficult with GSBS attributed to similarity of *A. marginale* with staining artifacts

and sometimes polymorphic attributes of *T. annulata*. On the contrary, diagnosis of *B. bigemina* had high S_p, PPV

which is attributed to the sampling of animals mostly after characteristic clinical symptoms of red urine. The Landis

and Koch (1977) have proposed the following as standards for strength of agreement for the kappa coefficient:

≤0=poor, 0.01–0.20=slight, 0.21–0.40=fair, 0.41–0.60=moderate, 0.61–0.80=substantial, and 0.81–1=almost perfect.

The Cohen's kappa coefficients indicates that diagnosis of *T. annulata* and *B. bigemina* are in substantial agreement

with the GSBS observation with obvious reason of sampling from animal with characteristic though overlapping

clinical symptoms.

Conclusion. A new method of multiplex PCR has been developed for simultaneous detection of *T. annulata, B.*

bigemina and *A. marginale* in cattle in a single run. This methods will be useful for researchers for diagnosis of co-

infection of tick transmitted diseases in cattle in endemic areas where possibilities of co-infection exists. Further

refinement in the methodology needs to be undertaken to diagnose other parasites like *T. orientalis*, *B. bovis, A.*

centrale, etc.

Conflict of Interest. There is no conflict to declare.

Author Contributions. PK, MK conceived the study, designed the primers, KS supervised the study PK, RRK

collected samples from field and initial screening. PK, PS and MK standardized the technique of multiplex PCR.

AK, PK, RRK and PS analysed the data statistically. PK, RRK, MK, PS and AK wrote the manuscript.

Acknowledgements. Authors acknowledge support and cooperation of Director, ICAR- Research Complex for

Eastern Region, Patna for providing necessary funds and facilities for undertaking this intra-mural project. Support

provided by project staff of ICAR Outreach program on Zoonotic Diseases is also duly acknowledged.

Funding: There is no external source of funding from any agency.

References.

Balakrishnan N (2017) Current scenario of tick-borne diseases in India-A review. Journal of Communicable
179   Diseases 49(2):6-13
180   Bilgiç HB, et al. (2013) Development of a multiplex PCR assay for simultaneous detection of Theileria annulata,
181   Babesia bovis and Anaplasma marginale in cattle. Experimental parasitology 133(2):222-229
182   El-Sayed A, Kamel M (2020) Climatic changes and their role in emergence and re-emergence of diseases.
183   Environmental science and pollution research international 27(18):22336–22352
184   doi:https://doi.org/10.1007/s11356-020-08896-w
185   Ganguly A, et al. (2017) Prevalence and haematobiochemical changes of tick borne haemoparasitic diseases in
186   crossbred cattle of Haryana, India. Indian Journal of Animal Sciences 87(5):552-557
187   Ganzinelli S, et al. (2020) Highly sensitive nested PCR and rapid immunochromatographic detection of Babesia
188   bovis and Babesia bigemina infection in a cattle herd with acute clinical and fatal cases in Argentina.
189   Transboundary and Emerging Diseases 67:159-164 doi:10.1111/tbed.13435
190   Jain NC (1999) Essentials of Veterinary Hematology. Lea and Febiger, Philadelphia
191   Jayalakshmi K, et al. (2019) Prevalence of haemoprotozoan diseases in cattle of Cauvery delta region of Tamil
192   Nadu. Journal of Parasitic Diseases 43(2):308–312 doi:https://doi.org/10.1007/s12639-019-01094-7
193   Kala SK, Deo BG, Kumari N (2018) Epidemiological aspects of theileriosis in cattle in and around Patna, Bihar,
194   India. International Journal of Current Microbiology Applied Sciences 7 1183-1191
195   Knight PA, et al. (1998) Identification of a Theileria annulata Antigen Expressed in Multiple Stages of the Parasite
196   Life Cycle. Experimental parasitology 90(1):110-121
197   Kumar B, Mondal D, Jithin M (2019) Evaluation of Reactive Oxidative Damage on Erythrocytic Cells Due to
198   Clinical Babesiosis in Lactating Cows. International Journal of Livestock Research 9(9):55-64
199   Kumar P, et al. (2021) Mixed infection of tick-borne haemo-parasites in water buffalo and associated pathological
200   responses and treatment. Indian Journal of Animal Research doi:10.18805/IJAR.B-4450
201   Kumari RR, Kumar R, Kumar P, Kumar M (2019) Emergence and Variations in Disease Ecology of Tick-Borne
202   Bovine Theileriosis in East India. International Journal of Livestock Research 9(11):12-25 doi:doi:
203   10.5455/ijlr.20190920061236
204   Landis JR, Koch GG (1977) An application of hierarchical kappa-type statistics in the assessment of majority
205   agreement among multiple observers. Biometrics 33(2):363-374 doi:doi:10.2307/2529786
206   Larcombe S, et al. (2019) The impact of tick-borne pathogen infection in Indian bovines is determined by host type
207   but not the genotype of Theileria annulata. Infection, Genetics and Evolution 75:103972
208   doi:https://doi.org/10.1016/j.meegid.2019.103972
209   Prabhakaran HS, Ghosh KK, Kumari RR, Kumar P, M. K (2021) Evaluation of sporozoite and macroschizont
210   antigen (Spm2) of Theileria annulata for its diagnostic potential. Ticks and Tick-Borne Diseases
211   14(4):101691 doi:https://doi.org/10.1016/j.ttbdis.2021.101691
212   Roy S, Bhandari, V., Barman, M., Kumar, P., Bhanot, V., Arora, J.S., Singh, S. and Sharma, P., (2021) Population
213   genetic analysis of the Theileria annulata parasites identified limited diversity and multiplicity of infection
214   in the vaccine from India. Frontiers in microbiology 11:3471
215   doi:https://doi.org/10.3389/fmicb.2020.579929

Tian Z, et al. (2018) Development of an indirect ELISA based on the recombinant Spm2 protein for detection of
tropical theileriosis. Acta Tropica 182:232-236
Trevethan R (2017) Sensitivity, Specificity, and Predictive Values: Foundations, Pliabilities, and Pitfalls in Research
and Practice. Frontiers in Public Health 5:307 doi:10.3389/fpubh.2017.00307
Turkson PK, Ganyo EY (2015) Relationship between haemoglobin concentration and packed cell volume in cattle
blood samples. Onderstepoort Journal of Veterinary Research 82(1):1-5

Table 1. List of primers designed from target genes for simultaneous diagnosis of *T. annulata, A. marginale* and *B.*
bigemina by multiplex PCR

Pathogens	Target Gene	Sequence (5′–3′)	Tm (°C)	Amplicon Size
T. annulata	*spm2*	TAAACCACAACATGTCCACAA	57.4	313bp
		AGTTACATATGGGTTTTCTTGTGTG	58.5	
B. bigemina	*apocytochrome b*	TTGGGCACTTCGTTATTTCC	59.9	205bp
		TGTTGCTCCCCAGTAACTCA	59.3	
A. marginale	*16S rRNA*	GGGGTAATGGCCTACCAAG	59.3	422bp
		CTACGAATTTCACCTCTACACTAGGA	59.3	

226 **Table 2.** Haematology of crossbred cattle diagnosed for tick transmitted pathogens based on GSBS

GSBS findings	Haemoglobin (g/L)	RBC count (×10^6/μL)	WBC count (×10^3/μL)	PCV (%)
T. annulata (n=7)	7.39[a]±0.43 (3.50-5.70)	4.20[a]±0.20 (1.64-3.24)	6.45[a]±0.45 (5.12-8.21)	14.71[a]±1.46 (9.0-21.0)
A. marginale (n=9)	7.69[a]±0.44 (5.8-9.6)	5.54[b]±0.28 (3.98-6.42)	5.99[a]±0.31 (4.89-7.12)	15.67[a]±1.50 (9.0-22.0)
Co-infections (n=7)	6.74[a]±0.37 (5.8-8.3)	4.54[a]±0.23 (3.88-5.62)	5.57[a]±0.31 (4.66-6.84)	12.43[a]±1.29 (9.00-18.0)
Negative (n=7)	10.94[b]±0.23 (9.9-11.8)	7.78[c]±0.34 (6.24-8.92)	7.85[b]±0.41 (6.38-9.45)	26.29[b]±0.75 (23.0-29.0)
Reference Range (Jain, 1993)	**80-150**	**5-10**	**4-12**	**24-46**

227 Values in column with different superscript varies significantly (P<0.05)

232 **Table 3.** Comparison of observation based on conventional GSBS examination and multiplex PCR developed

Sample No	GSBS			Multiplex PCR		
	A	T	B	A	T	B
1	+ve	-ve	-ve	+ve	+ve	-ve
2	+ve	+ve	-ve	+ve	+ve	-ve
3	+ve	+ve	-ve	+ve	+ve	-ve
4	+ve	+ve	-ve	+ve	+ve	-ve
5	-ve	+ve	-ve	+ve	+ve	-ve
6	+ve	-ve	-ve	-ve	-ve	+ve
7	-ve	+ve	+ve	-ve	+ve	+ve
8	-ve	+ve	+ve	-ve	+ve	+ve
9	+ve	-ve	+ve	-ve	-ve	+ve
10	-ve	+ve	-ve	-ve	-ve	+ve
11	+ve	+ve	-ve	-ve	+ve	-ve
12	+ve	-ve	-ve	+ve	-ve	-ve

233   Where, A= *A. marginale*; B= *B. bigemina*; T= *T. annulata*; +ve: Positive; -ve: Negative

235   **Table 4.** Comparison of sensitivity (S_e), specificity (S_p), positive predictive values (PPV), negative predictive values

236   (NPV) and Cohen's kappa coefficient of the GSBS observation with multiplex PCR

Pathogens	S_e (%)	S_p (%)	PPV (%)	NPV (%)	Kappa Coefficient
A	83	50	62.5	75	0.34
T	87.5	75	87.5	75	0.62
B	60	100	100	22.2	0.63
A+T+B	78.9	76.5	78.9	76.5	0.56

237   Where, A= *A. marginale*; B= *B. bigemina*; T= *T. annulata***;** Kappa coefficient of 0.6 and above is considered as in

238   agreement

Figure 1

A

C

Figure 1. **Diagnosis based on conventional GSBS.** Representative photographs were taken under oil immersion microscope 1000x magnification. **A**. Infected RBCs (Arrow) with *T. annulata* prioplasm; **B**. Infected RBCs with A. marginale (arrow); C. Infected RBCs with *B. bigemina* (paired pear-shaped bodies) and *T. annulata* (above arrow)

Figure 2

Figure 2. Multiplex PCR product agarose gel electrophoresis visualized under UV light of Gel Doc with positive amplification at 205, 313 and 422 bp for *B. bigemina, T. annulata* and *A. marginale*, respectively. **A.** Lane 1: Positive control; Lane 5: 100 bp Ladder; Remaining lanes: Samples 1-8; **B**. Lane 1:100 bp ladder; Lane 2: Negative control; Remaining lanes: Samples 9- 12