Here be dragons: important spatial uncertainty driven by climate data in

forecasted distribution of an endangered insular reptile

Heading title: High uncertainty driven by climate data in SDMs

Nicolas Dubos[1,2], Stephane Augros[3], Gregory Deso[4], Jean-Michel Probst[5], Jean-Cyrille Notter[6], Markus A.

Roesch[7]

1 INRAE (UMR TETIS), Maison de la télédétection, 500 rue Jean-François Breton 34093 Montpellier

Cedex 5, France.

2 Centre d'Ecologie et des Sciences de la Conservation (CESCO UMR 7204), Sorbonne Université, MNHN,

55 rue Buffon, 75005 Paris, France.

3 Eco-Med Océan Indien, 97400 Saint-Denis, France.

4 Association Herpétologique de Provence Alpes Méditerranée, Maison des Associations, 384 route de

Caderousse, F-84100 Orange, France.

5 Association Nature and Patrimoine, 1 rue des amarantes résidence Valeriane 2, Bat C appartement 15.

97490 Sainte Clotilde, La Réunion, France.

6 Parc National de La Réunion, 258 rue de La République, 97431, La Plaine Des Palmistes, La Réunion,

France.

7 Nature Océan Indien, 46 Rue des Mascarins, 97429 Petite-Ile, La Réunion, France

Abstract

The effect of future climate change is poorly documented in the tropics, especially in mountainous areas. Yet,

species living in these environments are predicted to be strongly affected. Newly available high-resolution

environmental data and statistical methods enable the development of forecasting models. Nevertheless, the

uncertainty related to climate models can be strong, which can lead to ineffective conservation actions.

Predicted studies aimed at providing conservation guidelines often account for a range of future climate

predictions (climate scenarios and global circulation models). However, very few studies considered

potential differences related to baseline climate data and/or did not account for spatial information (overlap)

in uncertainty assessments. We modelled the environmental suitability for *Phelsuma borbonica*, an

endangered reptile native to Reunion Island. Using two metrics of species range change (difference in overall

suitability and spatial overlap), we quantified the uncertainty related to the modelling technique (n = 10),

sample bias correction, climate change scenario, global circulation models (GCM) and baseline climate

(CHELSA *versus* Worldclim). Uncertainty was mainly driven by GCMs when considering overall suitability,

while for spatial overlap the uncertainty related to baseline climate became more important than that of

GCMs. The uncertainty driven by sample bias correction and variable selection was much higher when

assessed based on spatial overlap. The modelling technique was a strong driver of uncertainty in both cases.

We eventually provide a consensus ensemble prediction map of the environmental suitability of *P. borbonica*

to identify the areas predicted to be the most suitable in the future with the highest certainty. Predictive

studies aimed at identifying priority areas for conservation in the face of climate change need to account for

a wide panel of modelling techniques, GCMs and baseline climate data. We recommend the use of multiple

approaches, including spatial overlap, when assessing uncertainty in species distribution models.

Keywords: Baseline climate, CHELSA, *Phelsuma borbonica*, Reunion Island, Schoener's D overlap, Species

distribution models, Uncertainty, Worldclim

Introduction

Predictive studies of climate change effects are poorly documented in the tropics, especially in rare and endangered species (Pearson et al., 2014). Yet, tropical species are predicted to be more severely impacted by climate change than temperate species because they live closer to their thermal limit (Tewksbury, Huey, & Deutsch, 2008; Jiguet et al., 2010; Dubos et al., 2019). This is particularly true for species with a narrow distribution, a narrow niche, or living in highly heterogeneous environments such as mountainous areas (Raxworthy et al., 2008; Freeman & Class Freeman, 2014; Platts et al., 2014; Tang et al., 2018; Ahmadi et al., 2019; Cordier et al., 2020; Manes et al., 2021). Those species are difficult to model mostly because of data scarcity and statistical constrains related to low sample sizes and the lack of spatial coverage in environmental gradients. The recent availability of high-resolution environmental data along with methodological advances help filling this gap of knowledge (e.g., Dubos et al., 2021; Lannuzel et al., 2021). Forecasting models are useful to anticipate conservation management in the face of climate change. Specifically, species distribution models (SDMs) can be used to identify priority areas for protection (Leroy et al., 2014; Lannuzel et al., 2021), suitable conditions for habitat restoration and species (re)introductions and translocations (Minteer & Collins, 2010; Wilson, Roberts, & Reid, 2011; Adhikari, Barik, & Upadhaya, 2012; Draper, Marques, & Iriondo, 2019; Bellis et al., 2020; Butt et al., 2020; Westwood et al., 2020; Zhong et al., 2021).

A range of climate data is available for species distribution models, including several baseline climate data, some of which are used to predict climate change (e.g., Worldclim and CHELSA; Fick & Hijmans, 2017; Karger et al., 2017). Future climate predictions include multiple emission gas scenarios (Shared Socio-economic Pathways, SPP; as known as Representative Concentration Pathways, RCP) and global circulation models (GCMs). Both gas emission scenarios and GCMs can produce highly heterogeneous results in terms of predicted future distributions (Buisson et al., 2010; Baker et al., 2016). Most predictive studies included a range of scenarios and GCMs, but very few have considered potential uncertainties related to baseline climate data (Baker et al., 2016; Morales-Barbero & Vega-Álvarez, 2019; Datta, Schweiger, & Kühn, 2020; Ocon, 2020; Dubos, et al., 2021). This can induce a risk of misidentification of suitable environments, which can affect conservation prioritisation (Kujala et al., 2013; Baker et al., 2016; Muscatello, Elith, & Kujala, 2020) and subsequently lead to ineffective conservation actions (Converse & Sipe, 2021). Despite recent studies emphasising the need to account for multiple baseline climates (Morales-Barbero & Vega-Álvarez,

2019; Datta, Schweiger, & Kühn, 2020; Ocon, 2020), most studies aimed at providing conservation

guidelines do not assess the uncertainty related to bioclimatic input data.

Uncertainty assessments in SDMs can be made by comparing a range of modalities in model settings or data.

These comparisons can be based on differences in performance metrics such as the area under the receiving

operating characteristic curve (AUC) or the true skill statistic (TSS; Tessarolo et al., 2021). However, such

performance comparisons require independent data to be reliable. In absence of independent data,

uncertainty can be assessed based on differences in predicted species range change between model input

parameters, climate scenarios or climate data (e.g., Kujala et al., 2013; Baker et al., 2016; Muscatello, Elith,

& Kujala, 2020). Species range change can be estimated using the difference between current and future

summed suitability. This approach enables to identify which parameter drives the highest uncertainty in

terms of overall suitability across a region. However, it ignores the spatial component in suitability

differences. For instance, a difference in shift in distribution with no change in overall suitability would

remain undetected. Another approach is to discount the variation between model replicates from mean

suitability scores (Kujala et al., 2013). This method is highly relevant for conservation purposes since it

provides a spatially explicit map of the most consistently identified suitable areas but prevents from

disentangling the sources of uncertainty. A possible approach to assess each source of uncertainty while

accounting for spatial information is the use of overlap metrics such as Pearson's coefficient or similarity

indices (Muscatello, Elith, & Kujala, 2020; Dubos, Montfort, et al., 2021). To date, no study has tested for

potential differences in uncertainty assessments between these approaches.

In our study we generate a set of SDMs for the two subspecies of the Reunion Island day gecko, *Phelsuma*

borbonica borbonica and *P. borbonica mater* (subsequently referred to as *P. borbonica*), and test potential

differences in uncertainty assessments between the different approaches. Reunion Island is located in the

western Indian Ocean and has faced a number of alterations over the last century related to agricultural

practices (mostly sugarcane cultivation) and invasive species (Strasberg et al., 2005; Dubos, 2013; Fenouillas

et al., 2021; Irl et al., 2021). Its herpetofauna has already been strongly modified after multiple local

extinctions events and the arrival of allochthonous species (Cheke, 1987; Cheke & Hume, 2010; Sanchez &

Probst, 2016). Today only two native reptile species remain, the indigene *P. brobonica* and the endemic *P.*

inexpectata, while five reptile species having faced extinction in the recent past (Cheke & Hume, 2010).

Climate change can facilitate biological invasions and alters habitat suitability for locally adapted species

(Mainka & Howard, 2010; Gillard et al., 2017). To date, very few studies have focused on the potential

effects of climate change on the future of Reunion Island's biodiversity (but see Legrand et al., 2016). Dubos

et al. (2021) predicted the potential extinction of the endemic Reunion Island reptile, *P. inexpectata*, driven

by climate change.

The Reunion Island day gecko, *P. borbonica* is classified nationally as endangered due to its narrow

distribution. With high rates of land use change in Reunion Island, fire hazards and high pressure related to

introduced species (Macdonald & Cedex, 1991; Strasberg et al., 2005; Lagabrielle et al., 2009; Sanchez &

Probst, 2016), many populations are isolated and are exposed to local extinctions. This calls the urgent need

to identify priority areas for habitat conservation and restoration.

We assess for the first time the climatic niche of *P. borbonica* and the potential effects of climate change on

its future distribution. We quantify the uncertainty in species range change (SRC) driven by the statistical

methods (modelling technique, sample bias correction, variable selection) and climate data (RCPs, GCMs

and baseline data) using two approaches (difference in summed suitability and spatial overlap) in the

predicted future distribution of *P. borbonica*. We eventually provide a consensual map accounting for

uncertainty to guide conservation actions.

Methods

Occurrence data

Phelsuma borbonica is mostly distributed in the forested areas of the eastern, southern, and northern parts of

Reunion Island, from sea level to 2800 m (Meier, 1995; Mickaël Sanchez & Probst, 2017). Most

observations are made in intermediate altitudes, near trails, on artificial structures or at the edge of pristine or

disturbed forested areas (Augros et al., 2017). We obtained 5922 occurrence data (2648 after removing

redundancies) from the Système d'information de l'inventaire du patrimoine naturel (SINP) provided by the

Direction de l'Environnement, de l'Aménagement et du Logement (DEAL) and the Parc National de la

Réunion (PNRun) and some additional unpublished occurrence data from various sources (Biotope Océan

Indien, Cynorkis, Eco-Med Océan Indien, Nature Océan Indien). We removed 171 observations

corresponding to single observations with no evidence of persistence (presumably corresponding to

individuals transported out of their native range; Deso, 2001), translocated individuals, inaccurate

coordinates, or indices of past presence (e.g., subfossil clutches). To limit spatial autocorrelation, we

resampled one occurrence per occupied pixel at the resolution of the environmental data (3 arc seconds,

approximately 100m). The final sample included 1050 points when sampled from the Worldclim data, 1056

with CHELSA.

Land use data

We accounted for the habitat requirements of our model species by applying a filter to the climate data. We

obtained very high-resolution land cover categories (Urban, agricultural, natural, water) at 1.5m resolution

(resampled at 100m for computing purposes) derived from remote sensing (Dupuy, Gaetano, & Le Mézo,

2020; Fig. Sx). We removed agricultural and urbanised areas, because *P. borbonica* is only found in natural

and semi natural habitats, which enabled to minimise model complexity while remaining biologically

realistic and relevant for conservation applications.

Climate data

We used 19 bioclimatic variables for 30 arc sec (approximately 900m) resolution of the current climate data

and of the 2070 projections from CHELSA (Karger et al., 2017) and from Worldclim global climate data

(Fick & Hijmans, 2017). We removed isothermality (bio2) from the analysis because of a lack of variability

in Reunion Island. We decided to include all the remaining variables because both temperature and

precipitation are related to the species' biology, including those referring to indices of variability. We used

three Global Circulation Models (GCMs; i.e., BCC-CSM1-1, MIROC5 and HadGEM2-AO) and two

greenhouse gas emission scenarios (the most optimistic RCP26 and the most pessimistic RCP85; Fig. S3,

S4). To enable the filtering of unsuitable habitats, we downscaled the climate data to fit the resolution of the

land cover variables (3 arc sec) using bilinear interpolation.

Distribution modelling

We modelled and projected species distributions in R (version 4.0.3; R Core Team, 2020) with the Biomod2

R package (Thuiller et al., 2009), using 10 modelling techniques: generalised linear and generalised additive

models (GLM and GAM; Guisan, Edwards, & Hastie, 2002), classification tree analysis (CTA; Prasad,

Iverson, & Liaw, 2006), artificial neural network (ANN; Manel, Dias, & Ormerod, 1999), surface range

envelop (SRE, also known as BIOCLIM; Booth et al., 2014), flexible discriminant analysis (FDA; Manel,

Dias, & Ormerod, 1999), random forest (RF; Prasad, Iverson, & Liaw, 2006), multiple adaptive regression

splines (MARS; Leathwick et al., 2005), generalised boosting model (GBM; Elith, Leathwick, & Hastie,

2008) and maximum entropy (MaxEnt; Phillips & Schapire, 2006). We generated five different sets of

1050/1056 pseudo-absences (to equal the number of presences) and ran a first set of models using a random

pseudo-absence selection (Wisz & Guisan, 2009).

To accounting for sample bias, we reperformed all calculations using a non-random pseudo-absence

selection. Following Phillips et al. (2009), we produced five sets of pseudo-absences selected around the

presence points to reproduce the spatial bias of the sample. We used a geographic null model generated with

the dismo R package (Hijmans, 2012) and used it as a probability weight for pseudo-absence selection. Since

no independent data is available to assess the effect of sample bias correction, we used the relative overlap

index (ROI; Dubos, Préau, et al., 2021) based on Schoener's D overlap. The ROI enables to assess whether

the effect of correction is significant compared to the variability between model runs. It computes (1) the

mean overlap between the uncorrected and the corrected predictions (i.e., the absolute effect of correction),

and (2) the overlap between every pair of model replicate (between each pseudo-absence and cross

validation runs, individually for each modelling technique, i.e. model stochasticity). We computed the ROI

as follows:

$$ROI = \frac{\bar{D}_0 - \bar{D}(p_x, p_y)}{\bar{D}_0}$$

Where $\bar{D}_0$ is the mean overlap between model runs of the corrected group and $\bar{D}(p_x, p_y)$ is the mean

overlap between runs of the uncorrected and the corrected models. A value close to 0 represents a perfect

match between predictions (i.e. no effect of sample bias correction). The overlaps between uncorrected and

corrected groups tend to be significantly smaller than the overlaps between runs when the ROI approaches 1

(i.e. strong effect of sample bias correction).

For each individual baseline climate data, we selected one variable per group of inter-correlated variables to

avoid collinearity (Pearson's r > 0.7, Dormann et al., 2013) and assessed the relative importance of each

variable kept with 10 permutations per model replicate (total = 500 for each baseline data). The variables

included in the final models were those with a relative importance > 0.2 across at least 50% of model runs.

To determine whether variation in future predictions was driven by baseline climate *per se* or by differences

in the selected variables, we swapped the selected variables between baseline climates and repeated the

whole process.

Model evaluation−We spatially partitioned the data into five folds, with three runs of block cross-validation

(i.e., k-fold cross-validation; Fig. S1). We assessed model performance using the Boyce index, assumed to be

the best evaluation metric with pseudo-absence data (Leroy et al., 2018). A value of 1 means the models

reliably predicted the presence points while a value of 0 means that models did not perform better than

random. For ensemble models, we excluded models for which the Boyce index was below 0.5 (Gillard et al.,

2017). We verified that models are well informed for predictions on novel (future) data using clamping

masks.

Uncertainty analysis−We assessed the uncertainty in species range change (SRC) related to the modelling

technique, sample bias correction, climate scenarios, GCMs, baseline climate and variable selection. We

quantified SRC following two approaches. Firstly, we computed SRC as the difference between the summed

suitability scores of the baseline climate and the future predictions. Then we used linear models (LM,

assuming Gaussian errors), with SRC as a response variable, and the aforementioned sources of uncertainty

as explanatory variables, following Baker et al. (2016). We then assessed the proportion of deviance

explained by each source of uncertainty *f* as follows:

$$Pf = \frac{Df - D0}{D1}$$

where, Pf = proportion of deviance explained by factor f, $D1$ = deviance of full model, Df = deviance of full

model minus factor f, and $D0$ = deviance of null model.

Secondly, we quantified SRC using an overlap metric. We computed the Schoener's D overlap between

baseline projections and future projections using the ENMTools R package (Warren, Glor, & Turelli, 2010).

We eventually repeated the assessment of the proportion of deviance explained by the five aforementioned

sources of uncertainty using beta-regression GLM instead of LM, since overlap measures range continuously

between 0 and 1 (glmmTMB R package; Brooks et al., 2019).

In total, we computed 2000 models (10 modelling techniques * 5 pseudo-absence runs * 5 block cross-

validation runs * 2 modalities of sample bias correction * 2 baseline climate data * 2 modalities for variable

selection) for the current distribution, and 12000 projections on future climate data (2000 models * 3 GCMs

* 2 emission scenarios). To provide the most certain conservation guidelines, we built a consensus map of

the mean predictions across all simulations (of the corrected group and with the original selected variables)

after removing poorly performing models, discounted with the standard deviation (Kujala et al., 2013).

Results

Current distribution

Models predicted well the presence of *P. borbonica* (median Boyce index (Worldclim) = 0.81; median Boyce

index (CHELSA) = 0.78; Fig. S2, S3). A few points, corresponding to isolated populations, fell into areas

predicted as moderately suitable. Sample bias correction slightly increased the predictions of suitable areas,

but its effect on projections was not significant compared to within-model variation (ROI = 0.08; Fig. S4).

With the Worldclim baseline climate, the selected variables were mean annual temperature (Bio1) and winter

precipitation (Bio19; Fig. 1). With the CHELSA baseline climate, the selected variables were mean annual

temperature (Bio1) and summer precipitation (Bio18; Fig 2).

Fig. 1 Response of *Phelsuma borbonica* to the selected Worldclim bioclimatic variables. The black lines

represent the individual response curves for each iteration. The blue line represents the smoothed response

across iterations. Bio1: annual mean temperature (°C); Bio19: precipitation of coldest quarter (mm).

Fig. 2 Response of *Phelsuma borbonica* to the selected CHELSA bioclimatic variables. The black lines

represent the individual response curves for each iteration. The blue line represents the smoothed response

across iterations. Bio1: annual mean temperature (°C ×10); Bio18: precipitation of warmest quarter (mm).

Future distribution

Predictions differed strongly between modalities. Predictions derived from Worldclim baseline data indicated

a dramatic decline in climate suitability across the entire island, regardless of the scenario or GCM (Fig. 3).

The best predicted areas would shift upslope, but remain largely suboptimal. When based on CHELSA

baseline data, predictions were more optimistic but highly variable between GCMs (Fig. 4). The 'BCC-

CSM-1-1' GCM predicted little effect of climate change. The 'MIROC5' GCM indicated a slight decrease in

climate suitability in the north-western part of the current distribution area, while the HadGEM2-AO

predicted a strong decline in the eastern part.

Fig. 3 Current (a) and future (b) environmental suitability of *Phelsuma borbonica* in Reunion Island based

on Worldclim baseline climate data.

Fig. 4 Current (a) and future (b) environmental suitability of *Phelsuma borbonica* in Reunion Island based

on CHELSA baseline climate data.

Effect of variable selection

After swapping the selected variables between baseline climate data, model performances decreased only

slightly (median Boyce index (Worldclim) = 0.71; median Boyce index (CHELSA) = 0.70). The predicted

current distribution was overestimated with Worldclim, and slightly underestimated with CHELSA.

The differences between Worldclim and CHELSA in predictions of future climate suitability persisted.

Predictions based on Worldclim strongly differed between GCMs (Fig. 5) but were consistent when based on

CHELSA (Fig. 6).

Fig. 5 Current (a) and future (b) environmental suitability of *Phelsuma borbonica* in Reunion Island based on Worldclim baseline climate data using the selected variables from the CHELSA models.

Fig. 6 Current (a) and future (b) environmental suitability of *Phelsuma borbonica* in Reunion Island based on CHELSA baseline climate data using the selected variables from the Worldclim models.

Uncertainty analysis

The highest source of uncertainty differed according to the metric of species range change considered (Fig. 7). Overall change in suitability scores were the most variable between the modelling technique (SDM), followed by the GCM and the baseline climate. Uncertainty related to sample bias correction and variable selection were of lower magnitude. Regarding the amount of spatial information shared between current and future predictions (overlap), the highest source of uncertainty was the modelling technique, followed by variable selection and baseline climate. The uncertainty related to the GCM was lower than that of baseline climate when considering overlaps. Sample bias correction was the lowest source of uncertainty. The consensus maps indicate that the areas with the most consistent suitable conditions will be located at several places throughout the island (Fig. 8). The mean prediction maps also indicates that the best conditions will be largely suboptimal, from a maximum suitability score of 797 under current conditions to 362 by 2070.

Fig. 7 Proportion of deviance explained by five sources of uncertainty, quantified with two metrics of species

range change (SRC). SDM: modelling technique (species distribution model); GCM: global circulation

model; Correction: sample bias correction (non-random pseudo-absence selection); Baseline: baseline

climate data (Worldclim versus CHELSA); Overlap: Schoener's D overlap between current and future

projections; Suitability scores: difference between summed suitability scores of current and future

predictions.

Fig. 8 Consensus maps of current (a) and 2070 (b) priority areas for conservation for *Phelsuma borbonica*,

derived from predicted climate suitability. Current and future predictions (bottom) are the result of mean

projections (top left of each panel) discounted with inter-model variability (top right of each panel).

Discussion

We predicted the future climate suitability of *P. borbonica* while accounting for multiple climate models and

found a strong disagreement between future predictions derived from Worldclim and CHELSA baseline

climate data. The uncertainty related to both methodological aspects and input climate data depended on the

approach used to quantify species range change (summed suitability *versus* overlap). Mean projections

indicate an important decline in climate suitability by 2070.

Drivers of species distribution

The effect of temperature is consistent between simulations, with an optimal suitability estimated around

20°C on average throughout the year (mostly between 15 and 25°C, with a second peak near 10°C). The

identified temperature window corresponds to the overall thermal tolerance of tropical reptiles (Sunday,

Bates, & Dulvy, 2011).

Phelsuma borbonica is more present in areas with high precipitation, mostly corresponding to native

remnant rainforests. We presume that *P. borbonica* benefits from a higher availability of specific native tree

species (e.g., Arecaceae and Pandanaceae), which provide shelter, food and oviposition sites, as it is also found in other *Phelsuma* species from the Comoros and Seychelles (Noble et al., 2011; Augros et al., 2018; Augros, 2019). *P. borbonica* may also be adapted to cyclone activity, as this might be the case in Madagascar or in the Pacific islands (e.g., fallen trees used as hatching sites or open areas for thermoregulation; Ineich, 2010; Dubos et al., 2020). However, the species is mainly found in intact forests and may therefore use them as shelter to reduce cyclone impact instead.

Overall, *P. borbonica* can persist in very different habitats and environmental conditions, from cold and dry rocky slopes near the summits of the island (above 2.000 meters; Sanchez & Probst, 2017) to warm lowland and midland humid forests. We found a bimodality in the response curves to temperature and precipitation for several iterations. These were likely caused by the presence of isolated populations on mountain ridges and reflect the local adaptations to these specific environments. In Reunion, two subspecies were described, i.e. *P. borbonica borbonica* and *P. borbonica mater* (Meier, 1995; Probst & Deso, 2001). However, the two modalities we found do not correspond to these subspecies, which seem to share the same climatic niche instead. Preliminary unpublished genetic analyses suggest a strong isolation between these populations (Sanchez et al., 2015; unpubl. data). Further genetic analyses are needed to quantify the degree of isolation between populations.

Occurrence data which corresponded to isolated observations with no evidence of established populations (i.e., with no individual was observed after further surveys) were from areas that were predicted as unsuitable, a sign that our models identified well the species climatic niche.

 Predictions based on Worldclim indicated that suitable conditions can be met around the island at intermediate altitudes. It is possible that *P. borbonica* once occupied this whole area, but was extirpated because of intensive agriculture and habitat fragmentation in the western part of the island. A local extinction of the *P. borbonica* was already documented in Cheke (1987). This depletion was associated with the severe deforestation that occurred in the 18th century at the intermediate and lower belt of the island, and a high pressure from invasive alien species (i.e., the introduced wolf-snake *Lycodon aulicus*; Cheke & Hume, 2010). These combined factors may have dramatically decreased habitat suitability and habitat availability for *P. borbonica* in the past centuries, which may explain its absence in some of the predicted climatically suitable areas. The former presence of *P. borbonica* in the western part of the island is supported by the

presence of isolated remnant populations in the southwest (near Les Makes & Le Tampon) and recently

extinct populations in the northern lowlands (Cheke, 1987).

As shown by the response curves, *P. borbonica* persists over a relatively wide range of climatic conditions.

However, its presence remains localised throughout the island. To fully understand and explain the current

distribution patterns of *P. borbonica*, there is a need to also consider additional key factors such as micro-

habitat use and behaviour (Kearney, Shine, & Porter, 2009; Porter & Kearney, 2009). For instance, cold-

blooded species can respond to climate change by altering their period of activity and thermoregulation time

(Kearney, Shine, & Porter, 2009). Moreover, the availability of suitable oviposition sites, thermoregulation

sites with specific exposure to wind, sun or rain is a strong determinant of gecko occupancy (Ineich, 2010;

Bungard et al., 2014; Augros et al., 2017, 2018). Habitat structure is provided in native forests by tree species

from the Arecaceae and Pandanaceae families and abiotic features (e.g. sunny-exposed rocks, cliffs, man-

made structures; Petren & Case, 1998; Augros et al., 2017). In additions to climate change, habitat

modifications, such as further urbanisation and deforestation will strongly influence the future distribution of

P. borbonica in Reunion Island.

Biotic interactions play an important role in shaping species distributions (Araújo et al., 2007). The current

distribution of *P. borbonica* is strongly influenced by the occurrence of invasive alien species. The wolf

snake *Lycodon aulicus* is presumed to be the cause of historical local extinctions through predation (Cheke,

1987). The introduced Giant Madagascar day gecko *P. grandis* and the Gold-dust day gecko *P. laticauda* are

present throughout the island. Both species were reported from areas within the distribution of *P. borbonica*.

To date, there is no evidence of local extirpation of *P. borbonica* by these two species. However, *P. grandis*

was suspected to be the cause of local extinctions of *Phelsuma* species in Mauritius, presumably through

competitive exclusion and/or predation (Buckland et al., 2014). It is also the case for *P. laticauda* in French

Polynesia (Lund, 2015). *Phelsuma laticauda* is a rising cause of concern in the south of Reunion Island

where it possibly threatens the persistence of the critically endangered *P. inexpectata* (NOI, unpubl. data; but

see Porcel et al., 2021). Other introduced species may also affect *P. borbonica* through predation and/or

habitat alterations, including other reptiles (*Agama agama*, *Calotes versicolor*, *Furcifer pardalis*), rodents

(*Rattus rattus*, *Mus musculus*), ants (*Solenopsis geminata*), birds (*Pycnonotus jocosus, Acridotheres tristis*)

and plants (*Lantana camara*). The rate of invasion is increasing in Reunion Island (Fenouillas et al., 2021).

Further studies will need to account for biotic interactions to better understand the key drivers of the

distribution of *P. borbonica* and refine forecasted predictions.

Sources of uncertainty

We found a strong uncertainty driven by the modelling technique and GCMs. This is consistent with the

findings of Buisson et al. (2010) and Baker et al. (2016) and advocates the use of a wide range of modelling

techniques and GCMs for conservation planning. However, the uncertainty related to the baseline climate

data was stronger than that of GCM when considering the spatial overlap between current and future

predictions. When considering overall suitability, the importance of the baseline climate was probably

underestimated because it affected the suitability scores in a lower extent than GCMs, however, resulted in

higher discrepancies in spatial distribution of the future suitable conditions. The mismatch caused by the

baseline climate data can be due to the differences in temporal coverage, with Worldclim representing the

conditions of the 1960−1990 period while CHELSA was computed for 1979−2013. However, temporal

coverage cannot fully explain these discrepancies because future predictions (both for 2070) also strongly

differed, even when using the same predictors. Another factor potentially affecting these differences in

predictions is the higher accuracy in predictions for precipitation of the CHELSA climatic data, with more

topographic factors accounted for (Karger et al., 2017). This difference in accuracy may be exacerbated in

mountainous environments such as Reunion Island. These results suggest that the spatial component of

species range change should not be neglected when the aim is to identify priority areas for conservation.

More generally, the importance of drivers of uncertainty may be downplayed when ignoring spatial

information. We recommend the use of multiple approaches, including overlap estimations, in uncertainty

assessments of species range change.

Conservation considerations

The current distribution of *P. borbonica* is generally consistent with the predicted future suitable areas (see

Fig8b, 'Mean prediction' panel). We recommend that the areas identified with the highest certainty should be

prioritised for conservation and habitat restoration. In the context of Reunion Island, conservation actions are

drastically limited by land use policy as the available land is strongly disputed for by urbanisation and

agriculture planning. Overall, conservation efforts should be intensified in the forested uplands, where

predictions are favourable and anthropogenic pressure is the least. Nevertheless, the conservation of the

small, isolated populations of *P. borbonica* along some of the mountain ridges and in the west of Reunion

Island is of paramount importance, as these represent remnant populations that possibly form genetically

isolated entities (Sanchez et al., 2015; unpubl. data). Being supposedly more adapted to colder conditions,

mountain ridge populations may also be at even greater risk due to climate change (Raxworthy et al., 2008;

Freeman et al., 2018). We recommend the close monitoring of these populations for early detection of

potential signs of population declines.

Conservation actions should encompass a range of management strategies, including the protection of native

forests, restoration of degraded habitats, creation of artificial oviposition sites and the implementation of

sustainable agricultural practices. The control of invasive species represents an additional challenge, for

instance with *P. grandis* and *P. laticauda* currently in expansion throughout the island (Dubos, 2013; Porcel

et al., 2021).

This study stresses the need for proactive conservation actions given the high risk of extinction predicted by

some of our models. *Phelsuma borbonica* is already threatened by habitat loss and fragmentation, which will

likely increase in the future with human population. Future conservation actions will need to consider socio-

economic factors to prevent potential land use conflicts (Lagabrielle et al., 2011). This can be achieved by

involving stakeholders from urban and agricultural sectors and conservation practitioners into public

decision-making processes.

Depending on the climate data considered, our models predicted either a strong decline throughout the entire

island, upward, westward, or eastward shifts, or almost no change. Despite the high uncertainty, we

identified the areas with the most consistent predictions of suitable climate by 2070. In a context of urgent

decision making, we advocate the use of all the available tools to prevent possible extinctions in spite of the

apparent uncertainty. Forecasting models need to consider a wide range of methods and data, and assess the

variability between them in order to identify and mitigate potential sources of uncertainty and provide

relevant conservation guidelines.

Aknowledgements

We thank Boris Leroy for useful discussions. We are thankful to all the field workers that collected the data

used in this study. We thank Biotope Océan Indien and Cynorkis for providing additional data.

Data accessibility statement

All the occurrence data and the R scripts used in this study are available in online supporting information.

References

Adhikari, D., Barik, S. K., & Upadhaya, K. (2012). Habitat distribution modelling for reintroduction of Ilex khasiana Purk., a critically endangered tree species of northeastern India. *Ecol. Eng.* **40**, 37–43.

Ahmadi, M., Hemami, M. R., Kaboli, M., Malekian, M., & Zimmermann, N. E. (2019). Extinction risks of a Mediterranean neo-endemism complex of mountain vipers triggered by climate change. *Sci. Rep.* **9**, 1–12.

Araújo, M. B., & Luoto, M. (2007). The importance of biotic interactions for modelling species distributions under climate change. *Glob. Ecol. Biogeogr.* **16**, 743–753.

Augros, S. (2019). Atlas des amphibiens et des reptiles terrestres de l'archipel des Comores. *Inventaires & Biodiversité*. Biotope – Muséum national d'Histoire naturelle.

Augros, S., Faipoux, L., Bodin, M., Goff, A. Le, & Sanchez, M. (2017). Evidence for colonisation of anthropogenic habitats by the Réunion day gecko Phelsuma borbonica (Mertens , 1966) (Réunion Island , France): conservation implications **10**, 563–571.

Augros, S., Scherz, M. D., Wang-claypool, C. Y., Montfort, L., Glaw, F., & Hawlitschek, O. (2018). Comparative perch heights and habitat plant usage of day geckos (Phelsuma) in the Comoros Archipelago (Squamata : Gekkonidae). *Salamandra* **54**, 1–4.

Baker, D. J., Hartley, A. J., Butchart, S. H. M., & Willis, S. G. (2016). Choice of baseline climate data impacts projected species' responses to climate change. *Glob. Chang. Biol.* **22**, 2392–2404.

Bellis, J., Bourke, D., Maschinski, J., Heineman, K., & Dalrymple, S. (2020). Climate suitability as a predictor of conservation translocation failure. *Conserv. Biol.* **0**, 1–9.

Berghof, H. P. (2005). Taggeckos – Die Gatung Phelsuma – Pflege und Vermehrung. Natur and.

Booth, T. H., Nix, H. A., Busby, J. R., & Hutchinson, M. F. (2014). Bioclim: The first species distribution modelling package, its early applications and relevance to most current MaxEnt studies. *Divers. Distrib.* **20**, 1–9.

Brooks, M., Kristensen, K., Benthem, K. ,va., Magnusson, A., Berg, C., Nielsen, A., Skaug, H., Mächler, M., & Bolker, B. (2019). glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *R J.* **9**, 378.

Buckland, S., Cole, N. C., Aguirre-Gutiérrez, J., Gallagher, L. E., Henshaw, S. M., Besnard, A., Tucker, R. M., Bachraz, V., Ruhomaun, K., & Harris, S. (2014). Ecological effects of the invasive giant madagascar day gecko on endemic Mauritian geckos: Applications of binomial-mixture and species distribution models. *PLoS One* **9**.

Buisson, L., Thuiller, W., Casajus, N., Lek, S., & Grenouillet, G. (2010). Uncertainty in ensemble forecasting of species distribution. *Glob. Chang. Biol.* **16**, 1145–1157.

Bungard, M. J., Jones, C., Tatayah, V., & Bell, D. J. (2014). The habitat use of two species of Day Geckos (Phelsuma ornata and Phelsuma guimbeaui) and implications for conservation management in island ecosystems. *Herpetol. Conserv. Biol.* **9**, 551–562.

Butt, N., Chauvenet, A. L. M., Adams, V. M., Beger, M., Gallagher, R. V., Shanahan, D. F., Ward, M., Watson, J. E. M., & Possingham, H. P. (2020). Importance of species translocations under rapid climate change. *Conserv. Biol.* **00**, 1–9.

Cheke, A. S. (1987). An ecological history of the Mascarene Islands, with particular reference to extinctions and introductions of land vertebrates. In *Stud. Mascarene Isl. Birds*. Diamond (e., pp. 5–89. Cambridge.

Cheke, A. S., & Hume, J. P. (2010). Lost land of the dodo: the ecological history of Mauritius, Réunion and Rodrigues. Bloomsbury.

Converse, S. J., & Sipe, H. A. (2021). Finding the win-win strategies in endangered species conservation. *Anim. Conserv.* **24**, 161–162.

Cordier, J. M., Lescano, J. N., Rios, N. E., Leynaud, G. C., & Nori, J. (2020). Climate change threatens micro-endemic amphibians of an important South American high-altitude center of endemism. *Amphib. Reptil.* **41**, 233–243.

Datta, A., Schweiger, O., & Kühn, I. (2020). Origin of climatic data can determine the transferability of species distribution models. *NeoBiota* **59**, 61–76.

Deso, G. (2001). Note sur le transport insolite de Geckos verts, le cas du Phelsuma inexpectata. *Bull. Phaethon* **13**, 56.

Dormann, C. F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J. R. G., Gruber, B., Lafourcade, B., Leitão, P. J., Münkemüller, T., Mcclean, C., Osborne, P. E., Reineking, B., Schröder, B., Skidmore, A. K., Zurell, D., & Lautenbach, S. (2013). Collinearity: A review of methods to deal with it and a simulation study evaluating their performance. *Ecography (Cop.).* **36**, 27–46.

Draper, D., Marques, I., & Iriondo, J. M. (2019). Species distribution models with field validation, a key approach for successful selection of receptor sites in conservation translocations. *Glob. Ecol. Conserv.* **19**.

Dubos, N. (2013). New locality record for Phelsuma grandis (Sauria: Gekkonidae) in Reunion, in sympatry with the Critically Endangered Phelsuma inexpectata. *Herpetol. Notes* **6**, 309–311.

Dubos, N., Dehorter, O., Yves, P., & Viol, I. Le. (2019). Thermal constraints on body size depend on the population's position within the species' thermal range in temperate songbirds. *Glob. Ecol. Biogeogr.* **28**, 96–106.

Dubos, N., Montfort, F., Grinand, C., Nourtier, M., Deso, G., Razafimanahaka, J. H., Andriantsimanarilafy, R. R., Fanantenana, E., Razafindraibe, P., Jenkins, R., Crottini, A., & Associations, M. (2021). Are narrow-ranging species doomed to extinction? Probable dramatic and generalised decline in future climate suitability of highly threatened species. *bioRxiv*.

Dubos, N., Morel, L., Crottini, A., Freeman, K., Honoré, J., Lava, H., Noël, J., Porton, I., Rendryrendry, G., Rosa, G., & Andreone, F. (2020). High interannual variability of a climate - driven amphibian community in a seasonal rainforest. *Biodivers. Conserv.* **29**, 893–912.

Dubos, N., Préau, C., Lenormand, M., Papuga, G., Montsarrat, S., Denelle, P., Louarn, M. Le, Heremans, S., Roel, M., Roche, P., & Luque, S. (2021). Assessing the effect of sample bias correction in species distribution models. *arXiv*.

Dupuy, S., Gaetano, R., & Le Mézo, L. (2020). Mapping land cover on Reunion Island in 2017 using
satellite imagery and geospatial ground data. *Data Br.* **28**.

Elith, J., Leathwick, J. R., & Hastie, T. (2008). A working guide to boosted regression trees. *J. Anim. Ecol.*
77, 802–813.

Fenouillas, P., Ah-Peng, C., Amy, E., Bracco, I., Dafreville, S., Gosse, M., Ingrassia, F., Lavergne, C.,
Lequette, B., Notter, J. C., Pausé, J. M., Paye, G., Payet, N., Picot, F., Poungavanon, N., Strasberg, D.,
Thomas, H., Triolo, J., Turquet, V., & Rouget, M. (2021). Quantifying invasion degree by alien plants
species in Reunion Island. *Austral Ecol.* 1–13.

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: new 1-km spatial resolution climate surfaces for global
land areas. *Int. J. Climatol.* **37**, 4302–4315.

Freeman, B. G., & Class Freeman, A. M. (2014). Rapid upslope shifts in New Guinean birds illustrate strong
distributional responses of tropical montane species to global warming. *Proc. Natl. Acad. Sci. U. S. A.*
111, 4490–4494.

Freeman, B. G., Scholer, M. N., Ruiz-Gutierrez, V., & Fitzpatrick, J. W. (2018). Climate change causes
upslope shifts and mountaintop extirpations in a tropical bird community. *Proc. Natl. Acad. Sci. U. S. A.*
115, 11982–11987.

Guisan, A., Edwards, T. C., & Hastie, T. (2002). Generalized linear and generalized additive models in
studies of species distributions: setting the scene. *Ecol. Modell.* **157**, 89–100.

Hijmans, R. J. (2012). Cross-validation of species distribution models: removing spatial sorting bias and
calibration with a null model. *Ecology* **93**, 679–688.

Ineich, I. (2010). How habitat disturbance benefits geckos: Conservation implications. *Comptes Rendus -*
Biol. **333**, 76–82.

Irl, S. D. H., Schweiger, A. H., Steinbauer, M. J., Ah-Peng, C., Arévalo, J. R., Beierkuhnlein, C., Chiarucci,
A., Daehler, C. C., Fernández-Palacios, J. M., Flores, O., Kueffer, C., Maděra, P., Otto, R., Schweiger,
J. M. I., Strasberg, D., & Jentsch, A. (2021). Human impact, climate and dispersal strategies determine
plant invasion on islands. *J. Biogeogr.* 1–15.

Jiguet, F., Devictor, V., Ottvall, R., Van Turnhout, C., Van der Jeugd, H., & Lindström, A. (2010). Bird
population trends are linearly affected by climate change along species thermal ranges. *Proc. R. Soc. B*
Biol. Sci. **277**, 3601–8.

Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann, N. E., Linder,
H. P., & Kessler, M. (2017). Climatologies at high resolution for the earth's land surface areas. *Sci.*
Data **4**, 1–20.

Kearney, M., Shine, R., & Porter, W. P. (2009). The potential for behavioral thermoregulation to buffer
"cold-blooded" animals against climate warming. *Proc. Natl. Acad. Sci.* **106**, 3835–3840.

Kujala, H., Moilanen, A., Araújo, M. B., & Cabeza, M. (2013). Conservation Planning with Uncertain
Climate Change Projections. *PLoS One* **8**.

Lagabrielle, E., Rouget, M., Le Bourgeois, T., Payet, K., Durieux, L., Baret, S., Dupont, J., & Strasberg, D.
(2011). Integrating conservation, restoration and land-use planning in islands-An illustrative case study
in Réunion Island (Western Indian Ocean). *Landsc. Urban Plan.* **101**, 120–130.

Lagabrielle, E., Rouget, M., Payet, K., Wistebaar, N., Durieux, L., Baret, S., Lombard, A., & Strasberg, D.
(2009). Identifying and mapping biodiversity processes for conservation planning in islands: A case
study in Réunion Island (Western Indian Ocean). *Biol. Conserv.* **142**, 1523–1535.

Lannuzel, G., Balmot, J., Dubos, N., Thibault, M., & Fogliani, B. (2021). High-resolution topographic
variables accurately predict the distribution of rare plant species for conservation area selection in a
narrow-endemism hotspot in New Caledonia. *Biodivers. Conserv.* **30**, 963–990.

Leathwick, J. R., Rowe, D., Richardson, J., Elith, J., & Hastie, T. (2005). Using multivariate adaptive
regression splines to predict the distributions of New Zealand's freshwater diadromous fish. *Freshw.*
Biol. **50**, 2034–2052.

Legrand, B., Benneveau, A., Jaeger, A., Pinet, P., Potin, G., Jaquemet, S., & Le Corre, M. (2016). Current
wintering habitat of an endemic seabird of Réunion Island, Barau's petrel Pterodroma baraui, and
predicted changes induced by global warming. *Mar. Ecol. Prog. Ser.* **550**, 235–248.

Leroy, B., Bellard, C., Dubos, N., Colliot, A., Vasseur, M., Courtial, C., Bakkenes, M., Canard, A., & Ysnel,
F. (2014). Forecasted climate and land use changes, and protected areas: the contrasting case of spiders.
(B. Bradley, Ed.)*Divers. Distrib.* **20**, 686–697.

Leroy, B., Delsol, R., Hugueny, B., Meynard, C. N., Barhoumi, C., Barbet-Massin, M., & Bellard, C. (2018).
Without quality presence–absence data, discrimination metrics such as TSS can be misleading
measures of model performance. *J. Biogeogr.* **45**, 1994–2002.

Lund, I. (2015). Moorea's newest invasive species: the distribution and behavior of Phelsuma laticauda. *Biol.*
Geomorphol. Trop. Islands.

Macdonald, I. A. W., & Cedex, F. M. (1991). Effects of Alien Plant Invasions on Native Vegetation
Remnants on La Reunion (Mascarene Islands, Indian Ocean) Author (s): Ian A. W. Macdonald,
Christophe Thébaud, Wendy A. Strahm and Dominique Strasberg Published by : Cambridge University
Press **18**.

Manel, S., Dias, J. M., & Ormerod, S. J. (1999). Comparing discriminant analysis, neural networks and
logistic regression for predicting species distributions: A case study with a Himalayan river bird. *Ecol.*
Modell. **120**, 337–347.

Manes, S., Costello, M. J., Beckett, H., Debnath, A., Devenish-Nelson, E., Grey, K.-A., Jenkins, R., Khan, T.
M., Kiessling, W., Krause, C., Maharaj, S. S., Midgley, G. F., Price, J., Talukdar, G., & Vale, M. M.
(2021). Endemism increases species' climate change risk in areas of global biodiversity importance.
Biol. Conserv. **257**, 109070.

Meier, H. (1995). Neue Nachweise von Phelsuma borbonica auf Reunion, Maskarenen, mit dem Versuch
einer taxonomischen Einordnung. *Salamandra* **31**, 33–40.

Minteer, B. A., & Collins, J. P. (2010). Move it or lose it? the ecological ethics of relocating species under
climate change. *Ecol. Appl.* **20**, 1801–1804.

Morales-Barbero, J., & Vega-Álvarez, J. (2019). Input matters matter: Bioclimatic consistency to map more
reliable species distribution models. *Methods Ecol. Evol.* **10**, 212–224.

Muscatello, A., Elith, J., & Kujala, H. (2020). How decisions about fitting species distribution models affect
conservation outcomes. *Conserv. Biol.*

Noble, T., Bunbury, N., Kaiser-Bunbury, C. N., & Bell, D. J. (2011). Ecology and co-existence of two
endemic day gecko (Phelsuma) species in Seychelles native palm forest. *J. Zool.* **283**, 73–80.

Ocon, J. P. (2020). A Global Analysis of Tropical Dry Forest Extent and Cover Based on Climatic Definitions. University of California.

Pearson, R. G., Stanton, J. C., Shoemaker, K. T., Aiello-Lammens, M. E., Ersts, P. J., Horning, N., Fordham, D. A., Raxworthy, C. J., Ryu, H. Y., Mcnees, J., & Akçakaya, H. R. (2014). Life history and spatial traits predict extinction risk due to climate change. *Nat. Clim. Chang.* **4**, 217–221.

Petren, K., & Case, T. J. (1998). Habitat structure determines competition intensity and invasion success in gecko lizards. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 11739–11744.

Phillips, S. J., D., M., & Schapire, R. E. (2006). AMaximum Entropy Approach to Species Distribution Modeling Steven. *Ecol. Modell.* **190**, 231–259.

Platts, P. J., Garcia, R. A., Hof, C., Foden, W., Hansen, L. A., Rahbek, C., & Burgess, N. D. (2014). Conservation implications of omitting narrow-ranging taxa from species distribution models, now and in the future. *Divers. Distrib.* **20**, 1307–1320.

Porcel, X., Deso, G., Probst, J., & Dubos, N. (2021). Sympatrie entre le Gecko vert de Manapany Phelsuma inexpectata endémique de la Réunion et le Gecko vert poussière d ' or P . laticauda introduits au Domaine du café grillé : peuvent-ils cohabiter ? *Bull. Phaethon* **53**, 36–38.

Porter, W. P., & Kearney, M. (2009). Size, shape, and the thermal niche of endotherms. *Proc. Natl. Acad. Sci.* **106**, 19666–19672.

Prasad, A. M., Iverson, L. R., & Liaw, A. (2006). Newer classification and regression tree techniques: Bagging and random forests for ecological prediction. *Ecosystems* **9**, 181–199.

Probst, J.-M., & Deso, G. (2001). Le gecko vert des forêts Phelsuma borbonica. *Bull. Phaethon* **13**, 23–25.

Raxworthy, C. J., Pearson, R. G., Rabibisoa, N., Rakotondrazafy, A. M., Ramanamanjato, J.-B., Raselimanana, A. P., Wu, S., Nussbaum, R. a., & Stone, D. a. (2008). Extinction vulnerability of tropical montane endemism from warming and upslope displacement: a preliminary appraisal for the highest massif in Madagascar. *Glob. Chang. Biol.* **14**, 1703–1720.

Sanchez, M., Duval, T., Le Péchon, T., Cornuault, J., & Thébaud, C. (2015). Le gecko vert de Bourbon (Phelsuma borbonica) : répartition, écologie et génétique des populations, implications pour sa conservation. In S. Baret, E. Braun, B. Lequette, & D. Strasberg (Eds.), *Séminaire des Gest. la Conserv. la BIOdiversité à La Réunion, GECOBIO 1, 18 novembre 2015, Parc Natl. La Réunion*. pp. 26–31.

Sanchez, Mickaël, & Probst, J.-M. (2016). L'herpétofaune allochtone de l'île de La Réunion (Océan Indien): état des connaissances en 2015. *Bull. la Société Herpétologique Fr.* **160**, 49–78.

Sanchez, Mickaël, & Probst, J. (2017). Phelsuma borbonica Mertens , 1966 (Sauria : Gekkonidae) sur l'île de La Réunion . I . Répartition et habitats naturels. *Bull. la Société herpétologique Fr.* **162**, 17–30.

Strasberg, D., Rouget, M., Richardson, D. M., Baret, S., Dupont, J., & Cowling, R. M. (2005). An Assessment of habitat diversity and transformation on La Réunion Island (Mascarene Islands, Indian Ocean) as a basis for identifying broad-scale conservation priorities. *Biodivers. Conserv.* **14**, 3015–3032.

Sunday, J. M., Bates, A. E., & Dulvy, N. K. (2011). Global analysis of thermal tolerance and latitude in ectotherms. *Proceeding R. Soc. London B Biol. Sci.* **278**, 1823–1830.

Tang, Y., Winkler, J. A., Viña, A., Liu, J., Zhang, Y., Zhang, X., Li, X., Wang, F., Zhang, J., & Zhao, Z. (2018). Uncertainty of future projections of species distributions in mountainous regions. *PLoS One* **13**.

Tessarolo, G., Lobo, J. M., Rangel, T. F., & Hortal, J. (2021). High uncertainty in the effects of data
characteristics on the performance of species distribution models. *Ecol. Indic.* **121**, 107147.

Tewksbury, J. J., Huey, R. B., & Deutsch, C. A. (2008). Putting the Heat on Tropical Animals The Scale of
Prediction. *Science* **320**, 1296–1297.

Warren, D. L., Glor, R. E., & Turelli, M. (2010). ENMTools: A toolbox for comparative studies of
environmental niche models. *Ecography (Cop.)*. **33**, 607–611.

Westwood, R., Westwood, A. R., Hooshmandi, M., Pearson, K., LaFrance, K., & Murray, C. (2020). A field-
validated species distribution model to support management of the critically endangered Poweshiek
skipperling (Oarisma poweshiek) butterfly in Canada. *Conserv. Sci. Pract.* **2**, 1–15.

Wilson, C. D., Roberts, D., & Reid, N. (2011). Applying species distribution modelling to identify areas of
high conservation value for endangered species: A case study using Margaritifera margaritifera (L.).
Biol. Conserv. **144**, 821–829.

Wisz, M. S., & Guisan, A. (2009). Do pseudo-absence selection strategies influence species distribution
models and their predictions? An information-theoretic approach based on simulated data. *BMC Ecol.*
9.

Zhong, Y., Xue, Z., Jiang, M., Liu, B., & Wang, G. (2021). The application of species distribution modeling
in wetland restoration: A case study in the Songnen Plain, Northeast China. *Ecol. Indic.* **121**, 107137.
634