Sensitivity to Error During Visuomotor Adaptation is Similarly Modulated by Abrupt, Gradual and Random Perturbation Schedules

Susan K. Coltman[1,2,3], Robert J. van Beers[4,5], W. Pieter Medendorp[4], and Paul L. Gribble[2,3,6,7]

[1] Graduate Program in Neuroscience, Western University, London, ON, Canada
[2] Brain and Mind Institute, Western University, London, ON, Canada
[3] Department of Psychology, Western University, London, ON, Canada
[4] Donders Institute for Brain, Cognition and Behaviour, Radboud University, Nijmegen, The Netherlands
[5] Dept. of Human Movement Sciences, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands
[6] Department of Physiology and Pharmacology, Schulich School of Medicine & Dentistry, Western University, London, ON, Canada
[7] Haskins Laboratories, New Haven CT, USA

Acknowledgements: Canadian Institutes of Health Research (CIHR) and the Natural Sciences and Engineering Council of Canada (NSERC).

The authors declare no competing financial interests.

Please Address Correspondence to:

Paul L. Gribble
Brain and Mind Institute, Dept. of Psychology
Western University
1151 Richmond St.,
London, ON, Canada, N6A 3K7
Email: pgribble@uwo.ca

Running title: How prior experience modulates error sensitivity

Abstract

It has been suggested that sensorimotor adaptation involves at least two processes (i.e., fast and slow) that differ in retention and error sensitivity. Previous work has shown that repeated exposure to an abrupt force field perturbation results in greater error sensitivity for both the fast and slow processes. While this implies that the faster relearning is associated with increased error sensitivity, it remains unclear what aspects of prior experience modulate error sensitivity. In the present study, we manipulated the initial training using different perturbation schedules, thought to differentially affect fast and slow learning processes based on error magnitude, and then observed what effect prior learning had on subsequent adaptation. During initial training of a visuomotor rotation task, we exposed three groups of participants to either an abrupt, a gradual, or a random perturbation schedule. During a testing session, all three groups were subsequently exposed to an abrupt perturbation schedule. Comparing the two sessions of the control group who experienced repetition of the same perturbation, we found an increased error sensitivity for both processes. We found that the error sensitivity was increased for both the fast and slow processes, with no reliable changes in the retention, for both the gradual and structural learning groups when compared to the first session of the control group. We discuss the findings in the context of how fast and slow learning processes respond to a history of errors.

Keywords

human; motor learning; two-state model; error sensitivity; savings

New & Noteworthy

We investigated what aspects of prior experience modulate error sensitivity, within the

framework of a two-state model of short-term sensorimotor adaptation. We manipulated initial

training on a visuomotor adaptation reaching task using specific perturbation schedules that are

thought to differentially affect fast and slow learning processes, and we tested what effect these

had on subsequent adaptation. We found that sensitivity to adaptation error was similarly

modulated by abrupt, gradual, and random perturbation schedules.

Introduction

Adaptation is often defined as an error-driven process, in which the error experienced during a movement leads to a corrective adjustment in the motor output on the following movement (1-5). Behavioural measures of adaptation are well characterized by state-space models (1, 4), which represent trial-to-trial changes in movement as a function of how an error on a given trial affects motor output on the subsequent trial. The update from one trial to the next, or the change in motor output, is based on two parameters: a retention parameter which determines what proportion of motor output is retained from trial to trial, and an error sensitivity parameter which governs the proportion of error experienced on the current trial that is corrected for on the subsequent trial.

Variations of the state-space model are built on the assumption that adaptation is the product of multiple underlying processes with distinct timescales (3, 6-8). Researchers have begun to provide neural evidence to strengthen the theory that sensorimotor learning is supported by multiple processes (9, 10). An influential two-state model of short-term motor adaptation was proposed by Smith et al. (3) that proposed a fast process that learns quickly but has poor retention and a slow process that learns more slowly, but has strong retention.

The prevailing success of the two-state model continues to be that it accounts for the learning phenomenon known as savings, characterized as prior learning speeding up subsequent relearning (3, 11). While Smith et al. (3) initially argued that the reason for the fast relearning during a second introduction of the same perturbation was due to the resistance of the slow process to change, recent studies suggest that learning rate can be modified depending on factors such as the uncertainty of movement error (12, 13), size of movement error (14), and a history of movement errors (15-17).

The behavioural changes associated with savings suggest that some component of

memory from the initial training must lead to the faster relearning, but what is remembered and

recalled remains unclear (15-22). One perspective argues for the enhancement of an explicit

strategy (18, 20, 23), while the other side suggests that faster relearning is driven by the

experience of the motor errors (15-17, 24).

In support of the latter possibility, Herzfeld and colleagues (16) proposed that a history of

errors modulates the error sensitivity on each trial, systematically controlling how much the

motor system learns from the current motor error. They suggested that an error-based adaptation

model that provides for experience-dependent error sensitivity modification could account for

savings. Furthermore, Leow et al. (17) demonstrated that it is a memory of errors, not previous

actions, that is necessary for savings.

Recent work has shown that repeated exposure to the same force field perturbation results

in greater error sensitivity of both the fast and slow processes (15). While in Coltman et al. (15),

the error sensitivity terms for the fast and slow processes were held constant within a session, we

evaluated the theory of experience-dependent error sensitivity modulation in the context of

changes in error sensitivity from one learning session to the next. Although these results clearly

indicate that the motor system stored some component (i.e., memory) of prior training to speed

up subsequent learning, it remains unclear how the fast and slow learning processes contribute to

savings and what aspects of prior experience modulate error sensitivity. In other words, do both

fast and slow processes access a single stored component of prior training, or do they store

independent components? If the fast and slow processes depend on separate stored components

of prior training, then it may be possible to independently modulate characteristics of the fast and

slow processes (e.g., error sensitivity) by experimentally manipulating aspects of prior learning.

Recent findings demonstrate that participants do not adapt linearly in response to

different magnitudes of error (13, 14, 25). Additionally, Orban de Xivry and Lefevre (26)

propose that different perturbation schedules lead to distinct motor memories with different

attributes and neural representations (i.e., the amount of reorganization of the motor cortex). We

propose that perturbation schedules that are designed to produce learning using errors of

different magnitudes may have a differential effect on session-to-session changes to the fast

versus slow processes. In the present study, we manipulated initial training in a visuomotor

adaptation task using perturbation schedules which involved errors of different magnitudes, and

we tested what effect these different initial learning experiences had on subsequent adaptation,

and specifically on characteristics of the fast versus slow adaptation processes.

We asked one group of participants to counter a gradual perturbation schedule during

initial training. When a perturbation is gradually introduced, such that participants never

experience large errors, learning is believed to be more implicit in nature (26). We predicted that

when participants in this group were later tested on an abrupt perturbation, only the slow process

would be affected by the initial training, compared to a control group who were initially trained

using an abrupt perturbation. For a second group of participants, initial training was based on a

structural learning paradigm, involving a series of brief exposures to large, random perturbations

(27, 28). This perturbation schedule is thought to be based on explicit learning mechanisms (20,

29). During the original conception of the two-state model, Smith et al. (3) designed a rebound

paradigm to test the theorized characteristics of each process. Much like the brief reversal used in

the rebound paradigm, which includes large errors and limited exposure, the motor output during

this phase of the paradigm is believed to be dependent on the fast process (3). For this group we

predicted that when later tested on an abrupt perturbation, only the fast process would be affected

by the initial training, as compared to the control group.

We modelled perturbation-driven changes in movement with the state-space equations

proposed by Smith et al. (3), and focused on changes in the retention and error sensitivity

parameters. The model estimates function as a tool for understanding how the underlying

processes of adaptation were affected by the prior training. Substantiating the finding of Coltman

et al. (15), we confirm that repetition of the same visuomotor perturbation results in an increase

in error sensitivity for both processes, when comparing the two sessions of the control group. By

comparing the model estimates of participants in the gradual and structural learning groups to the

first session of the control, we expected to see changes in error sensitivity that depended on the

type of prior training participants experienced. Interestingly, however, we found that error

sensitivity of both the fast and slow processes was increased for both groups. The findings are

discussed in the context of storing and accessing a history of errors.

158   **Methods**

Participants. A total of 60 healthy young adults (age range 21-35; mean age ± sd 27.9 ± 4.2

years) participated in a visuomotor rotation experiment. Participants were recruited from the

online platform maintained by Prolific.co and received £11.25 for their participation. As part of

the Prolific platform, participants respond to a series of questions related to age, gender, health

and economic status. Based on this prescreen information, 24 participants identified as female

and 36 as male. Participants were recruited globally and reported being located in 17 different

countries (Estonia, Finland, France, Greece, Hungary, Israel, Italy, Mexico, Netherlands, Poland,

Portugal, Slovenia, South Africa, Spain, Sweden, United Kingdom and the United States). All

participants self-reported being right-handed and had normal or corrected-to-normal vision. The

protocol was approved by Western University's Research Ethics Board and all participants

indicated electronic consent.

Apparatus. Participants used a standard computer mouse and their own computer to access a

webpage hosted on a network computer located at Western Interdisciplinary Research Building.

The task was written in and controlled by JavaScript, running locally within the participants'

web browser.

Participants were asked to use a standard computer mouse and a standard credit or debit

card to complete a spatial calibration procedure. Participants were initially instructed how to turn

off the acceleration for the mouse, based on their operating system. Then, following an

instruction video, participants were asked to align the top of their mouse with the top of the

credit card. After a tone, they were instructed to move the mouse in a smooth and straight path,

aligning the top of their mouse with the bottom of the card. Participants were asked to hold still

while waiting for a second tone, indicating that they needed to realign the mouse with the top of

the card. This was repeated at two different speeds indicated in the video. When the calibration

procedure was successfully completed, participants watched an instructional video about the

experimental task.

The size and position of the stimuli were scaled based on a mouse calibration procedure.

Real-time position of the mouse was used to control the visual display and to provide on-line

visual feedback. The mouse speed was adjusted such that the distance from start position to

target was exactly 6 cm based on the calibration. While the physical target distance was always 6

cm, this translated to 300 pixels on screen. Therefore, the straight reach trajectory was 300

pixels, however a participant's view of this was potentially compressed or expanded relative to

190 the target value of 6 cm, depending on their monitor as well their viewing distance from the

191 monitor.

192 **Paradigm.** At the start of each trial, participants were instructed to click their mouse to begin. A

193 circular cursor (10 pixels radius) was virtually displayed on the participant's computer monitor

194 and was used to represent the position of the mouse on screen. The position of the mouse at the

195 start of the trial, represented the start position on screen. A small square (20 pixels by 20 pixels)

196 represented the target. The radial distance of the target from the start position was 300 pixels.

197 The target appeared at either 45°, 90°, or 135°, relative to the start position (where 6 cm directly

198 to the right of the start position represented 0°). The location of the target was randomized per

199 trial, per participant, such that each participant saw a different order of targets with an equal

200 number of presentations of each target over the course of a session.

201  Participants were instructed to make a straight movement from the start position to the

202 target, within a narrow temporal window. At the beginning of each trial the target appeared in

203 white. Participants were required to hold still at the start position for 500 ms, at which time the

204 target changed color to green, representing a "go" signal for participants to initiate a movement

205 to the target. In addition to the colour change of the target, a tone was used as a secondary "go"

206 signal. Participants needed to reach for the target and bring the centre of a red cursor

207 representing the position of their computer mouse within 10 pixels of the centre of the target

208 within 600–900 ms. If a participant's movement time was less than 600 ms, the target turned red

209 to indicate that the movement was "too fast". If the participant's movement time was within

210 600–900 ms, the target remained green to indicate that the movement was "good". If the

211 participant's movement time was greater than 900 ms, the target would turn blue to indicate "too

212 slow". Feedback related to movement time was displayed on the screen for 1000 ms before the

screen went blank and written instructions on screen indicated that the participant should return

the mouse to a comfortable starting position within their workspace. Participants were instructed

to try to obtain the "good" feedback as often as possible throughout the experiment.

To assist with making straight movements between the start position and the target using

a computer mouse, the first 20 trials of the first session represented a practice session for

participants. In these trials, a purple rectangle (50 pixels by 300 pixels), with two white lines on

either side was shown on screen, highlighting a straight path to the target. Participants were

instructed to keep the red cursor on the path, between the lines, toward the target. If the cursor

moved outside the path, the background colour changed from black to pink.

Participants were randomly assigned to one of three groups. Each group completed two

sessions (initial training and testing), separated by a 5-minute break (**Fig. 1**).

Each session included a total of 450 reaching movements, with a 1-minute mid-session break

halfway. The experimental paradigm for each session consisted of 4 epochs. The first epoch

(baseline) consisted of 70 trials in which participants were provided with veridical feedback of

the cursor position. The second epoch (adaptation) consisted of 300 trials in which a visuomotor

rotation was applied to the cursor feedback: an angular rotation was imposed on the cursor, such

that a hand movement aimed directly at a target produced a cursor movement that was rotated

radially about the start position and participants saw that their movement had generated an error.

Participants had to learn to counter the rotation by moving their hand in an equal and opposite

direction. With practice, participants adjusted their movements in such a manner that the visual

feedback produced straight trajectories from start position to the target. In the third epoch (error-

clamp; consisting of 30 trials) the task error was clamped to zero. During the clamp trials, the

angular position of the cursor relative to the start position was clamped to a straight line

connecting the start position to the target, while participants maintained control of the radial

distance of the cursor from the start position. Finally, in the fourth epoch (washout; consisting of

50 trials), participants were provided again with veridical feedback to bring performance back to

baseline.

During the adaptation epoch of the first session, participants experienced one of three

conditions: (1) a control learning group (n=20) experienced an abrupt 30° clockwise (CW)

rotation for all 300 trials during this phase (**Fig. 1; top**), (2) a gradual learning group, (n=20) in

which a rotation was increased linearly from 0° to 30° CW over 250 trials and then held at a

fixed 30° CW for another 50 trials (**Fig. 1; middle**), or (3) a structural learning group (n=20) in

which participants encountered random rotations, ranging from 60° counter-clockwise (CCW) to

60° CW in blocks of 6 trials with the same rotation (27, 28, 29; Fig. 1; bottom). In this group, we

deliberately set the average over all angles to zero, to prevent any accumulative learning. We

also excluded rotation sizes within 10° of the test rotation (30° CW) and its inverse (30° CCW).

We furthermore set the change in rotation angle to be equal to or greater than 15° to ensure the

errors were always large, which characteristically has the greatest influence on the fast process

(3, 29). During the second session, all three groups experienced an abrupt 30° CW rotation

during the adaptation epoch.

Data Analysis. The position of the cursor in both x (lateral) and y (sagittal), were sampled in

pixels at the refresh rate of their computer monitor (typically 60 Hz). Missed samples were

interpolated during analysis (less than 1 % of samples on average). In cases in which data were

acquired at higher sampling rates (for example because a participant's computer monitor refresh

rate exceeded 60 Hz), the data were down sampled to 60 Hz. Data were digitally smoothed using

a second-order low-pass Butterworth filter with a cut-off frequency of 15 Hz. All data were

stored for offline analysis using custom MATLAB R2020a (The MathWorks) scripts.

Movement trajectories were selected using an algorithm in which movement initiation

was defined as the time at which the tangential velocity of the mouse first exceeded 0.5 cm/s and

movement end was defined as the first time after peak velocity that tangential velocity fell below

0.5 cm/s, where peak velocity was defined as the fastest participants ever moved during the reach

movement. For each trial we computed the angle between the line connecting the start position

and the cursor position at peak velocity, and the line connecting the start position to the target.

We determined the average reach angle, per subject during the last 50 trials of the baseline epoch

and we subtracted this quantity from the reach angle measured on each trial.

Model fitting. Smith et al. (3) outlined a method for mathematically modelling an iterative

update of the states of the two proposed processes of short-term sensorimotor adaptation.

Essentially, the model involves fitting four parameters: an error sensitivity and a retention

parameter for both a fast and a slow process. The first parameter weighs the relative importance

of recalling previous motor commands, which is interpreted as the retention factor. The second

parameter is the sensitivity to error, which relates to the proportion of error that is corrected for

trial-to-trial (1, 3, 4, 30). The two important assumptions in this model are that the error

sensitivity is higher for the fast process compared with the slow process and that retention is

stronger for the slow process compared with the fast process (3). Adaptation can be decomposed

into a fast (**Eq. 1**) and a slow (**Eq. 2**) process, knowing that each state follows different learning

dynamics. The two processes are summed together to produce the overall output x (**Eq. 3**). Error,

denoted by e(n), arises on each trial n as the difference between the overall output xnet and the

task parameter r (i.e., the degree of the rotation; **Eq.4**).

$$x_f(n + 1) = A_f x(n) + B_f e(n) \tag{1}$$

$$x_s(n + 1) = A_s x(n) + B_s e(n) \tag{2}$$

$$x_{net}(n) = x_f(n) + x_s(n) \tag{3}$$

$$e(n) = r(n) - x_{net}(n) \tag{4}$$

285
Linear inequality constraints were defined in order to apply to standard two-state model

dynamics (31):

$$A_f \leq A_s + 0.001 \tag{5}$$

$$B_f \geq B_s + 0.001 \tag{6}$$

In order to approximate the four parameters (i.e., Af, As, Bf, and Bs), we fit the model to

the behavioral data (using the function *fmincon* in MATLAB r2020a) by minimizing the squared

difference between the estimated net output (xnet) of the model and the average participant reach

angle, measured on each trial. According to the methods described in Albert and Shadmehr (31),

we also included a mathematical formalization of visual error clamp trials and set breaks.

Statistical Design. Pairwise comparisons were performed with nonparametric bootstrap

hypothesis tests, as well as paired and unpaired t-tests. For statistical analyses that require

multiple comparisons, we used the Holm-Bonferroni correction (32). Statistical tests were

considered significant at $p < 0.05$. For all reported and depicted values, we report the mean and

SEM.

300   **Results**

Figure 2 shows the hand paths from one representative participant in the control group

during both sessions one and two, as well as one representative participant per group in session

two of the structural and gradual learning groups. During the baseline epoch (left column), these

paths are relatively straight to the target. The representative participants were all adapting to an

abrupt 30° CW rotation. During the early adaptation epoch (middle column) these movements

were initially deviated in the CW direction, with a corrective movement at the end of the

trajectory to bring the cursor to the target. In all three groups, participants adapted to the 30° CW

rotation by the late adaptation epoch (right column), reducing their movement errors and

resuming relatively straight hand paths to the target.

We used a kinematic behavioural measure to assess changes in performance. The primary

outcome measure for the study was reach angle at peak velocity, which was measured as the

angle between the straight line connecting the start position and the cursor position at peak

velocity and the straight line connecting the start position to the target. The control group of

participants adapted their movements to an abrupt 30° CW visuomotor rotation in both the first

and second session. **Figure 3A** shows the angle at peak velocity for all trials in each session,

averaged across participants in the control group. In both sessions, participants exhibited learning

during the adaptation epoch, decay during the visual error clamp epoch, and a return towards

baseline performance during the washout epoch. During the adaptation epoch we examined the

learning at two different time points: early (first fifty trials during adaptation) and late (last fifty

trials during adaptation; **Fig. 3B**). The mean angle in the early learning phase of the second

session ($M = 23.7$, $SD = 3.15$) was reliably greater than in the first session [$M = 19.9$, $SD =$

4.6; paired t-test, $t(19) = -6.2$, $P = 3.0\text{e-}06$], indicating savings. We did not detect a reliable

difference ($P = 0.08$) between sessions during late learning.

A second group of participants was exposed to a gradual perturbation schedule during

initial training. **Figure 4A** shows the angle at peak velocity for all trials in each session, averaged

across participants in the gradual learning group. Participants exhibited learning during the

adaptation epoch, decay during the visual error clamp epoch, and a return towards baseline

performance during the washout epoch. A final group of participants was exposed to a series of

brief exposures to large, random perturbations. Each participant in this group experienced a

different set of randomly varying rotations. **Figure 4B** illustrates the angle at peak velocity for

all trials in session one for four representative individual participants from the structural learning

group. We observed two participants who demonstrated learning within each block of six trials,

but who also appeared to have maintained a fraction of error throughout the adaptation epoch

(**Fig. 4B**, **S2** and **S8**). In addition to a participant who adapted quickly to the randomly changing

perturbation (**Fig. 4B**, **S18**), we observed a participant who qualitatively showed greater

reduction of error in the latter half of the adaptation epoch, compared to the early half (**Fig. 4B**,

S20).

Figure 5A shows the angle at peak velocity averaged across participants for all trials in

session one of the control group and session two of the structural and gradual learning groups.

When comparing the model estimates of participants in the gradual and structural learning

groups during the second session to the first session of the control group, we expected to see

changes in error sensitivity that depended on the type of prior training participants experienced.

To compare the changes in angle between the control, structural and gradual learning groups, we

examined learning during the adaptation epoch at two different time points: early (first fifty trials

during adaptation) and late (last fifty trials during adaptation; **Fig. 5B**). A one-way ANOVA

revealed a significant effect of mean angle between the control, structural and gradual learning

groups during early learning [$F(2,57) = 14.4$, $P = 8.8$ e-06].

Post hoc comparisons using Tukey HSD tests indicated that the mean angle for the

structural learning group ($M = 24.9$, $SD = 2.9$, $P = 2.9$ e-05) and the gradual learning group ($M =$

24.5, $SD = 2.9$, $P = 1.4$ e-04) were reliably higher than the mean angle for the control group (M

$= 19.9$, $SD = 4.6$). However, there was no reliable difference detected between the structural and

gradual learning groups ($P = 0.9$). During late learning, we did not detect a reliable difference in

mean angle among the groups ($P = 0.2$). Therefore, the structural and gradual learning groups

demonstrated fast learning when countering an abrupt 30° CW rotation, as compared to session

one of the control group. While the control group represented naive learners, the prior experience

from session one for the structural and gradual learning groups is suggested to have facilitated

the improved learning. Likewise, this was observed in the control group, in which participants

experienced a repetition of an abrupt rotation and demonstrated savings during the second

session.

360   Recent work suggests that error sensitivity in sensorimotor adaptation is likely not

constant, but rather can vary depending on prior experience (13, 14, 16, 33). We modelled

movement angle across each session with the state-space equations proposed by Smith et al. (3),

and focused on changes in the retention and error sensitivity parameters. The main objective of

this study was to compare the model parameters across groups learning to counter the abrupt 30°

CW rotation. To do this, we used the bootstrap procedure previously reported by Coltman et al.

(15). In this manner, we always fit the model to averaged group data for each resampled

population (15, 31). The estimated posterior distributions of each of the four two-state model

parameter values are depicted in **Fig. 6** for sessions one and two of the control group and session

two of the gradual and structural learning groups. To determine whether the difference between

the mean of each distribution was statistically reliable, we calculated the distribution of the

differences in individual samples. The *insets* in **Fig. 6** show the distribution of differences found.

Table 1 shows the mean and standard deviation for each of the two-state parameters for each

group.

We first compared parameter estimates from session one and session two for the control

group (**Fig. 6A**). Across all comparisons made between groups, we did not observe a reliable

difference in the retention parameters for either the fast or slow process. When participants

experienced repetition of the same abrupt rotation, we found a statistically reliable increase in the

error sensitivity parameter for both the fast (B_f, $P = 0.007$) and the slow (B_s, $P = 0.003$)

processes. Importantly, this comparison allowed us to demonstrate that our previous finding from

a force field adaptation task (15) was replicated in a visuomotor rotation task. Therefore, this

result suggests that both the fast and slow processes are responsive to a history of error and both

contribute to savings.

Next, we compared parameter estimates from session one of the control group with

session two of the structural learning group (**Fig. 6B**). Based on the theory of structural learning,

thought to be essential to capturing the initial rapid phase of learning, Braun et al. (27)

demonstrated that the benefit of knowing the underlying structure of a task is that it leads to

facilitated adaptation. For this group we predicted that when later tested on an abrupt

perturbation, only the fast process would be affected by the initial training, as compared to the

control group. When overall learning is decomposed into a fast and slow state, the initial rapid

phase of learning is dominated by the output of the fast process. Therefore, we assumed that such

practice would influence the fast process. In addition to a statistically reliable increase in the

error sensitivity parameter for the fast process (B_f, $P < 0.001$), we also found a statistically

reliable increase in the slow process error sensitivity (B_s, $P = 0.002$).

Learning is believed to be more implicit in nature when a perturbation is gradually

applied using small undetectable increases, so that participants never encounter large sensory

prediction errors(22, 26, 34). By exposing a group of participants to a gradual perturbation

schedule during initial training, we predicted that only the slow process would be influenced.

When we compared the parameter estimates from session one of the control group with session

two of the gradual learning group (**Fig. 6C**) we found the gradual learning group showed a

statistically reliable increase in the error sensitivity parameter for both the fast (B_f, $P = 0.009$)

and the slow (B_s, $P = 2$ e-04) processes.

Lastly, we compared parameter estimates between the structural and gradual learning

groups during session two (**Fig. 6D**). Our goal was to use two different adaptation schedules

thought to differentially affect fast and slow learning processes and test the idea that error

sensitivity for each process would be independently modulated. We expected that the error

sensitivity parameter for the fast process would be greater in the structural learning group

compared to the gradual learning group, while the error sensitivity parameter for the slow

process would be greater in the gradual learning group compared to the structural learning group.

The only statistically reliable difference was in the error sensitivity parameter for the fast process

that was larger for the structural learning group (B_f, $P = 0.02$).

From the bootstrap distributions we calculated the mean value for each parameter for

session one and session two of the control group, and session two of the structural and gradual

learning groups separately. Using these mean estimated parameter values, we used the two-state

model to simulate our experimental paradigm and generate simulated learning curves to visualize

the time course of the estimated fast and slow processes, as well as the simulated overall output.

Figure 7 demonstrates that the simulated learning curves are qualitatively in good agreement

with the measured behavioural data. The models explains 98 - 99 % of the variance in angle over

the course of learning (*control session 1*: $R^2 = 0.98$, $P = 2.2$ e-04; *control session 2*: $R^2 = 0.99$, P

$= 1.2$ e-04; *structural session 2*: $R^2 = 0.98$, $P = 2.3$ e-04; *gradual session 2*: $R^2 = 0.99$, $P = 1.6$ e-

04). The model effectively captures the initial improvement in learning during the adaptation

epoch, the decay during the visual error clamp epoch, as well as the subsequent return towards

baseline performance during the washout epoch.

Discussion

The integration of different perturbation schedules and two-state modelling of measured

behavioral data allowed us to test the role of prior experience on error sensitivity modulation

during subsequent adaptation. The modelling of the data in turn describes adaptation as an

interaction between error-sensitivity and retention. It has previously been shown in the context of

force field learning that repetition of the same perturbation results in increased error sensitivity

for both the fast and slow processes of adaptation (15). We substantiated this here by

demonstrating that sensitivity to errors is similarly increased for both the fast and slow processes

during the second session of a visuomotor rotation task. We found no reliable differences in the

retention parameter across conditions and sessions.

The behavioural changes associated with savings suggest that some component of

memory from the initial training must lead to the faster relearning, but what is remembered and

recalled remains unclear. In the context of the present study, how the fast and slow processes

individually contribute to savings, is not well known. To address this point, we used different

perturbation schedules that relied on errors of different magnitudes to determine whether the

underlying processes of adaptation could be independently manipulated, and whether an

independent memory would subsequently be formed. We expected to see differences in error

sensitivity depending on the type of prior training participants had received and therefore

compared the model parameter estimates of participants in the gradual and structural learning

groups to the first session of the control group, but we found that error sensitivity of both the fast

and slow processes was increased for both groups. Such a result might suggest that sensitivity to

error during visuomotor adaptation is modulated by abrupt, gradual and random perturbation

schedules.

As an alternative account, savings has previously been explained by the retrieval of

previous successful actions, reflecting the use of an explicit strategy (18-20, 23). Within the

framework of a two-state model, this theory suggests that savings is driven purely by the fast

process, without consideration of the contributions from the slow process (35). Several

researchers have argued that explicit cognitive strategies can account for a significant amount of

learning, particularly during the early phase of learning and relearning (36-38). The dissociation

of learning into implicit and explicit learning processes often relies on the use of verbal aiming

reports prior to reaching (18, 20, 23, 38). Recent findings, however, indicate that verbal aiming

reports could lead to an overestimated explicit contribution to adaptation (21, 39). In fact, Leow

et al. (21) demonstrated that the use of shortened preparation time, designed to prevent strategic

re-aiming, resulted in the estimated implicit learning being larger than that which was obtained

from verbal reports. Furthermore, Yin and Wei (22) provide supporting evidence that savings of

motor adaptation is possible without forming or recalling a cognitive strategy with the use of a

gradually introduced visuomotor rotation during initial learning. If savings is possible, with and

without an explicit strategy being formed during initial learning and predominant measures of

implicit and explicit processes may be confounding their mode of measurement, how reliable are

the findings suggesting savings is driven exclusively by an explicit process?

The debate about the contributions of explicit/implicit and fast/slow processes to savings

stems from a recent proposal that fast and slow processes reflect explicit and implicit learning

mechanisms, respectively (35). Motor memory development is thought to be based on two

components: recall, which involves retrieving past motor movements, and faster relearning,

which involves increased sensitivity to errors (3, 40). By using the framework of a two-state

model we focused on the dominant component of adaptation, which is sensory prediction error,

and found that motor memory was associated with an increase in error sensitivity. In addition to

sensory prediction error, there are other possible teaching cues that might drive adaptation. When

researchers are focused on the more cognitive aspects of learning, exploring the use of an explicit

strategy, the dominant component of adaptation may likely be the reinforcement of successful

actions. Future research may shed light on this debate by probing both proposed methods of dual

processing simultaneously during learning, and by assessing more directly their shared features.

Another long-standing question is how quickly implicit changes in learning emerge.

Huberdeau and colleagues (20) demonstrated that learning of an abrupt perturbation with only a

few trials is sufficient to cause savings via the explicit process, based on the belief that the fast

learning is too short for implicit learning to take its full effect. Ruttle et al. (41) however recently

confronted the long standing notion that implicit learning is slowly developing, typically

unfolding over tens of trials. By observing changes in both internal models and state estimates of

limb position as a characterization of implicit learning, they found that after only one to three

perturbed training trials participants had changes in both reach aftereffects and a shift in hand

localization. Taking this into account, it seems possible that the 6-trial repetition used in the

structural learning task, aiming to influence the fast process, may have simultaneously influenced

the slow process. For that reason, it is possible that a common component of all three

perturbation schedules used during initial training was that the slow process accounted for a

significant portion of the error reduction.

Albert et al. (33) recently investigated the persistence of residual errors during motor

adaptation in the context of implicit and explicit learning systems. Of importance to the present

study, they propose that it is the implicit learning system which maintains a history for prior

errors. Our results are consistent with this hypothesis that it is the implicit process that stored

some component of prior training. Given the suggestion that the history of errors is stored by

only one of the two proposed underlying processes, this finding would be lost if learning

behavior was represented using a single-state model. Nevertheless, one may ask whether a two-

state model was necessary to represent learning in the behavioral tasks tested in the present

study. To address this, we calculated AIC values for both single- and two-state models fits to the

behavioral data. We used the data associated with the four sessions of abrupt rotations (i.e., the

first and second sessions of the control group, and the second session of the gradual and

structural learning groups) and for each we estimated the overall output based on a single-state

model and separately using a two-state model. Based on the model with the lowest AIC value, in

all four cases the best-fit model describing the measured behavioural data was the two-state

model. As a follow-up to our initial question, we would further suggest that the stored memory is

accessible to both processes during subsequent learning. As it pertains to our findings, we would

argue that while the fast process may not maintain a history or errors, it does have access to this

information in subsequent learning as evident by the increased error-sensitivity for the fast

process during testing in all groups.

Alternatively, while the experimental design and two-state model, used in the present

study, account well for the results of savings, recent work looking at evoked recovery (42) posits

that memory formation is related to the storing of information about the dynamical and sensory

features of the environment is related to the context with which it is associated. Understanding

how contextual inference can be related to and accessed by each process of the two-state model

can shed light on future discussions about multiple processes underlying motor learning.

Table 1. *Two-state model parameters calculated from probability distribution*

	Fast Process		Slow Process	
	A (mean ± SD)	*B* (mean ± SD)	*A* (mean ± SD)	*B* (mean ± SD)
Control Session 1	0.86 (0.02)	0.17 (0.02)	0.996 (7e-04)	0.05 (0.006)
Control Session 2	0.87 (0.03)	0.28 (0.04)	0.994 (0.001)	0.07 (0.01)
Structural Session 2	0.80 (0.05)	0.40 (0.03)	0.993 (0.002)	0.10 (0.02)
Gradual Session 2	0.87 (0.02)	0.27 (0.04)	0.994 (0.001)	0.10 (0.02)

515
516

Figure Captions

Figure 1. Experimental design and perturbation schedule. The experiment was divided into two sessions, separated by a 5-min session break. Each session consisted of four blocks: 1) a baseline period of no rotation trials, 2) an adaptation period, 3) an error clamp period, and 4) a washout period. Participants were randomly assigned to one of three groups which differed in session one during the adaptation period: abrupt control group, gradual learning group, or structural learning group.

Figure 2. Hand trajectories from a representative participant in the control group during both session one (light blue) and two (dark blue), and one representative participant per group in session two of the structural (purple) and gradual (pink) learning groups. Baseline reaches were from the last three trials (from trial 68 to trial 70) during the baseline epoch. Early and late adaptation reaches were from the first (from trial 71 to 73) and last (from trial 368 to trial 370) three trials of the adaptation epoch, respectively. Participants saw a random ordering of the three possible targets (represented by the squares).

Figure 3. Control group. *A*: the average angle at peak velocity for all trials in *session 1* (light blue) and *session 2* (dark blue). The shaded region denotes ± SE. *B*: comparisons between *session 1* and *session 2* (dark blue) for the mean angle for the first 50 (early) and last 50 (late) trials of the adaptation epoch. Circles represent individual data.

Figure 4. Gradual and structural learning groups. *A*: the average angle at peak velocity for all trials in *session 1* of the gradual learning group. The shaded region denotes ± SE. *B*: the data from four representative individual participants (S2, *top left*, S8 *top right*, S18 *bottom left*, S20 *bottom right*).

Figure 5. *A*: the average angle at peak velocity for all trials in *session 1* for the control group (light blue) and *session 2* for the structural (purple) and gradual (pink) learning groups. The shaded region denotes ± SE. *B*: comparisons between groups for the mean angle for the first fifty (early) and last 50 (late) trials of the adaptation epoch. Circles represent individual data.

Figure 6. Probability distribution of the model parameters given the data. Light blue and dark blue represent *session 1* and *session 2* of the control group, respectively. Purple represents *session 2* of the structural learning group and Pink represents *session 2* of the gradual learning group. *Inset* represents the distribution of pairwise differences. The four model parameters of the two-state model are fast retention (A_f), slow retention (A_s), fast learning rate (B_f), and slow learning rate (B_s).

Figure 7. Model simulations. Parameter estimates for each session were based on the mean values from the bootstrap distributions shown in Fig. 6. The four parameters of the model are fast retention (Af), fast learning rate (Bf), slow retention (As), and slow learning rate (Bs).

References

1.   **Donchin O, Francis JT, and Shadmehr R**. Quantifying generalization from trial-by-trial behavior of adaptive systems that learn with basis functions: theory and experiments in human motor control. *Journal of Neuroscience* 23: 9032-9045, 2003.

2.   **Miall RC, and Wolpert DM**. Forward models for physiological motor control. *Neural networks* 9: 1265-1279, 1996.

3.   **Smith MA, Ghazizadeh A, and Shadmehr R**. Interacting adaptive processes with different timescales underlie short-term motor learning. *PLoS Biol* 4: e179, 2006.

4.   **Thoroughman KA, and Shadmehr R**. Learning of action through adaptive combination of motor primitives. *Nature* 407: 742-747, 2000.

5.   **Wolpert DM, and Kawato M**. Multiple paired forward and inverse models for motor control. *Neural networks* 11: 1317-1329, 1998.

6.   **Kording KP, Tenenbaum JB, and Shadmehr R**. The dynamics of memory as a consequence of optimal adaptation to a changing body. *Nature neuroscience* 10: 779-786, 2007.

7.   **Lee J-Y, and Schweighofer N**. Dual adaptation supports a parallel architecture of motor memory. *Journal of Neuroscience* 29: 10396-10404, 2009.

8.   **Coltman SK, and Gribble PL**. Time course of changes in the long-latency feedback response parallels the fast process of short-term motor adaptation. *J Neurophysiol* 124: 388-399, 2020.

9.   **Kim S, Ogawa K, Lv J, Schweighofer N, and Imamizu H**. Neural substrates related to motor memory with multiple timescales in sensorimotor adaptation. *PLoS biology* 13: e1002312, 2015.

10.   **Sarwary AM, Wischnewski M, Schutter DJ, Selen LP, and Medendorp WP**. Corticospinal correlates of fast and slow adaptive processes in motor learning. *Journal of neurophysiology* 120: 2011-2019, 2018.

11.   **Ebbinghaus H**. Memory: A Contribution to Experimental Psychology, eds. *HA Ruger and CE Bussenius (Trans)(Dover Publications, New York, 1964)(Original Work Published 1885)* 1913.

12.   **van Beers RJ**. Motor learning is optimally tuned to the properties of motor noise. *Neuron* 63: 406-417, 2009.

13.   **Wei K, and Kording K**. Relevance of error: what drives motor adaptation? *Journal of neurophysiology* 101: 655-664, 2009.

14.   **Marko MK, Haith AM, Harran MD, and Shadmehr R**. Sensitivity to prediction error in reach adaptation. *Journal of neurophysiology* 108: 1752-1763, 2012.

15.   **Coltman SK, Cashaback JG, and Gribble PL**. Both fast and slow learning processes contribute to savings following sensorimotor adaptation. *Journal of neurophysiology* 121: 1575-1583, 2019.

16.   **Herzfeld DJ, Vaswani PA, Marko MK, and Shadmehr R**. A memory of errors in sensorimotor learning. *Science* 345: 1349-1353, 2014.

17.   **Leow L-A, De Rugy A, Marinovic W, Riek S, and Carroll TJ**. Savings for visuomotor adaptation require prior history of error, not prior repetition of successful actions. *Journal of neurophysiology* 116: 1603-1614, 2016.

18.   **Avraham G, Morehead JR, Kim HE, and Ivry RB**. Reexposure to a sensorimotor perturbation produces opposite effects on explicit and implicit learning processes. *PLoS biology* 19: e3001147, 2021.

19.   **Huang VS, Haith A, Mazzoni P, and Krakauer JW**. Rethinking motor learning and
savings in adaptation paradigms: model-free memory for successful actions combines with
internal models. *Neuron* 70: 787-801, 2011.
20.   **Huberdeau DM, Haith AM, and Krakauer JW**. Formation of a long-term memory for
visuomotor adaptation following only a few trials of practice. *Journal of neurophysiology* 114:
969-977, 2015.
21.   **Leow L-A, Gunn R, Marinovic W, and Carroll TJ**. Estimating the implicit component
of visuomotor rotation learning by constraining movement preparation time. *Journal of
neurophysiology* 118: 666-676, 2017.
22.   **Yin C, and Wei K**. Savings in sensorimotor adaptation without an explicit strategy.
Journal of neurophysiology 123: 1180-1192, 2020.
23.   **Morehead JR, Qasim SE, Crossley MJ, and Ivry R**. Savings upon re-aiming in
visuomotor adaptation. *Journal of neuroscience* 35: 14386-14396, 2015.
24.   **Hanajima R, Shadmehr R, Ohminami S, Tsutsumi R, Shirota Y, Shimizu T, Tanaka
N, Terao Y, Tsuji S, and Ugawa Y**. Modulation of error-sensitivity during a prism adaptation
task in people with cerebellar degeneration. *Journal of neurophysiology* 114: 2460-2471, 2015.
25.   **Kim HE, Morehead JR, Parvin DE, Moazzezi R, and Ivry RB**. Invariant errors reveal
limitations in motor correction rather than constraints on error sensitivity. *Communications
Biology* 1: 1-7, 2018.
26.   **Orban de Xivry J-J, and Lefèvre P**. Formation of model-free motor memories during
motor adaptation depends on perturbation schedule. *Journal of neurophysiology* 113: 2733-2741,
2015.
27.   **Braun DA, Aertsen A, Wolpert DM, and Mehring C**. Motor task variation induces
structural learning. *Current Biology* 19: 352-357, 2009.
28.   **Braun DA, Mehring C, and Wolpert DM**. Structure learning in action. *Behavioural
brain research* 206: 157-165, 2010.
29.   **Bond KM, and Taylor JA**. Structural learning in a visuomotor adaptation task is
explicitly accessible. *Eneuro* 4: 2017.
30.   **Scheidt RA, Dingwell JB, and Mussa-Ivaldi FA**. Learning to move amid uncertainty.
Journal of neurophysiology 86: 971-985, 2001.
31.   **Albert ST, and Shadmehr R**. Estimating properties of the fast and slow adaptive
processes during sensorimotor adaptation. *Journal of Neurophysiology* 119: 1367-1393, 2018.
32.   **Holm S**. A simple sequentially rejective multiple test procedure. *Scandinavian journal of
statistics* 65-70, 1979.
33.   **Albert ST, Jang J, Sheahan HR, Teunissen L, Vandevoorde K, Herzfeld DJ, and
Shadmehr R**. An implicit memory of errors limits human sensorimotor adaptation. *Nature
Human Behaviour* 2021.
34.   **Criscimagna-Hemminger SE, Bastian AJ, and Shadmehr R**. Size of error affects
cerebellar contributions to motor learning. *Journal of neurophysiology* 103: 2275-2284, 2010.
35.   **McDougle SD, Bond KM, and Taylor JA**. Explicit and implicit processes constitute the
fast and slow processes of sensorimotor learning. *Journal of Neuroscience* 35: 9568-9579, 2015.
36.   **Mazzoni P, and Krakauer JW**. An implicit plan overrides an explicit strategy during
visuomotor adaptation. *Journal of neuroscience* 26: 3642-3645, 2006.
37.   **Taylor JA, and Ivry RB**. Flexible cognitive strategies during motor learning. *PLoS
computational biology* 7: e1001096, 2011.

38.   **Taylor JA, Krakauer JW, and Ivry RB**. Explicit and implicit contributions to learning
in a sensorimotor adaptation task. *Journal of Neuroscience* 34: 3023-3032, 2014.
39.   **de Brouwer AJ, Albaghdadi M, Flanagan JR, and Gallivan JP**. Using gaze behavior
to parcellate the explicit and implicit contributions to visuomotor learning. *Journal of*
neurophysiology 120: 1602-1615, 2018.
40.   **Mawase F, Bar-Haim S, and Shmuelof L**. Formation of Long-term Locomotor
memories is associated with functional connectivity changes in the cerebellar–thalamic–cortical
network. *Journal of Neuroscience* 37: 349-361, 2017.
41.   **Ruttle JE, Marius't Hart B, and Henriques DY**. Implicit motor learning within three
trials. *Scientific Reports* 11: 1-11, 2021.
42.   **Heald J, Lengyel M, and Wolpert D**. Contextual inference underlies the learning of
sensorimotor repertoires. *bioRxiv* 2020.
664

A

Perturbation
Session 1
Session 2

Early

Late

B

p = 0.000003

Session 1
Individual Data
Session 2
Individual Data

Early   Late

Adaptation Epoch