Computational fluid dynamics provide evidence for a compensatory suction

feeding like effect in the predatory strike of dragonfly larvae

Alexander Koehnsen[1], Martin Brede[2], Stanislav N Gorb[1] & Sebastian Büsse[1]*

[1] Functional Morphology and Biomechanics, Institute of Zoology, Kiel University, Germany

[2] Chair for Fluid Mechanics, University of Rostock, Germany

*corresponding author: sbuesse@zoologie.uni-kiel.de

Abstract

Most fast-moving aquatic predators face the challenge of bow wave formation. Water in front of the predator alarms or even displaces the prey. To mitigate the formation of such a bow wave, a strategy aiming at pressure reduction via suction has evolved convergently in several animal groups: compensatory suction feeding. The aquatic larvae of dragonflies and damselflies (Insecta: Odonata) are likely to face this challenge as well. They capture prey underwater using a fast-moving raptorial appendage, the so-called prehensile labial mask. Within dragonflies (Odonata: Anisoptera) two basic shapes of the prehensile labial mask have evolved, with an either flat and slender or concave distal segment. While the former is a pure grasping device, the latter is also capable of scooping up smaller prey and retaining it inside the cavity by arrays of bristle-like structures. The hydrodynamics of the prehensile labial mask was previously unknown. We used computational fluid dynamic (CFD) simulations of the distal segment of the mask, to investigate for the first time how different shapes of the mask impact their function. Our results suggest that both shapes are highly streamlined and generate a low-pressure area, likely leading to an effect analogous to compensatory suction feeding. This presents a vivid example of how convergent evolution enables very different animal groups to successfully deal with the challenges of their environment. It may also be an interesting concept for technical application in small scale grasping devices; e.g. for simple sampling mechanisms in small-sized autonomous underwater vehicles (µAUVs).

Keywords: convergent evolution, numeric flow analysis, ecomorphology, biomechanics, underwater grasping, Anisoptera, Insecta, Odonata

Abbreviations

α: angle of attack; ρ: density; μ: dynamic viscosity; C_{dw}: drag coefficient; C_{lw}: lift coefficient; F_d drag force; F_l lift force; PLM: prehensile labial mask; R: Reynolds number; v: Velocity;

Introduction

"Everything was an artefact of its function. That was what made evolution so gorgeous." (Corey 2014 p. 64). In this study, we explored the relationship between the form of a prey capturing device in dragonfly larvae and their hydrodynamic implications - which in the end, represents a paragon for convergent evolution.

Most aquatic predators face a challenge when it comes to prey capturing: fast underwater motion creates a bow wave in front of the predators head, displacing or alarming prey before capturing (Lauder and Prendergast 1992; Alfaro 2002; Holzman and Wainwright 2009).

The hydrodynamics of other underwater feeding and/or prey-capturing mechanisms have been studied primarily in vertebrates (Wainwright et al. 2015): for example snapping turtles (Lauder and Prendergast 1992), amphibious snakes (Alfaro, 2002) and especially fish (e.g. Edmonds et al. 2001; Yaniv et al. 2014; Van Wassenbergh 2015). Several strategies have evolved in these groups to mitigate bow wave formation. For example, many fish species rely on suction feeding (Muller and Osse 1984; Edmonds et al. 2001; Holzman and Wainwright 2009; Yaniv et al. 2014; Van Wassenbergh 2015) and some aquatic snakes use a lateral head movement (Young 1991). However, frogs (Carroño and Nishikawa 2010), snapping turtles (Lauder and Prendergast 1992) and some snakes project their head directly towards the prey (Alfaro 2002) potentially causing bow wave formation. This problem is overcome in these groups by a mechanism called compensatory suction feeding. For example, snapping turtles quickly open their mouth while projecting their head towards a target (Lauder and Prendergast 1992; Summers et al. 1998). By flooding the oral cavity, water is drawn from the bulk which is amassing in front of the accelerating head (Lauder and Prendergast 1992; Summers et al. 1998). The mechanism thereby avoids the formation of a bow wave, allowing for successful prey capturing (Lauder and Prendergast 1992; Summers et al. 1998). As this issue gets more difficult with smaller scale and smaller prey items/particles (China and Holzman 2014), it may be a factor influencing the prey capturing in smaller invertebrates as well. In aquatic arthropods, the hydrodynamics of the predatory strike of mantis shrimps and snapping shrimps have been investigated (McHenry et al. 2016; Koukouvinis et al. 2017), but these studies did not focus on the interaction between predator and prey item so far.

In the larvae of damselflies and dragonflies (Insecta: Odonata), a unique prey capturing mechanism has evolved: one mouthpart, the labium, is modified into a prehensile mask (PLM) (see figure 1); capable of rapid projection for prey capturing (Corbet, 1957; Pritchard, 1965; Olesen, 1979; Büsse et al. 2021). The prey spectra between larvae with these two different ecotypes of PLM differ significantly. The concave PLM can scoop up smaller prey items, which are trapped inside the premental cavity and then lead to the mouth for feeding (Pritchard 1964, 1965). The sole function of

the flat PLM is grasping relatively large prey with hook-like labial palps. (Tillyard 1917; Pritchard

1965).

The PLM in general is capable of rapid protraction (Pritchard 1965; Olesen 1979; Tanaka and Hisada

1980). Powered by a dual catapult mechanism, maximum accelerations of 20g and maximum

velocities of around 1 m/s have been measured (Büsse et al. 2021). At high speeds, the influence of

hydrodynamic forces on the movement becomes increasingly important, as substantial amounts of

drag incur (Vogel 1997; McHenry et al. 2016;). Yet, the hydrodynamics of the PLM have never been

studied previously. A few publications hypothesise about the impact of hydrodynamics on the

predatory strike in their discussion (Tanaka and Hisada 1980; Quenta Herrera et al. 2018), but to the

best of our knowledge, no study has put a focus on this specific topic.

In this study, we aimed to assess the two described premental shapes from a hydrodynamic point of

view. Using high-speed videography, we assessed the kinematics of the prementum in detail and

estimated the angle of attack during the strike for species of both ecotypes. This data was combined

with 3D models obtained from micro-computed tomography (µCT) and used to generate

computational fluid dynamics (CFD) simulations to resolve flow characteristics. With this approach,

we intend to shed light on the question of how the two differently shaped prementa differ in their

flow patterns, and how these differences might impact the respective feeding strategies.

Additionally, we investigated whether the PLM features a mechanism to mitigate bow wave

formation in front of the prey capturing device (Lauder and Prendergast, 1992; Alfaro 2002). This

aims to further understanding how simple, small-scale underwater prey capturing/grasping devices

function in general which in turn contributes to answering questions about the evolution of two

different shapes within Anisoptera. Additionally, the results of this study might be interesting for

potential technical applications (e.g. sample collection) by small-sized autonomous underwater

vehicles (µAUVs) (Shahinpoor 1992; Kim et al. 2005; Shi et al. 2010).

Material & Methods

Specimens

Larvae of *Anax imperator* LEACH, 1815 (Anisoptera: Aeshnidae) were collected in aquaculture ponds

in Oeversee, Schleswig-Holstein, Germany. Specimens of *Sympetrum* sp. (Anisoptera: Libellulidae)

were collected in a pond near Knoop, Schleswig-Holstein, Germany. All specimens were collected

with permission of the '*Landesamt für Landwirtschaft, Umwelt und ländliche Räume Schleswig-*

Holstein'. Specimens were kept in individual aquariums at 8°C to prevent premature emergence and

fed with chironomid larvae (Diptera: Chironomidae). Before the experiments, the specimens were

acclimatised to room temperature. Head and body length were measured as a scaling reference.

High-speed Videography and Motion Tracking

7 predatory strikes of 5 specimens of *Anax imperator* (Odonata: Anisoptera) and 6 strikes of 5

specimens of *Sympetrum* sp. (Odonata: Anisoptera) were recorded in a small aquarium (210 x 100 x

105 mm) using chironomid larvae as prey items. Recordings were made with a Photron Fastcam

SA1.1 (model 675K-M1; Photron, Pfullingen, Germany) equipped with a 105 mm/1:2.8 macro lens

(Sigma, Tokyo, Japan) mounted on a Manfrotto-055 tripod with Manfrotto-410 geared head

(Manfrotto, Spa, Italy). Illumination was provided using two Dedocol COOLT3 halogen spotlights

(Dedotec, Berikon, Switzerland). The footage was captured with a resolution of 1024 x 1024 pixels at

5400 frames per second and saved as a 16-Bit TIFF image stack. For further analysis, the prementum

was tracked using Adobe After Effects CS6 (Adobe Systems Software, San José, CA, USA) according to

Koehnsen et al. (2020). One Tracker (Tracker A) was placed at the prementum-postmentum joint and

a second tracker (Tracker B) was placed at the joint between prementum and labial palps

(highlighted in figure 1 and supplementary figure 1).

To validate the simulations, we additionally recorded two specimens of each species (two strikes per

specimen) with the previous setup. To visualize flow patterns, neutrally buoyant *Spirulina* algae were

added to the water and a light source was placed behind the aquarium. Adjusting the focal plane to

the centre line of the larva and using a shallow focal plane (fstop = 4.0), allowed only particles

flowing around the approximate centre line of the PLM to be in focus.

Data Analysis and Statistics

The tangential velocity was determined directly from the location data of tracker A, which is located

at the centre of rotation of the prementum. To assess the rotational velocity, a vector between the

two tracking points was established (AB). These points are also highlighted in figure 1 and

supplementary material 1. The angle of this vector at the timepoint t was compared to the

orientation of the vector at the beginning of the motion sequence (t_0). In both cases, data were fitted

using a polynomial regression with 11 degrees of freedom. Velocity was determined from the first

derivative. The tangential velocity was calculated from the rotational velocity with the length of the

prementum (without labial palps) as the radius r.

To calculate the angle of attack (α), we compared the orientation of the prementum (which is given

by the vector AB) to a second vector indicating the direction of movement. This vector is formed by

point B (the base of the prementum) and the location of point B five frames earlier, indicating the

direction of motion. α is the angle between these two vectors. The location of all points used is also

shown in supplementary figure S1. To account for the temporal shift between α and the directly

measured curve, α has been calculated as a moving average: once with the aforementioned

reference point, five frames earlier and once with a reference point five frames later. For each

timepoint, we used the mean of the two values.

We extracted data from the curve at ten standardised timepoints for CFD simulations. Initially, the

point of maximum total velocity (translational + tangential velocity) and the start and endpoint of

motion were determined, dividing the strike into an acceleration and deceleration phase. Six equally

spaced points (including starting point and point of maximum velocity) were used for the

acceleration phase, and four equally spaced points (including the endpoint) were used for the

deceleration phase.

All statistical analyses were performed in R Studio (v1.2.5001, the R Foundation for Statistical

Computing, Vienna, Austria). For the comparison of mean velocity and α, a Wilcoxon Rank Sum Test

was used, as the sample size was not equal. If not stated otherwise, all given values are median ±

interquartile range (IQR).

The Reynolds number of the prementum was calculated for both species according to the following

formula:

$$R = \frac{\rho\, v\, d}{\eta}. \tag{1}$$

with the maximum width of the prementum (without palps) as characteristic length (d). The velocity

(v) was determined from tracking data using the total velocity at the tip. Dynamic viscosity (η) and

density (ρ) were taken from literature data of freshwater at 25°C (Rumble et al. 2017) see

supplementary table S2).

The drag coefficient C_{dw} and lift coefficient C_{lw} was calculated for both species according to the

following formulas for drag coefficient and lift coefficient, respectively:

$$C_{dw} = \frac{2F_d}{\rho v^2 A_w} \tag{2}$$

$$C_{lw} = \frac{2F_l}{\rho v^2 A_w}. \tag{3}$$

As α (and thereby the frontal area) of the prementum differs over the duration of the predatory

strike, we used the total wetted area A_w as a reference area for the calculation of C_{dw} /C_{lw}.

Measurement of the reference area was based on the 3D models used for CFD simulation

(supplementary figure S4). The wetted area A_w was used instead of the more common projected

frontal area A_w, as the angle of attack, and thereby the projected frontal area would vary between

simulations, complicating the interpretation of the results. F_d represents the drag force obtained

from the CFD simulations, while the input value for velocity (v) was taken from tracking data at

maximum velocity and is identical to the data used for CFD simulation. The density (ρ) was taken

from literature data of freshwater at 25°C (supplementary table S2).

Micro-computed tomography (µCT)

Micro-computed tomography scans were used to obtain 3D models for CFD analysis. Therefore

specimens (n=1 for each species) were fixed in alcoholic Bouin solution (= Duboscq-Brasil) for tissue

preservation and subsequently stored in 70% ethanol. Samples were critical-point dried before

scanning (Leica EM CPD300 automated critical point dryer, Leica, Wetzlar, Germany) and scanned

with a SkyScan 1172 (Bruker micro-CT, Kontich, Belgium) with high-resolution settings (40 kV, 250 µA

and 0.25° rotation steps, performing a 360° scan). Segmentation and processing of the µCT-data

were carried out with Amira 6.0.1 (Thermo Fischer Scientific, Waltham, USA). 3D datasets were

retopologised, reoriented and rescaled using Blender (v2.8 Blender Foundation, Amsterdam,

Netherlands, www.blender.org). The prementum was scaled based on the average size of all

specimens (of the respective species) that were used for the high-speed video experiments.

Scanning Electron Microscopy (SEM)

Two specimens (n=2) of *Sympetrum* sp. were prepared with extended PLM using an expendable

custom-built brace (supplementary figure S5). One specimen with opened labial palps, one with

closed ones. The brace was designed using blender (v2.8, Blender Foundation, Amsterdam,

Netherlands, www.blender.org) and printed on a Prusa i3 Mk3 FDM 3D printer (Prusa Research s.r.o.,

Prague, Czech Republic) using polylactic acid filament (Prusament, Prusa Research s.r.o., Prague,

Czech Republic) due to its stability in ethanol. The samples were critical-point dried (Leica EM

CPD300, Leica, Wetzlar, Germany) and coated with a 10nm gold-palladium layer to (Leica Bal-TEC

SCD500, Leica, Wetzlar, Germany). We mounted the samples using a rotatable sample holder (Pohl

2010) and examined them using a Hitachi TM3000 (Hitachi Ltd., Tokyo, Japan) scanning electron

microscope with an acceleration voltage of 15kV. Obtained images were stitched into a single image

using Affinity Photo (Serif Europe Ltd. West Brigford, UK, https://affinity.serif.com).

Computational fluid dynamics

Meshing

Meshes were created using ICEM (2019 R2, Ansys Inc., Canonsburg Pa, USA). Models of the

prementum of *Anax* and *Sympetrum* were imported as '.stl files'. We oriented the model according to

the specified α and created a domain around the model with five times the model size as head and

ten times the model size as trail region. Three times the model size was used as the clearance

between model and lateral as well as top and bottom domain walls (see below for wall boundary

conditions).

We created unstructured tetrahedral meshes with ~4.6 million elements and 5 prism layers around

the model surface (maximum $y+$ was below 0.69 in all simulations for *Sympetrum* and below 2 in all

simulations for *Anax*). To increase mesh resolution around the model a density field was used.

Doubling the element count further changed the simulation results by <3%, therefore mesh

resolution was considered sufficient for our purposes (supplementary table S3). An exemplary mesh

for both species is shown in supplementary figure S4.

Boundary conditions and solution

For simulation setup and solving, we used CFX Pre (2019 R2, Ansys Inc., Canonsburg Pa, USA) with

fluid domain properties set to the values of water at 25°C. As boundary conditions, we set up the

lateral as well as top and bottom walls to be free slip walls. The rear wall was defined as an opening

with no ambient pressure difference. The front wall was defined as an inlet, with normal speed

settings based on either the tracking data in case of the first simulated experiment or to 0.4m/s in

the case of the second experiment. The model of the prementum was set up as a smooth no-slip

wall. We approximated turbulence using the κε turbulence model with ANSYS default low turbulence

settings. Simulations were solved using the RANS Solver Ansys CFX (2019 R2, Ansys Inc., Canonsburg

PA, USA). We used RMS (root-mean-square) error as convergence criterium with a remaining error of

1×10^{-4}. Increasing the convergence criteria by one order of magnitude changed the simulation result

by less than 5%.

Simulation setup

We performed three virtual experiments using CFD Simulations: 1) We sampled 10 timepoints from

each recorded strike and used the estimated α and flow velocity as simulation input. We used this

time series of stationary simulations as a quasi-dynamic approximation to the unsteady motion of

the PLM. We also neglected the rotational motion, but instead used the sum of rotational velocity

(tangential velocity at the tip of the prementum) and translational velocity, as we were most

interested in the area around the premental tip, where contact with the prey occurs. 2) To assess the

impact of α on the observed flow patterns, we performed 10 stationary simulations of each

prementum at a flow velocity of 0.4m/s (which is around the maximum observed flow speed), and

varied α in ten steps between 0° and 70°. 3) We performed a scaling experiment at an α of 30° and

0.4 m/s flow velocity. Here we scaled down the prementum of *Anax* to the size of the prementum of

Sympetrum (and vice versa).

Results

Kinematics and angle of attack

We recorded high-speed video footage of the predatory strike in *Anax* (flat-shaped) and *Sympetrum*

(spoon-shaped) and calculated the translational velocity (based on the motion of the base of the

prementum) and the tangential velocity (based on the rotation of the tip around the base). We

neither found a significant difference in maximum translational velocity between *Anax* and

Sympetrum (Wilcox rank-sum test, p=0.35) nor maximum tangential velocity (Wilcox rank-sum test,

p=0.13). We calculated the angle of attack (α) based on tracking data: the angle of the prementum

relative to the flow direction. Figure 2 shows the α (a) and the total velocity (translational +

tangential velocity) (b) for both species at 10 sample points. The results suggest that the median α is

higher in *Sympetrum* at every timepoint, yet this difference is only statistically significant at three

timepoints (TP$_1$ p=0.038, TP$_3$ p=0.003, TP$_4$ p=0.003 Wilcox rank-sum test). The velocity at each

timepoint varied widely between individuals (Fig 2 B) and we found no statistically significant

difference in strike velocity between both species at any timepoint (0.2 > p < 0.95, Wilcox rank-sum

test). The median maximum velocity is around 0.45 +- 0.32 m/s in *Anax* and 0.43 +- 0.30 m/s in

Sympetrum. Figure 2 A+B suggests that α decreases with an increased velocity and increases again

towards the end of the strike as the PLM slows down.

Reynolds number

We estimated the Reynolds number for the prementum of both *Anax* and *Sympetrum* models based

on kinematic data with the width of the prementum (without labial palps) as reference length *r*. We

calculated a Reynolds number of ~3700 for *Anax* and ~1800 for *Sympetrum* (supplementary table S2).

Computational fluid dynamics

Total forces on the prementum

We performed a total of 46 computational fluid dynamics (CFD) simulations of the prementum of

both species. We performed stationary simulations for each aforementioned timepoint (Fig. 2) and

calculated the total drag force (parallel to the direction of flow) and the total lift force (orthogonal to

the direction of flow) acting on the prementum (cf. Force diagram in figure 3). Results are shown in

figure 3 (3A data on *Anax*, 3B data on *Sympetrum*). In both species, forces increase with velocity, yet

the maximum velocity and maximum drag/lift force do not necessarily occur at the same timepoint.

The magnitude of forces is about twice as high in *Anax* (max. F_{drag} of 1.26 mN in *Anax* and 0.56 mN in

Sympetrum. Max. F_{lift} of 1.38 mN in *Anax* and 0.50 mN in *Sympetrum*). This also shows that the PLM

generates substantial amounts of lift. Lift forces are almost as high or even exceed (in *Anax*) the

amount of occurring drag at the respective timepoint.

Drag and lift coefficients

With an average prementum length of about 10 mm, the prementum of *Anax* is about three times

the length of the prementum in *Sympetrum*, hence the occurring drag forces are hardly comparable.

We, therefore, calculated the drag coefficient C_{dw} and lift coefficient C_{lw} for each timepoint (as C_{dw}

and C_{lw} change with α). Results are shown in figures 3 C and D. The drag coefficients of both PLM

start to diverge with an increased velocity (with exception of timepoint 5). During the phases with

high velocity, the drag coefficient in *Anax* is about half as high as in *Sympetrum*. Drag coefficients

increase again towards the end of the strike when velocity decreases and α increases. In both cases,

we see a decrease in the lift coefficient (concordant with a decreasing *α*) over the course of the

strike. In *Sympetrum,* we even see negative lift at timepoint 8, where *α* is lowest.

Pressure and flow speed profiles

We plotted contours of both the pressure (Fig. 4) and total flow speed (Fig.5) of four exemplary

timepoints along the median line of the prementum. Plots for all timepoints can be found in

supplementary figure S6. In both species, we see a high-pressure area below the prementum (as

indicated by reddish colour) and a low-pressure region above it (as indicated by a yellowish colour).

These areas are less pronounced around the prementum of *Anax* which is concurrent with its lower

drag coefficient. In the steps with lower velocity (such as 3 and 9), we also see smaller pressure

differences. Additional details can be inferred from the contours of the flow speed (Fig. 5). We see a

higher flow speed in low-pressure areas and a decrease in high-pressure areas. In *Anax,* the flow

separation is visible at higher α, whereas this does not occur in *Sympetrum* (see Fig. 5 step 3+5). The

separation of the flow is indicated by the blue area on the dorsal side representing a slow-moving

"bubble", where the fluid has detached from the surface.

Estimating the optimal angle of attack

To assess whether the shape of the prementum of either species is optimised towards a certain angle

of attack (α), we ran CFD simulations of the prementum at a velocity of 0.4 m/s and varied α within a

range from 0° to 70° (Fig. 4 A). We calculated the drag (C_{dw}) and lift (C_{lw}) coefficients for each

simulation (Fig. 4). Our results suggest that the relationship between α and C_{dw}/C_{dw} is different for

the two PLM shapes. The lowest C_{dw} for the prementum of *Anax* occurs at α of 10° (C_{dw}= 0.035), while

the minimum C_{dw} for the prementum of *Sympetrum* occurs at 25° (C_{dw}= 0.091). Additionally, the drag

coefficient is more sensitive to changes in α in *Anax* than in *Sympetrum*. We found a significant

increasing monotonic relationship between α and C_{dw} in *Anax* (Spearman's rank correlation, r_s=0.99,

p<2.2e-16), which was not present in *Sympetrum* (Spearman's rank correlation, r_s =0.51, p=0.114).

While at α of 10°, the drag coefficient is twice as high in *Sympetrum* than in *Anax*, the drag

coefficients are equal at α=30° and the increase continues at a steeper slope in *Anax.* The lift

coefficient increases monotonically with an increasing α in both species (*Anax*: Spearman's rank

correlation, r_s=0.75, p=0.010, *Sympetrum*: Spearman's rank correlation, r_s=0.99, p<2.2e-16). Yet while

the prementum of *Anax* starts to generate positive lift (directed into the dorsal direction) between 0°

and 10°, we see positive lift generation in *Sympetrum* later, between α=25° and 30°. Yet, the lift

coefficient starts to decrease again in *Anax* at α values higher than 45°, while it continues to increase

in *Sympetrum* until the second to last data point (60°).

Analysis of flow patterns

We visualised contours of both the pressure and total flow speed around the median line of the

respective prementum. Results of four exemplary timepoints are shown in figure 7 (pressure) and

figure 8 (flow speed). Results of all simulations are shown in supplementary material S7. The shape

and size of low- and high-pressure regions are dependent on α and the species. In *Anax,* the pressure

distribution is neutral at α=10° and we see the aforementioned pattern from α=20° onwards (which

corresponds with a positive lift coefficient, as shown in figure 6). At high α, the growth curve of the

lift coefficient flattens and decreases again beyond α=50°. At this time, the ventral side of the

prementum mask is almost perpendicular to the flow direction, incurring high amounts of pressure

drag. This also happens in *Sympetrum*, but at higher α (> 60°) and on a lesser scale (due to the

pressure difference between the high-pressure region and the surrounding fluid). In *Sympetrum,* we

see that the tip of the prementum (also known as ligula, which is tilted downwards) causes fluid to

amass in the premental cavity and form a high-pressure region. This corresponds to the higher drag

coefficients (Fig. 6). At α=30° and beyond, we see a ventral high-pressure region and a dorsal low-

pressure region, the latter is formed inside the premental cavity and both contribute to lift

generation. The flow speed is visualised in figure 8. We can see the flow separation on the dorsal side

of the *Anax* prementum. The point of separation is already visible at α=30° and moves towards the

tip with an increasing α. Separation of the flow is indicated by the slowly flowing region (blue). Flow

separation occurs in *Sympetrum* as well, yet to a lesser degree and only at α>=70°.

Scale comparison

To assess the impact of the different Reynolds numbers on the shapes (and thereby on the

comparability) of the two differently shaped PLMs, we ran an additional simulation of timepoint 5 of

the previous experiment for both species. We scaled down the prementum of *Anax* to the size of the

prementum of *Sympetrum* and vice versa. The results are shown in supplementary figure S8. In both

species, changing the scale leads to an increase in both the drag- and the lift coefficients.

Distribution of setae in Sympetrum

Four arrays of setae cover the inside of the prementum and the labial palps (Fig. 9 A, B). Two arrays

on each side of the prementum are protruded towards the centre of the premental cavity and angled

forwards and downwards into the cavity. The other two arrays are located on the labial palps and

protruded towards the centre of the premental cavity. When the labial palps are closed, the setae

cover the premental cavity completely (Fig. 9 B), but they leave an opening during the strike when

the labial palps are open (Fig. 9 A).

Experimental validation

We performed four additional high-speed recordings with particles (algae) in the fluid to visualize

actual flow patterns (Fig. 10, supplementary video S9). The results are consistent with our simulation

outputs. We see a high-pressure region below the prementum, which is indicated by outflow in the

region, but no bow wave is forming in either species and particles are flowing past the palps

over/into the prementum. Furthermore, algae are visibly retained by the setae (missing particles in

figure 10) and some non-steady effects can be observed (supp. video S9). After the palps have closed,

the PLM slows down, leading to a flow out of the premental cavity due to fluid inertia and a

pronounced bow wave.

Discussion

Kinematics of the predatory strike

We derived translational and rotational velocity, in an attempt to approximate the flow speed at the

tip of the prementum as closely as possible. Our results showed no significant difference between

both species.

We assessed the angle of attack (α) of the prementum for the first time. α is a typical characteristic

of airfoils and defined as the angle between the chord line of the airfoil, and the direction of flow

(e.g. Gracey 1958). In a more general way, α can be understood as the orientation of a streamlined

object towards the direction of flow. In both species, we see a decrease in median α as velocity

increases. Yet we found a higher α in *Sympetrum*. The trend is visible over all timepoints (Fig. 2) and

the difference is statistically significant at three timepoints. In general, a higher α leads to an increase

in the area exposed to the flow, and drag (and lift) coefficients increase (Fig. 6) (Hoerner 1965; Gur et

al. 2010).

Characteristics of drag

The most relevant factors responsible for the amount of drag are the shape of an object, its size and

orientation, as well as the velocity of the surrounding water. The drag/lift curves (Fig. 3) roughly

reflect the velocity profiles (Fig. 2 B). Yet velocity changes alone do not completely explain the

observed results (Fig. 3): for example, we find the maximum amount of drag and lift in *Sympetrum* at

timepoint 5, whereas the maximum velocity occurs at timepoint 6. This can be explained by α

changing over time. Changing the orientation changes flow patterns around the object (given that

the object is not a sphere), which changes both the amount of drag, as well as the drag coefficient

(Hoerner, 1965; Vogel, 1997). At timepoint 4, α is higher, leading to a higher amount of drag.

To assess the relationship between drag and α, we tested different values of α (0°-70°) at the same

velocity (Fig. 6). The drag coefficients are in a range that would be expected for a flat, streamlined

body (Hoerner 1965; Vogel 1997). In *Anax,* it increases proportionally with α as more and more

surface area is exposed to the flow (Fig. 6). The prementum of *Sympetrum* shows a different pattern.

The drag coefficient is minimal at an α of 30°. In general, the drag coefficient of the prementum of

Sympetrum is less responsive to changes of α as in *Anax* (Fig. 6). This corresponds to the observation

of a higher α during the predatory strike in *Sympetrum* (Fig. 2). We assume that these two different

minima are due to the different shapes of the prementum. If we look at longitudinal sections of the

prementum (Supplementary figure S4 A, B), the prementum of *Anax* is roughly represents a straight

slender shape. Yet in *Sympetrum* the anterior part (ligula) is angled downwards. Hence it is more

parallel to the flow at higher α. According to the simulation, we would expect an α of around 10° in

Anax, when strike velocity is high, and around 30° in *Sympetrum*. However, our kinematic data (Fig.

2) shows that the measured α is higher in both species in vivo, hence more factors are to be

considered.

Both prementa generate substantial amounts of dynamic lift. According to the principle of

Bernoulli/Venturi, lift is the result of a pressure difference between the upper and lower side of an

object (Hoerner, 1965; Vogel, 1997). This occurs in both prementa (Figs. 4, 7) and the amount of lift is

dependent on α (Fig. 6). The prementum of *Sympetrum* requires a higher α to achieve the same lift

coefficient as *Anax* but surpasses the lift coefficient of *Anax* at high α. For both species, lift

generation is rather low for values of α, where the drag coefficient is minimal. Higher α values*,*

observed during the strike, could thereby be explained by the necessity to generate lift, which is

rather counterintuitive. One would expect the prementum to generate as little lift as possible. Either

positive, as well as negative lift, would result in vertical motion of the prementum relative to the

direction of travel, likely complicating prey targeting and also requiring additional energy. The

amount of lift generated, however, is substantial and the calculated lift forces are almost as high as

the drag force (Fig. 3). We hypothesise that this is to avoid the formation of a bow wave in front of

the projecting prey capturing device, which may displace the prey item (Lauder and Prendergast

1992; Van Damme and Aerts 1997; Alfaro 2002). This behaviour would serve an analogous function

to the compensatory suction feeding observed in vertebrates (Lauder and Prendergast 1992; Van

Damme and Aerts 1997; Alfaro 2002; Holzman and Wainwright 2009), yet by deflecting the flow

instead of expanding a cavity.

The findings are concurrent with the contours of pressure in figures 4 and 7. As the prey items,

Sympetrum feeds on, tend to be rather small (especially compared to the usual prey of aeshnids)

(Pritchard 1964), countering bow wave formation might be more important for capturing smaller

prey, which would explain the more pronounced pressure difference between premental cavity and

surrounding fluid.

The prementum of *Anax* is about three times longer than the prementum of *Sympetrum*. To compare

these two shapes, we calculated the dimensionless drag coefficient (C_{dw}) (formula 2, see Materials

and Methods). C_{dw} only depends on the shape and orientation of the object (as long as the density of

the medium does not change and the Reynolds numbers are comparable). The Reynolds number of

both movements is within a similar order of magnitude (~3700 in *Anax* vs ~1800 in *Sympetrum*).

These values are comparable to the Reynolds numbers occurring in the flight of larger insects (Vogel

1997; Goyens et al. 2015). We consider the flow sufficiently similar to make the previous

comparisons. Yet both drag and lift coefficients are functions of the Reynolds number (Hoerner 1965;

Vogel 1997). When changing the size of the prementum, the drag and lift coefficients increase

(supplementary figure S7). The increase of the drag coefficient in *Anax* might be explained by the fact

that the streamlining is less effective at the smaller scale because the amount of viscous drag is

higher (Vogel 1997). Vice versa, the less streamlined prementum of *Sympetrum* has a higher impact

at the larger scale, which causes the drag coefficient to increase as well. In conclusion, the PLM might

be tuned towards its respective flow regime. Comparing the PLM from different larval instars might

be an interesting approach to test these hypotheses in future research.

Finally, we can combine the information on setae distribution with the result of our CFD simulations

(Fig. 9). Our SEM images, which are concordant with earlier studies on libellulids (Tillyard 1917;

Olesen 1979; Gupta et al. 1992) show that setae are covering the top of the premental cavity. With

open labial palps, not the entire area over the prementum is covered, which would be anticipated for

an efficient capturing device. Considering the streamlines inside the premental cavity (Fig. 9), the

low-pressure area at the prementum might also serve to deflect water with prey items deeper into

the premental cavity, where they are trapped. Once the labial palps close, the premental cavity is

completely covered by the seta arrays on the prementum and labial palps and prey cannot escape

(Fig 9B).

Limitations of the model & Validation

In this study, we used several simplifications/approximations which must be addressed. First of all,

we used a quasi-dynamic approach, to replicate the flow patterns during the actual predatory strike

with a series of stationary simulations. This is likely to underestimate the total amount of drag: due

to the added mass of fluid that sticks to the prementum due to viscous forces (Van Wassenbergh et

al. 2008). Additionally, we neglected the rotational component of the motion of the prementum but

took an approach using the sum of translational and rotational velocity at the tip of the prementum.

This is likely to overestimate the actual drag, as the base of the prementum is moving slower than

the tip. Yet we were not interested in calculating the actual drag with high precision, but rather in

estimating the impact of the PLM shape on prey capturing. Therefore, we were most interested in

fluid motion at the tip of the prementum, the point of contact between the predator and prey, which

is simulated with the highest accuracy and calculating α for each timeframe ensured that the

direction of flow is correct. Finally, it is important to note that we did not consider the drag

generated by the postmentum, to reduce simulation complexity, as it is also located downstream

from the prementum and the prey item. These aspects have to be kept in mind when interpreting

the results.

To verify the predicted flow patterns, we observed predatory strikes with algae as flow particles (Fig.

10, supplementary video S9). The results corroborate our simulations concerning bow wave

prevention. Outflow indicates a high-pressure region below the prementum and we observed fluid

passing over/into the prementum without obstruction. Furthermore, these findings suggest that

inertial effects of the fluid play little role in the prey capture event. We only see the ejection of fluid

from the cavity due to deceleration of the PRM after the labial palps have closed around the prey,

indicating the deceleration effects play little role in prey capturing (supplementary video 9).

However, to fully analyse the flow field around the PLM, more advanced and expensive methods, like

PIV or MRI flow measurements, would be necessary for future projects.

Conclusion

We showed that both prementum types (in *Anax* and *Sympetrum*) are well streamlined and generate

a low-pressure zone, likely to mitigate the formation of a high-pressure area in front of the

prehensile labial mask, a process analogous in function to compensatory suction feeding. We also

verified this hypothesis experimentally, showing there is no visible bow wave formation.

Furthermore, the prementum of *Sympetrum* is optimised towards higher angles of attack, possibly

due to the angled ligula. Both prementa are tuned towards their specific Reynolds number, which

raises the question: How does the shape of the labium and flow patterns change over different sizes

like in larval development? Finally, we found that the prementum of *Sympetrum* deflects the water

flow above the prementum downwards, likely to push prey items into the setae-armoured trap

located on the prementum. Both prementa function in a similar way, despite their different shapes.

The deep premental cavity is likely an innovation to trap prey, and possibly to further stabilise the

low-pressure region above the prementum. Our study explores the fluid mechanics of prey capturing

in odonate larvae for the first time and explains how fluid mechanics might facilitate prey capturing

in these aquatic predators. These findings also show an example of convergent evolution, as this bow

wave mitigation strategy represents a functional analogue to compensatory suction feeding known

from vertebrates, like turtles, salamanders, and fish.

Furthermore, we show, how small scale technical underwater grasping devices for differently sized

objects may function, which could be an interesting starting point for biomimetic applications.

Ethics: No permission of research ethics or animal ethics committees was necessary. All applicable regulations concerning the protection of free-living species were followed

Data accessibility: Raw data can be obtained from the corresponding author upon request.

Authors' contributions: AK, MB, SNG and SB designed the project and developed the concept of the study. AK and SB performed the high-speed video recordings and analysed the footage. AK and MB designed the setup for the CFD simulation and evaluated the results. AK performed the CFD simulations and SEM observations. SB performed CT scanning of the specimens. AK wrote the manuscript. All authors read and edited the manuscript, agreed to the content therein and approved the final version of the manuscript.

Competing interests: We declare we have no competing interests.

Funding: The work was funded and SB was directly supported through the DFG grants BU3169/1-1 and BU3169/1-2.

Acknowledgements: Finally, we would like to thank everyone, who supported us with this study: T. Büscher and H. Tramsen for support with statistics and data processing. B. Josten and H.-L. Troeger for support with the high-speed video experiments. Furthermore, we would like to thank S. Bodenstein, A. Georg, H. Jeske and K. Lehmann for providing dragonfly larvae for the study, O. Mudimo for providing *Spirulina*, and L. Hindenberg & H. Poggemann for proofreading. All members of the Functional Morphology and Biomechanics group at Zoological Institute of Kiel University are also greatly acknowledged for their continuous support.

References

Alfaro, M.E., 2002. Forward attack modes of aquatic feeding garter snakes. Funct. Ecol. 16, 204–215. https://doi.org/10.1046/j.1365-2435.2002.00620.x

Blanke, A., Büsse, S., Machida, R., 2015. Coding characters from different life stages for phylogenetic reconstruction: A case study on dragonfly adults and larvae, including a description of the larval head anatomy of *Epiophlebia superstes* (Odonata: Epiophlebiidae). Zool. J. Linn. Soc. 174, 718–732. https://doi.org/10.1111/zoj.12258

Büsse, S., Gorb, S.N., 2018. Material composition of the mouthpart cuticle in a damselfly larva (Insecta: Odonata) and its biomechanical significance. R. Soc. Open Sci. 5.

https://doi.org/10.1098/rsos.172117

Büsse, S., Hörnschemeyer, T., Gorb, S.N., 2017. The head morphology of *Pyrrhosoma nymphula*
larvae (Odonata: Zygoptera) focusing on functional aspects of the mouthparts. Front. Zool. 14,
1–13. https://doi.org/10.1186/s12983-017-0209-x

Büsse, S., Koehnsen, A., Rajabi, H., Gorb, S.N., 2021. A controllable dual-catapult system inspired by
the biomechanics of the dragonfly larvae's predatory strike. Science Robotics, 6(50).
https://doi.org/10.1126/scirobotics.abc8170

Carroño, C.A., Nishikawa, K.C., 2010. Aquatic feeding in pipid frogs: The use of suction for prey
capture. J. Exp. Biol. 213, 2001–2008. https://doi.org/10.1242/jeb.043380

China, V., Holzman, R., 2014. Hydrodynamic starvation in first-feeding larval fishes. Proc. Natl. Acad.
Sci. U. S. A. 111, 8083–8088. https://doi.org/10.1073/pnas.1323205111

Corbet, P.S., 1957. The Life-History of the Emperor Dragonfly *Anax imperator* Leach (Odonata :
Aeshnidae). J. Anim. Ecol. 26, 1–69.

Corey, J.S.A., 2014. Cibola Burn. Orbit, London, UK.

Edmonds, M.A., Motta, P.J., Hueter, R.E., 2001. Food capture kinematics of the suction feeding horn
shark, *Heterodontus francisci*. Environ. Biol. Fishes 62, 415–427.
https://doi.org/10.1023/A:1012205518704

Goyens, J., Van Wassenbergh, S., Dirckx, J., Aerts, P., 2015. Cost of flight and the evolution of stag
beetle weaponry. J. R. Soc. Interface 12. https://doi.org/10.1098/rsif.2015.0222

Gracey, Wi., 1958. Summary of Methods of Measuring Angle of Attack on Aircraft. NACA Tech. Note.

Gupta, A., Dey, S., Gupta, S., 1992. Cuticular structures on the labium of the larva of *Crocothemis*
servilia (Drury)(Anisoptera: Libellulidae). Odonatologica 21, 97–101.

Gur, O., Mason, W.H., Schetz, J.A., 2010. Full-configuration drag estimation. J. Aircr. 47, 1356–1367.
https://doi.org/10.2514/1.47557

Hoerner, S.F., 1965. Fluid-Dynamic Drag. published by the author, Midland Park, NJ.

Holzman, R., Wainwright, P.C., 2009. How to surprise a copepod: Strike kinematics reduce
hydrodynamic disturbance and increase stealth of suction-feeding fish. Limnol. Oceanogr. 54,
2201–2212. https://doi.org/10.4319/lo.2009.54.6.2201

Kim, B., Kim, D.H., Jung, J., Park, J.O., 2005. A biomimetic undulatory tadpole robot using ionic
polymer-metal composite actuators. Smart Mater. Struct. 14, 1579–1585.

https://doi.org/10.1088/0964-1726/14/6/051

Koehnsen, A., Kambach, J., Büsse, S., 2020. Step by step and frame by frame - workflow for efficient

motion tracking of high-speed movements in animals. Zoology.

https://doi.org/https://doi.org/10.1016/j.zool.2020.125800

Koukouvinis, P., Bruecker, C., Gavaises, M., 2017. Unveiling the physical mechanism behind pistol

shrimp cavitation. Sci. Rep. 1–12. https://doi.org/10.1038/s41598-017-14312-0

Lauder, B.Y.G. V, Prendergast, T., 1992. Kinematics of Aquatic Prey Capture in the Snapping Turtle

Chelydra serpentina. J. Exp. Biol. 164, 55–78.

McHenry, M.J., Anderson, P.S.L., Van Wassenbergh, S., Matthews, D.G., Summers, A.P., Patek, S.N.,

2016. The comparative hydrodynamics of rapid rotation by predatory appendages. J. Exp. Biol.

219, 3399–3411. https://doi.org/10.1242/jeb.140590

Muller, M., Osse, J.W.M., 1984. Hydrodynamics of suction feeding in fish. Trans. Zool. Soc. London

37, 51–135. https://doi.org/10.1111/j.1096-3642.1984.tb00068.x

Olesen, J., 1979. Prey capture in dragonfly nymphs (Odonata, Insecta): labial protraction by means of

a multi-purpose abdominal pump. Vidensk. Meddr dansk naturh. Foren. 141, 81–96.

Pohl, H., 2010. A scanning electron microscopy specimen holder for viewing different angles of a

single specimen. Microsc. Res. Tech. 73, 1073–1076. https://doi.org/10.1002/jemt.20835

Pritchard, G., 1965. Prey capture by dragonfly larvae (Odonata; Anisoptera). Can. J. Zool. 43.

Pritchard, G., 1964. The prey of dragonfly larvae (Odonata; Anisoptera) in ponds in Northern Alberta.

Can. J. Zool. 42, 785–800. https://doi.org/10.1139/z64-076

Quenta Herrera, E., Casas, J., Dangles, O., Pincebourde, S., 2018. Temperature effects on ballistic prey

capture by a dragonfly larva. Ecol. Evol. 1–9. https://doi.org/10.1002/ece3.3975

Rumble, J.R., Lide, D.R., Bruno, T.J., 2017. CRC handbook of chemistry and physics, 97th ed. CRC

Press, New York.

Shahinpoor, M., 1992. Conceptual design, kinematics and dynamics of swimming robotic structures

using ionic polymeric gel muscles. Smart Mater. Struct. 1, 91–94. https://doi.org/10.1088/0964-

1726/1/1/014

Shi, L., Guo, S., Asaka, K., 2010. A novel multifunctional underwater microrobot. 2010 IEEE Int. Conf.

Robot. Biomimetics, ROBIO 2010 873–878. https://doi.org/10.1109/ROBIO.2010.5723441

Snodgrass, R.E., 1955. The Dragonfly Larva. Smithson. Misc. Collect. Washingt. D.C.

Summers, A.P., Darouian, K.F., Richmond, A.M., Brainerd, E.L., 1998. Kinematics of aquatic and

terrestrial prey capture in Terrapene carolina, with implications for the evolution of feeding in

cryptodire turtles. J. Exp. Zool. 281, 280–287. https://doi.org/10.1002/(SICI)1097-

010X(19980701)281:4<280::AID-JEZ4>3.0.CO;2-K

Tanaka, Y., Hisada, M., 1980. The hydraulic mechanism of the predatory strike in dragonfly larvae. J.

Exp. Biol. 88, 1–19.

Tillyard, R.J., 1917. The biology of dragonflies. Igarss 2014 1–5. https://doi.org/10.1007/s13398-014-

0173-7.2

Van Damme, J., Aerts, P., 1997. Kinematics and functional morphology of aquatic feeding in

Australian snake-necked turtles (Pleurodira; Chelodina). J. Morphol. 233, 113–125.

https://doi.org/10.1002/(SICI)1097-4687(199708)233:2<113::AID-JMOR3>3.0.CO;2-7

Van Wassenbergh, S., 2015. A Solution Strategy to Include the Opening of the Opercular Slits in

Moving-Mesh CFD Models of Suction Feeding. Integr. Comp. Biol. 55, 62–73.

https://doi.org/10.1093/icb/icv031

Van Wassenbergh, S., Strother, J.A., Flammang, B.E., Ferry-Graham, L.A., Aerts, P., 2008. Extremely

fast prey capture in pipefish is powered by elastic recoil. J. R. Soc. Interface 5, 285–296.

https://doi.org/10.1098/rsif.2007.1124

Vogel, S., 1997. Life in moving fluids, 2nd Editio. ed. Princeton University Press, Princeton.

Wainwright, P.C., McGee, M.D., Longo, S.J., Patricia Hernandez, L., 2015. Origins, Innovations, and

Diversification of Suction Feeding in Vertebrates. Integr. Comp. Biol. 55, 134–145.

https://doi.org/10.1093/icb/icv026

Yaniv, S., Elad, D., Holzman, R., 2014. Suction feeding across fish life stages: Flow dynamics from

larvae to adults and implications for prey capture. J. Exp. Biol. 217, 3748–3757.

https://doi.org/10.1242/jeb.104331

Young, B.A., 1991. The influences of the aquatic medium on the prey capture system of snakes. J.

Nat. Hist. 25, 519–531. https://doi.org/10.1080/00222939100770321

Figures

Figure 1) 3D models of the different basic types of prehensile labial masks in dragonfly larvae (Anisoptera). Models are based on µCT scans of specimens of the aeshnid *Anax imperator* (left) which features a flat prementum for grasping large prey and the libellulid *Sympetrum* sp. (right) which features a spoon-shaped prementum also capable of scooping up smaller animals. A) dorsal view, B) lateral view; pm= postmentum, prm=prementum, lp=labial palps

Figure 2) The angle of attack and total velocity of the prementum over the course of the predatory strike. A) Angle of attack (α) of the prementum of each species for 10 timepoints extracted from strike data. We see a significantly higher α at three timepoints. (Wilcox Rank Sum Test, p<0.05. *Sympetrum* n=6, *Anax* n=7). B) Median strike velocity for both species at each timepoint. Velocity represents the sum of translational and rotational velocity components at the tip of the prementum. At no point, there is a significant difference in strike velocity between the two species (Wilcox Rank Sum Test, *Sympetrum* n=6, *Anax* n=7). Data extracted from High-speed footage of the predatory strike.

Figure 3) Both prehensile labial masks generate substantial amounts of lift and the drag coefficient (C_{dw}) is higher in *Sympetrum*. Top left corner: Force diagram showing the direction of drag and lift force. Please note that these directions are relative to the flow directions, not the orientation of the PLM. A, B) Total drag and lift force acting on the prementum for both species at each timepoint (red *Anax*, blue *Sympetrum*, lighter colours indicate lift data). C, D) C_{dw}/ C_{lw} based on the forces in A and B (same colour coding). Data points are results from CFD Simulations based on kinematic data from HS Videos (Figure 2). Lines are interpolations using a Savitzky-Golay filter (span=0.5). Please note that the drag/lift coefficient in *Anax* was zero at the first timepoint, this is an artefact due to the median velocity and forces of the prementum being too low to calculate meaningful coefficients.

Figure 4) Flow characteristics change during the predatory strike I. Contours of pressure around the median line of the prementum of *Anax* l) and *Sympetrum* r). Four exemplary timepoints are shown (results from all simulations are shown in supp. figure S5). Colour indicates fluid pressure around the prementum (the colour range is identical in all images). Note the formation of low-pressure regions on the dorsal side of both prementa (yellow areas). CFD Simulations were based on kinematic data from HS Videos (Figure 2).

Figure 5) Flow characteristics change during the predatory strike II. Contours of pressure around the median line of the prementum of *Anax* l) and *Sympetrum* r). Four exemplary timepoints are shown (results from all simulations are shown in Supp. figure S6). Colour indicates fluid speed around the prementum (the colour range is identical in all images). Note the flow separation at higher α in *Anax* (Step 3,5). This does not occur in *Sympetrum*. CFD Simulations were based on kinematic data from HS Videos (Figure 2).

Figure 6) Drag and Lift Coefficient vary with the angle of attack (α). Top left corner: force diagram showing the direction of drag and lift force. Please note that these directions are relative to the flow directions. A) drag coefficient of the prementum of *Anax* (red) and *Sympetrum* (blue) vs. α. Icons at the top of the plot show orientation of the prementum to flow (flow direction right to left). B) lift coefficient of the prementum of *Anax* (red) and *Sympetrum* (blue) vs. α. In both plots: points show values from simulations, lines are interpolations using Savitzky-Golay filter. Data from CFD Simulations run at 0.4 m/s flow velocity and α was varied between 0° and 70°.

Figure 7) Flow characteristics change with the angle of attack (α) I. Contours of pressure around the median line of the prementum of *Anax* l) and *Sympetrum* r). Four exemplary timepoints are shown (results from all simulations are shown in Supp. figure S7). Colour indicates fluid pressure around the prementum. Note the formation of low-pressure regions on the dorsal side of both prementa at higher α. In *Sympetrum* a high-pressure area forms in the premental cavity at low α (e.g. 10°). CFD Simulations were run at 0.4m/s fluid velocity and α was varied between 0° and 70° (Figure 6).

Figure 8 Flow characteristics change with the angle of attack (α) II. Contours of total flow speed around the median line of the prementum of *Anax* l) and *Sympetrum* r). Four exemplary timepoints are shown (Results from all simulations are shown in Supp. figure S6). Colour indicates flow speed around the prementum. Note the separation of flow occurring in the prementum of *Anax* at higher α. This does not occur in *Sympetrum.* CFD Simulations were run at 0.4m/s fluid velocity and α was varied between 0° and 70° (Figure 6).

Figure 9) Flow is directed to a low-pressure area where setae are located. A) Photograph of the PLM of *Sympetrum*. Four arrays of setae can be identified, one on each side of the premental cavity, one on each labial palp. Note that they do not completely cover the opening, but only the premental cavity. B) SEM Micrograph of the Prementum with closed labial palps. Note that setae now cover the entire premental cavity. C) Streamlines show that flow is directed downwards above the prementum, very likely into the setae. D) 3D Heatmap showing the pressure distribution around the prementum of *Sympetrum.* Only pressure exceeding ambient pressure is displayed. It shows no high-pressure region at the opening which might complicate prey capturing. E) Vector field in the median plane showing flow speed distribution around the prementum. Inside the premental cavity, the flow speed is increasing and directed into the cavity, which correlates with regions of lower pressure. CFD Simulations based on kinematic data at timepoint 6 (0.44m/s flow velocity and α = 44°).

Figure 10) Motion of the fluid during a predatory strike. A) Still frames from HS-Video show characteristic points during a predatory strike of *Anax* (l) and *Sympetrum* (r). *Spirulina* algae were used as particles to observe the surrounding water. Characteristic particles are highlighted to visualize changes in particle position. Particle positions from previous images are shown in a different colour depending on their age, with the newest position in red. Both series show very little bow wave formation, as particles move barely compared to the motion of the PLM. Especially in *Anax,* we can also observe motion against the direction of the PRM above it (indicating a low-pressure zone), while particles below the PRM are pushed down. The corresponding videos can be found in supplementary material S9.

Supplementary Material

S1) Supplementary Figure

Supplementary Figure S1) Exemplary still frames from two High-speed Videos showing the tracking points and calculated vectors. A) *Anax imperator* B) *Sympetrum* sp. Videos were recorded at 5400 fps. The formula for α calculation is shown in A), with B_{t0} being the location of the tracking point in the frame for which α is calculated, B_{t+5} being the location of the tracking point 5 frames later and B_{t-5} the location 5 frames earlier.

632   *S2) Supplementary Table*

R	ρ [kg/m^3]	l [mm]	v [ms^{-1}]	μ [Pa*s]	R
Anax	997.04	6.75	0.49	891	3701
Sympetrum	997.04	3.6	0.44	891	1773
C_{dw} (timepoint 6)	ρ [kg/m^3]	A_w [cm^2]	v [ms^{-1}]	F_d [mN]	C_{dw}
Anax	997.04	2.084	0.49	1.261	0.0051
Sympetrum	997.04	0.499	0.44	0.501	0.0106
C_{lw} (timepoint 6)	ρ [kg/m^3]	A_w [cm^2]	v [ms^{-1}]	F_l [mN]	C_{lw}
Anax	997.04	2.084	0.49	1.377	0.0552
Sympetrum	997.04	0.499	0.44	0.417	0.0865

Supplementary Table S2) Calculation of Reynolds Number and C$_{dw}$ and C$_{lw}$ for one exemplary timepoint.

634   *S3) Supplementary Table*

Parameter	Final Resolution	2x Resolution	Ratio
No. of elements	4,681,683	18,230,392	25.68
F_{drag} [N]	-0.000510598	-0.000514853	99.17
F_{lift} [N]	0.0004257	0.000417959	98.18

Supplementary Table S3) Exemplary results of mesh analysis for *Sympetrum*

646   *S4) Supplementary Figure*

Supplementary Figure S4) 3D meshes used for CFD simulations. A, B) Exemplary meshes for CFD Simulation as generated by ANSYS ICEM. A) *Anax imperator*, B), *Sympetrum* sp. Large images show a longitudinal section through the entire domain, insets provide a close-up around the model to show prism layers. C.D) 3D models of the prementum based on CT scans of the head of a specimen of each *Anax imperator* C) and *Sympetrum* sp. D). The models were retopologised to ensure a clean topology without holes or intersections. The wireframe shows surface mesh resolution.

647   *S5) Supplementary Figure*

Supplementary Figure S5) Expendable 3D printed brace for critical point drying of the prehensile labial mask in an opened state. The brace was designed in Blender and printed on a Prusa i3 Mk 3 FDM Printer using PLA Filament.

657   *S6) Supplementary Figure*

Supplementary Figure S6) Flow characteristics change during the predatory strike. Contours of pressure (left) and contours of total flow speed (right) around the median line of the prementum of *Anax* l) and *Sympetrum* r) at all timepoints. l) Colour indicates fluid pressure around the prehensile labial mask. Note the formation of low-pressure regions on the dorsal side of both labial masks (yellow areas). r) Colour indicates total flow speed around the prehensile labial mask (the colour range is identical in all images). Note the flow separation at higher α in *Anax* (Step 3,5). This does not occur in *Sympetrum*. CFD Simulations were based on kinematic data from HS Videos (Figure 2).

658   *S7) Supplementary Figure*

Supplementary Figure S7) Flow characteristics change with the angle of attack. Left) Contours of pressure around the median line of the prementum of *Anax* l) and *Sympetrum* r). Colour indicates fluid pressure around

the prehensile labial mask. Note the formation of low-pressure regions on the dorsal side of both labial masks at higher α. In *Sympetrum* a high-pressure area forms in the premental cavity at low α (e.g. 10°); right) Contours of total flow speed around the median line of the prementum of *Anax* l) and *Sympetrum* r). Colour indicates total flow speed around the prehensile labial mask (the colour range is set to a local range, to enhance pattern visibility). Note the separation of flow occurring in the prementum of *Anax* at higher α. This does not occur in *Sympetrum.* CFD Simulations were run at 0.4m/s fluid velocity and α was varied between 0° and 70° (Figure 6).

682   *S8) Supplementary Figure*

Supplementary Figure S8) Changing the size increases drag and lift coefficient. Scaling the prementum of *Anax* down to the size of the prementum of *Sympetrum* increases its drag and lift coefficient. Scaling the prementum of *Sympetrum* up also increases its drag and lift coefficient. The overall flow patterns are similar. We only see a lower degree of flow separation in the downscaled *Anax* prementum due to the lower Reynolds number.

684   *S9) Supplementary Video*

685   Video of predatory strikes with visualised flow.

686