Prediction of survival after partial hepatectomy using a physiologically based pharmacokinetic model of indocyanine green liver function tests

Adrian Köller [1], **Jan Grzegorzewski** [1], **Hans-Michael Tautenhahn** [2] **and Matthias König** [1,*]

[1] *Institute for Theoretical Biology, Institute of Biology, Humboldt University, Berlin, Germany*
[2] *Experimental Transplantation Surgery, Department of General, Visceral and Vascular Surgery, Jena University Hospital, Jena, Germany*

Correspondence*:
Matthias König
koenigmx@hu-berlin.de

2 ABSTRACT

The evaluation of hepatic function and functional capacity of the liver are essential tasks in
hepatology as well as in hepatobiliary surgery. Indocyanine green (ICG) is a widely applied test
compound that is used in clinical routine to evaluate hepatic function. Important questions for
the functional evaluation with ICG in the context of hepatectomy are how liver disease such as
cirrhosis alters ICG elimination, and if postoperative survival can be predicted from preoperative
ICG measurements. Within this work a physiologically based pharmacokinetic (PBPK) model of
ICG was developed and applied to the prediction of the effects of a liver resection under various
degrees of cirrhosis. For the parametrization of the computational model and validation of model
predictions a database of ICG pharmacokinetic data was established. The model was applied
(i) to study the effect of liver cirrhosis and liver resection on ICG pharmacokinetics; and (ii) to
evaluate the model-based prediction of postoperative ICG-R15 as a measure for postoperative
outcome. Key results are the accurate prediction of changes in ICG pharmacokinetics caused
by liver cirrhosis and postoperative changes of ICG-elimination after liver resection, as validated
with a wide range of data sets. Based on the PBPK model, individual survival after liver resection
could be classified, demonstrating its potential value as a clinical tool.

Keywords: Indocyanine Green, Hepatectomy, Liver Cirrhosis, Mathematical Model, Computational Model, Pharmacokinetics, Liver
Function, Liver Resection

1 INTRODUCTION

Determining liver function is a crucial task in hepatology, e.g., for liver disease diagnosis or evaluation
of pre- and postoperative functional capacity of the liver. An accurate assessment is especially relevant
in the context of liver surgery as postoperative complications are associated with reduced functional
capacity. A comprehensive characterization of the status of a patient and their liver is routinely performed
before liver surgery such as partial hepatectomy. This includes among others anthropometric factors (e.g.
age, sex, body weight), static liver function tests (e.g., ALT, AST, albumin, bilirubin, INR, prothrombin
time), cardiovascular parameters (e.g. cardiac output, blood pressure, hepatic blood flow) and lifestyle

factors (e.g. smoking, medication) as well as volumetric information obtained by CT or MRI scans. In
addition, quantitative evaluation of liver function is often performed via pharmacokinetic measurements of
test compounds specifically metabolized by the liver (dynamical liver function tests) such as methacetin
(LiMAx (Rubin et al., 2017)) and MBT (Gorowska-Kowolik et al., 2017)), caffeine (Renner et al., 1984),
galactose (Bernstein et al., 1960) and indocyanine green (ICG) (Sakka, 2018).

ICG is an inert, anionic, water-soluble, tricarbocyanine dye that is bound to plasma proteins. After
intravenous administration ICG is taken up exclusively by the liver and excreted unchanged into the bile. It
is not reabsorbed by the intestine and does not undergo enterohepatic circulation (Wheeler et al., 1958). As
a result, ICG is an ideal test compound to test hepatic uptake and biliary excretion. Based on the plasma
time course of ICG, pharmacokinetic parameters are calculated as a proxy for liver function, the most
common parameters being: (i) ICG retention ratio 15 minutes after administration (ICG-R15) [%]; (ii)
ICG plasma disappearance rate (ICG-PDR) [%/min]; (iii) ICG-clearance [ml/min]; and (iv) ICG half-life
(ICG-$t_{1/2}$) [min]. Reduced elimination of ICG by the liver is directly reflected by these parameters (Sakka,
2018).

Liver disease, especially advanced and more severe liver disease, is accompanied by a loss of liver
function which can be quantified with dynamical liver function tests. The effects of liver disease on
ICG-elimination have been studied extensively, e.g., in different stages of primary biliary cholangitis
(PBC) (Vaubourdolle et al., 1991). ICG elimination is reduced in Gilbert's disease (Martin et al., 1976) as
well as in patients with hepatic fibrosis and cirrhosis (Gadano et al., 1997). Interestingly, also non-liver
diseases can affect ICG parameters, e.g., ICG-clearance is significantly reduced in patients with chronic
pancreatitis (Andersen et al., 1999).

Liver cirrhosis is an end stage liver disease and highly relevant in the context of hepatobiliary surgery.
The most common causes for liver cirrhosis in Europe are alcohol abuse, chronic hepatitis B and/or C virus
infection or non-alcoholic fatty liver disease (Hackl et al., 2016). The pathological characteristics of liver
cirrhosis include degeneration of hepatocytes as well as a reduction of liver perfusion through increased
portal resistance. An additional characteristics is the formation of intrahepatic shunts, which bypass part
of the hepatic blood supply around the liver tissue. Consequently, no ICG can be extracted, resulting in
further reduced elimination (Schuppan and Afdhal, 2008).

The severity of cirrhosis can be described using the Child-Turcotte-Pugh-Score (CTP) (Child and Turcotte,
1964; Pugh et al., 1973). The CTP is an empiric, qualitative, dis-continuous classification of the severity
of the "hepatic functional reserve" (Botero and Lucey, 2003). Based on a set of parameters, the CTP
assigns a score from 5-15 to a cirrhotic patient, where the more severe a patients symptoms are the higher
the score is. These parameters are serum bilirubin and serum albumin concentrations, the International
Normalized Ratio (INR), amount of ascites in ultrasound and the degree of encephalopathy (Child and
Turcotte, 1964; Pugh et al., 1973). Patients are classified depending on their mortality risk as CTP-A (5-6
points, low risk), CTP-B (7-9 points, intermediate risk) or CTP-C (10-15 points, high risk). Differences in
ICG-elimination between cirrhotic patients and control subjects has been widely assessed (Caesar et al.,
1961; Burns et al., 1991; Gilmore et al., 1982; Figg et al., 1995; Møller et al., 2019; Mukherjee et al., 2006;
Pind et al., 2016). A good correlation between CTP-score and ICG-elimination has been reported with
ICG-elimination decreasing as CTP-score increases (Figg et al., 1995; Møller et al., 2019; Mukherjee et al.,
2006; Pind et al., 2016).

For many primary and secondary liver tumors, partial hepatectomy is the only curative treatment option.
Liver resection (partial hepatectomy) describes the surgical removal of a part of the liver. Hepatectomy

is an important procedure in general surgery with more than 20,000 liver resections in Germany per
year (Filmann et al., 2019). The procedure has been widely performed for the treatment of various liver
diseases, such as malignant tumors, benign tumors, calculi in the intrahepatic ducts, hydatid disease, and
abscesses (Jin et al., 2013). Despite advances in technology and increasing numbers of highly experienced
and specialized clinicians in the field of hepatobiliary surgery, postoperative morbidity and mortality is still
a major issue as complex resections are increasingly being performed in older and higher risk patients (Jin
et al., 2013). Major hepatectomy in the presence of cirrhosis is considered to be contraindicated due to the
high mortality rate. Only selected patients with stable Child A status or an ICG-R15 of less than 10% may
potentially be considered for extended liver resection. (Kitano and Kim, 1997).

A key challenge in HPB (hepato pancreato biliary) surgery is to predict functional capacity of the future
liver remnant and thus reduce morbidity and mortality after extended liver resection. As a result, the decision
whether or not a partial hepatectomy can be performed is often based on predictions of postoperative liver
function (in addition to the remnant liver volume), which are in turn based on preoperative assessment of
liver function (and volume). Understanding how cirrhosis alters liver function as measured via ICG is of
high clinical relevance. Elucidating how ICG parameters change with increasing CTP score would be a
valuable asset for the functional evaluation of patients with liver disease.

Important questions for the evaluation of liver function with ICG in the context of HPB surgery are
(i) how liver disease, especially cirrhosis, alter ICG elimination, and (ii) if postoperative survival can be
predicted from preoperative ICG measurements. Within this work a physiologically based computational
model of ICG pharmacokinetics was developed and applied to study these questions.

2 MATERIAL AND METHODS

2.1 Data

For the calibration and validation of the model a large data set of ICG measurements and physiological
data was established. All data is available via the pharmacokinetics database PK-DB (https://pk-
db.com) (Grzegorzewski et al., 2021). PK-DB was used to encode the information on (i) patient
characteristics (e.g. age, disease, medication), (ii) applied interventions (e.g. ICG dosing, route of
application); (iii) measured ICG time courses in plasma and (iv) ICG pharmacokinetic parameters (e.g.
ICG-PDR, ICG-R15, ICG-clearance).

2.2 Indocyanine Green Pharmacokinetic Parameters

Pharmacokinetic parameters of ICG were calculated from the plasma-concentration time courses using
non-compartmental methods (Urso et al., 2002). The elimination rate constant (k_{el}) was calculated by
fitting the concentration-decay-curve to an exponential function: $c(t) = c(0) \cdot e^{-k_{el} \cdot t}$. ICG-PDR is k_{el}
reported in [%/min]. Half-life ($t_{1/2}$) was calculated as $log(2)/k_{el}$, ICG-clearance as $CL = V_d \cdot k_{el}$, with
the apparent volume of distribution $V_d = D/(AUC_\infty \cdot kel)$. D is the applied dose of ICG and AUC_∞ is
the area under the plasma-concentration time curve AUC calculated via the trapezoidal rule, extrapolated
until infinity. ICG-R15 $= c(15)/c_{max}$ was calculated as the ratio between the plasma-concentration after
15 minutes and the maximum concentration c_{max}.

2.3 Model

107   The computational model is an ordinary differential equation (ODE) model encoded in the Systems
108 Biology Markup Language (SBML) (Hucka et al., 2019; Keating et al., 2020). It is defined as a set of
109 species (metabolites), compartments (organs and blood compartments) and reactions (processes such
110 as metabolic reactions and blood transport). The model was developed using sbmlutils (König, 2021b),

and cy3sbml (König and Rodriguez, 2019) and simulated using sbmlsim (König, 2021a) based on the high-performance SBML simulator libroadrunner (Somogyi et al., 2015).

2.4 Model Parameterization

Values for organ volumes and tissue blood flows were taken from literature (ICRP, 2002). A subset of model-parameters was determined by minimizing the residuals between model predictions and clinical data. This optimization-problem was solved using SciPy's least_squares method and differential evolution algorithm (Virtanen et al., 2020). For the objective cost function F depending on the parameters $\vec{p}$ a simple L2-Norm was used consisting of the sum of weighted residuals

$$F(\vec{p}) = 0.5 \cdot \sum_{i,k} \left(w_k \cdot w_{i,k} \cdot r_{i,k}(\vec{p}) \right)^2 = \sum_{i,k} \left(w_k \cdot w_{i,k} \cdot (y_{i,k} - m_{i,k}(\vec{p})) \right)^2 \tag{1}$$

where $r_{i,k} = (y_{i,k} - m_{i,k}(\vec{p}))$ is the residual of time point i in time course k for model prediction $m_{i,k}(\vec{p})$ and the corresponding data point $y_{i,k}$; $w_{i,k}$ is the weighting of the respective data point i in time course k based on the error of the data point and w_k = the weighting factor of time course k. Weighting of time courses was based on the number of subjects per study. The final parameter set given in Tab. 2 was determined using 250 runs of the local least square optimization. The data used for the parameter fit is listed in Tab. 1.

2.5 Uncertainty analysis

To evaluate the uncertainty of model predictions uncertainty analysis was performed for a subset of simulations. Each model parameter was changed individually by ±25%. From the set of resulting time courses the mean, standard deviation (SD) and minimum and maximum values at each time point were calculated. These uncertainty areas were displayed as shaded areas. Parameters corresponding to physical constants (such as molecular weights) and dosing were not varied in the uncertainty analysis, as well as parameters for conservation conditions such as the fractional blood flow through the lung (must be 1).

2.6 Classification

Classification models were developed which predict survival after hepatectomy (binary classification: Survivors/Non-Survivors) based on different sets of features (see Tab. 3). Classification was performed using scikit-learn (Pedregosa et al., 2011) with a C-support vector classifier using a polynomial kernel. For model training and evaluation a dataset of 141 patients with information on survival status, resection rate and preoperative ICG-R15 was used (Seyama and Kokudo, 2009; Wakabayashi et al., 2004). Cross-validation was performed using the ShuffleSplit method with 200 iterations and a train-test-ratio of 75%/25%. Based on the confusion matrix the following evaluation metrics were calculated: precision/positive predictive value (PPV), recall, balanced accuracy, F1 score, negative predictive value (NPV) and receiver operator curves (ROC).

For the classification models PBPK1 and PBPK2 the model parameter $f_{cirrhosis}$ was determined from preoperative ICG-R15 values using

$$f_{cirrhosis} = a \cdot \ln(b \cdot R15_{preop}) + c \tag{2}$$

with $R15_{preop}$ being the clinically measured preoperative ICG-R15 [dimensionless]. The coefficients $a = 0.312$, $b = 1.693$ and $c = 0.861$ were determined by fitting the curve to the predicted dependency of ICG-R15 on $f_{cirrhosis}$ (see Fig. 4C). This provided an estimate of individual liver disease (cirrhosis

degree). This estimated parameter allowed in combination with the resection rate to predict individual postoperative ICG-R15 values.

3 RESULTS

Within this work a PBPK model of ICG pharmacokinetics was developed and applied to study (i) how liver disease, especially cirrhosis, alter ICG elimination, and (ii) if postoperative survival can be predicted from preoperative ICG measurements in the context of partial hepatectomy.

3.1 Data

A wide range of heterogeneous data was curated for model building (parameterization) and subsequent model validation (comparison of model predictions to clinical data). An overview of the 29 studies with their respective clinical protocols is provided in (Tab. 1). All data is freely available from https://pk-db.com.

3.2 Model

A PBPK model for the prediction of ICG pharmacokinetics was developed. To simulate the whole-body distribution and hepatic elimination of ICG two models were coupled: (i) A whole-body model (Fig. 1A) describing the distribution of ICG in the body and to the organs via blood flow. (ii) A liver model (Fig. 1B) which describes hepatic uptake of ICG, biliary excretion of ICG and transport of ICG into the feces. ICG pharmacokinetic parameters (i.e. ICG-PDR, R15, clearance, half-life) were calculated from the resulting time course predictions of ICG in the venous plasma.

3.2.1 Distribution and blood flow

The distribution of ICG on the whole-body level is modeled using a network of blood flows representing the systemic circulation. From the venous blood, ICG is transported through the lung into the arterial blood from where it can reach the liver on two paths: (i) through the hepatic artery and (ii) through the gastrointestinal tract via the portal vein. Because the liver is the only tissue partaking in the elimination of ICG, all other organs (e.g. kidney, heart, adipose tissue, muscle, etc.) were pooled into the rest compartment. Each organ consists of a blood compartment (representing the vessels) and a tissue compartment. ICG transport via blood flow was implemented as irreversible transport. The transport v_i from compartment i to the next compartment is determined by the ICG concentration C_i in compartment i and a compartment-specific blood flow Q_i. Q_i is determined by the cardiac output Q_{CO} and a compartment specific fractional tissue blood flow fQ_i. Multiple conservation conditions hold in the model to ensure mass and flow balance. First the sum of blood flows from the arterial to the venous compartment must equal the sum of flows in the opposite direction: $Q_{CO} = Q_{lu} = Q_h + Q_{re}$. Flow into an organ must be equal to the flow out of the organ. E.g. hepatic venous blood flow must be equal to the sum of hepatic arterial and portal venous blood flow: $Q_h = Q_{ha} + Q_{po}$.

3.2.2 Hepatic metabolism and biliary excretion

The liver model (Fig. 1B) consists of three consecutive transport reactions of ICG. After ICG is taken up in the liver it is excreted into the bile. Both transport reactions are modeled as irreversible Michaelis-Menten-kinetics. From the bile, ICG is transported into the feces modeled via a first order kinetic. All transport kinetics scale with the liver volume V_{li}.

3.2.3 Parameter fitting

Parameter fitting of the model was performed using a subset of ICG time courses and extraction-ratio measurements (see Tab. 1). No ICG pharmacokinetic parameters were used in the model fitting. Overall, 5 model parameters were fitted (see Tab. 2). Two of them determine the import of ICG in the liver, three determine the subsequent excretion in the bile. The agreement between fit data and model predictions

188 improved substantially during parameter fitting and all training data with the exception of three simulations
189 (Meijer1988, Chijiiwa2000 and Burns1991) could be described very well after parameter fitting.

3.2.4 Modeling liver cirrhosis

The reference model, representing a healthy human subject, was adjusted to simulate cirrhosis by
including a combination of functional tissue loss (due to scarring and necrosis in cirrhosis) and the
formation of intrahepatic shunts, both key hallmarks of cirrhosis (Fig. 1C). The loss of functional liver
tissue was controlled via the parameter $f_{tissue\_loss} \in [0, 1)$ which defines the fraction of parenchymal
cell volume lost in the liver due to the disease. For modeling arteriohepatic and portosystemic shunts two
additional blood vessels were introduced into the model. They connect the hepatic artery and the portal
vein directly to the hepatic vein. As a result, a part of the portal venous and arterial blood bypasses the
active liver tissue and is shunted to the hepatic venous blood compartment, so that ICG can not be extracted
(corresponding to *in silico* shunts). The amount of blood that flows through the shunts is determined by the
parameter $f_{shunts} \in [0, 1)$, which defines the fraction of blood bypassing the liver. The remaining blood
$(1 - f_{shunts})$ reaches the liver tissue and ICG can be extracted. To simulate various degrees of cirrhosis the
parameters f_{shunts} and f_{tissue_loss} were varied in lockstep by coupling them into the parameter $f_{cirrhosis}$.
The following values for $f_{cirrhosis}$ were used: healthy - 0.0, mild cirrhosis - 0.41, moderate cirrhosis - 0.70,
severe cirrhosis - 0.82.

3.2.5 Modeling hepatectomy

206 The developed model allows to predict changes in ICG pharmacokinetic parameters after partial
207 hepatectomy (Fig. 1D). *In silico* liver resections were simulated by reducing the fractional liver volume
208 $FVli$ by up to 90% (corresponding to a resection rate of 90%). The absolute liver volume is determined
209 with the body weight BW via $FVli \cdot BW$. All liver resections were simulated under varying degrees of
210 cirrhosis as described above.

3.3 Healthy controls

In a first step the fitted model was evaluated with the data used for model calibration consisting of ICG
time courses in healthy subjects (Fig. 2). For the simulations infusion protocols and body weights were
adjusted as reported in the respective studies (see Tab. 1 for details). If no body weight was reported 75 kg
were assumed.

The model predictions for ICG plasma disappearance curves after an ICG bolus are in good agreement
with the clinical data (Andersen et al., 1999; Grundmann et al., 1992; Kamimori et al., 2000; Klockowski
et al., 1990; Niemann et al., 2000; Meijer et al., 1988) (Fig. 2A-E). In addition, more complex infusion
protocols as reported in Soons et al. (Soons et al., 1991) can also be described (Fig. 2F), infusion protocol
of three different infusion rates ($2.0 \rightarrow 0.5 \rightarrow 1.0$ mg/min, each for 40 minutes). Due to the high extraction-
ratio of ICG by the liver, the plasma concentration reaches steady state quickly after each change in the
infusion rate. Next, simulations of the biliary excretion rate of ICG after bolus administrations of 0.5, 1.0
and 2.0 mg/kg ICG were performed and the results were compared to clinical data (Meijer et al., 1988;
Chijiiwa et al., 2000) (Fig. 2G, H).

Finally, simulations of constant infusions were performed and compared to reported arterial and hepatic
vein time courses of ICG (Leevy et al., 1962) and ICG extraction ratios (Leevy et al., 1962; Grainger et al.,
1983) (Fig. 2I-L).

Overall, the model shows the ability to accurately predict ICG time courses for venous and arterial
plasma concentrations, for hepatic vein concentrations, the biliary excretion rate and extraction ratios when

compared to clinical data. Especially plasma time courses of ICG after ICG bolus and ICG infusion are
very well predicted by the model, even for varying administration protocols (dosing and infusion rates).

In a next step a systematic analysis of the dose dependency of ICG pharmacokinetic parameters was
performed (Fig. 3).A dose-dependency of the ICG parameters can only be observed if the ICG dose
exceeds 100 mg (much higher then the typically applied doses of 20 - 35 mg), resulting in a reduction in
ICG-clearance and ICG-PDR as well as an increase of ICG-R15 and ICG-$t_{1/2}$ (Fig. 3A-D). The model
predictions could be validated with clinical data (Martin et al., 1975, 1976; Meijer et al., 1988) (Fig. 3E-G).

3.4 Cirrhosis

To simulate changes of ICG pharmacokinetics in cirrhosis, hepatic tissue loss and shunts were included
in the model as described above. First, a systematic analysis of the effect of intrahepatic shunts (f_{shunts}),
functional tissue loss (f_{tissue_loss}) and the combination of both ($f_{cirrhosis}$) on ICG pharmacokinetic
parameters was performed (Fig. 4A-D). All three parameters were varied from 0 (no effect, healthy control)
to 0.9 (severe effect). ICG-clearance and ICG-PDR decrease with increasing $f_{cirrhosis}$ whereas ICG-R15
and ICG-$t_{1/2}$ increase. The loss of a fraction of functional liver tissue appears to have a smaller effect on
ICG pharmacokinetic parameters than shunting of an equal fraction of blood past the liver. When f_{shunts}
and f_{tissue_loss} are combined to $f_{cirrhosis}$ their effect on ICG pharmacokinetic parameters is additive. For
ICG-clearance and ICG-PDR the effects of both parameters combine to an almost linear dependency on
$f_{cirrhosis}$. The decrease in ICG-clearance and ICG-PDR with increasing cirrhosis and increase of ICG-R15
and ICG-$t_{1/2}$ with increasing cirrhosis can be observed over a wide range of applied ICG doses (Fig. 3A-D).

By varying the $f_{cirrhosis}$ parameter from 0 to 0.9 different degrees of cirrhosis were simulated and the
nonlinear relation between ICG-R20 and ICG-kel as well as ICG-R20 and ICG-$t_{1/2}$ could be predicted
(Fig. 4E,F). As seen in the systematic analysis (Fig. 4A-D) ICG-$t_{1/2}$ and ICG-R20 increase with cirrhosis
whereas ICG-kel decreases. The correlation between the ICG pharmacokinetic parameters is predicted
accurately by the model when compared to a clinical dataset that lacks information about the severity of
liver cirrhosis of its patients(Cherrick et al., 1960; Caesar et al., 1961). Next the ICG-PDR in cirrhotic
patients, acute and recovering hepatitis and control subjects after different doses of ICG (0.5 mg/kg and 5.0
mg/kg ICG) was compared to the model predictions. The clinical data shows higher ICG-PDR values after
an ICG dose of 0.5 mg/kg than after an ICG dose of 5.0 mg/kg (Leevy et al., 1967). In the model prediction
the ICG-PDR in acute and recovering hepatitis resembles that of mild to moderate cirrhosis (Fig. 4G).
ICG-clearances after a bolus administration and during a constant infusion show good positive correlation
in cirrhotic patients (Burns et al., 1991). This correlation is predicted accurately by the model (Fig. 4H).

Having evaluated and validated the effect of $f_{cirrhosis}$ on the model prediction of ICG parameters, we
were interested how the model $f_{cirrhosis}$ parameter compares to the *in vivo* estimation of cirrhosis degree
via the CTP-score (Fig. 5). As described above, the CTP-score is a semi-quantitative scoring system
that describes the severity of liver cirrhosis. An important step to apply the developed PBPK model in a
clinical setting, is the ability to adjust the model individually to the respective status of liver disease in
a patient. Therefore, the relationship between the $f_{cirrhosis}$ parameter and the CTP-Score was evaluated
using multiple datasets in which ICG pharmacokinetic parameters were reported in patient subgroups of
different CTP-Scores (Figg et al., 1995; Møller et al., 1998, 2019; Herold et al., 2001).

The clinical results of the ICG pharmacokinetic parameters in different CTP-classes were mapped onto
their respective systematic scan (Fig. 5A-D). The resulting $f_{cirrhosis}$ values were then compared between
the patient groups. Additional individual data is shown (Figg et al., 1995).

The resulting mapping between $f_{cirrhosis}$ and the CTP-classes shows a good positive correlation (Fig. 5E).
The $f_{cirrhosis}$ values for the controls groups are close to 0, increasing with the CTP-class. The relation
appears nonlinear, as $f_{cirrhosis}$ shows little difference between CTP-class B and C. The mappings of the
CTP-class to $f_{cirrhosis}$ for the different ICG parameters each give very similar results. From the mapping of
all four pharmacokinetic parameters a mean value of $f_{cirrhosis}$ was calculated for each CTP-class (Control:
$f_{cirrhosis}$ = 0.0; Mild cirrhosis: 0.41; Moderate cirrhosis: 0.70; Severe cirrhosis: 0.82). The resulting
values were used in all simulations of control, mild, moderate and severe cirrhosis, as well as in the above
described dose dependency analysis (Fig. 3A-D).

Only a single study reported the numerical CTP-score of the patient groups in combination with ICG-
clearance (Figg et al., 1995). All other studies instead used the CTP-classes (A, B, C) (Møller et al., 1998,
2019; Herold et al., 2001). With a dataset of individually reported CTP-scores in combination with ICG
pharmacokinetic parameters of cirrhotic patients, it would be possible to calculate the relationship of the
CTP-score on the $f_{cirrhosis}$ parameter more accurately. Such an improved mapping would allow to adjust
the model via the $f_{cirrhosis}$ parameter individually based on the respective severity of liver disease/cirrhosis
of the patient reported as CTP-score.

After establishing the CTP mapping the model was further validated via several comparisons with clinical
data of ICG time courses in cirrhotic and control subjects (Fig. 6).

Assuming moderate cirrhosis ($f_{cirrhosis}$ = 0.7), the model prediction of an ICG time course in a cirrhotic
patient agrees well with the clinical data (Burns et al., 1991) (Fig. 6A,B). The main alteration compared
to the healthy control is the slower disappearance rate resulting in higher ICG plasma concentrations.
The same effect is observed in steady state via a constant ICG infusion (Fig. 6C). Using the $f_{cirrhosis}$
values from the CTP mapping above, the steady state concentrations are predicted in agreement with the
clinical data (Caesar et al., 1961). Fig. 6D,E shows the relation between the hepatic venous and arterial ICG
concentrations and the extraction ratio in a cirrhotic subject. Here, $f_{cirrhosis}$ was set to 0.54 which allowed
to predict arterial and hepatic vein concentration as well as ICG extraction ratio. Finally, in Fig. 6F-H the
ICG extraction ratio predicted for controls and three different cirrhosis degrees was compared to clinical
data (Leevy et al., 1962; Gadano et al., 1997; Caesar et al., 1961). The extraction ratio in cirrhotic subjects
is reduced compared to healthy controls, as predicted by the model.

3.5 Hepatectomy

After validating the model predictions of ICG pharmacokinetics in liver cirrhosis, the model was applied
to liver surgery. To analyze the effect of partial hepatectomy on ICG elimination the change in ICG
pharmacokinetic parameters as a function of the resection rate was simulated (Fig. 7A-D). The scan was
performed for healthy controls as well as three different degrees of cirrhosis.

ICG-clearance and ICG-PDR are highest in the preoperative liver (resection rate = 0) and decrease with
increasing resection rate whereas ICG-$t_{1/2}$ and ICG-R15 are lowest in the healthy liver and increase with
increasing resection rate. The effect of varying the degree of cirrhosis is in accordance with the results
shown in (Fig. 4A-D). Importantly, increasing resection rate and increasing degree of cirrhosis affect
ICG pharmacokinetic parameters in the same manner. The dependencies of ICG-clearance, ICG-PDR,
ICG-$t_{1/2}$ and ICG-R15 on the resection rate are fairly linear up to 50-60% resection, and become much
more non-linear for higher resection rates.

For model validation the predictions were compared to clinical data of subjects undergoing hepatectomy.
For these simulations the resection rate was varied from 0 to 0.9. First, the relative change of ICG-PDR
after partial hepatectomy as a function of the resection rate was simulated (Fig. 7E). The model predicts a

nonlinear dependency of change in ICG-PDR on the remnant liver volume independent of the degree of cirrhosis. This prediction is in good agreement with the clinical data (Thomas et al., 2015; Stockmann et al., 2009). Furthermore, the correlation between measured postoperative ICG-kel and estimated remnant ICG-kel (ICG-kel · fractional liver remnant) was simulated under various degrees of cirrhosis (Fig. 7F). A good correlation can be observed. The model predictions were compared to three different data sets (Ohwada et al., 2006; Okochi et al., 2002; Sunagawa et al., 2021) and are in good agreement with them. In addition, all data sets are in good agreement with each other. The simulated correlation line is independent of the cirrhosis degree, but with increasing cirrhosis ICG-kel decreases. A large variability can be observed in the experimental data, but as our simulations indicate is most likely not due to the underlying liver disease (cirrhosis).

(Thomas et al., 2015) found significant correlation between post-hepatectomy ICG-PDR and intraoperative ICG-PDR measured under trial clamping of those parts of the liver that were to be removed. This was simulated by changing hepatic blood flow and liver volume in separate simulations but in the same intervals. This was performed for a healthy liver as well as three different degrees of cirrhosis (Fig. 7G). The predictions agree well with the clinical data and show that reducing hepatic blood flow (clamping of liver volumes which will be resected) has a very similar effect on ICG elimination as actually removing the respective liver volume via hepatectomy.

Finally, the correlation between preoperative and postoperative ICG-PDR for different resection rates and cirrhosis degrees was simulated and compared to clinical data (Fig. 7H). ICG-PDR is reduced in cirrhosis preoperatively as well as postoperatively. The model prediction agrees with the clinical data (Thomas et al., 2015).

Overall the predictions of liver resection in severely cirrhotic liver is not in good agreement with the clinical data. This reflects the fact that no resections are performed in severely cirrhotic liver due to high risk of postoperative complications. As a consequence, most of the liver resection are performed in mild to moderate cirrhosis. The model allows to perform these risky hepatectomies *in silico* and predict there effect.

In summary, the model allows to systematically predict the changes of ICG pharmacokinetic parameters in HPB surgery under various degrees of liver disease (cirrhosis).

3.6 Prediction of post-hepatectomy survival

An interesting application of the presented PBPK model is the prediction of postoperative outcome for patients undergoing hepatectomy. Preoperative ICG-R15 and the planned resection rate are key parameters included in the decision process whether a patient is eligible to receive liver resection surgery.

As shown above, the presented PBPK model accurately predicts ICG-R15 in liver cirrhosis as well as the changes in ICG-R15 following hepatectomy. As such, we were interested how a classification model based on the PBPK model prediction of postoperative ICG-R15 compares to classification approaches only using clinical data (preoperative ICG-R15, resection rate and calculated postoperative ICG-R15).

Five alternative classification models were developed to predict survival after partial hepatectomy using a dataset of 141 patients (109 survivors and 32 non-survivors) (Seyama and Kokudo, 2009; Wakabayashi et al., 2004). Three of the classification models were solely based on clinical data: (i) Data1A - using preoperative ICG-R15, (ii) Data1B - using the calculated postoperative ICG-R15 by multiplying the future liver remnant (1-resection rate) with the preoperative ICG-R15; and (iii) Data2 - using both the resection rate and the preoperative ICG-R15. In addition two classification models were developed using PBPK model predictions as input: (iv) PBPK1 - using the predicted postoperative ICG-R15. Hereby, the model

parameter $f_{cirrhosis}$ was estimated for every subject based on preoperative ICG-R15, and postoperative
ICG-R15 was predicted using $f_{cirrhosis}$ and the corresponding resection rate; and (v) PBPK2 - using the
resection rate and the estimated $f_{cirrhosis}$ model parameter. An overview of the classification results of
these five models is provided in Tab. 3, Fig. 8 and Fig. S1.

Both PBPK-based classifiers (PBPK1, PBPK2) as well as the Data2 classifier outperform the data-based
classifiers using a single feature (Data1A, Data1B) in predicting survival after partial hepatectomy.

When comparing the classification models using a single feature (Data1A, Data1B, PBPK1) the
physiological-based predicted postoperative ICG-R15 (PBPK1) clearly outperforms the preoperative
(Data1A) as well as calculated postoperative ICG-R15 (Data1B).

Fig. 8A shows the postoperative ICG-R15 in survivors and non-survivors predicted by the model as well
as corresponding preoperative ICG-R15 (inset).

The PBPK model predicted postoperative ICG-R15 is able to distinguish better between survivors and
non-survivors than the preoperative ICG-R15 as can be seen by the clearer separation of the histograms
(Fig. 8A) and the respective ROC curves (Fig. 8C). Both preoperative ICG-R15 as well as calculated
postoperative ICG-R15 are not very useful for the prediction of survival after partial hepatectomy, whereas
predicted postoperative ICG-R15 using PBPK1 is a very good measure to predict survival after partial
hepatectomy.

To determine possible cutoffs for predicted postoperative ICG-R15 based on the PBPK1 classifier the
dependency of evaluation metrics on the cutoff was analyzed (Fig. 8B). Balanced accuracy as well as
f1-score have a maximum at around 35%. The negative predictive value (NPV) as well as recall is 1 up to a
predicted postoperative ICG-R15 ¡ 20%.

Fig. 8D depicts how the predicted postoperative ICG-R15 depends on the resection rate and $f_{cirrhosis}$.
The data confirms that a cutoff value slightly below 40% would correctly predict most of the non-survivors
with a stricter cutoff of 20% avoiding any death after partial hepatectomy. Similar analysis of the data-based
single feature classification models failed to find a significant optimum of evaluation metrics for either
preoperative ICG-R15 or calculated postoperative ICG-R15 (see Supplemental Figure S1).

The two-feature classification models (PBPK2, Data2) show good performance in the survival prediction
comparable to PBPK1 (Fig. 8E, F). Whereas the one-dimensional PBPK1 classifier provides a simple
interpretation and cutoff value, the two dimensional classifiers are more difficult to interpret.

In summary, we developed a single-feature classification model based on a physiological-based model of
ICG elimination (PBPK1) which allows to predict post-hepatectomy survival solely based on preoperative
ICG-R15 input. Importantly, this computational model-based approach clearly outperforms data-based
approaches such as preoperative ICG-R15 and calculated postoperative ICG-R15.

4 DISCUSSION

In summary, a PBPK model for ICG based liver function evaluation was developed, validated, and applied
to the prediction of postoperative outcome after liver surgery, i.e., survival after partial hepatectomy. The
model takes into account physiological factors such as the degree of cirrhosis and the estimated functional
liver remnant, which allowed an accurate prediction of postoperative liver function in agreement with
clinical data. As such, the model has proven its potential of becoming a valuable clinical tool for the
planning hepato-pancreatico-biliary surgery.

The physiologically-based modeling approach allowed us to predict ICG pharmacokinetics data from
29 studies using only a small set of parameters and processes. The model accurately predicts changes in
ICG pharmacokinetic parameters in a wide range of conditions including varying degrees of cirrhosis.
Additionally, *in silico* hepatectomies with underlying cirrhosis are in good agreement with clinical data. As
an important note, all clinical data besides the time courses in healthy subjects used for model calibration
was used for model validation.

Overall, the clinical data shows large variability in ICG pharmacokinetic measurements, mostly due
to inter-individual differences (e.g. Fig. 7F). Possible explanations are differences in blood flow, plasma
proteins or protein amount or activity of the ICG transporter. An important next step would be a systematic
analysis of these possible causes of variability and account for these confounding factors in the model.

One main outcome of this study is a single-feature classification model based on a physiological-based
model of ICG elimination (PBPK1) which allows to predict post-hepatectomy survival solely based on
preoperative ICG-R15 and resection rate. One limitation is the relatively small sample size (n=141) using
retrospective data from Japanese subjects (Seyama and Kokudo, 2009; Wakabayashi et al., 2004). Further
validation of our results with a dataset consisting of Caucasian subjects would be highly relevant.

The developed classification models demonstrated the potential of using PBPK predicted postoperative
ICG-R15 values in the clinical decision process. Whereas the PBPK1 classifier provides a simple cutoff
based on the individual model prediction (Fig. 8AB), PBPK2 provides the dependency of predicted
postoperative ICG-R15 on resection rate and cirrhosis degree (Fig. 8D), both key factors for survival after
liver resection.

Comparing different approaches of predicting postoperative outcome after partial hepatectomy showed
the importance of taking resection rate into account. The data-based classifier combining resection rate
with preoperative ICG (Data2) allowed to achieve comparable classification results to the PBPK-based
classifiers. In contrast, the classification models Data1A and Data1B failed to achieve good prediction
results.

Clinical data supporting our results have been reported by Haegele et al. (Haegele et al., 2016) who
showed that an ICG-R15 >20% on postoperative day 1 predicted poor postoperative outcome, which
agrees well with the results shown in Fig. 8B. The cutoff of ICG-R15 <20% allows to identify low-risk
patients that are unlikely to have poor postoperative outcome after partial hepatectomy. This was confirmed
by the high negative and low positive predictive values (¿80% and 30% respectively), suggesting that
ICG-R15 is especially useful for the identification of low-risk patients. A recommendation was that subjects
with ICG-R15 20-40% should undergo a more careful evaluation of the treatment options and additional
information should be taken into consideration. In agreement with our results, Haegele et al. could show a
significant improvement of the prediction of postoperative survival when using postoperative ICG-R15
(AUC_{roc}=0.893) compared to preoperative ICG-R15 (AUC_{roc}=0.719). As a side note, the experimental
cutoff of >20% by Haegele et al. was determined in a Western population providing evidence that our
classification approach could be generally applied despite developed on data from Japanese subjects.

Multiple studies showed only moderate performance in predicting post-hepatectomy outcome using
preoperative ICG measurements. Gu et al. (Gu et al., 2020) found that preoperative ICG-R15 achieved
an AUC_{roc} of 0.657 (95% CI 0.576-0.739) and 0.640 (95% CI 0.445-0.836) for the prediction of post-
hepatectomy liver failure and 90-day mortality, respectively. Wong et al. (Wong et al., 2013) failed to achieve
any significant prediction of postoperative severe morbidity using preoperative ICG-R15 (AUC_{roc}=0.51,
95% CI 0.38-0.72). Wang et al. (Wang et al., 2018) found that preoperative ICG-R15 surpassed both

CTP-score and MELD for the prediction of severe post-hepatectomy liver failure, but with moderate AUC_{roc}=0.724 (95% CI 0.654-0.787). In summary, this data provides a strong argument for our approach of using predicted postoperative ICG-R15 values via a PBPK model for predicting postoperative survival, which allowed to improve the discriminatory power (see Fig. 8A).

Due to the high mortality rate, extended liver resection in the presence of cirrhosis is considered to be contraindicated. Recommendations are often that only selected patients with Child's A status or preoperative ICG-R15 of less than 10% undergo major hepatectomy (Kitano and Kim, 1997). As can be seen in Fig. 8A (inset), even such a strict cutoff can still result in mortality after hepatectomy. We suggest using a combination of resection rate and individual liver damage (cirrhosis degree) estimated from preoperative ICG-R15 instead of relying on preoperative ICG-R15 alone. As shown by the PBPK2 classifier (and indirectly by the PBPK1 classifier) this approach allows a better evaluation of postoperative risk.

A very interesting result is that our model accurately predicted intraoperative ICG measurements (during trial clamping) (Fig. 7G), in which the blood flow to the area to be resected was clamped. These intraoperative ICG measurements are in very good agreement with post-operative ICG measurements (Thomas et al., 2015). Such an approach would allow to measure the expected postoperative ICG-R15 intraoperatively. Additional clamping data showed that the postoperative hospital stay was significantly longer for intraoperatively clamped ICG-R15 >20% (27.5 ± 14.1 days) compared to <20% (17.9 ± 9.2 days). Due to the good correlation between intraoperative clamped and postoperative ICG measurements this data provides additional support for the proposed lower cutoff of predicted postoperative ICG-R15 <20%. It is likely that postoperative ICG-R15 is not only a good predictor for survival but for postoperative complications in general as indicated by the data from Akita et al. (Akita et al., 2008). Additional support comes from Haegele et al. which showed significantly reduced liver dysfunction (3.6 vs 42.9%, p=0.001), reduced severe morbidity (16.1 vs 42.9%, p=0.016) and hospitalization (7 vs 11 days, p=0.019) for POD1 ICG-R15 <20% vs POD1 ICG-R15 >20% (Haegele et al., 2016).

The presented PBPK model predicts postoperative ICG-R15 immediately after partial hepatectomy, corresponding to clinical measurements on the first postoperative day (POD1). Using postoperative ICG-R15 on POD1 is supported by Haegele et al. (Haegele et al., 2016) who showed that an ICG clearance test was able to predict poor postoperative outcome as early as POD1.

Importantly, due to the physiologically based modeling approach our predictions could easily be further individualized with the availability of respective data. A personalized risk prediction based on an individualized PBPK model could include (i) general information such as age, sex and ethnicity; (ii) physiological information such as body weight, body fat percentage, cardiovascular parameters and organ volumes; (iii) information regarding the liver specifically such as liver perfusion, liver volume and quantification of ICG protein amounts as well as assessment of liver disease such as degree of cirrhosis would be of high relevance. Such an individualization could substantially improve the models prediction of postoperative liver function and outcome in patients undergoing partial hepatectomy. Going forward, an important next step will be to evaluate the model in the clinical context using a high quality dataset reporting individual ICG time courses in combination with a subset of the above-mentioned additional clinical data.

CONFLICT OF INTEREST STATEMENT

All authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

AK and MK designed the study, developed the computational model, build the classifiers, implemented and performed the analysis, and wrote the initial draft of the manuscript. JG provided support with PK-DB and data curation. All authors discussed the results. All authors contributed to and revised the manuscript critically.

FUNDING

MK, AK and JG were supported by the Federal Ministry of Education and Research (BMBF, Germany) within the research network Systems Medicine of the Liver (LiSyM, grant number 031L0054). MK and AK were supported by the German Research Foundation (DFG) within the Research Unit Programme FOR 5151 "QuaLiPerF (Quantifying Liver Perfusion-Function Relationship in Complex Resection - A Systems Medicine Approach)" by grant number 436883643.

DATA AVAILABILITY STATEMENT

The datasets generated for this study can be found in PK-DB available from https://pk-db.com.

REFERENCES

Akita, H., Sasaki, Y., Yamada, T., Gotoh, K., Ohigashi, H., Eguchi, H., et al. (2008). Real-time intraoperative assessment of residual liver functional reserve using pulse dye densitometry. *World journal of surgery* 32, 2668–2674. doi:10.1007/s00268-008-9752-0

Andersen, V., Sonne, J., and Larsen, S. (1999). Antipyrine, oxazepam, and indocyanine green clearance in patients with chronic pancreatitis and healthy subjects. *Scandinavian journal of gastroenterology* 34, 813–817. doi:10.1080/003655299750025750

Bernstein, L. M., Wheeler, J. X., E E, B. O. N. D., Rohmsdahl, M., and Dougherty, N. (1960). The blood galactose disappearance curve as a test of liver function. *Gastroenterology* 39, 293–304

Botero, R. C. and Lucey, M. R. (2003). Organ allocation: model for end-stage liver disease, child-turcotte-pugh, mayo risk score, or something else. *Clinics in liver disease* 7, 715–27, ix. doi:10.1016/s1089-3261(03)00052-7

Burns, E., Triger, D. R., Tucker, G. T., and Bax, N. D. (1991). Indocyanine green elimination in patients with liver disease and in normal subjects. *Clinical science (London, England : 1979)* 80, 155–160

Caesar, J., Shaldon, S., Chiandussi, L., Guevara, L., and Sherlock, S. (1961). The use of indocyanine green in the measurement of hepatic blood flow and as a test of hepatic function. *Clinical science* 21, 43–57

Cherrick, G. R., Stein, S. W., Leevy, C. M., and Davidson, C. S. (1960). Indocyanine green: observations on its physical properties, plasma decay, and hepatic extraction. *The Journal of clinical investigation* 39, 592–600. doi:10.1172/JCI104072

Chijiiwa, K., Watanabe, M., Nakano, K., Noshiro, H., and Tanaka, M. (2000). Biliary indocyanine green excretion as a predictor of hepatic adenosine triphosphate levels in patients with obstructive jaundice. *American journal of surgery* 179, 161–166. doi:10.1016/s0002-9610(00)00274-9

Child, C. G. and Turcotte, J. G. (1964). Surgery and portal hypertension. *Major problems in clinical surgery* 1, 1–85

Figg, W. D., Dukes, G. E., Lesesne, H. R., Carson, S. W., Songer, S. S., Pritchard, J. F., et al. (1995). Comparison of quantitative methods to assess hepatic function: Pugh's classification, indocyanine green, antipyrine, and dextromethorphan. *Pharmacotherapy* 15, 693–700

Filmann, N., Walter, D., Schadde, E., Bruns, C., Keck, T., Lang, H., et al. (2019). Mortality after liver surgery in germany. *The British journal of surgery* 106, 1523–1529. doi:10.1002/bjs.11236

Gadano, A., Hadengue, A., Vachiery, F., Moreau, R., Sogni, P., Soupison, T., et al. (1997). Relationship
between hepatic blood flow, liver tests, haemodynamic values and clinical characteristics in patients
with chronic liver disease. *Journal of gastroenterology and hepatology* 12, 167–171. doi:10.1111/j.
1440-1746.1997.tb00401.x

Gilmore, I. T., Marigold, J. H., and Thompson, R. P. (1982). Half-life time or clearance of indocyanine
green in patients with liver disease. *Hepato-gastroenterology* 29, 55–57

Gorowska-Kowolik, K., Chobot, A., and Kwiecien, J. (2017). javax.xml.bind.jaxbelement@77416ce1, c
methacetin breath test for assessment of microsomal liver function: Methodology and clinical application.
Gastroenterology research and practice 2017, 7397840. doi:10.1155/2017/7397840

Grainger, S. L., Keeling, P. W., Brown, I. M., Marigold, J. H., and Thompson, R. P. (1983). Clearance and
non-invasive determination of the hepatic extraction of indocyanine green in baboons and man. *Clinical
science (London, England : 1979)* 64, 207–212

Grundmann, U., Ziehmer, M., Raahimi, H., Altmayer, P., Larsen, R., and Büch, H. P. (1992). [effect of the
volatile anesthetics halothane, enflurane and isoflurane on liver circulation in the human]. *Anasthesiologie,
Intensivmedizin, Notfallmedizin, Schmerztherapie : AINS* 27, 406–413. doi:10.1055/s-2007-1000324

Grzegorzewski, J., Brandhorst, J., Green, K., Eleftheriadou, D., Duport, Y., Barthorscht, F., et al. (2021).
Pk-db: pharmacokinetics database for individualized and stratified computational modeling. *Nucleic
acids research* 49, D1358–D1364. doi:10.1093/nar/gkaa990

Gu, J., Zhang, E., Liang, B., Zhang, Z., Chen, X., and Huang, Z. (2020). Effectiveness comparison of
indocyanine green retention test with the cirrhotic severity scoring in evaluating the pathological severity
of liver cirrhosis in patients with hepatocellular carcinoma and child-pugh grade a liver function. *World
journal of surgical oncology* 18, 79. doi:10.1186/s12957-020-01854-3

Hackl, C., Schlitt, H. J., Renner, P., and Lang, S. A. (2016). Liver surgery in cirrhosis and portal
hypertension. *World journal of gastroenterology* 22, 2725–2735. doi:10.3748/wjg.v22.i9.2725

Haegele, S., Reiter, S., Wanek, D., Offensperger, F., Pereyra, D., Stremitzer, S., et al. (2016). Perioperative
non-invasive indocyanine green-clearance testing to predict postoperative outcome after liver resection.
PloS one 11, e0165481. doi:10.1371/journal.pone.0165481

Herold, C., Heinz, R., Radespiel-Tröger, M., Schneider, H. T., Schuppan, D., and Hahn, E. G. (2001).
Quantitative testing of liver function in patients with cirrhosis due to chronic hepatitis c to assess disease
severity. *Liver* 21, 26–30. doi:10.1034/j.1600-0676.2001.210104.x

Hucka, M., Bergmann, F. T., Chaouiya, C., Dräger, A., Hoops, S., Keating, S. M., et al. (2019). The
systems biology markup language (sbml): Language specification for level 3 version 2 core release 2.
Journal of Integrative Bioinformatics

ICRP (2002). Basic anatomical and physiological data for use in radiological protection: reference values.
a report of age- and gender-related differences in the anatomical and physiological characteristics of
reference individuals. icrp publication 89. *Annals of the ICRP* 32, 5–265

Jin, S., Fu, Q., Wuyun, G., and Wuyun, T. (2013). Management of post-hepatectomy complications. *World
journal of gastroenterology* 19, 7983–7991. doi:10.3748/wjg.v19.i44.7983

Kamimori, G. H., Sirisuth, N., Greenblatt, D. J., and Eddington, N. D. (2000). The influence of the
menstrual cycle on triazolam and indocyanine green pharmacokinetics. *Journal of clinical pharmacology*
40, 739–744. doi:10.1177/00912700022009495

Keating, S. M., Waltemath, D., König, M., Zhang, F., Dräger, A., Chaouiya, C., et al. (2020). Sbml level 3:
an extensible format for the exchange and reuse of biological models. *Molecular systems biology* 16,
e9110. doi:10.15252/msb.20199110

Keiding, S., Ott, P., and Bass, L. (1993). Enhancement of unbound clearance of icg by plasma proteins,
demonstrated in human subjects and interpreted without assumption of facilitating structures. *Journal of*
hepatology 19, 327–344

Kitano, S. and Kim, Y. I. (1997). Icg clearance in assessing cirrhotic patients with hepatocellular carcinoma
for major hepatic resection. *HPB surgery : a world journal of hepatic, pancreatic and biliary surgery* 10,
182–183. doi:10.1155/1997/69231

Klockowski, P. M., Lener, M. E., Sirgo, M. A., and Rocci, M. L. (1990). Comparative evaluation of the
effects of isradipine and diltiazem on antipyrine and indocyanine green clearances in elderly volunteers.
Clinical pharmacology and therapeutics 48, 375–380. doi:10.1038/clpt.1990.165

König, M. (2021a). matthiaskoenig/sbmlsim: 0.1.14 - sbml simulation made easy doi:10.5281/zenodo.
4569155

König, M. (2021b). matthiaskoenig/sbmlutils: 0.4.12 - python utilities for sbml doi:10.5281/zenodo.
4595467

König, M. and Rodriguez, N. (2019). matthiaskoenig/cy3sbml: cy3sbml-v0.3.0 - SBML for Cytoscape
doi:10.5281/zenodo.3451319

Leevy, C. M., Mendenhall, C. L., Lesko, W., and Howard, M. M. (1962). Estimation of hepatic blood flow
with indocyanine green. *The Journal of clinical investigation* 41, 1169–1179. doi:10.1172/JCI104570

Leevy, C. M., Smith, F., Longueville, J., Paumgartner, G., and Howard, M. M. (1967). Indocyanine green
clearance as a test for hepatic function. evaluation by dichromatic ear densitometry. *JAMA* 200, 236–240

Martin, J. F., Mikulecky, M., Blaschike, T. F., Waggoner, J. G., Vergalla, J., and Berk, P. D. (1975).
Differences between the plasma indocyanine green disappearance rates of normal men and women.
Proceedings of the Society for Experimental Biology and Medicine. Society for Experimental Biology
and Medicine (New York, N.Y.) 150, 612–617

Martin, J. F., Vierling, J. M., Wolkoff, A. W., Scharschmidt, B. F., Vergalla, J., Waggoner, J. G., et al.
(1976). Abnormal hepatic transport of indocyanine green in gilbert's syndrome. *Gastroenterology* 70,
385–391

Meijer, D. K., Weert, B., and Vermeer, G. A. (1988). Pharmacokinetics of biliary excretion in man. vi.
indocyanine green. *European journal of clinical pharmacology* 35, 295–303

Mukherjee, S., Rogers, M. A. M., and Buniak, B. (2006). Comparison of indocyanine green clearance with
child's-pugh score and hepatic histology: a multivariate analysis. *Hepato-gastroenterology* 53, 120–123

Møller, S., Hillingsø, J., Christensen, E., and Henriksen, J. H. (1998). Arterial hypoxaemia in cirrhosis:
fact or fiction? *Gut* 42, 868–874. doi:10.1136/gut.42.6.868

Møller, S., la Cour Sibbesen, E., Madsen, J. L., and Bendtsen, F. (2019). Indocyanine green retention
test in cirrhosis and portal hypertension: Accuracy and relation to severity of disease. *Journal of*
gastroenterology and hepatology 34, 1093–1099. doi:10.1111/jgh.14470

Niemann, C. U., Henthorn, T. K., Krejcie, T. C., Shanks, C. A., Enders-Klein, C., and Avram, M. J.
(2000). Indocyanine green kinetics characterize blood volume and flow distribution and their alteration
by propranolol. *Clinical pharmacology and therapeutics* 67, 342–350. doi:10.1067/mcp.2000.104945

Ohwada, S., Kawate, S., Hamada, K., Yamada, T., Sunose, Y., Tsutsumi, H., et al. (2006). Perioperative
real-time monitoring of indocyanine green clearance by pulse spectrophotometry predicts remnant liver
functional reserve in resection of hepatocellular carcinoma. *The British journal of surgery* 93, 339–346.
doi:10.1002/bjs.5258

Okochi, O., Kaneko, T., Sugimoto, H., Inoue, S., Takeda, S., and Nakao, A. (2002). Icg pulse
spectrophotometry for perioperative liver function in hepatectomy. *The Journal of surgical research* 103,
109–113. doi:10.1006/jsre.2001.6328

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., et al. (2011). Scikit-learn:
Machine learning in Python. *Journal of Machine Learning Research* 12, 2825–2830

Pind, M.-L. L., Bendtsen, F., Kallemose, T., and Møller, S. (2016). Indocyanine green retention test (icg-r15)
as a noninvasive predictor of portal hypertension in patients with different severity of cirrhosis. *European*
journal of gastroenterology & hepatology 28, 948–954. doi:10.1097/MEG.0000000000000611

Pugh, R. N., Murray-Lyon, I. M., Dawson, J. L., Pietroni, M. C., and Williams, R. (1973). Transection
of the oesophagus for bleeding oesophageal varices. *The British journal of surgery* 60, 646–649.
doi:10.1002/bjs.1800600817

Renner, E., Wietholtz, H., Huguenin, P., Arnaud, M. J., and Preisig, R. (1984). Caffeine: a model compound
for measuring liver function. *Hepatology (Baltimore, Md.)* 4, 38–46. doi:10.1002/hep.1840040107

Rubin, T. M., Heyne, K., Luchterhand, A., Bednarsch, J., W R Vondran, F., Polychronidis, G., et al. (2017).
Kinetic validation of the limax test during 10 000 intravenous , javax.xml.bind.jaxbelement@10fb8f50,
c-methacetin breath tests. *Journal of breath research* 12, 016005. doi:10.1088/1752-7163/aa820b

Sakka, S. G. (2018). Assessment of liver perfusion and function by indocyanine green in the perioperative
setting and in critically ill patients. *Journal of clinical monitoring and computing* 32, 787–796. doi:10.
1007/s10877-017-0073-4

Schuppan, D. and Afdhal, N. H. (2008). Liver cirrhosis. *Lancet (London, England)* 371, 838–851.
doi:10.1016/S0140-6736(08)60383-9

Seyama, Y. and Kokudo, N. (2009). Assessment of liver function for safe hepatic resection. *Hepatology*
research : the official journal of the Japan Society of Hepatology 39, 107–116. doi:10.1111/j.1872-034X.
2008.00441.x

Somogyi, E. T., Bouteiller, J.-M., Glazier, J. A., König, M., Medley, J. K., Swat, M. H., et al. (2015).
libroadrunner: a high performance sbml simulation and analysis library. *Bioinformatics* 31, 3315–3321

Soons, P. A., De Boer, A., Cohen, A. F., and Breimer, D. D. (1991). Assessment of hepatic blood flow in
healthy subjects by continuous infusion of indocyanine green. *British journal of clinical pharmacology*
32, 697–704

Stockmann, M., Lock, J. F., Riecke, B., Heyne, K., Martus, P., Fricke, M., et al. (2009). Prediction of
postoperative outcome after hepatectomy with a new bedside test for maximal liver function capacity.
Annals of surgery 250, 119–125. doi:10.1097/SLA.0b013e3181ad85b5

Sunagawa, Y., Yamada, S., Kato, Y., Sonohara, F., Takami, H., Inokawa, Y., et al. (2021). Perioperative
assessment of indocyanine green elimination rate accurately predicts postoperative liver failure in patients
undergoing hepatectomy. *Journal of hepato-biliary-pancreatic sciences* 28, 86–94. doi:10.1002/jhbp.833

Thomas, M. N., Weninger, E., Angele, M., Bösch, F., Pratschke, S., Andrassy, J., et al. (2015).
Intraoperative simulation of remnant liver function during anatomic liver resection with indocyanine
green clearance (limon) measurements. *HPB : the official journal of the International Hepato Pancreato*
Biliary Association 17, 471–476. doi:10.1111/hpb.12380

Urso, R., Blardi, P., and Giorgi, G. (2002). A short introduction to pharmacokinetics. *European review for*
medical and pharmacological sciences 6, 33–44

Vaubourdolle, M., Gufflet, V., Chazouillères, O., Giboudeau, J., and Poupon, R. (1991). Indocyanine
green-sulfobromophthalein pharmacokinetics for diagnosing primary biliary cirrhosis and assessing
histological severity. *Clinical chemistry* 37, 1688–1690

Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., et al. (2020).
SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. *Nature Methods* 17, 261–272.
doi:10.1038/s41592-019-0686-2

Wakabayashi, H., Ishimura, K., Izuishi, K., Karasawa, Y., and Maeta, H. (2004). Evaluation of liver
function for hepatic resection for hepatocellular carcinoma in the liver with damaged parenchyma. *The*
Journal of surgical research 116, 248–252. doi:10.1016/j.jss.2003.09.015

Wang, Y.-Y., Zhao, X.-H., Ma, L., Ye, J.-Z., Wu, F.-X., Tang, J., et al. (2018). Comparison of the ability of
child-pugh score, meld score, and icg-r15 to assess preoperative hepatic functional reserve in patients
with hepatocellular carcinoma. *Journal of surgical oncology* 118, 440–445. doi:10.1002/jso.25184

Wheeler, H. O., Cranston, W. I., and Meltzer, J. I. (1958). Hepatic uptake and biliary excretion of
indocyanine green in the dog. *Proceedings of the Society for Experimental Biology and Medicine.*
Society for Experimental Biology and Medicine (New York, N.Y.) 99, 11–14

Wong, J. S.-W., Wong, G. L.-H., Chan, A. W.-H., Wong, V. W.-S., Cheung, Y.-S., Chong, C.-N., et al.
(2013). Liver stiffness measurement by transient elastography as a predictor on posthepatectomy
outcomes. *Annals of surgery* 257, 922–928. doi:10.1097/SLA.0b013e318269d2ec

Figure 1. Model overview: **A:** Whole-body model. The whole-body PBPK model for ICG consists of venous blood, arterial blood, lung, liver, gastrointestinal tract and rest compartment (accounting for organs not modeled in detail) and the systemic blood circulation connecting these compartments. **B:** Liver model. ICG in the liver plasma compartment is taken up into the liver tissue (hepatocytes). Subsequently hepatic ICG is excreted in the bile from where it is excreted in the feces. No metabolization of ICG takes place in the liver. **C:** Modeling liver cirrhosis. Liver cirrhosis was modeled as a combination of tissue loss and hepatic shunts (see main text for details). **D:** Modeling hepatectomy. Hepatectomy was modeled as a removal of tissue volume with corresponding vessels (see main text for details)

Figure 2. Model prediction of ICG time courses in healthy subjects: **A-D:** Venous concentration after bolus ICG administration (Andersen et al., 1999; Grundmann et al., 1992; Kamimori et al., 2000; Klockowski et al., 1990). **E:** Arterial concentration after bolus ICG administration (Niemann et al., 2000). **F:** Venous concentration during an ICG infusion protocol (2.0, 0.5, 1.0 mg/min, 40 minutes each) (Soons et al., 1991). **G, H:** Venous concentration and biliary excretion rate after 3 different ICG doses (0.5, 1.0, 2.0 mg/kg) (Meijer et al., 1988; Chijiiwa et al., 2000). **I:** Hepatic venous and arterial concentration during constant ICG infusion (Leevy et al., 1962). **J-L:** ICG extraction ratio during constant infusion (Leevy et al., 1962; Grainger et al., 1983).

Figure 3. Dose dependency of ICG pharmacokinetic parameters: **A-D:** Dose dependency of ICG pharmacokinetic parameters in controls and three different degrees of cirrhosis. **E-H:** Dose dependency of ICG-kel, ICG-clearance, ICG-$t_{1/2}$ in healthy subjects with clinical data (Martin et al., 1975, 1976; Meijer et al., 1988).

Figure 4. Dependency of ICG pharmacokinetic parameters on cirrhosis.: Simulations of 4 specific cirrhosis degrees are indicated by stars. **A-D:** Dependency of ICG pharmacokinetic parameters on the degree of shunting (green), degree of tissue loss (yellow) and degree of cirrhosis (black-blue-red). **E:** Correlation between ICG-R20 and ICG-$t_{1/2}$ in cirrhotic and control subjects (Cherrick et al., 1960). **F:** Correlation between ICG-R20 and ICG-kel in cirrhotic and control subjects (Caesar et al., 1961). **G:** Correlation between ICG-PDR after an ICG dose of 0.5 mg/kg and 5.0 mg/kg in control subjects and subjects with various liver diseases (Leevy et al., 1967). **H:** Correlation between ICG-clearance after a bolus administration and during a constant infusion of ICG in cirrhotic subjects (Burns et al., 1991).

Figure 5. Mapping of model cirrhosis degree on CTP-score: **A:** Mapping based on ICG clearance (Figg et al., 1995; Møller et al., 1998). **B:** Mapping based on ICG-kel (Møller et al., 2019; Herold et al., 2001). **C:** Mapping based on ICG-R15 (Møller et al., 2019). **D:** Mapping based on ICG-thalf (Møller et al., 2019). **E:** Resulting $f_{cirrhosis}$ values for each CTP-class combining the information from the mappings based on individual ICG pharmacokinetic parmeters (Control: $f_{cirrhosis}$ = 0.0; Mild cirrhosis: 0.41; Moderate cirrhosis: 0.70; Severe cirrhosis: 0.82).

Figure 6. Model prediction of ICG time courses in subjects with cirrhosis: **A, B:** Venous concentration after a bolus ICG administration in a healthy subject and a cirrhotic patient ($f_{cirrhosis}$ was set to 0.7 corresponding to moderate cirrhosis) (Burns et al., 1991). **C:** Venous concentration during a constant ICG infusion in healthy and cirrhotic subjects (Caesar et al., 1961). **D, E:** Hepatic venous and arterial ICG concentration and ICG extraction ratio in a cirrhotic patient ($f_{cirrhosis}$ was set to 0.54 corresponding to mild-moderate cirrhosis) (Keiding et al., 1993). **F-H:** ICG extraction ratio in cirrhotic, hepatitis and control subjects during a constant ICG infusion (Leevy et al., 1962; Gadano et al., 1997; Caesar et al., 1961).

Figure 7. Model prediction of ICG pharmacokinetic parameters in hepatectomy under varying degree of cirrhosis:**A-D:** Dependency of postoperative ICG pharmacokinetic parameters on the resected volume in 4 different degrees of cirrhosis. **E:** Dependency of postoperative change in ICG-PDR on the resected volume (Thomas et al., 2015; Stockmann et al., 2009). **F:** Correlation between the measured postoperative ICG-kel and the estimated postoperative ICG-kel (product of preoperative ICG-kel and the future liver remnant) (Ohwada et al., 2006; Okochi et al., 2002; Sunagawa et al., 2021). **G:** Correlation between postoperative ICG-PDR and intraoperative ICG-PDR during trial clamping (Thomas et al., 2015). **H:** Correlation between postoperative and preoperative ICG-PDR (Thomas et al., 2015). Simulations for **E-H:** were performed for varying resection rates in healthy subjects and three different degrees of cirrhosis.

Figure 8. Classification of survivors/non-survivors after hepatectomy: All classification models predict non-survival after hepatectomy. Classification data (n=141) from (Seyama and Kokudo, 2009; Wakabayashi et al., 2004). **A:** PBPK model based prediction of postoperative ICG-R15 in survivors and non-survivors. $f_{cirrhosis}$ was estimated for every subject based on preoperative ICG-R15 and post-operative ICG-R15 calculated using the respective resection rate (used in PBPK1). Corresponding measured preoperative ICG-R15 as inset (used in Data1A). **B:** Dependency of the evaluation metrics of the cross-validated classification model PBPK1 on the predicted postoperative ICG-R15 cutoff (mean ± SD). PPV: positive predictive value; NPV: negative predictive value **C:** ROC curve for the prediction of non-survival after hepatectomy using the complete dataset with cross-validation (mean ± SD) for classification models Data1A, Data1B and PBPK1. **D:** Predicted postoperative ICG-R15 and survival status depending on the resection rate and $f_{cirrhosis}$ (used in PBPK2). $f_{cirrhosis}$ was estimated for every subject individually based on preoperative ICG-R15 and post-operative ICG-R15 calculated using the respective resection rate. Shaded blue areas correspond to respective predicted postoperative ICG-R15. **E:** Decision boundary of the two-dimensional classification model PBPK2 based on the resection rate and $f_{cirrhosis}$ using the complete dataset. White area: predicted survivor; blue area: predicted non-survivor. **F:** ROC curve for the prediction of non-survival after hepatectomy using the complete dataset with cross-validation (mean ± SD) for classification models Data2 and PBPK2.

Table 1. Overview of curated clinical studies.

Study	PK-DB	PMID	Protocol	Body weight	Fit	Data used in fit	Description
Andersen1999	PKDB00386	10499483	Bolus: 0.5 mg/kg	Contr.: 83.4 kg; Panc. 58.2 kg	✓	ICG time course (plasma)	Pharmacokinetic time course of ICG-disappearance in chronic pancreatitis and healthy subjects.
Burns1991	PKDB00388	1848168	Bolus: 0.5 mg/kg; Infusion: 0.25 mg/min	NR	✓	ICG time course (plasma)	ICG pharmacokinetics in healthy subjects and patients with liver disease.
Caesar1961	PKDB00389	13689739	Bolus: 0.5 mg/kg; Infusion: 0.5 mg/min	NR	✓	ICG extraction-ratio	Measuring hepatic blood flow and assessing hepatic function by ICG pharmacokinetics.
Cherrik1960	PKDB00390	13809697	Bolus: 0.5 mg/kg	NR	-	-	ICG pharmacokinetics in healthy subjects and patients with liver disease.
Chijiiwa2000	PKDB00391	10773154	Bolus: 0.5 mg/kg	NR	✓	ICG time course (bile excretion)	Biliary excretion of ICG and ATP-dependency of ICG pharmacokinetics.
Figg1995	PKDB00393	8602375	Bolus: 0.5 mg/kg	NR	-	-	Comparison of methods to assess hepatic function including CTP-score and ICG-clearance
Gadano1997	PKDB00394	9083919	Infusion: 0.4 mg/min (controls); 0.8 mg/min (cirrhotics) with priming dose (24 mg controls; 12 mg cirrhotics).	NR	✓	-	ICG-clearance and extraction-ratio and their dependency on hepatic bloodflow in healthy subjects and in patients of hepatic fibrosis and cirrhosis.
Grainger1983	PKDB00395	6822056	Bolus: 0.25 mg/kg	NR	✓	ICG extraction-ratio	Non-invasive measurement of hepatic blood flow using ICG extraction-ratio.
Grundmann1992	PKDB00396	1482735	Bolus: 0.3 mg/kg	NR	✓	ICG time course (plasma)	Effect of anesthetics on ICG-pharmacokinetics and hepatic blood flow.
Herold2001	PKDB00397	11169069	Bolus: 0.5 mg/kg	NR	-	-	Comparison of liver function tests. Correlation between CTP-score and ICG-kel in healthy subjects and cirrhotic patients.
Kamimori2000	PKDB00299	10883415	Bolus: 0.5 mg/kg	58.6 ± 11.2 kg	✓	ICG time course (plasma)	Effect of the menstrual cycle on ICG pharmacokinetics.
Keiding1993	PKDB00402	8151094	Infusion: 0.08 mg/min	NR	-	-	Effect of changing plasma protein concentrations on ICG-extraction.
Klockowski1990	PKDB00403	2146057	Bolus: 0.5 mg/kg	NR	✓	ICG time course (plasma)	Effect of isradipine and diltiazem on ICG-pharmacokinetics.
Leevy1962	PKDB00404	14463639	Infusion: 0.3 and 1.5 mg/min/m^2 with priming dose (10 mg).	NR	✓	ICG extraction-ratio; ICG time course (hv and ar)	Estimation of hepatic blood flow using ICG.
Leevy1967	PKDB00405	6071462	Bolus: 0.5 mg/kg and 5.0 mg/kg	NR	-	-	Estimation of liver function with ICG. Dose dependency of ICG-PDR.
Martin1975	PKDB00406	1208580	Bolus: 0.5, 2.0, 3.5 and 5.0 mg/kg	NR	-	-	Differences in ICG-pharmacokinetics in men and women at varying ICG-doses.
Martin1976	PKDB00407	814028	Bolus: 0.5, 2.0, 3.5 and 5.0 mg/kg	NR	-	-	ICG-pharmacokinetics in patients with Gilbert's Syndrome.
Meijer1988	PKDB00408	3181282	Bolus: 0.5, 1.0 and 2.0 mg/kg	64.8 kg	✓	ICG time course (plasma and bile excretion)	Biliary excretion of ICG at different doses.
Moeller1998	PKDB00409	9691928	Infusion: Rate not reported	72.7 (40-115) kg	-	-	Arterial hypoxaemia in cirrhosis. Correlation between ICG-clearance and CTP-score.
Moeller2019	PKDB00410	30221390	Infusion: 0.2 mg/min with priming dose (2 mg), Bolus: 0.5 mg/kg	79.2 ± 18.5 kg	-	-	Correlation between ICG-pharmacokinetics and CTP-score.
Niemann2000	PKDB00414	10801242	Bolus: 10 mg	80 ± 17 kg	✓	ICG time course (plasma)	ICG- pharmacokinetic time course under administration of propranolol.
Ohwada2006	PKDB00412	16498606	Bolus: 20 mg	NR	-	-	Prediction of postoperative liver functional capacity based on ICG-pharamcokinetics.
Okochi2002	PKDB00415	11855925	Bolus: 20 mg	NR	-	-	Comparison of preoperative and postoperative ICG-pharmacokinetics. Prediction of postoperative liver functional capacity based on ICG-pharamcokinetics.
Seyama2009	PKDB00416	19208031	Bolus: 0.5 mg/kg	NR	-	-	Assessment of liver function for safe hepatic resection. Prediction of survival after hepatectomy based on ICG-R15.
Soons1991	PKDB00411	1768562	Infusion: 2.0, 0.5, 1.0 mg/min (consecutive)	72 ± 6 kg	✓	ICG time course (plasma)	Assessment of hepatic blood flow in healthy subjects by continuous infusion of ICG.
Stockmann2009	PKDB00417	19561474	Bolus: 0.5 mg/kg	NR	-	-	Prediction of postoperative liver functional capacity based on ICG-pharamcokinetics.
Sunagawa2021	PKDB00418	33052632	Bolus: 20 mg	NR	-	-	Prediction of postoperative liver failure based on ICG-kel measurements.
Thomas2015	PKDB00419	25581073	Bolus: 0.25 mg/kg	NR	-	-	Intraoperative prediction of postoperative liver functional capacity using trial-clamping and ICG-pharmacokinetics.
Wakabayashi2004	PKDB00420	15013363	Bolus: 0.5 mg/kg	NR	-	-	Correlation between preoperative ICG-R15 and estimated liver remnant. Prediction of survival based on ICG-R15.

Table 2. Overview of key model parameters.

Parameter	Description	Value	Unit	Fitted
BW	Body weight	75	kg	-
$COBW$	Cardiac output per body weight	0.83	ml/s/kg	-
QC	Cardiac output $(BW \cdot COBW)$	3.75	l/min	-
HCT	Hematocrit	0.51	-	-
$Fblood$	Fraction of organ volume that is blood vessels	0.02	-	-
$FVgi$	Fractional tissue volume gastrointestinal tract	0.0171	l/kg	-
$FVli$	Fractional tissue volume liver	0.0210	l/kg	-
$FVlu$	Fractional tissue volume lung	0.0076	l/kg	-
$FVve$	Fractional tissue volume venous blood	0.0587	l/kg	-
$FVar$	Fractional tissue volume arterial blood	0.0184	l/kg	-
LI_ICGIM_Vmax	V_{max} of liver import	3.70E-2	mmole/min/l	✓
LI_ICGIM_Km	K_m of liver import	2.17E-2	mM	✓
$LI\_ICGLI2CA\_Vmax$	V_{max} of bile excretion	9.44E-4	mmole/min/l	✓
$LI\_ICGLI2CA\_km$	K_m of bile excretion	1.24E-2	mM	✓
$LI\_ICGLI2BI\_Vmax$	V_{max} of bile transport	1.14E-4	1/min	✓

Table 3. Evaluation metrics for classification models of survival after hepatectomy. Evaluation metrics of classification model are values for the model fitted with the complete dataset. Mean $\pm$ SD of cross validation reported in brackets.

Classification Model	Data1A	Data1B	PBPK1	Data2	PBPK2
Features	Preoperative ICG-R15	Postoperative ICG-R15 (calculated)	Postoperative ICG-R15 (predicted)	Preoperative ICG-R15 & Resection rate	$f_{cirrhosis}$ & Resection rate
ROC AUC	0.663 (0.555 $\pm$ 0.072)	0.517 (0.481 $\pm$ 0.052)	0.862 (0.753 $\pm$ 0.090)	0.880 (0.767 $\pm$ 0.090)	0.879 (0.766 $\pm$ 0.092)
Balanced accuracy	0.562 (0.555 $\pm$ 0.072)	0.515 (0.481 $\pm$ 0.052)	0.761 (0.753 $\pm$ 0.090)	0.785 (0.767 $\pm$ 0.090)	0.805 (0.766 $\pm$ 0.092)
F1-score	0.267 (0.243 $\pm$ 0.152)	0.143 (0.118 $\pm$ 0.126)	0.611 (0.587 $\pm$ 0.136)	0.625 (0.593 $\pm$ 0.133)	0.650 (0.592 $\pm$ 0.139)
Precision (PPV)	0.462 (0.443 $\pm$ 0.295)	0.300 (0.192 $\pm$ 0.227)	0.550 (0.538 $\pm$ 0.152)	0.521 (0.515 $\pm$ 0.145)	0.542 (0.509 $\pm$ 0.153)
Recall	0.188 (0.182 $\pm$ 0.133)	0.094 (0.170 $\pm$ 0.273)	0.688 (0.681 $\pm$ 0.172)	0.781 (0.739 $\pm$ 0.170)	0.812 (0.747 $\pm$ 0.169)
NPV	0.797 (0.792 $\pm$ 0.067)	0.779 (0.745 $\pm$ 0.133)	0.901 (0.898 $\pm$ 0.057)	0.925 (0.912 $\pm$ 0.056)	0.935 (0.915 $\pm$ 0.054)