Article

Tensor Decomposition-based Unsupervised Feature Extraction Succeeded in Identification of Differentially Expressed Transcripts from Redundant *de novo* Transcriptome of *Planarian*

Makoto Kashima [1,†] **, Nobuyoshi Kumagai** [2]**, Hiromi Hirata**[3] **and Y-h. Taguchi** [4*]

1. Department of Biology, Aoyama Gakuin Univeristy, 150-8366, Tokyo, Japan
2. Department of Life Science, Gakushuin University, 171-8588, Tokyo Japan; nobuyoshi.kumagai@hirosaki-u.ac.jp
3. Department of Biology, Aoyama Gakuin Univeristy, 150-8366, Tokyo, Japan; hihirata@chem.aoyama.ac.jp
4. Department of Physics, Chuo University, 1-13-27 Kasuga, Bunkyo-ku, 112-8551, Tokyo, Japan; tag@granular.com
* Correspondence: tag@granular.com
† Current address: Department of Biomolecular Science, Faculty of Science, Toho University, Miyama 2-2-1, Funabashi, 274-8510, Chiba, Japan;makoto.kashima@sci.toho-u.ac.jp

Abstract: RNA-Seq data analysis of non-model organisms is often difficult because of the lack of a well-annotated genome. However, in non-model organisms, contigs can be generated by *de novo* assembling. This can result in a large number of transcripts, making it difficult to easily remove redundancy. A large number of transcripts can also lead to difficulty in the recognition of differentially expressed transcripts (DETs) between more than two experimental conditions, because *P*-values must be corrected by considering multiple comparison corrections whose effect is enhanced as the number of transcripts increases. Heavily corrected *P*-values often fail to take sufficiently small *P*-values as significant. In this study, we applied a recently proposed tensor decomposition (TD)-based unsupervised feature extraction (FE) to the RNA-seq data obtained for a non-model organism, planarian *Dugesia japonica*; Although we used de novo assembled transcriptome reference with high redundancy, we successfully obtained a larger number of transcripts whose expression was altered between normal and defective samples as well as during time development than those identified by a conventional method. TD-based unsupervised FE is expected to be an effective tool that can identify a substantial number of DETs, even when a poorly annotated genome is available.

Keywords: Planarian; Tensor Decomposition; Feature Selection; unsupervised learning; machine learning

1. Introduction

Identification of differentially expressed transcripts (DETs) [1,2] between more than two distinct experimental conditions is the starting point of RNA-seq data analysis, as DETs are expected to be related to the experimental conditions considered, for example, diseases. Once DETs are successfully identified, it is possible to study the biological properties that are enriched in a set through a statistical test such as gene ontology enrichment analysis [3]. Thus, unless DETs can be successfully identified, it is very difficult to make use of RNA-seq data to identify the biological processes that take place during experiments using RNA-seq datasets.

DET identification in model organisms has been well-established. Once short reads are successfully mapped to the genome, we can concentrate those mapped to well-annotated regions; for example, protein-coding genes. Then, the number of reads can be summed up within the considered regions, for example, the gene body, then it is possible to estimate the amount of expression of individual genes [4]. Another way is that reference transcriptome sequences are generated based on gene models [2,5], following by mapping short reads

onto it. Then, quantification is performed considering information of isoform and reads mapped onto several reference sequences [2,6–8]. The latter approaches tend to be superior in quantification accuracy than the former [9]. In contrast, DET identification of non-model organisms is not straightforward. Because of the lack of a well-annotated genome, no short reads can be mapped to it. Instead of the genome, it is possible to assemble the contigs by assembling *de novo*, and short reads can then be mapped toward these contigs. However, there is one problem with this strategy; it is never possible to be confident that the parts of the assembled contigs are unique. Some contigs may overlap with some parts of another contig [10,11]. This results in so-called multiple mapping, which drastically reduces the accuracy of the estimated amount of transcript [12]. In addition, the number of contigs can often be far larger than the true number of transcripts because of the above-mentioned redundancy of contigs. e.g. the number of contigs can often be in millions. This causes serious problems of drastic reduction of statistical power in DET identification. In popular DET identification methods such as DESeq2 [13], the statistical tests check how significant the observed distinct expression of a transcript is under the null hypothesis and calculate *P*-values, which must be corrected by considering multiple comparison corrections. As the number of multiple comparisons is equivalent to the number of contigs, too many contigs (e.g. millions) results in false negative identification of DETs.

To address this problem, we applied tensor decomposition (TD)-based unsupervised feature extraction (FE) [14] to the RNA-seq data. As proof of concept, here, we analyzed RNA-Seq data of a planarian *Dugesia japonica*, one of whose *de novo* trascriptome assembly contains as many as 2.8×10^5 [15], which is much larger than the expected number of true genes of the planarian [16]. TD based unsupervised FE could successfully select a larger number of contigs whose expression was altered during time development as well as was distinct between normal and defective samples than a conventional method DESeq2. This suggests the usefulness of TD-based unsupervised FE when it is applied to RNA-seq data analysis of non-model organisms, from which too many redundant contigs are often obtained.

2. Materials and Methods

2.1. Maintenance of planarian

A clonal strain of planarian *D. japonica*, a sexualizing super planarian (2n = 16), was used in this study [17]. The planarians were fed edible chicken liver once or twice a week. The planarians that were approximately 5 mm in length were starved for at least one week prior to the following experiments.

2.2. Double-stranded RNA (dsRNA) synthesis

dsRNA was synthesised as described by [18]. We prepared templates flanked by the T7 promoter for dsRNA synthesis by using polymerase chain reaction (PCR) for each EST (*Djpiwi A* [*Dj_aH_000_03609HH*], *DjpiwiB* [*Dj_aH_221_M14*], *DjpiwiC* [*Dj_aH_000_05977HH*], *Djhp1-1* [*Dj_aH_000_01636HH*], *Djima1-1* [*Dj_aH_313_F03*]). The primers for the PCR reaction were as follows (5′ to 3′): Zap Linker + T7: GATCACTAATACGACTCAC-TATAGGGGAATTCGGCACGAGG M13 Rev: GTTTTCCCAGTCACGACGTTGTAA.

2.3. Feeding RNA interference

RNA interference was conducted as described previously [18]. dsRNA-containing food consisting of 25 μL of chicken liver solution, 6 μL of 2% agarose, and 6.5 μL of 2.0 μg/μL dsRNA was synthesised in vitro. We fed the planarians twice at intervals of two days. Control animals were fed food containing eGFP dsRNA. Six individual planarians were sacrificed every day from day 1 to 16 after the second feeding.

2.4. Total RNA purification

Total RNA was extracted from each individual planarian using the "Direct-TRI" method [19]. Briefly, a planarian was lysed using TRI Reagent-LS (Molecular Research

Center, Cincinnati, OH, USA). Then, the lysate was purified using an AcroPrep Advance 96-well long tip filter plate for nucleic acid binding (Pall, Port Washington, NY, USA). RNA was eluted with 30 µL nuclease-free water.

2.5. RNA-Seq library preparation and sequencing

3' mRNA-Seq was conducted using the Lasy-Seq ver. 1.1 Protocol (https://sites.google.com/vie seq/) [20,21]. Briefly, 100 ng of total RNA was reverse transcribed using an RT primer with an index and SuperScript IV reverse transcriptase (Thermo Fisher Scientific, Waltham, MA, USA). All RT mixtures of the samples were pooled and purified using an equal volume of AMpure XP beads (Beckman Coulter, Brea, CA, USA) according to the manufacturer's instructions. Second-strand synthesis was conducted on the pooled samples using RNaseH (5 U/μL, Enzymatics, Beverly, MA, USA), and DNA polymerase I (10 U/μL, Enzymatics, Beverly, MA, USA). To avoid carryover of large amounts of rRNAs, the mixture was subjected to RNase treatment using RNase T1 (Thermo Fisher Scientific, Waltham, MA, USA). Then, purification was conducted with a 0.8× volume of AMpure XP beads. Fragmentation, end-repair, and A-tailing were conducted using a 5× WGS fragmentation mix (Enzymatics, Beverly, MA, USA). The adapter for Lasy-Seq was ligated using 5× Ligation Mix (Enzymatics, Beverly, MA, USA), and the adapter-ligated DNA was purified twice with a 0.8× volume of AMpure XP beads. After the optimisation of PCR cycles for library amplification by qPCR using Evagreen, 20× in water (Biotium, Fremont, CA, USA) and the QuantStudio5 Real-Time PCR System (Applied Biosystems, Waltham, MA, USA), the library was amplified using KAPA HiFi HotStart ReadyMix (KAPA BIOSYSTEMS, Wilmington, MA, USA) on the ProFlex PCR System (Applied Biosystems, Waltham, MA, USA). The amplified library was purified using an equal volume of AMpure XP beads. One microliter of the library was then used for electrophoresis using a Bioanalyzer 2100 with the Agilent High Sensitivity DNA kit (Agilent Technologies, Santa Clara, CA, USA) to check for quality. Sequencing of 150 bp paired-end reads was performed using HiSeq X Ten (Illumina, San Diego, CA, USA).

2.6. Mapping and gene quantification

Read 1 reads were processed with fastp (version 0.21.0) [22] using the following parameters: trim_poly_x -w 20 –adapter_sequence=AGATCGGAAGAGCACACGTCTGAACTCCAGTCA –adapter_sequence_r2= AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT-l 31. The trimmed reads were then mapped to a *D. japonica* reference sequence of GJEZ00000000 in the TSA repository , which is a de novo transcriptome assembly generated in a previous study [15] , using BWA mem (version 0.7.17-r1188) [23] with the default parameters. The read count for each gene was calculated with salmon using -l IU, which specifies the library type (version v0.12.0) [6]. All quantification results and sample information of the RNA-seq analysis were deposited as GSE174855 in the GEO repository.

2.7. Mapping to oxford genome

To see if the analysis of contig without genome is reasonable, we mapped the obtained short reads to genome, too. The reference transcriptome sequences were generated based on a draft genome [24] and gene models [25] by using gffread (v0.12.1) [5]. Then, mapping and quantification were conducted as described above.

2.8. Comparison of the de novo transcriptome assembly and oxford references

BLASTN anaysis (version 2.9.0+) [26] was performed using the oxford references as query and the de novo transcriptome assembly as reference. Best hit of each oxford reference sequence was refer to the corresponding contig in the de novo transcriptome assembly.

2.9. TD-based unsupervised FE

The number of reads mapped to individual contigs was formatted as a tensor, $x_{ijtk} \in \mathbb{R}^{N \times 6 \times T \times 6}$, which represents the expression of the ith contig ($N = 278, 167$) or gene ($N = 24,090$) of the kth biological replicates at the tth day after the jth RNAi was performed. The 14th dataset for contig was excluded because it included serious outliers. Thus, $t \leq 13$ corresponds to t days after the treatment, whereas $t \geq 15$ corresponds to $t + 1$ days after the treatment. Then $T = 15$ for contig. Since it was not a case for genes, $T = 16$.

x_{ijtk} is normalised such that $\sum_i x_{ijtk} = 0$ and $\sum_i x_{ijtk}^2 = N$. Then, we applied higher order singular value decomposition (HOSVD) [14] and got

$$x_{ijtk} = \sum_{\ell_1=1}^{6} \sum_{\ell_2=1}^{T} \sum_{\ell_3=1}^{6} \sum_{\ell_4=1}^{N} G(\ell_1 \ell_2 \ell_3 \ell_4) u_{\ell_1 j} u_{\ell_2 t} u_{\ell_3 k} u_{\ell_4 i} \qquad (1)$$

where $G \in \mathbb{R}^{6 \times T \times 6 \times N}$ is the core tensor that represents the weight of the product $u_{\ell_1 j} u_{\ell_2 t} u_{\ell_3 k} u_{\ell_4 i}$. $u_{\ell_1 j} \in \mathbb{R}^{6 \times 6}$. $u_{\ell_2 t} \in \mathbb{R}^{T \times T}$ and $u_{\ell_3 k} \in \mathbb{R}^{6 \times 6}$, and $u_{\ell_4 i} \in \mathbb{R}^{N \times N}$ are singular value matrices that are orthogonal matrices.

First, we needed to identify $u_{\ell_1 j}$, $u_{\ell_2 t}$, and $u_{\ell_3 k}$ that satisfy

- $u_{\ell_1 j}$ that exhibits distinction between normal and defective samples.
- $u_{\ell_2 k}$ with time (days) dependence.
- $u_{\ell_3 k}$ independent of individual biological replicates, k, i.e. with constant values.

After identifying ℓ_1, ℓ_2, and ℓ_3 that satisfy the above requirements, we sought a set of ℓ_4s, Ω_{ℓ_4}, associated with the larger $|G(\ell_1 \ell_2 \ell_3 \ell_4)|$s given ℓ_1, ℓ_2, and ℓ_3, because $u_{\ell_4 i}$s are expected to have larger absolute values for the genes whose expression is associated with the above requirement. Then, we attributed P-values to the ith contig by assuming that $u_{\ell_4 i}$ obeys a multiple Gaussian distribution (null hypothesis)

$$P_i = P_{\chi^2} \left[> \sum_{\ell_4 \in \Omega_{\ell_4}} \left(\frac{u_{\ell_4 i}}{\sigma_{\ell_4}} \right)^2 \right], \qquad (2)$$

where $P_{\chi^2}[> x]$ is the cumulative probability distribution of the χ^2 distribution when the argument takes larger values than x. The obtained P_is are corrected by the BH criterion [14], and contigs associated with an adjusted P_i less than 0.01 were selected.

2.10. DESeq2

DESeq2 [13] (version 1.30.0) was applied to either comparison between two groups of RNAi treatments, that is, *gfp*, *DjpiwiA* and others (*Djhp1-1, Djima1-1, DjpiwiB, DjpiwiC*) or that between two time periods, that is, $t = 1, 2, \cdots, 8$, and others ($t = 9, 10, 11, 12, 13, 15, 16$). We selected genes associated with adjusted P-values less than 0.01, which is the same as that used for TD-based unsupervised FE. All other parameters were taken as the default values.

2.11. GO term enrichment analysis

BLASTX (version 2.9.0+) [26] was used for annotation of planarian contigs/genes. All mouse protein data (GRCm3.8) [27] was used as BLAST subject sequences. Best hit of each transcript was used for the annotation of planarian transcripts. Then, GO term annotation to planarian transcripts was conducted by using biomaRt (version 2.46.0) [28],GO.db (version 3.12.1) [29], and org.Mm.eg.db (version 3.11.4) [30]. GO term enrichment analysis was conducted with a function in R: fisher.test. Then, the adjustment for multiple comparisons against P-values was performed using the Benjamini–Hochberg (BH) method [31] using the p.adjust function in R. Finally, we listed only GO terms of the most offspring with the adjusted P-value < 0.05.

3. Results

The RNA-Seq data were derived from RNAi experiments designed to identify transcripts related to regenerative defects caused by knockdown of *DjpiwiB* and *DjpiwiC* in a planarian *Dugesia japonica*. In addition to a control and three planarian *piwi* genes', we conducted RNAi experiments against two planarian genes, *Djhp1-1* and *Djima1-1*, because they are homologous genes in fruit fly are known to be partners of PIWI proteins [32,33]. The each targeted gene was significantly knocked-down under each RNAi treatment (Fig. 1).

Figure 1. log2(rpm+1) of RNAi target genes at 3, 4, and 5 days into the RNAi treatment. P-value were calculated with t-test.

First, we attempted to identify which $u_{\ell_1 j}$ is associated with the desired property. Figure 2 shows u_{2j} that represents distinction between the RNAi treatments. The phenotype of RNAi of *Djhp1-1*, *Djima1-1*, *DjpiwiB*, and *DjpiwiC* is distinct from that of *gfp* and *DjpiwiA*. While *gfp* is the control treatment and KD of *piwiA* is known not to affect (inhibit) the regeneration of planarians, RNAi of *Djhp1-1*, *Djima1-1*, *DjpiwiB*, and *DjpiwiC* causes regenerative defects [15,34,35]. Thus, u_{2j} coincides with the distinction between normal and defective samples.

Figure 2. u_{2j} that represent 2nd singular value vectors attributed to RNAi experiments that coincides with the distinction between normal and defective samples. That is, *gfp* and KD of *piwiA* is known not to affect (inhibit) the regeneration of planarians and RNAi of *Djhp1-1*, *Djima1-1*, *DjpiwiB*, and *DjpiwiC* causes regenerative defects [15,34,35].

Next, we attempted to identify which $u_{\ell_2 t}$ coincides with time development. Figure 3 shows u_{2t} that exhibits the desired time development. Several days after RNAi treatment, RNAi appeared to have an effect, which then gradually decreased. The previous study showed *DjpiwiB*-knocked-down planarians lose their regenerative ability at seven days after the RNAi treatment [15]. This time point corresponds to the first date when u_{2t} takes positive values (the sign of u_{2t} does not have any meaning because only the product of $G(\ell_1 \ell_2 \ell_3 \ell_4) u_{\ell_1 j} u_{\ell_2 t} u_{\ell_3 k} u_{\ell_4 i}$ matters, not individual terms in the product (see eq.(1)). Thus, this time development seemed to coincide with the experimental procedures. The reason why u_{2t} takes almost zero might be because it requires several days until RNAi starts to affect the transcriptome.

Figure 3. u_{2t} that represent 2nd singular value vectors attributed to days after the treatments. The time point where sign of u_{2t} changes coincides with time point when regenerative defects of *DjpiwiB* RNAi appears.

We also noticed that u_{1k} has almost constant values independent of the biological replicate, k (Fig. 4); this suggests that the expression of genes associated with u_{1k} is likely to be common among the six biological replicates.

Figure 4. u_{1k} that represent 1st singular value vectors attributed to biological replicates. The values indicate there is no bias between the biological replicates.

We then determine which ℓ_4 had the largest $|G(2,2,1,\ell_4)|$. Table 1 shows $G(2,2,1,\ell_4)$. $\ell_4 = 1,3,4$ had a larger $|G|$ value. It is not possible to employ all three ℓ_4s to compute P_i because $|G(1,1,1,1)|$ is much larger than $G(2,2,1,1)$ (not shown here). This means that u_{1i} is more coincident with $\ell_1 = \ell_2 = \ell_3 = 1$, which we are not interested in. Thus, we decided to employ $\ell_4 = 3,4$ to attribute P-values to the ith contig (that is, $\Omega_{\ell_4} = \{3,4\}$ in Eq.(2)). P_is are corrected as described in the Methods section, and contigs with adjusted P_i less than 0.01 were selected.

Table 1. $G(2,2,1,\ell_4)$

ℓ_1	$G(2,2,1,\ell_4)$
1	227.87614
2	-8.36258
3	-113.10063
4	250.20079
5	-43.37827
6	-64.02292
7	37.49047
8	12.84754
9	31.20391
10	-21.70284

After applying TD-based unsupervised FE to RNA-seq data obtained from Planarian as described in the Methods section, we obtained 155 contigs as those whose expression was altered during time development as well as distinct between normal and defective samples. To confirm whether we could successfully identify genes with altered expression, we applied statistical tests that validated significant dependence upon time as well as significant distinction between normal and defective samples. First, we applied the *t* test to determine whether gene expression was distinct between $t \leq 8$ and $t \geq 9$ (Table 2). There were at least non-zero genes expressed differently between $t \leq 8$ and $t \geq 9$.

Table 2. The number of contigs identified by the *t* test to be distinct between $t \leq 8$ and $t \geq 9$.

RNAi	netative	positive
gfp	89	66
Djhp1-1	96	59
Djima1-1	139	16
DjpiwiA	89	66
DjpiwiB	102	53
DjpiwiC	151	4

We also checked whether genes were expressed differently on individual days between normal and defective samples (Table 3). In this case, more than half of the genes were differentially expressed between normal and defective samples.

Table 3. The number of contigs identified by the *t* test to be distinct between normal and defective samples.

days	negative	positive
1	76	79
2	118	37
3	41	114
4	69	86
5	78	77
6	90	65
7	48	107
8	73	82
9	125	30
10	93	62
11	65	90
12	46	109
13	58	97
15	95	60
16	50	105

To compare the statistical power of TD-based unsupervised FE and conventional methods, a conventional method DESeq2 [13] was applied to either comparison between two groups of RNAi treatments, that is, *gfp*, *DjpiwiA* and others (*Djhp1-1*, *Djima1-1*, *DjpiwiB*, *DjpiwiC*) or that between two time periods, that is, $t = 1, 2, \cdots, 8$, and others ($t = 9, 10, 11, 12, 13, 15, 16$). Although we selected genes associated with adjusted *P*-values less than 0.01, which is the same as that used for TD-based unsupervised FE, DESeq2 identified as few as 10 and 5 contigs for those expressed differently between normal and defective samples, respectively.

Furthermore, in order to evaluate 155 contigs from the viewpoints of biological functions, we performed a Basic Local Alignment Search Tool (BLAST) match between 155 contigs and a well-annotated mouse protein database [27], followed by GO terms enrichment analysis. As a result, 14 GO terms of "Biological process", 10 GO terms of "Molecular function", and 4 GO terms of "Cellular component" were significantly enriched (FDR=0.05)

(Fig. 5). Consistent with the enrichment of GO terms "translation" and "extracellular spac", translation and extracellular matrix are thought to be important in planarian regeneration [36,37]. Consist with the enrichment of GO term "serine-type endopeptidase inhibitor activity", it was reported that transcripts coding for serine-type endopeptidase inhibitors are differentially expressed during planarian regeneration [38]. Thus, we concluded that our analysis successfully selected contigs related to regenerative ability in the planarian.

Figure 5. The results of GO term enrichment analysis of planarian transcripts selected by our method based on blast results against well annotated mouse proteins

4. Discussion

We employed unpopular TD-based unsupervised FE to identify DET despite the existence of many other conventional methods. This was to show that the problem is too difficult to resolve using other conventional methods. Therefore, we applied DESeq2 to the present dataset. As a result, we identified as few as 10 and 5 contigs for those expressed differently between normal and defective samples, respectively, and those expressed dependent upon time, and those expressed dependent upon time; these numbers are very low when compared with the number of contigs identified by TD-based unsupervised FE, 155. DESeq2 failed because there were as many as 278167 contigs, whereas the number of samples was as low as $6 \times 15 \times 6 = 540$; its ratio was as large as 5×10^2. It is a difficult problem to tackle using standard conventional methods designed for a much lower ratio of the number of features to the number of samples.

In order to further validate identified genes biologically, we also performed a BLAST
search against nt database of 155 contigs (see Additional file 1). We found that almost
half of the 155 contigs had a common match with two known long planarian transcripts;
AK388828.1 and AK389113.1, and that these are likely to be alternatively spliced transcripts
of the transcript. TRINITY_DN1947 (Fig. 6). Although we are unsure why Trinity [39] failed
to merge these redundant contigs into one, this suggests that TD-based unsupervised FE
can detect transcripts that share the same expression patterns. If individual contigs are alter-
native spliced short transcripts of a long transcript, which is likely to be TRINITY_DN1947,
there are similar expression patterns.

Figure 6. Multiple alignment of contigs identified by using BLAST search to be aligned to known
planarian transcripts, AK388828.1 or AK389113.1.

One possible objrction to the present study is that the coincidence of analyses between
with and without genome and gene models. If the analyses based upon contig differ from
those with genome, the present study is useless. To examine this possibility, we mapped
short read to genome [24,25] and repeated the above analyses. At first, we evaluated $u_{\ell t}$
and $u_{\ell_1 j}$ as well (Figs. 7 and 8) and we found that u_{2t} and $u_{\ell_1 j}, 2 \le \ell_1 \le 4$) coincide with
the desired property; in contrast to Fig. 2 where three genes are distinct from others in u_{2j},
three genes are distinct from others in $2 \le \ell_1 \le 4$ one by one and similar to Fig. 3, u_{2t} is
dependent on t. Furthermore, u_{1k} is independent of k, replicates, as in Fig. 4 (not show
here), we decided to investigate $|G(\ell_1, 2, 1, \ell_4)|, 2 \le \ell_1 \le 4$ to see which ℓ_4 is associated
with desired properties (Fig. 9) and employed $\ell_4 = 3$. P-values are attributed to *is* in x_{ijtk}
(i.e., genes) with eq. (2) and are corrected by BH method. We identified 89 genes associated
with the adjusted P-values less than 0.01. Since the number does not differ from that of
contigs, 155, so much, analyses with and without genome seem to coincide with each other.

Figure 7. $u_{\ell_1 j}$ $(1 \leq \ell_1 \leq 4)$ that represent 1st to 4th singular value vectors attributed to RNAi experiments that coincides with the distinction between normal and defective samples.

Figure 8. u_{2t} that represent 2nd singular value vectors attributed to days after the treatments. The time point where sign of u_{2t} changes coincides with time point when regenerative defects of *DjpiwiB* RNAi appears.

G(,2,2,1)

G(,3,2,1)

G(,4,2,1)

Figure 9. $|G(\ell_1, 2, 1, \ell_4)|$. Horizontal axes are ℓ_4.

To further confirm the coincidence of analyses between with and without genome, we draw Venn diagram (Fig. 10). Considering the large numbers of contigs ($N = 278,167$) or genes ($N = 24,090$), this numbed of overlaps is clearly significant. Thus, we can conclude the analysis with contigs coincide with that with genome well. We also compute AUC to predict the selected 155 contigs with P-values attributed to genes as follows. At first, all contigs are divided to two classes, either selected 155 contigs or others. Then, P-values attributed to genes are re-attributed to the corresponding contigs. These P-values re-attributed from genes to contigs are used to discriminate contigs between selected 155 contigs and others.

Figure 10. Venn diagram between the selected contigs and genes.

and found that AUC is 0.95. Thus not only selected genes/contigs, but also P-values themselves coincide with each other between with and without genome.

Finally, we have performed enriched analyses toward the selected 89 genes (Fig. 11). Although the number of GO terms increased from Fig. 5, there are some overlaps between Figs. 5 and 11. Thus, TD based unsupervised FE has ability to select biologically reasonable genes even without genome.

Figure 11. The results of GO term enrichment analysis of planarian genes selected by our method based on blast results against well annotated mouse proteins

Yet another objection might be small number of genes/contigs selected. Nevertheless, we already proposed the solution to improve the selected ones [40]. Since in the case when we need more genes/contigs selected, we can employ the strategy, the small number of genes/contigs selected is not a critical problem.

5. Conclusions

First, TD-based unsupervised FE could identify more contigs that are expressed distinctly between normal and defective samples, as well as being expressed in a time-dependent manner (days after RNAi) then DESeq2. GO term enrichment analysis also showed these contigs might be related to planarian regeneration.

In addition, half of the identified 155 transcripts were likely alternative spliced transcripts of a longer transcript. This supports the ability of TD-based unsupervised FE to select contigs that share similar expression patterns, since alternative spliced transcripts from longer transcripts are likely to share the same expression pattern. As a result, TD-based unsupervised FE is expected to be an effective tool to be applied to DET detection using the redundant contigs obtained by applying the *de novo* assembly of short reads from RNA-seq applied to non-model organisms that lack a well-annotated genome.

Furthermore, the results obtained with contigs coincide with those with genome and gene model; it is expected that TD based unsupervsied FE can be a promising tool to deal with contigs of organism without genome.

Supplementary Materials: The following supporting information can be downloaded at: BLAST search results toward 155 contigs (excel) and contigs to genes correspnding table (csv)

Author Contributions: M.K. and N.K. conceived and conducted the experiments, and all authors analysed the results and reviewed the manuscript.

Funding: This work was supported by KAKENHI [grant numbers 19H05270, 20H04848, and 20K12067] to Y.T. and KAKENHI [grant numbers 19K16149, 22K15127] to M.K.

Data Availability Statement: All quantification results and sample information of the RNA-seq analysis were deposited as GSE174855 in the GEO repository.

References

1. Taguchi, Y.H. Comparative Transcriptomics Analysis. In *Encyclopedia of Bioinformatics and Computational Biology*; Ranganathan, S.; Gribskov, M.; Nakai, K.; Schönbach, C., Eds.; Academic Press: Oxford, 2019; pp. 814–818. https://doi.org/https://doi.org/10.1016/B978-0-12-809633-8.20163-5.
2. Trapnell, C.; Roberts, A.; Goff, L.; Pertea, G.; Kim, D.; Kelley, D.R.; Pimentel, H.; Salzberg, S.L.; Rinn, J.L.; Pachter, L. Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nature Protocols* **2012**, *7*, 562–578. https://doi.org/10.1038/nprot.2012.016.
3. Young, M.D.; Wakefield, M.J.; Smyth, G.K.; Oshlack, A. Gene ontology analysis for RNA-seq: accounting for selection bias. *Genome Biology* **2010**, *11*, R14. https://doi.org/10.1186/gb-2010-11-2-r14.
4. Liao, Y.; Smyth, G.K.; Shi, W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* **2013**, *30*, 923–930, [https://academic.oup.com/bioinformatics/article-pdf/30/7/923/48921621/bioinformatics_30_7_9]. https://doi.org/10.1093/bioinformatics/btt656.
5. Pertea, G.; Pertea, M. GFF Utilities: GffRead and GffCompare [version 1; peer review: 3 approved]. *F1000Research* **2020**, *9*. https://doi.org/10.12688/f1000research.23297.1.
6. Patro, R.; Duggal, G.; Love, M.I.; Irizarry, R.A.; Kingsford, C. Salmon provides fast and bias-aware quantification of transcript expression. *Nature Methods* **2017**, *14*, 417–419, [1505.02710]. https://doi.org/10.1038/nmeth.4197.
7. Bray, N.L.; Pimentel, H.; Melsted, P.; Pachter, L. Near-optimal probabilistic RNA-seq quantification. *Nature Biotechnology* **2016**, *34*, 525–527. https://doi.org/10.1038/nbt.3519.
8. Li, B.; Dewey, C.N. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* **2011**, *12*, 323. https://doi.org/10.1186/1471-2105-12-323.
9. Sarantopoulou, D.; Brooks, T.G.; Nayak, S.; Mrčela, A.; Lahens, N.F.; Grant, G.R. Comparative evaluation of full-length isoform quantification from RNA-Seq. *BMC Bioinformatics* **2021**, *22*, 266. https://doi.org/10.1186/s12859-021-04198-1.
10. Raghavan, V.; Kraft, L.; Mesny, F.; Rigerte, L. A simple guide to de novo transcriptome assembly and annotation. *Briefings in Bioinformatics* **2022**, *23*, bbab563, [https://academic.oup.com/bib/article-pdf/23/2/bbab563/42805091/bbab563.pdf]. https://doi.org/10.1093/bib/bbab563.
11. Ungaro, A.; Pech, N.; Martin, J.F.; McCairns, R.J.S.; Mévy, J.P.; Chappaz, R.; Gilles, A. Challenges and advances for transcriptome assembly in non-model species. *PLOS ONE* **2017**, *12*, 1–21. https://doi.org/10.1371/journal.pone.0185020.
12. Hsieh, P.H.; Oyang, Y.J.; Chen, C.Y. Effect of de novo transcriptome assembly on transcript quantification. *Scientific Reports* **2019**, *9*, 8304. https://doi.org/10.1038/s41598-019-44499-3.
13. Love, M.I.; Huber, W.; Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology* **2014**, *15*. https://doi.org/10.1186/s13059-014-0550-8.
14. Taguchi, Y.H. *Unsupervised Feature Extraction Applied to Bioinformatics*; Springer International Publishing, 2020. https://doi.org/10.1007/978-3-030-22456-1.
15. Shibata, N.; Kashima, M.; Ishiko, T.; Nishimura, O.; Rouhana, L.; Misaki, K.; Yonemura, S.; Saito, K.; Siomi, H.; Siomi, M.C.; et al. Inheritance of a Nuclear PIWI from Pluripotent Stem Cells by Somatic Descendants Ensures Differentiation by Silencing Transposons in Planarian. *Developmental Cell* **2016**, *37*, 226–237. https://doi.org/10.1016/j.devcel.2016.04.009.
16. Tian, Q.; Guo, Q.; Guo, Y.; Luo, L.; Kristiansen, K.; Han, Z.; Fang, H.; Zhang, S. Whole-genome sequence of the planarian Dugesia japonica combining Illumina and PacBio data. *Genomics* **2022**, *114*, 110293. https://doi.org/https://doi.org/10.1016/j.ygeno.2022.110293.
17. Shibata, N.; Hayashi, T.; Fukumura, R.; Fujii, J.; Kudome-Takamatsu, T.; Nishimura, O.; Sano, S.; Son, F.; Suzuki, N.; Araki, R.; et al. Comprehensive gene expression analyses in pluripotent stem cells of a planarian, Dugesia japonica. *International Journal of Developmental Biology* **2012**, *56*, 93–102. https://doi.org/10.1387/ijdb.113434ns.
18. Rouhana, L.; Weiss, J.A.; Forsthoefel, D.J.; Lee, H.; King, R.S.; Inoue, T.; Shibata, N.; Agata, K.; Newmark, P.A. RNA interference by feeding in vitro-synthesized double-stranded RNA to planarians: Methodology and dynamics. *Developmental Dynamics* **2013**, *242*, 718–730. https://doi.org/10.1002/dvdy.23950.
19. Ujibe, K.; Nishimura, K.; Kashima, M.; Hirata, H. Direct-TRI: High-throughput RNA-extracting Method for all Stages of Zebrafish Development. *BIO-PROTOCOL* **2021**, *11*. https://doi.org/10.21769/bioprotoc.4136.

20. Kamitani, M.; Kashima, M.; Tezuka, A.; Nagano, A.J. Lasy-Seq: a high-throughput library preparation method for RNA-Seq and its application in the analysis of plant responses to fluctuating temperatures. *Scientific Reports* **2019**, *9*, 7091. https://doi.org/10.1038/s41598-019-43600-0.

21. Kashima, M.; Kamitani, M.; Nomura, Y.; Hirata, H.; Nagano, A.J. DeLTa-Seq: direct-lysate targeted RNA-Seq from crude tissue lysate. (8/15 words) Running Title: Development of direct-lysate targeted RNA-Seq method Corresponding Author. *bioRxiv* **2020**, p. 2020.09.15.299180. https://doi.org/10.1101/2020.09.15.299180.

22. Chen, S.; Zhou, Y.; Chen, Y.; Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **2018**, *34*, i884–i890. https://doi.org/10.1093/bioinformatics/bty560.

23. Li, H.; Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **2009**, *25*, 1754–1760, [1303.3997]. https://doi.org/10.1093/bioinformatics/btp324.

24. An, Y.; Kawaguchi, A.; Zhao, C.; Toyoda, A.; Sharifi-Zarchi, A.; Mousavi, S.A.; Bagherzadeh, R.; Inoue, T.; Ogino, H.; Fujiyama, A.; et al. Draft genome of Dugesia japonica provides insights into conserved regulatory elements of the brain restriction gene nou-darake in planarians. *Zoological Letters* **2018**, *4*, 24. https://doi.org/10.1186/s40851-018-0102-2.

25. García-Castro, H.; Kenny, N.J.; Iglesias, M.; Álvarez-Campos, P.; Mason, V.; Elek, A.; Schönauer, A.; Sleight, V.A.; Neiro, J.; Aboobaker, A.; et al. ACME dissociation: a versatile cell fixation-dissociation method for single-cell transcriptomics. *Genome Biology* **2021**, *22*, 89. https://doi.org/10.1186/s13059-021-02302-5.

26. Camacho, C.; Coulouris, G.; Avagyan, V.; Ma, N.; Papadopoulos, J.; Bealer, K.; Madden, T.L. BLAST+: architecture and applications. *BMC bioinformatics* **2009**, *10*, 1–9.

27. Waterston, R.H.; Pachter, L. Initial sequencing and comparative analysis of the mouse genome. *Nature* **2002**, *420*, 520–562.

28. Durinck, S.; Spellman, P.T.; Birney, E.; Huber, W. Mapping identifiers for the integration of genomic datasets with the R/Bioconductor package biomaRt. *Nature protocols* **2009**, *4*, 1184–1191.

29. Carlson, M.; Falcon, S.; Pages, H.; Li, N. GO. db: A set of annotation maps describing the entire Gene Ontology. *R package version* **2019**, *3*, 10–18129.

30. Carlson, M. org. Mm. eg. db: Genome wide annotation for Mouse. R package version 3.2. 3. *Bioconductor. London, United Kingdom: Genome Biology (BMC)* **2019**.

31. Benjamini, Y.; Hochberg, Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)* **1995**, *57*, 289–300.

32. Yashiro, R.; Murota, Y.; Nishida, K.M.; Yamashiro, H.; Fujii, K.; Ogai, A.; Yamanaka, S.; Negishi, L.; Siomi, H.; Siomi, M.C. Piwi Nuclear Localization and Its Regulatory Mechanism in Drosophila Ovarian Somatic Cells. *Cell Reports* **2018**, *23*, 3647–3657. https://doi.org/10.1016/j.celrep.2018.05.051.

33. Brower-Toland, B.; Findley, S.D.; Jiang, L.; Liu, L.; Yin, H.; Dus, M.; Zhou, P.; Elgin, S.C.; Lin, H. Drosophila PIWI associates with chromatin and interacts directly with HP1a. *Genes & Development* **2007**, *21*, 2300–2311. https://doi.org/10.1101/gad.1564307.

34. Hubert, A.; Henderson, J.M.; Cowles, M.W.; Ross, K.G.; Hagen, M.; Anderson, C.; Szeterlak, C.J.; Zayas, R.M. A functional genomics screen identifies an Importin-α homolog as a regulator of stem cell function and tissue patterning during planarian regeneration. *BMC Genomics* **2015**, *16*, 1–18. https://doi.org/10.1186/s12864-015-1979-1.

35. Zeng, A.; Li, Y.Q.; Wang, C.; Han, X.S.; Li, G.; Wang, J.Y.; Li, D.S.; Qin, Y.W.; Shi, Y.; Brewer, G.; et al. Heterochromatin protein 1 promotes self-renewal and triggers regenerative proliferation in adult stem cells. *Journal of Cell Biology* **2013**, *201*, 409–425. https://doi.org/10.1083/jcb.201207172.

36. Yang, Y.; Wang, P.; Jin, B.; Dong, Z.; Chen, G.; Liu, D. Screening of Potential Key Transcripts Involved in Planarian Regeneration and Analysis of Its Regeneration Patterns by PacBio Long-Read Sequencing. *Frontiers in genetics* **2020**, *11*, 580.

37. Sonpho, E.; Wootthichairangsan, C.; Ishida, M.; Inoue, T.; Agata, K.; Maleehuan, A.; Charngkaew, K.; Chomanee, N.; Moonsom, S.; Wongtrakoongate, P.; et al. ECM-body: A cell-free 3D biomimetic scaffold derived from intact planarian body. *Zoological science* **2020**, *37*, 307–313.

38. Sandmann, T.; Vogg, M.C.; Owlarn, S.; Boutros, M.; Bartscherer, K. The head-regeneration transcriptome of the planarian Schmidtea mediterranea. *Genome biology* **2011**, *12*, 1–19.

39. Haas, B.J.; Papanicolaou, A.; Yassour, M.; Grabherr, M.; Blood, P.D.; Bowden, J.; Couger, M.B.; Eccles, D.; Li, B.; Lieber, M.; et al. De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nature Protocols* **2013**, *8*, 1494–1512. https://doi.org/10.1038/nprot.2013.084.

40. Taguchi, Y.h.; Turki, T. Adapted tensor decomposition and PCA based unsupervised feature extraction select more biologically reasonable differentially expressed genes than conventional methods. *Scientific Reports* **2022**, *12*, 17438. https://doi.org/10.1038/s41598-022-21474-z.