Title: GIP: An open-source computational pipeline for mapping genomic

instability from protists to cancer cells

Authors: Gerald F. Späth[1] and Giovanni Bussotti[1,2,*]

Affiliations:

[1]Institut Pasteur, INSERM U1201, Unité de Parasitologie moléculaire et Signalisation, Paris,

France.

[2]Hub de Bioinformatique et Biostatistique - Département Biologie Computationnelle, Institut

Pasteur, 75015 Paris, France

*corresponding author: Giovanni Bussotti, giovanni.bussotti@pasteur.fr

Abstract:

One of the hallmarks of eukaryotic pathogens is the ability to genetically adapt to environmental

change, causing for instance frequent drug resistance. Even though genome instability has been

recognized as a key driver for microbial adaptation, most available computational tools focus

just on one mutation type or analytical step. To overcome this limitation and better understand

the role of genetic changes in enhancing microbial pathogenicity we established GIP, a novel,

powerful bioinformatic pipeline for comparative genome analysis across microbial populations.

GIP allows batch processing of whole genome sequencing datasets, including read alignment,

normalization, quantification of chromosomes, genes and genomic bins, and detection of single

nucleotide and structural variants. GIP produces a comprehensive summary report providing

sample statistics together with graphical representations of genomic features and tabulated

results. GIP further includes a tool-suite that enables downstream custom comparisons of

samples subsets. GIP is broadly applicable to different eukaryotic systems that exploit genome

instability for adaptation, including *Leishmania*, *Plasmodium, Candida*, and cancer.

25
26
27

One Sentence Summary:

GIP - a novel pipeline for detecting, comparing and visualizing genome instability.

Keywords:

Genome instability, aneuploidy, copy number variations, single nucleotide variation, structural

variation, genetic adaptation.

Main Text:

In recent years, the field of genomics has rapidly expanded with a fast increase in the number

of newly sequenced genomes (1). This surge is a direct consequence of the development of new

and ever more efficient, high-throughput capable sequencing technologies (2). On the one hand,

the improvement in long reads technology allowed the generation of high-quality genome

assemblies (3,4). On the other hand, the decreasing costs for short-reads sequencing and the

parallel increase in sequencing throughput propelled the exponential increase of available

whole genome sequencing (WGS) data (5). Thanks to these advances one can reasonably expect

WGS to rapidly become a key component of personalized medicine and clinical applications.

In this context several consortia-based projects have been established with the goal to produce

WGS for the study of different biological systems (5), and an number of publicly available

databases have been compiled or updated (6). Parallel to data availability, many bioinformatics

tools have been developed to perform specific genome analysis tasks (7,8). For instance, tools

such as Freebayes (9), CNVnator (10) and DELLY (11) have been respectively used for the

detection or characterization of DNA single nucleotide variants (SNVs), copy number

variations (CNVs), and structural variations (SVs), but their scope is limited to the analysis of

one genomic feature at the time. A number of integrative WGS pipelines and workflows have

been established combining the execution of multiple bioinformatics algorithms serving

different analysis steps (12). Even though continuous progress has been made, there is no

standardized or unified approach for genomic investigation. For the development of improved

WGS data analysis pipelines, several important requirements need be considered, including

portability, reproducibility, scalability and compatibility with high-performance computing

(HPC) clusters and remote cloud computing. Here we introduce a novel genome instability

pipeline (GIP) that fulfills all these requirements. GIP facilitates the genome-wide detection,

quantification, comparison and visualization of chromosome aneuploidies, gene CNVs, SNVs

and SVs. GIP is implemented in Nextflow (13), a workflow language that allows to execute

GIP seamlessly in local workstation, on an HPC or remotely in the cloud. All required

environment and software dependencies of GIP are fulfilled and provided with a Singularity

container, thus making GIP reproducible, easy-to-install and easy-to-use. GIP allows the use of

giptools, a tool-suit of R-based modules for genome data exploration, enabling the comparison

of sample sub-sets. GIP and giptools generate a summary report with publication-quality figures

and spreadsheet tables. GIP and giptools constitute a single framework for WGS analysis

suitable both for large scale batch analysis of individual genomes and comparison of samples

from different experimental conditions or origins. Lastly, a key strength of GIP and giptools is

the general applicability to different eukaryotic species. We already successfully applied GIP

on the analysis of *Leishmania* genomes (14-16). In this study, we validate the use of GIP and

giptools using WGS data from published datasets of the three major human pathogens

Leishmania infantum, *Plasmodium vivax* and *Candida albicans* and as well as three human

cancer cell lines.

Results:

The GIP workflow. GIP is a tool for scientific investigation compatible with Linux systems,

requiring minimal configuration and distributed as a self-contained package. GIP consists of 3

files: the Nextflow pipeline code, the configuration and the Singularity container files. The

minimum required input is a paired-end WGS data set and a reference genome assembly in the

standard fastq and FASTA format respectively. GIP analyses include (i) extracting genomic

features such as assembly gaps or repetitive elements, (ii) mapping the reads, (iii) evaluating

chromosome, gene and genomic bin copy numbers, (iv) identifying and visualizing copy

number variation with respect to the reference genome, (v) identifying and quantifying gene

clusters, (vi) detecting and annotating SNVs, (vii) measuring non-synonimous (N) and

synonymous (S) mutations for all genes, (viii) detecting SVs including tandem duplications,

deletions, inversions and break-ends translocations using split-read and read-pair orientation

information, (ix) producing a report file providing summary statistics, tables and visualizations

(**Fig. 1**). GIP allows to customize filtering and visualization options via the configuration file

(see methods). The output of GIP can be used as input for giptools, a tool-suite to compare

sample sub-sets and highlight chromosome copy number, gene copy number and SNV

differences.

Applying giptools on a *Leishmania infantum* case study. GIP permits the batch analysis of a

set of individual samples, where each sample is considered separately and compared only with

respect to the provided reference genome assembly. As a consequence, all variants and copy

number alterations detected in a sample merely reflect the differences between the sequenced

and the reference genomes. While this application may be sufficient in some circumstances,

research projects often involve downstream comparison between samples. Examples include

the comparison of gene or chromosome copy variation number between for example drug

resistant and drug susceptible samples, or the juxtaposition of SNVs detected in isolates from

different geographic areas. For this purpose, we developed giptools, a suit of thirteen modules

that allows to compare samples processed by GIP (**Table 1**). All modules in giptools are fully

embedded in the Singularity container and they are provided with their own documentation.

Table 1: giptools modules

Module name	Purpose
karyotype	Compare the chromosome sequencing coverage distributions
binCNV	Compare bin sequencing coverage in 2 samples
geCNV	Compare gene sequencing coverage in 2 samples
ternary	Compare gene sequencing coverage in 3 samples
ternaryBin	Compare bin sequencing coverage in 3 samples
SNV	Compare SNVs in multiple samples
binDensity	Density plot of bin sequencing coverage of many samples
geInteraction	Detect CNV genes in many samples and produce correlation-based networks
genomeDistance	Compare samples genomic distance
phylogeny	Extract the SNVs union and infer the phylogenetic tree
convergentCNV	Detect convergent CNV gene amplifications
overview	Overview of the sequencing coverage of chromosomes, genomic bins and genes
panel	Extract genomic information of a gene panel

To illustrate the type of exploratory data analyses and the biological questions that can be

addressed, we tested giptools on a previously analyzed dataset of seven clinical *Leishmania*

infantum isolates from Tunisia (14). *Leishmania* is the etiological agent of leishmaniasis, a life-

threatening human and veterinary disease affecting 12 million people worldwide (17). The

parasites were derived from patients affected with visceral leishmaniasis, expanded in cell

culture and their genomes were sequenced. The dataset includes four Glucantime drug

susceptible isolates and three isolates from relapsed patients, and thus comparison may inform

on genetic factors resulting in treatment failure. giptools allowed the detection and visualization

of pervasive intra-chromosomal CNVs across the thirty-six *Leishmania* chromosomes (**Fig.**

2A). Additionally, giptools enables targeted comparison of normalized genomic bin sequencing

coverage of sample pairs. We used giptools' `binCNV` module to compute the ratio between

corresponding genomic bins of the strains LIPA83 over ZK43, which correspond to a first-

episode and a relapse leishmaniasis isolate, respectively. giptools represents different

chromosomes as separate panels (**Fig. 2B**), as part of single genome-wide overview (**Fig. S1A**)

or as distinct plots (**Fig. S1B**). This analysis allowed the identification of 2 905 and 2 208 bins

that were respectively amplified or depleted in LIPA83 with respect to ZK43. The results are

returned by giptools as a Microsoft Excel table (.xlsx format) providing ratio scores at each

genomic position (**Table S1**). Likewise, giptools permits three-way comparisons of normalized

genomic bin sequencing coverage with ternary plots (**Fig. 2C**). We used this representation to

display the genomic bin relative abundance in samples ZK43, LIPA83 and ZK28. The analysis

shows important strain-specific differences in bin copy number that are visualized by shifts of

the signals out of the center. Similar to genomic bin analysis, giptools makes it possible to

compare the sequencing depth of annotated genes and thus the copy number in two or three

samples (**Fig. 2D** and **Fig. S2**). However, the determination of gene copy number might be

impeded by (i) short read length or fragment insert size, (ii) the complexity of the target genome,

or (iii) the presence of repetitive elements. The read map quality (MAPQ) score is a measure

that reflects how much each gene is supported by unambiguously mapped reads (high MAPQ)

in contrast to multimapping reads (low MAPQ). Together with the coverage, GIP also computes

the mean read MAPQ score for each gene and allows a different strategy to determine the copy

number of low MAPQ genes (see methods). The evaluation of LIPA83 and ZK43 gene

coverage ratio scores revealed 13 gene CNVs with a stringent MAPQ cutoff of 50 (**Table S2**).

The maximum normalized coverage value of 6.5 was observed for a putative amastin surface

glycoprotein (LINF_310009800). Other examples of gene CNVs in this set include the putative

surface antigen protein 2 (LINF_120013500) and the heat shock protein HSP33

(LINF_300021600) (**Table S2**). Genes falling below a user defined MAPQ score and sharing

high level of sequence similarity are assigned to the same gene cluster, and their measured

coverage scores are averaged across all members of the group. Low MAPQ scores can also be

associated with single genes, e.g. in the case of internal repetitive elements that cause multiple

ambiguous alignments inside the gene itself, or if mapping occurs in possibly misannotated

intergenic regions. The LIPA83/ZK43 comparison showed 27 CNV gene clusters, including

cluster clstr303 (3 genes with 'amastin-like' annotation) and cluster clstr16 (2 tb-292

membrane-associated protein-like proteins) (**Table S2**). These results demonstrate the power

of GIP and giptools to detect and compare intra-chromosomal CNVs in *Leishmania* at genomic

bin level. Conveniently, analogous two- or three-ways comparisons can be applied to reveal

copy number variations at individual gene or gene cluster levels.

Comparative genomics of a *Plasmodium vivax* WGS dataset. We next applied GIP and

giptools on other biological systems to demonstrate its broad applicability outside the

Leishmania field, including the human apicomplexan parasite *Plasmodium vivax*. *Plasmodium*

vivax is an obligate protist parasite and a human pathogen causing malarial illness. *Plasmodium*

vivax gives rise every year to 130 million clinical cases (18), and it is estimated that 2.5 billion

people are at risk of infection worldwide (19-21). We applied GIP and giptools to investigate

genomic variations across a sizeable dataset of 222 *Plasmodium vivax* genomes isolated from

clinical samples of 14 countries worldwide (22,23) (**Table S3**). The phylogenetic tree

reconstruction and PCA analyses (**Fig. 3A** and **B**) showed a high correlation between genotypes

and the geographic origin of the samples. However, we detected substantial genomic variability

between isolates collected at smaller geographical scale, with 14 555 SNVs (~42% of the total)

uniquely characterizing representative samples from five Ethiopian study sites (23) (**Fig. 3C**

and **D**). This result may reflect diverging evolutionary trajectories radiating from few founder

strains. At gene level we profiled the copy number variations of two gene panels. The first panel

accounts for 43 previously described genes encoding for potential erythrocyte binding proteins

suggested to operate at the interface of the parasite-host invasion process (23). The second panel

includes 2 drug resistance markers comprising the chloroquine resistance transporter

PVP01_0109300 and the multidrug resistance protein 1 PVP01_1010900, and 4 proteins

implicated in red blood cell invasion, such as the merozoite surface protein gene MSP7

PVP01_1219700, the reticulocyte binding protein gene 2c PVP01_0534300, the serine-repeat

antigen 3 PVP01_0417000, and the reticulocyte binding protein 2b PVP01_0800700) (24-33).

Read depth analysis indicated that 4 genes in the panel (PVP01_0623800, PVP01_1031400,

PVP01_1031200, PVP01_1031300) show a high degree of variability, with amplifications

observed in samples from distinct geographic areas thus suggesting a convergent evolutionary

process (**Fig. 3E**). Furthermore, 6 genes positioned on chromosome 14 are deleted in the Thai

strain PD0689_C as a result of the entire chromosome loss (nullisomy) (**Fig. 3E** and **F**). Finally,

the comparison of synonymous and non-synonymous SNVs in the panel genes revealed

important differences between sample groups. Our analysis indicates an overall higher number

of non-synonymous mutations in Ethiopian rather than Cambodian isolates, therefore

suggesting a stronger evolutionary pressure in the African strains (**Fig. 3G**). Taken together

these analyses well illustrate how GIP and giptools can be readily applied for bulk analysis of

Plasmodium vivax genomes to assess genome diversity, extract evolutionary information and

profile potential disease biomarkers.

Gene CNV analysis of *Candida albicans* evolutionary adapted strains. We next applied GIP

and giptools to the human fungal pathogen *Candida albicans*, an opportunistic yeast exhibiting

major genome plasticity (34-43) and causing hundreds of thousands of severe infections each

year (44). Candidemia, a bloodstream infection with *Candida*, are often associated with high

rates of morbidity and mortality, (15–50%) notwithstanding existing antifungal treatments

(45,46). We applied GIP and giptools to a *Candida albicans* WGS dataset described in a recent

study that covers five different progenitor strains (P75063, P75016, P78042, SC5314,

AMS3050) and investigates CNVs driving tolerance and resistance to anti-fungal azole drugs

(47). We analyzed nineteen samples, including (i) four clinical isolates (P75063, P75016,

P78042, SC5314), (ii) seven strains selected *in vitro* against the anti-fungal drug fluconazole

(FLC) (AMS4104, AMS4105, AMS4106, AMS4107, AMS4397, AMS4444, AMS4702), (iii)

four isogenic colonies adapted to the drug miconazole (AMS3051, AMS3052, AMS3053 and

AMS3054) together with their progenitor (AMS3050), and (iv) three colonies derived from a

miconazole-adapted population and isolated on a rich medium (AMS3092, AMS3093 and

AMS3094) (47-49). We confirmed gene amplification for the drug efflux pumps TAC1

(orf19.3188) and ERG11 (orf19.922), for the stress response proteins HSP70 (orf19.4980),

CGR1 (orf19.2722), ERO1 (orf19.4871), TPK1 (orf19.4892), ASR1 (orf19.2344), PBS2

(orf19.7388) and CRZ1 (orf19.7359), and for proteins involved in membrane and cell wall

integrity, including CDR3 (orf19.1313), NCP1 (orf19.2672), ECM21 (orf19.4887), MNN23

(orf19.4874), RHB1 (orf19.5994) and KRE6 (orf19.7363) (**Table S4**). Further, the cross

comparison of gene read depths across the nineteen genomes uncovered a novel set of 1 505

gene CNV showing correlating and anti-correlating gene dosage variations (**Fig. 4A** and **B**,

Table S4) that could be assigned to nine distinct correlation clusters (CC) (**Fig. S3A**, **Table**

S5). We verified the sequencing coverage of genomic regions encompassing gene CNVs,

including three regions amplified in fluconazole resistant strains (**Fig. 4B**, **Fig. S3B**) (47) and

a region whose amplification correlates with the level of miconazole resistance (47) (**Fig. S3C**),

as well as the loss of heterozygosity associated to the depletion of chromosome 3 left arm in

sample AMS3051 (**Fig. S3D**) (47). Eventually, by representing genes and absolute correlation

respectively as nodes and edges of a network, we identified 9 highly interconnected network

clusters (NC) (**Fig. 4C**, **Table S6**). NC7, NC8 and NC9 embody genes from individual

chromosomes, respectively chromosomes 1, 3 and 4. The most parsimonious explanation for

the high levels of correlation observed in these NCs (**Fig. 4C**) is the occurrence of sub-

chromosomal amplifications affecting several adjacent genes. A different scenario is pictured

for each of the remaining NCs (NC1-6) where the genes are located on different chromosomes

thus suggesting genetic interactions that causes coordinated changes in gene copy number.

Altogether, the reproducibility of previously published results validates GIP and giptools for

the analysis of fungal gene CNVs, and uncovers a role of epistatic CNV interactions similar to

Leishmania (in preparation) in driving genomic adaptation in *Candida albicans* populations

under drug selection.

Exploring instability of larger genomes using cancer cell lines as a benchmark. The larger

genome size, and the higher number of genes and WGS reads and can represent a challenge

when working with higher eukaryotes. For the purpose of comparison, the human genome is

~216 times larger than the one of *Candida albicans* we analyse in this study. Therefore, we

sought to evaluate the applicability of the GIP and giptools framework to human data and

utilized a panel of cancer cell lines as a test set. In our analyses we considered publicly available

WGS data of the cell lines T47D, NCI_H460 and K562 (50), which respectively derive from

human breast, lung and blood cancers. We detected 1 647 016 SNVs (**Supplementary Data 1**)

and allele frequency shifts indicating repeated loss of heterozygosity events (**Fig. 5**). The

coverage analysis indicates pervasive presence of CNVs both at chromosomal and sub-

chromosomal levels (**Fig. S4**) correlating with the SNV frequency patterns (**Fig. S5** and **S6**).

These analyses exemplify the application of GIP and giptools to larger, more complex genomes

involving minor parameter adjustments while relying on the same bioinformatic tool.

Discussion:

Genome instability is a key driver of evolution for microbial pathogens and cancer cells (51) and a major source of human morbidity. Here we introduce GIP and giptools, a unified framework for genotype profiling of biological systems exploiting genome instability for adaptation. Current analysis tools suffer from several limitations. A fist common limitation is that their application is limited to individual types of mutations, but are unable to provide genome-wide comprehensive reports covering multiple forms of genome instability. A second limitation is when analysis tools allow to identify variations with respects to a reference assembly, but do not support comparisons between samples. A third important limitation is the software applicability range, where genome analysis tools are tailored for the analysis of WGS data from a given species (52-56), but are not generally applicable to different biological systems. Such tools cannot be applied to investigate genome variations across multiple species and explore potentially conserved genomic adaptation mechanisms. A possible workaround is to use different tools for different species. However, each tool has its own parametrization and specificity, and as a consequence this strategy has the potential to introduce technical biases making unable to perform meaningful cross-species comparisons. Furthermore, especially when investigating non-model organisms, a tool may not be available for the analysis of your species of interest. A forth critical limitation—with potential serious consequences for clinical applications, is software portability and more in general reproducibility. The same bioinformatic application may be unsuited to run on different computer environments, or worse, if the software requirements are not carefully specified, it can generate irreproducible and potentially flawed results. Lastly, one more common limitation is posed by software scalability. In the WGS domain, with the rapid increase new samples made available and the enormous amount of data generated in each sequencing run, the CPU and memory resources of local workstation risk to quickly become inadequate for data analysis. Therefore, it is paramount that

WGS tools are implemented to run on high-performance computing (HPC) clusters and feature

remote cloud computing solutions.

To support and ease the study of genome instability and address all these common challenges

we developed GIP and giptools. GIP provides karyotype, gene CNV, SNV and SV batch

analyses. giptools enables custom sample comparisons, to explore differences and common

features between genomes. Our pipeline limits as much as possible the use of hardcoded

parametrization which could limit its use to a specific organism. Therefore, GIP's flexible

design makes it adapted for the genome analysis of both model and non-model organisms,

including *Leishmania* or human. In this study we exemplify the ability GIP and giptools in

performing genomic screenings of three major pathogenic eukaryotes and human cancer cell

lines. Thanks to the Singularity implementation all required software are embedded and

provided within the container. As a consequence, a user could easily recreate the same work

environment just by downloading the pipeline container, and reproduce exactly the same

publication-quality plots and tables presented in this study (see methods). All plots and table

presented in this study were directly generated with GIP and giptools with only minor manual

layout changes to fit multiple plots in the same figure and homogenize the font sizes. The only

exception is the plot showing sample PD0689_C chromosome 14 nullisomy (**Fig. 3F**), which

was generated visualizing with IGV (57) the coverage density track produced by GIP (see

methods). Finally, because of its Nextflow implementation GIP can be executed on a local

machine, on cluster resource manager or the cloud. GIP can be executed on individual samples

and with no additional effort on large WGS data sets for batch computation as shown for the

222 *Plasmodium vivax* genomes. While originally deployed for *Leishmania* genome analysis,

in this study we validate the use of our pipeline on other organisms reproducing expected

results. For example, in *Candida albicans* we confirmed the CNVs correlating to drug

resistance as well as a loss of heterozygosity event (**Fig. S3B-C** and **D**). Parallel to this we also

show how GIP and giptools can be used for data mining and scientific discovery. New findings

include (i) the discovery of the convergent amplification of erythrocyte binding proteins in

Plasmodium vivax strains sampled from distinct geographic areas (**Fig. 3E**), (ii) the detection

of a nullisomic strain (**Fig. 3F**), and (iii) the identification of correlated copy number variations

between genes positioned on separate chromosomes of *Candida albicans* adapting strains.

These results well illustrate how GIP and giptools can be applied to perform extended genomic

analyses in different biological systems and drive biomedical discovery. To conclude, we

believe that GIP and giptools represent a step forward toward reproducible research in

genomics, and provide a robust framework study how microbes and tumor cells harness genome

instability for environmental adaptation and fitness gain.

Figures

Fig. 1

GIP workflow. The schema on the left recapitulates the GIP inputs, processes and outputs (see

methods). Blue, orange, green and purple boxes indicate genome reference, read mapping,

quantification and variants computation steps, respectively. The panels on the right demonstrate

example plots included in the GIP report computed for individual samples. The "Karyotype"

plot shows the coverage distributions for each chromosome (y-axis). The "Genomic bins" plot

shows the genomic position (x-axis) and the normalized genomic bin sequencing coverage (y-

axis). The "Gene CNVs" plot shows the normalized gene sequencing coverage. The "Structural

variants" panel shows a Circos plot representing break-end translocations (black links in the

inmost part of the plot), and other possible structural variations in the outer tracks, including

insertions, duplications, deletions and inversions. The outmost track shows the normalized

sequencing coverage. The "Single nucleotide variants" plot shows on the x and y axes

respectively the genomic position and variant allele frequency of detected SNVs.

Fig. 2:

Comparing *Leishmania infantum* genomes with giptools. (**A**) Density plot representing the

genomic coverage of the seven *Leishmania infantum* isolates. The x-axis shows the log 10

normalized coverage of genomic bins. The y-axis reflects the genomic position. The thirty-six

different chromosomes are materialized as separate panels. The blue shading indicates the (2D)

kernel density estimates of genomic bins. The two red vertical lines mark the 1.5 and 0.5

coverage values. A selection of 50 000 bins with coverage > 1.5 or < 0.5 is shown as black dots.

(**B**) Scatterplot of the genomic bin normalized sequencing coverage ratio of samples LIPA83

over ZK43. The x any y axes show the ratio score and the genomic position respectively. Ratio

scores > 1.25 are labelled in orange and indicate genomic bin amplification. Ratio scores < 0.75

are labelled in blue and indicate genomic bin depletion. (**C**) Ternary comparison showing the

relative abundance of the genomic in samples LIPA83, ZK43 and ZK28. The axes report the

fraction of the bins normalized sequencing coverage in the three strains. The blue contour

indicates the log10 bin density. A subset of 5 000 bins is shown as black dots. Each given point

in the plot is adding up to 100. The density area at the center of the plot indicates bins with

equal copy number and thus a ~33 distribution across the three axes. (**D**) Scatterplot showing

the log10 normalized sequencing coverage of annotated genes in ZK43 (x-axis) and LIPA83

(y-axis). The red line indicates the bisector. Dots represent individual genes. (**E**) Sequencing

coverage ratio of gene clusters in samples LIPA83 and ZK43. Dots represent gene clusters. For

plots **D** and **E,** the ratio scores > 1.5 or < 0.5 are labelled in orange and blue respectively.

Fig. 3:

Plasmodium vivax genomic diversity. (**A**) Predicted maximum likelihood phylogenetic tree

reconstruction. (**B**) PCA analysis of the phylogenetic distances estimated from the tree in (**A**).

Each dot indicates a sample. The colour code reflects the geographic origin of the samples and

matches with the colours of the legend in (**A**). (**C**) Venn diagram comparing the SNVs of five

representative Ethiopian strains. (**D**) Pairwise scatterplot comparing the variant allele frequency

of all detected SNVs in the five Ethiopian strains. (**E**) Gene panel analysis. The x-axis reports

a set of genes of interest. The y-axis indicates the normalized mean gene coverage. The boxplots

demonstrate the coverage values distributions for each gene across all samples. Each dot

represents the coverage of the indicated gene in a given sample. Dot colours reflect the sample

geographic origin as in (**A**). (**F**) Reads per kilo base per million mapped reads (RPKM)

normalized sequencing coverage density track of sample PD0689_C. The boundaries of the 14

chromosomes are shown on the bottom. (**G**) Comparison of non-synonymous (N) and

synonymous (S) mutations between Ethiopia and Cambodia sample groups. Dots represent

genes. The x-axis represents the difference between the mean non-synonymous mutation count

in the two sample groups. The y-axis represents the difference between the mean synonymous

mutation count in the two sample groups. The dot size demonstrates the ratio of the mean

normalized sequencing coverage between the two sample groups for each gene. Red and blue

dot colors indicate genes belonging to the 43 genes panel (23) and the custom 6 genes panel

respectively.

Fig. 4:

Gene CNVs interactions. (**A**) All-vs-all normalized sequencing coverage correlation heatmap.

The heatmap is symmetrical along its diagonal and reports both on the rows and the columns

the detected gene CNVs. The colour scale indicates with green and pink high levels of positive

and negative Pearson correlation, respectively. The side ribbons demonstrate in different

colours the chromosome and the correlation cluster of each gene. (**B**) Gene CNV heatmap. The

columns and the rows report respectively the samples and the detected gene CNVs. The colour

scale indicates the normalized sequencing coverage of the genes. To ease visualization,

coverage values greater than 3 are reported as 3 (red). Black boxes highlight the genomic

regions shown in **Fig. S3B** (panels 1, 2 and 3) and **Fig. S3C**. The ribbons on the left indicate

the chromosome and the correlation cluster of each gene. Top ribbons indicate the genotype

and the strains resulting from the different evolutionary experiments. (**C**) Gene interaction

network. Nodes indicate gene CNVs. Edges reflect the absolute Pearson correlation value. The

closer the nodes are, the higher is the correlation. Only significant interactions (Benjamini-

Hochberg adjusted p-value < 0.01) are shown. The colour of the edges indicates in red and blue

respectively positive and negative correlations. The colour of the nodes denotes the predicted

network cluster for each gene.

Fig. 5:

SNV heterogeneity of cancer cell lines [...] tion. The y-axis indicates the variant allele frequency. T[...] they are coloured according to the sample. K562, green; NCI_H460, brown; T47D, grey. Different panels show different chromosomes.

 Supplementary figures

 Fig. S1:

Bin coverage ratio visualizations. The figure shows alternative representations produced for the

genomic bin normalized sequencing coverage ratio of samples ZK43 and LIPA83. Same layout

as **Fig. 2B**. (**A**) Whole genome overview. (**B**) All the individual chromosomes separately. The

plot shows the example of chromosome 33.

Fig. S2:

Two-ways and three-ways gene coverage comparisons. (**A**) Log10 normalized sequencing

coverage comparison of genes in ZK43 (x-axis) and LIPA83 (y-axis). Same layout a **Fig. 2D**

but representing the different chromosomes in individual panels. (**B**) Ternary plot showing the

relative abundance of genes (green dots) and gene clusters (red dots) in samples LIPA83, ZK43

and ZK28. The axes report the fraction of the genes and genes clusters normalized sequencing

coverage in the three strains. Each given point in the plot adding up to 100. Genes with equal

copy number are shown in the center of the ternary plot, while copy number variations are

visualized by shifts of the dots out of the center. The dots size reflects the fold change variation

between the maximum and the minimum observed normalized coverage value.

Fig. S3:

Copy number variatio­ ... (**A**) Gene CNV

correlation clusters. ... s and the scaled

normalized sequencing coverage of the genes in each cluster. Different panels indicate different

correlation clusters. (**B**) Scatterplots showing the bin normalized sequencing coverage (y-axis)

of three genomic regions (chr4:400 000-1 000 000; chr3:1 250 000-1 500 000; chr1:2 000 000-

2 500 000) in three different samples AMS4702, AMS4104 and AMS4107. The x-axis shows

the genomic position. Dots represent genomic bins. (**C**) Scatterplot showing the bin normalized

sequencing coverage of a genomic region (chr3:500 000-1 000 000) in five different samples

represented as separate panels. From top to bottom the samples are: AMS3050, AMS3053,

AMS3054, AMS3052 and AMS3051. The x and y axes show respectively the genomic position

and the normalized sequencing coverage. Dots represent bins. (**D**) Comparative analysis of

chromosome 3 SNVs. AMS3050 SNVs are shown in green, while AMS3051 SNVs are shown

in grey. The x and the y axes show respectively the genomic position and the variant allele

frequency.

Fig. S4:

Genomic coverage analysis of three cancer c[...]genomic position. The y-axis indicates the normalized [...]es. Dots demonstrate the genomic bins. To ease visualization, coverage values greater than 5 are reported as 5. Different panels show different chromosomes. The three panel rows indicate top to bottom the following samples: K562, NCI_H460 and T47D. (**B**) Chromosome coverage analysis. The x-axis reports the chromosomes. The y-axis reports the estimated somy score. The boxes show the somy score distributions. Green, brown, and grey boxes indicate respectively samples K562, NCI_H460 and T47D.

Fig. S5:

Chromosome 11 combined SNV and bin cove[...]ls allows

the simultaneous displaying of variant allele fr[...]coverage

(y-axis, right). Dots represent SNVs. The lines represent the normalized bin sequencing

coverage. The x-axis indicates the genomic position. Coverage values greater than 5 are shown

as 5. Green, brown, and grey colours indicate respectively samples K562, NCI_H460 and

T47D.

Fig. S6:

SNV frequency density plot. The x-[...]cy. The y-axis the

estimated kernel density between 0 a[...] grey slopes indicate respectively

samples K562, NCI_H460 and T47D.

Methods:

GIP and giptools. All results presented in this study were generated using GIP and giptools

version 1.0.9. GIP code is available upon request. The GIP configuration files (**Supplementary**

Data 2) and the giptools command options used to generate all results (**Supplementary Data**

3) are provided.

Read alignment. WGS reads were downloaded from the Sequence Read Archive (SRA) (58)

and the European Nucleotide Archive (ENA) (59) repositories and the Encyclopedia of DNA

Elements (ENCODE) dashboard (60) (**Table S3**). For *Leishmania infantum* the

GCA_900500625 genome reference and gene annotations available from the ENSEMBL

protists (61) server (release-48) were used. For *Candida albicans* the assembly 21 of the

SC5314 strain genome reference and gene annotations available from the *Candida* Genome

Database (CGD) (62) were used. For *Plasmodium vivax* the P01 reference genome and gene

annotations available from PlasmoDB (63) (release-50) were used. For the cancer cell lines the

human genome GRCh38 primary assembly and gene annotations available from ENSEMBL

(release-102) were used. The repetitive elements of reference genomes were soft-masked by

GIP using Red (64). WGS reads were mapped by GIP using BWA-mem (version 0.7.17)

(65,66) run with option -M to label shorter split hits as secondary. Then the alignment files

were sorted, indexed and reformatted by GIP using Samtools (version 1.8) (67). Finally, read

duplicates were removed by GIP using Picard MarkDuplicates

(http://broadinstitute.github.io/picard) (version 2.18.9) with the option

"VALIDATION_STRINGENCY=LENIENT." In the four considered datasets, WGS reads

were aligned against full assemblies, including unsorted contigs if present. However just the

canonical assembled chromosomes were considered for all downstream analyses (`chrs` option,

Supplementary Data 2). A minimum read alignment MAPQ score was adopted to select genes

for cluster analysis, and to call for SNVs and SVs (`MAPQ` option, **Supplementary Data 2**).

Altogether a total of 6 306 951 266 reads were aligned to the respective reference genomes.

The `giptools overview` module was run to gather the alignment statistics as estimated by

Picard CollectAlignmentSummaryMetrics (**Table S7**).

Genomic bins and genes quantification. GIP was used to evaluate the mean sequencing

coverage and the mean read MAPQ of genomic bins and genes. For genomic bins, GIP

partitioned the input genomes into adjacent windows of user defined lengths (`binSize` option,
Supplementary Data 2). The coverage GC-content score bias was corrected (`CGcorrect`
option, **Supplementary Data 2**) fitting a LOESS regression with a 5-fold cross validation to
optimize the model span parameter. A larger window length was utilized to bin the reference
genomes for Circos plot representations (`binSizeCircos` option, **Supplementary Data 2**). In
Fig. 1 ("Genomic bins" and "Gene CNVs" plots), **Fig. 2**, **Fig. S1**, **Fig. S2** and **Fig. 4** bins and
genes coverage scores were normalized by median chromosome coverage to highlight
amplifications or depletions with respect to the chromosome copy number. In **Fig. 1**
("Structural variants" plot), **Fig. S3B-C**, **Fig. 3E-G**, and **Fig. S4A** bins and genes coverage
scores were normalized by median genome coverage to account for sequencing library size
differences. GIP evaluated statistically significant copy number variant bins and genes (**Fig. 1**
"Genomic bins" and "Gene CNVs" plots) using a p-value threshold of 0.001
(`covPerBinSigOPT` and `covPerGeSigOPT` options, **Supplementary Data 2**). Estimated p-
values for bins and genes CNVs were corrected for multiple testing using the Benjamini –
Yekutieli (`--padjust BY`) and the Benjamini – Hochberg (`--padjust BH`) methods. The somy
scores shown in **Fig. 1** (karyotype plot) and **Fig. S4B** were computed multiplying the median
genome coverage normalized bin coverage by 2. GIP enabled the CNV analysis of genes
sharing high sequence identity by clustering the nucleotide sequences of the genes with low
mean MAPQ score into groups with cd-hit-est (version 4.8.1) (68) with options `-s 0.9 -c 0.9 -
r 0 -d 0 -g 1`. Then for each gene cluster GIP computed the mean gene coverage normalized by
median chromosome coverage (**Fig. 2E**, **Fig S2B**).

Sequencing coverage density estimates. GIP was used to convert the read alignment files
(.bam format) in binary data files reflecting sequencing coverage (.bigWig format). The
coverage file were produced using bamCoverage from the deepTools2 suite (69) (version 3.5.1)
with options "--normalizeUsing RPKM --ignoreDuplicates --binSize 10 --smoothLength 30"

(`bigWigOPT` option, **Supplementary Data 2**). The coverage track of sample PD0689_C was

visualized with IGV using the `Bar Chart`, `Autoscale` and windowing function `Mean`

options.

Single-nucleotide variant analysis. GIP was used to call SNVs using Freebayes (version

1.3.2) (`freebayesOPT` option, **Supplementary Data 2**) and filter its output

(`filterFreebayesOPT` option, **Supplementary Data 2**). Filters included the minimum allele

frequency (`--minFreq`), the minimum number of reads supporting the alternative alleles (`--

minAO`) and minimum mean mapping quality of for the reads supporting the reference (`--

minMQMR`) or the alternative allele (`--minMQM`). A higher number of reads supporting the

variants was requested for predictions positioned inside simple repeats of the same nucleotide

(homopolymers) (`--minAOhomopolymer`). The homopolymers were defined as the DNA

region spanning ±5 bases from the SNV (`--contextSpan 5`), with over 40% of identical

nucleotides (--homopolymerFreq 0.4 `). Further, GIP discarded SNVs with sequencing

coverage above or below 4 median absolute deviations (MADs) from the median chromosome

coverage (`--MADrange`). snpEff (version 4.3t) (70) was used to predict the impact of SNVs

on coding sequence. The predicted effects that GIP considered synonymous mutations are:

"synonymous_variant", "stop_retained_variant" and "start_retained". The predicted effects that

GIP considered non-synonymous mutations are: "missense_variant", "start_lost",

"stop_gained", "stop_lost" and "coding_sequence_variant". The phylogenetic tree was

computed by the giptools module `phylogeny` using IQtree2 (version 2.1.2) (71,72) with

options `--seqtype DNA --alrt 1000 -B 1000`. The Venn-diagram comparison considered the

strains QS0044_C, QS0001_C, QS0037_C, QS0016_C and SGH_358 that were sampled from

different locations in Ethiopia, respectively Habala, Badowacho, Arbaminch, Hawassa and

Jimma. The strains were selected to have comparable average genome coverage (23). To infer

the tree GIP considered the set of filtered SNV and adopted the IUPAC ambiguous notation for

the positions with allele frequency below 70%. The tree was visualized by giptools using the

ggtree R-package (73).

Analysis of structural variants. GIP was used to detect structural variants including insertions,

tandem duplications, deletions, inversions and break-end translocations with DELLY (version

0.8.7) (11). To reduce incorrect predictions the DELLY output was additionally filtered

(`filterDellyOPT` option, **Supplementary Data 2**). GIP discarded poor predictions with

DELLY label "LowQual" (`--rmLowQual`) and low median MAPQ score of mapping reads (`-

-minMAPQ`). SVs positioned in proximity of chromosome ends were removed (--

chrEndFilter) to limit false predictions caused by potential misassembled regions close to the

telomeric ends. To ease visualization and limit the analysis only to best supported SVs GIP

limited the output only to the top predictions (`--topHqPercentIns`, `--topHqPercentDel`, `--

topHqPercentDup` and `--topHqPercentInv`) based on the SV support score as in **Formula 1**,

where *DV*, *DR*, *RV*, and *RR* are respectively the number of high-quality variant pairs, reference

pairs, variant junction reads, and reference junction reads.

535
$$\frac{DV + RV}{DV + RV + DR + RR} * 100$$

Formula 1: SV support score.

The predicted structural variants were represented with Circos (version 0.69-9) (74).

Data and materials availability. All data is available in the main text or the supplementary

materials.

Acknowledgments:

Funding: This study was supported by a seeding grant from the Institut Pasteur International

Department to the LeiSHield Consortium and the EU H2020 project LeiSHield-MATI - REP-

778298-1.

Author contributions: Conceptualization, G.B. and G.F.S.; Methodology, G.B. and G.F.S.;

Software, G.B.; Formal Analysis, G.B., and G.F.S.; Investigation, G.B. and G.F.S.; Writing –

Original Draft, G.B. and G.F.S.; Writing – Review & Editing, G.B. and G.F.S.; Supervision,

G.B. and G.F.S.; Funding Acquisition, G.F.S. and G.B.

Competing interests: Authors declare no competing interests.

Supplementary Tables:

Table S1: Genomic bin coverage ratio of sample LIPA83 over ZK43.

Table S2: Gene coverage ratio of sample LIPA83 over ZK43.

Table S3: Sample information.

Table S4: *Candida albicans* gene CNVs.

Table S5: Gene correlation clusters in *Candida albicans*.

Table S6: Network analysis of gene CNVs in *Candida albicans*.

Table S7: Mapping statistics.

Supplementary Data:

Supplementary Data 1: Cancer cell lines SNVs.

Supplementary Data 2: GIP configuration files.

Supplementary Data 3: giptools commands.

References:

1. Muir, P., Li, S., Lou, S., Wang, D., Spakowicz, D.J., Salichos, L., Zhang, J., Weinstock, G.M., Isaacs, F., Rozowsky, J. *et al.* (2016) The real cost of sequencing: scaling computation to keep pace with data generation. *Genome Biol*, **17**, 53.

2. Pareek, C.S., Smoczynski, R. and Tretyn, A. (2011) Sequencing technologies and genome sequencing. *J Appl Genet*, **52**, 413-435.

3. Chin, C.S., Alexander, D.H., Marks, P., Klammer, A.A., Drake, J., Heiner, C., Clum, A., Copeland, A., Huddleston, J., Eichler, E.E. *et al.* (2013) Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods*, **10**, 563-569.

4. Chin, C.S., Peluso, P., Sedlazeck, F.J., Nattestad, M., Concepcion, G.T., Clum, A., Dunn, C., O'Malley, R., Figueroa-Balderas, R., Morales-Cruz, A. *et al.* (2016) Phased diploid genome assembly with single-molecule real-time sequencing. *Nat Methods*, **13**, 1050-1054.

5. Reuter, J.A., Spacek, D.V. and Snyder, M.P. (2015) High-throughput sequencing technologies. *Mol Cell*, **58**, 586-597.

6. Aurrecoechea, C., Barreto, A., Basenko, E.Y., Brestelli, J., Brunk, B.P., Cade, S., Crouch, K., Doherty, R., Falke, D., Fischer, S. *et al.* (2017) EuPathDB: the eukaryotic pathogen genomics database resource. *Nucleic Acids Res*, **45**, D581-D591.

7. Dolled-Filhart, M.P., Lee, M., Jr., Ou-Yang, C.W., Haraksingh, R.R. and Lin, J.C. (2013) Computational and bioinformatics frameworks for next-generation whole exome and genome sequencing. *ScientificWorldJournal*, **2013**, 730210.

8. Pabinger, S., Dander, A., Fischer, M., Snajder, R., Sperk, M., Efremova, M., Krabichler, B., Speicher, M.R., Zschocke, J. and Trajanoski, Z. (2014) A survey of tools for variant analysis of next-generation genome sequencing data. *Brief Bioinform*, **15**, 256-278.

9. Garrison, E. and Marth, G. (2012), *arXiv preprint arXiv:1207.3907 [q-bio.GN]*.

10. Abyzov, A., Urban, A.E., Snyder, M. and Gerstein, M. (2011) CNVnator: an approach to discover, genotype, and characterize typical and atypical CNVs from family and population genome sequencing. *Genome Res*, **21**, 974-984.

11. Rausch, T., Zichner, T., Schlattl, A., Stütz, A.M., Benes, V. and Korbel, J.O. (2012), *Bioinformatics*, Vol. 28, pp. i333-i339.

12. Hwang, K.B., Lee, I.H., Li, H., Won, D.G., Hernandez-Ferrer, C., Negron, J.A. and Kong, S.W. (2019) Comparative analysis of whole-genome sequencing pipelines to minimize false negative findings. *Sci Rep*, **9**, 3219.

13. Di Tommaso, P., Chatzou, M., Floden, E.W., Barja, P.P., Palumbo, E. and Notredame, C. (2017) Nextflow enables reproducible computational workflows. *Nat Biotechnol*, **35**, 316-319.

14. Bussotti, G., Benkahla, A., Jeddi, F., Souiai, O., Aoun, K., Spath, G.F. and Bouratbine, A. (2020) Nuclear and mitochondrial genome sequencing of North-African Leishmania infantum isolates from cured and relapsed visceral leishmaniasis patients reveals variations correlating with geography and phenotype. *Microb Genom*, **6**.

15. Bussotti, G., Gouzelou, E., Cortes Boite, M., Kherachi, I., Harrat, Z., Eddaikra, N., Mottram, J.C., Antoniou, M., Christodoulou, V., Bali, A. *et al.* (2018) Leishmania Genome Dynamics during Environmental Adaptation Reveal Strain-Specific Differences in Gene Copy Number Variation, Karyotype Instability, and Telomeric Amplification. *MBio*, **9**.

16. Prieto Barja, P., Pescher, P., bussotti, g., Dumetz, F., Imamura, H., Kedra, D., Domagalska, M., Chaumeau, V., Himmelbauer, H., Pages, M. *et al.* (2017), *Nat Ecol Evol*, Vol. 1, pp. 1961-1969.

17. Alvar, J., Velez, I.D., Bern, C., Herrero, M., Desjeux, P., Cano, J., Jannin, J., den Boer,
M. and Team, W.H.O.L.C. (2012) Leishmaniasis worldwide and global estimates of its
incidence. *PLoS One*, **7**, e35671.
18. Organization, W.H. (2018), WHO, Geneva. .
19. Price, R.N., Tjitra, E., Guerra, C.A., Yeung, S., White, N.J. and Anstey, N.M. (2007)
Vivax malaria: neglected and not benign. *Am J Trop Med Hyg*, **77**, 79-87.
20. Gething, P.W., Elyazar, I.R., Moyes, C.L., Smith, D.L., Battle, K.E., Guerra, C.A., Patil,
A.P., Tatem, A.J., Howes, R.E., Myers, M.F. *et al.* (2012) A long neglected world
malaria map: Plasmodium vivax endemicity in 2010. *PLoS Negl Trop Dis*, **6**, e1814.
21. Battle, K.E., Gething, P.W., Elyazar, I.R., Moyes, C.L., Sinka, M.E., Howes, R.E.,
Guerra, C.A., Price, R.N., Baird, K.J. and Hay, S.I. (2012) The global public health
significance of Plasmodium vivax. *Adv Parasitol*, **80**, 1-111.
22. Pearson, R.D., Amato, R., Auburn, S., Miotto, O., Almagro-Garcia, J., Amaratunga, C.,
Suon, S., Mao, S., Noviyanti, R., Trimarsanto, H. *et al.* (2016) Genomic analysis of
local variation and recent evolution in Plasmodium vivax. *Nat Genet*, **48**, 959-964.
23. Ford, A., Kepple, D., Abagero, B.R., Connors, J., Pearson, R., Auburn, S., Getachew,
S., Ford, C., Gunalan, K., Miller, L.H. *et al.* (2020) Whole genome sequencing of
Plasmodium vivax isolates reveals frequent sequence and structural polymorphisms in
erythrocyte binding genes. *PLoS Negl Trop Dis*, **14**, e0008234.
24. Singh, V., Gupta, P. and Pande, V. (2014) Revisiting the multigene families:
Plasmodium var and vir genes. *J Vector Borne Dis*, **51**, 75-81.
25. Rayner, J.C., Tran, T.M., Corredor, V., Huber, C.S., Barnwell, J.W. and Galinski, M.R.
(2005) Dramatic difference in diversity between Plasmodium falciparum and
Plasmodium vivax reticulocyte binding-like genes. *Am J Trop Med Hyg*, **72**, 666-674.
26. Rahul, C.N., Shiva Krishna, K., Pawar, A.P., Bai, M., Kumar, V., Phadke, S. and
Rajesh, V. (2014) Genetic and structural characterization of PvSERA4: potential
implication as therapeutic target for Plasmodium vivax malaria. *J Biomol Struct Dyn*,
32, 580-590.
27. Rahul, C.N., Shiva Krishna, K., Meera, M., Phadke, S. and Rajesh, V. (2015)
Plasmodium vivax: N-terminal diversity in the blood stage SERA genes from Indian
isolates. *Blood Cells Mol Dis*, **55**, 30-35.
28. Luo, Z., Sullivan, S.A. and Carlton, J.M. (2015) The biology of Plasmodium vivax
explored through genomics. *Ann N Y Acad Sci*, **1342**, 53-61.
29. Lin, J.T., Patel, J.C., Kharabora, O., Sattabongkot, J., Muth, S., Ubalee, R., Schuster,
A.L., Rogers, W.O., Wongsrichanalai, C. and Juliano, J.J. (2013) Plasmodium vivax
isolates from Cambodia and Thailand show high genetic complexity and distinct
patterns of P. vivax multidrug resistance gene 1 (pvmdr1) polymorphisms. *Am J Trop
Med Hyg*, **88**, 1116-1123.
30. Gunalan, K., Niangaly, A., Thera, M.A., Doumbo, O.K. and Miller, L.H. (2018)
Plasmodium vivax Infections of Duffy-Negative Erythrocytes: Historically Undetected
or a Recent Adaptation? *Trends Parasitol*, **34**, 420-429.
31. Costa, G.L., Amaral, L.C., Fontes, C.J.F., Carvalho, L.H., de Brito, C.F.A. and de
Sousa, T.N. (2017) Assessment of copy number variation in genes related to drug
resistance in Plasmodium vivax and Plasmodium falciparum isolates from the Brazilian
Amazon and a systematic review of the literature. *Malar J*, **16**, 152.
32. Cornejo, O.E., Fisher, D. and Escalante, A.A. (2014) Genome-wide patterns of genetic
polymorphism and signatures of selection in Plasmodium vivax. *Genome Biol Evol*, **7**,
106-119.
33. Chen, E., Salinas, N.D., Huang, Y., Ntumngia, F., Plasencia, M.D., Gross, M.L.,
Adams, J.H. and Tolia, N.H. (2016) Broadly neutralizing epitopes in the Plasmodium

vivax vaccine candidate Duffy Binding Protein. *Proc Natl Acad Sci U S A*, **113**, 6277-6282.

34. Zolan, M.E. (1995) Chromosome-length polymorphism in fungi. *Microbiol Rev*, **59**, 686-698.

35. Suzuki, T., Nishibayashi, S., Kuroiwa, T., Kanbe, T. and Tanaka, K. (1982) Variance of ploidy in Candida albicans. *J Bacteriol*, **152**, 893-896.

36. Sionov, E., Chang, Y.C. and Kwon-Chung, K.J. (2013) Azole heteroresistance in Cryptococcus neoformans: emergence of resistant clones with chromosomal disomy in the mouse brain during fluconazole treatment. *Antimicrob Agents Chemother*, **57**, 5127-5130.

37. Shin, J.H., Chae, M.J., Song, J.W., Jung, S.I., Cho, D., Kee, S.J., Kim, S.H., Shin, M.G., Suh, S.P. and Ryang, D.W. (2007) Changes in karyotype and azole susceptibility of sequential bloodstream isolates from patients with Candida glabrata candidemia. *J Clin Microbiol*, **45**, 2385-2391.

38. Selmecki, A., Forche, A. and Berman, J. (2010) Genomic plasticity of the human fungal pathogen Candida albicans. *Eukaryot Cell*, **9**, 991-1008.

39. Magee, B.B. and Magee, P.T. (2000) Induction of mating in Candida albicans by construction of MTLa and MTLalpha strains. *Science*, **289**, 310-313.

40. Gerstein, A.C., Fu, M.S., Mukaremera, L., Li, Z., Ormerod, K.L., Fraser, J.A., Berman, J. and Nielsen, K. (2015) Polyploid titan cells produce haploid and aneuploid progeny to promote stress adaptation. *mBio*, **6**, e01340-01315.

41. Croll, D. and McDonald, B.A. (2012) The accessory genome as a cradle for adaptive evolution in pathogens. *PLoS Pathog*, **8**, e1002608.

42. Chibana, H., Beckerman, J.L. and Magee, P.T. (2000) Fine-resolution physical mapping of genomic diversity in Candida albicans. *Genome Res*, **10**, 1865-1877.

43. Bravo Ruiz, G., Ross, Z.K., Holmes, E., Schelenz, S., Gow, N.A.R. and Lorenz, A. (2019) Rapid and extensive karyotype diversification in haploid clinical Candida auris isolates. *Curr Genet*, **65**, 1217-1228.

44. Brown, G.D. and Netea, M.G. (2012) Exciting developments in the immunology of fungal infections. *Cell Host Microbe*, **11**, 422-424.

45. Pfaller, M.A., Diekema, D.J., Turnidge, J.D., Castanheira, M. and Jones, R.N. (2019) Twenty Years of the SENTRY Antifungal Surveillance Program: Results for Candida Species From 1997-2016. *Open Forum Infect Dis*, **6**, S79-S94.

46. Pfaller, M.A., Castanheira, M., Messer, S.A., Moet, G.J. and Jones, R.N. (2010) Variation in Candida spp. distribution and antifungal resistance rates among bloodstream infection isolates by patient age: report from the SENTRY Antimicrobial Surveillance Program (2008-2009). *Diagn Microbiol Infect Dis*, **68**, 278-283.

47. Todd, R.T. and Selmecki, A. (2020) Expandable and reversible copy number amplification drives rapid adaptation to antifungal drugs. *Elife*, **9**.

48. Mount, H.O., Revie, N.M., Todd, R.T., Anstett, K., Collins, C., Costanzo, M., Boone, C., Robbins, N., Selmecki, A. and Cowen, L.E. (2018) Global analysis of genetic circuitry and adaptive mechanisms enabling resistance to the azole antifungal drugs. *PLoS Genet*, **14**, e1007319.

49. Hirakawa, M.P., Martinez, D.A., Sakthikumar, S., Anderson, M.Z., Berlin, A., Gujja, S., Zeng, Q., Zisson, E., Wang, J.M., Greenberg, J.M. *et al.* (2015) Genetic and phenotypic intra-species variation in Candida albicans. *Genome Res*, **25**, 413-425.

50. Dixon, J.R., Xu, J., Dileep, V., Zhan, Y., Song, F., Le, V.T., Yardimci, G.G., Chakraborty, A., Bann, D.V., Wang, Y. *et al.* (2018) Integrative detection and analysis of structural variation in cancer genomes. *Nat Genet*, **50**, 1388-1398.

51. Yao, Y. and Dai, W. (2014) Genomic Instability and Cancer. *J Carcinog Mutagen*, **5**.

714   52.   Abbey, D.A., Funt, J., Lurie-Weinberger, M.N., Thompson, D.A., Regev, A., Myers,
715   C.L. and Berman, J. (2014) YMAP: a pipeline for visualization of copy number
716   variation and loss of heterozygosity in eukaryotic pathogens. *Genome Med*, **6**, 100.

717   53.   Bogaerts, B., Delcourt, T., Soetaert, K., Boarbi, S., Ceyssens, P.J., Winand, R., Van
718   Braekel, J., De Keersmaecker, S.C.J., Roosens, N.H.C., Marchal, K. *et al.* (2021) A
719   bioinformatics WGS workflow for clinical Mycobacterium tuberculosis complex isolate
720   analysis, validated using a reference collection extensively characterized with
721   conventional methods and in silico approaches. *J Clin Microbiol*.

722   54.   Bogaerts, B., Winand, R., Fu, Q., Van Braekel, J., Ceyssens, P.J., Mattheus, W.,
723   Bertrand, S., De Keersmaecker, S.C.J., Roosens, N.H.C. and Vanneste, K. (2019)
724   Validation of a Bioinformatics Workflow for Routine Analysis of Whole-Genome
725   Sequencing Data and Related Challenges for Pathogen Typing in a European National
726   Reference Center: Neisseria meningitidis as a Proof-of-Concept. *Front Microbiol*, **10**,
727   362.

728   55.   Ellison, M.A., Walker, J.L., Ropp, P.J., Durrant, J.D. and Arndt, K.M. (2020)
729   MutantHuntWGS: A Pipeline for Identifying Saccharomyces cerevisiae Mutations. *G3*
730   *(Bethesda)*, **10**, 3009-3014.

731   56.   Quijada, N.M., Rodriguez-Lazaro, D., Eiros, J.M. and Hernandez, M. (2019) TORMES:
732   an automated pipeline for whole bacterial genome analysis. *Bioinformatics*, **35**, 4207-
733   4212.

734   57.   Thorvaldsdóttir, H., Robinson, J.T. and Mesirov, J.P. (2013), *Brief. Bioinformatics*, Vol.
735   14, pp. 178-192.

736   58.   Leinonen, R., Sugawara, H., Shumway, M. and Collaboration, I.N.S.D. (2011), *Nucleic*
737   *Acids Res.*, Vol. 39, pp. D19-21.

738   59.   Leinonen, R., Akhtar, R., Birney, E., Bower, L., Cerdeno-Tarraga, A., Cheng, Y.,
739   Cleland, I., Faruque, N., Goodgame, N., Gibson, R. *et al.* (2011) The European
740   Nucleotide Archive. *Nucleic Acids Res*, **39**, D28-31.

741   60.   Sloan, C.A., Chan, E.T., Davidson, J.M., Malladi, V.S., Strattan, J.S., Hitz, B.C.,
742   Gabdank, I., Narayanan, A.K., Ho, M., Lee, B.T. *et al.* (2016) ENCODE data at the
743   ENCODE portal. *Nucleic Acids Res*, **44**, D726-732.

744   61.   Howe, K.L., Contreras-Moreira, B., De Silva, N., Maslen, G., Akanni, W., Allen, J.,
745   Alvarez-Jarreta, J., Barba, M., Bolser, D.M., Cambell, L. *et al.* (2020) Ensembl
746   Genomes 2020-enabling non-vertebrate genomic research. *Nucleic Acids Res*, **48**,
747   D689-D695.

748   62.   Arnaud, M.B., Costanzo, M.C., Skrzypek, M.S., Binkley, G., Lane, C., Miyasato, S.R.
749   and Sherlock, G. (2005) The Candida Genome Database (CGD), a community resource
750   for Candida albicans gene and protein information. *Nucleic Acids Res*, **33**, D358-363.

751   63.   Bahl, A., Brunk, B., Crabtree, J., Fraunholz, M.J., Gajria, B., Grant, G.R., Ginsburg, H.,
752   Gupta, D., Kissinger, J.C., Labo, P. *et al.* (2003) PlasmoDB: the Plasmodium genome
753   resource. A database integrating experimental and computational data. *Nucleic Acids*
754   *Res*, **31**, 212-215.

755   64.   Girgis, H.Z. (2015) Red: an intelligent, rapid, accurate tool for detecting repeats de-
756   novo on the genomic scale. *BMC Bioinformatics*, **16**, 227.

757   65.   Li, H. (2013), *arXiv:1303.3997v1 [q-bio.GN]*.

758   66.   Li, H. and Durbin, R. (2009), *Bioinformatics*. 2009/05/20 ed, Vol. 25, pp. 1754-1760.

759   67.   Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
760   Abecasis, G., Durbin, R. and Subgroup, G.P.D.P. (2009), *Bioinformatics*, Vol. 25, pp.
761   2078-2079.

68.   Li, W. and Godzik, A. (2006), *Bioinformatics*, Vol. 22, pp. 1658-1659.

69.   Ramírez, F., Ryan, D.P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne,
S., Dündar, F. and Manke, T. (2016), *Nucleic Acids Res.*, Vol. 44, pp. W160-165.
70.   Cingolani, P., Platts, A., Wang le, L., Coon, M., Nguyen, T., Wang, L., Land, S.J., Lu,
X. and Ruden, D.M. (2012) A program for annotating and predicting the effects of
single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila
melanogaster strain w1118; iso-2; iso-3. *Fly (Austin)*, **6**, 80-92.
71.   Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von
Haeseler, A. and Lanfear, R. (2020) IQ-TREE 2: New Models and Efficient Methods
for Phylogenetic Inference in the Genomic Era. *Mol Biol Evol*, **37**, 1530-1534.
72.   Hoang, D.T., Chernomor, O., von Haeseler, A., Minh, B.Q. and Vinh, L.S. (2018)
UFBoot2: Improving the Ultrafast Bootstrap Approximation. *Mol Biol Evol*, **35**, 518-
522.
73.   Yu, G., Lam, T.T., Zhu, H. and Guan, Y. (2018) Two Methods for Mapping and
Visualizing Associated Data on Phylogeny Using Ggtree. *Mol Biol Evol*, **35**, 3041-
3043.
74.   Krzywinski, M., Schein, J., Birol, I., Connors, J., Gascoyne, R., Horsman, D., Jones,
S.J. and Marra, M.A. (2009), *Genome Res.*, Vol. 19, pp. 1639-1645.
780