Title: GIP: An open-source computational pipeline for mapping genomic

instability from protists to cancer cells

Authors: Gerald F. Späth[1] and Giovanni Bussotti[1,2,]*

Affiliations:

[1]Institut Pasteur, INSERM U1201, Unité de Parasitologie moléculaire et Signalisation, Paris,

France.

[2]Hub de Bioinformatique et Biostatistique - Département Biologie Computationnelle, Institut

Pasteur, 75015 Paris, France

*corresponding author: Giovanni Bussotti,   giovanni.bussotti@pasteur.fr

Abstract:

Genome instability has been recognized as a key driver for microbial and cancer adaptation and

thus plays a central role in many human pathologies. Even though genome instability

encompasses different types of genomic alterations, most available genome analysis software

are limited to just one kind mutation or analytical step. To overcome this limitation and better

understand the role of genetic changes in enhancing pathogenicity we established GIP, a novel,

powerful bioinformatic pipeline for comparative genome analysis. Here we show its application

to whole genome sequencing datasets of *Leishmania*, *Plasmodium, Candida*, and cancer.

Applying GIP on available data sets validated our pipeline and demonstrated the power of our

analysis tool to drive biological discovery. Applied to *Plasmodium vivax* genomes, our pipeline

allowed us to uncover the convergent amplification of erythrocyte binding proteins and to

identify a nullisomic strain. Re-analyzing genomes of drug adapted *Candida albicans* strains

revealed correlated copy number variations of functionally related genes, strongly supporting a

mechanism of epistatic adaptation through interacting gene-dosage changes. Our results

illustrate how GIP can be used for the identification of aneuploidy, gene copy number

variations, changes in nucleic acid sequences, and chromosomal rearrangements. Altogether,

GIP can shed light on the genetic bases of cell adaptation and drive disease biomarker

discovery.

29
30

One Sentence Summary:

GIP - a novel pipeline for detecting, comparing and visualizing genome instability.

Keywords:

Genome instability, aneuploidy, copy number variations, single nucleotide variation, structural

variation, genetic adaptation.

Main Text:

In recent years, the field of genomics has rapidly expanded with a fast increase in the number

of newly sequenced genomes (1). This surge is a direct consequence of the development of new

and ever more efficient, high-throughput capable sequencing technologies (2). On the one hand,

the improvement in long reads technology allowed the generation of high-quality genome

assemblies (3,4). On the other hand, the decreasing costs for short-reads sequencing and the

parallel increase in sequencing throughput propelled the exponential increase of available

whole genome sequencing (WGS) data (5). Thanks to these advances one can reasonably expect

WGS to rapidly become a key component of personalized medicine and clinical applications.

In this context several consortia-based projects have been established with the goal to produce

WGS for the study of different biological systems (5), and a number of publicly available

databases have been compiled or updated (6). Parallel to data availability, many bioinformatics

tools have been developed to perform specific genome analysis tasks (7,8). For instance, tools

such as Freebayes (9), CNVnator (10) and DELLY (11) have been respectively used for the

detection or characterization of DNA single nucleotide variants (SNVs), copy number

variations (CNVs), and structural variations (SVs), but their scope is limited to the analysis of

one genomic feature at the time. A number of integrative WGS pipelines and workflows have

been established combining the execution of multiple bioinformatics algorithms serving

different analysis steps (12). Even though continuous progress has been made, there is no

standardized or unified approach for genomic investigation. For the development of improved

WGS data analysis pipelines, several important requirements need be considered, including

portability, reproducibility, scalability and compatibility with high-performance computing

(HPC) clusters and remote cloud computing. Here we introduce a novel genome instability

pipeline (GIP) that fulfills all these requirements. GIP facilitates the genome-wide detection,

quantification, comparison and visualization of chromosome aneuploidies, gene CNVs, SNVs

and SVs. GIP is implemented in Nextflow (13), a workflow language that allows to execute

GIP seamlessly in local workstation, on an HPC or remotely in the cloud. All required

environment and software dependencies of GIP are fulfilled and provided with a Singularity

container, thus making GIP reproducible, easy-to-install and easy-to-use. GIP allows the use of

giptools, a tool-suit of R-based modules for genome data exploration, enabling the comparison

of sample sub-sets. GIP and giptools generate a summary report with publication-quality figures

and spreadsheet tables. GIP and giptools constitute a single framework for WGS analysis

suitable both for large scale batch analysis of individual genomes and comparison of samples

from different experimental conditions or origins. Lastly, a key strength of GIP and giptools is

the general applicability to different eukaryotic species. We already successfully applied GIP

on the analysis of *Leishmania* genomes (14-16). In this study, we validate the use of GIP and

giptools using WGS data from published datasets of the three major human pathogens

Leishmania infantum, *Plasmodium vivax* and *Candida albicans* and as well as three human

cancer cell lines. Furthermore, we demonstrate how the extensive and powerful analytical

approach operated by GIP and giptools can be used to find new biological signal that escaped

previous analyses.

Results:

The GIP workflow. GIP is a tool for scientific investigation compatible with Linux systems,

requiring minimal configuration and distributed as a self-contained package. GIP consists of

three files: the Nextflow pipeline code, the configuration and the Singularity container files.

The minimum required input is a paired-end WGS data set and a reference genome assembly

in the standard fastq and FASTA format, respectively. GIP analyses include (i) extracting

genomic features such as assembly gaps or repetitive elements, (ii) mapping the reads, (iii)

evaluating chromosome, gene and genomic bin copy numbers, (iv) identifying and visualizing

copy number variation with respect to the reference genome, (v) identifying and quantifying

gene clusters, (vi) detecting and annotating SNVs, (vii) measuring non-synonymous (N) and

synonymous (S) mutations for all genes, (viii) detecting SVs including tandem duplications,

deletions, inversions and break-ends translocations using split-read and read-pair orientation

information, and (ix) producing a report file providing summary statistics, tables and

visualizations (**Fig. 1**). GIP allows to customize filtering and visualization options via the

configuration file (see methods). The output of GIP can be used as input for giptools, a tool-

suite to compare sample sub-sets and highlight chromosome copy number, gene copy number

and SNV differences.

Applying giptools on a *Leishmania infantum* case study. GIP permits the batch analysis of a

set of individual samples, where each sample is considered separately and compared only with

respect to the provided reference genome assembly. As a consequence, all variants and copy

number alterations detected in a sample merely reflect the differences between the sequenced

and the reference genomes. While this application may be sufficient in some circumstances,

research projects often involve downstream comparison between samples. Examples include

the comparison of gene or chromosome copy variation number between for example drug

resistant and drug susceptible samples, or the juxtaposition of SNVs detected in isolates from

different geographic areas. For this purpose, we developed giptools, a suit of thirteen modules

that allows to compare samples processed by GIP (**Table 1**). All modules in giptools are fully

embedded in the Singularity container and they are provided with their own documentation.

Table 1: giptools modules

Module name	Purpose
karyotype	Compare the chromosome sequencing coverage distributions
binCNV	Compare bin sequencing coverage in 2 samples
geCNV	Compare gene sequencing coverage in 2 samples
ternary	Compare gene sequencing coverage in 3 samples
ternaryBin	Compare bin sequencing coverage in 3 samples
SNV	Compare SNVs in multiple samples
binDensity	Density plot of bin sequencing coverage of many samples
geInteraction	Detect CNV genes in many samples and produce correlation-based networks
genomeDistance	Compare samples genomic distance
phylogeny	Extract the SNVs union and infer the phylogenetic tree
convergentCNV	Detect convergent CNV gene amplifications
overview	Overview of the sequencing coverage of chromosomes, genomic bins and genes
panel	Extract genomic information of a gene panel

110 To illustrate the type of exploratory data analyses and the biological questions that can be

111 addressed, we tested giptools on a previously analyzed dataset of seven clinical *Leishmania*

112 *infantum* isolates from Tunisia (14). *Leishmania* is the etiological agent of leishmaniasis, a life-

113 threatening human and veterinary disease affecting 12 million people worldwide (17). The

parasites were derived from patients affected with visceral leishmaniasis, expanded in cell

culture and their genomes were sequenced. The dataset includes four Glucantime drug

susceptible isolates and three isolates from relapsed patients, and thus comparison may inform

on genetic factors resulting in treatment failure. giptools allowed the detection and visualization

of pervasive intra-chromosomal CNVs across the thirty-six *Leishmania* chromosomes (**Fig.**

2A). Additionally, giptools enables targeted comparison of normalized genomic bin sequencing

coverage of sample pairs. We used giptools' `binCNV` module to compute the ratio between

corresponding genomic bins of the strains LIPA83 over ZK43, which correspond to a first-

episode and a relapse leishmaniasis isolate, respectively. giptools represents different

chromosomes as separate panels (**Fig. 2B**), as part of single genome-wide overview (**Fig. S1A**)

or as distinct plots (**Fig. S1B**). This analysis allowed the identification of 2,905 and 2,208 bins

that were respectively amplified or depleted in LIPA83 with respect to ZK43. The results are

returned by giptools as a Microsoft Excel table (.xlsx format) providing ratio scores at each

genomic position (**Table S1**). Likewise, giptools permits three-way comparisons of normalized

genomic bin sequencing coverage with ternary plots (**Fig. 2C**). We used this representation to

display the genomic bin relative abundance in samples ZK43, LIPA83 and ZK28. The analysis

shows important strain-specific differences in bin copy number that are visualized by shifts of

the signals out of the center. Similar to genomic bin analysis, giptools makes it possible to

compare the sequencing depth of annotated genes and thus the copy number in two or three

samples (**Fig. 2D** and **Fig. S2**). However, the determination of gene copy number might be

impeded by (i) short read length or fragment insert size, (ii) the complexity of the target genome,

or (iii) the presence of repetitive elements. The read map quality (MAPQ) score is a measure

that reflects how much each gene is supported by unambiguously mapped reads (high MAPQ)

in contrast to multimapping reads (low MAPQ). Together with the coverage, GIP also computes

the mean read MAPQ score for each gene and allows a different strategy to determine the copy

number of low MAPQ genes (see methods). The evaluation of LIPA83 and ZK43 gene

coverage ratio scores revealed 13 gene CNVs with a stringent MAPQ cutoff of 50 (**Table S2**).

The maximum normalized coverage value of 6.5 was observed for a putative amastin surface

glycoprotein (LINF_310009800). Other examples of gene CNVs in this set include the putative

surface antigen protein 2 (LINF_120013500) and the heat shock protein HSP33

(LINF_300021600) (**Table S2**). Genes falling below a user defined MAPQ score and sharing

high level of sequence similarity are assigned to the same gene cluster, and their measured

coverage scores are averaged across all members of the group. Low MAPQ scores can also be

associated with single genes, e.g. in the case of internal repetitive elements that cause multiple

ambiguous alignments inside the gene itself, or if mapping occurs in possibly misannotated

intergenic regions. The LIPA83/ZK43 comparison showed 27 CNV gene clusters, including

cluster clstr303 (3 genes annotated as 'amastin-like') and cluster clstr16 (2 tb-292 membrane-

associated protein-like proteins) (**Table S2**). These results demonstrate the power of GIP and

giptools to detect and compare intra-chromosomal CNVs in *Leishmania* at genomic bin level.

Conveniently, analogous two- or three-ways comparisons can be applied to reveal copy number

variations at individual gene or gene cluster levels.

Comparative genomics of a *Plasmodium vivax* WGS dataset. We next applied GIP and

giptools on other biological systems to demonstrate its broad applicability outside the

Leishmania field, including the human apicomplexan parasite *Plasmodium vivax*. *Plasmodium*

vivax is a protist parasite and a human pathogen causing malaria. *Plasmodium vivax* gives rise

every year to 130 million clinical cases (18), and it is estimated that 2.5 billion people are at

risk of infection worldwide (19-21). We applied GIP and giptools to investigate genomic

variations across a sizeable dataset of 222 *Plasmodium vivax* genomes isolated from clinical

samples of 14 countries worldwide (22,23) (**Table S3**). The phylogenetic tree reconstruction

and PCA analyses (**Fig. 3A** and **B**) showed a high correlation between genotypes and the

geographic origin of the samples. However, we detected substantial genomic variability

between isolates collected at smaller geographical scale, with 14,555 SNVs (~42% of the total)

uniquely characterizing representative samples from five Ethiopian study sites (23) (**Fig. 3C**

and **D**). This result may reflect diverging evolutionary trajectories radiating from few founder

strains. At gene level we profiled the copy number variations of two gene panels. The first panel

accounts for 43 previously described genes encoding for potential erythrocyte binding proteins

suggested to operate at the interface of the parasite-host invasion process (23). The second panel

includes two drug resistance markers comprising the chloroquine resistance transporter

PVP01_0109300 and the multidrug resistance protein 1 PVP01_1010900, and four proteins

implicated in red blood cell invasion, such as the merozoite surface protein gene MSP7

PVP01_1219700, the reticulocyte binding protein gene 2c PVP01_0534300, the serine-repeat

antigen 3 PVP01_0417000, and the reticulocyte binding protein 2b PVP01_0800700) (24-33).

Read depth analysis indicated that four genes in the panel (PVP01_0623800, PVP01_1031400,

PVP01_1031200, PVP01_1031300) show a high degree of variability, with amplifications

observed in samples from distinct geographic (**Fig. 3E**). This convergence is sign of strong

natural selection, which further sustains the functional importance of these genes in the

infection process. Furthermore, 6 genes positioned on chromosome 14 are absent in the Thai

strain PD0689_C as a result of the loss of this chromosome (nullisomy) (**Fig. 3E** and **F**). Finally,

the comparison of synonymous and non-synonymous SNVs in the panel of genes revealed

important differences between sample groups. Our analysis indicates an overall higher number

of non-synonymous mutations in Ethiopian compared to Cambodian isolates, therefore

suggesting a stronger evolutionary pressure acting on the African strains (**Fig. 3G**). Taken

together these analyses well illustrate how GIP and giptools can be readily applied for bulk

analysis of *Plasmodium vivax* genomes to assess genome diversity, extract evolutionary

information and identify potential disease biomarkers.

Gene CNV analysis of *Candida albicans* evolutionary adapted strains. We next applied GIP

and giptools to the human fungal pathogen *Candida albicans*, an opportunistic yeast exhibiting

major genome plasticity (34-43) and causing hundreds of thousands of severe infections each

year (44). Candidemia, a bloodstream infection with *Candida*, are often associated with high

rates of morbidity and mortality (15–50%) notwithstanding existing antifungal treatments

(45,46). We applied GIP and giptools to a *Candida albicans* WGS dataset described in a recent

study that covers five different progenitor strains (P75063, P75016, P78042, SC5314,

AMS3050) and investigates CNVs driving tolerance and resistance to anti-fungal azole drugs

(47). We analyzed nineteen samples, including (i) four clinical isolates (P75063, P75016,

P78042, SC5314), (ii) seven strains selected *in vitro* against the anti-fungal drug fluconazole

(FLC) (AMS4104, AMS4105, AMS4106, AMS4107, AMS4397, AMS4444, AMS4702), (iii)

four isogenic colonies adapted to the drug miconazole (AMS3051, AMS3052, AMS3053 and

AMS3054) together with their progenitor (AMS3050), and (iv) three colonies derived from a

miconazole-adapted population and isolated on a rich medium (AMS3092, AMS3093 and

AMS3094) (47-49). GIP and giptools were able to reproduce previous observations of the

amplification of the genes for the drug efflux pumps TAC1 (orf19.3188) and ERG11

(orf19.922), for the stress response proteins HSP70 (orf19.4980), CGR1 (orf19.2722), ERO1

(orf19.4871), TPK1 (orf19.4892), ASR1 (orf19.2344), PBS2 (orf19.7388) and CRZ1

(orf19.7359), and for proteins involved in membrane and cell wall integrity, including CDR3

(orf19.1313), NCP1 (orf19.2672), ECM21 (orf19.4887), MNN23 (orf19.4874), RHB1

(orf19.5994) and KRE6 (orf19.7363) (**Table S4**). Furthermore, the powerful comparative

approach of our pipeline permitted the discovery of 1,505 genes showing correlating or anti-

correlating copy number variations (**Fig. 4A** and **B**, **Table S4**), which could be assigned to nine

distinct correlation clusters (CC) (**Fig. S3A**, **Table S5**) that escaped previous analyses. We

verified the sequencing coverage of genomic regions encompassing gene CNVs, including

three regions amplified in fluconazole resistant strains (**Fig. 4B**, **Fig. S3B**) (47) and a region

whose amplification correlates with the level of miconazole resistance (47) (**Fig. S3C**), as well

as the loss of heterozygosity associated to the depletion of chromosome 3 left arm in sample

AMS3051 (**Fig. S3D**) (47). Eventually, by representing genes and absolute correlation

respectively as nodes and edges of a network, we identified 9 highly interconnected network

clusters (NC) (**Fig. 4C**, **Table S6**). NC7, NC8 and NC9 embody genes from individual

chromosomes, respectively chromosomes 1, 3 and 4. The most parsimonious explanation for

the high levels of correlation observed in these NCs (**Fig. 4C**) is the occurrence of sub-

chromosomal amplifications affecting several adjacent genes. A different scenario is pictured

for each of the remaining NCs (NC1-6) where the genes are located on different chromosomes

thus suggesting genetic interactions that causes coordinated changes in gene copy number. The

gene ontology (GO) and metabolic pathway analyses revealed a significant functional

enrichment of genes expressed on the cell surface and involved in the interaction with the host

(NC2), gibberellin biosynthesis (NC3), transmembrane nucleobase transporters (NC4) and

gluconeogenesis (NC5) (**Table S7**). Altogether, GIP and giptools are validated by reproducing

previously published results, and beyond that can drive new biological findings as documented

by the discovery of a network of epistatic CNV interactions supporting genomic adaptation in

Candida albicans populations under drug selection.

Exploring instability of larger genomes using cancer cell lines as a benchmark. The larger

genome size, and the higher number of genes and WGS reads can represent a challenge when

working with higher eukaryotes. For the purpose of comparison, the human genome is ~216

times larger than the one of *Candida albicans* we analyse in this study. Therefore, we sought

to evaluate the applicability of the GIP and giptools framework to human data and utilized a

panel of genomes from cancer cell lines as a test set. In our analyses we considered publicly

available WGS data of the cell lines T47D, NCI_H460 and K562 (50), which respectively

derive from human breast, lung and blood cancers. The karyotype analysis revealed aneuploidy

for all chromosomes except chromosome 4 (**Fig. 5A**). The observed heterogeneity in read depth

across chromosomes, illustrated by large interquartile range in the boxplot, suggests sub-

chromosomal or episomal copy number variations, or the co-existence of karyotypically

different sub-populations. Indeed, the coverage analysis confirmed the pervasive presence of

CNVs both at chromosomal and sub-chromosomal levels (**Fig. S4**) with remarkable instability

observed for specific chromosomes, e.g. chromosomes 6, 9, 10 and 16 (**Fig. 5B**). Overall, we

detected 1,647,016 SNVs (**Supplementary Data 1**) and allele frequency shifts with respect to

the reference genome, suggesting haplotype selection and the preferential expression of distinct

alleles in different cell lines (**Fig. 5C** and **Fig. S5**). Furthermore, we identified repeated loss of

heterozygosity events and uneven distribution of SNVs that form "patches" of high frequency

correlating with chromosomal and sub-chromosomal CNVs (**Fig. 5D**, **Fig. 5E** and **Fig. S6**).

These results identify GIP and giptools as a powerful new platform to reveal loci, genes or

alleles that are under natural selection in cancer cells, thus allowing important new insight into

the genetic basis of tumor development, cancer cell evolution and drug resistance.

Discussion:

Genome instability is a key driver of evolution for microbial pathogens and cancer cells (51)

and a major source of human morbidity. Here we introduce GIP and giptools, an integrated

framework for the genotype profiling of biological systems exploiting genome instability for

adaptation. We document the power and versatility of GIP and giptools by performing genomic

screenings of three major pathogenic eukaryotes and human cancer cell lines. While originally

deployed for *Leishmania* genome analysis, in this study we validate the use of our pipeline on

other organisms reproducing expected results. For example, in *Candida albicans* we confirmed

the CNVs correlating to drug resistance as well as a loss of heterozygosity event (**Fig. S3B-C**

and **D**). Parallel to this we also show how GIP and giptools can be used for data mining and

scientific discovery. New findings include (i) the discovery of the convergent amplification of

erythrocyte binding proteins in *Plasmodium vivax* strains sampled from distinct geographic

areas (**Fig. 3E**), (ii) the detection of a nullisomic strain (**Fig. 3F**), (iii) the identification of

correlated copy number variations between genes positioned on separate chromosomes of

Candida albicans adapting strains, and (iv) the functional association of such genes, strongly

supporting a mechanism of epistatic interactions exerted through gene-dosage changes, and

corroborating previous reports on adapting *Leishmania* populations (52).

Importantly, GIP and giptools overcome key limitations of current analysis tools, such

as the breadth of analysis that is often limited to individual types of mutations, and the lack of

genome-wide, comprehensive reports. To ease genome instability investigations our pipeline

offers a single solution to karyotype, gene CNV, SNV and SV batch analyses, providing

summary reports and high-quality, genome-wide visualizations. Furthermore, many current

tools identify variations with respects to a reference assembly only, which leaves the between

samples comparisons to external tools that need installing, may be incompatible in terms of file

format, and may rely on different analytical assumptions. To address this limitation giptools

enables custom sample comparisons, to explore differences and common features between

genomes, and provides a vast choice of analytical tools with compatible features. Likewise,

current tools are often restricted to the analysis of data from one or few species only (53-57),

but are not generally applicable to different biological systems, which interferes with the

investigation of genome variations across multiple species and the exploration potentially

conserved genomic adaptation mechanisms. By contrast, our pipeline limits as much as possible

the use of hardcoded parametrization, which could limit its use to a specific organism.

Therefore, GIP's flexible design makes it adapted for the genome analysis of both model and

non-model organisms, including *Leishmania* or human.

Many current tools are further limited in software portability and reproducibility across

different computer environments, which can produce faulty results calling their clinical

application into question. Conversely, thanks to the Singularity implementation all required

software are embedded and provided within the software container. As a consequence, users

can easily recreate the same work environment just by downloading the pipeline container, and

reproduce exactly the same publication-quality plots and tables presented in this study. Lastly,

one more common limitation is posed by software scalability. In the WGS domain, with the

rapid increase new samples made available and the enormous amount of data generated in each

sequencing run, the CPU and memory resources of local workstation risk to quickly become

inadequate for data analysis. Therefore, it is paramount that WGS tools are implemented to run

on high-performance computing (HPC) clusters and feature remote cloud computing solutions.

Because of its Nextflow implementation GIP can be executed on a local machine, on cluster

resource manager or the cloud. GIP can be applied on individual samples and without additional

effort on large WGS data sets for batch computation as shown for the 222 *Plasmodium vivax*

genomes.

These results well illustrate how GIP and giptools can be applied to perform extended

genomic analyses in different biological systems and drive biomedical discovery. To conclude,

we believe that GIP and giptools represent a step forward toward reproducible research in

genomics, and provide a robust computational framework to study how microbes and tumor

cells harness genome instability for environmental adaptation and fitness gain.

Figures

Fig. 1

GIP workflow. The schema on the left recapitulates the GIP inputs, processes and outputs (see

methods). Blue, orange, green and purple boxes indicate genome reference, read mapping,

quantification and variants computation steps, respectively. The panels on the right demonstrate

example plots included in the GIP report computed for individual samples. The "Karyotype"

plot shows the coverage distributions for each chromosome (y-axis). The "Genomic bins" plot

shows the genomic position (x-axis) and the normalized genomic bin sequencing coverage (y-

axis). The "Gene CNVs" plot shows the normalized gene sequencing coverage. The "Structural

variants" panel shows a Circos plot representing break-end translocations (black links in the

inmost part of the plot), and other possible structural variations in the outer tracks, including

insertions, duplications, deletions and inversions. The outmost track shows the normalized

sequencing coverage. The "Single nucleotide variants" plot shows on the x and y axes

respectively the genomic position and variant allele frequency of detected SNVs.

Fig. 2:

Comparing *Leishmania infantum* genomes with giptools. (**A**) Density plot representing the

genomic coverage of the seven *Leishmania infantum* isolates. The x-axis shows the log 10

normalized coverage of genomic bins. The y-axis reflects the genomic position. The thirty-six

different chromosomes are materialized as separate panels. The blue shading indicates the (2D)

kernel density estimates of genomic bins. The two red vertical lines mark the 1.5 and 0.5

coverage values. A selection of 50 000 bins with coverage > 1.5 or < 0.5 is shown as black dots.

(**B**) Scatterplot of the genomic bin normalized sequencing coverage ratio of samples LIPA83

over ZK43. The x any y axes show the ratio score and the genomic position respectively. Ratio

scores > 1.25 are labelled in orange and indicate genomic bin amplification. Ratio scores < 0.75

are labelled in blue and indicate genomic bin depletion. (**C**) Ternary comparison showing the

relative abundance of the genomic in samples LIPA83, ZK43 and ZK28. The axes report the

fraction of the bins normalized sequencing coverage in the three strains. The blue contour

indicates the log10 bin density. A subset of 5,000 bins is shown as black dots. Each given point

in the plot is adding up to 100. The density area at the center of the plot indicates bins with

equal copy number and thus a ~33 distribution across the three axes. (**D**) Scatterplot showing

the log10 normalized sequencing coverage of annotated genes in ZK43 (x-axis) and LIPA83

(y-axis). The red line indicates the bisector. Dots represent individual genes. (**E**) Sequencing

coverage ratio of gene clusters in samples LIPA83 and ZK43. Dots represent gene clusters. For

plots **D** and **E,** the ratio scores > 1.5 or < 0.5 are labelled in orange and blue respectively.

Fig. 3:

Plasmodium vivax genomic diversity. (**A**) Predicted maximum likelihood phylogenetic tree

reconstruction. (**B**) PCA analysis of the phylogenetic distances estimated from the tree in (**A**).

Each dot indicates a sample. The colour code reflects the geographic origin of the samples and

matches with the colours of the legend in (**A**). (**C**) Venn diagram comparing the SNVs of five

representative Ethiopian strains. (**D**) Pairwise scatterplot comparing the variant allele frequency

of all detected SNVs in the five Ethiopian strains. (**E**) Gene panel analysis. The x-axis reports

a set of genes of interest. The y-axis indicates the normalized mean gene coverage. The boxplots

demonstrate the coverage values distributions for each gene across all samples. Each dot

represents the coverage of the indicated gene in a given sample. Dot colours reflect the sample

geographic origin as in (**A**). (**F**) Reads per kilo base per million mapped reads (RPKM)

normalized sequencing coverage density track of sample PD0689_C. The boundaries of the 14

chromosomes are shown on the bottom. (**G**) Comparison of non-synonymous (N) and

synonymous (S) mutations between Ethiopia and Cambodia sample groups. Dots represent

genes. The x-axis represents the difference between the mean non-synonymous mutation count

in the two sample groups. The y-axis represents the difference between the mean synonymous

mutation count in the two sample groups. The dot size demonstrates the ratio of the mean

normalized sequencing coverage between the two sample groups for each gene. Red and blue

dot colors indicate genes belonging to the 43 genes panel (23) and the custom 6 genes panel

respectively.

Fig. 4:

Gene CNVs interactions. (**A**) All-vs-all normalized sequencing coverage correlation heatmap.

The heatmap is symmetrical along its diagonal and reports both on the rows and the columns

the detected gene CNVs. The colour scale indicates with green and pink high levels of positive

and negative Pearson correlation, respectively. The side ribbons demonstrate in different

colours the chromosome and the correlation cluster of each gene. (**B**) Gene CNV heatmap. The

columns and the rows report respectively the samples and the detected gene CNVs. The colour

scale indicates the normalized sequencing coverage of the genes. To ease visualization,

coverage values greater than 3 are reported as 3 (red). Black boxes highlight the genomic

regions shown in **Fig. S3B** (panels 1, 2 and 3) and **Fig. S3C**. The ribbons on the left indicate

the chromosome and the correlation cluster of each gene. Top ribbons indicate the genotype

and the strains resulting from the different evolutionary experiments. (**C**) Gene interaction

network. Nodes indicate gene CNVs. Edges reflect the absolute Pearson correlation value. The

closer the nodes are, the higher is the correlation. Only significant interactions (Benjamini-

Hochberg adjusted p-value < 0.01) are shown. The colour of the edges indicates in red and blue

respectively positive and negative correlations. The colour of the nodes denotes the predicted

network cluster for each gene.

Fig. 5:

Cancer cell lines genome instability. Green, brown, and grey colours indicate respectively

samples K562, NCI_H460 and T47D. (**A**) Chromosome coverage analysis. The x-axis reports

the chromosomes. The y-axis reports the estimated somy score. The boxes show the somy score

distributions. (**B**) Sub-chromosomal copy number variation. Dots indicate genomic bins.

Different panels indicate different chromosomes. The panel columns indicate from left to right

four selected chromosomes: 6, 9, 10 and 16. The panel rows show top to bottom the samples

K562, NCI_H460 and T47D. The x-axis indicates the genomic position. The y-axis indicates

the normalized genomic bin sequencing coverage values. Coverage values greater than 5 are

reported as 5. (**C**) SNV frequency density plots. The four different panels represent different

selected chromosomes: 6, 9, 10 and 16. The x-axis reports the variant allele frequency. The y-

axis the estimated kernel density between 0 and 3. (**D**) SNV frequency scatter plots. The four

different panels represent different selected chromosomes: 6, 9, 10 and 16. The x-axis indicates

the genomic position. The y-axis indicates the variant allele frequency. (**E**) Chromosome 11

combined SNV and bin coverage plot. To ease visualization, giptools allows the simultaneous

displaying of variant allele frequencies (y-axis, left) and sequencing coverage (y-axis, right).

Dots represent SNVs. The lines represent the normalized bin sequencing coverage. The x-axis

indicates the genomic position. Coverage values greater than 5 are shown as 5.

Supplementary figures

Fig. S1:

Bin coverage ratio visualizations. The figure shows alternative representations produced for the

genomic bin normalized sequencing coverage ratio of samples ZK43 and LIPA83. Same layout

as **Fig. 2B**. (**A**) Whole genome overview. (**B**) All the individual chromosomes separately. The

plot shows the example of chromosome 33.

Fig. S2:

A

B

417 Two-ways and three-ways gene coverage comparisons. (**A**) Log10 normalized sequencing

418 coverage comparison of genes in ZK43 (x-axis) and LIPA83 (y-axis). Same layout a **Fig. 2D**

419 but representing the different chromosomes in individual panels. (**B**) Ternary plot showing the

420 relative abundance of genes (green dots) and gene clusters (red dots) in samples LIPA83, ZK43

421 and ZK28. The axes report the fraction of the genes and genes clusters normalized sequencing

422 coverage in the three strains. Each given point in the plot adding up to 100. Genes with equal

423 copy number are shown in the center of the ternary plot, while copy number variations are

424 visualized by shifts of the dots out of the center. The dots size reflects the fold change variation

425 between the maximum and the minimum observed normalized coverage value.

Fig. S3:

Copy number variati... (**A**) Gene CNV

correlation clusters. ... s and the scaled

normalized sequencing coverage of the genes in each cluster. Different panels indicate different

correlation clusters. (**B**) Scatterplots showing the bin normalized sequencing coverage (y-axis)

of three genomic regions (chr4:400,000-1,000,000; chr3:1,250,000-1,500,000; chr1:2,000,000-

2,500,000) in three different samples AMS4702, AMS4104 and AMS4107. The x-axis shows

the genomic position. Dots represent genomic bins. (**C**) Scatterplot showing the bin normalized

sequencing coverage of a genomic region (chr3:500,000-1,000,000) in five different samples

represented as separate panels. From top to bottom the samples are: AMS3050, AMS3053,

AMS3054, AMS3052 and AMS3051. The x and y axes show respectively the genomic position

and the normalized sequencing coverage. Dots represent bins. (**D**) Comparative analysis of

chromosome 3 SNVs. AMS3050 SNVs are shown in green, while AMS3051 SNVs are shown

in grey. The x and the y axes show respectively the genomic position and the variant allele

frequency.

Fig. S4:

Genomic coverage analysis of three cancer cell lines. The x-axis indicates the genomic position. The y-axis indicates the normalized genomic bin sequencing coverage values. Dots demonstrate the genomic bins. To ease visualization, coverage values greater than 5 are reported as 5. Different panels show different chromosomes. The three panel rows indicate top to bottom the following samples: K562, NCI_H460 and T47D.

 Fig. S5:

451 SNV frequency density plot. The x-[ ] y. The y-axis the

452 estimated kernel density between 0 a[ ] grey slopes indicate respectively

453 samples K562, NCI_H460 and T47D.

455   **Fig. S6:**

457   SNV heterogeneity of cancer cell lines   tion. The y-axis

458   indicates the variant allele frequency. T   they are coloured according

459   to the sample. K562, green; NCI_H460, brown; T47D, grey. Different panels show different

460   chromosomes.

Methods:

GIP and giptools. All results presented in this study were generated using GIP and giptools version 1.0.9. GIP code is maintained and freely distributed at the github page: https://github.com/giovannibussotti/GIP. giptools container is accessible from the Singularity cloud at https://cloud.sylabs.io/library/giovannibussotti/default/giptools. The GIP configuration files (**Supplementary Data 2**) and the giptools command options used to generate all results (**Supplementary Data 3**) are provided. The full documentation of GIP and giptools including a description of all options is available from https://gip.readthedocs.io/en/latest/.

Read alignment. WGS reads were downloaded from the Sequence Read Archive (SRA) (58) and the European Nucleotide Archive (ENA) (59) repositories and the Encyclopedia of DNA Elements (ENCODE) dashboard (60) (**Table S3**). For *Leishmania infantum* the GCA_900500625 genome reference and gene annotations available from the ENSEMBL protists (61) server (release-48) were used. For *Candida albicans* the assembly 21 of the SC5314 strain genome reference and gene annotations available from the *Candida* Genome Database (CGD) (62) were used. For *Plasmodium vivax* the P01 reference genome and gene annotations available from PlasmoDB (63) (release-50) were used. For the cancer cell lines the human genome GRCh38 primary assembly and gene annotations available from ENSEMBL (release-102) were used. The repetitive elements of reference genomes were soft-masked by GIP using Red (64). WGS reads were mapped by GIP using BWA-mem (version 0.7.17) (65,66) run with option -M to label shorter split hits as secondary. Then the alignment files were sorted, indexed and reformatted by GIP using Samtools (version 1.8) (67). Finally, read duplicates were removed by GIP using Picard MarkDuplicates (http://broadinstitute.github.io/picard) (version 2.18.9) with the option

"VALIDATION_STRINGENCY=LENIENT." In the four considered datasets, WGS reads

were aligned against full assemblies, including unsorted contigs if present. However just the

canonical assembled chromosomes were considered for all downstream analyses (`chrs` option,

Supplementary Data 2). A minimum read alignment MAPQ score was adopted to select genes

for cluster analysis, and to call for SNVs and SVs (`MAPQ` option, **Supplementary Data 2**).

Altogether a total of 6,306,951,266 reads were aligned to the respective reference genomes.

The `giptools overview` module was run to gather the alignment statistics as estimated by

Picard CollectAlignmentSummaryMetrics (**Table S8**).

Genomic bins and genes quantification. GIP was used to evaluate the mean sequencing

coverage and the mean read MAPQ of genomic bins and genes. For genomic bins, GIP

partitioned the input genomes into adjacent windows of user defined lengths (`binSize` option,

Supplementary Data 2). The coverage GC-content score bias was corrected (`CGcorrect`

option, **Supplementary Data 2**) fitting a LOESS regression with a 5-fold cross validation to

optimize the model span parameter. A larger window length was utilized to bin the reference

genomes for Circos plot representations (`binSizeCircos` option, **Supplementary Data 2**). In

Fig. 1 ("Genomic bins" and "Gene CNVs" plots), **Fig. 2**, **Fig. S1**, **Fig. S2** and **Fig. 4** bins and

genes coverage scores were normalized by median chromosome coverage to highlight

amplifications or depletions with respect to the chromosome copy number. In **Fig. 1**

("Structural variants" plot), **Fig. S3B-C**, **Fig. 3E-G**, and **Fig. S4A** bins and genes coverage

scores were normalized by median genome coverage to account for sequencing library size

differences. GIP evaluated statistically significant copy number variant bins and genes (**Fig. 1**

"Genomic bins" and "Gene CNVs" plots) using a p-value threshold of 0.001

(`covPerBinSigOPT` and `covPerGeSigOPT` options, **Supplementary Data 2**). Estimated p-

values for bins and genes CNVs were corrected for multiple testing using the Benjamini –

Yekutieli (`--padjust BY`) and the Benjamini – Hochberg (`--padjust BH`) methods. The somy

scores shown in **Fig. 1** (karyotype plot) and **Fig. S4B** were computed multiplying the median

genome coverage normalized bin coverage by 2. GIP enabled the CNV analysis of genes

sharing high sequence identity by clustering the nucleotide sequences of the genes with low

mean MAPQ score into groups with cd-hit-est (version 4.8.1) (68) with options `-s 0.9 -c 0.9 -

r 0 -d 0 -g 1`. Then for each gene cluster GIP computed the mean gene coverage normalized by

median chromosome coverage (**Fig. 2E**, **Fig S2B**).

Gene ontology and metabolic pathway enrichment. The FungiDB online tool (Release 52,

20 May 2021) (69) was used to evaluate the functional enrichment of network clusters genes.

For the gene ontology analysis, the biological process (BP), molecular function (MF) and

cellular compartment (CC) terms enrichments were tested, considering both computed and

curated evidences and a p-value cutoff of 0.05. For the metabolic pathway enrichment, both

KEGG (70) and MetaCyc (71) pathway sources were considered with a p-value cutoff of 0.05.

Terms and pathways with Benjamini – Hochberg adjusted p-values < 0.05 were considered

statistically significant.

Sequencing coverage density estimates. GIP was used to convert the read alignment files

(.bam format) in binary data files reflecting sequencing coverage (.bigWig format). The

coverage file were produced using bamCoverage from the deepTools2 suite (72) (version 3.5.1)

with options "--normalizeUsing RPKM --ignoreDuplicates --binSize 10 --smoothLength 30"

(`bigWigOPT` option, **Supplementary Data 2**). The coverage track of sample PD0689_C was

visualized with IGV using the `Bar Chart`, `Autoscale` and windowing function `Mean`

options.

Single-nucleotide variant analysis. GIP was used to call SNVs using Freebayes (version

1.3.2) (`freebayesOPT` option, **Supplementary Data 2**) and filter its output

(`filterFreebayesOPT` option, **Supplementary Data 2**). Filters included the minimum allele

frequency (`--minFreq`), the minimum number of reads supporting the alternative alleles (`--minAO`) and minimum mean mapping quality of for the reads supporting the reference (`--minMQMR`) or the alternative allele (`--minMQM`). A higher number of reads supporting the variants was requested for predictions positioned inside simple repeats of the same nucleotide (homopolymers) (`--minAOhomopolymer`). The homopolymers were defined as the DNA region spanning ±5 bases from the SNV (`--contextSpan 5`), with over 40% of identical nucleotides (--homopolymerFreq 0.4 `). Further, GIP discarded SNVs with sequencing coverage above or below 4 median absolute deviations (MADs) from the median chromosome coverage (`--MADrange`). snpEff (version 4.3t) (73) was used to predict the impact of SNVs on coding sequence. The predicted effects that GIP considered synonymous mutations are: "synonymous_variant", "stop_retained_variant" and "start_retained". The predicted effects that GIP considered non-synonymous mutations are: "missense_variant", "start_lost", "stop_gained", "stop_lost" and "coding_sequence_variant". The phylogenetic tree was computed by the giptools module `phylogeny` using IQtree2 (version 2.1.2) (74,75) with options `--seqtype DNA --alrt 1000 -B 1000`. The Venn-diagram comparison considered the strains QS0044_C, QS0001_C, QS0037_C, QS0016_C and SGH_358 that were sampled from different locations in Ethiopia, respectively Habala, Badowacho, Arbaminch, Hawassa and Jimma. The strains were selected to have comparable average genome coverage (23). To infer the tree GIP considered the set of filtered SNV and adopted the IUPAC ambiguous notation for the positions with allele frequency below 70%. The tree was visualized by giptools using the ggtree R-package (76).

Analysis of structural variants. GIP was used to detect structural variants including insertions, tandem duplications, deletions, inversions and break-end translocations with DELLY (version 0.8.7) (11). To reduce incorrect predictions the DELLY output was additionally filtered (`filterDellyOPT` option, **Supplementary Data 2**). GIP discarded poor predictions with

DELLY label "LowQual" (`--rmLowQual`) and low median MAPQ score of mapping reads (`-

-minMAPQ`). SVs positioned in proximity of chromosome ends were removed (--

chrEndFilter) to limit false predictions caused by potential misassembled regions close to the

telomeric ends. To ease visualization and limit the analysis only to best supported SVs GIP

limited the output only to the top predictions (`--topHqPercentIns`, `--topHqPercentDel`, `--

topHqPercentDup` and `--topHqPercentInv`) based on the SV support score as in **Formula 1**,

where *DV*, *DR*, *RV*, and *RR* are respectively the number of high-quality variant pairs, reference

pairs, variant junction reads, and reference junction reads.

$$\frac{DV + RV}{DV + RV + DR + RR} * 100$$

569   **Formula 1:** SV support score.

570   The predicted structural variants were represented with Circos (version 0.69-9) (77).

571   **Data and materials availability.** All data is available in the main text or the supplementary

572   materials.

Acknowledgments:

Funding: This study was supported by a seeding grant from the Institut Pasteur International

Department to the LeiSHield Consortium and the EU H2020 project LeiSHield-MATI - REP-

778298-1.

Author contributions: Conceptualization, G.B. and G.F.S.; Methodology, G.B. and G.F.S.;

Software, G.B.; Formal Analysis, G.B., and G.F.S.; Investigation, G.B. and G.F.S.; Writing –

Original Draft, G.B. and G.F.S.; Writing – Review & Editing, G.B. and G.F.S.; Supervision,

G.B. and G.F.S.; Funding Acquisition, G.F.S. and G.B.

Competing interests: Authors declare no competing interests.

Supplementary Tables:

Table S1: Genomic bin coverage ratio of sample LIPA83 over ZK43.

Table S2: Gene coverage ratio of sample LIPA83 over ZK43.

Table S3: Sample information.

Table S4: *Candida albicans* gene CNVs.

Table S5: Gene correlation clusters in *Candida albicans*.

Table S6: Network analysis of gene CNVs in *Candida albicans*.

Table S7: Gene ontology and metabolic pathway enrichment analyses.

Footnote: Terms and pathways with Benjamini – Hochberg adjusted p-values < 0.05 are

labeled in red.

Table S8: Mapping statistics.

Supplementary Data:

597 **Supplementary Data 1:** Cancer cell lines SNVs.

598 **Supplementary Data 2:** GIP configuration files.

599 **Supplementary Data 3:** giptools commands.

References:

1. Muir, P., Li, S., Lou, S., Wang, D., Spakowicz, D.J., Salichos, L., Zhang, J., Weinstock,
G.M., Isaacs, F., Rozowsky, J. *et al.* (2016) The real cost of sequencing: scaling
computation to keep pace with data generation. *Genome Biol*, **17**, 53.
2. Pareek, C.S., Smoczynski, R. and Tretyn, A. (2011) Sequencing technologies and
genome sequencing. *J Appl Genet*, **52**, 413-435.
3. Chin, C.S., Alexander, D.H., Marks, P., Klammer, A.A., Drake, J., Heiner, C., Clum,
A., Copeland, A., Huddleston, J., Eichler, E.E. *et al.* (2013) Nonhybrid, finished

microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods*, **10**,
563-569.

4. Chin, C.S., Peluso, P., Sedlazeck, F.J., Nattestad, M., Concepcion, G.T., Clum, A.,
612 Dunn, C., O'Malley, R., Figueroa-Balderas, R., Morales-Cruz, A. *et al.* (2016) Phased
613 diploid genome assembly with single-molecule real-time sequencing. *Nat Methods*, **13**,
614 1050-1054.

5. Reuter, J.A., Spacek, D.V. and Snyder, M.P. (2015) High-throughput sequencing
616 technologies. *Mol Cell*, **58**, 586-597.

6. Aurrecoechea, C., Barreto, A., Basenko, E.Y., Brestelli, J., Brunk, B.P., Cade, S.,
618 Crouch, K., Doherty, R., Falke, D., Fischer, S. *et al.* (2017) EuPathDB: the eukaryotic
619 pathogen genomics database resource. *Nucleic Acids Res*, **45**, D581-D591.

7. Dolled-Filhart, M.P., Lee, M., Jr., Ou-Yang, C.W., Haraksingh, R.R. and Lin, J.C.
621 (2013) Computational and bioinformatics frameworks for next-generation whole exome
622 and genome sequencing. *ScientificWorldJournal*, **2013**, 730210.

8. Pabinger, S., Dander, A., Fischer, M., Snajder, R., Sperk, M., Efremova, M., Krabichler,
624 B., Speicher, M.R., Zschocke, J. and Trajanoski, Z. (2014) A survey of tools for variant
625 analysis of next-generation genome sequencing data. *Brief Bioinform*, **15**, 256-278.

9. Garrison, E. and Marth, G. (2012), *arXiv preprint arXiv:1207.3907 [q-bio.GN]*.

10. Abyzov, A., Urban, A.E., Snyder, M. and Gerstein, M. (2011) CNVnator: an approach
628 to discover, genotype, and characterize typical and atypical CNVs from family and
629 population genome sequencing. *Genome Res*, **21**, 974-984.

11. Rausch, T., Zichner, T., Schlattl, A., Stütz, A.M., Benes, V. and Korbel, J.O. (2012),
631 *Bioinformatics*, Vol. 28, pp. i333-i339.

12. Hwang, K.B., Lee, I.H., Li, H., Won, D.G., Hernandez-Ferrer, C., Negron, J.A. and
633 Kong, S.W. (2019) Comparative analysis of whole-genome sequencing pipelines to
634 minimize false negative findings. *Sci Rep*, **9**, 3219.

13. Di Tommaso, P., Chatzou, M., Floden, E.W., Barja, P.P., Palumbo, E. and Notredame,
636 C. (2017) Nextflow enables reproducible computational workflows. *Nat Biotechnol*, **35**,
637 316-319.

14. Bussotti, G., Benkahla, A., Jeddi, F., Souiai, O., Aoun, K., Spath, G.F. and Bouratbine,
639 A. (2020) Nuclear and mitochondrial genome sequencing of North-African Leishmania
640 infantum isolates from cured and relapsed visceral leishmaniasis patients reveals
641 variations correlating with geography and phenotype. *Microb Genom*, **6**.

15. Bussotti, G., Gouzelou, E., Cortes Boite, M., Kherachi, I., Harrat, Z., Eddaikra, N.,
643 Mottram, J.C., Antoniou, M., Christodoulou, V., Bali, A. *et al.* (2018) Leishmania
644 Genome Dynamics during Environmental Adaptation Reveal Strain-Specific
645 Differences in Gene Copy Number Variation, Karyotype Instability, and Telomeric
646 Amplification. *MBio*, **9**.

16. Prieto Barja, P., Pescher, P., bussotti, g., Dumetz, F., Imamura, H., Kedra, D.,
648 Domagalska, M., Chaumeau, V., Himmelbauer, H., Pages, M. *et al.* (2017), *Nat Ecol*
649 *Evol*, Vol. 1, pp. 1961-1969.

17. Alvar, J., Velez, I.D., Bern, C., Herrero, M., Desjeux, P., Cano, J., Jannin, J., den Boer,
651 M. and Team, W.H.O.L.C. (2012) Leishmaniasis worldwide and global estimates of its
652 incidence. *PLoS One*, **7**, e35671.

18. Organization, W.H. (2018), WHO, Geneva. .

19. Price, R.N., Tjitra, E., Guerra, C.A., Yeung, S., White, N.J. and Anstey, N.M. (2007)
655 Vivax malaria: neglected and not benign. *Am J Trop Med Hyg*, **77**, 79-87.

20. Gething, P.W., Elyazar, I.R., Moyes, C.L., Smith, D.L., Battle, K.E., Guerra, C.A., Patil,
A.P., Tatem, A.J., Howes, R.E., Myers, M.F. *et al.* (2012) A long neglected world
malaria map: Plasmodium vivax endemicity in 2010. *PLoS Negl Trop Dis*, **6**, e1814.

21. Battle, K.E., Gething, P.W., Elyazar, I.R., Moyes, C.L., Sinka, M.E., Howes, R.E.,
Guerra, C.A., Price, R.N., Baird, K.J. and Hay, S.I. (2012) The global public health
significance of Plasmodium vivax. *Adv Parasitol*, **80**, 1-111.
22. Pearson, R.D., Amato, R., Auburn, S., Miotto, O., Almagro-Garcia, J., Amaratunga, C.,
Suon, S., Mao, S., Noviyanti, R., Trimarsanto, H. *et al.* (2016) Genomic analysis of
local variation and recent evolution in Plasmodium vivax. *Nat Genet*, **48**, 959-964.
23. Ford, A., Kepple, D., Abagero, B.R., Connors, J., Pearson, R., Auburn, S., Getachew,
S., Ford, C., Gunalan, K., Miller, L.H. *et al.* (2020) Whole genome sequencing of
Plasmodium vivax isolates reveals frequent sequence and structural polymorphisms in
erythrocyte binding genes. *PLoS Negl Trop Dis*, **14**, e0008234.
24. Singh, V., Gupta, P. and Pande, V. (2014) Revisiting the multigene families:
Plasmodium var and vir genes. *J Vector Borne Dis*, **51**, 75-81.
25. Rayner, J.C., Tran, T.M., Corredor, V., Huber, C.S., Barnwell, J.W. and Galinski, M.R.
(2005) Dramatic difference in diversity between Plasmodium falciparum and
Plasmodium vivax reticulocyte binding-like genes. *Am J Trop Med Hyg*, **72**, 666-674.
26. Rahul, C.N., Shiva Krishna, K., Pawar, A.P., Bai, M., Kumar, V., Phadke, S. and
Rajesh, V. (2014) Genetic and structural characterization of PvSERA4: potential
implication as therapeutic target for Plasmodium vivax malaria. *J Biomol Struct Dyn*,
32, 580-590.
27. Rahul, C.N., Shiva Krishna, K., Meera, M., Phadke, S. and Rajesh, V. (2015)
Plasmodium vivax: N-terminal diversity in the blood stage SERA genes from Indian
isolates. *Blood Cells Mol Dis*, **55**, 30-35.
28. Luo, Z., Sullivan, S.A. and Carlton, J.M. (2015) The biology of Plasmodium vivax
explored through genomics. *Ann N Y Acad Sci*, **1342**, 53-61.
29. Lin, J.T., Patel, J.C., Kharabora, O., Sattabongkot, J., Muth, S., Ubalee, R., Schuster,
A.L., Rogers, W.O., Wongsrichanalai, C. and Juliano, J.J. (2013) Plasmodium vivax
isolates from Cambodia and Thailand show high genetic complexity and distinct
patterns of P. vivax multidrug resistance gene 1 (pvmdr1) polymorphisms. *Am J Trop*
Med Hyg, **88**, 1116-1123.
30. Gunalan, K., Niangaly, A., Thera, M.A., Doumbo, O.K. and Miller, L.H. (2018)
Plasmodium vivax Infections of Duffy-Negative Erythrocytes: Historically Undetected
or a Recent Adaptation? *Trends Parasitol*, **34**, 420-429.
31. Costa, G.L., Amaral, L.C., Fontes, C.J.F., Carvalho, L.H., de Brito, C.F.A. and de
Sousa, T.N. (2017) Assessment of copy number variation in genes related to drug
resistance in Plasmodium vivax and Plasmodium falciparum isolates from the Brazilian
Amazon and a systematic review of the literature. *Malar J*, **16**, 152.
32. Cornejo, O.E., Fisher, D. and Escalante, A.A. (2014) Genome-wide patterns of genetic
polymorphism and signatures of selection in Plasmodium vivax. *Genome Biol Evol*, **7**,
106-119.
33. Chen, E., Salinas, N.D., Huang, Y., Ntumngia, F., Plasencia, M.D., Gross, M.L.,
Adams, J.H. and Tolia, N.H. (2016) Broadly neutralizing epitopes in the Plasmodium
vivax vaccine candidate Duffy Binding Protein. *Proc Natl Acad Sci U S A*, **113**, 6277-
6282.
34. Zolan, M.E. (1995) Chromosome-length polymorphism in fungi. *Microbiol Rev*, **59**,
686-698.
35. Suzuki, T., Nishibayashi, S., Kuroiwa, T., Kanbe, T. and Tanaka, K. (1982) Variance
of ploidy in Candida albicans. *J Bacteriol*, **152**, 893-896.
36. Sionov, E., Chang, Y.C. and Kwon-Chung, K.J. (2013) Azole heteroresistance in
Cryptococcus neoformans: emergence of resistant clones with chromosomal disomy in

the mouse brain during fluconazole treatment. *Antimicrob Agents Chemother*, **57**, 5127-
5130.

37.	Shin, J.H., Chae, M.J., Song, J.W., Jung, S.I., Cho, D., Kee, S.J., Kim, S.H., Shin, M.G.,
Suh, S.P. and Ryang, D.W. (2007) Changes in karyotype and azole susceptibility of
sequential bloodstream isolates from patients with Candida glabrata candidemia. *J Clin*
Microbiol, **45**, 2385-2391.

38.	Selmecki, A., Forche, A. and Berman, J. (2010) Genomic plasticity of the human fungal
pathogen Candida albicans. *Eukaryot Cell*, **9**, 991-1008.

39.	Magee, B.B. and Magee, P.T. (2000) Induction of mating in Candida albicans by
construction of MTLa and MTLalpha strains. *Science*, **289**, 310-313.

40.	Gerstein, A.C., Fu, M.S., Mukaremera, L., Li, Z., Ormerod, K.L., Fraser, J.A., Berman,
J. and Nielsen, K. (2015) Polyploid titan cells produce haploid and aneuploid progeny
to promote stress adaptation. *mBio*, **6**, e01340-01315.

41.	Croll, D. and McDonald, B.A. (2012) The accessory genome as a cradle for adaptive
evolution in pathogens. *PLoS Pathog*, **8**, e1002608.

42.	Chibana, H., Beckerman, J.L. and Magee, P.T. (2000) Fine-resolution physical mapping
of genomic diversity in Candida albicans. *Genome Res*, **10**, 1865-1877.

43.	Bravo Ruiz, G., Ross, Z.K., Holmes, E., Schelenz, S., Gow, N.A.R. and Lorenz, A.
(2019) Rapid and extensive karyotype diversification in haploid clinical Candida auris
isolates. *Curr Genet*, **65**, 1217-1228.

44.	Brown, G.D. and Netea, M.G. (2012) Exciting developments in the immunology of
fungal infections. *Cell Host Microbe*, **11**, 422-424.

45.	Pfaller, M.A., Diekema, D.J., Turnidge, J.D., Castanheira, M. and Jones, R.N. (2019)
Twenty Years of the SENTRY Antifungal Surveillance Program: Results for Candida
Species From 1997-2016. *Open Forum Infect Dis*, **6**, S79-S94.

46.	Pfaller, M.A., Castanheira, M., Messer, S.A., Moet, G.J. and Jones, R.N. (2010)
Variation in Candida spp. distribution and antifungal resistance rates among
bloodstream infection isolates by patient age: report from the SENTRY Antimicrobial
Surveillance Program (2008-2009). *Diagn Microbiol Infect Dis*, **68**, 278-283.

47.	Todd, R.T. and Selmecki, A. (2020) Expandable and reversible copy number
amplification drives rapid adaptation to antifungal drugs. *Elife*, **9**.

48.	Mount, H.O., Revie, N.M., Todd, R.T., Anstett, K., Collins, C., Costanzo, M., Boone,
C., Robbins, N., Selmecki, A. and Cowen, L.E. (2018) Global analysis of genetic
circuitry and adaptive mechanisms enabling resistance to the azole antifungal drugs.
PLoS Genet, **14**, e1007319.

49.	Hirakawa, M.P., Martinez, D.A., Sakthikumar, S., Anderson, M.Z., Berlin, A., Gujja,
S., Zeng, Q., Zisson, E., Wang, J.M., Greenberg, J.M. *et al.* (2015) Genetic and
phenotypic intra-species variation in Candida albicans. *Genome Res*, **25**, 413-425.

50.	Dixon, J.R., Xu, J., Dileep, V., Zhan, Y., Song, F., Le, V.T., Yardimci, G.G.,
Chakraborty, A., Bann, D.V., Wang, Y. *et al.* (2018) Integrative detection and analysis
of structural variation in cancer genomes. *Nat Genet*, **50**, 1388-1398.

51.	Yao, Y. and Dai, W. (2014) Genomic Instability and Cancer. *J Carcinog Mutagen*, **5**.

52.	Giovanni Bussotti, L.P., Pascale Pescher, Malgorzata Anna Domagalska, K.
Shanmugha Rajan, Tirza Doniger, Disha Gajanan Hiregange, Peter J Myler, Ron Unger,
Shulamit J Michaeli, Gerald F. Spaeth. (2021) Genome instability drives epistatic
adaptation in the human pathogen Leishmania. *bioRxiv*

53.	Abbey, D.A., Funt, J., Lurie-Weinberger, M.N., Thompson, D.A., Regev, A., Myers,
C.L. and Berman, J. (2014) YMAP: a pipeline for visualization of copy number
variation and loss of heterozygosity in eukaryotic pathogens. *Genome Med*, **6**, 100.

54. Bogaerts, B., Delcourt, T., Soetaert, K., Boarbi, S., Ceyssens, P.J., Winand, R., Van
Braekel, J., De Keersmaecker, S.C.J., Roosens, N.H.C., Marchal, K. *et al.* (2021) A
bioinformatics WGS workflow for clinical Mycobacterium tuberculosis complex isolate
analysis, validated using a reference collection extensively characterized with
conventional methods and in silico approaches. *J Clin Microbiol*.

55. Bogaerts, B., Winand, R., Fu, Q., Van Braekel, J., Ceyssens, P.J., Mattheus, W.,
Bertrand, S., De Keersmaecker, S.C.J., Roosens, N.H.C. and Vanneste, K. (2019)
Validation of a Bioinformatics Workflow for Routine Analysis of Whole-Genome
Sequencing Data and Related Challenges for Pathogen Typing in a European National
Reference Center: Neisseria meningitidis as a Proof-of-Concept. *Front Microbiol*, **10**,
362.

56. Ellison, M.A., Walker, J.L., Ropp, P.J., Durrant, J.D. and Arndt, K.M. (2020)
MutantHuntWGS: A Pipeline for Identifying Saccharomyces cerevisiae Mutations. *G3*
(Bethesda), **10**, 3009-3014.

57. Quijada, N.M., Rodriguez-Lazaro, D., Eiros, J.M. and Hernandez, M. (2019) TORMES:
an automated pipeline for whole bacterial genome analysis. *Bioinformatics*, **35**, 4207-
4212.

58. Leinonen, R., Sugawara, H., Shumway, M. and Collaboration, I.N.S.D. (2011), *Nucleic*
Acids Res., Vol. 39, pp. D19-21.

59. Leinonen, R., Akhtar, R., Birney, E., Bower, L., Cerdeno-Tarraga, A., Cheng, Y.,
Cleland, I., Faruque, N., Goodgame, N., Gibson, R. *et al.* (2011) The European
Nucleotide Archive. *Nucleic Acids Res*, **39**, D28-31.

60. Sloan, C.A., Chan, E.T., Davidson, J.M., Malladi, V.S., Strattan, J.S., Hitz, B.C.,
Gabdank, I., Narayanan, A.K., Ho, M., Lee, B.T. *et al.* (2016) ENCODE data at the
ENCODE portal. *Nucleic Acids Res*, **44**, D726-732.

61. Howe, K.L., Contreras-Moreira, B., De Silva, N., Maslen, G., Akanni, W., Allen, J.,
Alvarez-Jarreta, J., Barba, M., Bolser, D.M., Cambell, L. *et al.* (2020) Ensembl
Genomes 2020-enabling non-vertebrate genomic research. *Nucleic Acids Res*, **48**,
D689-D695.

62. Arnaud, M.B., Costanzo, M.C., Skrzypek, M.S., Binkley, G., Lane, C., Miyasato, S.R.
and Sherlock, G. (2005) The Candida Genome Database (CGD), a community resource
for Candida albicans gene and protein information. *Nucleic Acids Res*, **33**, D358-363.

63. Bahl, A., Brunk, B., Crabtree, J., Fraunholz, M.J., Gajria, B., Grant, G.R., Ginsburg, H.,
Gupta, D., Kissinger, J.C., Labo, P. *et al.* (2003) PlasmoDB: the Plasmodium genome
resource. A database integrating experimental and computational data. *Nucleic Acids*
Res, **31**, 212-215.

64. Girgis, H.Z. (2015) Red: an intelligent, rapid, accurate tool for detecting repeats de-
novo on the genomic scale. *BMC Bioinformatics*, **16**, 227.

65. Li, H. (2013), *arXiv:1303.3997v1 [q-bio.GN]*.

66. Li, H. and Durbin, R. (2009), *Bioinformatics*. 2009/05/20 ed, Vol. 25, pp. 1754-1760.

67. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
Abecasis, G., Durbin, R. and Subgroup, G.P.D.P. (2009), *Bioinformatics*, Vol. 25, pp.
2078-2079.

68. Li, W. and Godzik, A. (2006), *Bioinformatics*, Vol. 22, pp. 1658-1659.

69. Basenko, E.Y., Pulman, J.A., Shanmugasundram, A., Harb, O.S., Crouch, K., Starns,
D., Warrenfeltz, S., Aurrecoechea, C., Stoeckert, C.J., Jr., Kissinger, J.C. *et al.* (2018)
FungiDB: An Integrated Bioinformatic Resource for Fungi and Oomycetes. *J Fungi*
(Basel), **4**.

70. Kanehisa, M. and Goto, S. (2000) KEGG: kyoto encyclopedia of genes and genomes.
Nucleic Acids Res, **28**, 27-30.

71.   Caspi, R., Billington, R., Keseler, I.M., Kothari, A., Krummenacker, M., Midford, P.E.,
Ong, W.K., Paley, S., Subhraveti, P. and Karp, P.D. (2020) The MetaCyc database of
metabolic pathways and enzymes - a 2019 update. *Nucleic Acids Res*, **48**, D445-D453.

72.   Ramírez, F., Ryan, D.P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne,
S., Dündar, F. and Manke, T. (2016), *Nucleic Acids Res.*, Vol. 44, pp. W160-165.

73.   Cingolani, P., Platts, A., Wang le, L., Coon, M., Nguyen, T., Wang, L., Land, S.J., Lu,
X. and Ruden, D.M. (2012) A program for annotating and predicting the effects of
single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila
melanogaster strain w1118; iso-2; iso-3. *Fly (Austin)*, **6**, 80-92.

74.   Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von
Haeseler, A. and Lanfear, R. (2020) IQ-TREE 2: New Models and Efficient Methods
for Phylogenetic Inference in the Genomic Era. *Mol Biol Evol*, **37**, 1530-1534.

75.   Hoang, D.T., Chernomor, O., von Haeseler, A., Minh, B.Q. and Vinh, L.S. (2018)
UFBoot2: Improving the Ultrafast Bootstrap Approximation. *Mol Biol Evol*, **35**, 518-
522.

76.   Yu, G., Lam, T.T., Zhu, H. and Guan, Y. (2018) Two Methods for Mapping and
Visualizing Associated Data on Phylogeny Using Ggtree. *Mol Biol Evol*, **35**, 3041-
3043.

77.   Krzywinski, M., Schein, J., Birol, I., Connors, J., Gascoyne, R., Horsman, D., Jones,
S.J. and Marra, M.A. (2009), *Genome Res.*, Vol. 19, pp. 1639-1645.