The microtubule plus-end tracking protein Bik1 is required for chromosome congression

Alexander Julner [1], Marjan Abbasi [1], and Victoria Menéndez-Benito [1,2]

[1] Department of Biosciences and Nutrition, Karolinska Institutet, SE-141 83, Huddinge, Sweden.

[2] Correspondence: victoria.menendez.benito@ki.se

Abstract

During mitosis, sister chromatids congress on either side of the spindle equator to facilitate the correct partitioning of the genomic material. Chromosome congression requires a finely tuned control of microtubule dynamics by the kinesin motor proteins. In *Saccharomyces cerevisiae*, the kinesin proteins Cin8, Kip1, and Kip3 have pivotal roles in chromosome congression. It has been hypothesized that additional proteins that modulate microtubule dynamics are also involved. Here, we show that the microtubule plus-end tracking protein Bik1 - the budding yeast ortholog of CLIP-170 – is essential for chromosome congression. We find that nuclear Bik1 localizes to the kinetochores in a cell-cycle-dependent manner. Disrupting the nuclear pool of Bik1 with a nuclear export signal (Bik1-NES) leads to a slower cell cycle progression characterized by a delayed metaphase-anaphase transition. Bik1-NES cells have mispositioned kinetochores along the spindle in metaphase. Furthermore, using proximity-dependent methods, we identify Cin8 as an interaction partner of Bik1. Deleting CIN8 reduces the amount of Bik1 at the spindle. In contrast, Cin8 retains its typical bilobed distribution in Bik1-NES and does not localize to the unclustered kinetochores characteristic of Bik1-NES cells. Thus, we propose that Bik1 functions together with Cin8 to regulate kinetochore-microtubule dynamics for correct kinetochore positioning and chromosome congression.

26 **Keywords:** mitosis, kinetochore, chromosome, kinesins, genetic stability.

Introduction

Cell division relies on the accurate segregation of sister chromatids to generate functional daughter

cells. Before segregation, sister chromatids are positioned at the spindle equator through a process

known as chromosome congression that culminates with the metaphase plate formation in

metazoans (reviewed in (Maiato et al., 2017)). The establishment of a metaphase plate forces all the

chromosomes to segregate from the same starting position, thereby contributing to faithful

segregation. Despite the relevance of chromosome congression, its molecular mechanisms remain

unclear.

The budding yeast *Saccharomyces cerevisiae* is an appealing model organism to study spindle

dynamics because it is amenable to genetic engineering and has a relatively simple spindle. Budding

yeast has 16 chromosomes. After duplication, each sister chromatid is linked by a kinetochore to a

single kinetochore microtubule emanating from the spindle pole body (SPB), the yeast centrosome

(Winey & O'Toole, 2001). Each SPB also gives rise to four interpolar microtubules, with plus-ends that

overlap at the middle of the spindle (O'Toole et al., 1999; Peterson & Ris, 1976; Winey et al., 1995).

In prometaphase, kinetochores congress in two opposite clusters near the spindle equator,

equidistant from the SPBs. As in metazoan cells, this appears to occur prior to biorientation of sister

chromatids (Marco et al., 2013).

Several studies have shown that kinetochore microtubules are regulated in a length-dependent

manner (Gardner et al., 2005; Pearson et al., 2006; Sprague et al., 2003). Short microtubules tend to

grow, whereas long microtubules that extend towards the spindle equator are destabilized and

depolymerize. As sister chromatids are held together by cohesin, depolymerization of kinetochore

microtubules generates tension, mediated by the Ndc80 and Dam1 complexes' ability to track the

dynamic tips of the kinetochore microtubules (Powers *et al.*, 2009; Suzuki *et al.*, 2016).

Kinesin motors – proteins that move along microtubules powered by the hydrolysis of adenosine

triphosphate (ATP) – are involved in chromosome congression in several organisms (see Maiato *et*

al., 2017, Table 1). In budding yeast, the deletion of the kinesins Cin8, Kip3, and, to a lesser extent,

Kip1, results in declustering of kinetochores (Gardner et al., 2008; Marco et al., 2013; Tytell & Sorger,

2006; Wargacki et al., 2010). Cin8 and Kip3 have been shown to act as length-dependent microtubule

depolymerases, promoting catastrophe of long microtubules (Gardner et al., 2008; Su et al., 2011;

Varga et al., 2006).

It has long been speculated that other microtubule-associated proteins could play a role in

chromosome congression (Kops *et al.*, 2010). Specifically, microtubule plus-end tracking proteins

(+TIPs) are interesting candidates, because they accumulate at kinetochore microtubule plus-ends

and regulate their dynamics (reviewed in Akhmanova & Steinmetz, 2010). Bik1, the yeast ortholog of

CLIP-170, is a +TIP protein that exists in cytoplasmic and nuclear pools within the cell (Carvalho et al.,

2004). The cytoplasmic pool is important for mitotic spindle positioning through both the Kar9 and

the dynein pathways (reviewed in Miller *et al.*, 2006). First, Bik1 promotes phosphorylation of Kar9

for asymmetric localization at the old SPB (Moore et al., 2006). Kar9, together with the +TIP protein

Bim1 and the actin-guided myosin motor protein Myo2, mediates the capture of astral microtubules

at the bud cortex, contributing to spindle alignment and nuclear migration (Xiang, 2018). During

anaphase, Bik1 is transported by the kinesin Kip2 to the plus-ends of astral microtubules, where it

facilitates the recruitment of dynein (Carvalho *et al.*, 2004; Roberts *et al.*, 2014). Dynein is a minus

end-directed motor and will exert a pulling force on the spindle that results in nuclear migration to

the bud.

The nuclear pool of Bik1 is associated with the spindle at SPBs and kinetochores (Carvalho et al.,

2004; Lin et al., 2001) but its function is not well understood. Early studies found that deleting BIK1

results in cells with short or non-detectable astral microtubules (Berlin *et al.*, 1990). In contrast, cells

overexpressing Bik1 display short spindle microtubules and long astral microtubules. In this study,

we investigate the nuclear function of Bik1 and its role in chromosome congression and cell cycle

progression.

Results

Bik1 localizes to kinetochores in a cell cycle-dependent manner

We examined the localization of Bik1 in relation to tubulin, kinetochores and SPBs at different cell

cycle phases. We used cells expressing Bik1-superfolder GFP (sfGFP), mTurquoise-Tub1 (alpha

tubulin, microtubule marker), and either Ndc80-tagRFPT (kinetochore marker) or Spc42-tagRFPT (SPB

marker) (Figure 1A). In metaphase cells, nuclear Bik1 was primarily found between SPBs and

overlapping with kinetochores (Figure 1A, upper panels). By contrast, in anaphase cells, nuclear Bik1

was concentrated in patches along interpolar microtubules, while cytoplasmic Bik1 localized to astral

microtubules (Figure 1A, lower panels).

To confirm the metaphase association with kinetochores, we used cells expressing the APC/C co-

activator Cdc20 under control of the galactose inducible *GAL1* promoter. Shifting the carbon source

from galactose to glucose represses the expression of CDC20 and thereby arrests the cells at the

metaphase-to-anaphase transition. Bik1-sfGFP was also localized at kinetochore clusters in

metaphase-arrested cells (Figure 1B, upper panel), confirming what we observed in cycling cells. We

measured the distribution of Bik1-sfGFP in metaphase arrested cells and found that it had a bilobed

localization pattern like that of the kinetochore marker Ndc80-tagRFPT (Figure 1B, lower panel).

Bik1 in the cytoplasm localizes to astral microtubules and plays an important role in mitotic spindle

positioning (reviewed in Miller et al., 2006). To disrupt the kinetochore function of Bik1 while

maintaining its cytoplasmic function, we fused a strong nuclear export consensus sequence (NES)

(Kosugi et al., 2008) to the C-terminal end of Bik1-sfGFP or Bik1-5xFlag (collectively referred to as

Bik1-NES, Figure 1C). We imaged Bik1-NES cells expressing mTurquoise-Tub1, and Nc80-tagRFPT and

found that the mutant indeed excludes Bik1 from the nucleus (Figure 1D). To quantify the

effectiveness of the NES, we followed the localization of Bik1 in Bik1-sfGFP and Bik1-sfGFP-NES cells

expressing mTurquoise-Tub1. We synchronized the cells in G1 with α-factor, released them in fresh

media and imaged them at metaphase. As expected, Bik1-NES cells have significantly less sfGFP

signal at the spindle than Bik1 cells (Figure 1E-F).

We next assessed cell cycle progression in the absence of nuclear Bik1. We synchronized Bik1-sfGFP

and Bik1-sfGFP-NES cells in G1 with α-factor. Then, we released the cells from the G1 block and

prepared fixed samples at 10 min intervals. We scored the number of cells in G1/S, metaphase, or

anaphase according to the spindle length (mRuby2-Tub1) (Figure 2A). The strain expressing Bik1-

sfGFP had a sharp peak of metaphase cells at 40 min and a high number of cells with long spindles

50-60 min after the release. Bik1-sfGFP-NES cells, by contrast, displayed a broader peak of

metaphase cells that spans between 40-60 min post-release, and a 10 min delay in the anaphase

peak.

We then examined Bik1-sfGFP and Bik1-sfGFP-NES cells after G1 synchronization with α-factor by

time-lapse imaging (Figure 2B-E). In the Bik1-sfGFP strain, Bik1 was localized to the kinetochores

from G1 through metaphase. When the spindle elongates in anaphase, Bik1 was no longer associated

with the kinetochores and was found in patches along the interpolar microtubules instead (Figure

2B-C), as observed in fixed cells. The fluorescence intensity of Bik1-sfGFP at kinetochores was

greatest at metaphase when kinetochores cluster and reduced to near background levels in

anaphase (Figure 2D). In agreement with the analysis from fixed cells, we observed a slight delay in

cell cycle progression in Bik1-NES cells (Figure 2B, arrows). Our quantifications showed that the time

from the appearance of two kinetochore clusters to their segregation into mother and daughter cells

is approximately 10 min longer in Bik1-NES cells (Figure 2E).

Cells lacking nuclear Bik1 display aberrant kinetochore positioning

Because Bik1 localizes near the kinetochores in metaphase, we hypothesized that the prolonged

metaphase of Bik1-NES cells might be due to a dysregulation in kinetochore clustering. To test this

hypothesis, we analyzed the distribution of the kinetochore component Ndc80-sfGFP relative to the

SPBs (Spc42-tagRFPT) in cells expressing either Bik1-5xFlag or Bik1-5xFlag-NES arrested in metaphase

(PGal1-Cdc20) (Figure 3). While Bik1-5xFlag cells showed typical Ndc80 clusters, we observed cells

with long stretches of Ndc80 in the Bik1-5xFlag-NES mutant (Figures 3A-B). The unclustered

kinetochore phenotype was observed in 60% of the Bik1-NES metaphase cells (Figure 3C). Line scan

analysis of the Ndc80-sfGFP fluorescence relative to the SPBs showed that Bik1-NES cells have a

flattened distribution of Ndc80 with kinetochores mispositioned towards the spindle equator (Figure

3D). Overall, the defects in kinetochore-clustering suggest that some kinetochore-microtubules are

abnormally long in Bik1-NES at metaphase.

Bik1 interacts with Cin8 at the spindle

Kinesin motor proteins have been shown to play an essential role in chromosome congression. To

investigate the overall interaction of Bik1 with kinesins, we performed single-step affinity purification

of Bik1-Flag in metaphase arrested cells, followed by mass spectrometry analysis. Bik1-Flag co-

purified with the yeast kinesins Kip2, Cin8, Kip1, Kip3 and Kar3 (Table 1). Noticeably, the kinesin Cin8

appeared with a similar coverage as the known interactor Kip2 (Supplementary data S1).

The defect of Bik1-NES cells in kinetochore clustering shown in Figure 3A phenocopies CIN8 deletion

(Gardner et al., 2008; Tytell & Sorger, 2006; Wargacki et al., 2010). Therefore, we decided to study

the Bik1-Cin8 interaction in more detail. We investigated whether Cin8 and Bik1 interact *in vivo* using

proximity-dependent biotin identification (TurboID), a proximity labelling assay that can detect

transient or weak protein-protein interactions (Roux et al., 2012). We fused TurboID, a mutated

version of a bacterial biotin ligase BirA (Branon et al., 2018; Larochelle et al., 2019), to the C-terminus

of Bik1 (Figure 4A). We then Flag-tagged Cin8 or Kip2, a kinesin known to interact with Bik1 at astral

microtubules (Carvalho et al., 2004). Both Cin8 and Kip2 were recovered after streptavidin pulldown

when expressed in Bik1-TurboID cells, indicating that Bik1 and Cin8 are in proximity *in vivo* (Figure

4B).

We next looked at the interaction between Bik1 and Cin8 in individual cells by bimolecular

fluorescence complementation (BiFC) (Sung & Huh, 2007). Briefly, an N-terminal fragment of the

yellow fluorescent protein Venus was fused to the C-terminus of Bik1, and a complimentary C-

terminal Venus fragment was fused to the C-terminus of Cin8. We detected BiFC signal at the spindle,

predominantly in metaphase and anaphase cells (Figure 4 C-D). Interestingly, the BiFC signal in

metaphase cells appeared as 1-2 dots close to the ends of the spindle, while the signal in anaphase

cells appeared as a single dot around the spindle midzone (Figure 4 C).

We then investigated the dependence of Cin8 and Bik1 to localize to the spindle. In the Bik1-NES

mutant, Cin8 still forms two foci between SPBs and kinetochores (Figure 5A and B), even in cells with

unclustered kinetochores. This result was surprising as Cin8 has been shown to bind kinetochores

directly (Suzuki et al., 2018). Linescan analysis of Cin8-sfGFP localization in relation to SPBs showed

that Cin8 localization is unaltered in the absence of nuclear Bik1 (Figure 5B). Deletion of CIN8, on the

other hand, resulted in a significantly reduced Bik1 intensity at the spindle (Figure 5C). Overall, these

data suggest that a portion of nuclear Bik1 associates with and is dependent on Cin8 to bind the

spindle.

Discussion

Our results demonstrate that the +TIP protein Bik1 – the yeast homolog of CLIP-170 – is critical for

chromosome congression.

We have shown that nuclear Bik1 localizes to kinetochores from G1 to metaphase, when it presents a

bilobed distribution like the pattern of the kinesins Cin8, Kip1, and Kip3 (Gardner et al., 2008; Tytell &

Sorger, 2006; Wargacki et al., 2010). Similar to these kinesins, Bik1 stays at kinetochores until

anaphase, when it concentrates at interpolar microtubules.

We explored the functions of Bik1 at the mitotic spindle using a Bik1-NES mutant, which preserves

Bik1 localization at astral microtubules and its role in spindle positioning, revealing phenotypes that

would have been difficult to interpret in a deletion mutant. Bik1-NES cells have a slower cell cycle

progression, characterized by a longer metaphase. Others have shown that Bik1 is needed to

nucleate microtubules that capture kinetochores after centromere re-activation (Tanaka et al., 2005).

Thus, Bik1's function in kinetochore capture could contribute to the cell cycle delay we observe.

However, we do not detect unattached kinetochores in the Bik1-NES mutant. Instead, the

kinetochores are aligned in the spindle but unclustered in a high percentage (60%) of metaphase

cells. Therefore, we hypothesize that the defect in chromosome congression is the main cause of the

metaphase delay.

Previous studies have shown that the kinesins Cin8 and to a lesser extent Kip1 and Kip3 are required

for yeast chromosome congression (Gardner et al., 2008; Tytell & Sorger, 2006; Wargacki et al.,

2010). However, the deletion of each kinesin results in different types of clustering defects. In the

cin8Δ mutant, the kinetochores move closer to the spindle equator (Gardner et al., 2008; Tytell &

Sorger, 2006; Wargacki et al., 2010). The *kip1Δ* mutant has a similar, but weaker, phenotype

(Gardner et al., 2008; Tytell & Sorger, 2006). Conversely, *kip3Δ* mutant cells mislocalize the

kinetochores towards the poles (Wargacki et al., 2010). Ndc80 clusters in the Bik1-NES mutant

appear shifted towards the spindle equator (Figure 3D), as has been reported for cin8Δ, suggesting

they perform similar functions at the kinetochores.

We have also shown that Bik1 associates with Cin8 at microtubules *in vivo*, using proximity-

dependent assays, in metaphase and anaphase. Furthermore, we found that the spindle associated

Bik1 is severely decreased in *cin8Δ*. The reduced Bik1 recruitment to the spindle and kinetochores

explains why Bik1 cannot compensate for the lack of Cin8, which is known to cause severe defects in

assembling the mitotic spindle (Hoyt et al., 1992; Kotwaliwale et al., 2007). A previous study has

shown that Cin8 mediates chromosome congression by depolymerizing kinetochore microtubules in

a length-dependent manner (Gardner et al., 2008). Gardner and coworkers proposed that Cin8 binds

randomly to kinetochore microtubules and walks towards the plus-end to promote microtubule plus-

end disassembly directly. However, in Bik1-NES cells, Cin8 is insufficient to regulate chromosome

congression. Moreover, Cin8 does not localize to the unclustered kinetochores in Bik1-NES. Taken

together, these observations suggest that Bik1 is a binding partner of Cin8 that contributes to the

disassembly of microtubule plus-ends to facility chromosome congression at metaphase. There are at

least two other examples where plus-end directed kinesin motors transport Bik1 in the cells. Kip2

transports Bik1 to astral microtubule plus ends to ensure proper spindle alignment (Carvalho et al.,

2004; Caudron et al., 2008). Furthermore, it has been suggested that Bik1 forms a complex with the

kinesin Kip3 during mating to induce microtubule depolymerization for nuclear congression (Molk et

al., 2006).

An important, but unanswered, question is how Bik1 regulates the dynamics of kinetochore

microtubules plus-end at the molecular level. One possibility is that Bik1 directly promotes

kinetochore-microtubule depolymerization. *In vitro* microtubule polymerization assays have shown

that Bik1 inhibits microtubule polymerization (Blake-Hodek et al., 2010). In that *in vitro* experimental

setup, however, Bik1 did not bind the microtubule ends, and therefore the authors proposed that

Bik1 could sequester free tubulin subunits, thereby limiting the rate of microtubule growth.

Another possibility is that Bik1 increases the processivity of Cin8. Bik1 has been proposed to act as a

processivity factor for the kinesin Kip2 at astral microtubules (Hibbel et al., 2015). Likewise, Bik1

might be needed for Cin8 to reach the plus-ends of long kinetochore microtubules that extend

beyond bundled kinetochore microtubules. Cin8 is a bidirectional kinesin and has been shown to

preferentially move in a minus-end directed manner as single motors (Gerson-Gurwitz et al., 2011;

Pandey et al., 2021; Roostalu et al., 2011). Directional switching has been shown to occur when Cin8

motors are clustered (Pandey et al., 2021; Roostalu et al., 2011), and other kinesin-5 motors switch

directionality in response to steric crowding on MTs (Britto et al., 2016). It is possible that loss of Bik1

leads to ineffective plus-end directed movement of Cin8 on long kinetochore microtubules.

In summary, we propose that that Bik1 and Cin8 collaborate to depolymerize long kinetochore-

microtubules to promote kinetochore clustering.

Materials and Methods

Yeast strains and plasmids

All *S. cerevisiae* strains used in this study are derivatives of BY4741 and are listed in Table S1. The

epitope tagged alleles (sfGFP and tagRFPT), TurboID-3xmyc and BiFC were constructed at the

endogenous loci by standard PCR-based integration as described in (Longtine et al., 1998) and

confirmed by PCR, sequencing and microscopy. The plasmids used to generate the strains are listed

in Table S2. Strains with fluorescently tagged Tub1 were constructed as described in (Markus et al.,

2015). Briefly, the listed plasmids were linearized using BsaBI (New England Biolabs, R0537S). The

linearized plasmid was then integrated by at the endogenous TUB1 locus.

Growth conditions

Cells were grown at 30°C in YEP (1% yeast extract, 2% bacto-peptone) medium supplemented with

2% glucose (YEPD) unless otherwise stated. For G1 arrest, overnight cultures were diluted to OD_{600} =

0.1 in fresh media, grown for 2.5 hours and arrested by adding 3 µg/ml α-factor (Sigma, custom

peptide WHWLQLKPGQPMY) every 60 min for 120 min. Cells were released by washing away α-factor

with 1x culture volume of YEPD, resuspending in 1x culture volume of YEPD, and further grown at

30°C. For metaphase arrest, PGal1-Cdc20 cells were grown in YEP supplemented with 2% raffinose

and 2% galactose, and switched to YEPD for 120 min.

Fluorescence imaging

Cells were fixed in 4% formaldehyde on ice for 60 minutes, washed with PBS and stored in PBS at 4°C

until imaging. Cells were imaged in concanavalin A-coated 96 well glass bottom plates. Images were

acquired using a Nikon Ti microscope, equipped with a Zyla 4.2+ sCMOS camera and a 60x water

immersion objective (NA 1.2). Z-stack images (4-8 µm, 0.3 µm steps) were captured for fixed cells.

Images for BiFC and immunofluorescence were acquired using a Nikon Ti2 microscope, equipped

with a Prime BSI sCMOS camera and a 60x water immersion objective (NA 1.2).

Live-cell imaging

Cells were arrested in G1 as described. Arrested cells were released in low-fluorescence media (LFM)

containing 1.7 g/l yeast nitrogen base without ammonium sulfate, amino acids, folic acid and

riboflavin (MP-Biomedicals, 4030-512), and 1 g/l monosodium glutamate, supplemented with 2%

glucose and amino acids. Cells were left to settle on the bottom of concanavalin A-coated 96 well

glass bottom plates for 20-30 minutes inside the microscope incubator at 30°C before being imaged.

Images were captured every 5 min for 100-120 min.

Image analysis

Fluorescence intensity measurements were made from maximum intensity projections of image z-

stacks using FiJi (Schindelin et al., 2012). Integrated intensities were measured in a region of interest

drawn around spindles/kinetochores. Background correction was done by multiplying the mean

fluorescence of regions outside of the nucleus with the measured region's area and subtracting from

the measured integrated intensity.

Linescan analysis was performed by drawing a line between the centers of opposing SPB foci and

measuring the fluorescence intensities along the line. Background correction was done by

subtracting the mean fluorescence measured at the cytoplasm. Binned means were calculated for

each strain.

Immunofluorescence

PGal1-Cdc20 cells were arrested in metaphase and fixed by adding formaldehyde to 3.7% (vol/vol)

and incubating while shaking at 30°C for 30 min. The fixed cells were washed once with PBS and then

twice with SP buffer (0.1 M KPO4, 1.2 M sorbitol, pH 7.5). The cells were resuspended in 500 μl SP

buffer with 0.1% β-mercaptoethanol and 10 μl of 20 mg/ml Zymoylase-20T (Seikagaku). The cells

were incubated with soft rotation for 20 min at room temperature (RT). The resulting spheroplasts

were then washed twice in SP buffer and permeabilized in blocking solution (3% BSA 0.1% Triton X-

100 in PBS) for 30 min at RT. The spheroplasts were immunostained in suspension. First, cells were

incubated with 1:1000 mouse monoclonal anti-FLAG M2 antibody (Sigma, F1804) in blocking buffer

for 1 hour at RT. Cells were washed three times in PBS and then incubated with 1:100 donkey anti-

mouse Alexa Fluor 647 antibody (ThermoFisher, A-31571) for 45 min at RT. Cells were washed three

times in PBS, seed in a glass-bottom imaging plate coated with Concanavalin A, and imaged.

Sample preparation for mass spectrometry analysis

Bik1-5xFlag PGal1-Cdc20 cells were arrested in metaphase. Briefly, cells were grown overnight in 50

ml YEP with 2% galactose and 2% raffinose. Cells were diluted in 2 L YEP with 2% and 2% raffinose

and grown at 30°C to OD_{600} = 1. Then, cells were spun down and resuspended in 2 L YEPD with 2%

glucose and arrested at 30°C for 2 h. Cells were spun down at 6000 rom at 4°C, washed with cold

dH_2O, and spun down at 6000 rpm at 4°C. The cell pellet was diluted in a few drops of lysis buffer (50

mM Tris, pH 8.0, 150 mM NaCl, 5% glycerol, 0.1% Triton X-100). The cell suspension was snap frozen

by pipetting the drops directly into liquid nitrogen. The frozen pellets were then lysed using a

Freezer/Mill (SPEX™ sample prep freezer/mill), using 10 rounds consisting of 2 min grinding at 14

rpm, 2 min cooling). The yeast pellets were diluted in buffer EB1 (40 mM sodium hepes, 300 mM

NaCl, 0.5% Triton X-100, 2 mM $MgCl_2$, 5% glycerol and protease and phosphatase inhibitors (1mM

PMSF, 1mM leupeptin, 1mM peptasin A, 10mM NaF, 1mM sodium pyrophosphatase, 1mm sodium

orthovanadate), and clarified by centrifugation at 10 000 g for 10 min at 4 °C. The cleared lysate was

incubated for 30 min with magnetic Dynabeads (Thermo Fisher Scientific) coupled with anti-FLAG M2

antibody (Sigma-Aldrich). The Dynabeads were washed three times with buffer A (25 mM sodium

Hepes, 2 mM $MgCl_2$, 0.1% EDTA, 0.1% Triton X-100, 5% glycerol, 150 mM NaCl, protease and

phosphatase inhibitors (1 mM PMSF, 1 mM leupeptin, 1 mM peptasin A, 10mM NaF, 1 mM sodium

pyrophosphatase, 1 mm sodium orthovanadate) and four times with buffer B (25 mM Hepes, pH 8.0,

and 150 mM KCl).

Protein on-beads digestion

Proteins bound to the Dynabeads were resuspended and washed with 500 μL of 50 mM ammonium

bicarbonate (AmBic), pH 8.0, twice. After incubation at room temperature for 15 min, 50 μL of 50

mM AmBic was added. Protein disulfide bonds were reduced with 5 µL of 200 mM dithiothreitol at

37°C for 45 min and alkylated with 5 µL of 600 mM iodoacetamide at room temperature in dark for

30 min. Tryptic digestion was started by adding 1 µg of trypsin and incubation at 37°C overnight,

followed by adding another 500 ng of trypsin and incubation at 30°C for 3 h. The digestion was

stopped with 5 µL concentrated formic acid. The samples were then cleaned on a C18 StageTip

(Thermo Scientific) and dried using a speedvac (MiVac, Thermo Scientific).

PRLC-MS/MS analysis

Chromatographic separations of peptides were performed on a 50 cm long C18 EASY-spray column

connected to an Easy nanoLC-1000 UPLC system (Thermo Fisher Scientific). The gradient was 4-25%

of solvent B (98% acetonitrile, 0.1% formic acid) in 50 min and 25-40%B in 10 min at a flow rate of

300 nL/min. An Orbitrap Fusion mass spectrometer (Thermo Scientific) analyzed the eluted peptides

that were ionized in a nano-Easy electrospray source. The survey mass spectrum was acquired at the

resolution of 120,000 in the range of m/z 350-1800. MS/MS data of precursors were obtained with

30% normalized collision energy by higher-energy collisional dissociation (HCD) for ions with charge

z=2-7 at a resolution of 30,000 in 3 s cycle time.

Data analysis

Mascot v 2.5.1 search engine (MatrixScience Ltd., UK) was used for protein identification with the

following search parameters: up to two missed cleavage sites for trypsin, peptide mass tolerance 10

ppm, 0.05 Da for the HCD fragment ions. Carbamidomethylation of cysteine was specified as a fixed

modification, whereas oxidation of methionine, deamidation of asparagine and glutamine were

defined as variable modifications.

Acknowledgements

We thank C. Björkegren (Karolinska Institutet, Sweden) for reagents, and C. Boone and Z. Li for help

with construction of deletion mutants and critical reading of this manuscript. Microscopy was

performed at the LCI facility/Nikon Center of Excellence, Karolinska Institutet, supported by grants

from the Knut and Alice Wallenberg Foundation, Swedish Research Council, KI infrastructure, Centre

for Innovative Medicine and Jonasson center at the Royal Institute of Technology. Protein

identification and quantification were carried out by the Proteomics Biomedicum core facility,

Karolinska Institutet. This work was supported by grants from the Swedish Council (VR-NT 2017-04536), and STINT (Mobility Grant for Internationalization, MG2019-8484) to V.M.B. The authors declare no competing financial interests.

References

Akhmanova, A., & Steinmetz, M. O. (2010). Microtubule +TIPs at a glance. In *Journal of Cell Science* (Vol. 123, Issue 20). https://doi.org/10.1242/jcs.062414

Berlin, V., Styles, C. A., & Fink, G. R. (1990). BIK1, a protein required for microtubule function during mating and mitosis in Saccharomyces cerevisiae, colocalizes with tubulin. *Journal of Cell Biology*, *111*(6 I). https://doi.org/10.1083/jcb.111.6.2573

Blake-Hodek, K. A., Cassimeris, L., & Huffaker, T. C. (2010). Regulation of microtubule dynamics by Bim1 and Bik1, the budding yeast members of the EB1 and CLIP-170 families of plus-end tracking proteins. *Molecular Biology of the Cell*, *21*(12). https://doi.org/10.1091/mbc.E10-02-0083

Branon, T. C., Bosch, J. A., Sanchez, A. D., Udeshi, N. D., Svinkina, T., Carr, S. A., Feldman, J. L., Perrimon, N., & Ting, A. Y. (2018). Efficient proximity labeling in living cells and organisms with TurboID. *Nature Biotechnology*, *36*(9). https://doi.org/10.1038/nbt.4201

Britto, M., Goulet, A., Rizvi, S., Von Loeffelholz, O., Moores, C. A., & Cross, R. A. (2016). Schizosaccharomyces pombe kinesin-5 switches direction using a steric blocking mechanism. *Proceedings of the National Academy of Sciences of the United States of America*, *113*(47). https://doi.org/10.1073/pnas.1611581113

Carvalho, P., Gupta, M. L., Hoyt, M. A., & Pellman, D. (2004). Cell cycle control of kinesin-mediated transport of Bik1 (CLIP-170) regulates microtubule stability and dynein activation. *Developmental Cell*, *6*(6). https://doi.org/10.1016/j.devcel.2004.05.001

Caudron, F., Andrieux, A., Job, D., & Boscheron, C. (2008). A new role for kinesin-directed transport of

Bik1p (CLIP-170) in Saccharomyces cerevisiae. *Journal of Cell Science*, *121*(9).

https://doi.org/10.1242/jcs.023374

Gardner, M. K., Bouck, D. C., Paliulis, L. V., Meehl, J. B., O'Toole, E. T., Haase, J., Soubry, A., Joglekar,

A. P., Winey, M., Salmon, E. D., Bloom, K., & Odde, D. J. (2008). Chromosome Congression by

Kinesin-5 Motor-Mediated Disassembly of Longer Kinetochore Microtubules. *Cell*, *135*(5).

https://doi.org/10.1016/j.cell.2008.09.046

Gardner, M. K., Pearson, C. G., Sprague, B. L., Zarzar, T. R., Bloom, K., Salmon, E. D., & Odde, D. J.

(2005). Tension-dependent Regulation of Microtubule Dynamics at Kinetochores Can Explain

Metaphase Congression in Yeast. *Molecular Biology of the Cell*, *16*(8), 3764–3775.

https://doi.org/10.1091/mbc.e05-04-0275

Gerson-Gurwitz, A., Thiede, C., Movshovich, N., Fridman, V., Podolskaya, M., Danieli, T., Lakämper, S.,

Klopfenstein, D. R., Schmidt, C. F., & Gheber, L. (2011). Directionality of individual kinesin-5 Cin8

motors is modulated by loop 8, ionic strength and microtubule geometry. *EMBO Journal*,

30(24). https://doi.org/10.1038/emboj.2011.403

Hibbel, A., Bogdanova, A., Mahamdeh, M., Jannasch, A., Storch, M., Schäffer, E., Liakopoulos, D., &

Howard, J. (2015). Kinesin Kip2 enhances microtubule growth in vitro through length-

dependent feedback on polymerization and catastrophe. *ELife*, *4*(NOVEMBER2015).

https://doi.org/10.7554/eLife.10542

Hoyt, M. A., He, L., Kek Khee Loo, & Saunders, W. S. (1992). Two Saccharomyces cerevisiae kinesin-

related gene products required for mitotic spindle assembly. *Journal of Cell Biology*, *118*(1).

https://doi.org/10.1083/jcb.118.1.109

Kops, G. J. P. L., Saurin, A. T., & Meraldi, P. (2010). Finding the middle ground: How kinetochores

power chromosome congression. In *Cellular and Molecular Life Sciences* (Vol. 67, Issue 13).

https://doi.org/10.1007/s00018-010-0321-y

Kosugi, S., Hasebe, M., Tomita, M., & Yanagawa, H. (2008). Nuclear export signal consensus

sequences defined using a localization-based yeast selection system. *Traffic*, *9*(12).

https://doi.org/10.1111/j.1600-0854.2008.00825.x

Kotwaliwale, C. V., Frei, S. B., Stern, B. M., & Biggins, S. (2007). A Pathway Containing the Ipl1/Aurora

Protein Kinase and the Spindle Midzone Protein Ase1 Regulates Yeast Spindle Assembly.

Developmental Cell, *13*(3). https://doi.org/10.1016/j.devcel.2007.07.003

Larochelle, M., Bergeron, D., Arcand, B., & Bachand, F. (2019). Proximity-dependent biotinylation

mediated by TurboID to identify protein-protein interaction networks in yeast. *Journal of Cell*

Science, *132*(11). https://doi.org/10.1242/jcs.232249

Lin, H., De Carvalho, P., Kho, D., Tai, C. Y., Pierre, P., Fink, G. R., & Pellman, D. (2001). Polyploids

require Bik1 for kinetochore-microtubule attachment. *Journal of Cell Biology*, *155*(7).

https://doi.org/10.1083/jcb.200108119

Longtine, M. S., McKenzie, A., Demarini, D. J., Shah, N. G., Wach, A., Brachat, A., Philippsen, P., &

Pringle, J. R. (1998). Additional modules for versatile and economical PCR-based gene deletion

and modification in Saccharomyces cerevisiae. *Yeast*, *14*(10).

https://doi.org/10.1002/(SICI)1097-0061(199807)14:10<953::AID-YEA293>3.0.CO;2-U

Maiato, H., Gomes, A. M., Sousa, F., & Barisic, M. (2017). Mechanisms of chromosome congression

during mitosis. In *Biology* (Vol. 6, Issue 1). https://doi.org/10.3390/biology6010013

Marco, E., Dorn, J. F., Hsu, P. H., Jaqaman, K., Sorger, P. K., & Danuser, G. (2013). S. cerevisiae

chromosomes biorient via gradual resolution of syntely between S phase and anaphase. *Cell*,

154(5). https://doi.org/10.1016/j.cell.2013.08.008

Markus, S. M., Omer, S., Baranowski, K., & Lee, W. L. (2015). Improved Plasmids for Fluorescent

Protein Tagging of Microtubules in Saccharomyces cerevisiae. *Traffic*, *16*(7).

https://doi.org/10.1111/tra.12276

Miller, R. K., D'Silva, S., Moore, J. K., & Goodson, H. V. (2006). The CLIP-170 Orthologue Bik1p and

Positioning the Mitotic Spindle in Yeast. In *Current Topics in Developmental Biology* (Vol. 76).

https://doi.org/10.1016/S0070-2153(06)76002-1

Molk, J. N., Salmon, E. D., & Bloom, K. (2006). Nuclear congression is driven by cytoplasmic

microtubule plus end interactions in S. cerevisiae. *Journal of Cell Biology*, *172*(1).

https://doi.org/10.1083/jcb.200510032

Moore, J. K., D'Silva, S., & Miller, R. K. (2006). The CLIP-170 homologue Bik1p promotes the

phosphorylation and asymmetric localization of Kar9p. *Molecular Biology of the Cell*, *17*(1).

https://doi.org/10.1091/mbc.E05-06-0565

O'Toole, E. T., Winey, M., & McIntosh, J. R. (1999). High-voltage electron tomography of spindle pole

bodies and early mitotic spindles in the yeast Saccharomyces cerevisiae. *Molecular Biology of

the Cell*, *10*(6). https://doi.org/10.1091/mbc.10.6.2017

Pandey, H., Reithmann, E., Goldstein-Levitin, A., Al-Bassam, J., Frey, E., & Gheber, L. (2021). Drag-

induced directionality switching of kinesin-5 Cin8 revealed by cluster-motility analysis. *Science

Advances*, *7*(6). https://doi.org/10.1126/sciadv.abc1687

Pearson, C. G., Gardner, M. K., Paliulis, L. V., Salmon, E. D., Odde, D. J., & Bloom, K. (2006). Measuring

nanometer scale gradients in spindle microtubule dynamics using model convolution

microscopy. *Molecular Biology of the Cell*, *17*(9). https://doi.org/10.1091/mbc.E06-04-0312

Peterson, J. B., & Ris, H. (1976). Electron microscopic study of the spindle and chromosome

movement in the yeast Saccharomyces cerevisiae. *Journal of Cell Science*, *22*(2).

Powers, A. F., Franck, A. D., Gestaut, D. R., Cooper, J., Gracyzk, B., Wei, R. R., Wordeman, L., Davis, T.

N., & Asbury, C. L. (2009). The Ndc80 Kinetochore Complex Forms Load-Bearing Attachments to

Dynamic Microtubule Tips via Biased Diffusion. *Cell*, *136*(5).

https://doi.org/10.1016/j.cell.2008.12.045

Roberts, A. J., Goodman, B. S., & Reck-Peterson, S. L. (2014). Reconstitution of dynein transport to

the microtubule plus end by kinesin. *ELife*, *3*. https://doi.org/10.7554/elife.02641

Roostalu, J., Hentrich, C., Bieling, P., Telley, I. A., Schiebel, E., & Surrey, T. (2011). Directional

switching of the kinesin Cin8 through motor coupling. *Science*, *332*(6025).

https://doi.org/10.1126/science.1199945

Roux, K. J., Kim, D. I., Raida, M., & Burke, B. (2012). A promiscuous biotin ligase fusion protein

identifies proximal and interacting proteins in mammalian cells. *Journal of Cell Biology*, *196*(6).

https://doi.org/10.1083/jcb.201112098

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden,

C., Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P., &

Cardona, A. (2012). Fiji: An open-source platform for biological-image analysis. In *Nature*

Methods (Vol. 9, Issue 7). https://doi.org/10.1038/nmeth.2019

Sprague, B. L., Pearson, C. G., Maddox, P. S., Bloom, K. S., Salmon, E. D., & Odde, D. J. (2003).

Mechanisms of microtubule-based kinetochore positioning in the yeast metaphase spindle.

Biophysical Journal, *84*(6). https://doi.org/10.1016/S0006-3495(03)75087-5

Su, X., Qiu, W., Gupta, M. L., Pereira-Leal, J. B., Reck-Peterson, S. L., & Pellman, D. (2011).

Mechanisms Underlying the Dual-Mode Regulation of Microtubule Dynamics by Kip3/Kinesin-8.

Molecular Cell, *43*(5). https://doi.org/10.1016/j.molcel.2011.06.027

Sung, M. K., & Huh, W. K. (2007). Biomolecular fluorescence complementation analysis system for in

vivo detection of protein-protein interaction in Saccharomyces cerevisiae. *Yeast*, *24*(9).

https://doi.org/10.1002/yea.1504

Suzuki, A., Badger, B. L., Haase, J., Ohashi, T., Erickson, H. P., Salmon, E. D., & Bloom, K. (2016). How

the kinetochore couples microtubule force and centromere stretch to move chromosomes.

Nature Cell Biology, *18*(4). https://doi.org/10.1038/ncb3323

Suzuki, A., Gupta, A., Long, S. K., Evans, R., Badger, B. L., Salmon, E. D., Biggins, S., & Bloom, K. (2018).

A Kinesin-5, Cin8, Recruits Protein Phosphatase 1 to Kinetochores and Regulates Chromosome

Segregation. *Current Biology*, *28*(17). https://doi.org/10.1016/j.cub.2018.08.038

Tanaka, K., Mukae, N., Dewar, H., Van Breugel, M., James, E. K., Prescott, A. R., Antony, C., & Tanaka,

T. U. (2005). Molecular mechanisms of kinetochore capture by spindle microtubules. *Nature*,

434(7036). https://doi.org/10.1038/nature03483

Tytell, J. D., & Sorger, P. K. (2006). Analysis of kinesin motor function at budding yeast kinetochores.

Journal of Cell Biology, *172*(6). https://doi.org/10.1083/jcb.200509101

Varga, V., Helenius, J., Tanaka, K., Hyman, A. A., Tanaka, T. U., & Howard, J. (2006). Yeast kinesin-8

depolymerizes microtubules in a length-dependent manner. *Nature Cell Biology*, *8*(9).

https://doi.org/10.1038/ncb1462

Wargacki, M. M., Tay, J. C., Muller, E. G., Asbury, C. L., & Davis, T. N. (2010). Kip3, the yeast kinesin-8,

is required for clustering of kinetochores at metaphase. *Cell Cycle*, *9*(13).

https://doi.org/10.4161/cc.9.13.12076

Winey, M., Mamay, C. L., O'Toole, E. T., Mastronarde, D. N., Giddings, T. H., McDonald, K. L., &

McIntosh, J. R. (1995). Three-dimensional ultrastructural analysis of the Saccharomyces

cerevisiae mitotic spindle. *Journal of Cell Biology*, *129*(6).

https://doi.org/10.1083/jcb.129.6.1601

Winey, M., & O'Toole, E. T. (2001). The spindle cycle in budding yeast. In *Nature Cell Biology* (Vol. 3,

Issue 1). https://doi.org/10.1038/35050663

Xiang, X. (2018). Nuclear movement in fungi. In *Seminars in Cell and Developmental Biology* (Vol. 82).

https://doi.org/10.1016/j.semcdb.2017.10.024

Figure Legends

Figure 1. Bik1 localizes to kinetochores in a cell-cycle dependent manner. (A) Representative images of Bik1-sfGFP (green) localization in cycling cells expressing mTurquoise-Tub1 (blue) and Spc42-tagRFP-T (red, top panel), and Nc80-tagRFP-T (red, lower panel). **(B)** Top panel: Representative images of Bik1-sfGFP (green) localization in pGal1-Cdc20 metaphase-arrested cells expressing Nc80-tagRFP-T. Lower panel: Line scan analysis shows the distribution of Bik1-sfGFP and kinetochores (Ndc80-tagRFP-T) in metaphase arrested cells. The fluorescence intensity was measured along lines drawn between the kinetochore foci and normalized relative to the maximum values, and mean values are shown (n=26 cells). **(C)** Cartoon showing the design of Bik1-NES. In short, a strong nuclear export signal (NES) was fused to Bik1-sfGFP or Bik1-5xFlag to promote nuclear export. **(D)** Representative images of Bik1-sfGFP-NES (green) localization in cycling cells expressing mTurquoise-Tub1 (blue) and Ndc80-tagRFPT (red). **(E)** Quantification of Bik1-sfGFP or Bik1-sfGFP-NES localization in metaphase cells. Bik1-sfGFP and Bik1-sfGFP-NES cells expressing mRuby2-Tub1 were arrested in G1 with α-factor and subsequently released in fresh media. Cells were fixed every 10 minutes for 120 minutes. Left: Representative cells at metaphase. Time after release = 40 min (Bik1-sfGFP), and 50 min (Bik1-sfGFP-NES). Right: Boxplots of the integrated fluorescence intensity of Bik1-sfGFP and Bik1-sfGFP-NES at the spindle (mRuby2-Tub1) in metaphase cells (n=55 cells). All images shown are maximum intensity projections of Z-stacks (9 μm, 1 μm steps). Dashed lines represent the cell outlines based on bright field images. Scale bars: 4 μm.

Figure 2. Bik1-NES delays spindle elongation at metaphase/anaphase transition. (A) Cells from (1E) were scored for cell cycle progression based on bud size and spindle length (mRuby-Tub1). Means of 3 independent experiments are shown (n=100 cells each). Error bars SD. **(B-E)** Time-lapse imaging of cells expressing Bik1-sfGFP or Bik1-sfGFP-NES (green) and Ndc80-tagRFP-T (red). Cells were arrested in G1 with α-factor, subsequently released in fresh medium under the microscope. Images were taken at 5 min intervals. **(B)** Representative time-lapse images. Arrowheads indicate the appearance

of separated kinetochore clusters. Arrows indicate segregated kinetochores in mother/bud. The time

(min) elapsed from appearance of separated kinetochore clusters is indicated by the numbers above

the frames. Images shown are maximum intensity projections of Z-stacks (9 μm, 1 μm steps). **(C)**

From the upper panel of Figure 2B, zoomed-in images of the kinetochores (Ndc80-tagRFP-T) at

metaphase and anaphase. **(D)** Boxplots of the integrated Bik1-sfGFP fluorescence intensity at

kinetochores (Ndc80-tagRFP-T) at metaphase (meta) and anaphase (ana) (n= 27 cells). **(E)**

Quantification of average time from the appearance of separated kinetochore clusters to segregated

kinetochores in mother/bud in Bik1-sfGFP and Bik1-sfGFP-NES cells from 2 independent experiments

(n = 57-59 cells). Error bars SD. p-value = 0.001691 using Welch Two Sample t-test.

Figure 3. Nuclear Bik1 is needed for kinetochore clustering at metaphase. (A) Representative

immunofluorescence staining for Bik1-5xFlag and Bik1-5xFlag-NES (magenta) in metaphase arrested

pGal1-Cdc20 cells expressing Spc42-sfGFP (red) and Ndc80-TagRFP-T (green). Images shown are

maximum intensity projections of Z-stacks (4.95 μm, 0.3 μm steps). **(B)** Representative images of

Bik1-5xFlag and Bik1-5xFlag-NES arrested as in (A), including zoomed-in images of the kinetochores

(Ndc80-sfGFP). Images shown are maximum intensity projections of Z-stacks (8.7 μm, 0.3 μm steps).

(C) Quantification of kinetochore (Ndc80-tagRFPT) unclustered phenotype in Bik1-5xFlag and Bik-

5xFlag-NES cells arrested in metaphase as shown in (B). Bar plots show means from 3 independent

experiments (n=100 cells each). Error bars, SD. p=0.0006884 using Welch Two Sample t-test **(D)** Line

scan analysis shows the distribution of kinetochores (Ndc80sfGFP) and spindle pole bodies (SPBs,

Spc42-tagRFP-T) in metaphase arrested cells. Fluorescence intensity was measured along lines drawn

between the SPBs (Spc42-TagRFP-T) and normalized relative to the maximum values, and mean

values are shown (n=100 cells). Scale bars: 4 μm.

Figure 4. Bik1 interacts with Cin8 at metaphase (A-B) Identification of Cin8 as a Bik1 interaction

partner using TurboID. Cartoon outlining the principle of the TurboID assay. Briefly, a biotin ligase

was fused to endogenous Bik1 (Bik1-TurboID) in cells expressing the protein of interest tagged with

Flag (Prey). Biotinylated substrates were purified using streptavidin beads, and inputs and pull-down

lysates were analyzed by immunoblot for the indicated proteins. **(B)** Western blot analysis of total

cell extracts (input) and streptavidin pull-downs prepared from Bik1-TurboID-3xmyc cells expressing

Kip2-5xFlag or Cin8-5xFlag as prey, and control strains, using anti-Myc (top) and anti-Flag (bottom)

antibodies. (*) unspecific band. **(C-D)** Bimolecular fluorescence complementation (BiFC) analysis of

Bik1 and Cin8 interaction. **(C)** Representative images of cycling cells expressing Bik1-VN and Cin8-VC

in G1, metaphase and anaphase. Images shown are single focal planes. **(D)** Proportion of cells in (C)

with BIFC signal based on Venus fluorescence (3 independent experiments, n=50 cells per

experiment and cell cycle stage). Error bars, SD.

Figure 5. Dependence of Cin8 and Bik1 to localize to the spindle (A-B) Cin8 localizes at the spindle in

Bik1-NES cells, but does not co-localize with unclustered kinetochores. **(A)** Cin8 localization at the

metaphase spindle does not require nuclear Bik1. Representative images of pGal1-Cdc20 cells

arrested in metaphase expressing Bik1-5xFlag or Bik1-5xFlag-NES, Cin8-sfGFP, and the markers

Ndc80-TagRFP-T (top panel) or Spc42-TagRFP-T (lower panel). Images shown are maximum intensity

projections of Z-stacks (8.7 μm, 0.3 μm steps). **(B)** Line scan analysis of Cin8 distribution (Cin8-sfGFP)

in metaphase arrested cells). Top panel: Cin8 distribution relative to the kinetochores (Ndc80-

tagRFP-T). The fluorescence intensity of Cin8-sfGFP and Ndc80-tagRFPT was measured along lines

drawn between the SPBs and normalized relative to the maximum values, mean values are shown

(n= 26 cells). Low panel: Cin8 distribution relative to the SPBs (Spc42-tagRFP-T). The fluorescence

intensity of Cin8-sfGFP and Spc42-tagRFPT was measured along lines drawn between the SPBs and

normalized relative to the maximum values. Mean values are shown (n= 26cells). **(C)** Cin8 is needed

for Bik1-sfGFP Bik1-sfGFP localization at kinetochores. CIN8 and cin8Δ cells expressing Bik1-sfGFP and

Ndc80-TagRFP-T were arrested in G1 using α-factor, released in fresh media at 25°C and fixed every

10 minutes for 130 minutes. Left: Representative images of cells in metaphase (CIN8, 60 min after

release; cin8Δ, 110 min after release) Images shown are maximum intensity projections of Z-stacks

(6.6 μm, 0.6 μm steps). Right: Boxplots of the integrated fluorescence intensity of Cin8-sfGFP at the

spindle (based on Ndc80-TagRFP-T localization) in metaphase cells (n=100 cells each). Scale bars: 4

μm.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Table 1. A subset of the protein interactors of Bik1-FLAG (See Data S1 for a complete list).

Name	% Coverage	Unique peptides
STU2	72.33	104
BIM1	67.00	21
CIN8	50.67	46
KIP2	49.33	28
KAR9	47.33	26
KIP1	38.67	28
KAR3	38.67	23
CIK1	22.67	10
VIK1	21.00	10
KIP3	15.33	6

Table S1. Strains used in this study. All strains are derived from BY4741

Strain	Genotype
VMB-212	MATα BIK1-sfGFP:Ura3 SPC42-tagRFP-T:KanMX Leu2:HIS3p:mTurquoise2-TUB1
VMB-154	MATα BIK1-sfGFP:Ura3 NDC80-tagRFP-T:KanMX Leu2:HIS3p:mTurquoise2-TUB1
VMB-185	MATa BIK1-sfGFP:Ura3 NDC80-tagRFP-T:KanMX
VMB-293	MATa NatN2:PGAL1-CDC20 BIK1-sfGFP:Ura3 NDC80-tagRFP-T:KanMX
VMB-156	MATα BIK1-sfGFP-NES:NatMX NDC80-tagRFP-T:KanMX Leu2:HIS3p:mTurquoise2-TUB1
VMB-166	MATa BIK1-sfGFP:Ura3 Leu2:HIS3p:mRuby2-TUB1
VMB-209	MATa BIK1-sfGFP-NES:NatMX Leu2:HIS3p:mRuby2-TUB1
VMB-455	MATa HygroMX:PGAL1-CDC20 BIK1-5xFlag-NES:NatMX NDC80-sfGFP:Ura3 SPC42-tagRFP-T:KanMX
VMB-458	MATa HygroMX:PGAL1-CDC20 BIK1-5xFlag:NatMX NDC80-sfGFP:Ura3 SPC42-tagRFP-T:KanMX
VMB-433	MATa KIP2-5xFlag:KanMX
VMB-422	MATa BIK1-TurboID-3xMyc:NatMX
VMB-431	MATa BIK1-TurboID-3xMyc:NatMX KIP2-5xFlag:KanMX
VMB-393	MATa CIN8-5xFlag:KanMX
VMB-428	MATa BIK1-TurboID-3xMyc:NatMX CIN8-5xFlag:KanMX
VMB-448	MATa HygroMX:PGAL1-CDC20 CIN8-sfGFP:Ura3 NDC80-tagRFP-T:KanMX
VMB-453	MATa HygroMX:PGAL1-CDC20 BIK1-5xFlag-NES:NatMX CIN8-sfGFP:Ura3 NDC80-tagRFP-T:KanMX
VMB-459	MATa HygroMX:PGAL1-CDC20 BIK1-5xFlag:NatMX CIN8-sfGFP:Ura3 SPC42-tagRFP-T:KanMX
VMB-457	MATa HygroMX:PGAL1-CDC20 BIK1-5xFlag-NES:NatMX CIN8-sfGFP:Ura3 SPC42-tagRFP-T:KanMX
VMB-308	MATa BIK1-sfGFP:Ura3 cin8Δ:natMX NDC80-tagRFP-T:KanMX
VMB-288	MATa NatN2:PGAL1-CDC20 BIK1-5xFlag:KanMX
VMB-494	MATa BIK1-VN173:His3MX CIN8-VC155:KanMX Leu2:HIS3p:mTurquoise2-TUB1

Table S2. Plasmids used in this study

Plasmid	Relevant insert	Source
p44873	pFA6a-link-yosuperfolderGFP-CaURA3	Gift from Wendell Lim & Kurt Thorn (Addgene: 44873) PMID: 23844123
p44906	pFA6a-link-yoTagRFP-T-Kan	Gift from Wendell Lim & Kurt Thorn (Addgene: 44906) PMID: 23844123
p50641	pHIS3p:mTurquoise2-Tub1+3'UTR::LEU2	Gift from Wei-Lih Lee (Addgene: 50641) PMID: 25711127
p50645	pHIS3p:mRuby2-Tub1+3'UTR::LEU2	Gift from Wei-Lih Lee (Addgene: 50645) PMID: 25711127
pYM-N23	ScGAL1 promoter::pAgTEF-nat-tScADH1	PCR-Toolbox Knop (Euroscarf) MID: 15334558
p15983	pFA6a-5xFlag-KanMX6	Gift from Eishi Noguchi (Addgene: 15983) PMID: 18729046
	pFA6a-5xFlag-NatMX6	This study (subclonned from p15983)
pFB1434	pFA6a-Turbo-ID-3Myc-NatMX6	Gift from François Bachand (Addgene: 126050) PMID: 31064814
P30555	pFA6a-VN-His3MX6	Gift from Sung MK & Huh WK. (Euroscarf) PMID: 17534848
P30560	pFA6a-VC-kanMX6	Gift from Sung MK & Huh WK. (Euroscarf) PMID: 17534848