Defining protein variant functions using high-complexity mutagenesis libraries and

enhanced mutant detection software ASMv1.0

Xiaoping Yang[1,9], Andrew L. Hong[2,5,6,9], Ted Sharpe[3,9], Andrew O. Giacomelli[2,7,9,10], Robert E.

Lintner[4,11], Douglas Alan[1], Thomas Green[1], Tikvah K. Hayes[2,7], Federica Piccioni[1,12], Briana

Fritchman[1], Hinako Kawabe[1], Edith Sawyer[1], Luke Sprenkle[1], Benjamin P. Lee[5,6], Nicole S.

Persky[1], Adam Brown[1], Heidi Greulich[2,7], Andrew J. Aguirre[2,7], Matthew Meyerson[2,7,8], William

C. Hahn[2,7], Cory M. Johannessen[2,13], and David E. Root[1,*]

Affiliations:

[1]Genetic Perturbation Platform, [2]Cancer Program, [3]Data Science Platform, and [4]Genomics

Platform, Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA

[5]Department of Pediatrics, Emory University, Atlanta, GA 30322, USA

[6]Aflac Center for Cancer and Blood Disorders, Children's Healthcare of Atlanta, Atlanta, GA

30322, USA

[7]Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA 02215, USA

[8]Department of Genetics, Harvard Medical School, Boston, MA 02115, USA

[9]These authors contributed equally

[10]Current Address: Ontario Cancer Princess Margaret Cancer Centre, 610 University Avenue,

Toronto, ON, M5G 2M9 Canada

[11]Current Address: RAN Biotechnologies, 100 Cummings Center, Ste 434J, Beverly, MA 01915,

USA

[12]Current address: Merck Research Laboratories, 33 Avenue Louis Pasteur, Boston, MA 02115,

USA

[13]Current address: Novartis Institutes for Biomedical Research, Oncology Innovative Targets and

Technologies, 250 Massachusetts Ave, Cambridge, MA 02139, USA

Correspondence should be addressed to D.E.R. (droot@broadinstitute.org)

Abstract

Saturated variant libraries that encode all single-amino-acid substitutions allow one to test all

variants of a gene simultaneously in a single experiment. Because a variant differs from its

reference at a single codon-position located anywhere along the coding sequence, variant

detection presents a profound technical challenge. A popular but expensive brute-force approach

is to divide a library into sub-libraries for direct sequencing. Here we present our approach that

features enhanced mutant-detection software that calls variants reliably from massive shotgun

sequencing data, and methodologies that enable one to design, construct and screen a single

saturated library per gene of any size.

Keywords: saturation mutagenesis, deep mutational scanning (DMS), variants, log-fold change,

base miscall, DNA polymerase error rate, next-generation sequencing (NGS).

Background

An open reading frame (ORF) variant library of a gene of interest provides a powerful

means of systematically characterizing the functions of over thousands of single-amino-acid

substitutions in a single experiment. The functional characterization of variants of an ORF has

historically been confined to alanine scans[1, 2], random mutagenesis by error-prone PCR[3] or

mutator *E. coli* strains[4], site-directed mutagenesis[5] , or transposon insertional

mutagenesis[6]. Advances in DNA oligonucleotide and gene synthesis[7] and in next-generation

sequencing[8] have accelerated our ability to assess all possible single-amino-acid changes for

every amino acid position of a protein. The resulting variant collections are known as saturation

mutagenesis libraries, or deep mutational scanning libraries. A well-executed screen of such a

library, that is, a pool of ORF variants, produces a comprehensive dictionary that maps variants

to phenotype strength.

As illustrated in **Figure 1**, a typical pooled screen allows one to test the functions of

thousands of genetic perturbation agents in a single experiment. Pooled screens rely on

introduction of one genetic perturbation cassette, e.g., an shRNA, a CRISPR sgRNA or an ORF

variant, per cell into the genome using delivery by a retrovirus. The perturbed cell population is

then subjected to selection pressure, for example, growth-over-time proliferation, treatment with

a drug, or flow-sorting for a reporter marker. Eventually, the collected cell population samples

are processed to measure the abundance of each library perturbation in each of the samples. The

perturbagen counts are the basis to assess the enrichment or depletion of each library

perturbation by comparing samples that have undergone selection and samples that have been

treated with reference conditions (e.g., pre-selection or vehicle treatment samples). In shRNA or

CRISPR screens, the functional elements, hairpins or guides respectively, serve as the perfect

barcodes, short and unambiguous, for screening sample deconvolution. In contrast, ORF variant

library screens are confronted with challenges in variant detection.

The complexity of a saturated ORF variant library is a function of the ORF length and the

number of desired variants per codon position. As illustrated in **Supplementary Figure 1**, at a

given amino acid position, the template amino acid is typically mutated to 19 other amino acids,

a stop codon, and when design rules allow, some synonymous codons (**Methods**). Factoring in

the amino acid positions to be mutagenized (N), the library is a collection of approximately 20*N

variants, or alleles, each differing from the template ORF at a single amino acid position. The

coordinates of all variations in a library are spread out along the full-length ORF. Thus,

capturing the variant-defining signature as subtle as a single codon variation located anywhere

along the full-length ORF sequence is one of the foremost technical challenges. Over the years,

admirable efforts and steady progress have been made in dividing a library into sub-libraries

each with mutational regions of initially 25[9, 10], then 47[11], 100[12], and recently 170 [13]

amino acids. These sub-libraries were then screened and sequenced separately. In some cases,

these sub-libraries were further divided into tiles to allow the direct next-generation sequencing

readout of variants[13]. The benefits of these approaches are limited to and deemed practical to

small ORFs only. Our methodology, on the other hand, emphasizes robust variant-calling tools

that can process massive shotgun NGS data to detect and accurately quantify variants, and

consequently allows one to screen a single saturated variant library for an ORF of any size (e.g.,

~600-bp *KRAS[14]*, ~1200-bp *TP53[15]* and *SMARCB1*, ~1400-bp region of *PDE3A[16]*,

~1700-bp *SHOC2[17]*, ~1800-bp *PPM1D[18]*, ~2200-bp *EZH2*, ~3600-bp *EGFR*) in a single

experiment (**Fig. 1b**). Here, we present our methods and strategies as we draw from our ever-

evolving knowledge from designing and constructing over 20 libraries and analyzing the

resulting screening data (**Supplementary Table 1**). Furthermore, we have made our library

designer, variant-calling software, and data analysis tools freely available (**Supplementary**

Table 2). We hope that this study will help lower the access barrier to this technology and pave a

path for sharing library resources within the scientific community.

Results

Overview of variant-detecting software ASMv1.0.

Despite differences in screening protocols, all ORF variant library screens converge at the quantification of library alleles in each of the screening samples (**Fig. 1b**). Variant quantification requires resolution of subtle differences among variants. Improving upon our earlier version of variant-calling software, we developed Analyze Saturation Mutagenesis v1.0 (ASMv1.0; see **Methods** and **Supplementary Fig. 2**), to more accurately process the NGS data and measure variant abundance.

ASMv1.0 is a part of the Genome Analysis Toolkit (GATK v4.2.0.0). For each screening sample, this program analyzes NGS reads aligned to the reference sequence that includes the ORF template sequence and flank sequences defined by the PCR primers used to amplify the full-length ORF from genomic DNA. It counts the number of observed molecules that have a set of mutations. ASMv1.0 observes this mutational landscape without regard to the library design. Because the NGS reads are typically far shorter than the full-length ORF and because each planned variant in the library carries a single codon change from the template ORF, most aligned reads will be wild-type, and serve only to indicate the depth of read coverage across a codon position. The relative abundance of reads that do contain mutations gives information on the identity and abundance of the variant defined by those mutations.

The ASMv1.0 variant-calling process is followed by a process that uses an R toolkit to parse ASMv1.0 output files of all screen samples into a single datafile (see **Methods**). Throughout these processes, the toolkit preserves the variant description, which unambiguously defines the called variant and which alone allows the resctruction of mutations in the variant. In

our experience, the detail-rich variant descriptions and the annotations derived from them are

invaluable datasets that allow us to perform a myriad of in-depth data analyses.

The enhanced performance of ASMv1.0.

The early version of variant detection software, ORFCallv1.0, was successfully used in

analyzing many saturation mutagenesis screens[15, 19-21]. ORFCallv1.0, however, tallies the

intended variants in the space of a single codon, and it ignores the association of an intended

codon change and any *in-cis* unintended changes. For a library that has prevalent unintended

errors associated with an intended change, ignoring the presence of these errors can mask the

true phenotype of the intended variant by the phenotypes caused by the unintended changes. As

an improvement, our new software, ASMv1.0, calls variants in the context of full-length read

pairs, allowing us to separate the variants that bear only the intended sequence change from those

that also harbor extra changes. Furthermore, in ASMv1.0, a pair of reads needs to pass through

sequential filters such as minimum base quality, minimum read length, and requirement of

consistency in overlapping sequences of a pair of reads, before being admitted for variant calling

(**Supplementary Fig. 2**). **Supplementary Table 3** summarizes the enhanced functionalities

from ORFCallv1.0 to ASMv1.0.

We used the screens performed in Giacomelli *et al*[15] to compare the two methods.

Specifically, the ectopically expressed *TP53* variants were screened in isogenic p53^{WT} and

p53^{NULL} A549 lung cancer cells, with selection pressure from either nutlin-3 or etoposide. Under

nutlin-3 treatment, the screen of p53^{NULL} cells enriched *TP53* loss-of-function (LOF) variants,

and depleted wild-type-like variants (**Fig. 2**). Under etoposide treatment, the p53^{NULL} cells

enriched wild-type-like *TP53* variants and depleted the *TP53* LOF variants (**Supplementary Fig.**

3). Under nutlin-3 treatment, the screen of p53^{WT} cells enriched *TP53* variants exhibiting

dominant-negative effects (DNE) (**Supplementary Fig. 4**). Using both the ORFCallv1.0 and

ASMv1.0 variant calling methods, we reprocessed the next-generation sequencing data from

these three screens. The phenotypic strength of each variant, measured as log-fold change (LFC)

in the pre- and post-treatment samples, produced similar heatmaps, indicating that the two

variant-calling algorithms are largely consistent with each other (**Fig. 2a, Supplementary Figs.**

3a, 4a). However, we also noted that several variants exhibited substantial differences in LFC

calculated using each algorithm; we define the differences as delta(LFC)=LFC^{ASM}-LFC^{ORFCall} in

Figure 2b, Supplementary Figure 3b and **4b.**

In a perfect library that consists solely of intended variants, ORFCallv1.0 and ASMv1.0

should be equal in performance. However, it is inevitable that some planned codon changes are

physically associated with one or more unintended nucleotide changes in the same molecule. We

define the 'variant purity' of a planned variant as the fraction of pure intended molecules over all

molecules that harbor the codon change of interest, whether or not they harbor additional

nucleotide changes. When the purity of a planned variant is high, as in a high-quality library,

both versions of the software will call variants with similar accuracy. However, when the variant

purity is low, the enrichment and depletion scores measured by ORFCallv1.0 can be misleading,

as they reflect the sum of effects of all species that happened to have a given codon change. We

computed the purity of every planned variant using the reference samples of the Giacomelli

screens, and for each purity bin we computed the Pearson correlation coefficient of two sets of

LFCs, one using ORFCallv1.0, the other using ASMv1.0 (panel **c** of **Fig. 2, Supplementary**

Figs. 3,4). The two methods correlate well when the variant purity is high and diverge as the

variant purity decreases.

We therefore reasoned that ASMv1.0 is an upgrade over ORFCallv1.0, particularly

considering how unintended *in cis* changes are handled. As a way of validating the higher

performance of ASMv1.0 over ORFCallv1.0, we compared the Giacomelli screen LFC scores

called by both versions of the software with the results from the Kotler *et al*[12] that used a small

variant library that focused on the p53 DNA binding domain (DBD) and screened in p53[NULL]

H1299 cells (**Fig. 2d**).

Kotler *et al* targeted p53 DBD with 4 libraries, each covering the variants of a 47-amino

acid region of the ORF. The Kotler screens were processed by direct sequencing readout of the

mutational region of each of the 4 libraries. Furthermore, Kotler *et al* processed their screen with

stringent analysis thresholds by requiring (1) at least 80bp overlap of the two reads in a read pair,

(2) perfect sequence agreement in the overlapping region, and (3) minimum read coverage

of >200 reads for each variant. These stringent thresholds and the direct readout of variant counts

rendered the Kotler dataset the best available for, independently, validating the ASMv1.0

algorithm.

Similar to the Giacomelli p53[NULL] A549/nutlin-3 screen, the Kotler p53[NULL] H1299 cell

screen enriched for *TP53* alleles that exhibit LOF, and depleted wild-type-like variants. From the

7890-variant Giacomelli p53[NULL] A549/nutlin-3 screen data, we extracted a 2964-variant subset

that covered the same mutational space as the Kotler H1299 screen. Among these 2964 variants,

we identified 76 variants from the Giacomelli p53[NULL]/nutlin-3 screen that were called

differently by ORFCallv1.0 and ASMv1.0, with abs(delta(LFC))>1 (**Fig. 2d, right**). Of these

differentially scored variants, the LFC scores called by two versions of the software were

individually compared with the Kotler screen scores (**Fig.1e**). The ASMv1.0 LFC scores for the

Giacomelli p53[NULL]A549/nutlin-3 screen agree with the Kotler calls better than the ORFCallv1.0

LFC scores do (**Fig. 2e**).

ASMv1.0-enabled analysis reveals that the secondary mutations (errors) in the library are

near the targeted mutation and introduced by errors in oligo synthesis.

The ability of ASMv1.0 to detect molecules with intended or unintended changes along

the entire read pair allows one to conduct in-depth analyses of each saturated ORF variant

library. Thus, in addition to the variant purity discussed earlier, we can also tally and characterize

the unintended nucleotide changes to profile the errors introduced during library construction.

For example, we were able to determine that in the *TP53* library, 84% of molecules are pure and

planned variants (**Fig. 3a left**), and the errors in the library are predominantly single nucleotide

changes.

Taking one step further with the ASMv1.0 output, we investigated the proximity

distribution of errors in relation to the intended codon change. In **Figure 3**, we compared

libraries made by two different methods (see **Methods**): the *TP53* library made by MITE

(Mutagenesis by Integrated TilEs)[21], and the *PDE3A* library synthesized by Twist Bioscience.

The MITE method involved oligo synthesis to cover the length of 'tiles' that are 90 nucleotides

long. The Twist Bioscience technology involved the synthesis of oligos that vary, depending on

GC content, between 30-50 nucleotides and with the variant-encoding bases in the middle. In

Figure 3b, we profiled the error rate by distance, in nucleotides, between the locations of the

error and the planned change. This distance profile yields an important observation that have

shaped our screen deconvolution sequencing strategy, which will be discussed in detail later in

this section:

The stretches of frequent errors in libraries produced by the two methods, ~90 bases for

the MITE method and ~30 bases for the Twist Bioscience method, correspond to the length of

chemically synthesized DNA oligonucleotides involved in library production. This indicates that

the errors in the libraries are concentrated near the planned codon changes and sequencing reads

as short as 100 bases can detect both intended and unintended changes in the library.

Furthermore, it reasons that we can focus on the reads that carry expected variant

signatures to tally variant counts. Without exception, each full-length ORF variant in the library

consists of nucleotides inherited from mutation-introducing DNA strands (e.g., primers) and

nucleotides synthesized by high-fidelity DNA polymerase. The errors in the library are dictated

predominantly by the former, as it has much higher error rate than DNA polymerase. A high-

fidelity DNA polymerase with an error rate of 10^{-6} errors per cycle per amplified nucleotide will

produce a small number of erroneous molecules (1.1% for a 1kb ORF; 4.3% for a 4kb ORF),

which amount to a small fraction of those introduced from oligo synthesis (10% or 18% between

two library production methods) (**Supplementary Table 4**). We therefore argue that by focusing

on reads that carry expected mutations, and with NGS reads of 100 bases or longer, we

effectively capture the errors in the library.

ASMv1.0-enabled analysis allows assessing the effect of miscalls (artifactual errors)

introduced by processes involving PCR.

While the errors discussed above are true errors that are present in the libraries, it is

important to address the artifactual errors in the form of miscalls introduced by the variant

quantification process itself. In the screen deconvolution process, PCR was used to replicate the

ORF DNA from genomic DNA (up to 30 PCR cycles) and used again in NGS sample

preparation (12 cycles). The errors introduced by these PCR steps are artifacts. We define

'miscalls' as the artifactual and mispresented base calls relative to what are in the samples. The

error rate of DNA polymerases can be as low as one error per one million bases synthesized in a

PCR cycle. When we, in parallel with the screening samples, processed and sequenced the

reference *TP53* ORFs that are clonal, pure, and without any variants, the sequencing results are

striking – at each nucleotide position, we observed miscalls at a rate of ~0.0003-0.0006 miscalls

per nucleotide read-out (**Fig. 4a, right**), or 3-6 miscalls per 10,000 sequenced bases.

While this miscall rate may appear small, it is a significant factor in variant detection. As

an example, illustrated in **Supplementary Figure 5**, these miscalls will lead to (1) inflated

counts of variants with codons that are one-nucleotide away from the template codon (we refer to

this variant type as '1-nt delta' codon), and (2) the artifactually narrowed log-fold change range

of 1-nt delta codons (**Supplementary Fig. 5,** legend**). Indeed, in the *TP53* library screen with

p53^{NULL} cells/nutlin-3, we observed the inflation of counts (**Fig. 4b**), and the suppression in fold-

change (**Fig. 4c**), in the 1-nt delta variant group relative to 2- or 3-nt delta variants. This effect of

PCR errors on 1-nt delta codon changes has been observed, without exception, in all libraries we

have processed, and the effect magnifies as ORF length increases.

A recent report[22] presented software DiMSum that used 1-nt delta codons to model

PCR/NGS errors as a method to assess library quality. We, however, took a direct approach and

opted to mitigate the effects of PCR/NGS miscall errors by implementing library design rules

and adding control experiments. First, we set an important rule for saturated ORF variant library

design - minimizing the use of codons that are one nucleotide away from the reference codons.

In our experience, to achieve 'saturated' amino acid coverage in a library, we cannot completely

avoid variant codons that are one nucleotide away from a reference codon. But we can minimize

the use of 1-nt delta codons, reducing them from 15% to 3% of the library members. Secondly,

dealing with these 3% 1-nt delta variants, we added a clonal template plasmid sample and

processed it as if it were a screening sample. One may opt to use the clonal template data to

monitor or even correct the screen sample data (**Fig. 4b**, right).

Silent variants are used as a measure of screen baseline.

Another rule we have implemented for library design is the inclusion of silent variants

with codons that are 2- or 3-nt different from the reference codon. ORF variant library screens

often produce many hits. As a result, one cannot always rely on the large portion of the library

producing no phenotype to serve as the baseline. It is therefore essential to include silent codon

changes in the library; such nucleotide changes allow us to identify and quantify the silent alleles

in parallel with missense and nonsense variants (see rows marked as 'B' in all long-fold-change

heatmaps). Often, codons encoding an amino acid differ from one another at the wobble base. As

a result, many silent codons are one nucleotide away from the template codon. Requiring that

any admitted silent variants have a 2- or 3-nt difference relative to the reference codon results in

silent variants comprising about 2% of the library.

In the *SMARCB1* library, we included nearly all possible silent variants, including 1-nt

delta silent variants. As expected, due to the artifactual miscalls discussed earlier, the abundance

of 1-nt delta silent variants is inflated relative to the abundance of 2- and 3-nt delta silent variants

(**Supplementary Fig. 6a**). For the same reason, the 1-nt delta silent variants show an

artifactually narrowed log-fold change range, compared with 2- and 3-nt delta groups

(**Supplementary Fig. 6b**). In our current library designer download (**Supplementary Table 2**),

we avoid 1-nt delta silent variants and minimize 1-nt delta missense and nonsense variants.

A justification for using next-generation sequencing (NGS) over long-read sequencing (LRS).

An often-debated topic is how saturation mutagenesis screen samples should be sequenced. It would seem evident that screen samples should be sequenced with long reads to widen the space of variant characterization and to capture all mutations present on each molecule. However, the technology that can capture the readout of full-length ORF sequences[23, 24] is currently cost-prohibitive for most labs, and more importantly, undesirable for variant detection due to low concordance rate of base calls[25]. For example, in a study by Wenger *et al*.[26], circular consensus sequencing (CCS), the best of its kind, has a concordance rate of 99.772%. This is equivalent to a Q26 Phred score, producing one discordance per 439-base readout. This discordance rate is higher than the error rate in commercial DNA oligonucleotide synthesis, the key contributor of the real errors in the libraries. Since the measurement discordance rate is higher than the underlying error rate which it is expected to measure, the CCS read attrition rate will be high. As a result, one may be forced to either throw out many reads that show secondary mutations or focus locally on the expected codon changes and ignore mutations in the distance.

It is however encouraging that the long-read sequencing technology is advancing at a steady and rapid pace, and its applications are actively explored. In the recent work by Frank *et al*.[27], long read sequencing was used to associate a set of unique external barcodes to each of ~7000 variants in a saturation variant library, so that the screen deconvolution work can be achieved by direct next-generation sequencing of the barcodes. However, for four prevailing

reasons, we favor a sequencing platform capable of producing NGS read pairs of 150 bases per

read with a Phred score of Q30 or more for each base call, which is comfortably achievable by

many NGS platforms: (1) We have demonstrated that the errors in our libraries are near the

planned codon changes (**Figure 3, right**), and a 150-base read is adequate to detect both the

variant-defining sequences and library errors from DNA oligonucleotide synthesis. (2) We have

shown that it is unnecessary to sequence an ORF variant far into its regions synthesized by high-

fidelity DNA polymerase during library construction, as the errors introduced by DNA

polymerase in regions outside of a 100-base window around the planned codon changes are rare

and negligible. Therefore, we rightfully focus on reads that bear a planned codon change, which

effectively represents the variant's error-harboring region, that is, the region that was chemically

synthesized. (3) We have demonstrated that PCR-introduced miscalls produce predominantly 1-

nt delta codon changes. As we minimize the use of 1-nt delta codon variants, the artifactual

variants resulting from the miscalls are mostly unintended variants. By ignoring reads carrying

only unintended variations, we largely removed the effects of sequencing miscalls. (4) ASMv1.0

allows users to set adequate base quality threshold, for example, at Phred score Q37 with

Illumina Novaseq NGS. With that, we can comfortably achieve <1 discordance per 5000-base

NGS readout. As we focus on the reads that carry an intended-variant signature, at this low

discordance rate, >94% (0.9998^{300}) of 150-base read pairs used to call variants are free of

miscalls.

Discussion

Variant detection software ASMv1.0 enabled a cascade of in-depth data analyses that

allowed us to explore, for instance, variant purity, the rate and nature of library errors, and the

artifactual miscalls. The results of these analyses have shaped and reshaped our strategies at

many key steps of the projects, on both laboratory and computational fronts. We believe that the

analytical and laboratory technical blueprint presented here, along with our library designer,

variant-calling software, and data analysis tools will help enable the scientific community to

fully utilize this technology in research applications.

It should be noted that the strong hits among the 1-nt delta variants may still score well,

even with the miscall-related log-fold change suppression. The issue of 1-nt delta codon variants

in processing screening samples stems from the fact that we sequence the ORF to quantify the

variants. The effect of miscalls on 1-nt delta variant counts becomes less of an issue for short

ORFs (e.g., *KRAS*). As illustrated in **Supplementary Figure 5**, the longer the ORF, for each

readout of a variant-defining codon, the more copies of the template codon need to undergo

PCR-based amplification and NGS, and consequently the more counts of the artifactual 1-nt

delta codons will result. Our suggestion of minimizing 1-nt delta codon variants in ORF variant

libraries is to insulate the true variant counts in the screening samples from PCR-introduced

artifactual miscalls. If one prefers 1-nt delta codons in the variant libraries, we recommend

adding the clonal template ORF as a control sample and processing it in parallel with the

screening samples. This control sample may be used as a count-correction factor (as shown in

Fig. 4b, right).

The recent establishment of an open-source platform, MaveDB, to enable variant

function data sharing, is a significant development[28]. We further advocate that the scientific

community find a way to share saturation mutagenesis reagents and resources. In many cases, the

content of a saturated ORF variant library is well defined by the gene reference sequence. For a

given gene, the library designed by one lab will be highly similar to one done by another lab. As

a perfect example of resources sharing, Boettcher *et al*[29], using the *TP53* saturated ORF

variant library created by Giacomelli *et al*[15], quickly demonstrated that the dominant-negative

activity of *TP53* missense mutations in DNA binding domain was selected for in myeloid

malignancies. If the scientific community can come to an agreement for library sharing and

repurposing, the available resources could be used to cover more genes, as opposed to building

redundant libraries. Imagine what a collection of saturated ORF variant libraries of 1000 disease-

related genes could do!

Conclusions

Saturated ORF variant libraries are powerful tools that enable one to exhaustively test all

possible protein variants simultaneously in a single experiment. Each experiment produces a

comprehensive dictionary of variants and functions. Yet, this technology has high access barriers

as it demands expertise across many disciplines, and the costs of producing and utilizing the

libraries are prohibitive to many academic labs. This report presents the scientific community a

full blueprint and a complete set of tools of conducting saturation mutagenesis projects. We are

certain that it lowers the technical barrier of this technology.

Methods

ORF expression vector. The vector we use to clone most of our saturated ORF variant libraries

is a lenti-viral vector pMT025 (**Supplementary Fig. 7**). It is suited for restriction/ligation

cloning using NheI/BamHI, or NheI/MluI restriction enzyme pairs. The vector has a puromycin

resistance selection marker driven by an SV40 promoter. After the cloning, the ORF expression

is driven by EF1a. To facilitate cloning, we require that the ORF sequences be properly flanked

with constant sequences (in which restriction enzyme sites for cloning are highlighted):

GGTTCAAAGTTTTTTTCTTCCATTTCAGGTGTCGTGAG**GCTAGC**GCCACC [ORF with

ATG start and stop]**GGATCC**CGGGACTAGT**ACGCGT**TAAGTCGACAATC

Template ORF plasmid. The starting library we are using is either in a pUC57 plasmid form

(libraries made via MITE method), or arrive as linear fragments (libraries purchased from Twist

Bioscience). In the cases where the ORF reference sequence contains restriction sites needed for

cloning, the template ORF sequences need to be modified to remove these sites. It is important

that prior to making the library, screening assay development work is done with control clones in

the same vector as the vector for the future library. It is equally important that a clonal plasmid

of the template ORF is first cloned into the vector, made into virus, tested for infection, genomic

DNA prep, and PCR with the gDNA. This clonal plasmid will be used as the template for

mutagenesis.

Library design. A saturation mutagenesis designer written in R is available. The template ORF

sequence is first checked for absence of restriction sites that are used in cloning: either

NheI/BamHI or NheI/MluI. The ORF template may need to be modified in order to satisfy these

requirements. The designer then steps through from N-terminus to C-terminus of the template

ORF one amino acid position at a time to select variant codons. For a given template codon there

are 63 possible variant codons to choose from at each amino acid position. These 63 codons are

divided into two bins: 54 codons that have 2- or 3-nucleotide differences relative to the template

codon (2-3-nt delta), and 9 codons with a single-nucleotide difference (1-nt delta). The codons

that would violate the restriction sites exclusion rule are removed from both bins. In designing

missense or nonsense variants, we first select from the 2-3-nt delta codon bin, and then if needed

from the 1-nt delta codon bin. If within a bin there are multiple codon choices available for an

amino acid variant, the codon is picked by sampling the available codons with probability

weightings according to the codon usage frequency of the organism (*Homo sapiens* in all our

experiments to date, http://www.kazusa.or.jp/codon/cgi-bin/showcodon.cgi?species=9606). For

silent codon changes we pick only from 2-3-nt delta bin. In the end, each of the designed variant

codons substitutes for its corresponding template codon in the full-length reference ORF as one

member of the library.

Mutagenized ORF Library cloning. Saturation mutagenesis library production requires that

variant codons be encoded into pools of oligos. These oligos are then either amplified and

assembled into cloneable linear fragments using the Mutagenesis by Integrated TilEs (MITE)

method or delivered as a linear fragment pool (Twist Biosciences). For MITE method, once the

linear fragments bearing the mutagenized codons are produced, they are then cloned into a high

copy entry vector for amplification and sequence validation. For Twist Bioscience method, the

linear fragment pool is a collection of properly flanked full-length ORFs that are ready for

cloning into pMT025 ORF expression vector.

MITE method: The region of the open reading frame to be mutated is partitioned into

90-base tiles. Oligos representing all mutations in the space of a tile were synthesized. Each 90

base tile is flanked by 30 base sequences that are complementary to the reference ORF. Oligos

for adjacent tiles should be synthesized on separate arrays. Oligo tile pools were amplified by

emulsion PCR using primers to the 30 base flanking regions and purified on a 2% agarose gel.

Entry vector cloning is performed on a per tile basis by linearizing the entry vector at the

appropriate tile position using Phusion polymerase (New England Biolabs) and primers to the

regions flanking the tiles. The linearized vectors were purified on 1% agarose gel and DpnI

treated. The DpnI-treated linear plasmid backbones were then mixed with the relevant PCR

amplified tile and assembled by *in vitro* recombination with the NEBuilder HiFi DNA Assembly

Master Mix (New England BioLabs). The assembly reactions were purified, electroporated into

TG1 *E. coli* cells (Lucigen), and recovered for one hour at 37˚C in Recovery Media (Lucigen).

Aliquots from the transformations were used to inoculate overnight cultures of LB containing 25

µg/mL of Zeocin (ThermoFisher). Cells were harvested by centrifugation and plasmid DNA was

isolated using the QIAGEN Midiprep Plus kit (QIAGEN). Plasmid pools, each corresponding to

a tile, were verified by sequencing Nextera XT (Illumina). A pool of all pUC57-tile libraries was

made with equal weight per variant. This pool was subjected to restriction digest (e.g., NheI and

BamHI). After purification by 1% agarose gel, the excised linear fragment library is ready for

cloning ligation with an expression vector that was pre-processed with the compatible restriction

enzyme pair.

Twist Bioscience method: The mutagenesis library made by Twist Bioscience was

delivered as linear fragments of the full-length ORF flanked with adapters for cloning. The

fragment library was first digested with NheI and BamHI overnight, and then underwent ligation

with pMT025 vector pre-processed with the same restriction enzyme pair.

Cloning ligation: For either method, the ligation was carried out with 5:1 insert-to-vector

molar ratio, using T7 DNA ligase at room temperature for 2 hours. The ligation was cleaned up

with isopropanol precipitation. The resulting DNA pellet was used to transform Stbl4 bacterial

cells. Plasmid DNA (pDNA) was extracted from the harvested colonies using QIAGEN Maxi

Prep Kits. The resulting pDNA library was sequenced via the Illumina Nextera XT platform to

determine the distribution of variants within the library.

Virus preparation and titer measurement. Lentivirus was produced and quantified by the

Broad Institute Genetic Perturbation Platform (GPP). The detailed protocol is available at

http://www.broadinstitute.org/rnai/public/resources/protocols. Briefly, 293T viral packaging

cells were transfected using TransIT-LT1 transfection reagent (Mirus Bio) with the pDNA

library, a packaging plasmid containing *gag*, *pol* and *rev* genes (psPAX2, Addgene), and an

envelope plasmid containing VSV-G (pMD2.G, Addgene). Media was changed 6-8 hours after

transfection, and virus was harvested 24 hours thereafter. The virus titer was measured by

infecting 3000 A549 cells in 96-w cell culture plate with a series of diluted standard virus of

known titer for a standard curve, and a series of diluted library virus of unknown titer to find data

points that lie in the linear range of the standard curve. Twenty-four hours post infection, the

cells were subjected to puromycin selection for 2 days. The surviving cells were quantified with

alamarBlue.

Determine cell numbers to start and finish the screen. In general, infections were performed,

per infection replicate, to aim a representation on average of 1000 cells carrying a given variant

throughout the entire experiment. At the time of splitting for drug-treatment and reference

sample pair, the same rules apply. For positive selection screens, where a majority of the library

will be depleted and a small number of cells carrying variants that are a subset of the library will

survive, and consequently a lower cell number will be collected. In negative selection screen

where the majority of the library will be present, the end-point samples should be of at least 1000

cells per variant (i.e., per library member).

Genomic DNA preparation. Upon the completion of the screen, cell pellets were collected.

Genomic DNA was extracted from cell pellets using the Machery Nagel NucleoSpin Blood Kit

(Midi, Clontech Cat. #740954.20; XL Maxi, Clontech Cat. #740950). The detailed protocol is

available at http://www.broadinstitute.org/rnai/public/resources/protocols.

Design PCR primers. The integrated ORF in genomic DNA will be amplified by PCR. The

PCR products will be shotgun sheared with transposon, index labeled, and sequenced with next-

generation sequencing technology. Each sequenced fragment is generated by 2 transposon

insertion events so that both ends of the fragment are tagged with a transposon. Therefore, the

ends of the gDNA PCR products will be less frequently represented in the NGS library and will

have shallow coverage. For this reason, we advise that the PCR primers be designed in such way

that there is a ~100 bp extra sequence at each end leading up to the mutated ORF region. For

pMT025 vector, we use these 2 primers:

Forward: 5'-ATTCTCCTTGGAATTTGCCCTT-3'

Reverse: 5'-CATAGCGTAAAAGGAGCAACA-3'

Determine the number of PCR reactions per sample. For positive screen end-point samples,

the number of cells that need to go through ORF-extracting PCR is determined in such a way

that for each library variant, gDNA from 500 or more cells will be subjected to PCR. For

reference samples and negative selection screen end-point samples, the cells/variant ratio should

be maintained at 1000 or more. PCR is sensitive to the amount of gDNA input and prior to

screening, these conditions must be optimized. For example, in 50ul PCR reactions, we test

different amounts of gDNA, using a series of 250ng, 500ng, 1ug, 2ug. This titration will help to

determine the highest concentration of DNA that can be used to produce robust PCR products.

In general, for human cell lines, we assume a cell produces 6.6 pg DNA. Here is the formula to

determine how many PCR reactions are needed for a sample:

$$\text{PCR reactions per sample} = 6.6 * \frac{library\ size}{ng\ of\ gDNA\ per\ PCR\ well}$$

ORF amplification from gDNA by PCR. PCR reactions were set up in 96-well plates

according to the optimized PCR condition and number of PCR reactions per sample calculated

above. The volume of each PCR reaction was 50 μL and contained an optimized amount of

gDNA. Q5 DNA polymerase (New England Biolabs) was used as the DNA polymerase. All PCR

reactions for each gDNA sample were pooled, concentrated with a PCR cleanup kit (QIAGEN),

and separated by gel electrophoresis. Bands of the expected size were excised, and DNA was

purified, first using a QIAquick kit (QIAGEN), and then an AMPure XP kit (Beckman Coulter).

Nextera sequencing. Nextera reactions were performed according to the Illumina Nextera XT

protocol. For each sample, we set up 6 Nextera reactions, each with 1 ng of purified ORF DNA.

Each reaction was indexed with unique i7/i5 index pairs. After the limited-cycle PCR step, the

Nextera reactions were purified with AMPure XP kit. All samples were then pooled and

sequenced using an Illumina sequencing platform chosen among Miseq, Hiseq, or NovaSeq,

according to the number of reads needed as calculated below.

Determine number of reads needed for the screen. We need to sequence for every variant

500x (for positive selection screen samples) or 1000x (for negative selection screen samples, or

reference samples) the full-length PCR product. Thus, the number of total reads depends on ORF

size, library size, read length, read quality etc. Below are the formulas:

For negative selection screen samples, or reference samples:

516
$$Reads\ per\ sample = 7000 * \frac{(ORF\ length\ in\ NT)\char`\^2}{read\ length * read\ assembly\ rate}$$

518   Where '*read length*' is typically 300, as we use paired-end sequencing reads, 150

519   bases each, and '*read assembly rate*' is in range of 0.7-0.9.

521   For a positive selection screen sample:

523
$$Reads\ per\ sample = 3500 * \frac{(ORF\ length\ in\ NT)\char`\^2}{read\ length * read\ assembly\ rate}$$

Determine the optimal parameters for variant-calling software 'ASMv1.0' The next-

generation sequences are from transposon-sheared fragments, which are not uniform in size

(average fragment size: 140 base pairs, data not shown). For fragments longer than 2x read

length (e.g., 2 x 150 = 300 bases), the pair of reads are disjoint reads; for fragments shorter than

the read length (e.g., 150 bases), the reads are fully overlapping and each extends into the

transposon adaptor located in the opposite end; for fragments sized between 1x and 2x read

length (e.g., 150-300 bases), the pair of reads overlap but do not have transposon sequences. The

variant calling software first aligns reads to the reference sequence. The transposon tails are then

clipped off at the coordinate determined by the start of its paired read. It then applies the ***min-q***

(minimum base quality) filter to trim the end of the aligned reads to ensure there are at least 15

bases from the trimmed end inwards with base quality above the min-q parameter. The ***min-***

read-length filter admits the trimmed reads that meet the threshold. All failed reads go into the

low-q bin. Next, the filter of ***min-flank-length*** is applied that requires reference bases at end of

the trimmed reads. A min-flank-length of 2 or more bases prevents cases in which a changed

codon is split between 2 Nextera fragments. It also prevents reads ending with transposon

sequences from being called as variants in cases for which transposon clipping fails to work due

to a lack of paired reads. Finally, a variant is called only when the underlying base changes are

above the min-q threshold. In the end, what matters are reads defined as 'wild-type' and reads

defined as 'called variants.'

To tune our software parameters, we used the sequencing data of a saturation mutagenesis

plasmid DNA library for *PDE3A* gene. This library was built by mutagenizing a region of 1425

base pairs using Twist Bioscience services. The errors in the library are concentrated in within 30

bases of a planned codon change (**Fig. 2**). These errors can be explained by mutagenesis primer

synthesis errors. The codon changes are located in the middle of the mutagenesis primers,

sandwiched between two 20-30 base pair flanks. In next-generation sequencing, the transposon

sheared fragments are, on average, 140 base pairs, similar to the sequencing read length. The

close proximity of variants, planned and unintended, indicates that these variants reside on the

same transposon-sheared fragment and on the same sequencing read. Therefore, each library

ORF, after transposon shearing, produces one variant-bearing fragment; the rest of the fragments

are all wild-type. That is, if we ignore miscalls introduced by PCR, the wild-type/variant read

ratio of this library should be: 1425/140 -1 = 9.2. This ratio, 9.2, represents the highest ratio we

could observe, as any PCR miscalls would only shift this ratio lower.

In determining the PCR miscall rate, we processed a clonal plasmid of pMT025-PDE3A. The

sequencing of this clonal plasmid DNA, which had undergone the same PCR process as the

saturation mutagenesis plasmid DNA library, showed error rate of 0.00037 miscalls/sequenced

nucleotide. Thus, when we factor in PCR miscalls, the read ratio of wild-type/variant becomes

5.9:

Start with miscall-free wild-type/variant read count ratio 9.2:1, the miscalls switch

(9.2*140*0.00037)=0.48 from wild-type group to variant group, producing a new wild-

type/variant read count ratio:

$$(9.2 - 0.48)/(1 + 0.48) = 5.9$$

By processing the same plasmid DNA library sequencing data set with various parameters, we

were able to identify the optimal parameters that produce a wild-type/variant ratio of 5.9, and

yield reasonable output of both wild-type and variants: min-q = 30, min-read-length = 40, min-

flank-length = 2.

NGS Data processing and analysis: The data from the NGS pipeline are in .fastq format. Each

NGS cluster produced 2 long reads (150 bases each), and 2 index reads (each 8 bases long). The

2 indexes identify a screen sample, so that a large number of samples can be pooled and

sequenced in bulk. Saturated ORF variant library screen data are typically 4-5TB in data size.

Generous computer storage is needed to process the data. The reads need to go through the

filtering process (**Supplementary Fig. 2**) prior to variant calling.

Read filtering: The reads from an NGS fragment are trimmed (1) by base quality, (2) by

removing non-ORF transposon sequences, (3) by requiring that the outermost nucleotide

variations in the reads have, at minimum, a number of reference bases as specified by the min-

flank-length parameter, and (4) by requiring the trimmed reads be no shorter than the min-read-

length (see **Supplementary Fig. 2**)

Variant calling**:** There are two versions of variant calling software, ORFCallv1.0, the earlier

version; and ASMv1.0, the current version. The reads that have passed the filtering step are used

differently by ORFCallv1.0 and ASMv1.0:

i)   ORFCallv1.0 (an earlier version software; Giacomelli *et al*[15]): Variants are tallied

588   in the context of a single codon space. Without regard of the association of the

589   multiple changes physically in a single molecule, ORFCallv1.0 essentially scans from

590   N- to C-termini, one amino acid position at a time, and sums up the occurrence of

591   codons or amino acids at that position.

593   ii)   ASMv1.0: This is a molecule-centric approach and an enhanced version over

594   ORFCallv1.0. The NGS dataset is analyzed in its entirety, with the following new

595   capabilities: (1) The trimmed reads are evaluated full-length; that is, variants are

596   called in the context of entire read (or read pair). The programmed variant is called

597   when 2 conditions are met: the detection of the programmed codon changes, and the

598   absence of any *additional* nucleotide variations throughout the entire read or read

599   pair. (2) Indel detection: This feature was lacking in ORFCallv1.0. This new feature

600   of ASMv1.0 allows us to process indel libraries.

602   In ASMv1.0, mutations in a molecule are depicted nucleotide-wise by position along

603   the PCR product, followed by the nature of the nucleotide change. For example, 'a

604   change from G to A at position 1140' is described as '**1140:G>A';** 'a deletion of an

605   A at position 1141' is '**1141:A>-**'; 'an insertion of a T after position 1142' is '**1143:-**

606   **>T**'.

608   Also with ASMv1.0, a host of annotations formatted as a datafile can be derived from

609   the variant descriptions. For example, a *TP53* variant with description '**1140:G>A,**

610   **1141:A>C, 1142:G>C**' is annotated as (a pair of brackets marks an annotation):

611   [E343T] [Intended] [one codon change] [343:GAG>ACC] [missense] [substitution].

612   Another *TP53* variant with description '**1140:G>A, 1141:A>C, 1142:G>C,**

613   **1145:G>T**' is: **[**E343T] [Unintended] [two codon changes] [343:GAG>ACC,

614   344:CTG>CTT] [substitution] [missense/nonsense] etc. The first variant is an

intended variant that matches a designed variant of the library. The second variant is

the same amino acid call as the first variant, but it is annotated, rightfully, as an

unintended variant carrying an error 4-nucleotide away from the intended codon

change - this error results in a silent codon change.

Parsing data into a form ready for hit calling**:** The output files from the above variant-calling

software are still large in size (over 100MB), large in numbers of files, and not readily useable.

These output files need to be parsed, aggregated and annotated. For this purpose, we built a

parser called ASMv1.0_Parser, which was written in R, and is available for download (see the

links listed in **Supplementary Table 2**). The Parser Pak consists of a 'readme' file detailing the

instructions, and R codes and training data sets. Through the parser, all data files will be

organized into a single .csv file, one column per screen sample. These .csv files include a short

version that is a subset of the planned variants only, and a complete version that includes all

called variants, including planned/unplanned, insertion/deletion/substitution, etc. In addition, we

also report the ORFCallv1.0 version of data.

Scaling: We use read counts from the NGS data as the quantitative measurements of variants

defined by 'changes' relative to the reference sequence. Reads that entirely match the reference

sequence are not informative, as they are, as a group, the aggregate of all variants whose changes

are located outside of the region covered by the read. At any amino acid position, the total

coverage, specifically the sum of counts of variants passing through the position and the sum of

counts of the wildtype reads, should be consistent throughout the ORF. But that is not actually

the case due to the transposon insertion bias that results in unevenness of total read coverage

(**Supplementary Fig. 8**). In cases that address the relative abundance among variants, as in

Supplementary Figure 9, the raw counts need to be scaled to the fraction of the total read

coverage across the codon position. However, in cases that compare the variant abundance

change between two samples, such as computing fold-changes in screening samples, scaling the

raw counts may not be necessary, as the positional raw count variations caused by transposon

insertion bias will cancel out between two samples when computing fold-changes

(**Supplementary Fig. 10**).

Acknowledgements

We thank our collaborators of saturation mutagenesis projects, Alice Berger, Jesse

Boehm, Ben Ebert, Laura Evans, Eejung Kim, Shengwu Liu, Julie-Aurore Losman, Huong Ly,

Peter Miller, Utthara Nayar, and Nikhil Wagle for demonstrating the diverse utilities of the

technology, Broad Institute Genomics Platform for providing NGS support, Nancy Tran and

Cong Zhu for the early work on saturation mutagenesis method, Stephanie Milczarek, Kirsten

Milenkowic, Michael Zagieboylo, and Jonah Schwartz for conducting lab work for some

projects, Leon Yang for critical reading of the manuscript. This work was supported by NCI U01

CA176058 (WCH), the Mark Foundation (WCH), American Cancer Society Professorship

(MM), American Cancer Society Mentored Research Scholar Grant MRSG-18-202-01 (ALH),

Department of Defense W81XWH-18-PRCRP-CDA (ALH), Team Lick Cancer (ALH), Unravel

Foundation (ALH), and Wong Family Award (ALH)

Author Contributions

X.Y., D.E.R., A.L.H., A.O.G., and T.S. wrote the manuscript. W.C.H., M.M., H.G.,

A.J.A., R.E.L., N.P, M.M, A.B., and F.P. reviewed and edited the manuscript. A.L.H., X.Y., T.S,

and D.E.R. designed the new variant-calling software ASMv1.0. T.S. wrote the variant-calling

software ORFCallv1.0 and ASMv1.0. X.Y., A.L.H., D.E.R., and T.H. optimized the run

parameters of ASMv1.0. X.Y. wrote R codes to design saturated ORF variant libraries and to

parse the output of the variant-calling software. X.Y. conducted data analysis with input from

D.E.R., A.L.H., A.O.G., and assistance from B.P.L. D.A and T.G. adapted the variant-calling

software into a saturation mutagenesis data pipeline. X.Y., A.L.H., A.O.G, R.E.L., B.P.L., B.F,

H.K., E.S., F.P., and L.S. designed and conducted wet-lab experiments. All authors contributed

to and approved the manuscript.

Competing Interests

W.C.H. is a consultant for ThermoFisher, Solasta, MPM Capital, iTeos, Jubilant

Therapeutics, Tyra Therapeutics, RAPPTA Therapeutics, Frontier Medicines, KSQ Therapeutics

and Paraxel. A.J.A has consulted for Oncorus, Inc., Arrakis Therapeutics, and Merck & Co., Inc,

and has research funding from Mirati Therapeutics, Syros, Deerfield, Inc., and Novo Ventures

that is unrelated to this work. A.O.G. is a share and option holder of 10X Genomics.

References

1.   Lefèvre F, Rémy MH, Masson JM: **Alanine-stretch scanning mutagenesis: a simple**
and efficient method to probe protein structure and function. *Nucleic acids research*
1997, **25:**447-448.
2.   Cunningham BC, Wells JA: **High-resolution epitope mapping of hGH-receptor**
interactions by alanine-scanning mutagenesis. *Science (New York, NY)* 1989,
244:1081-1085.
3.   McCullum EO, Williams BAR, Zhang J, Chaput JC: **Random mutagenesis by error-**
prone PCR. *Methods in molecular biology (Clifton, NJ)* 2010, **634:**103-109.
4.   Muteeb G, Sen R: **Random mutagenesis using a mutator strain.** *Methods in molecular*
biology (Clifton, NJ) 2010, **634:**411-419.
5.   Kato S, Han S-Y, Liu W, Otsuka K, Shibata H, Kanamaru R, Ishioka C: **Understanding**
the function–structure and function–mutation relationships of p53 tumor
suppressor protein by high-resolution missense mutation analysis. *Proceedings of the*
National Academy of Sciences 2003, **100:**8424.

6.   Chen L, Jenjaroenpun P, Pillai AMC, Ivshina AV, Ow GS, Efthimios M, Zhiqun T, Tan
TZ, Lee S-C, Rogers K, et al: **Transposon insertional mutagenesis in mice identifies**
human breast cancer susceptibility genes and signatures for stratification.
Proceedings of the National Academy of Sciences 2017, **114:**E2215.
7.   Hughes RA, Ellington AD: **Synthetic DNA Synthesis and Assembly: Putting the**
Synthetic in Synthetic Biology. *Cold Spring Harbor perspectives in biology* 2017,
9:a023812.
8.   Heather JM, Chain B: **The sequence of sequencers: The history of sequencing DNA.**
Genomics 2016, **107:**1-8.
9.   Fowler DM, Araya CL, Fleishman SJ, Kellogg EH, Stephany JJ, Baker D, Fields S:
High-resolution mapping of protein sequence-function relationships. *Nature Methods*
2010, **7:**741-746.
10.   Fowler DM SJ, Fields S. : **Measuring the activity of protein variants on a large scale**
using deep mutational scanning. Nature Protocols. 2014.
11.   Kotler E, Shani O, Goldfeld G, Lotan-Pompan M, Tarcic O, Gershoni A, Hopf TA,
Marks DS, Oren M, Segal E: **A Systematic p53 Mutation Library Links Differential**
Functional Impact to Cancer Mutation Pattern and Evolutionary Conservation. *Mol*
Cell 2018, **71:**873.
12.   Mighell TL, Evans-Dutson S, O'Roak BJ: **A Saturation Mutagenesis Approach to**
Understanding PTEN Lipid Phosphatase Activity and Genotype-Phenotype
Relationships. *Am J Hum Genet* 2018, **102:**943-955.
13.   Weile J, Kishore N, Sun S, Maaieh R, Verby M, Li RJ, Fotiadou I, Kitaygorodsky J, Wu
YZ, Holenstein A, et al: **Shifting landscapes of human MTHFR missense-variant**
effects. *American Journal of Human Genetics* 2021, **108:**1283-1300.
14.   Awad MM, Liu S, Rybkin, II, Arbour KC, Dilly J, Zhu VW, Johnson ML, Heist RS, Patil
T, Riely GJ, et al: **Acquired Resistance to KRAS(G12C) Inhibition in Cancer.** *N Engl*
J Med 2021, **384:**2382-2393.
15.   Giacomelli AO, Yang X, Lintner RE, McFarland JM, Duby M, Kim J, Howard TP,
Takeda DY, Ly SH, Kim E, et al: **Mutational processes shape the landscape of TP53**
mutations in human cancer. *Nature Genetics* 2018, **50:**1381-1387.
16.   Garvie CW, Wu X, Papanastasiou M, Lee S, Fuller J, Schnitzler GR, Horner SW, Baker
A, Zhang T, Mullahoo JP, et al: **Structure of PDE3A-SLFN12 complex reveals**
requirements for activation of SLFN12 RNase. *Nat Commun* 2021, **12:**4375.
17.   Kwon JJ, Hajian B, Bian Y, Young LC, Amor AJ, Fuller JR, Fraley CV, Sykes AM, So J,
Pan J, et al: **Structure-function analysis of the SHOC2-MRAS-PP1C**
holophosphatase complex. *Nature* 2022, **609:**408-415.
18.   Miller PG, Sathappa M, Moroco JA, Jiang W, Qian Y, Iqbal S, Guo Q, Giacomelli AO,
Shaw S, Vernier C, et al: **Allosteric inhibition of PPM1D serine/threonine**
phosphatase via an altered conformational state. *Nat Commun* 2022, **13:**3778.
19.   Persky NS, Hernandez D, Do Carmo M, Brenan L, Cohen O, Kitajima S, Nayar U,
Walker A, Pantel S, Lee Y, et al: **Defining the landscape of ATP-competitive inhibitor**
resistance residues in protein kinases. *Nat Struct Mol Biol* 2020, **27:**92-104.
20.   Majithia AR, Tsuda B, Agostini M, Gnanapradeepan K, Rice R, Peloso G, Patel KA,
Zhang X, Broekema MF, Patterson N, et al: **Prospective functional classification of all**
possible missense variants in PPARG. *Nat Genet* 2016, **48:**1570-1575.

21.   Melnikov A, Rogov P, Wang L, Gnirke A, Mikkelsen TS: **Comprehensive mutational**
737   **scanning of a kinase in vivo reveals substrate-dependent fitness landscapes.** *Nucleic*
738   *Acids Res* 2014, **42:**e112.
739   22.   Faure AJ SJ, Baeza-Centurion P, Lehner B.: **DiMSum: an error model and pipeline for**
740   **analyzing deep mutational scanning data and diagnosing common experimental**
741   **pathologies.** *Genome Biol* 2020, **21:**207.
742   23.   Legnini I, Alles J, Karaiskos N, Ayoub S, Rajewsky N: **FLAM-seq: full-length mRNA**
743   **sequencing reveals principles of poly(A) tail length control.** *Nature Methods* 2019,
744   **16:**879-886.
745   24.   Starr TN, Greaney AJ, Hilton SK, Ellis D, Crawford KHD, Dingens AS, Navarro MJ,
746   Bowen JE, Tortorici MA, Walls AC, et al: **Deep Mutational Scanning of SARS-CoV-2**
747   **Receptor Binding Domain Reveals Constraints on Folding and ACE2 Binding.** *Cell*
748   2020, **182:**1295-1310.e1220.
749   25.   Marx V: **Long road to long-read assembly.** *Nat Methods* 2021, **18:**125-129.
750   26.   Wenger AM, Peluso P, Rowell WJ, Chang P-C, Hall RJ, Concepcion GT, Ebler J,
751   Fungtammasan A, Kolesnikov A, Olson ND, et al: **Accurate circular consensus long-**
752   **read sequencing improves variant detection and assembly of a human genome.**
753   *Nature Biotechnology* 2019, **37:**1155-1162.
754   27.   Frank F, Keen MM, Rao A, Bassit L, Liu X, Bowers HB, Patel AB, Cato ML, Sullivan
755   JA, Greenleaf M, et al: **Deep mutational scanning identifies SARS-CoV-2**
756   **Nucleocapsid escape mutations of currently available rapid antigen tests.** *Cell* 2022,
757   **185:**3603-3616 e3613.
758   28.   Esposito D, Weile J, Shendure J, Starita LM, Papenfuss AT, Roth FP, Fowler DM, Rubin
759   AF: **MaveDB: an open-source platform to distribute and interpret data from**
760   **multiplexed assays of variant effect.** *Genome biology* 2019, **20:**223-223.
761   29.   Boettcher S, Miller PG, Sharma R, McConkey M, Leventhal M, Krivtsov AV,
762   Giacomelli AO, Wong W, Kim J, Chao S, et al: **A dominant-negative effect drives**
763   **selection of TP53 missense mutations in myeloid malignancies.** *Science*
764   2019, **365:**599.
765

Figure legends

Figure 1. Pooled screens with ORF variant libraries versus shRNA/sgRNA libraries. A

plasmid pool of thousands of genetic perturbation agents is first packaged into lentiviral

particles. Cells were then transduced with the pool of virus at an infection rate (the proportion of

infected cells over total cells) of 30-50% to ensure that most of the infected cells get a single

genetic perturbation. The perturbed cell population is then subjected to either selection pressure,

or treatment under a reference condition. Upon the completion of the screen, the remaining cell

population is harvested and processed for enrichment or depletion of each library perturbation by

comparing a sample that has undergone selection and a sample that has been treated under a

reference condition. **(a) A typical pooled screen using shRNA/sgRNA libraries.** For

shRNA/sgRNA, the collected cells are first processed for genomic DNA (gDNA), which serve as

the PCR template to amplify the perturbagen barcodes. These barcodes are in fact the functional

elements themselves, the hairpins of shRNA or guides of sgRNA. These barcodes are short

enough to be read straight out by a 50-base NGS read. **(b) ORF variant library screens.** In an

ORF variant library, however, the variant signatures are subtle as each member of the library

differs from the template by a single codon variation, and the variant signatures of the library are

spread out across the entire ORF. Consequently, the detection of variant requires (1) PCR-

amplification and then shotgun shearing of the full-length ORF, (2) next-generation sequencing

of the resulting fragments, (3) robust variant detecting tools to align the reads to the template,

then identify, evaluate, annotate and tally the variant-defining reads. The NGS reads consist of

variant signature-carrying reads (shown as red arrows) and wild-type reads (black arrows). The

variant-calling software processes the NGS reads to detect variant signature-carrying reads,

which is a minority of the total reads, and records the counts of each variant. These counts are

the basis for downstream analyses that map the variants to the phenotype strength in the form of

log2-fold change (LFC).

Figure 2. Reanalysis of *TP53* saturation mutagenesis screen using ASMv1.0. Screens most

often involve the comparison of a sample subjected to selective pressure with a sample processed

under a reference condition. This Giacomelli *et al*[15] p53[NULL] A549 cell/nutlin-3 screen

enriches *TP53* variants that exhibit loss-of-function (LOF), and depletes wild-type-like variants.

(**a**) Heatmaps: each variant is placed in a 2-dimensional grid, where codon position is presented

along the horizontal dimension and the amino acid mutation along the vertical dimension. We

assigned the letter 'X' to represent nonsense mutations, and 'B' to represent silent codon

changes. The value of the log fold change (LFC) between treatment and reference samples is the

heat index. We processed the same original NGS data through the two versions of software and

presented the resulting log fold change (LFC). The pair of heatmaps are nearly identical to the

naked eye. To help inspect all variants one by one, the third heatmap used the

delta(LFC)=(LFC^{ASM} – LFC^{ORFCall}) as the heat index. This heatmap revealed variants that were

scored differently by the two versions of software. (**b**) Scatter plot comparing all LFC values

produced by the two versions of software, each dot representing a planned variant. The colors are

scaled by variant purity. (**c**) Demonstrating the impact of variant purity on the improved

performance of the new ASMv1.0 software. The new software's massive output allows us to

compute 'variant purity' defined by the fraction of pure planned variant over all variants that

harbor the planned variant, with or without unintended changes in the same sequencing reads.

We binned the variants by variant purity and in each purity bin we assessed the Pearson

correlation coefficient between two sets of LFC values produced by the two versions of software.

It is clear that high variant purity results in similar LFC values, whereas as the purity decreases,

the two sets of fold-changes diverge. The new software's ability to detect and tally variants that

carry additional unintended changes is called for when the variant purity is low. Similarly, we

analyzed two other Giacomelli *et al*[15] screens of the *TP53* saturated ORF variant library and

the results are presented in **Supplementary Figure 3** and **4**. (**d,e**) Validation of ASMv1.0 using

the published data by Kotler *et al.*[11] The Kotler dataset includes a screen of p53^{NULL}H1299

cells with a library covering *TP53* DBD. We first took a subset of Giacomelli library to cover the

same *TP53* region as Kotler screens, resulting in a total of 2964 variants (**d**, **left**). We then

identified 76 variants that recorded most different LFC scores by our two versions of

software (**d, right**). We then compared the Kotler scores against our LFC scores called by

ORFCallv1.0 (**e, left**) and ASMv1.0 (**e, middle**). We observed that between two versions of

software, ASMv1.0 calls are clearly more in agreement with Kotler calls (**e, right**).

Figure 3. The errors in our saturated ORF variant libraries are low in number and local

relative to the location of the planned codon change. In screens, the mutational spaces of

libraries depend on how many planned variants can be detected in the final sequencing data. A

library with a high impurity rate will require a larger experimental effort. (**a**) **Breakdown of the**

planned variants and those with errors. Going through all called variants that contain a

planned codon change, we identified the number of unintended nucleotide changes. This bar-

chart shows that in the pMT025-TP53 library, 84% molecules are planned variants without

additional mutations. Of 16% imperfect molecules, the majority are those with 1-2 extra

nucleotide changes. (**b**) **Major errors are in the vicinity of a planned change.** We first pinned

down the planned change in a read and then travelled in both directions and tallied the errors as a

function of the distance to the pinned-down codon change. In this figure, we compared two

libraries made with different technologies, *TP53* library by MITE and *PDE3A* by Twist

Bioscience. The error distribution profile along the distance to a planned variant is strikingly

explainable by the oligo synthesis scheme involved in these technologies. MITE technology uses

synthesized oligos of 150 bases long, of which 90 bases (called TILE) go into the final products,

whereas Twist Bioscience introduced errors through oligo primers with the intended codon

changes placed in the middle and flanked with 15-25 bases on either side. The major errors are

concentrated within ~100 bases (MITE method) or ~30 bases (Twist Bioscience method) from a

planned codon change. This is important because the close proximity of the errors to the

intended codon change renders the planned change and errors all detectable in a single read (e.g.,

150 bases) or read pair (e.g., 2x150=300 bases).

Figure 4. The effect of PCR and sequencing errors on variant quantification. (a) A clonal

TP53 reference plasmid (i.e. all sequences are wild-type) was processed as if it were a screening

sample. The base call at each position is predominantly, but not absolutely, the reference

nucleotide. The non-reference calls represent artifactual variants that are not actually in the

library. The heatmap on the left shows tallied counts of called nucleotides at each nucleotide

position; a blank spot represents zero counts. The heatmap on the right depicts the rate of

miscalls. Blank spots in this heatmap are either the wild-type nucleotide or nucleotides of zero

counts. (**b**) The miscalls resulting from PCR errors inflate the abundance measurement of

variants whose variant-defining codon is 1-nt away from the reference codon. In the *TP53* library

screen data, predominately affected are variants of 1-nt (not 2- or 3-nt) difference from the

reference codons. As a result, the variants of 1-nt delta codons as a group appear more abundant

than groups of 2- or 3-nt delta codons (**left**). The artificial 'variant' calls resulting from PCR

errors can be corrected with abundance adjustment by subtracting the miscall rate as determined

by the clonal sample experiment (**right**). The variant abundance measurements of the 2-nt and 3-

nt delta codons are not affected by the miscalls. (**c**) **Suppression of fold-changes in 1-nt delta**

codon variants. PCR errors in the workflow inflate the counts of variants that differ from the

template codon by one nucleotide. Using the pMT025-TP53 library as an example, 1-nt delta

variants showed much narrower fold-change values than 2-nt and 3-nt delta variants. Thus, in the

library design stage, when we have choices of 2- or 3-nt delta codons, we pick them over 1-nt

delta codons. However, we do admit 1-nt delta codons when they are the only choices available;

without them, the library would have gaps in coverage. The fold-change suppression in the 1-nt

delta group leads to false negative calls. In order to mitigate this error, correction of variant

counts with data from a carefully constructed clonal plasmid control experiment may be helpful

for 1-nt delta variants. Altogether, we have arrived at an important rule for saturated ORF

variant library design: minimizing the use of codons that are 1-nt away from the reference

codons.

Figure 1

a

shRNA/sgRNA library screens

plasmid library   viral particles   transduction   genomic DNA   NGS

Direct readout
50-base reads

reference condition

under selection

~20 bases

log-fold change
LFC

b

ORF variant library screens

reference condition

under selection

PCR-amplification of full-length ORF
Shotgun shearing
150-base paired reads

ORF
0.5 - 4 kb

variant-calling
tool

variant-calling
tool

LFC

Figure 2

Figure 3

a

b

Figure 4