**Defining protein variant functions using high-complexity mutagenesis libraries and**

**enhanced mutant detection software ASMv1.0**

Xiaoping Yang[1,9], Andrew L. Hong[2,5,6,9], Ted Sharpe[3,9], Andrew O. Giacomelli[2,7,9,10], Robert E.

Lintner[4,11], Douglas Alan[1], Thomas Green[1], Tikvah K. Hayes[2,7], Federica Piccioni[1,12], Briana

Fritchman[1], Hinako Kawabe[1], Edith Sawyer[1], Luke Sprenkle[1], Benjamin P. Lee[5,6], Nicole S.

Persky[1], Adam Brown[1], Heidi Greulich[2,7], Andrew J. Aguirre[2,7], Matthew Meyerson[2,7,8], William

C. Hahn[2,7], Cory M. Johannessen[2,13], and David E. Root[1,*]

Affiliations:

[1]Genetic Perturbation Platform, [2]Cancer Program, [3]Data Science Platform, and [4]Genomics

Platform, Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA

[5]Department of Pediatrics, Emory University, Atlanta, GA 30322, USA

[6]Aflac Center for Cancer and Blood Disorders, Children's Healthcare of Atlanta, Atlanta, GA

30322, USA

[7]Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA 02215, USA

[8]Department of Genetics, Harvard Medical School, Boston, MA 02115, USA

[9]These authors contributed equally

[10]Current Address: Ontario Cancer Princess Margaret Cancer Centre, 610 University Avenue,

Toronto, ON, M5G 2M9 Canada

[11]Current Address: RAN Biotechnologies, 100 Cummings Center, Ste 434J, Beverly, MA 01915,

USA

[12]Current address: Merck Research Laboratories, 33 Avenue Louis Pasteur, Boston, MA 02115,

USA

[13]Current address: Novartis Institutes for Biomedical Research, Oncology Innovative Targets and

Technologies, 250 Massachusetts Ave, Cambridge, MA 02139, USA

Correspondence should be addressed to D.E.R. ([droot@broadinstitute.org](mailto:droot@broadinstitute.org))

**Abstract**

Pooled variant expression libraries can test thousands of variants of a gene in a single

multiplexed experiment. In a library encoding all single-amino-acid substitutions of a protein,

each variant differs from its reference only at a single codon-position located anywhere along the

coding sequence, such that identifying these variants by sequencing to read out the outcomes of

the screen is a major technical challenge. A popular but expensive brute-force approach is to

divide the pool of variants into much smaller sub-libraries in each of which only a small region is

mutated to allow direct sequencing readout of variant abundance. Here we present an approach

to screen very large variant libraries with mutations spanning a wide region in a single pool,

including library design criteria and mutant-detection algorithms that permits reliable calling and

counting of variants from large-scale sequencing data.

**Keywords**: saturation mutagenesis, deep mutational scanning (DMS), variants, open reading

frame (ORF), log-fold change (LFC), base miscall, DNA oligonucleotide synthesis error, DNA

polymerase error, next-generation sequencing (NGS).

**Background**

A pooled open reading frame (ORF) variant expression library for a gene of interest can

be screened for cellular phenotypes, providing a powerful means to systematically characterize

the functions of thousands of single-amino-acid substitutions in a single experiment. Advances in

DNA oligonucleotide and gene synthesis [1] and in next-generation sequencing [2] have enabled

production and screening of large variant pools. Pools encompassing all possible single-amino-

acid changes for every amino acid position of a protein are known as saturation mutagenesis or

deep mutational scanning (DMS) libraries. Screening such a library in cells can map each variant

to phenotype in a single experiment.

As illustrated in **Figure 1**, a typical pooled screen tests the functions of thousands of

genetic perturbation agents in a single experiment. In general, cell-based pooled screens rely on

introduction of one genetic perturbation cassette, e.g., a CRISPR sgRNA, an shRNA, or an ORF

variant, per cell into the genome using delivery by a retrovirus, so that a single perturbation

cassette is inserted into genomic DNA of each cell. The perturbed cell population is then

subjected to selection pressure, for example, treatment with a drug, flow-sorting for reporter

expression, or simply growth in culture. Post-selection cell samples are collected and processed

to measure the abundance of library members in each sample by sequencing. Enrichment or

depletion of each library perturbation in samples that have undergone selection versus samples

under reference conditions (e.g., pre-selection or vehicle treatment samples) reveals the impact

of the perturbations. For CRISPR screens, the functional elements themselves, the sgRNAs,

serve as ideal short unambiguous barcodes to use for screen "deconvolution", the measurement

of enriched and depleted members of the library. The deconvolution of CRISPR libraries is a

direct readout that simply counts sgRNA-matching sequencing reads. In contrast, ORF variant

library screens are confronted with greater challenges for variant detection.

The complexity of a site-saturated ORF variant library is a function of the ORF length

and the number of desired variants per codon position. As illustrated in **Supplementary Figure**

**1**, at each amino acid position, the template amino acid is typically mutated to 19 other amino

acids, a stop codon, and when possible synonymous codons that can serve as loss-of-function

(LOF) and negative controls respectively (**Methods**). Factoring in the amino acid positions to be

mutagenized (N), the library is a collection of approximately 20*N variants, or alleles, each

differing from the template ORF at a single amino acid position. Capturing differences among

the variant-defining signatures as subtle as a single codon variation located anywhere along the

full-length ORF sequence is a significant technical challenge. A popular approach has been to

employ multiple sub-libraries, each with short mutated regions with lengths of 25 [3, 4], then 47

[5], 100 [6], and recently as many  as 170 [7] amino acids. These sub-libraries were screened and

sequenced separately. Each of these sub-libraries covers a mutated region short enough to be

covered by a single next-generation sequencing read for direct readout of variants. While

practical for small ORFs, for longer genes this approach requires an unwieldy number of mini-

libraries and screens. Our methodology, on the other hand, permits screening of very large pools

in a single experiment using robust variant-calling tools that can accurately quantify variants

from sequencing data generated by amplifying and randomly shearing the entire ORF. A single

site-saturated variant library is thus adequate a wide range of ORF lengths, e.g., ~600-bp *KRAS*

[8]*, ~1200-bp *TP53* [9] and *SMARCB1,* ~1400-bp catalytic region of *PDE3A* [10], ~1700-bp

*SHOC2* [11], ~1800-bp *PPM1D* [12], ~2200-bp *EZH2*, ~3600-bp *EGFR*) (**Fig. 1b**). Here, we

present methods and strategies developed in the process designing, constructing and screening

more than 20 variant libraries (**Supplementary Table 1**). We have made available our library

designer, variant-calling software, and data analysis tools (**Supplementary Table 2**). We

anticipate that these tools and methods will facilitate variant screening for longer ORFs.

**Results**

**Variant calling with ASMv1.0**

The challenge of calling variants correctly from PCR-amplified, fragmented, and

sequenced genomic DNA is due to the multiple factors that lead to unexpected or misattributed

variant sequences. In addition to the intended, library-designed codon sequence modifications

versus the reference sequence, each ORF molecule in our library is subject to two kinds of

unintended changes: oligonucleotide chemical synthesis errors and DNA polymerase errors.

Although the detailed synthesis strategies can vary, ORF variant libraries are generally all

produced using chemically synthesized oligonucleotide strands that encode the variant-defining

codon modifications and that constitute part of the final full ORF library sequences together with

unmodified reference (often "wild-type") portions of the ORF that are enzymatically synthesized

by high-fidelity DNA polymerases in a PCR process. To fully characterize the composition of

the initial library, and of the variant distributions from the genomic DNA of screened cells, it is

desirable to first detect empirically all sequence variations between the reads and the reference

sequence, without consideration of the intended library design. In general, these variations will

include nucleotide substitutions, insertions, deletions and combinations thereof. Some variations

exactly match design-intended variants, while others carry unexpected nucleotide changes in

addition to, or instead of, the expected variation. Aside from the initial library production process,

the process of extracting and sequencing ORF molecules from the library pDNA or the genomic

DNA of screened cells, often called "library deconvolution", is itself subject to errors from

imperfect PCR amplification, and from sequencing miscalls (e.g., a Phred Q20 base-call has a 99%

chance of being correct). Our enhanced variant-detection software aimed to distinguish the

intended variants from those arising from errors of synthesis of the initial library or from the

library "deconvolution" process, and to assess library quality and deconvolution miscall rates.

Improving upon our earlier version of variant-calling strategy and the associated software

(ORFCallv1.0), we developed Analyze Saturation Mutagenesis v1.0 (ASMv1.0; see **Methods**

and **Supplementary Fig. 2**) to more accurately interpret pooled variant NGS data and measure

variant abundance. ASMv1.0 is now included in the Genome Analysis Toolkit (GATK v4.2.0.0).

NGS reads are each first aligned to the reference (or template) ORF to produce binary alignment

map (BAM) files that constitute the input to ASMv1.0. ASMv1.0 then performs all the steps

required, starting from per-sample read-and-reference alignments, to produce a list of observed

nucleotide sequence variations in relation to the reference ORF and the abundance of each of

these nucleotide sequence variations for each sample.  Notably, the list of observed nucleotide

sequence variations is independent of any prior expectations of the library's variant composition.

As shown in **Supplementary Figure 2** , the key steps in this process are: (1) trimming from

reads the flanking sequences introduced during fragmentation and barcoding of the ORF DNA,

(2) combining overlapping paired-end reads when possible to obtain a "per-molecule" single

sequence result per read pair, (3) applying base quality, minimum read length and other

exclusion filters prior to variant calling, (4) identifying bona fide sequence variations, if any, for

each read or read pair versus the reference (substitutions, insertions, deletions, or a set thereof ),

and (5) counting examples of each set of observed nucleotide sequence variations to quantify the

abundance of it in each screening sample.  Because the NGS reads are far shorter than the full-

length ORF and because each planned variant in the library typically carries only a single codon

change from the template ORF, most observed aligned reads will exactly match the reference

sequence. These perfectly reference-matching reads are used to assess the depth of read coverage

across the reference ORF and this read coverage is used to normalize the variant abundance.  The

reads that contain mutations provide the identity and abundance of each variant sequence that is

defined by the mutations. Because the ASMv1.0 algorithm identifies empirically observed

variants in an entirely data-driven manner that does not rely on expectations from the library

design, the output can be used not only to filter for the results for designed ORF variants (i.e., the

intended variants), but also to assess the library quality and characterize its actual composition,

which will always differ to some extent from the intended design.

**The enhanced performance of ASMv1.0**

ASMv1.0 incorporates several enhancements versus an earlier variant-detection process,

implemented in the software package ORFCallv1.0.  ORFCallv1.0 is a simple and effective

variant detection tool designed strictly for single amino acid substitution variant libraries, and it

was employed to interpret a number of saturation mutagenesis screen results [9, 13-15].

ORFCallv1.0 quantifies variant abundance by first aligning reads to the reference sequence, then

recording at each codon position the count of each of 64 possible codons. Note that ORFCallv1.0

tallies the codon read counts at each amino acid position by staying within the 3-nucleotide

codon window and ignoring any additional changes located in the same read. If a library has high

rate of synthesis errors (e.g., due to oligonucleotide chemical synthesis errors), these errors will

lead to high frequency of unintended changes that are combined, in the same molecule, with an

intended codon change. Ignoring the presence of these unintended changes in the manner of the

ORFCallv1.0 algorithm can be costly when these additional nucleotide changes produce

phenotypes that differ from the phenotype of the pristine intended variant. Because ASMv1.0

calls variants in the context of full-length read pairs, incorporating all nucleotide variations

observed between the read pair and the reference sequence including all combinations of

substitutions, insertions and deletions, it can separate pure variants that bear only the intended

sequence changes from those that also harbor extra changes. **Supplementary Table 3**

summarizes the enhanced functionalities from ORFCallv1.0 to ASMv1.0.

We used the screens performed in Giacomelli *et al* [9] to compare the two variant-

detecting methods. Ectopically expressed *TP53* variants were screened in isogenic p53$^{WT}$ and

p53$^{NULL}$ A549 lung cancer cells, with selection pressure from either nutlin-3 or etoposide. Under

nutlin-3 treatment, the screen of p53$^{NULL}$ cells enriched *TP53* loss-of-function (LOF) variants

and depleted wild-type-like variants (**Fig. 2**). Under etoposide treatment, the p53$^{NULL}$ cells

enriched wild-type-like *TP53* variants and depleted the *TP53* LOF variants (**Supplementary Fig.**

**3**). Under nutlin-3 treatment, the screen of p53$^{WT}$ cells enriched *TP53* variants exhibiting

dominant-negative effects (DNE) (**Supplementary Fig. 4**). Using both the ORFCallv1.0 and

ASMv1.0 variant-calling methods, we reprocessed the next-generation sequencing data from

these three screens. The phenotypic strength of each variant, measured as log-fold change (LFC)

between pre- and post-treatment samples, produced similar heatmaps, indicating that the two

variant-calling algorithms are very consistent with each other for most variants (**Fig. 2a,**

**Supplementary Figs. 3a, 4a**). However, we noted that a small subset of variants (~3% of the

library) exhibited substantial differences between the LFC values (delta(LFC)=LFC$^{ASM}$-

LFC$^{ORFCall}$) calculated using these two algorithms (**Fig. 2b, Supplementary Figs. 3b** and **4b).**

In a perfect library consisting solely of intended variants, ORFCallv1.0 and ASMv1.0

should be equal in performance. However, owing to errors introduced during library synthesis

such as oligonucleotide synthesis errors and DNA polymerase errors incurred in the production

of the ORF variant materials, it is inevitable that planned codon changes are sometimes

physically accompanied by one or more unintended accidental nucleotide changes in the same

molecule. We define the 'variant purity' of a designed variant as the ratio of molecules that

exhibit exclusively the intended mutation to the sum all molecules that harbor the intended codon

change, regardless of whether they also harbor additional nucleotide changes. When the purity of

a planned variant is high, both versions of the software will call and quantify variants similarly;

however, when the variant purity is low the enrichment and depletion scores measured by

ORFCallv1.0 can be misleading. We computed the purity of every planned variant using the

reference samples (samples without selection pressure) of the Giacomelli screens, and for each

purity bin we computed the Pearson correlation coefficient of two sets of LFCs, one using

ORFCallv1.0, the other using ASMv1.0 (**Fig. 2c, Supplementary Figs. 3, 4**). As expected, the

two methods correlate well when the variant purity is high and diverge as the variant purity

decreases.

Work by Kotler *et al* [5] offered an additional way of testing the performance of

ASMv1.0 and ORFCallv1.0. We compared the Giacomelli screen LFC scores called by both

versions of the software with the results from the Kotler *et al* [5], which used a smaller variant

library that was focused on the p53 DNA binding domain (DBD) and screened in p53$^{NULL}$

H1299 cells. Kotler *et al* targeted p53 DBD with four sub-libraries, each covering a 47-amino

acid region of the ORF. The four sub-libraries were screened separately, and the screen samples

were processed by direct sequencing readout of the mutational region of each of the 4 libraries.

Like the Giacomelli p53$^{NULL}$ A549/nutlin-3 screen, the Kotler p53$^{NULL}$ H1299 cell screen

enriched for *TP53* alleles that exhibit LOF and depleted wild-type-like variants. From the 7890-

variant Giacomelli p53$^{NULL}$ A549/nutlin-3 screen data, we extracted a 2964-variant subset that

covered the same amino acid mutational space as the four combined Kotler H1299 screens.

These 2964 Kotler scores and Giacomelli scores, using either ASMv1.0 or ORFCallv1.0,

exhibited good correlation (Pearson coefficient 0.65 (p-value=0), and 0.62 (p-value=2.4e-317)

respectively), despite the fact that the screens were performed using different variant detection

methods, different libraries, different cell lines, and different conditions consisting in one case of

simple growth in culture and in the other nutlin-3 treatment (**Supplementary Fig. 5**).

Interestingly, when focusing on p53 DNA binding domain, both Kotler and Giacomelli data

exhibit a bimodal pattern representing the enriched p53 loss-of-function variants

(**Supplementary Fig. 5, left,** upper right quadrant) and the depleted p53 wild-type-like variants

(**Supplementary Fig. 5, left,** lower left quadrant). There is therefore a natural cutoff to classify

each of the 2964 variants as having either LOF or WT-like activity. **Supplementary Figure 5**

suggests that we may use score 0.3 as the cutoff for classification**.** Consistent with the high

similarity noted above between the ASMv1.0 and ORFCallv1.0 scores for most variants, the

agreement between Kotler and Giacomelli classifications is similar whether using the ASMv1.0

method (2519 out of 2964 variants, 85% agreement) or the ORFCallv1.0 method (2482 out of

2964 variants, 84% agreement). This is to be expected for a library that has mostly high purity

variants for which ASMv1.0 and ORFCallv1.0 show high agreement. However, focusing on the

small subset of 76 variants that ASMv1.0 and ORFCallv1.0 scored very differently in the

Giacomelli data (abs(delta(LFC)) >1, **Fig. 2d, right**), the ASMv1.0 classification agrees with the

Kotler classification much better (69 out of 76 variants, 91%) than the ORFCallv1.0

classification does (42 out of 76 variants, 55%, **Fig. 2e**).

**229**   **ASMv1.0-enabled analysis reveals that the secondary mutations (errors) in the library are**

**230**   **near the targeted mutation and are introduced by errors in DNA oligonucleotide synthesis**

**231**   　　The ability of ASMv1.0 to detect molecules with intended or unintended changes along

**232**   the entire read pair allows in-depth analyses of ORF variant library composition. Thus, in

**233**   addition to the variant purity discussed earlier, we can also tally and characterize the unintended

**234**   nucleotide changes to profile the errors introduced during library construction. For example, we

**235**   were able to determine that in the *TP53* library, 84% of molecules are pure and planned variants

**236**   (**Fig. 3a, top**), and the errors in the library are predominantly single nucleotide changes.

**237**   　　Using ASMv1.0 output, we also investigated the proximity distribution of errors in

**238**   relation to the intended codon change. In **Figure 3**, we compared libraries made by two different

**239**   methods: a *TP53* library made by MITE (Mutagenesis by Integrated TilEs) [15], and a *PDE3A*

**240**   library synthesized by Twist Bioscience [10]. The MITE method used chemically synthesized

**241**   oligonucleotide strands to encode 90-nucleotide variant 'tiles' that were subsequently

**242**   incorporated with the rest ORF segments to form the full-length ORF sequence. These ORF

**243**   reference segments were prepared by high-fidelity DNA polymerase used in PCR. The Twist

**244**   Bioscience technology involved the chemical synthesis of oligonucleotides varying in length

**245**   between 30-50 nucleotides, depending on GC content, with the variant-encoding bases located in

**246**   the middle of the oligonucleotide; these oligonucleotides were used as the PCR primers and

**247**   became part of the ORF sequences of the library.  We profiled the library errors (i.e., unintended

**248**   nucleotide changes) by the number of bases between the locations of the error and the intended

**249**   codon change (**Fig. 3b).** We observed that the stretches of frequent errors in libraries produced

**250**   by the two methods, ~90 bases for the MITE method and ~30 bases for the Twist Bioscience

**251**   method, corresponding to the length of chemically synthesized DNA oligonucleotides involved

in library production. Since most of the errors in an ORF variant are concentrated near the

intended codon change, sequencing reads as short as 100 bases can detect most of the unintended

changes that accompany each intended change.

It is important to note that all nucleotides in variant ORF libraries come from either of

two sources: (1) chemically synthesized oligonucleotides that encode and introduce the

mutations (e.g., PCR primers or oligonucleotide tiles that encode variants), or (2) nucleotides

copied from the reference template by high-fidelity DNA polymerase. The errors in these

libraries are therefore expected to arise predominantly from oligonucleotide chemical synthesis

errors (~2 errors per kb), since DNA polymerase (Q5, Fushion, or Ex Taq) nucleotide

misincorporation rates are far lower [16]. A high-fidelity DNA polymerase with an error rate of

$10^{-6}$ errors per cycle per amplified nucleotide will produce only a small fraction of miscopied

molecules. For example, with that error rate a 1kb-ORF, after amplification with 12 PCR cycles,

will yield 1.1%  error-containing molecules due to DNA polymerase misincorporation. In

contrast, oligonucleotide chemical synthesis errors contribute far higher proportions of ORF

molecules with errors, 10% by Twist Bioscience method and 18% by MITE method

(**Supplementary Table 4**). In this case, NGS reads of 100 bases or longer that carry the expected

mutations, which mark those reads as those covering chemically synthesized nucleotides, will

effectively also capture most of the accompanying synthesis errors in the library.

**ASMv1.0-enabled analysis assesses the effect of miscalls (artifactual errors) introduced by**

**processes involving PCR.**

The synthesis errors discussed above are incorporated into the variant libraries and the

resulting variants, albeit not part of the intended library design, are screened in cell populations

to which the libraries are applied. It is important also to consider library assessment errors in the

form of miscalls introduced by the variant quantification process itself.  To deconvolute the

variant abundances from a pooled library, PCR was used to replicate the ORF DNA from the

genomic DNA of the library-transduced cells (up to 30 PCR cycles) and was used again in to

barcode samples in preparation for NGS (12 cycles). We define 'miscalls' as the artifactual

nucleotide calls that do not accurately reflect a sequence that was actually present in the sample.

In this work, miscall rates are driven by the error rate of DNA polymerases used for PCR in

extracting ORF from gDNA and in the subsequent sequencing sample preparation, and also the

sequencing error rates which are reported as Phred Q scores. To directly assess the rate of

miscalls in the deconvolution of a variant library, we processed and sequenced a clonal pure

reference *TP53* ORF sample that should therefore produce only reads that perfectly match the

reference sequence. We observed miscalls at a rate of ~3-6 miscalls per 10,000 sequenced bases

(**Fig. 4a, right**).

The miscall rate is sufficiently large to be a significant factor in variant detection for

large complex libraries. As an example, illustrated in **Supplementary Figure 6**, miscalls lead to

(1) inflated counts of variants with codons that are one-nucleotide different from the template

codon (we refer to this variant type as '1-nt delta' codon), and (2) as a result of these inflated

counts, artifactually reduced log-fold changes between samples of 1-nt delta codons that are truly

present in the library (**Supplementary Fig. 6,** legend). Indeed, in the *TP53* library screen with

p53$^{NULL}$ cells/nutlin-3, we observed the inflation of counts (**Fig. 4b**), and the suppression in fold-

change (**Fig. 4c**), in the 1-nt delta variant group relative to 2- or 3-nt delta variants. This

contribution of PCR errors to 1-nt delta codon change counts has been observed in every library

we have processed (~20 to date), and the effect increases as ORF length increases (data not

shown).

The effects of PCR/NGS miscalls on the quantification of variants reshaped our strategy

in library design rules and adding control experiments. First, we avoid the use of codons that are

one nucleotide away from the reference codons whenever possible. For a typical ORF sequence,

to achieve all possible single site amino acid substitutions, it is sometimes necessary to use

codons that deviate from the corresponding reference codons by one nucleotide. As result of this

design rule that limits the use of 1-nt delta codons, variants of 1-nt delta codons can be reduced

to approximately 3% (from otherwise ~15%) of the library members. Secondly, to better

interpret the results of those 3% 1-nt delta variants, we perform the deconvolution process on a

control sample that is a single clone of the reference ORF and therefore contains only one

uniform sequence. This sample is processed exactly as if it were a screening sample. These

clonal reference ORF data may be used to identify the 1-nt delta variants that are most affected

by miscalls, or can even be used to apply a correction to the screen sample data based on the

background miscall rates determined from this sample (**Fig. 4b**, **right**).

**Silent variants are used as a measure of screen baseline.**

The inclusion of silent variants provides a set of controls that serve as independently

measured replicates of the reference amino acid sequence. Like the missense- and nonsense-

variant sequences, in order to reduce contamination by miscalls, these silent mutations should

employ codons that are 2- or 3-nt different from the reference codon. Often, codons encoding an

amino acid differ from one another at the wobble base. As a result, many of the possible silent

variant codons are one nucleotide away from the template codon but these are less useful as

controls. Silent variants with 2- or 3-nt delta differences relative to their respective reference

codon results in silent variants comprising about 2% of the library. As demonstrated by the

bimodal split of phenotypes in *TP53* DBD library screens discussed earlier, ORF variant library

screens often produce a high rate of "hit" variants that deviate from the reference phenotype. As

a result, in contrast to many other common pooled screening situations, one cannot always rely

on the large portion of the library matching the reference phenotype and serving as a reference

baseline, making the silent codon change controls a particularly important reference-point for

interpreting the screen results. The silent changes LFC results are displayed in the row marked

by '**B**' in the heatmaps in **Figure 2a**.

In the *SMARCB1* library, we included nearly all possible silent variants, including 1-nt

delta silent variants. This case provides an example of the consequences of single-nucleotide

miscalls on the reference sequence that masquerade as 1-nt library variants. As expected, due to

the miscalls discussed earlier, the apparent abundance of the library's 1-nt delta silent variants is

inflated by those miscalls relative to the abundance of 2- and 3-nt delta silent variants

(**Supplementary Fig. 7a**). As a consequence, relative to 2- and 3-nt delta groups, the 1-nt delta

silent variants show an artificially suppressed log-fold change range when comparing among

samples, because every sample across which log-fold change comparisons are made has the same

contribution of counts due to miscalls of the reference sequence (**Supplementary Fig. 7b**),

equalizing the counts for these 1-nt variants across samples. In our current library designer

download (**Supplementary Table 2**), we avoid 1-nt delta silent variants and minimize 1-nt delta

missense and nonsense variants.

**Using next-generation sequencing (NGS) versus long-read sequencing (LRS) for DMS**

**deconvolution.**

Long-read sequencing technologies have the potential to capture all mutations present on each ORF-spanning DNA molecule. However, the technology that can capture the readout of full-length ORF sequences [17, 18] has significant limitations for variant detection due to low concordance rate of base calls [19]. For example, in a study by Wenger *et al* [20], circular consensus sequencing (CCS), the best of its kind, has a concordance rate of 99.772%. This is equivalent to a Q26 Phred score, producing one miscall per 439-base readout. This sequencing-error rate is higher than the error rate in commercial DNA oligonucleotide synthesis, the key contributor of the real errors in the libraries as discussed earlier. As a result, one may be forced to either throw out many reads that show secondary mutations or focus locally on the expected codon changes and ignore mutations in the distance.

It is encouraging that long-read sequencing technology is advancing at a steady and rapid pace, and its applications are actively explored. For example, based on circular consensus sequencing (CCS), by concatenating multiple DNA molecules, MAS-seq has shown the potential of increasing the CCS throughput by 15 to 30-fold while maintaining the sequencing phredQ [21]. In the recent work by Frank *et al* [22], long read sequencing was used to associate a set of unique external barcodes to each of ~7000 variants in a saturation variant library, so that the screen deconvolution work can be achieved by direct next-generation sequencing of the barcodes. However, there are several factors that create a high bar for long-read sequencing to match the results obtained with NGS paired-end reads of 150 bases each with a Phred score >Q30, which is comfortably achievable by many NGS platforms: (1) We have demonstrated that the errors in our

libraries are near the intended codon changes (**Fig. 3b**), and a 150-base read is adequate to detect

both the variant-defining sequences and library errors from DNA oligonucleotide synthesis. We

have shown that it is unnecessary to sequence an ORF variant into the regions synthesized by

high-fidelity DNA polymerase during library construction, because the errors introduced by

DNA polymerase in regions outside of a 100-base window around the intended codon changes

are rare enough to be inconsequential. ASMv1.0 output offers many choices to filter the results

in various ways, such as to retain only the reads that bear a library-designed codon change and

that effectively represents the variant's error-bearing region, i.e., the region that was chemically

synthesized. (2) We have demonstrated that PCR-introduced miscalls produce predominantly 1-

nt delta codon changes. Because we minimize the use of 1-nt delta codon variants, the artifactual

variants resulting from miscalls are mostly unexpected variants that were not part of the library

design. In this case, ignoring reads carrying only unexpected variations largely removes the

effects of sequencing miscalls. (3) Even with high base-quality threshold, NGS currently

produces adequate read yield for sequencing screen samples of variant libraries. ASMv1.0 allows

users to set adequate base quality threshold, for example, at Phred score Q37 with Illumina

Novaseq NGS.  With this filter, we can comfortably achieve <1 miscall per 5000-base NGS

readout. At this low miscall rate, >94% ($0.9998^{300}$) of 150-base read pairs used to call variants

are free of miscalls.

**Discussion**

Variant detection software ASMv1.0 implements in-depth data analyses that allowed

exploration of variant library composition and the fidelity of the screen sample deconvolution

process. The types and occurrence frequency of unintended species that arise during the variant

library production process and the overall variant purity of each designed library variant were

empirically assessed for multiple ORF libraries. Miscalls arising from PCR and sequencing in

the sample deconvolution process were identified and their contributions to different libraries

compared. In addition to this general characterization of libraries and the deconvolution process,

ASMv1.0 offers a means to filter the deconvolution data of screen result  for those variants

expected by design, or to identify high abundance variants independent of the library design.

The results of these analyses have informed the choice of strategies at multiple steps including

library design, screen setup, and data analysis. We believe that the analytical principle and screen

strategy presented here, along with our library designer, variant-calling software, and data

analysis tools will help enable the scientific community to fully utilize this technology in

research applications.

Although, as noted, producing the desired amino acid changes using 2 or 3-nt changes

versus the reference sequence is superior to using 1-nt changes versus the reference sequence, it

should be noted that the strong hits among the 1-nt delta variants may still score well, even with

some suppression of the apparent log-fold change results due to miscalls of reference sequence

that match the 1-nt variant. The effect of miscalls on 1-nt delta variant counts becomes less of an

issue for short ORFs (e.g., *KRAS*). As illustrated in **Supplementary Figure 6**, the longer the

ORF, the more copies of the reference codon will undergo PCR-based amplification and NGS,

the more copies of each actual designed 1-nt variant will be also read out, and consequently the

more counts of the 1-nt delta miscalls will occur per correct call of the 1-nt variant molecules

truly present. To better interpret any 1-nt delta codons that in a variant library, we recommend

adding the clonal template ORF as a control sample and processing it in parallel with the

screening samples. This control sample may be used as a count-correction factor (as shown in

**Fig. 4b**, right) or simply to flag those 1-nt delta variant sequence that occur at high rates due to

miscalls of the reference sequence, contaminating and therefore suppressing any fold-changes in

that variant between screen arms.

The recent establishment of an open-source platform, MaveDB, to enable variant

function data sharing, is a significant development [23] and is a good step towards more

widespread sharing of saturation mutagenesis data, reagents and resources. A well-designed

variant library should be useful for many different screens of that same genes, provided that the

choice of reference sequence upon which the design is based is satisfactory for each application.

For example, Boettcher *et al* [24], using the *TP53* site-saturated variant library created by

Giacomelli *et al* [9], quickly demonstrated that the dominant-negative activity of *TP53* missense

mutations in DNA binding domain was selected-for in myeloid malignancies. A shared inventory

of pooled gene variant libraries would be a powerful tool for variant-to-function mapping and

common practices for the design composition and readout of these libraries will enhance their

value.

**Conclusions**

Complex ORF variant libraries with all possible single-amino-acid substitutions are

powerful tools that enable one to exhaustively test all possible protein variants simultaneously in

a single experiment. Each experiment produces extensive dictionary of variants and functions.

This technology has high access barriers as it demands expertise across many disciplines, and the

costs of producing and utilizing the libraries are large. This report explores some factors that are

important for screening very large variant pools efficiently and presents a set of well-tested

practices and tools to conduct pooled saturation mutagenesis screens.

**Methods**

Experimental methods for the published data employed in this study are provided in the

original publications. However, methods for experimental steps for large-pool variant screening

that incorporate our current best practices, based on experience from over 20 variant screens

(**Supplementary Table 1**) are provided here along with the analysis and computational methods

and the data-driven evaluations of library compositions and variant screen error modes that are

presented in this study.

**ORF expression vector.** A lentiviral delivery expression vector that we have frequently used to

clone site-saturated variant libraries is pMT025 (**Supplementary Fig. 7**). This vector was used

for the libraries and screens that are re-analyzed in this report and numerous other ORF variant

libraries (**Supplementary Table 1**), It is suited for restriction/ligation cloning using

NheI/BamHI, or NheI/MluI restriction enzyme pairs. The vector has a puromycin resistance

selection marker driven by an SV40 promoter. After the cloning, the ORF expression is driven

by EF1α. To facilitate cloning, the ORF sequences are flanked with constant sequences carrying

the highlighted restriction enzyme sites for cloning:

GGTTCAAAGTTTTTTTCTTCCATTTCAGGTGTCGTGAG**GCTAGC**GCCACC [ORF with

ATG start and stop ]**GGATCC**CGGGACTAGT**ACGCGT**TAAGTCGACAATC

To note, two library construction techniques are described here, one in which the ORF

variant sequences originate as pooled double-stranded DNA fragments encompassing the entire

ORF, purchased from Twist Bioscience, and the other method in which the variants are

assembled, per 'tile,' from a pool of single-stranded DNA oligonucleotides to create a plasmid

DNA pool of the variants in a pUC57 vector. While the resulting variant expression libraries, the

screening and data processing of these libraries are the same, these two methods command some

differences in library design and library cloning.

**Template ORF plasmid**. An ORF variant library is built from a common template sequence to

which mutations are introduced. This reference sequence may be some version of the "wild-type"

coding sequence of a gene, or it may itself already possess known functional mutations to which

additional mutations are added to form the library. In this paper, we call this template sequence

the "reference sequence". It is important at the outset of library design to ensure that the chosen

reference sequence will be fully compatible with all the requirements of library construction and

screening. For example, in the cases where a preliminary choice of ORF reference sequence

contains restriction sites needed for the library cloning, the reference sequence needs to be

modified with silent mutations to remove these sites. If the screening assay has been developed

using expression of the initial reference sequence, then any changes to this sequence necessitate

ensuring that these changes have no impact on the reference sequence phenotype and assay.  In

general, we recommend that, prior to library construction, the screening assay be fully vetted

using the *exact* reference sequence and expression vector that the library will employ, even if

that requires repeating assay development experiments that were previously performed using a

similar vector or ORF sequence. It is equally important that a clonal plasmid of the template

ORF is cloned into the vector, made into virus, tested for transduction, selection condition and

genomic DNA prep. Finally, the genomic DNA is tested for PCR condition to be used in future

screen deconvolution. Twist Bioscience uses this clonal plasmid as the PCR template to produce

the mutagenesis library.

Two library construction techniques are described here, one in which the ORF variant sequences

originate as pooled linear oligonucleotides encompassing the entire ORF, purchased from Twist

Biosciences, and another method in which the variants are assembled from 90-nt tiles using a

method described below to create a pool of the variants in a pUC57 plasmid.

**Library design**. A saturation mutagenesis designer written in R is available (see the links listed

in **Supplementary Table 2**). The reference ORF sequence is first checked to ensure that the

restriction sites to be used in cloning are excluded, and otherwise the sequence must be modified

with silent changes to eliminate any occurrences of the cloning sites. NheI/BamHI and

NheI/MluI cloning sites have been successfully employed for numerous ORF libraries. The

designer algorithm steps through from N-terminus to C-terminus of the template ORF one amino

acid position at a time to select variant codons. For a given template codon there are 63 possible

variant codons from which to choose for each amino acid position. These 63 codons are divided

into two bins: 54 codons that have 2- or 3-nucleotide differences relative to the template codon

(2-3-nt delta), and 9 codons with a single-nucleotide difference (1-nt delta). The codons that

would violate the restriction sites exclusion rule are removed from both bins. In designing

missense or nonsense variants, we first select from the 2-3-nt delta codon bin, and then in cases

for which no 2 or 3-nt delta option is available, from the 1-nt delta codon bin. If within a bin

there are multiple codon choices available for an amino acid variant, the codon is picked by

sampling the available codons with probability weightings according to the codon usage

frequency of the organism (*Homo sapiens* in all our experiments to date,

http://www.kazusa.or.jp/codon/cgi-bin/showcodon.cgi?species=9606). For silent codon changes

we pick only from 2-3-nt delta bin and do not include a silent change variant in the library for

those residues for which no 2 or 3-nt delta option is available. In the end, each of the designed

variant codons substitutes for its corresponding template codon in the full-length reference ORF

as one member of the library.

**Mutagenized ORF Library cloning**. Saturation mutagenesis library production requires that

variant-defining codons be encoded in DNA oligo strands, which will be chemically synthesized.

For the MITE method, once the 150-nt DNA oligo strands bearing the mutagenized codons are

produced, they are first amplified by PCR and then cloned into a high copy number entry vector,

pUC57. For Twist Bioscience method, the DNA oligo strands serve as PCR primers to introduce

codon changes into full-length ORF template to produce a pool of properly flanked and variant-

carrying full-length ORFs that are ready for cloning into pMT025 ORF expression vector.

**MITE method**: The region of the open reading frame to be mutated is partitioned into

90-base tiles. Oligos representing all mutations in the space of a tile were synthesized. Each 90

base tile is flanked by 30 base sequences that are complementary to the reference ORF. Oligos

for adjacent tiles should be synthesized on separate arrays. Oligo tile pools were amplified by

emulsion PCR using primers to the 30 base flanking regions and purified on a 2% agarose gel.

Entry vector cloning is performed on a per tile basis by linearizing the entry vector at the

appropriate tile position using Phusion polymerase (New England Biolabs) and primers to the

regions flanking the tiles. The linearized vectors were purified on 1% agarose gel and DpnI

treated. The DpnI-treated linear plasmid backbones were then mixed with the relevant PCR

amplified tile and assembled by *in vitro* recombination with the NEBuilder HiFi DNA Assembly

Master Mix (New England BioLabs). The assembly reactions were purified, electroporated into

TG1 *E. coli* cells (Lucigen), and recovered for one hour at 37˚C in Recovery Media (Lucigen).

Aliquots from the transformations were used to inoculate overnight cultures of LB containing 25

µg/mL of Zeocin (ThermoFisher). Cells were harvested by centrifugation and plasmid DNA was

isolated using the QIAGEN Midiprep Plus kit (QIAGEN). Plasmid pools, each corresponding to

a tile, were verified by sequencing Nextera XT (Illumina). A pool of all pUC57-tile libraries was

made with equal weight per variant. This pool was subjected to restriction digest (e.g., NheI and

BamHI). After purification by 1% agarose gel, the excised linear fragment library is ready for

cloning ligation with an expression vector that was pre-processed with the compatible restriction

enzyme pair.

**Twist Bioscience method**: The full-length variant pool made by Twist Bioscience was

delivered as linear fragments of the full-length ORF flanked with adapters for cloning. This

fragment library was first digested with NheI and BamHI overnight, and then underwent ligation

with pMT025 vector pre-processed with the same restriction enzyme pair.

**Cloning ligation:** For either cloning method, the ligation was carried out with 5:1 insert-

to-vector molar ratio, using T7 DNA ligase at room temperature for 2 hours. The ligation was

cleaned up with isopropanol precipitation. The resulting DNA pellet was used to transform Stbl4

bacterial cells. Plasmid DNA (pDNA) was extracted from the harvested colonies using QIAGEN

Maxi Prep Kits. The resulting pDNA library was sequenced via the Illumina Nextera XT

platform to determine the distribution of variants within the library.

**Virus preparation and titer measurement.** Lentivirus was produced and quantified by the

Broad Institute Genetic Perturbation Platform (GPP). The detailed protocol is available at

http://www.broadinstitute.org/rnai/public/resources/protocols. Briefly, 293T viral packaging

cells were transfected using TransIT-LT1 transfection reagent (Mirus Bio) with the pDNA

library, a packaging plasmid containing *gag*, *pol* and *rev* genes (psPAX2, Addgene), and an

envelope plasmid containing VSV-G (pMD2.G, Addgene). Media was changed 6-8 hours after

transfection, and virus was harvested 24 hours thereafter. The virus titer was measured by

infecting 3000 A549 cells per well in 96-w cell culture plate with a series of diluted standard

virus of known titer for a standard curve, and a series of diluted library virus of unknown titer to

find data points that lie in the linear range of the standard curve. Twenty-four hours post

infection, the cells were subjected to puromycin selection for 2 days. The surviving cells were

quantified with alamarBlue.

**Determine cell numbers required for library representation.** The number of cells transduced

per replicate and the number of cells seeded per replicate at each passage was typically an

average of 1000 cells per variant in the library pool. This minimum cell number was preserved

during cell culture passaging. For positive selection screens, a majority of the cell population is

depleted and only the subset of cells that carry survival-conferring variants will survive. In these

cases, adequate cell number representation of the full library must be maintained until the

selection pressure is applied, but as the selection pressure takes effect, the cell numbers are

reduced, and the resulting smaller cell population, enriched for survival-conferring variants is

collected from these selected arms at the end of the screen. For negative selection screens, most

of the library will be present at the end of the screen, and the end-point samples should comprise

at least 1000 cells per variant (i.e., per library member).

**Genomic DNA preparation.** Upon the completion of the screen, cell pellets were collected.

Genomic DNA was extracted from cell pellets using the Machery Nagel NucleoSpin Blood Kit

(Midi, Clontech Cat. #740954.20; XL Maxi, Clontech Cat. #740950). The detailed protocol is

available at http://www.broadinstitute.org/rnai/public/resources/protocols.

**Design PCR primers**. The ORF sequence is amplified from the genomic DNA by PCR. The

PCR products will be randomly sheared with transposon, index labeled, and sequenced with

next-generation sequencing technology. Each sequenced fragment is generated by 2 transposon

insertion events so that both ends of the fragment are tagged with a transposon. Therefore, the

ends of the initial gDNA PCR product will be less frequently represented in the NGS library and

will have shallow coverage. For this reason, we advise that the PCR primers be designed include

an extra ~100 bp of sequence at each end beyond the region of ORF that is mutated in the library.

For the pMT025 vector carrying libraries whose variants span entire ORF insert, we use these 2

primers:

Forward: 5'-ATTCTCCTTGGAATTTGCCCTT-3'

Reverse: 5'-CATAGCGTAAAAGGAGCAACA-3'

**Determine the number of PCR reactions per sample.** For positive screen end-point samples,

the number of cells that need to go through ORF-extracting PCR is determined in such a way

that for each library variant, gDNA template from 500 or more cells will be amplified by PCR.

For reference samples and negative selection screen end-point samples, the cells/variant ratio

should be maintained at 1000 or more. PCR is sensitive to the amount of gDNA input and prior

to screening, these conditions must be optimized. For example, in 50ul PCR reactions, we test

different amounts of gDNA, using, for example, a series of 250 ng, 500 ng, 1 µg, 2 µg. This

titration will determine the optimal concentration of gDNA that can be used to produce robust

PCR products. The gDNA loaded into each PCR reaction should not exceed the amount that can

be successfully amplified for the reaction conditions and scale employed. Diploid human cell

lines have roughly 6.6 pg DNA per cell. Here is the formula to determine how many PCR

reactions are needed for a sample, again making sure not to exceed the maximum ng of gDNA

per well that can be amplified:

$$\text{PCR reactions per sample} = 6.6 * \frac{total\ number\ of\ variants\ in\ the\ library}{ng\ of\ gDNA\ per\ PCR\ well}$$

**ORF amplification from gDNA by PCR.** PCR reactions were set up in 96-well plates

according to the optimized PCR condition and number of PCR reactions per sample calculated

above. The volume of each PCR reaction was 50 µL and contained an optimized amount of

gDNA. Q5 DNA polymerase (New England Biolabs) was used as the DNA polymerase. All PCR

reactions for each gDNA sample were pooled, concentrated with a PCR cleanup kit (QIAGEN),

and separated by gel electrophoresis. Bands of the expected size were excised, and DNA was

purified, first using a QIAquick kit (QIAGEN), and then an AMPure XP kit (Beckman Coulter).

**Nextera sequencing.** Nextera reactions were performed according to the Illumina Nextera XT

protocol. Each sample was split into 4 Nextera reactions, each with 1 ng of purified ORF PCR

products. Each reaction was indexed with unique i7/i5 index pairs. After the limited-cycle PCR

step, the Nextera reactions were purified with AMPure XP kit. All samples were then pooled and

sequenced using an Illumina sequencing platform chosen among Miseq, Hiseq, Nextseq, or

NovaSeq, according to the number of reads needed as calculated below.

**Determine number of reads needed for the screen.** We need to sequence the Nextera-

produced fragment molecules for every variant. The number of total reads required depends on

ORF length, total number of variants in the library, read length, read quality etc. The minimum

number of read required can be estimated as follows:

For negative selection screen samples, or reference samples:

$$\text{Reads per sample} = 7000 * \frac{(ORF\ length\ in\ NT)^{\wedge}2}{read\ length * read\ assembly\ rate}$$

632            Where '*read length*' is typically 300, as we use paired-end sequencing reads, 150

633    bases per read, and '*read assembly rate*' is in range of 0.7-0.9.

635    For a positive selection screen sample:

$$\text{Reads per sample} = 3500 * \frac{(ORF\ length\ in\ NT)^{\wedge}2}{read\ length * read\ assembly\ rate}$$

**Determine the optimal parameters for variant-calling software ASMv1.0.** NGS is performed

on transposon-sheared fragments which are not uniform in size. The Nextera conditions as

recommended typically achieve an average fragment size of approximately 140 base pairs, data

not shown). For fragments longer than 2x read length (e.g., 2 x 150 = 300 bases), the pair of

reads are disjoint (non-overlapping); for fragments shorter than the read length (e.g., 150 bases),

the reads are fully overlapping and each extends into the transposon adaptor located in the

opposite end; for fragments sized between 1x and 2x read length (e.g., 150-300 bases), the pair

of reads overlap but do not extend into the transposon sequences. The variant calling software

first aligns reads to the reference sequence. The transposon tails are then clipped off at the

coordinate determined by the start of its paired read. It then applies the ***min-q*** (minimum base

quality) filter to trim the end of the aligned reads to ensure there are at least 15 bases from the

trimmed end inwards with base quality above the min-q parameter. The ***min-read-length*** filter

admits the trimmed reads that meet the selected length threshold. All failed reads go into the

low-q bin. Next, the filter of ***min-flank-length*** is applied that requires reference bases at end of

the trimmed reads. A min-flank-length of 2 or more bases excludes cases in which an altered

codon is split between 2 Nextera fragments. It also prevents erroneous variant calls due to reads

that end with transposon sequences for which transposon clipping failed to work due to a lack of

paired read overlap. Finally, a variant is called only when the underlying base changes are above

the min-q threshold.

To tune our software parameters, we used the sequencing data of a saturation mutagenesis

plasmid DNA library for *PDE3A* gene. This library was built by mutagenizing a region of 1425

base pairs, using a variant oligonucleotide pool produced by Twist Bioscience. The errors in the

library are concentrated in within 30 bases of a planned codon change (**Fig. 3**). These errors can

be explained by mutagenesis primer synthesis errors. The codon changes are located in the

middle of the mutagenesis primers, sandwiched between two 20-30 base pair flanks. In next-

generation sequencing, the transposon sheared fragments are, on average, 140 base pairs, similar

to the sequencing read length. Characteristic to the way how the variant libraries were

synthesized, the close proximity of codon modifications, including the planned and the

unintended, may indicate that these codon modifications reside on the same transposon-sheared

fragment and on the same sequencing read. Therefore, each library ORF, after transposon

shearing, produces one variant-bearing fragment; the rest of the fragments are all wild-type. That

is, if we ignore the miscalls introduced by PCR, the wild-type/variant read ratio of this library

should be: 1425/140 -1 = 9.2. This ratio, 9.2, represents the highest ratio we could observe, as

any PCR miscalls would only shift this ratio lower.

In determining the PCR miscall rate, we processed a clonal plasmid of pMT025-PDE3A. The

sequencing of this clonal plasmid DNA, which had undergone the same PCR process as the

saturation mutagenesis plasmid DNA library, showed error rate of 0.00037 miscalls/sequenced

nucleotide.

Start with miscall-free wild-type/variant read count ratio 9.2:1, At this miscalls rate, one expects

fewer reference reads versus variant reads, namely (9.2*140*0.00037)=0.48 of the expected

reference reads incur miscalls and appear as variants, predicting a wild-type/variant read count

ratio:

$$(9.2 - 0.48)/(1 + 0.48) = 5.9$$

By processing the same plasmid DNA library sequencing data set with various parameters, we

were able to identify the optimal parameters that produce a wild-type/variant ratio of 5.9, and

yield reasonable output of both wild-type and variants: min-q = 30, min-read-length = 40, min-

flank-length = 2.

**NGS Data processing and analysis:** The data from the NGS pipeline are in .fastq format. Each

NGS cluster produced 2 long reads (150 bases each), and 2 index reads (each 8 bases long). The

2 indexes were used to label the screen samples, so that a large number of samples can be pooled

and sequenced together. Site-saturated variant library screen data file sizes are typically 4-5TB.

Generous computer storage is needed to process the data, especially as the reads go through the

filtering process (**Supplementary Fig. 2**) prior to variant calling.

<u>Read filtering:</u> The reads from an NGS fragment are trimmed (1) by base quality, (2) by

removing non-ORF transposon sequences, (3) by requiring that mutated nucleotides are flanked

by, at minimum, the number of reference-matching bases that is specified by the min-flank-

length parameter, and (4) by requiring the trimmed reads be no shorter than the min-read-length

(see **Supplementary Fig. 2**).

<u>Variant calling</u>**:** There are two versions of variant calling software, ORFCallv1.0, the earlier

version; and ASMv1.0, the current version. The reads that have passed the filtering step are

processed differently by ORFCallv1.0 and ASMv1.0:

i)        ORFCallv1.0 (an earlier version software; Giacomelli *et al*[9]): Variants are tallied in

702                the context of a single codon space. Without regard of the association of the multiple

703                changes physically in a single molecule, ORFCallv1.0 essentially scans from N- to C-

termini, one amino acid position at a time, and sums up the occurrence of codons or

amino acids at each position.

ii)    ASMv1.0**:** This is a molecule-centric approach that is more sophisticated than the

approach of ORFCallv1.0. The NGS dataset is analyzed in its entirety, with the

following new capabilities: (1) The trimmed reads are evaluated full-length; that is,

variants are called in the context of entire read (or read pair) and by nucleotides, not

amino acids. The programmed variant is called when 2 conditions are met: (i) a

library-designed nucleotide change are detected (in 1-3 nucleotides within a codon

position), and (ii) there are zero *additional* nucleotide variations throughout the entire

read or read pair. (2) Indel detection: This feature was lacking in ORFCallv1.0. This

new feature of ASMv1.0 allows us to process libraries that include indel variants.

In ASMv1.0, mutations in a molecule are depicted at the nucleotide level by position

along the entire pre-framentation PCR product. They are annotated by listing

positions of change followed by the change itself. For example, 'a change from G to

A at position 1140' is described as '**1140:G>A';** 'a deletion of an A at position 1141'

is '**1141:A>-**'; 'an insertion of a T after position 1142' is '**1143:->T**'.

Also, with ASMv1.0, a host of annotations formatted as a datafile can be derived

from the variant descriptions.  For example, a *TP53* variant with description

'**1140:G>A, 1141:A>C, 1142:G>C**' is annotated as (a pair of brackets marks an

annotation): [E343T] [Intended] [one codon change] [343:GAG>ACC] [missense]

[substitution].  Another *TP53* variant with description '**1140:G>A, 1141:A>C,**

**1142:G>C, 1145:G>T**' is: [E343T] [Unintended] [two codon changes]

[343:GAG>ACC, 344:CTG>CTT] [substitution] [missense/nonsense] etc. These two

variants encode the same amino-acid variant. The first variant is an intended variant

that matches a designed nucleotide-variant of the library. The second variant is the

same amino acid call as the first variant, but it is annotated, rightfully, as an

unintended variant carrying an extra nucleotide change 4-nucleotide away from the

intended codon change - this extra nucleotide change results in a silent codon change.

Parsing data into a form ready for hit calling**:** The output files from the above variant-calling

software are still typically quite large in size (over 100MB), large in numbers of files, and not

readily useable. These output files need to be parsed, aggregated and annotated. For this purpose,

we built a parser called ASMv1.0_Parser, which was written in R, and is available for download

(see the links listed in **Supplementary Table 2**). The Parser Pak consists of a 'readme' file

detailing the instructions, and R codes and training data sets.  Through the parser, all data files

are organized into a single .csv file, one column per screen sample. These .csv files include a

short version that is a subset of the planned variants only, and a complete version that includes

all called variants, including planned/unplanned, insertion/deletion/substitution, etc. In addition,

we also report the ORFCallv1.0 version of data.

Scaling: We use read counts from the NGS data as the quantitative measurements of variants.

Reads that entirely match the reference sequence are not informative, since they are presumably

all reflect the unmutated, uninteresting parts of an amplified ORF molecule (the majority) that

did NOT contain the mutation that defined that specific ORF variant. At any amino acid position,

the coverage is given by the count of all variant sequences aligned to that position plus the count

of the wildtype reads that align to that position. The coverage should be consistent throughout

the ORF. But that is not actually the case due to the transposon insertion bias that results in

unevenness of total read coverage (**Supplementary Fig. 9**). To determine the relative abundance

among variants, as in **Supplementary Figure 10**, the raw counts should first be scaled to the

fraction of the total read coverage across the codon position. However, in cases that compare the

variant abundance change between two samples, such as computing fold-changes in screening

samples, scaling the raw counts may not be necessary, as the positional raw count variations

caused by transposon insertion bias will cancel out between two samples when computing fold-

changes (**Supplementary Fig. 11**).

**Acknowledgements**

We thank our collaborators of saturation mutagenesis projects, Alice Berger, Jesse

Boehm, Ben Ebert, Laura Evans, Eejung Kim, Shengwu Liu, Julie-Aurore Losman, Huong Ly,

Peter Miller, Utthara Nayar, and Nikhil Wagle for demonstrating the diverse utilities of the

technology, Broad Institute Genomics Platform for providing NGS support, Nancy Tran and

Cong Zhu for the early work on saturation mutagenesis method, Stephanie Milczarek, Kirsten

Milenkowic, Michael Zagieboylo, and Jonah Schwartz for construction of some of the variant

libraries referenced in this study, Leon Yang for critical reading of the manuscript, and to all the

members of Broad's Genetic Perturbation Platform for their help and support. This work was

supported by NCI U01 CA176058 (WCH), the Mark Foundation (WCH), American Cancer

Society Professorship (MM), American Cancer Society Mentored Research Scholar Grant

MRSG-18-202-01 (ALH), Department of Defense W81XWH-18-PRCRP-CDA (ALH), Team

Lick Cancer (ALH), Unravel Foundation (ALH), and Wong Family Award (ALH), and the

Functional Genomics Consortium (D.E.R.).

**Author Contributions**

X.Y., D.E.R., A.L.H., A.O.G., and T.S. wrote the manuscript. W.C.H., M.M., H.G.,

A.J.A., R.E.L., N.P, M.M, A.B., and F.P. reviewed and edited the manuscript. A.L.H., X.Y., T.S

and D.E.R. designed the new variant-calling software ASMv1.0. T.S. wrote the variant-calling

software ORFCallv1.0 and ASMv1.0. X.Y., A.L.H., D.E.R., and T.H. optimized the run

parameters of ASMv1.0. X.Y. wrote R codes to design site-saturated variant libraries and to

parse the output of the variant-calling software. X.Y. conducted data analysis with input from

D.E.R., A.L.H., A.O.G., and assistance from B.P.L.  D.A and T.G. adapted the variant-calling

software into a saturation mutagenesis data pipeline. X.Y., A.L.H., A.O.G, R.E.L., B.P.L., B.F,

H.K., E.S., F.P., and L.S. designed and conducted wet-lab experiments. All authors contributed

to and approved the manuscript.

**Competing Interests**

W.C.H. is a consultant for ThermoFisher, Solasta, MPM Capital, iTeos, Jubilant

Therapeutics, Tyra Therapeutics, RAPPTA Therapeutics, Frontier Medicines, KSQ Therapeutics

and Paraxel. A.J.A has consulted for Oncorus, Inc., Arrakis Therapeutics, and Merck & Co., Inc,

and has research funding from Mirati Therapeutics, Syros, Deerfield, Inc., and Novo Ventures

that is unrelated to this work. A.O.G. is a share and option holder of 10X Genomics. D.E.R.

receives research funding from members of the Functional Genomics Consortium (Abbvie, BMS,

Jannsen, Merck, Vir), and is a director of Addgene, Inc.

**References**

1.    Hughes RA, Ellington AD: **Synthetic DNA Synthesis and Assembly: Putting the**
**Synthetic in Synthetic Biology.** *Cold Spring Harbor perspectives in biology* 2017,
**9:**a023812.
2.    Heather JM, Chain B: **The sequence of sequencers: The history of sequencing DNA.**
*Genomics* 2016, **107:**1-8.
3.    Fowler DM, Araya CL, Fleishman SJ, Kellogg EH, Stephany JJ, Baker D, Fields S:
**High-resolution mapping of protein sequence-function relationships.** *Nature Methods*
2010, **7:**741-746.
4.    Fowler DM SJ, Fields S. : **Measuring the activity of protein variants on a large scale**
**using deep mutational scanning. Nature Protocols.** 2014.
5.    Kotler E, Shani O, Goldfeld G, Lotan-Pompan M, Tarcic O, Gershoni A, Hopf TA,
Marks DS, Oren M, Segal E: **A Systematic p53 Mutation Library Links Differential**
**Functional Impact to Cancer Mutation Pattern and Evolutionary Conservation.** *Mol*
*Cell* 2018, **71:**873.
6.    Mighell TL, Evans-Dutson S, O'Roak BJ: **A Saturation Mutagenesis Approach to**
**Understanding PTEN Lipid Phosphatase Activity and Genotype-Phenotype**
**Relationships.** *Am J Hum Genet* 2018, **102:**943-955.
7.    Weile J, Kishore N, Sun S, Maaieh R, Verby M, Li RJ, Fotiadou I, Kitaygorodsky J, Wu
YZ, Holenstein A, et al: **Shifting landscapes of human MTHFR missense-variant**
**effects.** *American Journal of Human Genetics* 2021, **108:**1283-1300.
8.    Awad MM, Liu S, Rybkin, II, Arbour KC, Dilly J, Zhu VW, Johnson ML, Heist RS, Patil
T, Riely GJ, et al: **Acquired Resistance to KRAS(G12C) Inhibition in Cancer.** *N Engl*
*J Med* 2021, **384:**2382-2393.
9.    Giacomelli AO, Yang X, Lintner RE, McFarland JM, Duby M, Kim J, Howard TP,
Takeda DY, Ly SH, Kim E, et al: **Mutational processes shape the landscape of TP53**
**mutations in human cancer.** *Nature Genetics* 2018, **50:**1381-1387.
10.   Garvie CW, Wu X, Papanastasiou M, Lee S, Fuller J, Schnitzler GR, Horner SW, Baker
A, Zhang T, Mullahoo JP, et al: **Structure of PDE3A-SLFN12 complex reveals**
**requirements for activation of SLFN12 RNase.** *Nat Commun* 2021, **12:**4375.
11.   Kwon JJ, Hajian B, Bian Y, Young LC, Amor AJ, Fuller JR, Fraley CV, Sykes AM, So J,
Pan J, et al: **Structure-function analysis of the SHOC2-MRAS-PP1C**
**holophosphatase complex.** *Nature* 2022, **609:**408-415.
12.   Miller PG, Sathappa M, Moroco JA, Jiang W, Qian Y, Iqbal S, Guo Q, Giacomelli AO,
Shaw S, Vernier C, et al: **Allosteric inhibition of PPM1D serine/threonine**
**phosphatase via an altered conformational state.** *Nat Commun* 2022, **13:**3778.
13.   Persky NS, Hernandez D, Do Carmo M, Brenan L, Cohen O, Kitajima S, Nayar U,
Walker A, Pantel S, Lee Y, et al: **Defining the landscape of ATP-competitive inhibitor**
**resistance residues in protein kinases.** *Nat Struct Mol Biol* 2020, **27:**92-104.
14.   Majithia AR, Tsuda B, Agostini M, Gnanapradeepan K, Rice R, Peloso G, Patel KA,
Zhang X, Broekema MF, Patterson N, et al: **Prospective functional classification of all**
**possible missense variants in PPARG.** *Nat Genet* 2016, **48:**1570-1575.

15.    Melnikov A, Rogov P, Wang L, Gnirke A, Mikkelsen TS: **Comprehensive mutational**
**scanning of a kinase in vivo reveals substrate-dependent fitness landscapes.** *Nucleic*
*Acids Res* 2014, **42:**e112.
16.    Masaki Y, Onishi Y, Seio K: **Quantification of synthetic errors during chemical**
**synthesis of DNA and its suppression by non-canonical nucleosides.** *Sci Rep* 2022,
**12:**12095.
17.    Legnini I, Alles J, Karaiskos N, Ayoub S, Rajewsky N: **FLAM-seq: full-length mRNA**
**sequencing reveals principles of poly(A) tail length control.** *Nature Methods* 2019,
**16:**879-886.
18.    Starr TN, Greaney AJ, Hilton SK, Ellis D, Crawford KHD, Dingens AS, Navarro MJ,
Bowen JE, Tortorici MA, Walls AC, et al: **Deep Mutational Scanning of SARS-CoV-2**
**Receptor Binding Domain Reveals Constraints on Folding and ACE2 Binding.** *Cell*
2020, **182:**1295-1310.e1220.
19.    Marx V: **Long road to long-read assembly.** *Nat Methods* 2021, **18:**125-129.
20.    Wenger AM, Peluso P, Rowell WJ, Chang P-C, Hall RJ, Concepcion GT, Ebler J,
Fungtammasan A, Kolesnikov A, Olson ND, et al: **Accurate circular consensus long-**
**read sequencing improves variant detection and assembly of a human genome.**
*Nature Biotechnology* 2019, **37:**1155-1162.
21.    Al'Khafaji AM, Smith JT, Garimella KV, Babadi M, Sade-Feldman M, Gatzen M,
Sarkizova S, Schwartz MA, Popic V, Blaum EM, et al: **High-throughput RNA isoform**
**sequencing using programmable cDNA concatenation.** *bioRxiv*
2021**:**2021.2010.2001.462818.
22.    Frank F, Keen MM, Rao A, Bassit L, Liu X, Bowers HB, Patel AB, Cato ML, Sullivan
JA, Greenleaf M, et al: **Deep mutational scanning identifies SARS-CoV-2**
**Nucleocapsid escape mutations of currently available rapid antigen tests.** *Cell* 2022,
**185:**3603-3616 e3613.
23.    Esposito D, Weile J, Shendure J, Starita LM, Papenfuss AT, Roth FP, Fowler DM, Rubin
AF: **MaveDB: an open-source platform to distribute and interpret data from**
**multiplexed assays of variant effect.** *Genome biology* 2019, **20:**223-223.
24.    Boettcher S, Miller PG, Sharma R, McConkey M, Leventhal M, Krivtsov AV,
Giacomelli AO, Wong W, Kim J, Chao S, et al: **A dominant-negative effect drives**
**selection of <em>TP53</em> missense mutations in myeloid malignancies.** *Science*
2019, **365:**599.
871

**Figure legends**

**Figure 1. Pooled screens with ORF variant libraries versus sgRNA libraries.** For either

screen type, a plasmid pool of genetic perturbation agents is first packaged into lentiviral

particles. Cells are transduced with the pooled virus at an infection rate of 30-50% to ensure that

most of the infected cells get a single genetic perturbation. The perturbed cell population is

divided into selection and reference arms. As the screen completes, the cell populations are

harvested, and the collected screening samples are processed to quantify the relative abundance

of the library perturbations for each sample, i.e., "deconvolution." The relative enrichment or

depletion of each perturbation in treatment versus reference condition is assessed differently for

sgRNA versus ORF screens. **(a) A typical pooled screen using sgRNA libraries.** From the

gDNA of collected cells, the guide sequence itself is PCR-amplified to serve as a barcode of the

perturbation that was introduced to each cell. The sgRNA barcodes are short enough to be read

straight out by a short NGS read. **(b) ORF variant library screens.** For an ORF variant library,

the entire ORF is amplified from gDNA to serve as a 'barcode' of the variant. The distinguishing

feature among barcodes is the mutation of a single codon position located anywhere along the

entire ORF. The following steps serve to quantify the relative abundance of variants: (1) PCR-

amplification to extract ORFs from genomic DNA, (2) random shearing of the full-length ORF,

(3) next-generation sequencing of the resulting fragments, (4) application of variant detecting

tools to identify and tally the reads. The NGS reads consist of variant signature-carrying reads

(shown as red arrows) and wild-type reads (black arrows). The variant-calling software processes

these NGS reads to determine the relative abundance of each variant and enables downstream

analyses that map variants to phenotype strength, gauged by log-fold changes (LFCs).

**Figure 2. Reanalysis of *TP53* saturation mutagenesis screen using ASMv1.0.** The

Giacomelli *et al* [9] p53[NULL] A549 cell/nutlin-3 screen enriches *TP53* variants that exhibit loss-

of-function (LOF), and depletes wild-type-like variants. (**a) LFC heatmaps depicting**

**enrichment or depletion of all variants in this library.** Each variant is placed in a 2-

dimensional grid, where the horizontal dimension specifies amino acid position, and the vertical

dimension specifies the amino acid substitution. 'X' designates nonsense mutations, and 'B'

designates silent codon changes. The heat index is log-fold change (LFC) between treatment and

reference samples. The same original NGS screen data were analyzed using ORFCallv1.0 (top)

and ASMv1.0 (middle), yielding mostly similar results. The small subset of variants for which

the two analyses give different results are highlighted in the third heatmap (bottom) that shows

delta(LFC)=(LFC$^{ASM}$ – LFC$^{ORFCall}$) as the heat index. (**b**) **Scatter plot comparing LFC values**

**of all variants as assessed by the two versions of software.** The color index of each data point

is the purity of the corresponding variant. (**c**) The percentile of "Variant purity", calculated by

ASMv1.0, was used to group variants into five bins. For each bin of variants the Pearson

correlation coefficient between the ORFCallv1.0 and ASMv1.0-computed LFCs is plotted on the

y-axis. (**d,e**) **Testing ASMv1.0 using the published data by Kotler *et al* [5].** Despite the noted

differences, the Kotler screen of p53$^{NULL}$H1299 cells with a library covering the *TP53* DNA

binding domain provides an intriguing comparison to the larger Giacomelli pooled screen. We

first took a 2964-variant subset of Giacomelli library to match the *TP53* amino acid variants of

the Kotler library. Of these 2964 shared variants, the 3-way comparisons are shown, with

ORFCall and ASM scores as x- and y-axis, Kotler score as the color index (**d**, **left**). We then

identified 76 variants that were scored most differently in the Giacomelli screen processed by the

two versions of software (**d, right**). We finally compared the 76 Kotler scores against the

corresponding Giacomelli LFC scores called by ORFCallv1.0 (**e, left**) and ASMv1.0 (**e, right**).

**Figure 3. Unexpected mutations in the site-saturated variant libraries are rare and tend to**

**co-localize with planned mutations. (a) Breakdown of the planned variants versus those**

**with additional mutations.** ASMv1.0 output records information that allows us to identify the

number of unintended nucleotide changes that are associated to planned codon changes. The

number of additional unexpected nucleotide changes were counted and the abundance

breakdown in fraction is shown in the bar-chart. The pure variants are presented by the bar with

'0' extra nucleotide changes.  (**b**) **Unexpected mutations are concentrated in the vicinity of a**

**planned change.** Again, the ASMv1.0 output contains information enables us to tally any

unexpected nucleotide changes as a function of their distance, in nucleotides, from the planned

codon change.  For all called variants that contained a planned codon change and additional

changes, we tallied the relative abundance (counts scaled against reference counts, i.e., read

coverage) and summed the relative abundance as a function of the distance between the error and

the planned codon change, The error distribution profile with distance to a planned codon variant

is shown for two libraries made with different technologies, *TP53* library by MITE and *PDE3A*

by Twist Bioscience. The MITE technology uses synthesized DNA strands of 150 bases long, of

which 90 bases (called TILE) go into the final products, while Twist Bioscience introduced

desired codon changes through oligonucleotide primers with the intended codon changes placed

in the middle and flanked with 15-25 bases on either side. The major errors are concentrated

within ~100 bases (MITE method) or ~30 bases (Twist Bioscience method) from a planned

codon change.

**Figure 4. Miscalls due to PCR and sequencing errors and their effects on variant**

**quantification and LFC. (a)** A clonal *TP53* reference plasmid (i.e., all sequences are wild-type)

was processed as if it were a screening sample, subjected to PCR amplification, NGS sample

preparation and NGS sequencing read out. As expected, the base call at each position is

predominantly, but not absolutely, the reference nucleotide. The non-reference calls represent

artifactual variants that were not actually present in the sample being measured. The heatmap on

the left shows tallied counts of called nucleotides at each nucleotide position. The heatmap on

the right depicts the rate of miscalls; blank spots in the heatmap are the wild-type nucleotides. (**b**)

**Abundance distributions of 3 codon types in a reference screening sample.** *TP53* variants

were grouped by the number of differing nucleotides between the variant signature codon and the

reference codon; the 3 groups are called 1-nt-delta, 2-nt-delta and 3-nt-delta variants. The

miscalls resulting from PCR/NGS errors inflate the abundance measurement of variants whose

variant-defining codon is 1-nt away from the reference codon. In the *TP53* library screen data,

predominately affected are variants whose signature codon is 1-nt (but not 2- or 3-nt) away from

the reference codon. As a result, the variants of 1-nt delta codons as a group appear more

abundant than groups of 2- or 3-nt delta codons (**left**). The artifactual 'variant' abundance

resulting from PCR/NGS errors can be corrected with abundance adjustment by subtracting the

miscall rate as determined by a control experiment using a clonal sample from every screening

sample (**right**). The variant abundance measurements of the 2-nt and 3-nt delta codons are not

affected by the miscalls and the corrections do not appreciably alter the abundance distributions

of these 2 codon types. (**c**) **Suppression of fold-changes in 1-nt delta codon variants.** Three

scatter plots, each for one of the three codon types, depict LFC versus variants ordered by codon

positions. A density plot on the right depicts the LFC score distribution within each of the three

codon types. Using the pMT025-*TP53* library as an example, 1-nt delta variants (data series

colored in red) showed narrower fold-change values than 2-nt and 3-nt delta variants.

## Figure 1

### a

**shRNA/sgRNA library screens**

### b

**ORF variant library screens**

Figure 2

Figure 3

**a**

### *TP53* (MITE)

**b**

### *TP53* (MITE)

### *PDE3A (Twist Bioscience)*

### *PDE3A (Twist Bioscience)*

# Figure 4

**a**

**count**

base calls

**miscall rate**

base calls