Peripheral CB1 receptor blockade acts as a memory enhancer through an adrenergic-dependent mechanism

Sara Martínez-Torres[1,2,3], Araceli Bergadà-Martínez[1], Jorge Ortega[4,5], Lorena Galera-López[1], Arnau Hervera[2,3], Antonio Ortega-Álvaro[1], Floortje Remmers[6], Emma Muñoz-Moreno[7,8], Guadalupe Soria[7,8], José Antonio del Río [2,3], Beat Lutz[6], Jose Ángel Ruíz-Ortega[4], José Javier Meana[4,5], Rafael Maldonado[*1,9], Andrés Ozaita[*1,9]

[1]Laboratory of Neuropharmacology-NeuroPhar, Department of Experimental and Health Sciences. Universitat Pompeu Fabra, Barcelona, Spain.
[2]Molecular and Cellular Neurobiotechnology, Institute for Bioengineering of Catalonia. Parc Científic de Barcelona, Barcelona, Spain.
[3]Centro de Investigación Biomédica en Red sobre Enfermedades Neurodegenerativas; Department of Cell Biology, Physiology and Immunology, Universitat de Barcelona; Institute of Neuroscience, University of Barcelona, Barcelona, Spain.
[4]Department of Pharmacology, Faculty of Medicine and Dentistry, University of the Basque Country, Leioa, Spain.
[5]Centro de Investigación Biomédica en Red de Salud Mental CIBERSAM, Spain; Biocruces Bizkaia Health Research Institute, Spain.
[6]Institute of Physiological Chemistry, University Medical Center of the Johannes Gutenberg University of Mainz, Mainz, Germany.
[7]Experimental 7T MRI Unit (IDIBAPS), Barcelona, Spain.
[8]Laboratory of Surgical Neuroanatomy, Faculty of Medicine and Health Sciences, Institute of Neurosciencies, University of Barcelona, Barcelona, Spain
[9]IMIM Hospital del Mar Research Institute, Barcelona, Spain.

* Corresponding authors: Andrés Ozaita, Laboratori de Neurofarmacologia, Facultat de Ciències de la Salut i de la Vida, Universitat Pompeu Fabra, Parc de Recerca Biomèdica de Barcelona, C/ Doctor Aiguader 88, 08003 Barcelona, Spain. Phone: +34-93-3160823; Fax: + 34-93-3160901; e-mail: andres.ozaita@upf.edu
Rafael Maldonado, Laboratori de Neurofarmacologia, Facultat de Ciències de la Salut i de la Vida, Universitat Pompeu Fabra, Parc de Recerca Biomèdica de Barcelona, C/ Doctor Aiguader 88, 08003 Barcelona, Spain. Phone: +34-93-3160824; Fax: + 34-93-3160901; e-mail: rafael.maldonado@upf.edu

Keywords: memory persistence; CB1 cannabinoid receptor; peripheral-to-central, adrenergic system, vagus nerve, locus coeruleus

Abstract

Peripheral inputs to the brain continuously shape its function and adjust non-emotional memory, but the mechanisms involved are not fully understood. Cannabinoid type-1 receptors (CB1Rs), widely distributed in the organism, are well recognized players in memory performance and their systemic modulation significantly influence memory function. By assessing non-emotional memory in mice, we found a relevant role of peripheral CB1R in memory persistence. Indeed, peripherally restricted CB1R antagonist AM6545 showed a mnemonic effect occluded in adrenalectomized mice, after peripheral adrenergic blockade, or when vagus nerve was chemogenetically inhibited. Genetic CB1R deletion in dopamine β-hydroxylase-expressing cells enhanced memory persistence, supporting a role of peripheral CB1Rs modulating the adrenergic tone. Notably, while brain connectivity was slightly affected by peripheral CB1R inhibition, *locus coeruleus* activity and extracellular norepinephrine in the hippocampus, were increased, and intra-hippocampal β-adrenergic blockade prevented AM6545 mnemonic effects. Together, we disclose a novel peripheral mechanism relevant for non-emotional memory persistence modulation.

Introduction

Most everyday lived events create non-emotional memories with different lifespans. These are generally short-lived recalls that may fade away or, with the intervention of the hippocampus and further recall, may persist through unpredictable periods, even when all short- and long-lived memories were created from similar sensorial stimuli (Morris, 2006). Therefore, the persistence of stimuli-driven memories involves a discrimination/selection of worth-memorizing stimuli, which may occur at or around the time of encoding (Kandel et al., 2014; Yonelinas et al., 2019). Such memory persistence is enhanced in emotionally arousing experiences, where encoding is combined with the natural stress-coping response, which leads to a significant increase in memory persistence for the sensory information recorded at that time (De Quervain et al., 2016). Instead, the mechanisms involved in memory persistence for non-emotional memories are not well understood, although they are most relevant for everyday life events.

The endocannabinoid system (ECS), highly expressed in the central nervous system (CNS) and peripheral tissues (Kano et al., 2009; Maccarrone et al., 2015), plays a key role in learning and memory (Kano et al., 2009). The cannabinoid type-1 receptor (CB1R) is heavily expressed in the brain (Pacher et al., 2006), predominantly localized at presynaptic sites of different neuronal cell types, where it suppresses neurotransmitter release depending on local synaptic activity (Castillo et al., 2012). Exogenous compounds with agonist properties for CB1R contribute to memory impairment (Niyuhire et al., 2007; Puighermanal et al., 2012) in the novel object-recognition test (NORT) in mice as a model for non-emotional memory , while pharmacological or genetic CB1R inactivation increase memory persistence (Maccarrone et al., 2002; Reibaud et al., 1999). Although the mechanisms involved are largely unknown, such regulation of memory by CB1R

blockade was previously assumed to occur solely through centrally located receptors (Zanettini et al., 2011).

We set to challenge this dogma by studing the mnemonic effect of a peripherally restricted CB1R antagonist, AM6545 (Tam et al., 2010). Peripheral CB1R had been previously found relevant for the amnesic effects associated to stress at the time of memory consolidation (Busquets-Garcia et al., 2016), but their contribution to memory enhancement had not been assess previously.

We found that peripheral blockade of CB1Rs on adrenergic cells enhanced non-emotional memory persistence through an adrenergic mechanism involving the vagus nerve, where, as a result, brain connectivity was slighltly reduced while norepinephrine release in the hippocampus was enhanced, all contributing to the enhancement on memory persistence.

Results

CB1R inhibition enhances memory persistence in the novel object-recognition memory test

Novel object-recognition memory is a labile non-emotional type of memory that can be revealed 3 h or 24 h after the familiarization session, when naive mice readily discriminate novel and familiar objects (Fig. 1a). Notably, discrimination values significantly decrease when novel object-recognition memory is assessed 48 h after the familiarization phase (one-way ANOVA, interaction: $F_{(2,19)} = 8.55$, $p = 0.002$; *post hoc* Tukey, 3 h vs 48 h p = 0.003; 24h vs 48h p = 0.007) (Fig. 1a). We used this non-emotional memory paradigm assayed 48 h after the familiarization phase to evaluate the role of CB1R inhibition in memory persistence. We found that acute post-familiarization treatment with a low dose of the systemic CB1R specific antagonist rimonabant (1 mg/kg, i.p.) showed higher memory persistence than vehicle-treated mice (Student's t-test: $p = 0.02$) (Fig. 1b). In addition, heterozygous mice for the *Cnr1* gene (CB1HZ) also showed enhanced memory persistence (Student's t-test: $p = 0.004$) compared to their respective wild-type littermates (Fig. 1c), indicating that such a modulation in memory persistence is CB1R dependent. Post-familiarization administration of the peripherally-restricted CB1R antagonist AM6545 (1 mg/kg, i.p.) also enhanced novel object-recognition memory at 48 h (Student's t-test: $p = 0.002$) (Fig. 1d) pointing to the involvement of a peripheral mechanism. No differences in total exploration time were detected between genotypes or pharmacological treatments in any of the experimental groups above (Supplementary Fig. 1) discarding differences between groups in exploratory behavior. Furthermore, AM6545 treatment did not affect locomotor activity analyzed for 120 min post-administration (Supplementary Fig. 2) discarding a major unspecific effect of the treatment on behavioral responses.

Enhanced memory persistence of peripheral CB1R inhibition involves a peripheral adrenergic mechanism

We hypothesized that a peripherally located tissue such as the adrenal glands, that express CB1R (Hillard, 2015) could be a relevant player modulating memory consolidation (McIntyre et al., 2012). Therefore, we evaluated the effect of post-familiarization AM6545 administration in bilaterally adrenalectomized mice. Memory persistence enhancement by AM6545 was significantly blocked in mice without adrenal glands (two-way ANOVA, interaction: $F_{(1,22)} = 4.89$, $p = 0.037$; *post hoc* Tukey, naive-VEH vs naive-AM6545, $p = 0.022$; naive-AM6545 vs ADX-AM6545, $p = 0.026$) (Fig. 2a), supporting the role of CB1R blockade in this peripheral tissue. Adrenal glands release glucocorticoids and catecholamines into the blood, both relevant for memory (McIntyre et al., 2012). In addition, both rimonabant (Wade et al., 2006) and high doses of AM6545 have been observed to increase circulating corticosteroids (Roberts & Hillard, 2020). To figure out which hormones were responsible for the mnemonic effects produced by peripheral CB1R blockade, mice were pre-treated after the familiarization phase with the glucocorticoid receptor antagonist mifepristone (50 mg/kg) or the peripherally-restricted β-adrenergic receptor antagonist sotalol 20 min before AM6545 injection (Fig. 2b). Mifepristone pre-treatment did not prevent enhancement of memory persistence by AM6545 (two-way ANOVA, interaction: $F_{(1,21)} = 0.038$, $p = 0.845$; mifepristone/vehicle effect: $F_{(1,21)} = 0.707$, $p = 0.409$; AM6545/vehicle effect: $F_{(1,21)} = 25.11$, $p<0.001$) (Fig. 2c). In contrast, mice pre-treated with sotalol did not show the memory improvement observed in AM6545-treated mice (two-way ANOVA, interaction: $F_{(1,31)} = 7.58$, $p = 0.009$; *post hoc* Tukey, Saline-VEH vs Saline-AM6545 $p = 0.01$; Saline-AM6545 vs Sotalol-AM6545 $p = 0.001$) (Fig. 2d).

On the light of these data, we assessed whether inhibition of CB1R exclusively in dopamine beta-hydroxylase cells (DBH+ cells), the main cells responsible for circulating levels of epinephrine/norepinephrine, could mimic the mnemonic effect of systemic and peripheral CB1R antagonists. We used a combination of genetic and pharmacological approaches to show that conditional knock-out mice lacking the CB1R in DBH+ cells (DBH-CB1KO mice) displayed enhanced novel object-recognition memory persistence compared to wild-type controls (Student's t-test: $p = 0.04$) (Fig. 2e). Interestingly, enhanced memory persistence in DBH-CB1KO mice was abolished by sotalol administration (two-way ANOVA, interaction: $F_{(1,22)} = 10.47$, $p = 0.003$; *post hoc* Tukey, Saline-WT vs Saline-DBH-CB1KO $p = 0.01$; Saline- DBH-CB1KO vs Sotalol-DBH-CB1KO $p = 0.04$ (Fig. 2f), pointing to a relevant role of CB1R-modulated peripheral adrenergic/noradrenergic tone in memory persistence. Sotalol pre-treatment similarly prevented the cognitive improvement elicited by systemically-acting rimonabant supporting the relative prominence of peripheral CB1R blockade in such effect of rimonabant (two-way ANOVA, interaction: $F_{(1,37)} = 9.408$, $p = 0.004$; *post hoc* Tukey, Saline-VEH vs Saline-rimonabant $p = 0.01$; Saline-rimonabant vs Sotalol-rimonabant $p = 0.001$) (Fig. 2g). No differences in total exploration time were detected in the memory test between genotypes or pharmacological treatments in any of the experimental groups above (Supplementary Fig. 3).

Chemogenetic vagus nerve inhibition prevents memory improvement produced by peripheral CB1R blockade

Epinephrine administration enhances hippocampal dependent memory in rodents by the activation of β-adrenergic receptors (Dornelles et al., 2007). Such mnemonic effects have been hypothesized to occur either by the activation of β-adrenergic receptors in the liver

and the subsequent increase of blood glucose levels, or by the activation of β-adrenergic receptors on the afferent fibers of the vagus nerve (Gold & Korol, 2012).

To further elucidate whether any of these two mechanisms could be involved in the mnemonic effects produced by acute AM6545 administration, we first measured blood glucose levels after AM6545 administration in mice. No differences in blood glucose levels were observed in mice treated with AM6545 in comparison to vehicle-treated group (Fig. 3a).

To assess the involvement of vagus nerve fibers as a major link between peripheral AM6545 effects and memory performance, we used a chemogenetic approach to selectively reduce the neural activity of vagus nerve. We first confirmed that AM6545 effect on memory persistence was maintained in control mice receiving bilateral vagus nerve injection of AAV5-mCherry which were treated with clozapine N-oxide (CNO, 3 mg/kg, i.p.) (Fig. 3b), indicating that surgery, vagus nerve infection with adeno-associated viral vectors, and CNO administration were compatible with the assessment of memory persistence. However, chemogenetic inhibition in animals with bilateral vagus nerve injection of AAV5-hM4Di prevented AM6545-memory enhancement (Fig. 3c) revealing a role of vagus nerve afferents in the mnemonic effect of AM6545. As expected, no differences were observed neither when memory was evaluated 24 h after familiarization in AAV5-hM4Di mice nor in exploratory behavior during the memory test (Supplementary Fig. 4). Thus, AM6545 administration improves memory consolidation through an adrenergic mechanism involving the vagus nerve.

Central effects of AM6545 administration

We first analyzed the pattern of c-Fos expression, as a marker of neuronal activity, focusing on brain regions relevant for memory performance. Samples were obtained 90

min after receiving AM6545 or vehicle once mice came out of the familiarization phase in the Novel object recognition test (NORT). Brain areas included CA1 and CA3 hippocampal regions, dentate gyrus, prelimbic and infralimbic prefrontal cortex, the locus coeruleus and basal, lateral and central regions of the amygdala. No significant differences in c-Fos expression were observed between AM6545- and vehicle-treated mice in the areas analyzed (Supplementary Fig.5). However, network analysis using the Pearson correlation coefficient for each pair of regions revealed that AM6545 treatment produced a significant alteration over the functional connectivity of these brain areas compared to the vehicle condition. (Fig. 4a-b and Supplementary Fig.6). To compare positive and negative connectivity between groups we calculated Z-scores starting from the Pearson r correlation values. AM6545 treatment did not significantly modified neither positive nor negative correlations (Fig. 4c-d).

We further explored the central effects after peripheral CB1R blockade by performing resting state functional magnetic resonance imaging (rsfMRI). After 1h of AM6545 administration, global connectivity metrics showed significant enhancement of AM6545 treatment in whole brain local efficiency (Fig. 5a) (uncorrected $p<0.05$) indicating an enhanced ability of parallel transmission of information. Regional connectivity analysis showed significant increases of nodal efficiency values in AM6545- treated mice compared to vehicle in left frontal cortex (Kruskal-Wallis: $p<0.01$), right hippocampus (Kruskal-Wallis: $p = 0.023$) and right globus pallidus (Kruskal-Wallis: $p = 0.023$) (Fig. 5 b-d), although these differences did not reach significance when corrected for multiple comparisons (FDR), indicating a subtle effect of AM6545 enhancing brain connectivity. Seed-based analysis of rsfMRI with seed in the posterior brainstem, which contains the nucleus tractus solitarius (NTS), a region receiving vagal afferents, exhibited a different pattern of connectivity in AM6545-treated animals compared to vehicle. Significant

differences of this connectivity pattern were observed in areas corresponding to hippocampus, frontal cortex, cingulate cortex, brainstem nuclei such as pontine reticular nucleus, different preolivary and spinal trigeminal nuclei and cerebellar nuclei (Fig. 5e). In these regions AM6545-treated animals showed strong negative correlation values with the seed BOLD signal, especially in hippocampus, that is not observed in vehicle treated animals (Supplementary Fig.7).

Increased hippocampal norepinephrine mobilization by peripheral CB1R blockade induces memory improvement

Next, we measured the firing rate at *locus coeruleus* neurons after CB1R peripheral modulation with AM6545. Systemic AM6545 produced a non-significant enhancement of basal firing rates compared to vehicle-treated mice (Fig. 6a). To assess the functional significance of the basal firing rate, we performed extracellular microdialysis analysis in the hippocampus after systemic AM6545 treatment. Analysis of norepinephrine (NE), dopamine (DA) and serotonin (5-HT) extracellular levels after AM6545 administration revealed a specific transient increase in NE in comparison to the vehicle-treated mice (two-way repeated measure ANOVA, interaction: $F_{(1,14)} = 2.19$ p = 0.009) (Fig. 6b and Supplementary Fig.8). We then tested whether β-adrenergic receptors in the hippocampal region were involved in the increased memory persistence mediated by AM6545. We found that local intra-hippocampal microinjection of propranolol at a dose that did not affect memory performance on its own (1 μg per 0.5 μl per side, see Supplementary Fig. 9) blocked the mnemonic effects produced by systemic AM6545 administration (two-way ANOVA, interaction: $F_{(1,119)} = 5.03$, p = 0.03; *post hoc* Tukey, Saline-VEH vs Saline-AM6545 p = 0.004; Saline-AM6545 vs Propranolol-AM6545 p = 0.01) (Fig. 6c), without affecting the exploratory behavior on the memory test (Fig. 6d). These data

indicate the functional relevance of noradrenergic hippocampal activation in the effect of peripheral CB1R blockade over memory persistence.

Discussion

Our study identifies a relevant role of the peripheral ECS in modulating non-emotional memory persistence through the mobilization of central and peripheral adrenergic/noradrenergic mechanisms.

We choose to study novel object-recognition memory, a hippocampal-dependent test (Cohen & Stackman, 2015) since this is a model of non-emotional memory and memory persistence might be modulated by post-training manipulation. In this regard, our group has observed deficits in novel object-recognition memory by CB1R agonists (Puighermanal et al., 2009), endocannabinoid build-up (Busquets-Garcia et al., 2011) or stress (Busquets-Garcia et al., 2016), all administered after the familiarization phase. In the present study, we analyzed novel object-recognition memory recall 48 h after the familiarization phase since at that time mice usually show reduced signs of novel object discrimination compared to shorter intervals. Using this 48 h interval, we found that overall CB1R blockade, through pharmacological or genetic approaches, significantly increased discrimination indexes. In agreement, a number of previous studies had demonstrated that both pharmacological or genetic CB1R inactivation enhanced memory in different hippocampal-dependent tasks (Jacob et al., 2012; Lichtman, 2000; Maccarrone et al., 2002; Reibaud et al., 1999; Takahashi et al., 2005; Wolff & Leander, 2003). We found that the peripherally restricted CB1R neutral antagonist AM6545 administered after the familiarization phase significantly enhanced object-recognition memory persistence in mice. Actually, this drug, compared to the systemic antagonist rimonabant, shows no significant blood-brain barrier permeability at doses ten times higher than the one we used in the present study (Tam et al., 2010), suggesting that the AM6545 effects in our study derive from a peripheral mechanism. Notably, adrenalectomized mice treated with AM6545 did not present an enhancement in memory

persistence, pointing to a relevant role of adrenal glands in this response. Epinephrine/norepinephrine and corticosteroids secreted by the adrenal glands have a significant impact in memory consolidation (McIntyre et al., 2012; Roozendaal & McGaugh, 2011; Yang et al., 2013). We then gathered evidence that pointed to the mobilization of the adrenergic transmission, compared to the mobilization of corticosteroids, by peripheral CB1R blockade given that AM6545-induced memory persistence enhancement was prevented by the peripherally restricted β-adrenergic receptor antagonist sotalol, but not by the corticosteroid receptor antagonist mifepristone. Such differential modulation by adrenal hormones by AM6545 is indicative of a rather specific mechanism of peripheral CB1R blockade for potentiating the consolidation of non-emotional memory, since sotalol has been found ineffective in modulating fear conditioning (Lee et al., 2001) and mifepristone was previously found to reduce contextual memory in a fear conditioning paradigm (Zhou et al., 2010) but it did not modify NORT results in our experimental conditions. Notably, the fact that mice lacking CB1R in DBH+ cells (Busquets-Garcia et al., 2016) (DBH-CB1KO) reproduce a similar phenotype in enhanced memory persistence than constitutive CB1HZ mice, and our finding that sotalol prevents such a mnemonic phenotype on DBH-CB1KO mice and after systemic rimonabant treatment, points to a relevant effect of the peripheral adrenergic CB1R in the systemic effect of CB1R inhibition, and an undisclosed role of peripheral CB1R signaling in memory persistence.

The study of c-Fos expression after AM6545 administration allowed to confirm no major differences in neuronal stimulation due to the treatment, but significant alterations in functional connectivity between areas analyzed. Although c-Fos may not reveal the real activity of brain regions given its limited expression and the time constrains of the analysis (McReynolds et al., 2018), similar approaches to ours have been successfully

performed to acquire a glimpse of brain functional connectivity between discrete areas (Silva et al., 2019; Tanimizu et al., 2017). The subtle alterations in brain connectivity revealed by our biased c-Fos inter-regional correlation analysis were further corroborated by rsfMRI using an unbiased approach to study connectivity. The brain's intrinsic functional organization associated to the resting state has been proposed to determine the ability to create flexible and suitable behavioral outcomes to cognitive demands (Sala-Llonch et al., 2012). Indeed, enhancements in nodal efficiency, a representation for the ability of information propagation between a node and the remaining nodes in the network, were observed specifically in the frontal cortex, hippocampus and globus pallidus, a result parsimonious with the engagement of these regions in long-term memory consolidation (Kitamura et al., 2017; Tanimizu et al., 2017) as consequence of AM6545 treatment.

A growing body of evidence has demonstrated the efficacy of epinephrine administration, which does not cross the blood-brain barrier, to enhance hippocampal-dependent memory in rodents (Dornelles et al., 2007; Talley et al., 2000) by enhancing central noradrenergic signaling. Our data reveal that AM6545 administration did modestly modify brain activity to produce a reduction in brain network connectivity and a partial increase in basal LC firing and norepinephrine extracellular levels in the hippocampal region, suggesting increased activity of LC projections to the hippocampus. These finding are compatible with the activation of the vagus nerve by increasing peripheral adrenaline/noradrenaline, since it was demonstrated that epinephrine administration results in increased vagal nerve firing that was sensitive to sotalol administration (MIYASHITA & WILLIAMS, 2006). Vagus nerve stimulation has been found to enhance memory retention in mouse models (Vázquez-Oliver et al., 2020) as well as in humans (Ghacibeh et al., 2006). LC activity may be partially determined by the vagal nerve afferents but is well stablished to be

physiologically engaged on novelty and it is considered to prime the persistence of hippocampal-based long-term memories (Niels Hansen, 2017; Sara, 2009). Moreover, norepinephrine in the hippocampus facilitates the storage of new memories through the regulation of neural excitability and synaptic plasticity (Hagena et al., 2016). Although norepinephrine can bind to both α- or β-adrenergic receptors, synaptic information and plasticity in the hippocampus would depend largely on the activation of β-adrenergic receptors (A. Kemp & Manahan-Vaughan, 2008). Indeed, LC stimulation modulates hippocampal synaptic strength (N. Hansen & Manahan-Vaughan, 2015; A. Kemp & Manahan-Vaughan, 2008; Anne Kemp & Manahan-Vaughan, 2012), spike coupling of CA1 pyramidal neurons (Bacon et al., 2020) and improves memory through a β-adrenergic-dependent mechanism (Hagena et al., 2016; N. Hansen & Manahan-Vaughan, 2015). In agreement, we observed that the intra-hippocampal blockade of β-adrenergic receptors with propranolol prevented the increase of novel object-recognition memory persistence produced by AM6545.

Altogether, our study identifies peripheral CB1R as a relevant target to enhance memory persistence through central and peripheral adrenergic/noradrenergic mechanism that engages the vagus nerve. This novel peripheral mechanism involving CB1R could underlie previously described effects of systemic CB1R inhibitors assumed to be associated to direct interaction with central targets.

Materials and Methods

Animals

Young adult (10-12 weeks old) male Swiss albino (CD-1) mice (Charles River, France) were used for pharmacological approaches on behavioural, microdialysis, electrophysiological and biochemical experiments. For genetic approaches to reduce CB1R expression, heterozygous mice for the *Cnr1* gene and their wild-type littermates in C57BL/6J genetic background were used (Zimmer et al., 1999). Conditional KO mice for the *Cnr1* gene lacking CB1R exclusively in DβH-expressing cells were generated as previously detailed (Busquets-Garcia et al., 2016) in C57BL/6J genetic background.

Mice were housed in Plexiglas cages (2-4 mice per cage) and maintained in temperature $(21 \pm 1\,°C)$ and humidity $(55 \pm 10\%)$ controlled environment. Food and water were available *ad libitum*. All the experiments were performed during the light phase of a 12 h cycle (light on at 8 am; light off at 8 pm). Mice were habituated to the experimental room and handled for 1 week before starting the experiments. All behavioural experiments were conducted by an observer blind to experimental conditions.

All animal procedures were conducted following "Animals in Research: Reporting Experiments" (ARRIVE) guidelines and standard ethical guidelines (Kilkenny et al., 2010) (European Directive 2010/63/EU). Animal procedures were approved by the local ethical committee (Comitè Ètic d'Experimentació Animal-Parc de Recerca Biomèdica de Barcelona, CEEA-PRBB).

Drugs and treatments

Rimonabant (Axon Medchem) and mifepristone (Sigma-Aldrich) were dissolved in 5% ethanol, 5% cremophor-EL and 90% saline (0.9% NaCl). AM6545 (Tocris-Bio-Techne) was dissolved in 0.26% DMSO, 4.74% ethanol, 5% cremophor-EL and 90% saline.

Sotalol (Sigma-Aldrich) was dissolved in 90% saline. Propranolol (Sigma-Aldrich) was dissolved in 90% saline. For the activation of the inhibitory designer receptors exclusively activated by designer drugs (hM4Di-DREADD), clozapine N-oxide (CNO) (Enzo Life Sciences, NY) was dissolved 0.9% sterile saline. All intraperitoneally (i.p.)-injected drugs were administered in a volume of 10 ml/kg of body weight at the doses and time points indicated.

Viral vectors

We used the following vectors: AAV-hM4Di-DREADD (AAV5- hSyn-hM4D(Gi)-mCherry) and AAV-control-DREADD (AAV5-hSyn-mCherry) from Addgene.

To detect the viral expression in all the experiments we visualized through epifluorescence microscopy the reporter mCherry in the nodose ganglion.

Bilateral adrenalectomy

Mice were anesthetized by isoflurane inhalation, 5% v/v induction and 3% v/v for maintenance, with oxygen (0.8 L/min). A small incision of 1 cm was made in the left and right flanks, and the adrenal glands were identified and removed from the surrounding tissue. Wounds were closed in two layers using 4/0 silk sutures (Alcon). All animals were given access to saline after surgery to ensure adequate salt balance. The experiments were resumed following a recovery period of 10 d.

Bilateral intra-hippocampal cannula implantation

Bilateral intra-hippocampal cannula implantation was performed as previously described (Busquets-Garcia et al., 2018).

Intra-hippocampal drug administration

After familiarization phase of NORT, mice received a bilateral intra-hippocampal injection of 0.50 μl of propranolol (0.5, 1 or 2 μg) or saline at a constant rate of 0.25 μl/min by using a microinfusion pump during 2 min. The injection cannula projected 1.00 mm below the ventral tip of the implanted guide cannula. The displacement of an air bubble inside the length of the polyethylene tubing that connected the Hamilton syringe to the injection needle was used to monitor the microinjections. After infusion, the injection cannula was left for an additional period of 2 min to allow the fluid to diffuse and to prevent reflux before withdrawal.

Surgery and virus vector microinjection

Mice were anesthetized with a mixture of ketamine hydrochloride (Imalgène; Merial Laboratorios S.A.) and medetomidine hydrochloride (Domtor; Esteve, Spain) dissolved in sterile 0.9% physiological saline and administered intraperitoneally (i.p, 75 mg/kg, and 1 mg/kg of body weight respectively) and placed in a supine position. A small incision from the larynx to approximately the beginning of the sternum was made in the left and right flanks. We expose the submaxillary glands by blunt dissection, and we open the midline to expose the trachea. We identify the vagus nerve and the carotid artery within the set of neurovascular fibers parallel to the trachea, and we proceed to isolate the nerve from the artery using a spinal hook, taking great care not to damage the artery.

Once isolated, we introduce a pair of blunt-tipped forceps to keep the nerve fixed, between the nerve and the rest of the tissue. We then proceed to inoculate 1 μl of a viral

suspension (AAV) using a Hamilton syringe (34G, 45º bevel) very slowly. First, we introduce the needle into the nerve about 3mm, and as we inoculate, we move the needle back, before completely withdrawing the needle, we rotate the syringe slightly to avoid transfer. We remove the holding forceps and return the nerve to its initial position. We repeat the same procedure on the nerve on the other side. Once the inoculations are finished, we close the exposed area in layers and close the skin incision by means of a simple stitch suture with 4/0 braided silk.

The animals are then administered for 48 hours with analgesic and antibiotic. A possible nerve injury is also monitored by analysis of urination and defecation capacity.

Behavioural studies

NORT was performed following a protocol previously described (Puighermanal et al., 2009).

Locomotor activity

Locomotor activity was assessed for 120 min after acute administration of AM6545. Individual locomotor activity boxes ($9 \times 20 \times 11$ cm) (Imetronic) were used in a low luminosity environment (5 lux). The total activity and the total number of rearings were detected by infrared sensors to detect locomotor activity and infrared plane to detect rearings.

Tissue preparation for immunofluorescence

Mice were deeply anesthetized by i.p. injection (0.2 ml/10 g of body weight) of a mixture of ketamine/xylazine (100 mg/kg and 20 mg/kg, respectively) prior to intracardiac perfusion of cold 4% paraformaldehyde in 0.1 M phosphate buffer, pH7.4 (PB). Brains

were removed and post-fixed overnight at 4 °C in the same fixative solution. The next day, brains were moved to PB at 4 °C. Coronal brain sections of 30 μm were made on a freezing microtome and stored in a 5% sucrose solution at 4 ºC until use.

Immunofluorescence and cell quantification

Immunofluorescence and cell quantification were performed as previously described (Navarro-Romero et al., 2019). Antibodies against c-Fos (anti-cFos, rabbit, 1:1000, sc-72, Santa Cruz Biotechnology) were used as primary antibody and anti-rabbit (donkey, Alexa Fluor-488, A21206, 1:600, Life Technologies) and anti-rabbit (rabbit, Alexa 555, 1:1000, Jackson ImmunoResearch) as secondary antibodies respectively.

Network construction and graph analysis

Correlation matrices and circos plot were generated using Pearson *r* values from interregional correlations of c-fos expression data. Pearson *r* values were converted to Z-scores using Fisher's transformation to determine effects of AM6545 on positive and negative functional connectivity and compare across experimental groups. All graph analysis was performed in R (v4.0) using the DescTools (v0.99.40; Andri Signorell et mult. al., 2021) and the ggcorrplot (v0.1.3; Alboukadel Kassambara, 2019) packages.

Image acquisition

Magnetic resonance image was performed in the in-house Magnetic Resonance Core Facility (IDIBAPS 7T MRI Unit). Experiments were conducted on a 7.0T BioSpec 70/30 horizontal animal scanner (Bruker BioSpin, Ettlingen, Germany), equipped with an actively shielded gradient system (400 mT/m, 12 cm inner diameter). The receiver coil is a surface coil for the mouse brain. Animals were sedated (induction dose of 4%

isofluorane in a mixture of 30% O2 and 70% N2) and placed in supine position in a Plexiglas holder with a nose cone for administering anesthetic gases and were fixed using tooth and ear bars and adhesive tape. Eyes were protected from dryness with Siccafluid 2.5 mg/g ophthalmologic fluid. Once placed in the holder and keeping isoflurane to 1.5 %, a subcutaneous bolus (0.3 mg/kg) of medetomidine (Domtor®, Orion Pharma) was injected. For the next 15 min, the % of isofluorane was progressively decreased until 0.5 %. Then, a continuous perfusion of 0.6 mg/kg/h medetomidine started and was maintained until the end of the acquisition session. After completion of the imaging session, 1 µl/gr of atipamezol (Antisedan®, Orion Pharma) and saline were injected to reverse the sedative effect and compensate fluid loss.

Localizer scans were used to ensure accurate position of the head at the isocenter of the magnet. Anatomical T2 RARE images were acquired in coronal orientation with effective TE=33 ms TR=2.3 s, RARE factor=8, voxel size = 0.08 x 0.08 mm² and slice thickness = 0.5 mm. rs-fMRI was acquired with an EPI sequence with TR=2 s, TE=19.4, voxel size 0.21x0.21 mm² and slice thickness 0.5 mm. 420 volumes were acquired resulting in an acquisition time of 14 min.

Image analysis and processing

Two approaches were used to evaluate functional connectivity. On the one hand, whole-brain connectivity was evaluated using global and regional network metrics (Rubinov & Sporns, 2010) and on the other hand seed-based analysis was performed to evaluate connectivity of posterior brainstem with the rest of the brain. For both approaches, rs-fMRI was preprocessed, including slice timing, motion correction by spatial realignment using SPM8, correction of EPI distortion by elastic registration to the T2-weighted volume using ANTs (Avants et al., 2008), detrend, smoothing with a full-width half

maximum (FWHM) of 0.6mm, frequency filtering of the time series between 0.01 and 0.1 Hz and regression by motion parameters. All these steps were performed using NiTime (http://nipy.org/nitime). Brain parcellation was performed by registration of a mouse brain atlas (Ma et al., 2008) to the T2-RARE acquisition of each subject using ANTs diffeomorphic registration. Region parcellation was then registered from the T2-weighted volume to the preprocessed mean rs-fMRI. Whole-brain functional brain network was estimated considering the gray matter regions obtained by parcellation as the network nodes. Connectivity between each pair of regions was estimated as the Fisher-z transform of the correlation between average time series in each region. Network organization was quantified using regional and global graph metrics. Strength, local efficiency and clustering coefficient was computed for every region and whole brain connectivity was quantified by average strength, global and local efficiency and average clustering coefficient (Rubinov & Sporns, 2010).To perform the seed-based analysis, posterior brainstem containing the nuclei of the solitary tract was manually drawn in each animal. Average time series in the seed was computed and correlated with each voxel time series, resulting in a correlation map describing the connectivity of this region with the rest of the brain. Voxel-wise differences between the correlation maps were estimated using fsl-randomize, that performs non-parametric permutation test including family-wise error correction on neuroimaging data (Winkler et al., 2014).

In vivo microdialysis

Animals were anesthetized with isofluorane (1.5 - 2.5 % v/v for induction and maintenance) and placed in a Kopf stereotaxic frame. Intracerebral probe (cuprophan membrane of 2 mm) was implanted in the hippocampus and fixed to the skull. The coordinates were AP -3.4 mm, L +2.6 mm, DV –4.2 mm (Franklin and Paxinos, 1997).

The next day, mice were placed in a plastic bowl and connected to a fraction collection system for freely-moving animals (Raturn, BASi, USA). The input tube of the dialysis probe was connected to a syringe pump (BeeHive and BabyBee, BASi), which delivered a modified cerebrospinal fluid (CSF) containing NaCl 148 mM, KCl 2.7 mM, CaCl2 1.2mM and MgCl2 0.85 mM (pH 7.4) to the probe at a rate of 1 µl/min. The output tubes from the animals were attached to a refrigerated fraction collector (HoneyComb, BASi). Samples were collected every 35 min for the analysis of the different neurotransmitters on vials containing 5 µl of acetic acid 0.1 M. Eight baseline samples were collected from each animal, but only the last 6 ones were used for subsequent analysis.

NA, DA and 5-HT chromatographic analysis

Neurotransmitter concentrations were measured immediately after samples collection by Ultra Performance Liquid Chromatography (UPLC) coupled to an electrochemical detector (Alexys analyser, Antec Leyden, Holland). The mobile phase consisted of 100 mM phosphoric acid, 100 mM citric acid, 0.1 mM EDTA, 950-1500 mg/l 1-octanesulfonic acid (OSA), 5% v/v acetonitrile; the pH was adjusted to 6.0 with 50% NaOH/ 45% KOH solution. The flow rate of the mobile phase was 0.075 ml/min and the temperature for the analytical column (Acquity UPLC BEH C18, 1.7 µm, 1x100 mm; Waters, Milford, USA) was 37 ºC.

In vivo electrophysiological recording in the locus coeruleus

Mice were anaesthetized with chloral hydrate (400 mg/ kg, i.p.) and placed in the stereotaxic frame with the skull positioned horizontally. A burr hole was drilled and the recording electrode was placed 1.5 mm posterior to lambda and 0.2–1.2 mm from the midline and lowered into the LC usually encountered at a depth of between 2.7 and 4.0

mm from the brain surface (Gobbi et al., 2007) catheter (Terumo Surflo1; Teruma Medical Products) was then inserted in the peritoneum for additional administrations of anaesthetic by a pump and systemic drug. The body temperature was maintained at 37 ºC for the entire experiment using a heating pad.

Single-unit extracellular recordings of mouse LC neurons were performed as previously described (Gobbi et al., 2007). The recording electrode was filled with 2% solution of Pontamine Sky Blue in 0.5% sodium acetate and broken back to a tip diameter of 1–2 mm. The electrode was lowered into the brain by means of a hydraulic microdrive (model 640; David Kopf Instruments). LC neurons were identified by standard criteria, which included spontaneous activity displaying a regular rhythm and firing rate between 0.5 and 5 Hz, characteristic spikes with a long-lasting (>2 ms), positive–negative waveform action potentials and the biphasic excitation–inhibition response to pressure applied on contralateral hind paw (paw pinch), as previously described in mice (Gobbi et al., 2007) and rats (Cedarbaum & Aghajanian, 1978). The extracellular signal from the electrode was pre-amplified and amplified later with a high-input impedance amplifier and then monitored on an oscilloscope and on an audio monitor. This activity was processed using computer software (Spike2 software; Cambridge Electronic Design) and firing rate was calculated. Basal firing rate and other electrophysiological parameters were measured for 3 min. Changes in firing rate were expressed as percentages of the basal firing rate (mean firing rate for 3 min prior to drug injection) and were measured after 35 min until the end of the experiment. Only 1 cell was studied in each animal when any drug was administered.

Statistical analysis

Results are reported as mean ± standard error of the mean (s.e.m). Data analysis were performed using GraphPad Prism software (GraphPad Software). Statistical comparisons were evaluated using unpaired Student's t-test for 2 groups comparisons or two-way ANOVA for multiple comparisons. Subsequent Tukey *post hoc* was used when required (significant interaction between factors). Comparisons were considered statistically significant when $p < 0.05$.

Acknowledgements

We thank Dulce Real, Marta Linares and Francisco Porrón for expert technical assistance, and the Laboratory of Neuropharmacology-NeuroPhar for helpful discussion.

S.M-T. was the recipient of a predoctoral fellowship (Generalitat de Catalunya) [FI-B00531 2016], A.B-M. was supported by predoctoral fellowship (Generalitat de Catalunya) [FI_B0052 2020]. This study was supported by Ministerio de Economía, Innovación y Competitividad (MINECO), Spain (#RTI2018-099282-B-I00B to A.O., #SAF2017-84060-R to R.M.; Generalitat de Catalunya, Spain (2017SGR-669 to R.M.); ICREA (Institució Catalana de Recerca i Estudis Avançats, Spain) Academia to A.O. and R.M. Grant "Unidad de Excelencia María de Maeztu", funded by the MINECO (#MDM-2014-0370). FEDER, European Commission funding is also acknowledged.

This research was supported by PRPSEM Project with ref. RTI2018-099773-B-I00 from (MCIU/FEDER/AEI), the CERCA Program, and the Commission for Universities and Research of the Department of Innovation, Universities, and Enterprise of the Generalitat de Catalunya (SGR2017-648) to JADR. IBEC is the recipients of a Severo Ochoa Award of Excellence from MINECO (CEX2018-000789-S). SMT and AH were supported by Severo Ochoa program at IBEC

References

Avants, B. B., Epstein, C. L., Grossman, M., & Gee, J. C. (2008). Symmetric diffeomorphic image registration with cross-correlation: Evaluating automated labeling of elderly and neurodegenerative brain. *Medical Image Analysis*, *12*(1), 26–41. https://doi.org/10.1016/j.media.2007.06.004

Bacon, T. J., Pickering, A. E., & Mellor, J. R. (2020). Noradrenaline release from locus coeruleus terminals in the hippocampus enhances excitation-spike coupling in ca1 pyramidal neurons via β-adrenoceptors. *Cerebral Cortex*, *30*(12), 6135–6151. https://doi.org/10.1093/cercor/bhaa159

Busquets-Garcia, A., Gomis-González, M., Salgado-Mendialdúa, V., Galera-López, L., Puighermanal, E., Martín-García, E., Maldonado, R., & Ozaita, A. (2018). Hippocampal Protein Kinase C Signaling Mediates the Short-Term Memory Impairment Induced by Delta9-Tetrahydrocannabinol. *Neuropsychopharmacology*, *43*(5), 1021–1031. https://doi.org/10.1038/npp.2017.175

Busquets-Garcia, A., Gomis-González, M., Srivastava, R. K., Cutando, L., Ortega-Alvaro, A., Ruehle, S., Remmers, F., Bindila, L., Bellocchio, L., Marsicano, G., Lutz, B., Maldonado, R., & Ozaita, A. (2016). Peripheral and central CB1 cannabinoid receptors control stress-induced impairment of memory consolidation. *Proceedings of the National Academy of Sciences of the United States of America*, *113*(35), 9904–9909. https://doi.org/10.1073/pnas.1525066113

Busquets-Garcia, A., Puighermanal, E., Pastor, A., de la Torre, R., Maldonado, R., & Ozaita, A. (2011). Differential role of anandamide and 2-arachidonoylglycerol in memory and anxiety-like responses. *Biological Psychiatry*, *70*(5), 479–486. https://doi.org/10.1016/j.biopsych.2011.04.022

Castillo, P. E., Younts, T. J., Chávez, A. E., & Hashimotodani, Y. (2012). Endocannabinoid Signaling and Synaptic Function. In *Neuron* (Vol. 76, Issue 1, pp. 70–81). Neuron. https://doi.org/10.1016/j.neuron.2012.09.020

Cedarbaum, J. M., & Aghajanian, G. K. (1978). Afferent projections to the rat locus coeruleus as determined by a retrograde tracing technique. *The Journal of Comparative Neurology*, *178*(1), 1–15. https://doi.org/10.1002/cne.901780102

Cohen, S. J., & Stackman, R. W. (2015). Assessing rodent hippocampal involvement in the novel object recognition task. A review. *Behavioural Brain Research*, *285*, 105–117. https://doi.org/10.1016/j.bbr.2014.08.002

De Quervain, D., Schwabe, L., & Roozendaal, B. (2016). Stress, glucocorticoids and memory: Implications for treating fear-related disorders. In *Nature Reviews Neuroscience* (Vol. 18, Issue 1, pp. 7–19). Nature Publishing Group. https://doi.org/10.1038/nrn.2016.155

Dornelles, A., de Lima, M. N. M., Grazziotin, M., Presti-Torres, J., Garcia, V. A., Scalco, F. S., Roesler, R., & Schröder, N. (2007). Adrenergic enhancement of consolidation of object recognition memory. *Neurobiology of Learning and Memory*, *88*(1), 137–142. https://doi.org/10.1016/j.nlm.2007.01.005

Ghacibeh, G. A., Shenker, J. I., Shenal, B., Uthman, B. M., & Heilman, K. M. (2006). The influence of vagus nerve stimulation on memory. *Cognitive and Behavioral Neurology*, *19*(3), 119–122. https://doi.org/10.1097/01.wnn.0000213908.34278.7d

Gobbi, G., Cassano, T., Radja, F., Morgese, M. G., Cuomo, V., Santarelli, L., Hen, R., & Blier, P. (2007). Neurokinin 1 receptor antagonism requires norepinephrine to increase serotonin function. *European Neuropsychopharmacology*, *17*(5), 328–338. https://doi.org/10.1016/j.euroneuro.2006.07.004

Gold, P. E., & Korol, D. L. (2012). Making memories matter. *Frontiers in Integrative*

Neuroscience, *6*(NOV). https://doi.org/10.3389/fnint.2012.00116

Hagena, H., Hansen, N., & Manahan-Vaughan, D. (2016). β-Adrenergic Control of Hippocampal Function: Subserving the Choreography of Synaptic Information Storage and Memory. *Cerebral Cortex*, *26*(4), 1349–1364. https://doi.org/10.1093/cercor/bhv330

Hansen, N., & Manahan-Vaughan, D. (2015). Locus Coeruleus Stimulation Facilitates Long-Term Depression in the Dentate Gyrus That Requires Activation of -Adrenergic Receptors. *Cerebral Cortex*, *25*(7), 1889–1896. https://doi.org/10.1093/cercor/bht429

Hansen, Niels. (2017). The Longevity of Hippocampus-Dependent Memory Is Orchestrated by the Locus Coeruleus-Noradrenergic System. *Neural Plasticity*, *2017*, 1–9. https://doi.org/10.1155/2017/2727602

Hillard, C. J. (2015). Endocannabinoids and the Endocrine System in Health and Disease. In *Handbook of experimental pharmacology* (Vol. 231, pp. 317–339). https://doi.org/10.1007/978-3-319-20825-1_11

Jacob, W., Marsch, R., Marsicano, G., Lutz, B., & Wotjak, C. T. (2012). Cannabinoid CB1 receptor deficiency increases contextual fear memory under highly aversive conditions and long-term potentiation in vivo. *Neurobiology of Learning and Memory*, *98*(1), 47–55. https://doi.org/10.1016/j.nlm.2012.04.008

Kandel, E. R., Dudai, Y., & Mayford, M. R. (2014). The molecular and systems biology of memory. In *Cell* (Vol. 157, Issue 1, pp. 163–186). Cell Press. https://doi.org/10.1016/j.cell.2014.03.001

Kano, M., Ohno-Shosaku, T., Hashimotodani, Y., Uchigashima, M., & Watanabe, M. (2009). Endocannabinoid-Mediated Control of Synaptic Transmission. *Physiological Reviews*, *89*(1), 309–380. https://doi.org/10.1152/physrev.00019.2008

Kemp, A., & Manahan-Vaughan, D. (2008). -Adrenoreceptors Comprise a Critical Element in Learning-Facilitated Long-Term Plasticity. *Cerebral Cortex*, *18*(6), 1326–1334. https://doi.org/10.1093/cercor/bhm164

Kemp, Anne, & Manahan-Vaughan, D. (2012). Passive spatial perception facilitates the expression of persistent hippocampal long-term depression. *Cerebral Cortex (New York, N.Y. : 1991)*, *22*(7), 1614–1621. https://doi.org/10.1093/cercor/bhr233

Kilkenny, C., Browne, W., Cuthill, I. C., Emerson, M., Altman, D. G., & NC3Rs Reporting Guidelines Working Group. (2010). Animal research: Reporting in vivo experiments: The ARRIVE guidelines. *British Journal of Pharmacology*, *160*(7), 1577–1579. https://doi.org/10.1111/j.1476-5381.2010.00872.x

Kitamura, T., Ogawa, S. K., Roy, D. S., Okuyama, T., Morrissey, M. D., Smith, L. M., Redondo, R. L., & Tonegawa, S. (2017). Engrams and circuits crucial for systems consolidation of a memory. *Science (New York, N.Y.)*, *356*(6333), 73–78. https://doi.org/10.1126/science.aam6808

Lee, H. J., Berger, S. Y., Stiedl, O., Spiess, J., & Kim, J. J. (2001). Post-training injections of catecholaminergic drugs do not modulate fear conditioning in rats and mice. *Neuroscience Letters*, *303*(2), 123–126. https://doi.org/10.1016/S0304-3940(01)01733-5

Lichtman, A. H. (2000). SR 141716A enhances spatial memory as assessed in a radial-arm maze task in rats. *European Journal of Pharmacology*, *404*(1–2), 175–179.

Ma, Y., Smith, D., Hof, P. R., Foerster, B., Hamilton, S., Blackband, S. J., Yu, M., & Benveniste, H. (2008). In vivo 3D digital atlas database of the adult C57BL/6J mouse brain by magnetic resonance microscopy. *Frontiers in Neuroanatomy*, *2*(APR). https://doi.org/10.3389/neuro.05.001.2008

Maccarrone, M., Bab, I., Bíró, T., Cabral, G. A., Dey, S. K., Di Marzo, V., Konje, J. C., Kunos, G., Mechoulam, R., Pacher, P., Sharkey, K. A., & Zimmer, A. (2015). Endocannabinoid signaling at the periphery: 50 years after THC. *Trends in Pharmacological Sciences*, *36*(5), 277–296. https://doi.org/10.1016/j.tips.2015.02.008

Maccarrone, M., Valverde, O., Barbaccia, M. L., Castañé, A., Maldonado, R., Ledent, C., Parmentier, M., & Finazzi-Agrò, A. (2002). Age-related changes of anandamide metabolism in CB1 cannabinoid receptor knockout mice: correlation with behaviour. *The European Journal of Neuroscience*, *15*(7), 1178–1186.

McIntyre, C. K., McGaugh, J. L., & Williams, C. L. (2012). Interacting brain systems modulate memory consolidation. *Neuroscience & Biobehavioral Reviews*, *36*(7), 1750–1762. https://doi.org/10.1016/j.neubiorev.2011.11.001

McReynolds, J. R., Christianson, J. P., Blacktop, J. M., & Mantsch, J. R. (2018). What does the Fos say? Using Fos-based approaches to understand the contribution of stress to substance use disorders. In *Neurobiology of Stress* (Vol. 9, pp. 271–285). Elsevier Inc. https://doi.org/10.1016/j.ynstr.2018.05.004

MIYASHITA, T., & WILLIAMS, C. (2006). Epinephrine administration increases neural impulses propagated along the vagus nerve: Role of peripheral β-adrenergic receptors. *Neurobiology of Learning and Memory*, *85*(2), 116–124. https://doi.org/10.1016/j.nlm.2005.08.013

Morris, R. G. M. (2006). Elements of a neurobiological theory of hippocampal function: the role of synaptic plasticity, synaptic tagging and schemas. *The European Journal of Neuroscience*, *23*(11), 2829–2846. https://doi.org/10.1111/j.1460-9568.2006.04888.x

Navarro-Romero, A., Vázquez-Oliver, A., Gomis-González, M., Garzón-Montesinos, C., Falcón-Moya, R., Pastor, A., Martín-García, E., Pizarro, N., Busquets-Garcia, A., Revest, J. M., Piazza, P. V., Bosch, F., Dierssen, M., de la Torre, R., Rodríguez-Moreno, A., Maldonado, R., & Ozaita, A. (2019). Cannabinoid type-1 receptor blockade restores neurological phenotypes in two models for Down syndrome. *Neurobiology of Disease*, *125*, 92–106. https://doi.org/10.1016/j.nbd.2019.01.014

Niyuhire, F., Varvel, S. A., Martin, B. R., & Lichtman, A. H. (2007). Exposure to Marijuana Smoke Impairs Memory Retrieval in Mice. *Journal of Pharmacology and Experimental Therapeutics*, *322*(3), 1067–1075. https://doi.org/10.1124/jpet.107.119594

Pacher, P., Bátkai, S., & Kunos, G. (2006). The Endocannabinoid System as an Emerging Target of Pharmacotherapy. *Pharmacological Reviews*, *58*(3), 389–462. https://doi.org/10.1124/pr.58.3.2

Puighermanal, E., Busquets-Garcia, A., Maldonado, R., & Ozaita, A. (2012). Cellular and intracellular mechanisms involved in the cognitive impairment of cannabinoids. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *367*(1607), 3254–3263. https://doi.org/10.1098/rstb.2011.0384

Puighermanal, E., Marsicano, G., Busquets-Garcia, A., Lutz, B., Maldonado, R., & Ozaita, A. (2009). Cannabinoid modulation of hippocampal long-term memory is mediated by mTOR signaling. *Nature Neuroscience*, *12*(9), 1152–1158. https://doi.org/10.1038/nn.2369

Reibaud, M., Obinu, M. C., Ledent, C., Parmentier, M., Böhme, G. A., & Imperato, A. (1999). Enhancement of memory in cannabinoid CB1 receptor knock-out mice. *European Journal of Pharmacology*, *379*(1), R1-2.

Roberts, C. J., & Hillard, C. J. (2020). Peripherally restricted cannabinoid type 1 receptor (CB1R) antagonist, AM6545, potentiates stress-induced hypothalamic–pituitary–adrenal axis activation via a non-CB1R mechanism. In *Endocrine* (Vol. 72, Issue 1). Springer. https://doi.org/10.1007/s12020-020-02549-1

Roozendaal, B., & McGaugh, J. L. (2011). Memory modulation. *Behavioral Neuroscience*, *125*(6), 797–824. https://doi.org/10.1037/a0026187

Rubinov, M., & Sporns, O. (2010). Complex network measures of brain connectivity: Uses and interpretations. *NeuroImage*, *52*(3), 1059–1069. https://doi.org/10.1016/j.neuroimage.2009.10.003

Sala-Llonch, R., Peña-Gómez, C., Arenaza-Urquijo, E. M., Vidal-Piñeiro, D., Bargalló, N., Junqué, C., & Bartrés-Faz, D. (2012). Brain connectivity during resting state and subsequent working memory task predicts behavioural performance. *Cortex*, *48*(9), 1187–1196. https://doi.org/10.1016/j.cortex.2011.07.006

Sara, S. J. (2009). The locus coeruleus and noradrenergic modulation of cognition. *Nature Reviews Neuroscience*, *10*(3), 211–223. https://doi.org/10.1038/nrn2573

Silva, B. A., Burns, A. M., & Gräff, J. (2019). A cFos activation map of remote fear memory attenuation. *Psychopharmacology*, *236*(1), 369–381. https://doi.org/10.1007/s00213-018-5000-y

Takahashi, R. N., Pamplona, F. A., & Fernandes, M. S. (2005). The cannabinoid antagonist SR141716A facilitates memory acquisition and consolidation in the mouse elevated T-maze. *Neuroscience Letters*, *380*(3), 270–275. https://doi.org/10.1016/j.neulet.2005.01.049

Talley, C. E., Kahn, S., Alexander, L. J., & Gold, P. E. (2000). Epinephrine fails to enhance performance of food-deprived rats on a delayed spontaneous alternation task. *Neurobiology of Learning and Memory*, *73*(1), 79–86. https://doi.org/10.1006/nlme.1999.3920

Tam, J., Vemuri, V. K., Liu, J., Bátkai, S., Mukhopadhyay, B., Godlewski, G., Osei-Hyiaman, D., Ohnuma, S., Ambudkar, S. V., Pickel, J., Makriyannis, A., & Kunos, G. (2010). Peripheral CB1 cannabinoid receptor blockade improves cardiometabolic risk in mouse models of obesity. *Journal of Clinical Investigation*, *120*(8), 2953–2966. https://doi.org/10.1172/JCI42551

Tanimizu, T., Kono, K., & Kida, S. (2017). Brain networks activated to form object recognition memory. *Brain Research Bulletin*. https://doi.org/10.1016/j.brainresbull.2017.05.017

Vázquez-Oliver, A., Brambilla-Pisoni, C., Domingo-Gainza, M., Maldonado, R., Ivorra, A., & Ozaita, A. (2020). Auricular transcutaneous vagus nerve stimulation improves memory persistence in naïve mice and in an intellectual disability mouse model. *Brain Stimulation*, *13*(2), 494–498. https://doi.org/10.1016/j.brs.2019.12.024

Wade, M. R., Degroot, A., & Nomikos, G. G. (2006). Cannabinoid CB1 receptor antagonism modulates plasma corticosterone in rodents. *European Journal of Pharmacology*, *551*(1–3), 162–167. https://doi.org/10.1016/j.ejphar.2006.08.083

Winkler, A. M., Ridgway, G. R., Webster, M. A., Smith, S. M., & Nichols, T. E. (2014). Permutation inference for the general linear model. *NeuroImage*, *92*(100), 381–397. https://doi.org/10.1016/j.neuroimage.2014.01.060

Wolff, M. C., & Leander, J. D. (2003). SR141716A, a cannabinoid CB1 receptor antagonist, improves memory in a delayed radial maze task. *European Journal of Pharmacology*, *477*(3), 213–217.

Yang, C., Liu, J.-F., Chai, B.-S., Fang, Q., Chai, N., Zhao, L.-Y., Xue, Y.-X., Luo, Y.-X., Jian, M., Han, Y., Shi, H.-S., Lu, L., Wu, P., & Wang, J.-S. (2013). Stress

within a restricted time window selectively affects the persistence of long-term memory. *PloS One*, *8*(3), e59075. https://doi.org/10.1371/journal.pone.0059075

Yonelinas, A. P., Ranganath, C., Ekstrom, A. D., & Wiltgen, B. J. (2019). A contextual binding theory of episodic memory: systems consolidation reconsidered. In *Nature Reviews Neuroscience* (Vol. 20, Issue 6, pp. 364–375). Nature Publishing Group. https://doi.org/10.1038/s41583-019-0150-4

Zanettini, C., Panlilio, L. V, Alicki, M., Goldberg, S. R., Haller, J., & Yasar, S. (2011). Effects of endocannabinoid system modulation on cognitive and emotional behavior. *Frontiers in Behavioral Neuroscience*, *5*, 57. https://doi.org/10.3389/fnbeh.2011.00057

Zhou, M., Bakker, E. H. M., Velzing, E. H., Berger, S., Oitzl, M., Joëls, M., & Krugers, H. J. (2010). Both mineralocorticoid and glucocorticoid receptors regulate emotional memory in mice. *Neurobiology of Learning and Memory*, *94*(4), 530–537. https://doi.org/10.1016/j.nlm.2010.09.005

Zimmer, A., Zimmer, A. M., Hohmann, A. G., Herkenham, M., & Bonner, T. I. (1999). Increased mortality, hypoactivity, and hypoalgesia in cannabinoid CB1 receptor knockout mice. *Proceedings of the National Academy of Sciences of the United States of America*, *96*(10), 5780–5785. https://doi.org/10.1073/pnas.96.10.5780

Figures

Fig. 1. Pharmacological or genetic inhibition of CB1R improves memory persistence in the novel object-recognition test. a Discrimination index values obtained at 3 h, 24 h and 48 h after the familiarization phase (n = 5 - 8) **b-d** Discrimination index values in NORT at 48 h **b** after acute post-familiarization treatment with vehicle (VEH) or rimonabant (RIM) (1 mg/kg) (n = 7 - 11), **c** in CB1HZ and WT mice (n = 6 - 8), **d** after acute post-familiarization treatment with vehicle (VEH) or AM6545 (1 mg/kg) (n = 7 - 8). Data are expressed as mean ± s.e.m. *p < 0.05, **p < 0.01 by one-way ANOVA followed by Tukey *post hoc* or Student's t-test.

Fig. 2. AM6545 enhances novel object-recognition memory through a peripheral β-adrenergic mechanism. **a** Discrimination index values obtained in the NORT performed at 48 h of adrenalectomized (ADX) or naive mice treated with vehicle (VEH) or AM6545 (1 mg/kg) (n = 6 - 8). **b** Schematic representation of pre-treatment and treatment after the familiarization phase. **c, d** Discrimination index values obtained in the NORT performed at 48 h of mice treated with vehicle (VEH) or AM6545 (1 mg/kg) after pre-treatment with **c** vehicle (VEH) or mifepristone (50 mg/kg) (n = 6 - 7) and **d** saline or sotalol (10 mg/kg) (n = 8 - 10). **e** Discrimination index values in WT or DBH-CB1KO mice in NORT at 48 h **f** and after saline or sotalol (10 mg/kg) treatment (n = 6 - 8) **g** Discrimination index values for mice pre-treated with saline or sotalol (10 mg/kg) prior to rimonabant (RIM) (1 mg/kg) or vehicle (VEH) in the NORT at 48 h (n = 9 - 11) Data are expressed as mean ± s.e.m. *p < 0.05 (treatment effect) #p < 0.05, ##p < 0.01 (pre-treatment effect) by two-way ANOVA test followed by Tukey *post hoc*.

Fig. 3. AM6545 memory improvement depends on vagus nerve activation. a Glucose blood levels for 120 min after acute vehicle (VEH) or AM6545 (1 mg/kg) administration in mice (n = 6 – 7) **b,c** Discrimination index values in NORT after acute vehicle (VEH) or AM6545 (1 mg/kg) and CNO (3 mg/kg) administration in mice infected with **b** AAV5-mCherry-control or **c** AAV5-mCherry-hM4Di in the vagus nerve (n = 6 – 7).

Fig. 4. Circos plot and z-score analysis of c-Fos expression correlation analysis in different brain regions after acute AM6545 administration. Network graphs of c-Fos expression based on the Pearson coefficient in **a** vehicle and **b** AM6545 (1 mg/kg) between different brain areas, including prelimbic cortex (PL), infralimbic cortex (IL), *cornu ammonis* 1 (CA1), dentate gyrus (DG), *cornu ammonis* 3 (CA3), *locus coeruleus* (LC), basal amygdala (BA), central amygdala (CeA) and lateral amygdala (LA). Significant correlations are delimited with black lines. **c,d** Z-score values of the **c** positive and **d** negative Pearson *r* correlation coefficients between the different brain regions analysed.

Fig. 5. Effects of AM6545 administration on brain functional connectivity by resting state functional magnetic resonance imaging (rsfMRI). a Whole brain local efficiency and **b-d** nodal efficiencies of **b** left frontal cortex, **c** right hippocampus and **d** right globus pallidus. **e** Statistical map showing significant differences between connectivity of posterior brainstem with the rest of the brain in vehicle and AM6545 treated animals (comparison vehicle>AM6545; p<0.01). Data are expressed as mean ± s.e.m. *p < 0.05, **p < 0.01, ***p < 0.001.

Fig. 6. Acute AM6545 treatment increases central noradrenergic activity. a Percentage of mean firing rate in the LC after acute vehicle (VEH) or AM6545 (1 mg/kg) administration respect to baseline values (n = 2 - 3) **b** Percentage of extracellular norepinephrine (NE) levels in the hippocampus after acute vehicle (VEH) or AM6545 (1 mg/kg) administration respect to baseline values (n = 6 - 7). The arrow indicates the time of administration. **c** Discrimination index values and **d** total exploration time obtained in the NORT performed at 48 h mice treated with AM6545 (1 mg/kg) or vehicle (VEH) after bilateral intrahippocampal injection of saline or propranolol (1 μg/μl 0.5 μl per side) (n = 5 - 6). Data are expressed as mean ± s.e.m. For microdialysis *p < 0.05 two-way repeated measures ANOVA followed by Tukey post hoc was performed. For NORT **p < 0.01 (treatment effect) #p < 0.05 (pre-treatment effect) by two-way ANOVA test followed by Tukey *post hoc*.

Supplementary Fig. 1. Total exploration times in NORT after pharmacological or genetic CB1R inhibition. a Total exploration times obtained at 3 h, 24 h and 48 h after the familiarization phase (n = 5 - 8) **b-d** Total exploration times in NORT at 48h **b** after acute post-familiarization treatment with vehicle (VEH) or rimonabant (RIM) (1 mg/kg) (n = 7 - 11) **c** in CB1HZ and WT mice (n = 6 - 8) **d** after acute post-familiarization treatment with vehicle (VEH) or AM6545 (1 mg/kg) (n = 7 - 8). Data are expressed as mean ± s.e.m.

Supplementary Fig. 2. Locomotor activity after acute AM6545 administration. Total activity in locomotor activity boxes for 120 min by mice treated with vehicle (VEH) or AM6545 (1 mg/kg) (n = 6). Data are expressed as mean ± s.e.m.

Supplementary Fig. 3. Total exploration times of systemically or peripheral CB1R blockade are not affected. a Total exploration times obtained at 3 h, 24 h and 48 h after the familiarization phase (n = 5 - 8) **b-d** exploration times in NORT at 48h **b** after acute post-familiarization treatment with rimonabant vehicle (VEH) or rimonabant (RIM) (1 mg/kg) (n = 7 - 11) **c** in HZCB1R and WT mice (n = 6 - 8) **d** after acute post-familiarization treatment with vehicle (VEH) or AM6545 (1 mg/kg) (n = 7 - 8). Data are expressed as mean ± s.e.m.

Supplementary Fig. 4. Chemogenetic targeting of the vagus nerve does not modify the behavioral responses in the NORT. a Discrimination index values and **b** total exploration times in NORT at 24 h after the familiarization phase and after CNO (3 mg/kg) and vehicle (VEH) or AM6545 (1 mg/kg) in mice infected with AAV-hm4Di in the vagus nerve and i.p. CNO (3 mg/kg) administration (n = 10 - 11) **c,d** total exploration times in NORT at 48 h after CNO (3 mg/kg) and vehicle (VEH) or AM6545 (1 mg/kg) administration in mice infected with **c** AAV-mCherry (n = 6 - 8) or **d** AAV-hM4Di (n = 6 - 8). **e** Representative images of mCherry expression detected in nodose ganglion of mice after AAV5-mCherry injection in the vagus nerve (left image scale bar = 100μm, right image scale bar = 30μm). Data are expressed as mean ± s.e.m.

Supplementary Fig. 5. Effects of acute AM6545 treatment in c-Fos expression in specific regions of the CNS. Representative immunohistochemical staining of c-Fos positive cells 90 min after acute vehicle (VEH) or AM6545 (1 mg/kg) administration in mice (n = 5-9). Brain regions evaluated: prelimbic cortex (PL), infralimbic cortex (IL), *cornu ammonis* 1 (CA1), dentate gyrus (DG), *cornu ammonis* 3 (CA3), *locus coeruleus* (LC), basal amygdala (BA), central amygdala (CeA) and lateral amygdala (LA). Data are expressed as mean ± s.e.m.

Supplementary Fig. 6. Correlation heat map of c-Fos expression in different brain regions after acute AM6545 administration. Pearson correlation analysis of c-Fos expression in **a** vehicle and **b** AM6545 (1 mg/kg) between different brain areas, including prelimbic cortex (PL), infralimbic cortex (IL), *cornu ammonis* 1 (CA1), dentate gyrus (DG), *cornu ammonis* 3 (CA3), *locus coeruleus* (LC), basal amygdala (BA), central amygdala (CeA) and lateral amygdala (LA). *p < 0.05, **p < 0.01, ***p < 0.001.

Supplementary Fig. 7. Brainstem connectivity map in AM6545-treated mice: only areas where BOLD signal is negatively correlated to the brainstem signal are shown. Color represents z-Fisher of the correlation coefficient.

Supplementary Fig. 8. Acute AM6545 administration does not modify other monoamine extracellular levels in the hippocampus. a-b Extracellular **a** serotonin (5-HT) and **b** dopamine (DA) levels in the hippocampus after acute AM6545 (1 mg/kg) or vehicle (VEH) administration respect to baseline values (n = 6 - 7). Arrow indicates the time of administration. Points represent mean ± s.e.m and are expressed as percentages of baseline.

Supplementary Fig. 9. Intrahippocampal propranolol injection. a Cannula placement in the dorsal hippocampus. A brain coronal section from a representative mouse showing cannula placement in the dorsal hippocampus. Brain slices were stained with cresyl violet. **b** Discrimination index values of different doses of intrahippocampal propranolol infusion or saline in the NORT at 24 h (n = 4). Data are expressed as mean ± s.e.m. **p < 0.01 by one-way ANOVA followed by Tukey *post hoc*.