The Myogenesis Program Drives Clonal Selection and Drug Resistance in Rhabdomyosarcoma

Anand G. Patel[1,2]*, Xiang Chen[3]*, Xin Huang[3]*, Michael R. Clay[4], Natalia Komorova[5], Matthew J. Krasin[6], Alberto Pappo[2], Heather Tillman[7], Brent A. Orr[7], Justina McEvoy[1], Brittney Gordon[1], Kaley Blankenship[2], Colleen Reilly[1], Xin Zhou[3], Jackie L. Norrie[1], Asa Karlstrom[1], Jiyang Yu[3], Dominik Wodarz[8], Elizabeth Stewart[1,2] and Michael A. Dyer[1]

Departments of [1]Developmental Neurobiology, [2]Oncology, [3]Computational Biology, [6]Radiation Oncology and [7]Pathology at St. Jude Children's Research Hospital, Memphis, Tennessee 38105, USA

[4]Department of Pathology, University of Colorado School of Medicine, Aurora, Colorado 80045, USA

[5]Department of Mathematics, University of California, Irvine, California 92697, USA

[8]Department of Population Health and Disease Prevention, Program in Public Health, Susan and Henry Samueli College of Health Sciences, University of California, Irvine, California 92697 USA

*These authors contributed equally

Correspondence and requests for materials should be addressed to:
Michael A. Dyer
Department of Developmental Neurobiology, MS-323
St. Jude Children's Research Hospital
262 Danny Thomas Place, Memphis, TN 38105-3678, USA
Phone: (901) 595-2257
Fax: (901) 595-3143
E-mail: michael.dyer@stjude.org

Abstract

Rhabdomyosarcoma (RMS) is a pediatric cancer with features of skeletal muscle; patients with

unresectable or metastatic RMS fare poorly due to high rates of disease recurrence. Here, we use

single cell and single nucleus RNA-sequencing to show that RMS tumors recapitulate the

spectrum of embryonal myogenesis. Using matched patient samples from a clinical trial and

orthotopic patient-derived xenografts (O-PDXs), we show chemotherapy eliminates the most

proliferative component with features of myoblasts; after treatment, the quiescent immature

population with features of paraxial mesoderm expands to reconstitute the developmental

hierarchy of the original tumor. We discovered that this paraxial mesoderm population is

dependent on EGFR signaling and is sensitive to EGFR inhibitors. Taken together, this data

serves as a proof-of-concept that targeting each developmental state in RMS is an effective

strategy for improving outcomes by preventing disease recurrence.

Introduction

Rhabdomyosarcoma (RMS) is the most common pediatric soft tissue sarcoma and has molecular, cellular and histopathologic features of developing skeletal muscle[1–3]. The alveolar form of RMS (ARMS) is more differentiated than the embryonal form (ERMS) and each subtype has distinct genomic and epigenomic landscapes[2,4,5]. For newly diagnosed RMS patients, the overall survival rate is 70% using multiagent chemotherapy combined with radiation and/or surgical resection[6,7]. Unfortunately, a subset of patients experience disease recurrence after treatment completion; for those patients, overall survival rate drops below 20%[8]. Genomic studies have shown that clonal selection occurs with disease recurrence, but no recurrent genetic lesion has been identified that contributes to survival of the rare clones of cells for RMS[2,5,9]. This raises the possibility that other, non-genetic mechanisms may contribute to drug resistance and disease recurrence in RMS. To explore this possibility, we performed single cell (sc) and single nucleus (sn) RNA-seq of RMS patient tumors and matched orthotopic patient-derived xenografts (O-PDXs). We also performed lentiviral barcode labeling to trace the clonal expansion of individual tumor cells during normal growth and in response to treatment. Taken together, these studies showed that individual tumor cells transition through myogenesis and the underlying myogenic developmental hierarchy contributes to clonal selection with treatment. We used the developmental program in RMS to identify therapeutic vulnerabilities that could be exploited to reduce disease recurrence. Overall, this study reveals a developmental hierarchy with RMS and introduces a novel approach to treating pediatric cancers, wherein targeting specific developmental states that are destined to persist during therapy can be used to improve treatment efficacy.

Results

RMS tumors have developmental heterogeneity

Skeletal muscle develops from the mesodermal cells of the somites during embryogenesis and

undergoes stepwise differentiation, which is typified by the expression of myogenic regulatory

factors[10,11] (MRFs; Fig. 1A,B). RMS tumors have features of skeletal muscle including

myofibers and heterogenous expression of proteins such as myogenin (MYOG)[3,12]. To further

investigate the transcriptomic heterogeneity within RMS, we performed droplet-based single-cell

RNA-sequencing (scRNA-seq) (Extended Data Tables 1 and 2). We obtained fresh ERMS and

ARMS patient tumor tissue (Fig. 1C,D) following surgical resection and generated single-cell

suspensions (>90% viable cells) for 3'-directed scRNA-seq (10x Genomics). Inference of

somatic copy number alterations was used to distinguish malignant cells from non-malignant

cells[13,14] (Extended Data Fig. 1).

13   Single-cell analysis showed there were distinct populations of cells expressing

transcription factors characteristic of paraxial mesoderm (*MEOX2, PAX3*), myoblasts (*MYF5,*

MSC) and myocytes (*MYOG, MEF2C*) (Fig. 1E,F and data not shown). The proportion of *MYOG*

expressing cells in the scRNA-seq data was consistent with the proportion measured by

immunohistochemical staining (IHC) (Fig. 1C-F). The ARMS sample had fewer tumor cells

expressing the early paraxial mesoderm MRF MEOX2 (2.1%) than the ERMS sample (29.4%),

and more cells expressing the late myocyte MRF MYOG (75.2% versus 25.6%; Fig. 1E,F). RNA

velocity analysis, which leverages the simultaneous measurement of spliced and unspliced RNA

transcripts to generate a model of the future state of cells[15], showed unidirectional transit of cells

from the paraxial mesoderm through myoblast to the myocyte state in the ERMS tumor (Fig.

1G,H). Non-malignant cells including monocytes, fibroblasts, lymphocytes and vascular

endothelial cells were readily identifiable in our scRNA-seq dataset (Extended Data Fig. 1).

The rarity of childhood cancers limits the ability to obtain fresh tissue samples for

scRNA-seq. To increase the number of evaluable tumors, we validated single-nucleus RNA-

sequencing (snRNA-seq) of frozen tumor tissue and adapted our computational pipeline to

accommodate data generated from snRNA-seq[14]. Specifically, we compared scRNA-seq from

fresh tumors (SJRHB030680_R1 and SJRHB031320_D1) (Fig. 1E,F) to snRNA-seq of matched

frozen tumor specimens (Extended Data Fig. 2). As shown previously for neuroblastoma[14], we

were able to recover more cells of the tumor microenvironment (TME) by snRNA-seq compared

to data generated by scRNA-seq (Extended Data Fig. 2). To extend our single cell transcriptional

profiling, we performed snRNA-seq on 18 RMS tumors (12 ERMS and 6 ARMS) (Extended

Data Tables 1 and 2). In total, 122,731 nuclei were analyzed from the patient tumors. As for the

fresh tumors, copy number inference was used to distinguish malignant nuclei (111,474) from

the normal nuclei (11,257) in the TME. The malignant nuclei were integrated using Conos, an

approach that leverages inter-sample mappings to generate a unified graph for the identification

of communal cell clusters[16] (Fig. 2A). Leiden clustering identified 7 clusters, which we

combined into 1 mesoderm, 4 myoblast and 2 myocyte signature groups based on expression of

MRFs (Fig. 2B). The 2 myocyte populations were distinguished by expression of genes involved

in muscle differentiation and function (Extended Data Table 3). The 4 myoblast populations

were distinguished by ribosomal genes ($p=4.3 \times 10^{-40}$) and muscle differentiation genes

($p=0.0005$) (Extended Data Table 3). We identified 954 differentially expressed genes, of which

945 were cluster-type specific (Extended Data Table 3). Extracellular matrix and cell adhesion

pathways were enriched in the paraxial mesoderm-like tumor cells, ribosome biosynthesis

pathways were enriched in the myoblast-like cells and pathways involved in muscle function

were enriched in the myocyte-like cells (Extended Data Table 3). While all the tumors had a

mixture of cells with mesoderm, myoblast, and myocyte signatures, ARMS tumors contained

significantly fewer cells with the mesodermal gene expression signature (p=0.008; unpaired t-

test) and were skewed towards the myocyte signature (Fig. 2C and Extended Data Fig. 3). One

ERMS tumor (SJRHB010928_R1) was notable in that it contained a majority (97%) of tumor

cells with the mesodermal signature (Extended Data Fig. 3A,B). This sample was collected

during treatment (Extended Data Table 1) suggesting that mesodermal cells are more resistant to

treatment than the other cell populations. The proliferating cells were significantly enriched in

the myoblast population (p<0.0001; one-way ANOVA with multiple comparisons) (Fig. 2E,F).

All data can be viewed using an interactive viewer at: https://pecan.stjude.cloud/static/RMS-

scrna-atlas-2020/.

The same approach was used to cluster the non-malignant cells within the TME

(Extended Data Figure 3C-F). Comparing normal cell populations between ERMS and ARMS

showed that fibroblasts in ARMS were significantly enriched in pathways involved in

extracellular matrix synthesis and organization as well as cell adhesion. In addition, *SFRP2* and

SFRP4 were significantly ($p<1\mathrm{x}10^{-90}$) enriched in fibroblasts from ARMS (45% and 56% of

cells, respectively) relative to ERMS (1% and 3%, respectively) (Extended Data Table 4). A

previous pan-cancer analysis showed that *SFRP2* and *SFRP4* represent a tightly regulated

transcriptional program in cancer stroma that correlates with poor prognosis, EMT and

angiogenesis across multiple cancers[17]. The *HLA-A,B,C,E* and *B2M* and *CD74* genes were

significantly upregulated in lymphocytes from ARMS and *HLA-DRA, DRB1* and *DPB1* were

significantly upregulated in monocytes from ERMS (Extended Data Table 4).

We next investigated the spatial heterogeneity of malignant subpopulations using single

and multiplex immunohistochemistry (IHC) on 12 patient tumor specimens. Consistent with our

transcriptomic findings, there was heterogenous expression of MEOX2, MYF5 and MYOG

protein (Fig. 2G). The proportion of immunopositive cells were correlated with the proportion of

each population from the sc/snRNA-seq (Fig. 2H). Double IHC showed that these proteins were

expressed in a mutually exclusive pattern consistent with the distinct clusters of mesoderm,

myoblast and myocyte populations in RMS tumors from sc/snRNA-seq (Fig. 2I,J and data not

shown).

Developmental indexing of RMS using embryonic snRNA-seq data

To extend our analysis of the developmental trajectory of RMS beyond MRFs, we analyzed our

RMS data within the context of early muscle development using a single-nucleus atlas of

organogenesis from mouse embryos at E9.5, E10.5, E11.5, E12.5, and E13.5[18]. We extracted

data from the skeletal muscle lineage and performed trajectory analysis on half of the data to

generate a training dataset (Fig. 3A-D). We then adapted latent cellular analysis (LCA)[19] to

calculate the similarity in the latent cellular space between cells in the remaining half of the

skeletal muscle dataset to cells used for training; a normalized muscle developmental index was

then calculated for each nucleus within the validation dataset (Fig. 3E,F). The developmental

index increased with embryonic age as expected within the validation dataset (Fig. 3E,F).

Using this unsupervised developmental indexing approach, we confirmed that individual

RMS tumors have cellular heterogeneity that reflects normal myogenesis. For example, in

SJRHB030680_R1, an ERMS tumor, we identified a broad range of developmental indices

within the malignant components of the tumor (Fig. 3G). In contrast, in SJRHB031320_D1, an

ARMS tumor, the range of developmental indices was narrower and more skewed toward later

stages of myogenesis (Fig. 3H). Using our entire patient cohort of 18 tumors, we were able to

generalize these findings to RMS tumors - ERMS tumors had a wide diversity of developmental

indices while ARMS tumors narrowly centered with developmental indices from later stages of

murine myogenesis (Fig. 3I).

O-PDXs and organoids recapitulate clonal heterogeneity in RMS

We have previously established a panel of RMS O-PDXs and shared those models through the

Childhood Solid Tumor Network[9]. These O-PDXs encompass the clinical and molecular

diversity of RMS, and have previously undergone bulk genomic, transcriptomic, proteomic and

epigenomic analyses[2,4,9]. We expanded our single-cell transcriptomic profiling to include the O-

PDXs that correspond to the 18 patient tumors profiled here (Extended Data Table 2 and

https://pecan.stjude.cloud/static/RMS-scrna-atlas-2020/). We performed the same analyses,

including developmental indexing (Fig. 3J). All 3 cell types (mesoderm, myoblast, and myocyte)

were preserved in the O-PDXs in the snRNA-seq and IHC analysis (Extended Data Fig. 4 and

data not shown). As expected, the O-PDXs lacked normal cells from the patient TME but

contained infiltration of murine monocytes (Extended Data Fig. 4). The patient tumor that was

collected during treatment and was enriched in cells with the mesodermal signature

(SJRHB010928_R1) re-established the developmental hierarchy in the O-PDX

(SJRHB010928_X1) (Extended Data Fig. 4).

　　　　To complement the O-PDXs, we also evaluated the transcriptomic heterogeneity of ex

vivo organoids derived from the O-PDXs (Supplemental Methods). Malignant cells within

organoids shared the cellular diversity seen in the originating patient tumor and O-PDX by single

cell transcriptional profiling (Extended Data Fig. 4). IHC for MEOX2, MYF5 and MYOG for

the organoids showed a similarity to their matched patient tumor and corresponding O-PDX

(Extended Data Fig. 4 and data not shown).

RMS cells transition through developmental states

RNA velocity analysis (see Fig. 1G,H) suggests that individual RMS tumor cells may transition

through developmental stages from mesoderm to myoblast and myocyte (Fig. 4A). Alternatively,

it is possible that there are distinct clones of cells that are restricted to their developmental stage

(Fig. 4B). To distinguish between these two possibilities, we used a lentiviral barcoding library

that incorporates a unique oligonucleotide barcode into the 3'-untranslated region of blue

fluorescent protein (BFP)[20,21] (Fig. 4C,D). We infected 15 of the O-PDX models with the

barcode library and analyzed the barcode distribution in vivo by scRNA-seq. Following scRNA-

seq library generation, the barcode is retrievable by a separate PCR amplification step. In each of

the tumors that we analyzed, individual barcodes were found across all tumor cell types

(mesoderm, myoblast and myocytes) (Fig. 4E-G and Extended Data Table 5). Taken together,

these lineage tracing data, RNA-velocity analyses and genetic clonal analysis are consistent with

a model in which individual ERMS tumor cells can transition through developmental stages. The

same was true for ARMS tumors but the population of cells with paraxial mesoderm gene

expression signature was lower so some barcodes were found only in the myoblast and myocyte

population (Extended Data Table 5).

Tumor cell heterogeneity reflects differential enhancer activity

Several of the MRF genes that are turned on and off as cells transition through developmental

stages have core regulatory circuit super-enhancers[4] (CRC-SEs) (Fig. 4H and Extended Data

Table 6). For example, *MEOX2* and *NFIX* (mesoderm), *PAX7* and *CREB5* (myoblast) and

FOXO1 and *SOX6* (myocyte) each have CRC-SEs (Extended Data Table 6). To determine if the

chromatin accessibility of those CRC-SEs changes as individual cells transition through the

myogenic differentiation program, we performed droplet-based single-cell assay of transposase-

accessible chromatin sequencing (scATAC-seq) on 7 O-PDX tumors. We integrated scATAC-

seq and scRNA-seq profiles to investigate the chromatin accessibility of CRC-SEs for MRFs in

developmentally distinct subpopulations (Fig. 4H-J and Extended Data Table 6). Transferring

cell labels between scRNA-seq data and scATAC-seq data in SJRHB010927_X1 enabled us to

identify cell-type specific enhancer regions in *MYOD1*, *MSC*, *MEOX2* and several other

myogenic genes (Fig. 4H-J and Extended Data Table 6). These regions correspond to previously

reported core regulatory circuit domains for those genes[4,22]. Analysis of all 7 O-PDX tumors

showed CRC-SEs that change in their chromatin accessibility in tumor cells with mesoderm

(*MEOX2, SMAD3*), myoblast (*CREB5, PAX7*), and myocyte (*MYOD1, FOXO1*) features

(Extended Data Table 6 and Extended Data Fig. 5). Collectively, these scATAC-seq studies

indicate that heterogeneity of developmental states within RMS tumors is reflected in chromatin

dynamics for myogenic CRC-SEs and genes.

The mesoderm-like RMS cells are drug resistant

Current chemotherapeutic regimens for RMS include drugs that target proliferating cells. The

myoblast population has the highest proportion of dividing cells in the patient tumors, the O-

PDXs, and the ex vivo organoids (Fig. 5A,B and data not shown). In a pair of matched ERMS

samples obtained before and during treatment, SJRHB000026_R2 and SJRHB000026_R3

(Extended Data Fig. 3A), we noted that the post-treatment sample was skewed towards

mesoderm signature-expressing cells (28.6% post-treatment versus 3.4% pre-treatment) with a

concomitant reduction in cells expressing the myocyte signature (1.4% post-treatment versus

31.4% pre-treatment). Additionally, one ERMS patient tumor, SJRHB010928_R1, was obtained

during treatment with fewer than 5% viable cells by histology; in this sample, the majority

(96.8%) of remaining viable cells expressed the mesoderm signature (Extended Data Fig. 3).

Taken together, these data suggest that the myoblast population may be more sensitive to

chemotherapy and the mesoderm-like population is more likely to survive treatment. To

investigate this trend further, we evaluated matched formalin-fixed paraffin embedded (FFPE)

tissue from 11 patients obtained at diagnosis and mid-treatment on a single therapeutic clinical

trial, RMS13 (NCT01871766). We quantitated the number of cells in each sample expressing

MEOX2 and MYOG (Extended Data Table 7). There was a significant enrichment in MEOX2

immunopositive cells in the post-treatment tumors relative to the matched pre-treatment RMS

samples and a corresponding decrease in MYOG immunopositive cells (Fig. 5C).

To model clonal selection in the laboratory, we generated longitudinal samples from

repeat biopsy of O-PDXs treated with a standard drug combination used to treat patients with

RMS (vincristine (VCR) and irinotecan (IRN)) at clinically relevant doses and schedules[4,9] (Fig.

5D). For each O-PDX (SJRHB000026_X1, SJRHB013758_X1, SJRHB011_X,

SJRHB013757_X1 and SJRHB013759_X14), biopsies were performed at multiple timepoints

(before treatment (day 0), day 3, day 7, day 14 and day 21 of the first course) when sufficient

tumor was present to sample (Fig. 5E,F). We also collected tumor biopsies after the tumors

recurred. A portion of each biopsy underwent formalin-fixation for IHC staining for MEOX2,

MYF5 and MYOG (Fig. 5G,H and data not shown). The remaining biopsy portion was utilized

for quantitative RT-PCR for 21 genes expressed in mesoderm, myoblast and myocyte-like RMS

tumor cells or snRNA-seq. In total, 250 biopsies were collected and 6,480 qRT-PCR reactions

were performed (Extended Data Table 8-13). As in patient samples, the myoblast and myocyte

populations were sensitive to treatment and the mesoderm tumor cells population was enriched

(Fig. 5I-K and Extended Data Table 8-13). Moreover, the normally quiescent mesoderm-like

cells re-entered the cell cycle to initiate myogenesis (Fig. 5K).

Taken together, our data suggest that ERMS tumor cells transition through distinct states

that represent progressive stages of myogenesis. These different states (paraxial mesoderm,

myoblast, myocyte) have distinct proliferation properties and differential sensitivity to

chemotherapy. To further refine our understanding of the cellular heterogeneity of ERMS

tumors, their developmental trajectory and clonal selection with treatment, we developed a

mathematical model that follows the fate of cells in both 3-dimensional space and time.

Experimentally determined barcode distribution in each compartment was used to develop the

model (Fig. 5L), and barcode diversity was tracked over time. We assumed that upon cell

division, cells maintain their barcodes and we included barcoded and non-barcoded cells to

reflect the in vivo experiments. The relative proportion of different division types (self-

renewing/differentiating) in the mesodermal compartment determines whether the tissue remains

in homeostasis and influences the degree of clonal diversity loss over time. To parameterize the

model, we used experimental data from 10 barcoded ERMS xenografts. The fraction of dividing

cells and distribution of cells across compartments was determined from the sc/snRNA-seq data.

Our ERMS model predicts a decrease in clonal diversity (as measured by barcode diversity) over

time and clonal selection with treatment for individual tumors (Fig. 5M-P). To test this

experimentally, we performed scRNA-seq on a barcoded ERMS tumor (SJRHB000026_X1)

after initial labeling and a subsequent passage in mice with and without clinically relevant

chemotherapy (vincristine+irinotecan). As predicted by the three-compartment model, there was

a decrease in clonal diversity over time and clonal selection with treatment (Fig. 5O-Q).

Additional iterations of modeling and comparison to in vivo barcode distribution data are

consistent with differential cytotoxicity across the cellular populations (mesoderm, myoblast,

myocyte). In particular, a subset of the mesoderm-like cells are proliferating thereby making

them sensitive to chemotherapy. Based on our model, partial elimination of the mesoderm-like

population is required to account for the clonal selection we observe in vivo in mice and in

patients.

EGFR is a therapeutic vulnerability in paraxial mesoderm RMS cells

Having shown that the paraxial mesoderm RMS cells are more quiescent and drug resistant than

the myoblast population, we set out to identify therapeutic vulnerabilities unique to this

population using a systems biology algorithm, NetBID (data-driven Network-based Bayesian

Inference of Drivers)[23,24]. NetBID, which was originally developed for bulk -omics data, was

adapted to analyze snRNA-seq profiles of our panel of 18 RMS patient tumors. We first used the

SJARACNe algorithm[25] to reverse engineer cell type–specific interactomes for each of the 5

major cell types from the integrated snRNA-seq profiles (Fig. 6A). With a focus on signaling

drivers, we used the cell type–specific interactomes of 2,543 genes/proteins and inferred their

network activities in each nucleus using the NetBID algorithm. We then performed differential

activity analysis to identify cell type–specific therapeutic vulnerabilities in the RMS tumor cells

with the mesodermal signature. EGFR was significantly activated in the mesoderm population

compared to myoblasts ($p=4.4 \times 10^{-135}$) and myocytes ($p=1.8 \times 10^{-174}$) and the network was rewired

as cells transition through the developmental hierarchy (Fig. 6B). EGFR network activity was

also significantly higher in ERMS relative to ARMS ($p=5.4 \times 10^{-36}$) (Fig. 6C,D). These data are

consistent with previous integrated epigenetic/proteomic analyses for differential pathway

activity in ERMS and ARMS[4]. In addition, previous studies have shown heterogenous

expression of EGFR protein in FFPE samples of RMS[26–28]. To validate these data, we performed

IHC for EGFR alone and in combination with markers of each cell population. We used the 5B7

monoclonal antibody which has been previously shown to correlate with EGFR inhibitor

(EGFRi) responsiveness in lung cancer[29]. There was co-localization of EGFR with MEOX2 in 2-

color IHC and EGFR was mutually exclusive with MYOG (Fig. 6E,F).

To determine if EGFR is a therapeutic vulnerability in RMS, we exposed 3D ERMS

organoids that contain all 3 cell populations (Extended Data Fig. 6) to two different EGFRi's

(gefitinib and afatinib) with increasing concentrations of SN-38, the active metabolite of

irinotecan (Supplemental Information). The EGFRi's alone had no effect on overall organoid

viability as measured with CellTiter-Glo 3D which is not surprising given the low percentage of

mesoderm-like cells in the organoids (Extended Data Fig. 6). However, when the proliferating

myoblast population was targeted with increasing concentrations of SN-38, the addition of

EGFRi's significantly reduced viability (Extended Data Fig. 6). In a representative ERMS O-

PDX (SJRHB013758_X1), there was a significant improvement in outcome when afatinib or

gefitinib was combined with IRN+VCR (Fig. 6H,I). For a second ERMS O-PDX with a low

percentage of mesoderm-like cells, SJRHB010927_X1, O-PDXs responded robustly to

IRN+VCR therapy alone, reinforcing the importance of including standard-of-care treatment in

preclinical testing (Extended Data Fig. 6).

Discussion

In conclusion, we have discovered that RMS tumor cells can transition through different stages of myogenesis from an immature quiescent paraxial mesoderm state through a highly proliferative myoblast stage and into a more differentiated myocyte state. Not only do cells undergo changes in gene expression during these developmental transitions but super-enhancer chromatin accessibility is also dynamic. While proliferating cells can be identified in tumor cell population in patient tumors and O-PDXs, the most proliferative cells are in the myoblast stage. It is therefore not surprising that broad spectrum chemotherapeutic regimens that target rapidly dividing cells efficiently reduce tumor volume by killing the myoblast-like RMS tumor cells. The immature paraxial mesoderm-like RMS tumor cells are more quiescent and can survive therapy and then expand to repopulate the myogenic lineage found in the primary tumor. These observations are consistent with decades of clinical research showing that different combinations of broad- spectrum chemotherapy or intensification of existing regimens have failed to improve outcomes for children with RMS[6,7]. Those different treatment approaches are killing the rapidly dividing myoblast-like cells and the resistant mesoderm-like cells survive and contribute to disease recurrence. By focusing our investigation on the mesoderm-like cells, we identified a dependence on EGFR that can be exploited with EGFRi in vivo. While there were only a small number of mesoderm-like cells in ARMS tumors, we discovered a dramatic upregulation of EGFR during treatment suggesting that EGFRi's may be useful for both types of rhabdomyosarcoma. Our study shows that treatment for RMS and possibly other pediatric solid tumors should focus on total clonal elimination rather than continuing to target just the most proliferative cell population that makes up the bulk of the tumor. Such an approach may reduce disease recurrence and improve survival and quality of life for children with solid tumors.

Figure Legends

Figure 1: Single-cell RNA-sequencing (scRNA-seq) reveal a developmental hierarchy within RMS. A-B, During fetal myogenesis, mesodermal cells of the somite migrate to form skeletal muscle throughout the body (A). During that migration, these cells undergo stepwise differentiation typified by the transient expression of myogenic regulatory factors[10] (B). **C-D,** Photomicrographs of an embryonal RMS tumor, SJRHB030680_R1 (C) and an alveolar RMS tumor, SHRHB031320_D1. *Left,* H&E staining. *Right,* Myogenin (MYOG) immunohistochemistry (IHC) with 20X magnification. *inset,* 80X magnification. **E-F,** UMAP visualization of 3,973 malignant cells from SJRHB030680_R1 (E) and 2,414 malignant cells from SJRHB031320_D1 (F). Cells are colored based on expression of *MEOX2* (*left*), *MYF5* (*center*), and *MYOG* (*right*). **G-H,** RNA velocity analysis of SJRHB030680_R1 (G) and SJRHB031320_D1 (H). Abbreviations: ERMS, embryonal rhabdomyosarcoma; UMAP, uniform manifold approximation and projection. Scale bars: C,D, 100 μm.

Figure 2: Identification of major cell clusters within patient RMS tumors using single-nucleus RNA-sequencing. A, LargeVis visualization of snRNA-seq of 111,474 nuclei from 18 integrated patient RMS tumors, colored based on sample. **B,** Heatmap showing expression of myogenic regulatory factor expression across seven Leiden clusters. **C,** Boxplot showing the percentage of malignant nuclei within each muscle developmental state for each tumor. The center line demarcates the median value with rectangle showing interquartile range (IQR) between the first and third quartiles. The vertical bars extending from the rectangles indicate maximum and minimum values with the exception of outliers that exceed more than 3 times the IQR. **D-E,** LargeVis visualization of Leiden clustering of snRNA-seq grouped based on

expression of mesoderm, myoblast, or myocyte myogenic regulatory factors (D) or colored by

predicted cell cycle phase (E). **F,** Plot of the proportion of proliferating cells (S/G2/M phase) in

each group, estimated using gene signatures associated with G1, S, and G2/M phases[13]. Circles

are ERMS and squares are ARMS. Center line and rectangle indicate the median and IQR as in

panel (C). Vertical bars indicate the maximum and minimum values with the exception of

outliers that exceed more than 3 times the IQR. **G,** Immunohistochemistry image of an ERMS

tumor, SJRHB013758_D2 stained with antibodies against MEOX2 (left), MYF5 (center) and

MYOG (right). **H,** Quantitation of the percentage of cells positive for MEOX2 (blue), MYF5

(green), or MYOG (red) immunohistochemical staining (*x* axis) compared to percentage of cells

within each developmental state as determined by snRNA-seq (*y* axis). **I-J,** Dual staining of

MEOX2 (purple) and MYOG (brown) within SJRHB013758 (I) with magnified view (J).

Abbreviations: ERMS, embryonal rhabdomyosarcoma; ARMS, alveolar rhabdomyosarcoma.

Scale bars: G, 10 μm.

Figure 3: Developmental indexing of patient RMS tumors and orthotopic patient-derived

xenografts. A, UMAP plot of 1.5 million nuclei from the Mouse Organogenesis Cell Atlas[18],

downsampled to 100,000 nuclei. Clusters are colored based on trajectory. **B,** UMAP plot of

576,560 nuclei from the mesenchymal trajectory with identification of the skeletal myogenesis

sub-trajectory. Nuclei are colored based on Leiden cluster. **C,** UMAP plot of 58,573 nuclei of the

skeletal muscle sub-trajectory with computational clustering that identifies nuclei from early

mesodermal progenitors, paraxial mesoderm, myoblasts, myocytes and myotubes. **D,** Heatmap of

aggregated transcription from each cluster demonstrating expression of myogenic regulatory

factors and additional mesodermal markers. **E,** Violin plot of projected developmental indices of

embryonic skeletal muscle data separated by mouse embryonic stage. **F,** UMAP plot of

developmental indices within the embryonic skeletal muscle sub-trajectory. **G-H,** Application of

developmental indices to an ERMS tumor, SJRHB030680_R1 (G) and an ARMS tumor,

SJRHB031320_D1 (H). **I-J,** Developmental indices of 18 patient RMS tumors (I) or 18 O-PDXs

(J). Abbreviations: ERMS, embryonal rhabdomyosarcoma; ARMS, alveolar rhabdomyosarcoma;

UMAP, uniform manifold approximation and projection.

Figure 4: Developmental status in ERMS is plastic and associated with chromatin

accessibility at core regulatory superenhancer regions. A-B, Two competing models of tumor

heterogeneity within RMS. In the first model, RMS cells transition across developmental states

(A); in the alternate model, genetically distinct clones are restricted to muscle developmental

states (B). **C,** Schematic of the lentiviral barcode plasmid.[20,21] An 18-mer of random nucleotides

is incorporated into the 3'-untranslated region of a blue fluorescent protein (BFP) tag, enabling

barcode recovery from scRNA-seq libraries. **D,** Plot of frequency of individual barcodes for

subsequent passages of an individual ERMS O-PDX, SJRHB00026_X1. **E-F.** UMAP plot of an

ERMS O-PDX SJRHB013758_X2, colored based on developmental stage (E), or with 3 specific

barcodes highlighted (F). **G,** Quantitation of the developmental state diversity of all tumor cells

within SJRHB013758_X2, and from the 5 most prevalent barcoded clones. **H,** ChIP-seq and

chromHMM of *MYOD1* in an ERMS O-PDX, SJRHB10927_X1. Scales are indicated on the left,

and a previously identified CRC-SE[4] is highlighted in blue. **I,** Comparison of H3K27

trimethylation in various pediatric O-PDXs. OS, osteosarcoma; EWS, Ewing sarcoma; LPS,

liposarcoma; HGS, high-grade sarcoma; NB, neuroblastoma. **J,** Single-cell ATAC-seq of

SJRHB010927_X1 at the *MYOD1* locus; cell identities were defined via gene activity estimation,

and dataset integration with scRNA-seq data[30]. Abbreviations: ERMS, embryonal

rhabdomyosarcoma; ARMS, alveolar rhabdomyosarcoma; UMAP, uniform manifold

approximation and projection; RMS, rhabdomyosarcoma; OS, osteosarcoma; EWS, Ewing

sarcoma; LPS, liposarcoma; HGS, high grade sarcoma; NB, neuroblastoma.

Figure 5. Chemotherapy treatment of ERMS selects for mesoderm developmental stages.

A-B, Bar plots showing percentage of cells predicted to be dividing within each developmental

stage for patient tumors (a) and O-PDXs (b). **C,** Plots showing immunopositivity for MEOX2

(left) and MYOG (right) in patient samples from RMS13 obtained before treatment ("diagnosis")

and during therapy ("mid-treatment"). **D,** Treatment schema for VI therapy of mice bearing RMS

O-PDXs. Needle biopsies were performed at days 0, 3, 7, 14, and 21 or when tumors were large

enough to sample. **E,** Photograph of needle biopsy of an orthotopically-injected xenograft. **F-H,**

Photograph of tissue obtained by a biopsied O-PDX (F), which was fixed and stained using H&E

(G) or MYOG (H). **I,** Plot showing longitudinal expression of *MEOX2* by qRT-PCR during

treatment. There is an increase in *MEOX2* during chemotherapy (days 7,14,21) but the

proportion resets to basal levels after 28 days. This was verified by IHC (lower panel). **J,**

Boxplot of all biopsies for ERMS tumor bearing mice for the untreated and treated samples. The

plot is an integration of expression of 6 genes (*MEOX2, PAX3, EGFR, CD44, DCN, POSTN*)

expressed as normalized relative fold. The center line demarcates the median value with

rectangle showing interquartile range (IQR) between the first and third quartiles. The vertical

bars extending from the rectangles indicate maximum and minimum values with the exception of

outliers that more than 3 times the IQR. **K,** Relative proportion of nuclei in each developmental

state for longitudinal biopsies of a single O-PDX, determined using snRNA-seq of biopsied

tissue. **L,** Diagram of the mathematical model of ERMS developmental heterogeneity. **M-N,**

Simulated average population size for an untreated ERMS tumor (M) or a treated ERMS tumor

(N) briefly exposed to an antiproliferative agent (gray bar). Average population size over 524

simulations are shown, standard error bars are too small to see. **O-P,** Simulated time course of

barcode dynamics for an ERMS tumor that was either untreated (O) or briefly treated (P;

duration of treatment in grey bar). Each curve represents a different barcoded lineage. One

realization of the stochastic dynamics is shown. Insets under each graph show spatial

distributions of bar codes (color coded) in myoblast cells at an early and late stage of tumor

growth (O) and pre- and post-therapy (P). **Q,** Temporal development of the average entropy

index (measure of barcode diversity) during barcoded ERMS tumor growth, either untreated or

briefly treated (grey bar). Average entropy values over 524 simulations ± standard errors (dashed

lines) are shown. *Inset,* bar plot comparing the initial entropy index to the final entropy index of

untreated or treated tumors in experiments. Model parameters were: average value of L_{mes}

$=0.0035$ ($r_1=1.5$, $r_2=0.0001$), $L_{blast} =0.0045$, $P_{mes} =0.55$, $P_{blast} =0.49$, D=0.035, $\alpha_{mes} =0.0014$, α_{blast}

$=0.0035$. The parameter units are per minute. Abbreviations: ERMS, embryonal

rhabdomyosarcoma; ARMS, alveolar rhabdomyosarcoma; VCR, vincristine; IRN irinotecan.

Figure 6. Mesoderm-like ERMS cells are uniquely vulnerable to EGFR blockade. A,

Schematic workflow of NetBID algorithm to identify cell type–specific drivers from snRNA-seq

data. **B**, Volcano plot of differential activity analysis of signaling drivers in ERMSmesoderm vs.

other cell types. **C-D**, EGFR NetBID activity in different developmental states from snRNA-seq

data (C) and inferred from bulk RNA-seq of patient tumors (D). **E-F,** Dual IHC staining of

ERMS patient tumor, SJRHB030680_R1, combining EGFR (brown) with either MEOX2 (E) or

MYOG (F) in purple. **G,** Schedules of drugs used for preclinical study. **H,** Survival curves for

each treatment group for a ERMS tumor O-PDX (SJRHB01378_X1). **I,** Representative image of

bioluminescence for mice treated on the study, scale bar is photons/sec/cm^2 str. Scale bars: E,F,

10 μm. Abbreviations: VCR, vincristine; IRN, irinotecan; ERMS, embryonal

rhabdomyosarcoma; ARMS, alveolar rhabdomyosarcoma.

1 **Materials and Methods**

2 *See Supplementary Information.*

3
4 **Acknowledgements**
5

We thank A. McArthur for editing the manuscript. This work was supported by Cancer Center

Support (CA21765) and grants to M.A.D. from the National Institutes of Health (EY014867,

EY018599, and CA168875). The content is solely the responsibility of the authors and does not

necessarily represent the official views of the National Institutes of Health. Research was also

supported by American Lebanese Syrian Associated Charities. M.A.D. was also supported by the

Howard Hughes Medical Institute, Alex's Lemonade Stand, and Tully Family and Peterson

Foundations. A.G.P. was supported by Alex's Lemonade Stand, Hyundai Hope on Wheels, and

Damon Runyon Cancer Research Foundations.

Author Contributions
16
A.G.P, X.C. and M.A.D designed the study. A.G.P processed fresh and frozen tissue for single-

cell/nucleus RNA-seq and ATAC-seq. B.G., K.B., and E.S. assisted in patient sample and

xenograft tissue accrual for molecular analyses. A.G.P., X.C., X.H., and J.Y. performed

computational analysis of data. M.R.C., B.A.O., and H.T. supervised immunohistochemical

staining and pathology review. N.K. and D.W. performed mathematical modeling of tumors.

M.J.K. and A.P. wrote and supervised the RMS13 clinical trial. A.K. manages the Childhood

Solid Tumour Network. J.M. and K.B. assisted in molecular analysis. C.R. and X.Z. developed

the single-cell viewer portal. K.B. and E.S. performed the preclinical testing and tumor biopsies.

A.G.P. and J.L.N. developed the tumor organoid model.

Competing Interest Statement

The authors declare no competing financial interests.

Supplementary Information

Supplementary Information is available for this paper.

Corresponding Author

Correspondence and requests for materials should be addressed to Michael Dyer at

michael.dyer@stjude.org.

Data and Source Code Availability

NetBID: https://jyyulab.github.io/NetBID

All processed single-cell and single-nucleus RNA-sequencing data are publicly accessible via an

online data portal (https://pecan.stjude.cloud/static/RMS-scrna-atlas-2020).

All raw sequencing data will be uploaded and be publicly available at the time of publication

through GEO accession GSE174376.

References

1. Ries, L. *et al. Cancer Incidence and Survival among Children and Adolescents: United States SEER Program 1975-1995*. (National Cancer Institute, SEER Program).

2. Chen, X. *et al.* Targeting oxidative stress in embryonal rhabdomyosarcoma. *Cancer Cell* **24**, 710–724 (2013).

3. Kahn, H. J. *et al.* Immunohistochemical and electron microscopic assessment of childhood rhabdomyosarcoma. Increased frequency of diagnosis over routine histologic methods. *Cancer* **51**, 1897–1903 (1983).

4. Stewart, E. *et al.* Identification of Therapeutic Targets in Rhabdomyosarcoma through
Integrated Genomic, Epigenomic, and Proteomic Analyses. *Cancer Cell* **34**, 411–426 (2018).

5. Shern, J. F. *et al.* Comprehensive Genomic Analysis of Rhabdomyosarcoma Reveals a
Landscape of Alterations Affecting a Common Genetic Axis in Fusion-Positive and Fusion-
Negative Tumors. *Cancer Discov* **4**, 216–231 (2014).

6. Pappo, A. S. & Dirksen, U. Rhabdomyosarcoma, Ewing Sarcoma, and Other Round Cell
Sarcomas. *J Clin Oncol* **36**, 168–179 (2017).

7. Arndt, C. A. S., Bisogno, G. & Koscielniak, E. Fifty Years of Rhabdomyosarcoma studies on
both sides of the Pond and Lessons Learned. *Cancer Treat Rev* **68**, 94–101 (2018).

8. Pappo, A. S. *et al.* Survival After Relapse in Children and Adolescents With
Rhabdomyosarcoma: A Report From the Intergroup Rhabdomyosarcoma Study Group. *J Clin*
Oncol **17**, 3487–3493 (1999).

9. Stewart, E. *et al.* Orthotopic patient-derived xenografts of paediatric solid tumours. *Nature*
549, 96–100 (2017).

10. Chal, J. & Pourquié, O. Making muscle: skeletal myogenesis *in vivo* and *in vitro*.
Development **144**, 2104–2122 (2017).

11. Buckingham, M. & Rigby, P. W. J. Gene Regulatory Networks and Transcriptional
Mechanisms that Control Myogenesis. *Dev Cell* **28**, 225–238 (2014).

12. Sebire, N. J. & Malone, M. Myogenin and MyoD1 expression in paediatric
rhabdomyosarcomas. *J Clin Pathol* **56**, 412–416 (2003).

13. Tirosh, I. *et al.* Single-cell RNA-seq supports a developmental hierarchy in human
oligodendroglioma. *Nature* **539**, 309–313 (2016).

14. Slyper, M. *et al.* A single-cell and single-nucleus RNA-Seq toolbox for fresh and frozen
human tumors. *Nat Med* **26**, 792–802 (2020).

15. Manno, G. L. *et al.* RNA velocity of single cells. *Nature* **560**, 494–498 (2018).

16. Barkas, N. *et al.* Joint analysis of heterogeneous single-cell RNA-seq dataset collections. *Nat*
Methods **16**, 695–698 (2019).

17. Vincent, K. M. & Postovit, L.-M. A pan-cancer analysis of secreted Frizzled-related proteins:
re-examining their proposed tumour suppressive function. *Sci Rep* **7**, 42719 (2017).

18. Cao, J. *et al.* The single-cell transcriptional landscape of mammalian organogenesis. *Nature*
566, 496–502 (2019).

19. Cheng, C. *et al.* Latent cellular analysis robustly reveals subtle diversity in large-scale single-cell RNA-seq data. *Nucleic Acids Res* **16**, e143 (2019).

20. Adamson, B. *et al.* A Multiplexed Single-Cell CRISPR Screening Platform Enables Systematic Dissection of the Unfolded Protein Response. *Cell* **167**, 1867–1882 (2016).

21. Dixit, A. *et al.* Perturb-Seq: Dissecting Molecular Circuits with Scalable Single-Cell RNA Profiling of Pooled Genetic Screens. *Cell* **167**, 1853–1866 (2016).

22. Gryder, B. E. *et al.* PAX3-FOXO1 Establishes Myogenic Super Enhancers and Confers BET Bromodomain Vulnerability. *Cancer Discov* **7**, 884–899 (2017).

23. Du, X. *et al.* Hippo/Mst signalling couples metabolic state and immune function of CD8α+ dendritic cells. *Nature* **558**, 141–145 (2018).

24. Wijaya, J. *et al.* An ABC Transporter Drives Medulloblastoma Pathogenesis by Regulating Sonic Hedgehog Signaling. *Cancer Res* **80**, 1524–1537 (2020).

25. Khatamian, A., Paull, E. O., Califano, A. & Yu, J. SJARACNe: a scalable software tool for gene network reverse engineering from big data. *Bioinformatics* **35**, 2165–2166 (2018).

26. Ganti, R. *et al.* Expression and genomic status of EGFR and ErbB-2 in alveolar and embryonal rhabdomyosarcoma. *Modern Pathol* **19**, 1213–1220 (2006).

27. Grass, B. *et al.* Immunohistochemical detection of EGFR, fibrillin-2, P-cadherin and AP2beta as biomarkers for rhabdomyosarcoma diagnostics. *Histopathology* **54**, 873–879 (2009).

28. Wachtel, M. *et al.* Subtype and Prognostic Classification of Rhabdomyosarcoma by Immunohistochemistry. *J Clin Oncol* **24**, 816–822 (2006).

29. Mascaux, C. *et al.* EGFR Protein Expression in Non–Small Cell Lung Cancer Predicts Response to an EGFR Tyrosine Kinase Inhibitor—A Novel Antibody for Immunohistochemistry or AQUA Technology. *Clin Cancer Res* **17**, 7796–7807 (2011).

30. Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888–1902 (2019).

31. Stewart, E. *et al.* The Childhood Solid Tumor Network: A new resource for the developmental biology and oncology research communities. *Dev Biol* **411**, 287–293 (2016).

32. Smith, T., Heger, A. & Sudbery, I. UMI-tools: modeling sequencing errors in Unique Molecular Identifiers to improve quantification accuracy. *Genome Res* **27**, 491–499 (2017).

33. Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol* **36**, 411–420 (2018).

34. Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-seq data using regularized negative binomial regression. *Genome Biol* **20**, 296 (2019).

35. Becht, E. *et al.* Dimensionality reduction for visualizing single-cell data using UMAP. *Nat Biotechnol* **37**, 38–44 (2019).

36. Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression data analysis. *Genome Biol* **19**, 15 (2018).

37. Bergen, V., Lange, M., Peidli, S., Wolf, F. A. & Theis, F. J. Generalizing RNA velocity to transient cell states through dynamical modeling. *Nat Biotechnol* 1–7 (2020) doi:10.1038/s41587-020-0591-3.

38. Aran, D. *et al.* Reference-based analysis of lung single-cell sequencing reveals a transitional profibrotic macrophage. *Nat Immunol* **20**, 163–172 (2019).

39. Traag, V. A., Waltman, L. & Eck, N. J. van. From Louvain to Leiden: guaranteeing well-connected communities. *Sci Rep* **9**, 5233 (2019).

40. Saelens, W., Cannoodt, R., Todorov, H. & Saeys, Y. A comparison of single-cell trajectory inference methods. *Nat Biotechnol* **37**, 547–554 (2019).

41. Street, K. *et al.* Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics. *BMC Genomics* **19**, 477 (2018).

42. Hanawa, H. *et al.* Comparison of Various Envelope Proteins for Their Ability to Pseudotype Lentiviral Vectors and Transduce Primitive Hematopoietic Cells from Human Blood. *Mol Ther* **5**, 242–251 (2002).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A
ERMS mesoderm (n=19,317)
ERMS myoblast (n=43,667)
ARMS myoblast (n=29,040)
ERMS myocyte (n=10,674)
ARMS myocyte (n=7,757)

B
-log₁₀ (p-value) vs differential activity

EGFR
HAS2
GFPT2
ADAMTS12
PTGFR
USP14
SETDB1
ROR2
RAB1A
PLCB4
PIK3C2A
PDGFRB
PDE8A
OXSR1
NRP1
MTR
MMP2
MAT2A
GNGT1
ENPP2
EIF2B5
DDX5
CHD1L
BACE1
DUSP8
CHSY3
B4GALT5

C
EGFR pathway activity
p=4.4x10⁻¹³⁵
p=1.8x10⁻¹⁷⁴
mesoderm | myoblast | myocyte | myoblast | myocyte
ERMS | ARMS

D
EGFR network activity
p=5.4x10⁻³⁶
ERMS (n=52) | ARMS (n=25)

E MEOX2 EGFR

F MYOG EGFR

G
VCR+IRN
afatinib
gefitinib
VCR+IRN+afatinibᵁ
VCR+IRN+gefitinibᵁ
VCR+IRN+afatinibᴹ
VCR+IRN+gefitinibᴹ

week 1 | week 2 | week 3 | week 4 | week 5 | week 6

★ VCR
🥚 IRN
▬ afatinib
▬ gefitinib

H
SJRHB013758_X1 non-moribund (proportion) vs time (days)

placebo
afatinib
gefitinib
VCR+IRN
VCR+IRN+afatinibᵁ
VCR+IRN+gefitinibᵁ
VCR+IRN+afatinibᴹ
VCR+IRN+gefitinibᴹ

I
placebo | afatinib | gefitinib | VCR+IRN
VCR+IRN+afatᵁ | VCR+IRN+gefᵁ | VCR+IRN+afatᴹ | VCR+IRN+gefᴹ