The anatomical logic of smell

Mayra L. Ruiz Tejada Segura[1,2,3,#]; Eman Abou Moussa[4,#]; Elisa Garabello[5,6]; Thiago S. Nakahara[7]; Melanie Makhlouf[4]; Lisa S. Mathew[4]; Filippo Valle[5]; Susie S.Y. Huang[4]; Joel D. Mainland[8,9]; Michele Caselle[5]; Matteo Osella[5]; Stephan Lorenz[4,10]; Johannes Reisert[8]; Darren W. Logan[10]; Bettina Malnic[7]; Antonio Scialdone[1,2,3,†,*]; Luis R. Saraiva[4,8,11, †,*]

[1] Institute of Epigenetics and Stem Cells, Helmholtz Zentrum München; Feodor-Lynen-Strasse 21, 81377 München, Germany

[2] Institute of Functional Epigenetics, Helmholtz Zentrum München; Ingolstädter Landstraße 1, 85764 Neuherberg, Germany

[3] Institute of Computational Biology, Helmholtz Zentrum München; Ingolstädter Landstraße 1, 85764 Neuherberg, Germany

[4] Sidra Medicine; PO Box 26999; Doha, Qatar.

[5] Physics Department, University of Turin and INFN; via P. Giuria 1, 10125 Turin, Italy

[6] Department of Civil and Environmental Engineering, Cornell University; Ithaca, NY 14853, USA

[7] Department of Biochemistry, University of São Paulo; São Paulo, Brazil.

[8] Monell Chemical Senses Center; 3500 Market Street, Philadelphia, PA 19104, USA.

[9] Department of Neuroscience, University of Pennsylvania; Philadelphia, PA 19104, USA.

[10] Wellcome Sanger Institute; Wellcome Genome Campus, Hinxton, Cambridge, CB10 1SD, UK.

[11] College of Health and Life Sciences, Hamad Bin Khalifa University; Doha, Qatar.

[#] Equal contribution (co-first)
† These authors jointly supervised this work
*Correspondence and requests for materials should be addressed to A.S. (email: antonio.scialdone@helmholtz-muenchen.de) or L.R.S. (email: saraivalmr@gmail.com; twitter: @saraivalab)

Abstract

The sense of smell helps us navigate the environment, but its anatomical logic remains unknown. The spatial location of odorant receptor genes (*Olfrs*) in the nose is widely thought to be independent of the structural diversity of the odorants they detect. Using spatial transcriptomics, we created a genome-wide 3D atlas of the mouse olfactory mucosa (OM), and identified key genes differentially expressed in space. Expression maps reveal that *Olfrs* are distributed in a continuous and overlapping fashion over five broad zones in the OM. The spatial locations of *Olfrs* correlate with the mucus solubility of the odorants they recognize. Thus, we provide direct evidence for the chromatographic theory of olfaction, and elucidate the basic logic for the peripheral representation of smell.

One-Sentence Summary

Chemical solubility is organized topographically in the mouse nose

Keywords

Spatial transcriptomics; 3D; olfaction; olfactory epithelium; RNA-seq; odorant

Main text

The functional logic underlying the topographic organization of primary receptor neurons and their receptive fields is well-known for all sensory systems, except olfaction (*1*). The mammalian nose is constantly flooded with odorant cocktails. Powered by a sniff, air enters the nasal cavity, until it reaches the olfactory mucosa (OM). There, myriad odorants activate odorant receptors (*Olfrs*) present in the cilia of olfactory sensory neurons (OSNs), triggering a complex cascade of events that culminate in the brain and result in odor perception (*1, 2*). In mice, most mature OSNs express a single allele of one out of ~1100 Olfr genes (*Olfrs*) (*3-6*). Olfrs employ a combinatorial strategy to detect odorants, which maximizes their detection capacity (*4, 7*). OSNs expressing the same *Olfr* share similar odorant response profiles (*4, 7*), and drive their axons to the same glomeruli in the olfactory bulb (*8-10*). Thus, *Olfrs* define functional units in the olfactory system, and function as genetic markers to discriminate between different mature OSN subtypes (*6, 11*).
Another remarkable feature of the OSN subtypes is their spatial distribution in the OM. Early studies postulated that OSNs expressing different *Olfrs* are spatially segregated into four broad areas within the OM called 'zones' (*12, 13*), but recent work has suggested up to nine overlapping zones (*14-16*). The functional relevance of these zones is not fully understood. These studies, combined, have sampled only ~10% of the intact mouse *Olfr* repertoire. We do not currently understand the full complexity of the OM and, most importantly, lack an unbiased and quantitative definition of zones. In effect, the exact number of zones, their anatomical boundaries, their molecular identity and their potential functional relevance are yet to be determined.

One hypothesis is that the topographic distribution of *Olfrs*/OSN subtypes evolved because it plays a key role in the process of *Olfr* choice in mature OSNs and/or in OSN axon guidance (*17, 18*). An alternative hypothesis is that the spatial organization of *Olfrs*/OSN subtypes is tuned to maximize the detection and discrimination of odorants in the peripheral olfactory system (*12*). Interestingly, the receptive fields of mouse OSNs vary with their spatial location (*19*), which in some cases correlate with the patterns of odorant sorption in the mouse OM – this association was proposed as the '*chromatographic hypothesis*' decades before the discovery of the *Olfrs* (*20*), and later rebranded as the '*sorption hypothesis*' in olfaction (*21, 22*). While some studies lend support to these hypotheses (reviewed in (*23*)), others question their validity (*24, 25*). Thus, the logic underlying the peripheral representation of smell in the peripheral olfactory system still remains unknown, and it is subject of great controversy (*23, 26*).

Spatial transcriptomics, which combines spatial information with high-throughput gene expression profiling, has started to expand our knowledge of complex tissues, organs, or even entire organisms (*27-31*). In this study, we employed a spatial transcriptomics approach to create an 3D map of gene expression of the mouse nose,

90 and combined it with machine learning and chemoinformatics to reveal the
91 anatomical logic of smell in mouse.
92
A high-resolution spatial transcriptomic map of the mouse olfactory mucosa
We adapted the RNA-seq tomography (Tomo-seq) method (*30*) to create a spatially
resolved genome-wide transcriptional atlas of the mouse nose. We obtained
cryosections (35 µm) collected along the dorsal-ventral (DV), anterior-posterior (AP),
and lateral-medial-lateral (LML) axes (n=3 per axis) of the OM (Fig. 1A), and
performed RNA-seq on individual cryosections (gMethods). After quality control (see
Methods, fig. S1A-D and table S1), we computationally refined the alignment of the
cryosection along each axis, and we observed a high correlation between biological
replicates (see Methods and fig. 1B). Hence, we combined the three replicates into a
single series of spatial data including 54, 60 and 56 positions along the DV, AP and
LML axis, respectively (see Methods and Fig. 1C). On average, we detected >18,000
genes per axis, representing a total of 19,249 unique genes for all axes combined (Fig.
1D). Molecular markers for all canonical cell types known to populate the mouse OM
were detected among all axes (Fig. 1E), and were expressed at the expected levels (*6*).
Next, we verified the presence of a spatial signal with the Moran's I (*32*) (fig. S1E).
Given the left/right symmetry along the LML axis (see Methods; Fig. 1C), the data
was centered and averaged on the two sides (see Methods) – henceforth the LML axis
will be presented and referred to as the lateral-medial (LM) axis. We could reproduce
the expression patterns in the OM for known spatial markers, including the
dorsomedial markers *Acsm4* and *Nqo1* (*33, 34*), and the complementary ventrolateral
markers *Ncam2* and *Reg3g* (*35, 36*) (Fig. 1F, fig. S1F).
114
Spatial differential gene expression analysis identifies cell type-specific
expression patterns and functional hotspots in the OM
Having established the accuracy of our dataset, we investigated cell-type specific
patterns. Previous studies showed that the vast majority of genes displaying spatially
segregated expression patterns across the OM are expressed in mature OSNs (*6, 13,*
14, 16, 33, 34, 36-48), but a small fraction are expressed in sustentacular, globose
basal, olfactory ensheathing, Bowman's gland, and respiratory epithelial cells (*36, 42,*
47-53). Despite this progress, our knowledge on what genes display zonal expression
patterns, and what cell types they are primarily expressed in is still very limited.
124
125 To identify axis-specific differentially expressed genes (hereafter referred to as spatial
126 DEGs), we applied the Ljung-Box test to the autocorrelation function calculated on
127 the binarized expression values (fig. S2A, and Methods), and obtained a total of
128 12,303 DEGs for the 3 axes combined (FDR<0.01; Fig. 2A).
129

To add cell-type resolution to the spatial axes, we combined our data with a single-cell
RNA-seq (scRNA-seq) dataset from 13 cell types present in the mouse OM (*54*). We
catalogued spatial DEGs based on their expression in mature OSNs (mOSNs) versus
the 12 other cell types (non-mOSNs) (Fig. 2B,C; table S2). This led to the
identification of 456 spatial DEGs expressed exclusively in non-mOSNs, which are
associated with gene ontology (GO) terms such transcription factors, norepinephrine
metabolism, toxin metabolism, bone development, regulation of cell migration, T-cell
activation, and others (table S2). Some of these genes are expressed across many cell
types, but others are specific to a single cell type (Fig. 2C, table S2). For instance,
there are cell-specific markers with known spatial expression patterns, such as the
sustentacular cells and Bowman's glands markers *Cyp2g1* and *Gstm2* (*36*), the neural
progenitor cell markers *Eya2* and *Hes6* (*45*), and the basal lamina and olfactory
ensheathing cell markers *Aldh1a7* and *Aldh3a1* (*47*) (table S2). Additionally, we found
previously unknown genes with zonal expression patterns along a single axis or
multiple axes, such as the ribosomal protein *Rps21* that is primarily expressed in
horizontal basal cells (HBCs), the extracellular proteinase inhibitor *Wfdc18* in
microvillous cells type 1 (MVC1s), the fibroblast growth factor *Fgf20* in immature
sustentacular cells (iSCs), and the adapter protein *Dab2* in mature sustentacular cells
(mSCs) (fig. S2 B,C). These results suggest differences either in their expression levels
and/or in the relative abundance of cell types across the OM.
150
The successful identification of thousands of spatial DEGs, prompted us to examine
their distribution patterns along each axis, and the putative functions associated with
such spatial clusters of gene expression. We started by using uniform manifold
approximation and projection (UMAP) (*55*) and hierarchical clustering to visualize
and cluster all spatial DEGs along the three cartesian axes. This analysis uncovered
nine patterns of expression in the DV and AP axes each, and five patterns in the LM
axis (Fig. 2D-E). These patterns include variations of four major shapes:
monotonically increasing (/), monotonically decreasing (\), U-shape (∪), and inverted
U-shape (∩) (Fig. 2E). The latter two patterns present clear maximum/minimum at
different positions along the axis – for example, the brown, green, pink, magenta, and
black AP clusters show a similar inverted ∪ pattern, but their maximum moves along
the axis (Fig. 2E). As expected, several known genes with zonal expression patterns
are expressed in the cluster mimicking their respective expression pattern in the tissue.
For example, the dorsomedial markers *Acsm4* and *Nqo1* belong to the turquoise
clusters in both the DV and LM axes, while the ventrolateral marker *Reg3g* belongs to
the blue cluster from the DV axis (Fig. 1F, S1F; table S3).
GO enrichment analysis on the axis-specific DEGs yielded enriched terms for 14 of
the 23 spatial clusters (table S3). For example, the turquoise AP cluster displaying a
monotonically increasing pattern (Fig. 2E) yielded GO terms broadly associated with

the molecular machinery of mOSNS – such as axonal transportation, RNA
processing, protein modification and quality control, ribosomal regulation and
regulation of histone deacetylation (fig. S2D and table S3). Interestingly, the brown
DV cluster, which displays a monotonically decreasing expression pattern (Fig. 2E)
had similar GO term enrichment (fig. S2E and table S3). These results raise the
hypothesis that mOSN activity is enriched in the dorsoposterior region of the OM,
consistent with previous results (*56, 57*).
We then extended our analysis to the remaining clusters, and found additional GO
terms shared between several clusters among the three different axes. For example,
the GO terms enriched in the dorsomedial region (turquoise DV, pink AP and LM
green) suggest that this region is involved in the OM detoxification (Fig. S2F-H).
Moreover, the enrichment in terms related to immune system in the anteromedial
section along the AP axis (yellow, black and magenta AP clusters, and turquoise LM
cluster; table S3) hints at a role of this area in defending OM from pathogenic
invaders. Finally, the ventral portion of the DV (red DV cluster) was associated with
terms related to cilia movement and function, consistent with both the location and
functions of the respiratory epithelium (*36*) (table S3).

Next, we further explored the relationships between the genes populating each
cluster. We found that ventral genes (blue DV cluster) tend to reach a peak in
expression in the anterior area of the OM (yellow AP cluster) more often than
expected by chance (Fig. 2F). We also observed that medial genes (turquoise LM
cluster) are more highly expressed in the dorsal (magenta DV cluster) and anterior
regions (black, yellow and magenta AP cluster), while genes peaking in the lateral
region (brown LM cluster) tend to be ventral (red DV cluster; Figure 2F). These
conclusions hold even when we exclude olfactory receptor genes from the analysis
(fig. S2I-K).

These associations between the clusters of DEGs along different axes strongly
suggest that the presence of complex 3D expression patterns in OM is not restricted
to either *Olfrs* or mOSNs. Moreover, our data allowed us to uncover spatially
restricted functional hotspots within the OM.

A 3D transcriptomic atlas of the mouse OM

Our Tomo-seq data enables the quantification of gene expression estimates along the
three cartesian axes of the OM. Here, we take this analysis one step further and
generate a fully browsable tridimensional (3D) gene expression atlas of the mouse
OM. To this aim, first we reconstructed the 3D shape of OM based on publicly
available images of OM sections (made available by (*58*), see Methods). We then fed
the resulting shape information combined with the gene expression data along the

three cartesian axes into the iterative proportional fitting (IPF) algorithm (*30, 59*) (Fig.
3A). Our resulting 3D atlas of the OM faithfully reproduced the known 3D pattern of
the dorsomedial marker *Acsm4* (*34*) (Fig. 3B). In addition, we confirmed the predicted
3D expression patterns for two novel spatial DEGs we identified: *Cytl1* and *Moxd2*
(table S3) using in situ hybridization (Fig. 3C-F). *Cytl1* is expressed mainly along the
septum (Fig. 3C-D), consistent with the role *Cytl1* plays in osteogenesis,
chondrogenesis, and bone/cartilage homeostasis (*60, 61*). *Moxd2*, a monooxygenase
dopamine hydroxylase-like protein putatively involved in neurotransmitter
metabolism (*62*), is expressed mostly in a small ventrolateral patch (Fig. 3E-F).
We have made this new 3D gene expression atlas of the mouse OM available to the
scientific community via a web-portal (at the address: http://atlas3dnose.helmholtz-
muenchen.de:3838/atlas3Dnose), providing access to all the spatial transcriptomic
data in a browsable, user-friendly format.

Topographical expression patterns of *Olfrs*

In our combined dataset we detected a total of 959 unique *Olfrs* (Fig. 4A), of which
we confidently reconstructed the spatial expression patterns for 689 differentially
expressed in space (FDR<0.01; Fig. 4B) – a number six times larger than the
combined 112 *Olfrs* characterized by previous ISH studies (*10, 12, 14, 16*). To define
Olfr expression in 3D space in a rigorous, unbiased and quantitative way, we ran a
Latent Dirichlet Allocation algorithm (see Methods)(*63*) on the 689 spatially
differentially expressed *Olfr*, which suggested the presence of five zones (see Methods
and fig. S3A). Next, we visualized the spatial distribution of these five zones in our
3D model of the OM, with colors representing the probability that a given spatial
position belongs to each zone (Fig. 4C). These zones extend from the dorsomedial-
posterior to the lateroventral-anterior region, consistent with the previously described
zones (*12-14*).

Using the same mathematical framework, we can compute the probabilities that the
expression pattern of each *Olfr* gene belongs to a given zone, i.e., the "degree of
belonging" (DOB, table S4). The DOBs represent a decomposition of the expression
patterns in terms of the five zones shown in Fig. 4C and quantitatively describe the
changes in patterns of genes with overlapping areas of expression (e.g., see fig. S3B).
The width of the distribution of DOBs across the five zones, which can be measured
with entropy, can distinguish genes whose patterns mostly fit in a single zone from
those spanning multiple zones (see Methods and fig. S3C). These analyses can be
extended also to non-*Olfr* genes, allowing the identification of marker genes for each
zone (e.g., *Moxd2* for zone 2; see fig. 3E-F, S3D and Methods). To visualize the global
distribution of the 689 *Olfrs*, we applied the diffusion map algorithm (*64*) to their
DOBs. This showed that the genes are approximately distributed along a continuous

line spanning the five zones and without clear borders between zones (fig. 4D) (*14-*
16). With the diffusion pseudo-time algorithm (*65*), we calculated an index (hereafter
referred to as "3D index") that tracks the position of each *Olfr* gene along the 1D
curve in the diffusion map and represents its expression pattern (fig. 4E).
While this approach yielded an index for the 689 *Olfr* genes used to build the diffusion
map, there were ~700 *Olfrs* that could not be analyzed, either because they were too
lowly expressed or not detected at all in our dataset (Fig. 4A). However, since the
spatial expression patterns for some *Olfrs* are partly associated with their
chromosomal/genomic coordinates (*66-68*), we hypothesized that we could train a
machine learning algorithm to predict the 3D indexes for the ~700 *Olfrs* missing from
our dataset. Thus, we trained a Random Forest algorithm on the 3D indexes of 681
spatially differentially expressed *Olfrs* in our dataset using 9 genomic features as
predictors (e.g., chromosomal position, number of *Olfrs* in cluster, distance to nearest
known enhancer, etc; see Methods and Fig. 4F). The five most important predictors
were features associated with chromosomal position, distance to the closest *Olfr*
enhancer (*69*), length of the *Olfr* cluster, position in the *Olfr* cluster, and phylogenetic
class of *Olfrs* (Fig. 4F).
265
We assessed the performance of the algorithm with a cross-validation test (see
Methods and Fig. S4A), and then predicted the 3D indexes for the 697 *Olfrs* missing
reliable expression estimates in our dataset. Importantly, we found that our 3D
indexes strongly correlate with the zonal indexes inferred using ISH in (*14*)(rho=0.88,
p=3×10^{-27}; Fig. 4G) and in two other studies (*16, 66*)(fig. S4B,C). Furthermore, we
performed ISH for two *Olfrs* that have not been characterized before, *Olfr309* and
Olfr727, which were correctly predicted to be expressed primarily in zone 2 and 4
respectively (Fig. 4H-M and table S4)
274
Taken together, through multiple unsupervised and supervised computational
methods, we have quantitatively defined five spatial expression domains in the OM
(called zones), and have provided accurate 3D spatial indexes for 1386 *Olfrs*, which
represents ~98% of the annotated *Olfrs*.

The anatomical logic of smell
Next, we used our 3D, quantitative reconstruction of the OM to address whether
there is an underlying logic linked to *Olfrs* zonal distribution. We first compiled a list
of known 738 Olfr-ligand pairs, representing 153 *Olfrs* and 221 odorants (Fig. 5A,
table S5). Interestingly, we found that *Olfr* pairs sharing at least one common ligand
have more similar expression patterns (i.e., more similar 3D indexes) than *Olfrs*
detecting different sets of odorants (Fig. 5B). This observation is consistent with the

hypothesis that the *Olfr* zonal expression depends, at least partially, on the properties
of the odorants they bind to.
Previous studies showed that the sorption patterns of the odorants in the mouse OM
depend on their mucus solubility, which can be quantified by the air/mucus partition
coefficient (*21, 70*). In particular, odorants with higher mucus solubility reach a higher
concentration in the antero-dorsomedial region of the OM (*12, 21*). Hence, we
hypothesized that the Olfr zonal expression patterns correlate with the mucus
solubility of the cognate odorants, in such a way to mimic the sorption pattern of the
odorants.
To test this hypothesis, we compared the air/mucus partition coefficients (K_{am}) of
221 odorants to the average 3D indexes of the *Olfrs* detecting them (see Methods,
table S5). Strikingly, we found a positive correlation (rho=0.55, p-value=$7x10^{-10}$; Fig.
5C), indicating that the most soluble odorants (lower K_{am}) preferentially activate *Olfrs*
in the most antero-dorsomedial OM region (zone 1) of the OM, while the least
soluble odorants (higher K_{am}) activate *Olfrs* in the postero-ventrolateral OM region
(zones 4-5). In other words, gradients of odorants sorption (as defined by their K_{am})
correlate with the gradients of *Olfr* expression from zone 1 to zone 5, consistent with
the *chromatographic/sorption* hypothesis in olfaction (*20, 21*). This is exemplified by the
plots in Fig. 5D, illustrating the predicted average expression levels across OM
sections of the *Olfrs* binding to five odorants with different values of K_{am}. These
results show, for the first time, a direct association between *Olfr* spatial patterns and
the sorption patterns of their cognate ligands in the OM, providing a potential
explanation for the physiological function of the zones in the OM.

Discussion

Past studies yielded inconclusive and sometimes contradictory views on the basic logic
underlying the peripheral representation of smell, partly because the topographic
distribution of OSN subtypes and their receptive fields still remained vastly uncharted,
data on Olfr-ligand pairs was scarce, and the many pitfalls associated with electro-
olfactogram recordings used to study spatial patterns of odor recognition in the nose
(*26, 71, 72*).
Here, we combined RNA-seq and computational approaches that utilize unsupervised
and supervised machine learning methods to discover and quantitatively characterize
spatial expression patterns in the OM. We created the first 3D transcriptional map of
the mouse OM, which allowed us to spatially characterize thousands of genes,
including ~98% of the annotated *Olfrs*. We identified and validated by ISH several
new spatial marker genes, and a clustering analysis pinpointed the OM locations
where specific functions related to, e.g., the immune response, might be carried out.
We also identified five broad Olfr expression zones in the OM, which were
mathematically defined and used to decompose the expression patterns of all genes.

Our analysis enabled us to answer fundamental and longstanding questions about the
rationale behind the spatial organization of the peripheral olfactory system.
Specifically, we provide evidence to the hypothesis that the spatial zones increase the
discriminatory power of the olfactory system by distributing receptors in the areas of
the OM that are more likely to be reached by their cognate ligands, based on their
solubility in mucus.
The quantitative framework we built for this dataset will facilitate interrogation of
gene expression patterns via an online tool we provide, and help answer important
questions on the spatial patterns in the nose. Moreover, our approach could be easily
applied to spatial transcriptomic data collected from other tissues, or the same tissue
across multiple developmental stages. Results from this study will also serve as a
template to start answering other important questions about olfaction, such as
whether Olfr spatial expression maps can also encode maps of odor perception.
Because the general molecular mechanisms of olfaction, zonal organization of Olfrs,
and components of olfactory perception are conserved in mammals (*73-77*), findings
from our and other subsequent studies can likely be extrapolated to other mammals,
including humans. Finally, the functional logic underlying the topographic
organization of primary receptor neurons and their receptive fields in smell is now
starting to be exposed.
344

345   **Figures**

FIGURE 1

Figure 1. Application of TOMO-seq to mouse OM. (A) Experimental design.
TOMO-seq was performed on 9 tissue samples, from which 3 were sliced along the
Dorsal-Ventral axis, 3 along the Posterior-Anterior axis and 3 along the Lateral-
Medial-Lateral axis. (B) Boxplots showing the distributions of Spearman's correlation
coefficients (Rho) between replicates in each axis. (C) Heatmaps showing Spearman's
correlation between gene expression patterns at different positions along the three
axes. (D) Number of detected genes along each axis separately or across the whole
dataset. Genes were considered as detected when they had at least one normalized
count in at least 10% of the samples from one axis. (E) Heatmaps of $\log_{10}$ normalized
expression (after combining the three replicates per axis) of OM canonical markers
along the 3 axes (RPM = reads per million; mOSNs = mature Olfactory Sensory
Neurons; iOSNs = immature Olfactory Sensory Neurons; GBCs = Globose Basal
Cells; HBCs = Horizontal Basal Cells; SUSs = Sustentacular cells; RES = Respiratory
epithelium cells). (F) Normalized expression of canonical OM spatial marker genes
along the 3 axes. Red line showing fits with local polynomial models.

FIGURE 2

Figure 2. Genes with non-random spatial patterns across different cell types in the OM. (A) Venn diagram showing the numbers of spatial differentially expressed genes (DEGs) along each axis. (B) Bar plot showing the $\log_{10}$ number of spatial DEGs that are mOSN-specific ("mOSNs"), or that are detected only in cell types other than mOSNs ("other"). (C) Heatmap of log10 mean expression per cell type of genes that are not expressed in mOSNs, but only in other OM cell types (INP = Immediate Neuronal Precursors; GBC = Globose Basal Cells; mOSNs = mature Olfactory sensory neurons; iOSNs = immature Olfactory Sensory Neurons; MVC = Microvillous Cells; iSC = Immature Sustentacular Cells; mSC = Mature Sustentacular Cells; HBCs = Horizontal Basal Cells). (D) UMAP plots of spatial DEGs along the three axes. Each gene is colored according to the cluster it belongs to. (E) Normalized average expression patterns of spatial DEGs clusters along the three axes. (F) Heatmap showing the log2 enrichment over the random case for the intersection between lists of genes belonging to different clusters (indicated by colored circles) across pairs of axes.

FIGURE 3

Figure 3. The 3D reconstruction of the OM. (A) Schematic of 3D shape reconstruction strategy. Images of 2D slices along the AP axis of the OM were piled together to build an in silico 3D model of OM, which can also be used to visualize in silico sections. This 3D model, together with the gene expression data along each axis, was the input of the iterative proportional fitting algorithm, which allowed us to estimate a 3D expression pattern for any gene. (B) Reconstruction of the 3D expression pattern of the gene *Acsm4* in the OM, visualized in 3D and in OM slices taken along the AP axis. (C) Reconstruction of the 3D expression pattern of the gene *Cytl1* in the OM. (D) In-situ hybridization experiment validating *Cytl1* spatial expression pattern reconstructed in panel D. (E) Reconstruction of the 3D expression pattern of the gene *Moxd2* in the OM. (F) In-situ hybridization experiment validating *Moxd2* spatial expression pattern reconstructed in panel E.

FIGURE 4

Figure 4. Zonal organization of the OM. (A) Number of *Olfr* genes detected in our data and in an OM bulk RNA-seq data (*6*) (B) Venn diagram of spatially differentially expressed *Olfrs* per axis. (C) Visualization of the five zones estimated with a Latent Dirichlet Allocation algorithm. The colors indicate the probability (scaled by its maximum value) that a position belongs to a given zone. (D) Diffusion map of *Olfrs*. Genes are colored based on the zone they fit in the most. DC, diffusion component. (E) Same as panel D, with *Olfrs* colored by their 3D index. (F) We fitted a Random Forest algorithm to the 3D indexes of 681 spatially differentially expressed *Olfr* using nine genomic features as predictors. After training, the Random Forest was used to predict the 3D indexes of 697 *Olfrs* that have too low levels in our data. (G) 3D indexes versus the indexes of 80 Olfrs estimated in (*14*) from ISH data. Black circles indicate *Olfrs* detected in our dataset; green circles are *Olfrs* whose indexes were predicted with Random Forest. (H, K) Predicted expression patterns of *Olfr309* and *Olfr727*. (I,L) Degrees of belonging for *Olfr309* and *Olfr727*. (J,M) In-situ hybridization for *Olfr309* and *Olfr727*.

FIGURE 5

414
415

Figure 5. Physiological role of the zones. (A) Circular network illustrating the pairs
of Olfrs and ligands that we found in literature. (B) Box plots showing the
distributions of the absolute value of 3D index differences between pairs of Olfrs
sharing at least one ligand versus pairs of Olfrs without cognate ligands in common.
The difference between the two distributions is statistically significant ($p < 2.2 \times 10^{-16}$,
Wilcoxon Rank-Sum test). (C) Scatter plot illustrating the correlation between
odorants' air / mucus partition coefficients and the average 3D indexes of their
cognate Olfrs. Only odorants for which we know at least two cognate Olfrs (110) are
used here. Odorants are colored according to the zone they belong to (defined as the
zone with the highest average degree of belonging computed over all cognate
receptors). Five odorants, indicated on the right-hand side, are highlighted in the plot
by slightly larger circles. (D) Average expression patterns of the cognate Olfr for the
five odorants highlighted in panel C.

METHODS

Ethics statement

The use and care of animals used in this study was approved by the Internal Animal Care and Use Committee (IACUC) of Monell Chemical Senses Center, by the IACUC of the University of São Paulo, and by the Wellcome Trust Sanger Institute Animal Welfare and Ethics Review Board in accordance with UK Home Office regulations, the UK Animals (Scientific Procedures) Act of 1986.

Dissection of the olfactory mucosa, cryosections, and RNA-sequencing

All mice used were C57BL/6J (The Jackson Laboratory, Stock # 00664), 8-9 weeks old and group-housed. The olfactory mucosa (OM) of 9 mice was carefully dissected, and all the surrounding tissue (including glands and bone) removed. The OMs were then embedded in OCT (Tissue Tek), immediately frozen in dry-ice and kept at -80°C. Each OM was then cryosectioned along each of the 3 cartesian axes: dorsal-ventral (DV, N=3), anterior-posterior (AP, N=3), or lateral-medial-lateral (LML, N=3). Every second cryosections (35 μm thick) was collected into 1.5 mL eppendorf tubes containing 350 μl RLT Plus Buffer (Qiagen) supplemented with 1% 2-mercaptoethanol, immediately frozen in dry-ice and kept at -80C until extraction. RNA was extracted using the RNeasy Plus Micro Kit (Qiagen), together with a genomic DNA eliminator column and a 30-minute incubation with DNAse I (Qiagen). Reverse transcription and cDNA pre-amplification were performed using the SMART-Seq v4 Ultra Low Input RNA Kit for Sequencing (Clontech/Takara). cDNA was harvested and quantified with the Bioanalyzer DNA High-Sensitivity kit (Agilent Technologies). Libraries were prepared using the Nextera XT DNA Sample Preparation Kit and the Nextera Index Kit (Illumina). Multiplexed libraries were pooled and paired-end 150-bp sequencing was performed on the Illumina HiSeq 4000 platform at Sidra Medicine, except for one library (DV-I) for which 125-bp paired-end sequencing was performed on the Illumina HiSeq 2500 platform at the Wellcome Sanger Institute. For the remaining eight libraries, 150-bp sequencing was performed on the Illumina HiSeq 4000 platform. The raw data are available through ArrayExpress under accession number E-MTAB-10211.

RNA-seq data mapping and gene counting

Reads were aligned to the mm10 mouse genome (release 99). The sequences of the genes "*Xntrpc*" and "*Capn5*" were removed from the genome files as in (*6*). The alignment was performed with the software STAR version 2.7.3a (*78*). Genome indexes were generated using STAR --runMode genomeGenerate with default parameters. Then, alignment of reads was performed with the following options: --runThreadN 48 --outSAMunmapped Within --outFilterMultimapNmax 1000 --outFilterMismatchNmax 4 --outFilterMatchNmin 100 --alignIntronMax 50000 --

alignMatesGapMax 50500 --outSAMstrandField intronMotif --outFilterType BySJout.
The resulting SAM files were converted to bam format and sorted using samtools
(version 0.1.19-44428cd)(*79*). The multi mapping reads were eliminated using the
same software (samtools view -q 255). Finally, the reads for each gene were counted
using htseq-count (version 0.11.2) with the options -m intersection-nonempty -s no -i
gene_name -r pos (*80*).
475

Quality Control
476

We excluded all the samples that fulfilled any of these criteria: they had less than 50%
mapped reads, less than 4,000 detected genes, more than 20% mitochondrial reads,
less than 10,000 total number of reads, or did not express any of the 3 canonical OSN
markers *Omp*, *Cnga* and *Gnal*. This resulted in ~51 good-quality sections along the DV
axis (~84% out of the collected sections), ~76 (~91% of total) along the AP axis and
~59 (~93% of total) along the LML axis, as averaged across the three replicates per
axis.
484

Data normalization
485

Gene expression counts were normalized by reads-per-million (RPM), then genes
detected in only one replicate and genes that were detected in less than 10% of all
samples along one axis were eliminated. To check the similarity between replicates, we
calculated Spearman correlations between the transcriptional profiles of sections along
each axis (using the top 1000 Highly Variable Genes per axis). Close positions had the
most similar transcriptional profiles (Fig. 1C). Afterwards, the 3 replicates for each
axis were aligned as follows: the top 3,000 highly variable genes (HVGs) from each
replicate were identified using the method implemented in the scran library in R (*81*)
and the intersection of these 3 groups was used in the next steps. For the replicates'
alignment, we took as reference the replicate with the smallest number of slices. We
used a sliding window approach that identified the range of consecutive positions on
each replicate along which the average value of the Spearman's correlation coefficient
computed with the reference replicate over the HVG was maximum (mean
Spearman's Rho= 0.80, p<0.05). To mitigate batch effects, the level of every gene was
scaled in such a way that their average value in each replicate was equal to the average
calculated across all replicates. After this scaling transformation, the data was then
averaged between replicates. Once the 3 biological replicates were combined, we had
54 sections along the DV axis, 60 along the AP and 56 along the LML. Along the
LML axis a symmetric pattern of expression is expected around the central position,
where the septal bone is located. To confirm this in our data, first we identified the
central position by analyzing the expression pattern of neuronal markers like *Cnga2*,
Omp and *Gnal*, whose expression is lowest in the area around the septal bone. Indeed,
all three marker genes reach a minimum at the same position along the LML axis
(slice 28), which we considered to be the center. The expression patterns of ~90% of

genes on either side of the central position show a positive correlation, and ~70%
reach statistical significance (Spearman's correlation computed on the highly variable
genes having more than 50 normalized counts in at least 3 slices), further supporting
the hypothesis of the bilateral symmetry. Hence, after replicates were averaged, LML
axis was made symmetric averaging positions 1:28 and 56:29. Moreover, *Olfrs* were
normalized by the geometric mean of neuronal markers *Omp*, *Gnal* and *Cnga2*, as done
previously (*11*).
517
518 To verify the presence of a spatial signal, we calculated the Moran's I and the
519 associated p-values for the top 100 Highly Variable genes along each axis using the
520 "Moran.I" function from the "ape" library in R with default parameters (*82*). The p-
521 values of the genes along each axis were combined with the Simes' method (*83*) using
522 the function combinePValues from the scran R library (Fig. S1E).
523
Identification of differentially expressed genes and gene clustering
Before testing for differential expression along a given axis, we filtered out genes
whose expression levels had low variability. To this aim, for each gene we estimated
their highest and lowest expression by taking the average of its three highest and three
lowest values respectively. Then, we considered for downstream analyses only the
genes that meet either of these two criteria: the highest expression value is greater
than or equal to 5 normalized counts and the fold-change between the highest and
lowest value is greater than 2; or the difference between the highest and the lowest
value is greater than or equal to 4 normalized counts.
The expression levels of the genes were binarized according to whether their value
was higher or lower than their median expression along the axis. Finally, we used the
"ts" function in R to transform the binarized expression values into time series
objects, and we applied on them the Ljung-Box test (Box.test function in R with
lag=(axis length)-10) which identifies genes with statistically significant
autocorrelations, i.e., with non-random expression patterns along an axis. The
resulting p-values were adjusted using the FDR method and genes with an FDR <
0.01 were considered as differentially expressed. For the next steps, the $\log_{10}$
normalized expression of differentially expressed genes along each axis was fitted with
a local regression using the locfit function in the R library locfit (*84*). Smoothing was
defined in the local polynomial model term of the locfit model using the function "lp"
from the same library with the following parameters: nn = 1 (Nearest neighbor
component of the smoothing parameter) and deg = 2 (degree of polynomial). The
fitted expression values of these genes along each axis were normalized between 0 and
1. Clustering was performed separately for each axis on the fitted and normalized
patterns of the differentially expressed genes. We used the R function "hclust" to
perform hierarchical clustering on the gene expression patterns, with a Spearman's

correlation-based distance (defined as $\sqrt{0.5\,(1-\rho)}$) and the "average" aggregation
method. The number of clusters were defined with the cutreeDynamic function from
the dynamicTreeCut R library, with the parameters minClusterSize = 50, method =
"hybrid" and deepSplit = 0. To visualize the data in two dimensions, we applied the
UMAP dimensionality reduction algorithm (umap function in the R library umap with
default options; see Fig. 2D) (*55, 85*). To analyze the relationship between the
expression patterns of genes along different axes, we computed the intersections of
the gene clusters between any pair of axes. The expected number of elements in each
intersection under the assumption of independent sets is given by:

559
$$|A \cap B|_{exp} = \frac{|A|\,|B|}{|A \cup B|}$$

where A and B indicate the sets of genes in two clusters identified along two different
axes and $|\cdot|$ indicates the cardinality of a set (i.e., the number of its elements). The
ratio between the observed and the expected number of elements in the intersection
$|A \cap B|_{obs}\,/\,|A \cap B|_{exp}$ quantifies the enrichment/depletion of genes having a given
pair of patterns across two axes with respect to the random case. The $\log_2$ values of
$(1 + |A \cap B|_{obs}\,/\,|A \cap B|_{exp})$ are shown in Fig. 2F

Combining Tomo-seq with single-cell RNA-seq data
The TPM (transcripts per million)-normalized single cell RNA-seq (scRNA-seq) data
collected from mouse olfactory epithelium available from (*54*) was used to identify
cell-type specific genes. To this aim, we computed the average expression level for
each cell type in the scRNA-seq dataset for all the differentially expressed genes that
we identified in our TOMO-seq data. The genes with an average expression above
100 TPM in mOSNs and below 10 TPM in all other cell types were considered
mOSN-specific. Conversely, genes with an average expression above 100 TPM in any
of the non-mOSN cell types and below 10 in mOSNs were considered to be specific
for non-mOSN cells.

Gene Ontology (GO) enrichment analysis
GO Enrichment analyses were performed using the GOrilla online tool (http://cbl-
gorilla.cs.technion.ac.il) with the option "Two unranked lists of genes (target and
background lists)". For each axis, we used as background list the list of the genes we
tested.

3D spatial reconstruction
The olfactory mucosa shape was obtained from publicly available images of the
mouse nasal cavity along the posterior to the anterior axis published in (*58*). The area
of the slices corresponding to the OM was manually selected and images of their

silhouettes were made. Those images were then transformed into binary matrices
having 1's in the area occupied by the OM and 0's in the remaining regions. The
binary matrices were rescaled in order to match the spatial resolution in our dataset,
which is composed of 54 voxels along the DV axis, 56 along the LML axis and 60
along the AP axis. Finally, matrices were piled in a 3D array in R to obtain an in-silico
representation of the 3D shape of the OM, which, in total, was composed of 77,410
voxels. To perform the 3D reconstruction of the expression pattern for a given gene,
first we normalized its expression levels by the volume of the slice at each
corresponding position along the three axes, which was estimated using our 3D in
silico representation of the OM. Then, we rescaled the data in such a way that the sum
of the expression levels along each axis was equal to the average expression computed
across the whole dataset. This rescaled dataset together with the binary matrix
representing the 3D OM shape was used as input of the Iterative Proportional Fitting
algorithm, which produced an estimation of the expression level of a gene in each
voxel (*30*). Iterations stopped when the differences between the true and the
reconstructed 1D values summed across the three axes was smaller than 1.
604
Definition of zones by topic modelling
In order to identify zones, we fitted a Latent Dirichlet Allocation (LDA) (*86*)
algorithm to the 3D gene expression patterns (in log10 scale) of the differentially
expressed *Olfrs* (689 *Olfrs* x 77,410 voxels).
The LDA algorithm was originally employed for document classification: based on
the words included in each document, LDA can identify "topics", in which the
documents can then be classified. Using this linguistic analogy, in our application of
LDA, we considered the genes as "documents", and the spatial locations as "words",
with the matrix of gene expression levels being the analogous of the "bag-of-word"
matrix (*63*). In this representation, the zones are the equivalent of "topics", and they
are automatically identified by LDA. We used the LDA implementation included in
the R package "Countclust (*87*) ", developed by based on the "maptpx" library (*88*),
which performs a maximum a posteriori estimation to for model fitting. LDA was run
for all possible numbers of topics K ∈ [2,9]. The following parameters were chosen:
convergence tolerance = 0.1; max time optimization step = 180 seconds; n_init = 3.
For each number of topics k, three independent runs were performed with different
starting points, in order to avoid biases due to the choice of the initial condition. We
estimated the number of topics by computing the log-likelihood for each value of K ∈
[2,9]. As seen in fig. S3A, while the log-likelihood is a monotonically increasing
function of the number of topic (as expected), for a number of topics around ~5 it
shows a "knee" and starts to increase more slowly. This suggests that ~5 is the
optimal number of topics needed to describe the complexity of the data without
overfitting. Hence, we fix a number of topics equal to 5; however, we also verified

that all our conclusions remain substantially unaffected if a different number of topics
(e.g., 4 or 6) is chosen.
After running LDA with K=5, we retrieved the model output, which consists of two
probability distributions: the first is P(position| k) with k ∈ [1,5], which is the
conditional probability distribution defining the topic k; the second probability
distribution is P(k | gene), namely the probability distribution that quantifies the
degrees of belonging of a given gene to the topics k∈[1,5]. With these probability
distributions, we can identify the spatial positions that form each topic and how the
different topics can be combined to generate the spatial expression pattern of each
gene.
Being a generative model, once trained, LDA can also decompose into topics the
spatial expression patterns of genes that were not used during the training procedure.
We exploited this feature of LDA to estimate the degrees of belonging of non-
olfactory receptor genes. To this aim, we utilized an algorithm based on the python
gensim library Lda.Model.inference function (*89*), using as input the estimated
probability distribution P(position | k) with k ∈ [1,5]. The model fitting was
performed using the Open Computing Cluster for Advanced data Manipulation
(OCCAM), the High-Performance Computer designed and managed in collaboration
between the University of Torino and the Torino division of the Istituto Nazionale di
Fisica Nucleare (*90*).
648
Definition of *Olfr* 3D indexes via diffusion pseudo-time
As explained in the section above, we can describe the spatial expression pattern of
each gene through a set of five numbers, which represent the degrees of belonging to
the five topics identified by LDA. We applied a diffusion map (*64*) to the degrees of
belonging of the *Olfrs* to visualize them in two dimensions by using the
"DiffusionMap" function from the "destiny" R package (*91*) (with
distance="rankcor" and default parameters). In this two-dimensional map, the *Olfrs*
are approximately distributed along a curve that joins the most dorsal/medial genes
(those in zones 1-2) with those that are more ventral/lateral (zones 3-5). To track the
position of the genes along this curve, we computed a diffusion pseudo-time (DPT)
coordinate (*65*) with the "DPT" function from the "destiny" R package (taking as
starting point the gene with the smallest first diffusion component among the genes
suggested by the function find_tips from the same package). In order to make the
indexes go from Dorsal to Ventral, as in previous studies (*14*), we reversed the order
of the DPT coordinates by substracting the maximum coordinate from all coordinates
and multiplying them by (-1). By doing this, we obtained for each *Olfr* an index, which
we called 3D index, representing its spatial expression pattern in the 3D space: more
dorsal/medial genes (zones 1-2) have smaller 3D indexes than *Olfrs* expressed in the
ventral/lateral regions (zones 3-5).

Prediction of zone index for undetected *Olfrs* with Random Forest

We fitted a Random Forest model to the 3D indexes of 681 of the 689 *Olfrs* we
characterized with our dataset (i.e., those that are located in genomic clusters). The
following nine features of each *Olfr* were used as predictors: genomic position (i.e.,
gene starting position divided by chromosome length); genomic cluster; genomic
cluster length; number of *Olfrs* in the genomic cluster; number of enhancers in the
genomic cluster; cluster position (i.e., starting position of the cluster divided by the
chromosome length); distance to the closest enhancer; gene position within the cluster
(i.e., the distance of the gene starting position from the end of the cluster divided by
the cluster length); and phylogenetic class.
These features were computed using the mm10 mouse genome in Biomart (*92*), while
the list of enhancers and the genomic clusters assigned to each *Olfr* were taken from
(*69*). The Random Forest model was fitted with the function "randomForest" (R
library "randomForest" (*93*), with option "na.action = na.omit"). Afterwards, we
performed a cross-validation test with the function "rf.crossValidation" from the
"rfUtilities" package (*94*) with default parameters. Over 100 cross-validation iterations,
the root mean square error (RMSE) was $\lesssim 10\%$ of the mean 3D index. The feature
importance was computed with the "importance" function from the randomForest
library with default parameters. Finally, the Random Forest model trained on the 681
Olfrs was used to predict the 3D indexes of 697 *Olfrs* that were too lowly expressed or
were undetected in our dataset. Overall, we were able to compute or predict with
Random Forest a 3D index for all the *Olfrs* annotated in the mouse genome, except
for 28 of them that do not have any genomic cluster assigned.

Odorant information and Olfr-ligand pairs

All odorant structures and associated CAS numbers were retrieved from either Sigma-
Aldrich (www.sigmaaldrich.com) or PubChem (https://pubchem.ncbi.nlm.nih.gov).
A comprehensive catalog of the cognate mouse Olfr-ligand pairs was collected (last
update: March 2021) by combining data from the ODORactor database (*95*) and
additional literature sources (*4, 7, 74, 96-123*).
This catalog includes 738 Olfr-ligand interactions for a total of 153 Olfrs and 221
odorants. These 153 *Olfrs* include 100 spatial *Olfrs* in our dataset and for which we
have 3D indexes, and 49 additional *Olfrs* with predicted 3D indexes (see above). Next,
we checked whether *Olfrs* pairs sharing at least one cognate ligand have more similar
spatial expression patterns than pairs not sharing ligands. To do this, we computed
the absolute values of the differences between the 3D indexes (Δ) of 1706 pairs of
ORs sharing at least one odorant and 9922 pairs of ORs that are known to bind to
different odorants (Fig. 5B). The two sets of Δ values were significantly different
(Mann-Whitney U test, p-value $< 2.2 \times 10^{-16}$).

Estimation of the air/mucus partition coefficients

For each odorant, we estimated the air/mucus partition coefficients (K_{am}) as done previously (*21, 70*). Briefly, we calculated the air/water partition coefficients (K_{aw}) for each odorant from the Henry's Law constants obtained using the HENRYWIN model in the US EPA Estimation Program Interface (EPI) Suite (version 4.11; www.epa.gov/oppt/exposure/pubs/episuite.htm). Then, we computed the air/mucus partition coefficients (K_{am}) according to the formula:

$$Log(K_{am}) = 0.524 \cdot Log(K_{aw}) \cdot Log(K_{ow})$$

where K_{ow} indicates the octanol/water partition coefficient, which were obtained using the KOWWIN model in the EPI Suite.

Finally, we plotted in Fig. 5C the $Log(K_{am})$ of each odorant versus the mean 3D index of the cognate *Olfrs*, using only the 110 odorants for which we could find two or more cognate receptors. These two quantities have a Spearman's correlation coefficient equal to 0.55 (p-value=7×10^{-10}). The correlation remained significant even when we removed *Olfrs* whose indexes were predicted with the Random Forest (Spearman's correlation = 0.69, p-value= 2×10^{-3}), or when only class II *Olfrs* were included (Spearman's correlation = 0.5, p-value= 1.3×10^{-7}).

In-situ hybridization

In-situ hybridization was basically performed as previously described (Ibarra-Soria et al. 2017). 12-week-old male C57BL/6J mice were perfused with 4% paraformaldehyde and decalcified in RNase-free 0.45M EDTA solution (in 1x PBS) for two weeks. Decalcified heads were cryoprotected in RNase-free 30% sucrose solution (1x PBS), dried, embedded in OCT embedding medium, and frozen at -80°C. Sequential 16 mm sections were prepared with a cryostat and the sections were hybridized to digoxigenin-labeled cRNA probes prepared from the different genes using the following oligonucleotides: Cytl (5'–AAAGACACTACCTCTGTTGCTGCTG – 3' and 5' – GTAAGCAGAGACCAGAAAGAAGAGTG – 3'), Moxd2 (5' – TGTACCTTTCTCCCACTCCCTATTGTC – 3' and 5'– CCCATGCAACTGGAGATGTTAATTCTG – 3'), Olfr309 (5'– TACAATGGCCTATGACCGCTATGTG – 3' and 5' – TCCTGACTGCATCTCTTTGTTCCTG – 3') , Spen (5'– GGTGGGAAACTTACCGGAGAACGTG – 3' and 5' – TGCTGCTGATGGAGTCACTACTG – 3'), Olfr727 (5' – CGCTATGTTGCAATATGCAAGCCTC – 3' and 5'– GCTTTGACATTGCTGCTTTCACCTC – 3'). The PCR products were cloned into pGEM-T Easy vector and the probes were obtained by in vitro transcription of the

747 plasmids, using SP6 or T7 RNA Polymerases (Ambion) and DIG RNA Labeling mix
748 (Roche).
749
Data availability

751 The processed data can be visualized and downloaded from the website
752 http://atlas3dnose.helmholtz-muenchen.de:3838/atlas3Dnose.

Acknowledgments

754 We would like to thank Maria-Elena Torres-Padilla, Bob Datta, and the members of
755 the Saraiva and Scialdone Labs for the constructive feedback, and helpful comments.
756 We are thankful to Giorgia Greco for insightful discussions about chemical properties
757 of ligands. This work was supported by the Helmholtz Association (Scialdone Lab)
758 and Sidra Medicine (SDR400040).
759

Author contributions

M.L.R.T.S. analyzed data and wrote the initial version of the manuscript.
E.A.M. performed the RNA-seq experiments and analyzed data.
T.S.N., L.S.M., and S.L. performed experiments.
M.M., S.S.Y.H., J.D.M., F.V., M.C., and M.O. analyzed data.
E.G., J.R., D.W.L., and B.M. analyzed data, and helped write the manuscript.
A.S. and L.R.S. conceived and supervised the project, analyzed data, and wrote the
final version of the manuscript.
768
769

References

1. E. Kandel, J. Schwartz, T. Jessell, S. Siegelbaum, A. Hudspeth, *Principles of neural science*. (McGraw-Hill, New York, ed. 5th, 2013), pp. l, 1709 p.
2. L. Buck, R. Axel, A novel multigene family may encode odorant receptors: a molecular basis for odor recognition. *Cell* **65**, 175-187 (1991).
3. A. Chess, I. Simon, H. Cedar, R. Axel, Allelic inactivation regulates olfactory receptor gene expression. *Cell* **78**, 823-834 (1994).
4. B. Malnic, J. Hirono, T. Sato, L. B. Buck, Combinatorial receptor codes for odors. *Cell* **96**, 713-723 (1999).
5. N. K. Hanchate *et al.*, Single-cell transcriptomics reveals receptor transformations during olfactory neurogenesis. *Science* **350**, 1251-1255 (2015).
6. L. R. Saraiva *et al.*, Hierarchical deconstruction of mouse olfactory sensory neurons: from whole mucosa to single-cell RNA-seq. *Scientific reports* **5**, 18178 (2015).
7. K. Nara, L. R. Saraiva, X. Ye, L. B. Buck, A large-scale analysis of odor coding in the olfactory epithelium. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **31**, 9179-9191 (2011).
8. P. Mombaerts *et al.*, Visualizing an olfactory sensory map. *Cell* **87**, 675-686 (1996).
9. K. J. Ressler, S. L. Sullivan, L. B. Buck, Information coding in the olfactory system: evidence for a stereotyped and highly organized epitope map in the olfactory bulb. *Cell* **79**, 1245-1255 (1994).
10. R. Vassar *et al.*, Topographic organization of sensory projections to the olfactory bulb. *Cell* **79**, 981-991 (1994).
11. X. Ibarra-Soria *et al.*, Variation in olfactory neuron repertoires is genetically controlled and environmentally modulated. *eLife* **6**, (2017).
12. K. J. Ressler, S. L. Sullivan, L. B. Buck, A zonal organization of odorant receptor gene expression in the olfactory epithelium. *Cell* **73**, 597-609 (1993).
13. R. Vassar, J. Ngai, R. Axel, Spatial segregation of odorant receptor expression in the mammalian olfactory epithelium. *Cell* **74**, 309-318 (1993).
14. K. Miyamichi, S. Serizawa, H. M. Kimura, H. Sakano, Continuous and overlapping expression domains of odorant receptor genes in the olfactory epithelium determine the dorsal/ventral positioning of glomeruli in the olfactory bulb. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **25**, 3586-3592 (2005).
15. J. Strotmann, I. Wanner, J. Krieger, K. Raming, H. Breer, Expression of odorant receptors in spatially restricted subsets of chemosensory neurones. *Neuroreport* **3**, 1053-1056 (1992).
16. B. Zapiec, P. Mombaerts, The Zonal Organization of Odorant Receptor Gene Choice in the Main Olfactory Epithelium of the Mouse. *Cell reports* **30**, 4220-4234 e4225 (2020).
17. E. Bashkirova *et al.*, Homeotic Regulation of Olfactory Receptor Choice via NFI-dependent Heterochromatic Silencing and Genomic Compartmentalization. *bioRxiv*, 2020.2008.2030.274035 (2020).
18. D. M. Coppola, C. T. Waggener, S. M. Radwani, D. A. Brooks, An electroolfactogram study of odor response patterns from the mouse olfactory

epithelium with reference to receptor zones and odor sorptiveness. *J Neurophysiol* **109**, 2179-2191 (2013).

19. M. Ma, G. M. Shepherd, Functional mosaic organization of mouse olfactory receptor neurons. *Proceedings of the National Academy of Sciences of the United States of America* **97**, 12869-12874 (2000).

20. M. M. Mozell, The spatiotemporal analysis of odorants at the level of the olfactory receptor sheet. *J Gen Physiol* **50**, 25-41 (1966).

21. J. W. Scott, L. Sherrill, J. Jiang, K. Zhao, Tuning to odor solubility and sorption pattern in olfactory epithelial responses. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **34**, 2025-2036 (2014).

22. T. A. Schoenfeld, T. A. Cleland, Anatomical contributions to odorant sampling and representation in rodents: zoning in on sniffing behavior. *Chemical senses* **31**, 131-144 (2006).

23. L. Secundo, K. Snitz, N. Sobel, The perceptual logic of smell. *Curr Opin Neurobiol* **25**, 107-115 (2014).

24. D. M. Coppola, E. Fitzwater, A. D. Rygg, B. A. Craven, Tests of the chromatographic theory of olfaction with highly soluble odors: a combined electro-olfactogram and computational fluid dynamics study in the mouse. *Biol Open* **8**, (2019).

25. T. Abaffy, A. R. Defazio, The location of olfactory receptors within olfactory epithelium is independent of odorant volatility and solubility. *BMC research notes* **4**, 137 (2011).

26. S. M. Kurian *et al.*, Odor coding in the mammalian olfactory epithelium. *Cell Tissue Res* **383**, 445-456 (2021).

27. M. Asp, J. Bergenstrahle, J. Lundeberg, Spatially Resolved Transcriptomes-Next Generation Tools for Tissue Exploration. *Bioessays* **42**, e1900221 (2020).

28. M. Asp *et al.*, A Spatiotemporal Organ-Wide Gene Expression and Cell Atlas of the Developing Human Heart. *Cell* **179**, 1647-1660 e1619 (2019).

29. G. Peng *et al.*, Spatial Transcriptome for the Molecular Annotation of Lineage Fates and Cell Identity in Mid-gastrula Mouse Embryo. *Dev Cell* **36**, 681-697 (2016).

30. J. P. Junker *et al.*, Genome-wide RNA Tomography in the zebrafish embryo. *Cell* **159**, 662-675 (2014).

31. K. Achim *et al.*, High-throughput spatial mapping of single-cell RNA-seq data to tissue of origin. *Nature biotechnology* **33**, 503-509 (2015).

32. C. Schmal, J. Myung, H. Herzel, G. Bordyugov, Moran's I quantifies spatio-temporal pattern formation in neural imaging data. *Bioinformatics* **33**, 3072-3079 (2017).

33. F. Gussing, S. Bohm, NQO1 activity in the main and the accessory olfactory systems correlates with the zonal topography of projection maps. *Eur J Neurosci* **19**, 2511-2518 (2004).

34. Y. Oka *et al.*, O-MACS, a novel member of the medium-chain acyl-CoA synthetase family, specifically expressed in the olfactory epithelium in a zone-specific manner. *Eur J Biochem* **270**, 1995-2004 (2003).

35.   M. Alenius, S. Bohm, Identification of a novel neural cell adhesion molecule-related gene with a potential role in selective axonal projection. *J Biol Chem* **272**, 26083-26086 (1997).

36.   T. T. Yu *et al.*, Differentially expressed transcripts from phenotypically identified olfactory sensory neurons. *J Comp Neurol* **483**, 251-262 (2005).

37.   S. D. Liberles, L. B. Buck, A second class of chemosensory receptors in the olfactory epithelium. *Nature* **442**, 645-650 (2006).

38.   R. Pacifico, A. Dewan, D. Cawley, C. Guo, T. Bozza, An olfactory subsystem that mediates high-sensitivity detection of volatile amines. *Cell reports* **2**, 76-88 (2012).

39.   H. J. Fulle *et al.*, A receptor guanylyl cyclase expressed specifically in olfactory sensory neurons. *Proceedings of the National Academy of Sciences of the United States of America* **92**, 3571-3575 (1995).

40.   P. L. Greer *et al.*, A Family of non-GPCR Chemosensors Defines an Alternative Logic for Mammalian Olfaction. *Cell* **165**, 1734-1748 (2016).

41.   J. H. Cho, M. Lepine, W. Andrews, J. Parnavelas, J. F. Cloutier, Requirement for Slit-1 and Robo-2 in zonal segregation of olfactory sensory neuron axons in the main olfactory bulb. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **27**, 9094-9104 (2007).

42.   J. F. Cloutier *et al.*, Neuropilin-2 mediates axonal fasciculation, zonal segregation, but not axonal convergence, of primary accessory olfactory neurons. *Neuron* **33**, 877-892 (2002).

43.   Y. Yoshihara *et al.*, OCAM: A new member of the neural cell adhesion molecule family related to zone-to-zone projection of olfactory and vomeronasal axons. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **17**, 5830-5842 (1997).

44.   I. Tietjen, J. Rihel, C. G. Dulac, Single-cell transcriptional profiles and spatial patterning of the mammalian olfactory epithelium. *Int J Dev Biol* **49**, 201-207 (2005).

45.   I. Tietjen *et al.*, Single-cell transcriptional analysis of neuronal progenitors. *Neuron* **38**, 161-175 (2003).

46.   S. S. Wang, J. W. Lewcock, P. Feinstein, P. Mombaerts, R. R. Reed, Genetic disruptions of O/E2 and O/E3 genes reveal involvement in olfactory receptor neuron projection. *Development* **131**, 1377-1388 (2004).

47.   E. M. Norlin *et al.*, Evidence for gradients of gene expression correlating with zonal topography of the olfactory sensory map. *Mol Cell Neurosci* **18**, 283-295 (2001).

48.   D. M. Juilfs *et al.*, A subset of olfactory neurons that selectively express cGMP-stimulated phosphodiesterase (PDE2) and guanylyl cyclase-D define a unique olfactory signal transduction pathway. *Proceedings of the National Academy of Sciences of the United States of America* **94**, 3388-3395 (1997).

49.   P. M. Heron, A. J. Stromberg, P. Breheny, T. S. McClintock, Molecular events in the cell types of the olfactory epithelium during adult neurogenesis. *Molecular brain* **6**, 49 (2013).

50. C. E. Peluso, W. Jang, U. C. Drager, J. E. Schwob, Differential expression of components of the retinoic acid signaling pathway in the adult mouse olfactory epithelium. *J Comp Neurol* **520**, 3707-3726 (2012).

51. G. K. Whitby-Logan, M. Weech, E. Walters, Zonal expression and activity of glutathione S-transferase enzymes in the mouse olfactory mucosa. *Brain Res* **995**, 151-157 (2004).

52. A. Miyawaki, H. Homma, H. Tamura, M. Matsui, K. Mikoshiba, Zonal distribution of sulfotransferase for phenol in olfactory sustentacular cells. *EMBO J* **15**, 2050-2055 (1996).

53. C. D. Duggan, S. DeMaria, A. Baudhuin, D. Stafford, J. Ngai, Foxg1 is required for development of the vertebrate olfactory system. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **28**, 5229-5239 (2008).

54. R. B. Fletcher *et al.*, Deconstructing Olfactory Stem Cell Trajectories at Single-Cell Resolution. *Cell Stem Cell* **20**, 817-830 e818 (2017).

55. E. Becht *et al.*, Dimensionality reduction for visualizing single-cell data using UMAP. *Nat Biotechnol*, (2018).

56. V. Vedin, M. Molander, S. Bohm, A. Berghard, Regional differences in olfactory epithelial homeostasis in the adult mouse. *J Comp Neurol* **513**, 375-384 (2009).

57. M. D. Nickell, P. Breheny, A. J. Stromberg, T. S. McClintock, Genomics of mature and immature olfactory sensory neurons. *J Comp Neurol* **520**, 2608-2629 (2012).

58. A. W. Barrios, G. Nunez, P. Sanchez Quinteiro, I. Salazar, Anatomy, histochemistry, and immunohistochemistry of the olfactory subsystems in mice. *Front Neuroanat* **8**, 63 (2014).

59. S. E. Fienberg, An Iterative Procedure for Estimation in Contingency Tables. *The Annals of Mathematical Statistics* **41**, 907-917, 911 (1970).

60. S. Zhu *et al.*, Protein Cytl1: its role in chondrogenesis, cartilage homeostasis, and disease. *Cell Mol Life Sci* **76**, 3515-3523 (2019).

61. Y. Shin, Y. Won, J. I. Yang, J. S. Chun, CYTL1 regulates bone homeostasis in mice by modulating osteogenesis of mesenchymal stem cells and osteoclastogenesis of bone marrow-derived macrophages. *Cell Death Dis* **10**, 47 (2019).

62. C. J. Goh, D. Choi, D. B. Park, H. Kim, Y. Hahn, MOXD2, a Gene Possibly Associated with Olfaction, Is Frequently Inactivated in Birds. *PloS one* **11**, e0152431 (2016).

63. L. Liu, L. Tang, W. Dong, S. Yao, W. Zhou, An overview of topic modeling and its current applications in bioinformatics. *Springerplus* **5**, 1608 (2016).

64. L. Haghverdi, F. Buettner, F. J. Theis, Diffusion maps for high-dimensional single-cell analysis of differentiation data. *Bioinformatics* **31**, 2989-2998 (2015).

65. L. Haghverdi, M. Buttner, F. A. Wolf, F. Buettner, F. J. Theis, Diffusion pseudotime robustly reconstructs lineage branching. *Nat Methods* **13**, 845-848 (2016).

66. L. Tan, X. S. Xie, A Near-Complete Spatial Map of Olfactory Receptors in the Mouse Main Olfactory Epithelium. *Chemical senses* **43**, 427-432 (2018).

67.   X. Zhang *et al.*, High-throughput microarray detection of olfactory receptor gene
expression in the mouse. *Proceedings of the National Academy of Sciences of*
the United States of America **101**, 14168-14173 (2004).
68.   S. L. Sullivan, M. C. Adamson, K. J. Ressler, C. A. Kozak, L. B. Buck, The
chromosomal distribution of mouse odorant receptor genes. *Proceedings of the*
National Academy of Sciences of the United States of America **93**, 884-888
(1996).
69.   K. Monahan *et al.*, Cooperative interactions enable singular olfactory receptor
expression in mouse olfactory neurons. *Elife* **6**, (2017).
70.   A. D. Rygg, B. Van Valkenburgh, B. A. Craven, The Influence of Sniffing on
Airflow and Odorant Deposition in the Canine Nasal Cavity. *Chem Senses* **42**,
683-698 (2017).
71.   J. W. Scott, P. E. Scott-Johnson, The electroolfactogram: a review of its history
and uses. *Microsc Res Tech* **58**, 152-160 (2002).
72.   L. Secundo *et al.*, Individual olfactory perception reveals meaningful nonolfactory
genetic information. *Proceedings of the National Academy of Sciences of the*
United States of America **112**, 8750-8755 (2015).
73.   Daniel M. Bear, J.-M. Lassance, Hopi E. Hoekstra, Sandeep R. Datta, The
Evolving Neural and Genetic Architecture of Vertebrate Olfaction. *Current*
Biology **26**, R1039-R1049 (2016).
74.   L. R. Saraiva *et al.*, A transcriptomic atlas of mammalian olfactory mucosae
reveals an evolutionary influence on food odor detection in humans. *Sci Adv* **5**,
eaax0396 (2019).
75.   L. F. Horowitz, L. R. Saraiva, D. Kuang, K. H. Yoon, L. B. Buck, Olfactory
receptor patterning in a higher primate. *J Neurosci* **34**, 12241-12252 (2014).
76.   J. E. R. Octura, K. I. Maeda, Y. Wakabayashi, Structure and zonal expression of
olfactory receptors in the olfactory epithelium of the goat, Capra hircus. *J Vet*
Med Sci **80**, 913-920 (2018).
77.   D. Manoel *et al.*, Deconstructing the mouse olfactory percept through an
ethological atlas. *Curr Biol*, (2021).
78.   A. Dobin *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-
21 (2013).
79.   H. Li *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics*
25, 2078-2079 (2009).
80.   S. Anders, P. T. Pyl, W. Huber, *HTSeq - A Python framework to work with high-*
throughput sequencing data. (2014).
81.   A. T. Lun, D. J. McCarthy, J. C. Marioni, A step-by-step workflow for low-level
analysis of single-cell RNA-seq data with Bioconductor. *F1000Res* **5**, 2122
(2016).
82.   E. Paradis, J. Claude, K. Strimmer, APE: Analyses of Phylogenetics and
Evolution in R language. *Bioinformatics* **20**, 289-290 (2004).
83.   R. J. Simes, An Improved Bonferroni Procedure for Multiple Tests of
Significance. *Biometrika* **73**, 751-754 (1986).
84.   C. Loader. (2007).
85.   L. McInnes, J. Healy, J. Melville, UMAP: Uniform Manifold Approximation and
Projection for Dimension Reduction. 2018.

86.   D. M. Blei, A. Y. Ng, M. I. Jordan, Latent dirichlet allocation. *J. Mach. Learn. Res.* **3**, 993–1022 (2003).

87.   K. K. Dey, C. J. Hsiao, M. Stephens, Visualizing the structure of RNA-seq expression data using grade of membership models. *PLoS Genet* **13**, e1006599 (2017).

88.   M. Taddy, paper presented at the Proceedings of the Fifteenth International Conference on Artificial Intelligence and Statistics, Proceedings of Machine Learning Research, 2012.

89.   R. Rehurek, P. Sojka, paper presented at the Proceedings of the LREC 2010 Workshop on New Challenges for NLP Frameworks, Valletta, Malta, 2010.

90.   M. Aldinucci *et al.*, OCCAM: a flexible, multi-purpose and extendable HPC cluster. *Journal of Physics: Conference Series* **898**, 082039 (2017).

91.   P. Angerer *et al.*, destiny: diffusion maps for large-scale single-cell data in R. *Bioinformatics* **32**, 1241-1243 (2016).

92.   R. J. Kinsella *et al.*, Ensembl BioMarts: a hub for data retrieval across taxonomic space. *Database (Oxford)* **2011**, bar030 (2011).

93.   A. Liaw, M. Wiener, Classification and Regression by randomForest. *R News* **2**, 18--22 (2002).

94.   T. A. Rather, S. Kumar, J. A. Khan, Multi-scale habitat modelling and predicting change in the distribution of tiger and leopard using random forest algorithm. *Scientific reports* **10**, 11473 (2020).

95.   X. Liu *et al.*, ODORactor: a web server for deciphering olfactory coding. *Bioinformatics* **27**, 2302-2303 (2011).

96.   B. Malnic, P. A. Godfrey, L. B. Buck, The human olfactory receptor gene family. *Proceedings of the National Academy of Sciences of the United States of America* **101**, 2584-2589 (2004).

97.   T. Bozza, P. Feinstein, C. Zheng, P. Mombaerts, Odorant receptor expression defines functional units in the mouse olfactory system. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **22**, 3033-3043 (2002).

98.   R. C. Araneda, Z. Peterlin, X. Zhang, A. Chesler, S. Firestein, A pharmacological profile of the aldehyde receptor repertoire in rat olfactory epithelium. *J Physiol* **555**, 743-756 (2004).

99.   P. A. Godfrey, B. Malnic, L. B. Buck, The mouse olfactory receptor gene family. *Proceedings of the National Academy of Sciences of the United States of America* **101**, 2156-2161 (2004).

100.   E. Shirokova *et al.*, Identification of specific ligands for orphan olfactory receptors. G protein-dependent agonism and antagonism of odorants. *J Biol Chem* **280**, 11807-11815 (2005).

101.   K. Kajiya *et al.*, Molecular bases of odor discrimination: Reconstitution of olfactory receptors that recognize overlapping sets of odorants. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **21**, 6018-6025 (2001).

102.   Y. Oka, M. Omura, H. Kataoka, K. Touhara, Olfactory receptor antagonism between odorants. *Embo J* **23**, 120-126 (2004).

103. I. Gaillard *et al.*, A single olfactory receptor specifically binds a set of odorant molecules. *Eur J Neurosci* **15**, 409-418 (2002).
104. M. Q. Nguyen, Z. Zhou, C. A. Marks, N. J. Ryba, L. Belluscio, Prominent roles for odorant receptor coding sequences in allelic exclusion. *Cell* **131**, 1009-1017 (2007).
105. W. B. Floriano, N. Vaidehi, W. A. Goddard, 3rd, M. S. Singer, G. M. Shepherd, Molecular mechanisms underlying differential odor responses of a mouse olfactory receptor. *Proceedings of the National Academy of Sciences of the United States of America* **97**, 10712-10716 (2000).
106. Y. Oka *et al.*, Odorant receptor map in the mouse olfactory bulb: in vivo sensitivity and specificity of receptor-defined glomeruli. *Neuron* **52**, 857-869 (2006).
107. S. E. Repicky, C. W. Luetje, Molecular receptive range variation among mouse odorant receptors for aliphatic carboxylic acids. *J Neurochem* **109**, 193-202 (2009).
108. X. Grosmaitre *et al.*, SR1, a mouse odorant receptor with an unusually broad response profile. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **29**, 14545-14552 (2009).
109. X. Grosmaitre, A. Vassalli, P. Mombaerts, G. M. Shepherd, M. Ma, Odorant responses of olfactory sensory neurons expressing the odorant receptor MOR23: a patch clamp analysis in gene-targeted mice. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 1970-1975 (2006).
110. Y. Oka, Y. Takai, K. Touhara, Nasal airflow rate affects the sensitivity and pattern of glomerular odorant responses in the mouse olfactory bulb. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **29**, 12070-12078 (2009).
111. H. Saito, M. Kubota, R. W. Roberts, Q. Chi, H. Matsunami, RTP family members induce functional expression of mammalian odorant receptors. *Cell* **119**, 679-691 (2004).
112. H. Zhuang, H. Matsunami, Synergism of accessory factors in functional expression of mammalian odorant receptors. *J Biol Chem* **282**, 15284-15293 (2007).
113. K. Yoshikawa, K. Touhara, Myr-Ric-8A enhances G(alpha15)-mediated Ca2+ response of vertebrate olfactory receptors. *Chemical senses* **34**, 15-23 (2009).
114. T. Abaffy, H. Matsunami, C. W. Luetje, Functional analysis of a mammalian odorant receptor subfamily. *J Neurochem* **97**, 1506-1518 (2006).
115. A. Dunkel *et al.*, Nature's chemical signatures in human olfaction: a foodborne perspective for future biotechnology. *Angew Chem Int Ed Engl* **53**, 7124-7143 (2014).
116. K. Yoshikawa, H. Nakagawa, N. Mori, H. Watanabe, K. Touhara, An unsaturated aliphatic alcohol as a natural ligand for a mouse odorant receptor. *Nat Chem Biol* **9**, 160-162 (2013).
117. Y. Jiang *et al.*, Molecular profiling of activated olfactory neurons identifies odorant receptors for odors in vivo. *Nat Neurosci* **18**, 1446-1454 (2015).

118. H. Saito *et al.*, Immobility responses are induced by photoactivation of single glomerular species responsive to fox odour TMT. *Nature communications* **8**, 16011 (2017).

119. B. von der Weid *et al.*, Large-scale transcriptional profiling of chemosensory neurons identifies receptor-ligand pairs in vivo. *Nat Neurosci* **18**, 1455-1463 (2015).

120. M. Shirasu *et al.*, Olfactory receptor and neural pathway responsible for highly selective sensing of musk odors. *Neuron* **81**, 165-178 (2014).

121. Y. Yu *et al.*, Responsiveness of G protein-coupled odorant receptors is partially attributed to the activation mechanism. *Proceedings of the National Academy of Sciences of the United States of America* **112**, 14966-14971 (2015).

122. E. M. Jones *et al.*, A Scalable, Multiplexed Assay for Decoding GPCR-Ligand Interactions with RNA Sequencing. *Cell Syst* **8**, 254-260 e256 (2019).

123. P. Pfister *et al.*, Odorant Receptor Inhibition Is Fundamental to Odor Encoding. *Curr Biol* **30**, 2574-2587 e2576 (2020).