A CRISPRi/a platform in iPSC-derived microglia uncovers regulators of disease states

Nina M. Dräger[1], Sydney M. Sattler[1,=], Cindy Tzu-Ling Huang[2,=], Olivia M. Teter[1,3], Kun Leng[1,4,5], Sayed Hadi Hashemi[6], Jason Hong[1], Giovanni Aviles[1], Claire D. Clelland[2,7], Lihong Zhan[2], Joe C. Udeochu[2], Lay Kodama[2,5,8], Andrew B. Singleton[9,10], Mike A. Nalls[9,10,11], Justin Ichida[12,13,14], Michael E. Ward[15], Faraz Faghri[6,9,10,11], Li Gan[2,7,16,*], Martin Kampmann[1,17,18,*]

[1] Institute for Neurodegenerative Diseases, University of California, San Francisco, San Francisco, CA, USA

[2] Gladstone Institute of Neurological Disease, San Francisco, CA, USA

[3] UC Berkeley-UCSF Graduate Program in Bioengineering, University of California San Francisco, San Francisco, CA, USA

[4] Biomedical Sciences Graduate Program, University of California, San Francisco, San Francisco, CA, USA

[5] Medical Scientist Training Program, University of California, San Francisco, San Francisco, CA, USA

[6] Department of Computer Science, University of Illinois at Urbana-Champaign, Urbana, IL, USA

[7] Department of Neurology, University of California, San Francisco, CA, 94158, USA.

[8] Neuroscience Graduate Program, University of California, San Francisco, San Francisco, CA, USA

[9] Center for Alzheimer's and Related Dementias, National Institutes of Health, Bethesda, MD, USA

[10] Laboratory of Neurogenetics, National Institute on Aging, National Institutes of Health, Bethesda, MD, USA

[11] Data Tecnica International, LLC, Glen Echo, MD, USA

[12] Department of Stem Cell Biology and Regenerative Medicine, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA

[13] Eli and Edythe Broad CIRM Center for Regenerative Medicine and Stem Cell Research at USC, Los Angeles, CA, USA

[14] Zilkha Neurogenetic Institute, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA

[15] National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, MD, USA

[16] Helen and Robert Appel Alzheimer's Disease Research Institute, Brain and Mind Research Institute, Weill Cornell Medicine, New York, NY, USA

[17] Department of Biochemistry and Biophysics, University of California, San Francisco, San Francisco, CA, USA

[18] Chan Zuckerberg Biohub, San Francisco, CA, USA

[=] Equal contribution

*Corresponding authors: lig2033@med.cornell.edu, martin.kampmann@ucsf.edu.

ABSTRACT

Microglia are emerging as key drivers of neurological diseases. However, we lack a systematic understanding of the underlying mechanisms. Here, we present a screening platform to systematically elucidate functional consequences of genetic perturbations in human iPSC-derived microglia. We developed an efficient eight-day protocol for the generation of microglia-like cells based on the inducible expression of six transcription factors. We established inducible CRISPR interference and activation in this system and conducted three screens targeting the "druggable genome". These screens uncovered genes controlling microglia survival, activation and phagocytosis, including neurodegeneration-associated genes. A screen with single-cell RNA sequencing as the readout revealed that these microglia adopt a spectrum of states mirroring those observed in human brains and identified regulators of these states. A disease-associated state characterized by SPP1 expression was selectively depleted by CSF1R inhibition. Thus, our platform can systematically uncover regulators of microglia states, enabling their functional characterization and therapeutic targeting.

INTRODUCTION

Historically, neuroscience has investigated brain function and disease through a neuron-centric lens, relegating glia to the sidelines. Neuroinflammation has typically been viewed as a secondary, reactive aspect of disease. More recently, however, key roles have emerged for glial cell types, including microglia, the innate immune cells of the brain. It is now widely accepted that microglia have a central role in brain development and homeostasis as well as in the pathogenesis of many brain disorders[1]. Over the last decade, human genetics have pointed to a central role for microglia in brain diseases such as Alzheimer's Disease (AD)[2], where specific disease-associated genetic variants likely act in microglia, redefining them as potential drivers of AD. To understand the molecular mechanisms underlying the role of microglia in disease and target them therapeutically, it is necessary to bridge the gap between disease-associated genetic variants and changes in microglial function.

A major challenge is that microglia adopt a large number of distinct functional states in health and disease. In homeostatic states, microglia survey their local environment, phagocytose myelin and cell debris, and monitor neuronal activity[3]. In disease states, microglia can play beneficial roles, but they are also responsible for an increased production of proinflammatory cytokines, an exacerbated inflammatory response and secrete toxic factors to directly or indirectly damage neurons[4, 5]. Eventually, microglia exhibit pathological features, such as mitochondrial and endolysosomal dysfunction, impaired phagocytosis, and increased production of reactive oxygen species (ROS)[6].

Microglial states in health and disease are actively being mapped on the molecular level in mice and humans[7-13]. However, we do not systematically understand how these distinct microglial states contribute to brain function and disease, or the molecular mechanisms regulating these states.

A promising approach to tackle these questions is enabled by CRISPR-based functional
genomics in differentiated human cell types[14]. Pooled CRISPR interference (CRISPRi) and
CRISPR activation (CRISPRa) screens enable scalable modeling of changes in gene expression
and genetic screens to uncover regulatory mechanisms. When combined with induced pluripotent
stem cell (iPSC) technology, they enable the investigation of cell-type specific biology in human
cells, including those derived from patients[14]. We recently provided a proof of principle for this
strategy by establishing CRISPRi and CRISPRa platforms for genetic screens in iPSC-derived
neurons[15, 16]. However, such screens have not previously been implemented in iPSC-derived
microglia due to challenges inherent in available differentiation protocols. Pooled CRISPR
screens rely on lentiviral transduction to introduce libraries of single guide RNAs (sgRNAs), but
mature microglia are difficult to transduce with lentivirus. This problem could be overcome by
introducing sgRNAs at the iPSC stage. However, most existing protocols are lengthy and aim to
recapitulate human microglia ontogeny[17-23], resulting in population bottlenecks during
differentiation, which can skew the representation of the sgRNA library.
103
To overcome these challenges, we developed a different approach for the generation of iPSC-
derived microglia by generating a human iPSC line inducibly expressing six transcription factors
that enable the generation of microglia-like cells in a rapid and efficient eight-day protocol.
These induced-transcription factor microglia-like cells (iTF-Microglia) resemble other iPSC-
derived microglia in their expression profiles, response to inflammatory stimuli, phagocytic
capabilities, and capacity to be co-cultured with iPSC-derived neurons[17-23]. By integrating
inducible CRISPRi/a machinery into this cell line, we developed a genetic screening system that
enables robust knockdown and overexpression of endogenous genes in human microglia. Using
this platform, we conducted pooled CRISPRi and CRISPRa screens for modifiers of survival,
phagocytosis and inflammatory activation, which uncovered microglia-specific genes controlling
these phenotypes. A screen with single-cell RNA sequencing as the readout revealed that these
microglia adopt a spectrum of states mirroring those observed in human brains, and pinpointed
regulators of specific states, which can enable the functional characterization and therapeutic
targeting of these states.
118
119
120 **RESULTS**
121
122 **Inducible transcription factors enable rapid and scalable production of microglia-like cells**
123 **(iTF-Microglia)**
124
High-throughput genetic CRISPRi/a screens are a powerful discovery tool in human iPSC-
derived neurons[15, 16]. However, such screens have not yet been conducted in iPSC-derived
microglia. One obstacle has been the fact that until very recently[24], existing differentiation
protocols involved a long, multi-step procedure, which recapitulates the human microglia
ontogeny. We set out to create a fast, robust and scalable differentiation protocol to differentiate
iPSCs to microglia-like cells for use in CRISPR screens. To this end, we developed a strategy
based on direct cell fate conversion by overexpression of microglia fate-determining
transcription factors.
133

Based on transcriptomic and developmental data[25-27], we selected six transcription factors highly
expressed in human microglia. PU.1 and interferon regulatory factor-8 (IRF-8) are known to be
crucial for microgliogenesis[28]. MAFB increases during microglia development and promotes an
anti-inflammatory phenotype[29], while CEBPα and CEBPβ[30] and IRF-5[31] regulate a wide range
of inflammatory mediators. We engineered an iPSC line with two integrated cassettes of three
transcription factors each. Cassettes for the doxycycline-inducible expression of transgenic
PU.1/CEBPβ/IRF5 and MAFB/CEBPα/IRF8 were integrated into the CLYBL and AAVS1 safe
harbor loci, respectively, in the WTC11 iPSC line (Fig. 1a).

We established a simple three-step protocol to differentiate these iPSCs into microglia-like cells,
which we will refer to as iTF-Microglia, in only 8 days (Fig. 1b). After doxycycline induction of
transcription factor expression on Day 0, media was supplemented with cytokines GM-CSF and
IL-34 on Day 2 to promote differentiation and survival. On Day 4, the media was additionally
supplemented with the cytokines M-CSF and TGF-β. iTF-Microglia reached a fully ramified
morphology on Day 8 and maintained excellent viability for at least another 8 days (Fig. 1b). We
generally have continued doxycycline supplementation beyond Day 8; however, this is not
necessary for survival (Extended Data Fig. 1a,b). We confirmed robust inducible expression of
the transgenic transcription factors (Fig. 1c).

The canonical microglia markers GPR34 and IBA1 were expressed in the iTF-Microglia at Day
8 of differentiation (Fig. 1d). To confirm the cell type identity of iTF-Microglia, we conducted
RNA-Seq and compared transcript levels of iPSC markers and microglia markers in Day-9 and
Day-15 iTF-iMicroglia to the parental iPSCs (Fig. 1e, Supplementary Table 1). As expected, the
expression of iPSC markers was drastically reduced in iTF-Microglia at Day 9 and Day 15,
whereas microglia markers were induced. Some markers, such as *P2RY12*, *CSF1R*, *CYBB* and
CD14 slightly increased their expression from Day 9 to Day 15, indicating further incremental
maturation from Day 9 to Day 15. While the transcriptomic signature of our microglia was
distinct from primary human microglia (Fig. 1f, Extended Data Fig. 1c), it was comparable to
that of several other iPSC-derived microglia protocols.

In conclusion, our transcriptomic and immunofluorescence results indicate robust expression of
microglia markers in iTF-Microglia. Importantly, our novel differentiation strategy is compatible
with large-scale pooled sgRNA screens, whereas classical protocols create population
bottlenecks (Extended Data Fig. 1d).

Functional characterization of iTF-Microglia

Next, we asked whether iTF-Microglia recapitulated cellular functions of human microglia.
Microglia are the professional phagocytes in the brain, enabling them to clear neuronal debris,
prune synapses and engulf pathogens. Using live-cell imaging and flow cytometry, we found that
iTF-Microglia robustly phagocytose fluorescent beads (Extended Data Fig. 2a) and rat
synaptosomes labeled with the pH-sensitive fluorescent dye pHrodo (Fig. 2a, b, Extended Data
Fig. 2b). As expected, phagocytosis could be attenuated by the actin polymerization inhibitor
Cytochalasin D, since phagocytosis depends on actin dynamics (Fig. 2a, b, Extended Data Fig.
2c).

180
Microglia express pattern recognition receptors such as Toll-like receptors that mediate the
inflammatory response to pathogen-associated patterns including bacterial-derived
lipopolysaccharide (LPS). To test the inflammatory response of iTF-Microglia, we stimulated
them with LPS for 24 h and evaluated morphological changes after staining for F-actin. LPS-
stimulated iTF-Microglia were less ramified, and instead displayed the ameboid morphology
characteristic of activated microglia (Fig. 2c, Extended Data Fig. 2d). In addition to the observed
morphological changes, we examined transcriptomic alterations after LPS challenge by RNA-
Seq (Fig. 2d, Supplementary Table 2). As anticipated, many of the highly upregulated genes
were immune response genes such as *C3*, *CXCL10*, *IL32* and *SAA1*. Moreover, several
upregulated genes were members of the NF-κB pathway. Downregulated genes included
TREM2, markers of homeostatic microglia, such as *P2RY13*, and members of the TGF-β
signaling pathway, such as *SLC40A1*. Transcriptomic changes in response to LPS were
substantially overlapping with those observed in iPSC-derived microglia we generated following
an alternative, previously published[22] protocol (Extended Data Fig.2e, Supplementary Table 2).
195
To examine cytokine secretion of iTF-Microglia, we measured the abundance of 36 cytokines
secreted in standard culture conditions or following LPS stimulation. Control buffer-treated iTF-
Microglia secreted most cytokines at low levels, but higher levels of CCL2 and CXCL1,
suggesting the presence of activated cells under control conditions (Fig. 2e), consistent with
previous reports suggesting that even primary microglia become partially activated when
cultured[32]. When stimulated with LPS, levels of most secreted cytokines increased; the most
increased cytokine levels where IL-6 with a 14-fold increase and IL-8 and CXCL10, both
increased over 4-fold (Fig. 2e).
204
During human development, microglia precursors enter the developing brain and mature together
with neurons into fully functional microglia. To test if neurons can promote iTF-Microglia
maturation, we co-cultured iTF-Microglia with iPSC-derived glutamatergic neurons (iNeurons)
in medium optimized for survival and functionality of both cell types (see Methods for details).
Day 8 iTF-Microglia expressing GFP differentiated in mono-culture were co-cultured with
iNeurons for one week. Remarkably, the co-cultured iTF-Microglia displayed a pronounced
ramified morphology (Fig. 2f). In conclusion, we show that iTF-Microglia effectively
phagocytose synaptosomes, respond to LPS and can be co-cultured with iPSC-derived neurons.
213
214
215 **Durable gene knockdown and overexpression by CRISPRi and CRISPRa in iTF-Microglia**
216
Next, we established CRISPRi and CRISPRa in iTF-Microglia to enable robust knockdown and
overexpression of endogenous genes, as well as large-scale loss- and gain-of-function genetic
screens. Following the strategy, we previously established in human iPSC-derived neurons[15, 16],
we stably integrated constitutive CRISPRi machinery, inducible CRISPRi machinery, or
inducible CRISPRa machinery into safe-harbor loci of iPSCs also engineered with the inducible
microglial transcription factors (Fig. 3a). We confirmed a normal karyotype for the resulting
monoclonal cell lines (Extended Data Fig. 3). Inducible CRISPRi/a systems enable flexible
timing of the onset of gene perturbation in cells already expressing sgRNAs. This feature is
particularly important for experiments in microglia: it enables lentiviral delivery of sgRNAs to

occur in iPSCs, which are much more amenable to lentiviral infection than microglia, without
prematurely affecting genes that may be relevant for differentiation. In the constitutive CRISPRi
line, the expression cassette contains a CAG promotor-driven dCas9-BFP-KRAB. In the
inducible CRISPRi cassette, this CRISPRi machinery is flanked on both the N and the C termini
with dihydrofolate reductase (DHFR) degrons. In the absence of the small molecule
trimethoprim (TMP), DHFR degrons cause proteasomal degradation of fused proteins. Addition
of TMP stabilizes the degron-tagged CRISPRi machinery. The inducible CRISPRa machinery
consists of a DHFR-dCas9-VPH construct, which is similarly stabilized in the presence of TMP.
234
To validate CRISPRi activity, we transduced iPSCs with a lentiviral construct expressing a
sgRNA targeting the transferrin receptor gene (*TFRC*) or a non-targeting control (NTC) sgRNA.
In cells expressing the constitutive CRISPRi machinery, knockdown of *TFRC* was robust in
iPSCs and iTF-Microglia both on the protein level (Fig. 3b, Extended Data Fig. 4a) and mRNA
level (Extended Data Fig. 4c,e). In cells expressing the inducible CRISPRi machinery, TFRC
knockdown was completely dependent on the presence of TMP, and effective on the mRNA and
protein levels, albeit with reduced knockdown compared to the constitutive CRISPRi system
(Fig. 3c,d, Extended Data Fig. 4 b,d,f). For additional target genes we tested, we found examples
of excellent around 80% knockdown with both the constitutive and the inducible system for
INPP5D (Extended Data Fig. 4g,h), but also an example of a gene (*PICALM*) that was
effectively knocked down by 90% with the constitutive CRISPRi (Extended Data Fig. 4i), but
not by inducible CRISPRi (Extended Data Fig. 4j). Despite these limitations of our current
inducible CRISPRi system, we decided to use it for the studies presented in this paper, since it
enabled us to induce CRISPRi knockdown only upon differentiation, rather than in the iPSC
state, thus reducing the likelihood of recovering phenotypes due to effects in iPSCs or on
differentiation itself.
251
252 Next, we validated the functionality of the inducible CRISPRa machinery by testing the
253 induction of the endogenous gene *CXCR4*. We observed a robust and tightly inducible increase
254 of *CXCR4* levels in iPSCs and iTF-Microglia on the mRNA level (Extended Data Fig. 4l,m) and
255 the protein level (Fig. 3e, Extended Data Fig. 4k).
256
257
258 **Identification of modifiers of microglial survival/proliferation by CRISPRi screens.**
259
Our first application of the inducible CRISPRi iTF-Microglia platform was to identify modifiers
of microglia survival and proliferation in a pooled genetic screen (Fig. 4a). First, we transduced
the iPSCs with our next-generation lentiviral CRISPRi sgRNA library targeting the "druggable
genome"[33]. This library consists of sgRNAs targeting 2,325 genes encoding kinases,
phosphatases, and other classes of druggable proteins with five sgRNAs per gene and 500 non-
targeting control sgRNAs. After library transduction, the iPSCs were differentiated into iTF-
Microglia by addition of doxycyline and TMP was added to induce CRISPRi activity. iTF-
Microglia were collected before differentiation (Day 0) and on Day 15 post-induction.
Frequencies of cells expressing each sgRNA were determined by next-generation sequencing
(Fig. 4a, Supplementary Table 3).
270

We compared the results from the iTF-Microglia survival screen to our previously published[15]
CRISPRi survival screen in iPSC-derived neurons (Fig. 4b) and iPSCs (Extended Fig. 5a). We
found that genes affecting microglial survival, neuronal survival and iPSC survival were largely
distinct. Knockdown of cholesterol biogenesis enzymes and V-ATPase subunits drastically
reduced neuronal but not microglial survival (Fig. 4b). Conversely, knockdown of members of
the colony stimulating factor (CSF) receptor family (*CSF1R*, *CSF2RB*, *CSF2RA*) strongly
reduced survival of microglia but not neurons (Fig. 4b) or iPSCs (Extended Fig. 5a), consistent
with their role in the development and survival of microglia and macrophages[34-37]. We validated
CSF1R essentiality in a time-course experiment (Fig. 4c). The toxicity of CSF1R knockdown
became pronounced only in differentiated iTF-Microglia (from Day 8 onwards), consistent with
the microglia-specific role of CSF1R.
282
Interestingly, the knockdown of several genes, including *CDK8* and *TGFBR2,* increased
abundance of iTF-Microglia in our screen (Fig. 4b). However, we found that *CDK8* and
TGFBR2 knockdown resulted in decreased levels of microglia marker IBA1 (Extended Data Fig.
5b,c), suggesting disrupted microglial differentiation. Indeed, inhibition of TGF-β signaling has
been shown to compensate for loss of *Oct4* pluripotency signaling[38] and microglia have been
shown to be absent in TGF-β1-deficient mice[39]. *CDK8* expression has been shown to correlate to
the stem cell pluripotency state and loss of CDK8 could cause iPSCs to differentiate into a non-
microglia state[40]. This disruption of the microglia differentiation does not seem specific to our
iTF-Microglia differentiation protocol, since knockdown of *CDK8* also decreased IBA1 levels in
iPSC-derived microglia we generated using a non-transcription factor-based differentiation
protocol[22] (Extended Data Fig. 5d,e).
294
To test whether *CDK8* and *TGFBR2* knockdown would also act in differentiated microglia, in
addition to their effect on differentiation, we induced their CRISPRi knockdown on Day 8
(Extended Data Fig. 5f). Induction of *CDK8* and *TGFBR2* knockdown in fully differentiated
iTF-Microglia did not result in proliferation, and in the case of *TGFBR2* knockdown even in a
very slight decrease in survival. By contrast, knockdown of *CSF1R* in Day 8 iTF-Microglia
reproduced the phenotype observed in the initial screen (Extended Data Fig. 5f).
301
302
303 **Identification of modifiers of microglial activation by CRISPRi screens.**
304
In a second screen, we aimed to identify modifiers of inflammatory activation of microglia. For
this screen, we chose cell surface levels of CD38 as a readout for microglial activation. CD38,
also known as cyclic ADP ribose hydrolase, is a plasma membrane glycoprotein of all brain
cells[41]. CD38 expression and its enzymatic activity increases after LPS and interferon-gamma
treatment in primary microglia[42]. Likewise, we observed transcript-level upregulation of CD38
in response to LPS treatment in iTF-Microglia and microglia we differentiated based on a
different protocol[22] (Extended Data Fig. 2e). Similarly, cell-surface levels of CD38 protein
increased upon LPS treatment based on flow cytometry (Extended Data Fig. 5g). CD38 plays
several roles in microglial activation, including in the secretion of proinflammatory cytokines[43]
and in activation-mediated cell death[42]. Altogether, these data suggest that CD38 is both a
marker and an important effector for the activation of microglia and is therefore a suitable
marker for a screen for inflammation modifiers.

The screen for modifiers of microglial activation was conducted as shown in Figure 4d. Briefly,
iPSCs expressing the inducible CRISPRi machinery were transduced with the pooled sgRNA
library described above. The cells were then differentiated into iTF-Microglia, stained for cell
surface CD38 using a fluorescently tagged antibody and subjected to FACS sorting into CD38[low]
and CD38[high] populations. Frequencies of cells expressing each sgRNA were identified in these
populations using next-generation sequencing (Supplementary Table 3).

This CRISPRi screen identified several genes regulating cell surface levels of CD38.
Knockdown of two transcriptional regulators, *CDK12* and *MED1*, significantly increased CD38
surface levels in the screen (Fig. 4e) and in validation experiments (Fig. 4f). CDK12 is known to
be involved not only in cell cycle progression but also in TNF[44] and noncanonical NF-κB[45]
signaling. While these previous reports may suggest a pro-inflammatory role of CDK12, our
findings suggest that the role of CDK12 may be more nuanced or context-dependent, and we
designated it for further investigation (see below). Another class of hits whose knockdown
increased CD38 levels were members of the mitochondrial Complex I (NADH:ubiquinone
oxidoreductase) NDUFA8 and NDUFS5 (Fig. 4e). Knockdown of components of this complex
have previously been shown to promote an inflammatory state in macrophages[46], validating our
findings.

Taken together, our large-scale CRISPRi screens in iTF-Microglia uncovered microglia-specific
survival modifiers and novel modulators of inflammatory activation, demonstrating the ability of
the iTF-Microglia screening platform to identify microglia-specific biology.

Modifiers of synaptosome phagocytosis by microglia

Microglial phagocytosis is central to brain homeostasis from development through aging[47].
Dysfunction in efferocytosis, the phagocytosis of dead cells, debris, protein aggregates and in
synaptic pruning, the phagocytic elimination of neuronal synapses by microglia, have been
implicated in neurodegenerative and psychiatric diseases[48-51]. To uncover regulators of
microglial phagocytosis, we conducted parallel CRISPRi and CRISPRa screens in iTF-Microglia
transduced with sgRNA libraries targeting the "druggable genome". After 1.5 hours of
incubation with pHrodo Red-labeled synaptosomes isolated from rat brains, iTF-Microglia were
sorted via FACS based on the pHrodo Red fluorescence signal (Fig. 5a), and screens were
analyzed as described for the CD38 FACS-based screen.

There was little overlap between CRISPRi and CRISPRa hits (Extended Data Fig. 6a,
Supplementary Table 3), confirming our previous findings from screens in diverse biological
contexts that overexpression and knockdown screens can provide complementary insights[16, 52, 53].
The underlying reasons include (i) that unlike overexpression screens, knockdown screens can
only yield phenotypes for genes expressed in the cell type under investigation, and (ii) that
knockdown of a single element of a pathway or multi-subunit complex can cause a loss-of-
function phenotype, whereas overexpression of a single element is generally not sufficient to
elicit a gain-of-function phenotype for an entire pathway or multi-subunit complex. A prominent
exception was the actin-binding protein *PFN1*, coding mutations in which cause ALS[54]. *PFN1*

had opposing phenotypes on synaptosome phagocytosis upon CRISPRi repression and CRISPRa
induction (Fig. 5b). Unexpectedly, knockdown of *CSF1R* increased phagocytosis (Fig. 5b), even
though, as we had previously found (Fig. 4b) its knockdown decreased iTF-Microglia survival.
Another remarkable hit was the Alzheimer's disease risk factor *INPP5D*, knockdown of which
slightly increased phagocytosis. Overexpression of *CD209*, a C-type lectin receptor present on
the surface of macrophages and dendritic cells, greatly increased synaptosome phagocytosis (Fig.
5b). We validated these phenotypes from the primary CRISPRi screen individually in iTF-
Microglia (Extended Data Fig. 6b) and in iPSC-derived microglia generated an alternative
protocol[22] (Extended Data Fig. 6c).
372
We further investigated the CRISPRa hits *PFN1* and *CD209*. We validated upregulation of both
genes by qPCR (Extended Data Fig. 6e). Pattern-recognition receptor CD209 has previously
been shown to regulate phagocytic capacity in macrophages[55]. We wondered if *CD209* was
additionally a substrate-specific phagocytosis regulator. To directly test the effect of *CD209* on
substrate specificity, we adapted our phagocytosis assay to simultaneously test uptake of two
separate substrates, pHRodo-Red labeled synaptosomes and yellow-green (YG) fluorescently
labeled beads (Fig. 5c). Using this approach, we challenged iTF-Microglia with either
synaptosomes or beads alone, or a mixture of both. Consistent with the screen result,
overexpression of *CD209* increased phagocytosis of synaptosomes. Interestingly, it only changed
bead phagocytosis by 10% (Fig. 5c), suggesting substrate specificity. However, when
challenging iTF-Microglia with a mixture of beads and synaptosomes, bead phagocytosis was 2-
fold increased compared to control iTF-Microglia. This finding suggests that presence of
synaptosomes might stimulate general phagocytosis via *CD209*.
386
In addition to decreased synaptosome phagocytosis, we observed increased F-actin levels in
iTF-Microglia as a consequence of *PFN1* overexpression induced by CRISPRa (Fig. 5d),
consistent with previous finding that moderate overexpression of PFN1 induces long stress fiber-
like actin cables[56]. This process could disturb orchestrated actin polymerization at the membrane
and thus decrease phagocytosis. In addition to direct effects on the actin cytoskeleton, *PFN1*
knockdown has also been reported to result in anti-inflammatory changes[57]. Indeed, we observed
transcriptional changes in immune-related genes and AD risk genes upon *PFN1* overexpression
in iTF-Microglia (Fig. 5e, Supplementary Table 4).
395
396 In conclusion, our complementary CRISPRi and CRISPRa screens identify known as well as
397 novel phagocytosis modulators in microglia, which validated in iPSC-derived microglia
398 generated using an alternative protocol.
399
400
401 **Single-cell transcriptomics reveal distinct states of iTF-Microglia**
402
Several genes had CRISPRi phenotypes in more than one of the large-scale screens that we
conducted (Extended Data Fig. 6f, Supplementary Table 3). We therefore hypothesized that
some hit genes were not dedicated factors required for specific microglial processes, but rather
more global regulators of distinct functional states. To test this hypothesis and gain more detailed
insights into the mechanisms by which genes affect microglial functions, we selected 39 hit
genes of interest, most of which had phenotypes in more than one of the large-scale primary

screens (Extended Data Fig. 6f, Supplementary Table 3) for characterization in a CROP-seq
screen, which couples CRISPRi perturbation to single-cell RNA sequencing. We introduced a
library of 81 sgRNAs (two sgRNAs targeting each selected gene (only a single sgRNA was
included for *DBF4* due to an error) and four non-targeting control sgRNAs; Supplementary
Table 5) into iPSCs, induced iTF-Microglia differentiation and CRISPRi activity, and performed
single-cell RNA sequencing of 58,302 iTF-Microglia on Day 8 (Fig. 6a, Supplementary Table
6).
416
Unsupervised clustering and Uniform Manifold Approximation and Projection (UMAP)
dimensional reduction of the single-cell transcriptomes uncovered distinct clusters (Fig. 6b). In
one cluster, a high proportion of transcripts mapped to mitochondrial transcripts, suggesting
damaged or dying cells; this cluster was removed from downstream analysis (Extended Data Fig.
7a). Two clusters exclusively contained cells expressing sgRNAs targeting *CDK8* or *TGFBR2*
(Fig. 6b and Extended Data Fig. 7b). These cells expressed high levels of the pluripotency
marker *SOX2*, but low levels of the microglia marker *CSF1R* (Extended Data Fig. 7c). Together
with our previous experiments showing reduced IBA1 levels for iTF-Microglia targeting *CDK8*
and *TGFBR2* (Extended Data Fig. 5b,c), these findings suggest that microglia differentiation was
disrupted in those cells. We removed those clusters from further analysis and retained the
remaining cluster, which was characterized by high levels of *CSF1R* expression (Extended Data
Fig. 7c). Importantly, 92.4% of cells expressing NTC sgRNAs were part of the cluster with high
levels of *CSF1R* expression, confirming the high efficiency of our microglial differentiation
protocol for unperturbed cells.
431
Unsupervised clustering and UMAP dimensional reduction of the remaining 19,834 iTF-
Microglia revealed 9 transcriptionally distinct clusters (Fig. 6c, Extended Data Fig. 7d).
Microglia heterogeneity in response to different environmental conditions in the brain has been
extensively studied[58], but we were surprised to observe a wealth of distinct transcriptional states
in the cultured iTF-Microglia. Importantly, NTC sgRNAs are represented in cells in every
cluster, suggesting the observed heterogeneity is an innate quality of the iTF-Microglia
(Extended Data Fig. 7d).
439
Principal component analysis identified two major biological axes broadly defining these states.
The first principal component (PC-1) corresponded to a polarized axis of inflammatory
activation: starting from a central homeostatic state (cluster 6), one direction was defined by
interferon-induced gene expression, whereas the other direction was defined by induction of
chemokines (Fig. 6d and Extended Data Fig. 7e). The second principal component (PC-2)
captures markers of proliferation, mainly in cluster 7 (Fig. 6e and Extended Data Fig. 7f).
446
To further interpret each transcriptional microglia state, we performed differential gene
expression analysis across the clusters (Supplementary Table 7) and named each cluster
according to characteristic transcriptomic signatures. Figure 6f highlights the top three genes
selectively expressed by cells in each cluster. Clusters 1 and 2 are both defined by high
expression of interferon-induced genes and the complement gene *C3*. Cluster 1 is uniquely
defined by high expression of chemokine *CXCL10.* Subsets of microglia characterized by
upregulation of interferon response genes have been described in mouse models of
neurodegeneration[8].

Cluster 3 is defined by the high expression of *SPP1* (Fig. 6g), which encodes secreted
phosphoprotein 1, also known as osteopontin, a multifunctional protein acting both as part of the
extracellular matrix and as a secreted cytokine[59]. Importantly, SPP1 is upregulated in several
disease-associated microglia states in both human patients and mouse disease models. These
include disease-associated microglia (DAM)[9] and activated response microglia (ARM)[60] in AD
mouse models, and late-response microglia in CK-p25 mouse models of neurodegeneration[8].
SPP1-positive microglia states are also enriched in multiple sclerosis (MS) patients and mouse
models[12] and enriched in microglia in the aging human brain[13]. Furthermore, SPP1 is highly
expressed in glioma-associated microglia in mice and humans, where high expression of SPP1 is
associated with poor prognosis[61]. Using flow cytometry, we found that SPP1-positive microglia
have a slightly increased phagocytic activity, whereas CCL13-positive microglia have
substantially decreased phagocytic activity (Fig. 6h). Integration of our iTF-Microglia dataset
with a recent scRNA-seq dataset containing 16,242 human microglia from control and
Alzheimer's Disease patient brains[7] showed conservation of the SPP1-positive microglia state
(Fig. 6i). The proportion of SPP1-positive microglia was substantially increased in Alzheimer's
disease patients compared to controls (Fig. 6j). Notably, it remains to be determined how the
SPP1+ microglia state affects the pathogenesis of different diseases, since SPP1 has been linked
to both pro-inflammatory and anti-inflammatory responses[59]. This question has been challenging
to address since we have lacked tools to manipulate the SPP1+ state of microglia *in vivo*.

Cluster 4 is defined by expression of pro-apoptotic p53 signaling genes, and Cluster 5 by
expression of metallothionines. Cluster 6 is defined by the absence of interferon response genes
or chemokines, and thus we interpreted it as representing homeostatic microglia. Cluster 7 is
characterized by the expression of proliferation markers such as *TOP2A* and *MKI67*. Cluster 8
and 9 are characterized by the expression of high levels of chemokines such as *CCL2* and *CCL3*.
Cluster 9 is uniquely defined by high expression of *CCL13* (Fig. 6g). Such chemokine signatures
have recently been found to be a hallmark of human microglia not observed in mice[62].

Taken together, single-cell RNA sequencing revealed that many important features of microglia
diversity observed in human brains and in disease states are recapitulated in our iTF-Microglia *in*
vitro model.

CROP-seq uncovers regulators of microglial cell states

We next identified the differentially expressed genes (DEGs) caused by CRISPRi knockdown of
each gene targeted in the CROP-Seq screen (Extended Data Fig. 8, Supplementary Table 8). As
expected, knockdown of functionally related genes shared common DEG signatures. For
example, knockdown of *CSF1R*, *CSF2RA* and *CSF2RB* resulted in an upregulation of genes
encoding the major histocompatibility complex as well as *CD36*, *CD74* and *CD68* (Extended
Data Fig. 8, Supplementary Table 8), which are markers of phagocytic microglia and could
possibly explain the increased phagocytic capacity we observed in response to *CSF1R*
knockdown (Fig. 5b, Extended Data Fig. 6b).

Given the surprising heterogeneity of iTF-Microglia, we investigated if CRISPRi knockdown of specific genes could control microglial cell states. Indeed, cells containing sgRNAs targeting genes such as *CSF1R*, *CDK12* and *MAPK14* were enriched or depleted from specific clusters (Fig. 7A), and more generally, knockdown of many genes specifically affected the frequency of cell states (Fig. 7b, Extended Data Fig. 9a, Supplementary Table 9).

Knockdown of *CDK12* shifted cells into cluster 9 (CCL13+, chemokine) (Fig. 7a-c). To validate this phenotype, we used a flow-cytometry approach in which secretion of CCL13 was inhibited with the transport inhibitor GolgiPlug. CCL13 levels in were increased over 2-fold with knockdown of *CDK12* (Fig. 7d), confirming our screen results. Next, we asked if the shift into cluster 9 might also have functional consequences for the iTF-Microglia. We measured synaptosome phagocytosis in CCL13^{high} cells (representative of the cells in cluster 9) and CCL13^{low} cells (representative of other clusters) in *CDK12* knockdown iTF-Microglia. As observed already in our phagocytosis screen (Fig. 5b), knockdown of *CDK12* decreased synaptosome phagocytosis, both in the CCL13^{low} and CCL13^{high} population (Fig. 7e). *CDK12* knockdown caused transcriptional downregulation of phosphatidylserine recognition receptors in both CCL13^{low} and CCL13^{high} cells, which may contribute to the decreased phagocytic activity (Extended Data Fig. 9b). Phagocytosis was even further decreased in the CCL13^{high} population (Fig. 7e), suggesting that microglia in the CCL13+ state have lower phagocytic capacity, as we showed previously (Fig. 6h). Knockdown of *CDK12* also had some cluster-specific effects (Extended Data Fig. 9b), highlighting the complex effects of gene perturbation in both shifting occupancy of cells between defined functional states, but also affecting cellular pathways in both general and state-specific ways. Interestingly, *MED1* had very similar knockdown phenotypes to *CDK12*, and both genes encode factors associated with general transcription by RNA polymerase II.

We next turned our attention to regulators of the disease-relevant SPP1-positive cluster 3. Knockdown of *MAPK14* and *CSF1R* had dramatically opposing effects, increasing and decreasing occupancy in the SPP1 cluster, respectively (Fig. 7a,b,f). Using GolgiPlug treatment to block secretion of SPP1, we validated these phenotypes by flow cytometry: knockdown of *MAPK14* increased the population of SPP1+ cells more than 6-fold (Fig. 7g), whereas CSF1R knockdown greatly diminished the proportion of SPP1+ cells (Fig. 7h).

Based on these results of genetic perturbations, we asked if pharmacological targeting of the same hits would similarly modulate the abundance of the SPP1+ state. Indeed, inhibition of MAPK14 with Skepinon-L increased the fraction of SPP1+ microglia in a time-dependent manner at nontoxic concentrations (Fig 7i,j)

Given that pharmacological inhibition of CSF1R has shown beneficial effects in several neurodegenerative mouse models, and was observed by us and others to selectively affect subpopulations of microglia in mice[63-65], we tested if pharmacological inhibition of CSF1R would reduce the proportion of SPP1+ microglia. While the CSF1R inhibitor PLX3397 showed dose-dependent toxicity in iTF-Microglia (Fig. 7k), low concentrations of CSF1R inhibitor that were nontoxic or low toxic to bulk iTF-Microglia selectively depleted SPP1+ iTF-Microglia (Fig. 7l). Thus, both pharmacological and genetic inhibition of CSF1R can decrease the proportion of SPP1+ cells.

546
In conclusion, our CROP-seq screen enabled deep characterization of the hit genes that our
primary screens identified and revealed the existence of a wealth of microglia cell states and
their regulators. To enable the scientific community to further explore this large dataset, we
implemented additional functionality in the CRISPRbrain data commons
(https://www.crisprbrain.org/) we previously described[16]. Specifically, interactive three-
dimensional UMAP representations and heatmaps enable the selective investigation of cells by
expression levels of genes of interest, sgRNA identity, and cluster membership.
554
555
556 **DISCUSSION**
557
In this study, we described a novel platform for large-scale, multimodal CRISPRi/a-based
genetic screens in human iPSC-derived microglia. We demonstrated the power of this platform
in multiple large-scale screens. While CRISPR knockout strategies are commonly used for loss-
of-function screens, the partial knockdown achieved by CRISPRi enables a more nuanced
characterization of the function of essential genes. For example, we uncovered a selective
vulnerability of microglia in the SPP1+ state to partial knockdown of the microglia-essential
gene *CSF1R*. The use of human microglia (as opposed to mouse primary microglia) enabled us
to recapitulate microglia features found in human but not mouse brain, such as a state
characterized by a chemokine signature[62]. iPSC technology will also make it possible to conduct
CRISPRi/a screens in patient-derived cells to identify modifiers of phenotypes linked to genetic
risk variants[14].
569
570 Notwithstanding, there are several areas for future optimization of our iTF-Microglia platform.
571 Improved inducible CRISPRi/a machinery with more potent gene repression and activation in
572 fully differentiated iTF-Microglia would enable the induction of CRISPRi/a at later stages during
573 differentiation to avoid false-positive hits that affect microglial differentiation, such as *CDK8*
574 and *TGFBR2* (Fig. 4b, Extended Data Fig. 5b-f).
575
Another goal for future technology development is further acceleration and enhancement of the
microglial maturation. One potential concern about sustained expression of transgenic
transcription factors is that this could promote certain microglial states over others. A protocol in
which transcription factor expression is discontinued after day 8 (Extended Data Fig. 1a) can
mitigate this concern. As with all currently available *in vitro* culture systems, microglia are
slightly activated in monoculture and lose their unique homeostatic brain signature[32]. Previous
research has shown that iPSC-microglia become more homeostatic in co-culture with neurons[21],
which is compatible with our own observation of enhanced ramification of iTF-Microglia in
neuronal co-culture (Fig. 2f). Alternatively, optimizing the set of transcription factors used to
generate iTF-Microglia may result in improved abundance of homeostatic microglia. CRISPRa
screens in our current platform are a scalable strategy to identify additional transcription factors
to promote microglial maturation and homeostasis, leading to ever more faithful models of
human microglia.
589
While there is room for future improvements, our current platform already uncovered new
insights into microglial biology. We identified several genes associated with neurodegenerative

diseases, including *PFN1* and *INPP5D*, as modulators of phagocytosis in microglia (Fig. 5), thus
pointing to a possible cellular mechanism by which variants in these genes contribute to disease.
Coding mutations in profilin 1 (*PFN1*) gene cause amyotrophic lateral sclerosis (ALS)[54]. *PFN1*
is a small actin-binding protein that promotes formin-based actin polymerization and regulates
numerous cellular functions, but how mutations in *PFN1* cause ALS is unclear. The actin
cytoskeleton is known to be important for the physiological functions of microglia, including
migration and phagocytosis. We observed that *PFN1* overexpression disrupts the actin
cytoskeleton in iTF-Microglia with higher levels of F-actin. Recently, a study has shown that
PFN1 is also involved in microglia activation, since knockdown of *PFN1* inhibited M1
proinflammatory microglial polarization and promoted anti-inflammatory M2 microglia
polarization after oxygen and glucose deprivation[57]. Introducing the ALS-associated mutations in
the *PFN1* gene in iPSCs will shed light on the impact of these specific mutations on the function
of different relevant cell types, such as iPSC-derived neurons and microglia.
605
Genetic variants in the *INPP5D* locus are associated with an increased susceptibility to AD[66] and
cerebrovascular function as well as tau and Aβ levels in the cerebrospinal fluid of AD patients[67].
INPP5D encodes the lipid phosphatase SHIP1, which is selectively expressed in brain microglia.
SHIP1 inhibits signal transduction initiated by activation of immune cell surface receptors, such
as TREM2[68]. Intriguingly, *INPP5D* expression increases with AD progression, predominantly in
plaque-associated microglia, and correlates with plaque density[69]. Given the results from our
phagocytosis screen, *INPP5D* overexpression might result in microglia with deficient phagocytic
capacity, resulting in increased Aβ deposition and neurodegeneration. Concordant with the
findings from our genetic screen, a recent study found that pharmacological SHIP1/2 inhibitors
promote microglial phagocytosis *in vitro* and *in vivo*[70].
616
617 Our single-cell RNA sequencing screen revealed that iTF-Microglia adopt a spectrum of states,
618 including states mirroring those observed in human brains, such as the SPP1-positive state. Even
619 though our protocol uses overexpression of transcription factors to generate microglia-like cells,
620 our combination of six transcription factors does not specify a single state, but recapitulates the
621 intrinsic plasticity of cell states that is a hallmark of microglia.
622
Our CROP-Seq screen identified genes controlling the distribution of iTF-Microglia across
distinct cell states. Knockdown or pharmacological inhibition of *MAPK14* strongly promoted
adoption of the disease-associated SPP1-positive state. Previous work suggested a functional
connection between SPP1 and MAPK14 in cancer cells, where SPP1 can activate the p38 MAPK
signaling pathway, which comprises MAPK14[71]. MAPK14 was also recently predicted to be a
unique network regulator in DAM[72]. However, our identification of MAPK14 as a regulator of
the SPP1+ state is novel and enhances our understanding of modulators of microglia cell states.
630
We found that the SPP1-positive microglia state can be selectively depleted by genetic and
pharmacological inhibition of CSF1R. CSF1R inhibitors have beneficial effects in mouse models
of diseases including AD[73, 74], tauopathy[65] and MS[75]. Intriguingly, CSF1R inhibition reduced
SPP1 expression in the MS model, while homeostatic genes such as *TMEM119* and *P2RY12*
were increased[75], paralleling our finding that the SPP1 microglia state is selectively vulnerable to
CSF1R inhibition. Additionally, disruption of CSF1-CSF1R signaling downregulated SPP1 in
the cerebellum[76]. Combining CSF1R depletion and single cell profiling has enabled us

previously to elucidate the differential effects of CSF1R inhibitors on microglia subtypes[63].
Following CSF1R inhibition, we found an enrichment of microglia states with elevated markers
of inflammatory chemokines and proliferation and interestingly, in concordance with our
findings in iTF-Microglia here, an upregulation of cell surface receptor CD74[63]. Others have
reported compensatory upregulation of TREM2/β-catenin and IL-34 in microglia following
conditional CSF1R KO[77]; however, we did not find consistent upregulation of these factors in
our iTF-microglia (Supplementary Table 9). Based on our new finding that CSF1R inhibition at
low doses that are nontoxic to most microglia selectively depletes the SPP1+ population in iTF-
Microglia, low-dose CSF1R inhibition might also give us a tool to study the SPP1+ population in
mouse disease models.
648
649   Taken together, our results have provided pharmacological strategies to either promote or deplete
650   SPP1+ microglia. This will make it possible to determine the role of SPP1+ microglia in
651   different diseases, where they may play either beneficial or detrimental roles, and to manipulate
652   this disease-associated microglial state for therapeutic benefit.
653
We anticipate that the screening platform we describe here can be broadly applied to screen for
other microglia-related phenotypes, and to systematically identify regulators of different
microglia states. Using iPSCs derived from patients with familial or sporadic diseases will enable
the identification of potential therapeutic targets that can correct cellular phenotypes. For
example, the APOE ε4 variant reduces the ability of iPSC-microglia to clear Aβ[78] and
dysregulates cholesterol biogenesis[79], and CRISPR screens have the potential to elucidate the
underlying molecular mechanisms and uncover potential remedies. Introduction of microglia into
co-cultures or brain organoids can provide a screening platform to investigate their interactions
with other brain cell types, such as synaptic pruning of neurons. Finally, transplantation of iTF-
Microglia into postnatal, immune-deficient, humanized mice could result in microglia with an *ex*
vivo human microglial gene signature, including more homeostatic microglia, and enable the
investigation of factors controlling the interaction of microglia with a model for diseased brain
environment[80-83].
667
668

ACKNOWLEDGEMENTS
We thank Ruilin Tian for discussions about CROP-Seq, Chris Bohlen for advice on synaptosome
protocols, Jessica Mella, Karinna Vivanco, Capria Rinaldi, Gabriel Sturm, Yaqiao Li, Carlo
Condello, and Miranda Sullivan for contributions to preliminary studies. We thank Avi Samelson
for feedback on the manuscript and members of the Kampmann lab for discussions.
This research was supported by NIH grants DP2 GM119139 and U01 MH115747 to MK, U54
NS100717 to LG and MK, R01 AG051390 to LG, F30 AG066418 to KL, F30 AG062043 to LK,
and funded by the Center for Alzheimer's and Related Dementias (CARD) of the National
Institutes of Health (NIH) under Award Number ZO1 AG000534-02, and funded, in part,
through the Intramural Research Program of the National Institutes of Neurological Disorders
and Stroke (MEW). The research was also supported by an NSF Graduate Research Fellowship
to OT, Tau Consortium Investigator Awards (Rainwater Charitable Foundation) to JKI, LG and
MK, a Chan Zuckerberg Initiative Ben Barres Early Career Acceleration Award to MK. MK. is a
Chan Zuckerberg Biohub Investigator. Development of new features for CRISPRbrain was
supported, in part, by a collaboration among the Kampmann Lab, UCSF and Data Tecnica

International, LLC and funding from the Center for Alzheimer's and Related Dementias,
National Institutes of Health.
686
687
688
AUTHOR CONTRIBUTIONS
The iTF-Microglia differentiation strategy was developed and characterized by CH and LG with
contributions from CC, LZ, JCO, LK, JI and MW. Additionally, ND, SS, OT, KL, JH and MK
contributed to the optimization and characterization of iTF-Microglia. CRISPR-based functional
genomics studies were designed, conducted and analyzed by ND, SS and MK with contributions
from OT, KL, GA and JH. SS led the computational analysis of all screens and RNA-Seq
experiments. SHH and FF developed new features for the CRISPRbrain data commons with
critical input from SS, ND and MK and feedback from MAN and ABS. ND, SS, CH, OT and
MK created the Figures. ND, SS and MK conceptualized and wrote the manuscript with input
from the other authors. All authors reviewed and approved the final manuscript.
699
COMPETING INTEREST STATEMENT
MN consults for Neuron23. MN and FF participated in this work in part due to a competitively
awarded consulting contract between Data Tecnica International LLC and the National Institutes
of Health (USA). JKI is a cofounder of AcuraStem, Inc. and Modulo Bio, and serves on the
scientific advisory board of Spinogenix. LG is a founder of Aeton Therapeutics. MK has filed a
patent application related to CRISPRi and CRISPRa screening (PCT/US15/40449), serves on the
scientific advisory boards of Engine Biosciences, Casma Therapeutics, and Cajal Neuroscience
and is a consultant to Modulo Bio. The remaining authors declare no competing interests.
708
709 **METHODS**
710
Human iPSC culture. Human iPSCs (male WTC11 background, PMID 24509632) were
cultured in StemFlex™ Basal Medium (Gibco; Cat. No. A33493-01) on BioLite Cell Culture
Treated Dishes (Thermo Fisher Scientific; assorted Cat. No.) coated with Growth Factor
Reduced, Phenol Red-Free, LDEV-Free Matrigel Basement Membrane Matrix (Corning; Cat.
No. 356231) diluted 1:100 in Knockout DMEM (GIBCO/Thermo Fisher Scientific; Cat. No.
10829-018). StemFlex was replaced every other day or every day once 50% confluent. When
70%-80% confluent, cells were passaged by aspirating media, washing with DPBS (Gibco; Cat.
No. 14190-144), incubating with StemPro Accutase Cell Dissociation Reagent (GIBCO/Thermo
Fisher Scientific; Cat. No. A11105-01) at 37°C for 7 min, diluting Accutase 1:5 in StemFlex,
collecting cells in conicals, centrifuging at 220g for 5 min, aspirating supernatant, resuspending
cell pellet in StemFlex supplemented with 10 nM Y-27632 dihydrochloride ROCK inhibitor
(Tocris; Cat. No. 125410), counting, and plating onto matrigel-coated plates at the desired
number. Human iPSCs studies at the University of California, San Francisco (UCSF) were
approved by the Human Gamete, Embryo and Stem Cell Research Committee.
725
Human CRISPR iTF-iPS cell line generation. The two donor plasmids for inducible
expression of six codon-optimized transcription factors were constructed using the plasmid
pUCM (GENEWIZ). Human iPSCs (WTC11), acquired from Dr. Bruce Conklin (Gladstone
Institute, San Francisco), were engineered to express PU.1, CEBPβ, and IRF5 under a

doxycyline-inducible system in the CLYBL safe harbor locus and MAFB, CEBPα, and IRF8 in
the AAVS1 safe harbor locus using TALEN-based editing as previously described[15]. Clones
were selected using both neomycin and puromycin, thus generating the cell line we termed iTF-
iPSCs. Next, iTF-iPSCs were transfected with either pC13N-dCas9-BFP-KRAB[15], pRT029-
CLYBL-CAG-DHFR-dCas9-BFP-KRAB-NLS-DHFR[15], or pRT043-CLYBL-DDdCas9VPH-
GFP[16] to generate constitutive CRISPRi, inducible CRISPRi, or inducible CRISPRa iTF-iPS cell
lines, respectively, in the CLYBL safe harbor locus using the same TALEN-editing method.
After transfection, BFP-positive (CRISPRi) or GFP-positive (CRISPRa) iTF-iPSCs were
repeatedly enriched via FACS sorting (BD FACS Aria Fusion).
To generate monoclonal cell lines, 5,000 polyclonal CRISPR-iTF-iPSCs were plated on 10-cm
dishes to enable isolation of individual clones under direct visualization with an inverted
microscope (Evos FL, Thermo Fisher Scientific) in a tissue culture hood via manual scraping.
Monoclonal cell lines were tested for iTF-Microglia differentiation capability and CRISPRi/a
activity.
744
Human iPSC-derived iTF-Microglia cell culture and differentiation. iTF-iPSCs were grown
in StemFlex until reaching at least 50% confluency and were grown for at least 24h without
ROCK inhibitor. They were dissociated and centrifuged as described above and pelleted cells
were resuspended in Day 0 differentiation medium containing the following: Essential 8™ Basal
Medium (Gibco; Cat. No. A15169-01) as a base, 10 nM ROCK inhibitor, and 2 μg/ml
Doxycycline (Clontech; Cat. No. 631311). iTF-iPSCs were counted and seeded onto double
coated plates (Poly-D-Lysine-precoated Bio plates (Corning, assorted Cat. No.) + Matrigel
coating) with the following seeding densities: 10,000 cells/well for 96-well plate, 0.1
million/well for 12-well plate, 0.15 million/well for 6-well plate, 2 million/dish for 10-cm dish,
and 8 million/dish for 15-cm dish. On day 2, media was replaced with Day 2 differentiation
media containing Advanced DMEM/F12 Medium (Gibco; Cat. No. 12634-010) as a base
medium containing the following: 1X Antibiotic-Antimycotic (Anti-Anti) (Gibco; Cat. No.
15240-062), 1X GlutaMAX™ (Gibco; Cat. No. 35050-061), 2 μg/ml doxycycline, 100 ng/ml
Human IL34 (Peprotech; Cat. No. 200-34) and 10 ng/ml Human GM-CSF (Peprotech; Cat. No.
300-03). Two days later, on day 4, the medium was replaced with iTF-Microglia medium,
containing Advanced DMEM/F12 as a base medium and the following: 1X Anti-Anti, 1X
GlutaMAX, 2 μg/ml doxycycline, 100 ng/ml Human IL-34 and 10 ng/ml Human GM-CSF, 50
ng/ml Human M-CSF (Peprotech; Cat. No. 300-25) and 50 ng/ml Human TGFB1 (Peprotech;
Cat. No. 100-21C). On Day 8, the media was replaced with fresh iTF-Microglia medium. iTF-
Microglia can be cultured for at least 12 more days in iTF-Microglia medium with full medium
changes every 3-4 days. Cells were assayed on day 8, day 9 or day 15 in most experiments.
When differentiating the inducible CRISPRi/a iTF-Microglia, the media was supplemented with
50 nM trimethoprim (MP Biomedical, LLC; Cat. No. 195527) and changed every two days to
maintain strong knockdown/overexpression.
For dissociation, iTF-Microglia were washed once with PBS before adding TrypLE Express
(Gibco; Cat. No. 12605-028) and incubating for 10 min at 37 °C. Cells were diluted 1:3 in
Advanced DMEM/F12 and spun down at 220g for 5 min before resuspending in appropriate
media.
773
Doxycycline removal assay after Day 8 of differentiation

10,000 iTF-iPSCs were seeded into 96-well Flat Clear Bottom White Polystyrene Poly-D-Lysine
Coated Microplates (Corning; Cat. No. 3843) and differentiated into iTF-Microglia as described
above. At Day 8 of the differentiation, the media of iTF-Microglia was replaced with either i)
full media change of iTF-Microglia medium containing 2 µg/ml doxycycline, or ii) full media
change of iTF-Microglia medium containing no doxycycline, or iii) half media changes of iTF-
Microglia medium containing no doxycycline. This media-replacing paradigm was repeated
every three days until Day 15. Microglia survival was assessed by performing the CellTiter Glo
2.0 (Promega; Cat. No. G9242) assay according to the manufacturer's instructions.
Luminescence signal was recorded with the M5 plate reader (SpectroMax).
784
Differentiation and culture of iPSC-derived microglia following the protocol by Brownjohn
and colleagues. Brownjohn iPSC-Microglia (Brownjohn-iMG) were differentiated from dCas9-
KRAB iPSCs (AICS-0090, Allen Cell Collection) using the published protocol[22] with minor
modifications. In brief, iPSCs (cultured in Stem Flex media with colonies at 60-80% confluency)
were dissociated to single cells with Accutase, collected and plated at 10,000 cells per well in 96-
well ultra-low attachment, round bottom plates (Corning; Cat. No. 7007) in 100 µl embryoid
body medium (10 mM ROCK inhibitor, 50 ng/mL BMP-4 (Peprotech; Cat. No. 120-05), 20
ng/mL SCF (Peprotech; Cat. No. 300-07), and 50 ng/mL VEGF (Peprotech; Cat. No. 100-20) in
E8 medium), and then subjected to centrifugation at 300g for 3 min. Embryoid bodies were
cultured for 4 days, with a half medium change after 2 days. On day 4, embryoid bodies were
carefully collected and transferred into a 15 ml conical tube, and left to settle at the bottom. The
embryoid media was aspirated and 15 to 20 embryoid bodies were plated per well in 6-well
plates and cultured in 3 mL hematopoetic medium (2 mM GlutaMax, 1x Anti-Anti, 55 mM 2-
mercaptoethanol (BioRad; Cat. No. 1610710), 100 ng/mL M-CSF, and 25 ng/mL Human IL-3
(Peprotech; Cat. No. 200-03) in X-Vivo 15 (Lonza; Cat. No. BE02-060F). Two thirds of the
media was exchanged every 3-4 days. Microglia progenitors were harvested from suspension
after 14-21 days and plated onto PDL-coated plates in microglia maturation media (2 mM
GlutaMax, 1x Anti-Anti, 100 ng/mL IL-34, and 10 ng/mL GM-CSF in Advanced RPMI-1640
(Gibco; Cat. No. 12633012)). Microglia progenitors were further differentiated for 8 days with
full medium change every 2–3 days before using them for experiments.
805
iTF-Microglia coculture with iNeurons. iPSC-derived neurons (iNeurons) were differentiated
from WTC11 iPSCs engineered to express NGN2 under a doxycycline-inducible system in the
AAVS1 safe harbor locus as previously described[15, 84] with minor modifications as follows:
iPSCs were maintained and dissociated as described above and replated on Matrigel coated
dishes in N2 Pre-Differentiation Medium. After three days, hereafter Day 0, the pre-
differentiated neurons were dissociated to single cells with Accutase, collected, and plated at
10,000 cells per well in PDL-coated 96-well plates in BrainPhys Neuronal Medium (BrainPhys
(STEMCELL Technologies; Cat. No. 05790) as the base, 0.5x N2 supplement (Thermo Fisher;
Cat. No. 17502-048), 0.5x B27 Supplement (GIBCO/Thermo Fisher Scientific; Cat. No. 17504-
044), 10ng/mL NT-3 (PeproTech; Cat. No. 450-03), 10ng/mL BDNF, 1mg/mL Mouse laminin
(Thermo Fisher; Cat. No. 23017-015), and 2mg/mL doxycycline. On Day 3, a full media change
was performed. On day 7, half the media was removed, and an equal volume of BrainPhys
Neuronal Medium was added. On day 14, half the media was removed and an equal volume of
BrainPhys Neuronal Medium containing Day 8 iTF-Microglia expressing Lck-mNeonGreen and
supplemented with 2X the cytokines of the iTF-Microglia medium was added. 3,000 iTF-

Microglia were added to each well and immunostaining experiments were performed after one day.

Lentiviral transduction of iPSCs with sgRNA constructs. Individual or pooled sgRNAs were introduced into CRISPRi- or CRISPRa-iPSCs via lentiviral delivery using TransIT Lenti Reagent (Mirus Bio LLC; Cat. No. MIR 6600) according to manufacturer's protocol. Cells were selected with 2 μg/ml Puromycin (Gibco; Cat. No. A11138-03) for 2 - 4 days and recovered 2 - 4 days until MOI >0.9 as determined by flow cytometry of sgRNA-BFP fluorescence. sgRNA protospacer sequences are provided in Supplementary Table 10.

qPCR. To quantify *TFRC, INPP5D or PICALM* knockdown or *CXCR4, CD209 or PFN1* overexpression, lysed cell pellets from human iPSCs or iTF-Microglia were thawed on ice, and total RNA was extracted using the Quick-RNA Miniprep Kit (Zymo; Cat. No. R1054). cDNA was synthesized with the SensiFAST cDNA Synthesis Kit (Bioline; Cat. No. 65054). Samples were prepared for qPCR in technical triplicates in 5 μL reaction volumes using SensiFAST SYBR Lo-ROX 2X master mix (Bioline; Cat. No. BIO-94005), custom qPCR primers from Integrated DNA Technologies used at a final concentration of 0.2 μM, and cDNA diluted at 1:3. Quantitative real-time PCR was performed on an Applied Biosystems QuantStudio 6 Pro Real-Time PCR System with the following Fast 2-Step protocol: (1) 95° C for 20 s; (2) 95° C for 5 s (denaturation); (3) 60°C for 20 s (annealing/extension); (4) repeat steps 2 and 3 for a total of 40 cycles; (5) 95°C for 1 s; (6) ramp 1.92°C/s from 60°C to 95°C to establish melting curve. Expression fold changes were calculated using the ΔΔCt method normalizing to housekeeping gene *GAPDH*. Primer sequences are provided in Supplementary Table 10.

Cell surface protein staining for flow cytometry. Dissociated and resuspended iTF-Microglia were blocked for 15 min with 1:20 Human FC Block (BD Biosciences; Cat. No. 564220) and then stained with 1:66 PE/Cy7 anti-human CD184 (CXCR4) (BioLegend; Cat. No. 306514) for CRISPRa validation or 1:66 PE-Cy7 anti-human CD71 (TFRC) (BioLegend; Cat. No. 334112) for CRISPRi validation for 30 min in the dark. For the CD38 screen and validation experiments, iTF-Microglia were stained with 1:200 anti-hCD38 PE (R&D Systems; Cat. No. FAB2404P). Cells were washed twice with DPBS before analyzing them by flow cytometry using the BD LSRFortessa X14 (BD Biosciences). Flow cytometry data was analyzed using FlowJo (FlowJo, version 10.7.1), raw median fluorescence intensity values of CD184, CD71 and CD38 stained cells were normalized to non-stained control samples and data was plotted as fold change using Prism 8 (GraphPad, version 8.4.2).

Intracellular protein staining for flow cytometry. iTF-Microglia were treated for 6 h with 1:2000 GolgiPlug™ (BD; Cat. No. 555029) or DMSO as control before dissociating. Cells were fixed and permeabilized with the eBioscience Intracellular Fixation and Permeabilization Buffer Set (Invitrogen; Cat. No. 88-8824-00) according to the manufacturer's instructions. Cells were stained with 1:75 Anti-Hu Osteopontin (SPP1) eFluor 660 (eBioscience; Cat. No. 50-9096-42) or 1:75 Human CCL13 488 (R&D Systems; Cat. No. IC327G) or their isotype controls Mouse IgG1 Control Alexa Fluor 488 conjugated (R&D Systems; Cat. No. IC002G) and Mouse IGG1 kappa Isotype (eBioscience; Cat. No. 50-4714-82) over night at 4 °C. Cells were washed twice with DPBS before analyzing them by flow cytometry using the BD FACS CelestaTM (BD Biosciences) or the BD FACSAria Fusion. Flow cytometry data was analyzed using FlowJo, raw

867 median fluorescence intensity values of Osteopontin (SPP1) and CCL13 stained cells were
868 normalized to isotype-control samples and data was plotted as fold change using Prism 8. The
869 gating strategy used to determine the percentage of SPP1-positive cells is shown in
870 Supplementary Fig. 1c).
871
Immunohistochemistry. iTF-Microglia monocultures and co-cultures were differentiated in
PDL-coated 96-well plates. They were fixed with 4% Paraformaldehyde (Electron Microscopy
Sciences; Cat. No. 15710) for 10 min at room temperature. After washing with DPBS 3 times,
cells were permeabilized and blocked with 5% normal goat serum (Vector Laboratories; Cat. No.
S-1000-20) with 0.01% Triton X-100 (TEKnova; Cat. No. T1105) in PBS for 1 hr at room
temperature. Cells were then incubated with primary antibodies diluted in blocking buffer at 4 °C
overnight. After that, cells were washed with DPBS 3 times and incubated with secondary
antibodies diluted in blocking buffer for 1 hr at room temperature. Cells were then washed with
DPBS 3 times and stained with 10 μg/ml Hoechst 33342 (Thermo Fisher Scientific; Cat. No.
H3570) for 10 min. Cells were imaged using a confocal microscope (Leica SP8) or an IN Cell
Analyzer 6000 (GE; Cat. No. 28-9938-51). Primary antibodies used for immunofluorescence in
this study were as follows: anti-mouse 1:150 GPR34 (R&D Systems; Cat. No. MAB4617), anti-
rabbit 1:1000 IBA1 (Wako; Cat. No. 019-19741), anti-rabbit 1:200 TFRC (abcam; Cat. No.
ab84036), anti-rabbit 1:1000 synaptophysin (Synaptic Systems; Cat. No. 101 004). Secondary
antibodies used in this study were as follows: goat anti-rabbit IgG Alexa Fluor 555 (1:500
dilution; abcam; Cat. No. ab150078), goat anti-mouse IgG Alexa Fluor 488 (1:500 dilution;
abcam; Cat. No. ab150113) and goat anti-chicken IgG Alexa Fluor 647. F-actin was stained
using ActinGreenTM 488 (Invitrogen; Cat. No. R37110) according to the manufacturer's
protocol.
891
Synaptosome isolation and pHrodoRed labelling. Synaptosomes were isolated from fresh
Innovative Grade US Origin Rat Sprague Dawley Brain (Innovative Research, Inc.; Cat. No.
IGRTSDBR) with the Syn-PER™ Synaptic Protein Extraction Reagent (Thermo Scientific™;
Cat. No. 87793) according to the manufacture's protocol with minor changes. Briefly, 10 mL of
Syn-PER Reagent supplemented with 1x protease inhibitor cOmplete Mini, EDTA free (Roche;
Cat. No. 11836170001) and 1x phosphatase inhibitor PhosSTOP (Roche; Cat. No. 4906845001)
were added per gram of brain tissue. Dounce homogenization was performed on ice and
homogenate was transferred to a conical tube and centrifuged at $1200 \times g$ for 10 minutes at 4°C.
The pellet was discarded, the supernatant was transferred to a new tube, and the centrifugation
step was repeated. The supernatant was then centrifuged at $15,000 \times g$ for 20 minutes at 4°C.
The supernatant was removed and the wet pellet was weighed. The synaptosome fractions were
resuspended at a concentration of 50 mg/ml. 3 μM pHrodo™ Red, succinimidyl ester (pHrodo™
Red, SE) (ThermoFisher Scientific; Cat. No. P36600) was added to the synaptosome fraction and
incubated for 45 min at room temperature in the dark. After diluting the solution 1:10 in DPBS,
the synaptosomes were spun down at $2500 \times g$ for 5 min. The supernatant was removed and then
the synaptosomes were washed two times with DPBS. The pHrodo-labeled synaptosomes were
resuspended in microglia iTF-Microglia medium at a stock concentration of 50 mg/ml and
directly used for phagocytosis assays or frozen in synaptosome freezing media (5% DMSO in
Advanced DMEM/F12) for later use.
911

Phagocytosis assays. Day 8 iTF-Microglia were used for all phagocytosis assays. iTF-Microglia medium was prepared with pHRodoRed-labeled synaptosomes at a concentration of 1 mg/ml or 0.5 μl/ml media of Fluoresbrite Carboxylate YG 1.0 Micron Microspheres (15702-10; Cat. No. 15702-10). After replacing the media with the substrate media, iTF-Microglia were incubated for 1.5 h in the incubator if not otherwise stated. Cells were washed twice with DPBS, dissociated, resuspended in ice-cold DPBS, and analyzed via flow cytometry. Where indicated, actin polymerization was inhibited by pretreating cells with 5 μM Cytochalasin D (Invitrogen; Cat. No. PHZ1063) for 30 min before the addition of phagocytic substrate media. For analyzing phagocytic capabilities within microglia clusters, pHRodoRed-labeled synaptosomes at a concentration of 1 mg/ml were added to iTF-Microglia for 1.5h. Microglia were washed 3x with PBS before incubating them in iTF-Microglia media supplemented with 1:2000 GolgiPlug™ (BD; Cat. No. 555029) for 4h. Cells were dissociated, fixed, and stained for CCL13 and SPP1 as described above. Flow cytometry data was analyzed using FlowJo, raw median fluorescence intensity values of phagocytosing cells were normalized to no-substrate control samples and data plotted as fold change using Prism 8.

Human cytokine array. Day 8 iTF-Microglia were treated with 100 ng/ml LPS (Millipore Sigma; Cat. No. LPS25) or DPBS control. After 24 hours, the supernatant was collected and processed using the Proteome Profiler Human Cytokine Array Kit (R&D Systems; Cat. No. ARY005B), according to manufacturer's instructions. For analysis of the signals, Fiji (version 2.0.0) was used to measure the integrated pixel density for each pair of duplicate dots representing a cytokine. Background signal was measured from negative control dots and then subtracted from each dot. The relative change in cytokine levels as a result of LPS-treatment was obtained by comparing corresponding cytokine signals across multiple arrays performed in tandem.

Live-cell imaging. iTF-iPSCs transduced with individual sgRNAs as described above were passaged and differentiated into iTF-Microglia in the 96-well format described above. Starting on day 2 of differentiation, and continuing every two days until day 15, iTF-Microglia were stained with 10 μg/ml Hoechst 33342 for 10 min at 37 °C, washed with PBS, and imaged with the IN Cell Analyzer 6000. Using the same 96-well format as described above, Day 8 iTF-Microglia were stained for F-actin using 25 nM SiR-actin (Cytoskeleton, Inc; Cat. No. CY-SC001) probe, diluted in iTF-Microglia medium, with a 4 hour incubation at 37 °C. A full media change with iTF-Microglia medium was completed before imaging using the IN Cell Analyzer 6000.

CellTiter Glo assay after pharmaceutical inhibition of CSF1R or MAPK14. 10,000 iTF-iPSCs were seeded into 96-well Flat Clear Bottom White Polystyrene Poly-D-Lysine Coated Microplates (Corning; Cat. No. 3843) and differentiated into iTF-Microglia. For CSF1R inhibition, cells were treated with the CSF1R inhibitor PLX3397 at Day 8 (ApexBio; Cat. No. B5854) at indicated concentrations or DMSO control for 24h before performing the CellTiter Glo 2.0 (Promega; Cat. No. G9242) assay according to the manufacturer's instructions. For MAPK14 inhibition, cells were treated with the MAPK14 inhibitor Skepinone-L at Day 8 (Selleckchem; Cat. No. 1221485831) at indicated concentrations or DMSO control for 24h before performing the CellTiter Glo 2.0 assay. Luminescence signal was recorded with the M5 plate reader (SpectroMax).

958
CRISPR SCREENS

Large-scale survival-based and FACS-based screens. Inducible CRISPRi and CRISPRa iTF-
iPSCs were infected with pooled CRISPRi or CRISPRa sgRNA libraries[33] targeting the
druggable genome and selected for lentiviral integration with puromycin, as described above.
Day 0 iTF-iPSCs, with a cell count corresponding to an average 1000x coverage per library
element, were differentiated into iTF-Microglia as described above with constant TMP
supplementation for dCas9 stabilization.
For the survival screens, Day 0 iPSCs and Day 15 iTF-Microglia were lifted with Accutase or
TryplE Express, respectively. Lifted cells were harvested and subjected to sample preparation for
next-generation sequencing as described below.
For the CD38-activation screen, Day 8 iTF-Microglia were dissociated with TripleE and then
blocked and stained with anti-PE-CD38 as described in the cell surface staining section. Cells
were sorted into high and low signal population corresponding to the top 30% and the bottom
30% of the CD38-PE signal distribution (gating strategy shown in Supplementary Fig. 1a).
For the phagocytosis FACS screen, Day 15 iTF-Microglia were incubated with PhRodo-Red
synaptosomes as described in phagocytosis assay section. Cells were then dissociated with
TryplE and sorted into high and low signal population corresponding to the top 30% and the
bottom 30% of the PhRodoRed signal distribution (gating strategy shown in Supplementary Fig.
1b). Based on simulations, we previously found that this sorting strategy is optimal for hit
detection in FACS-based screens[85].
Cells were subjected to sample preparation for next-generation sequencing as previously
described[15]. Briefly, for each screen sample, genomic DNA was isolated using a Macherey-
Nagel Blood L kit (Macherey-Nagel; Cat. No. 740954.20). sgRNA-encoding regions were
amplified and sequenced on an Illumina HiSeq-4000.

Quant-Seq. Cell culture medium was aspirated, cells were washed once with DPBS, and RNA
lysis buffer was added directly to wells containing either Day 0 iTF-iPSCs, Day 15 iTF-
Microglia +/- 50 ng/ml 24h LPS treatment, Brownjohn-iMG +/- 100 ng/ml 24h LPS treatment,
or Day 15 iTF-Microglia. For assessing transcriptomic effects after *PFN1* overexpression, two
different *PFN1* sgRNAs and NTC sgRNAs were transduced into inducible CRISPRa iPSCs and
cells were differentiated to Day 8 iTF-Microglia. Biological triplicates for each condition
(approximately 0.15 Mio cells each) were pelleted, snap frozen, and stored at -80°C. RNA was
extracted using the Quick-RNA Miniprep Kit (Zymo; Cat. No. R1055). Libraries were prepared
from total RNA (250-473 ng per sample) using the QuantSeq 3' mRNA-Seq Library Prep Kit for
Illumina (FWD) (Lexogen; Cat. No. 015UG009V0252) following the manufacturer's
instructions. Library amplification was performed with 14 total PCR cycles. mRNA-seq library
concentrations (mean of 1.13 ± 0.66 ng/uL) were measured with the Qubit dsDNA HS Assay Kit
(Invitrogen; Cat. No. Q32851) on a Qubit 2.0 Fluorometer. Library fragment-length distributions
(mean of 287 ± 28 bp) were quantified with High Sensitivity D5000 Reagents (Agilent
Technologies; Cat. No. 5067-5593) on the 4200 TapeStation System. The libraries were
sequenced on an Illumina NextSeq 2000 instrument with single-end reads.

1001

CROP-Seq. A pooled sgRNA library consisting of 2 sgRNAs per targeted gene and 4 non-
targeting control sgRNAs was designed to target 39 genes which were selected hit genes from

iTF-Microglia survival and FACS-based screens (Supplementary Table 5; only 1 sgRNA for
gene *DBF4* due to technical error). Briefly, top and bottom strands of sgRNA oligos were
synthesized (Integrated DNA Technologies) and annealed in an arrayed format, pooled in equal
amounts, and ligated into our optimized CROP-seq vector, as previously described[15].
Inducible CRISPRi-iTF-iPSCs were infected with the pooled sgRNA library at <0.15 MOI and
then selected for lentiviral integration. Next, iTF-iPSCs were differentiated into iTF-microglia
and cultured with the addition of TMP. Day 8 iTF-Microglia were washed 3X with DPBS,
dissociated with TrypLE, and resuspended in nuclease-free water before loading onto four wells
of the 10x Chromium Controller (10x Genomics, v3.1) according to the manufacturer's protocol,
with 35,000 cells recovered per sample as the target. Sample preparation was performed using
the Chromium Next GEM Single Cell 3′ Reagent Kits version 3.1 (10x Genomics, cat. no. PN-
1000121) according to the manufacturer's protocol, reserving 10-30 ng full-length cDNA to
facilitate sgRNA assignment by amplifying sgRNA-containing transcripts using hemi-nested
PCR reactions adapted from a previously published approach[15, 86]. cDNA fragment analysis was
performed using the 4200 TapeStation System and sgRNA enrichment libraries were separately
indexed and sequenced as spike-ins alongside the whole-transcriptome scRNA-seq libraries
using a NovaSeq 6000 using the following configuration: Read 1: 28; i7 index: 8; i5 index: 0;
Read 2: 91.
1022
1023 **COMPUTATIONAL AND STATISTICAL ANALYSIS**
1024
Primary CRISPR screen analysis. Primary screens were analyzed using our previously
published MAGeCK-iNC bioinformatics pipeline[15], available at
https://kampmannlab.ucsf.edu/mageck-inc. Briefly, raw sequencing reads from next-generation
sequencing were cropped and aligned to the reference using Bowtie version 0.12.9[87] to determine
sgRNA counts in each sample. The quality of each screen was assessed by plotting the
$\log_{10}$(counts) per sgRNA on a rank order plot using ggplot2 version 3.3.3[88]. Raw phenotype
scores and significance P values were calculated for target genes, as well as for 'negative-
control-quasi-genes' that were generated by random sampling with replacement of five non-
targeting control sgRNAs from all non-targeting control sgRNAs. The final phenotype score for
each gene was calculated by subtracting the raw phenotype score by the median raw phenotype
score of 'negative-control-quasi-genes' and then dividing by the standard deviation of raw
phenotype scores of 'negative-control-quasi-genes'.
A 'Gene Score' was defined as the product of phenotype score and $-\log_{10}$(P value). Hit genes
were determined based on the Gene Score cutoff corresponding to an empirical FDR of 10%.
Volcano plots of gene scores were generated using ggplot2 version 3.3.3[88].
1040
RNA-seq analysis. Alignment and mapping were performed using Salmon version 1.4.0[89], (the -
-noLengthCorrection flag was used for QuantSeq samples) and either the human reference
genome GRCh38 (Gencode, release 37), or a custom GRCh38 reference genome containing the
references for each 3TF transgene integrated in iTF-iPSCs, to obtain transcript abundance
counts. Tximport version 1.18.0[90] was used to obtain gene-level count estimates. Genes with
zero counts across all samples were removed from the analysis. To visualize differences in gene
expression across samples, a list of gene symbols corresponding to microglia markers, microglia
activation markers, and iPSC markers were compiled from a previous publication[91]; the
normalized counts of each of these genes were then standardized across samples (*i.e.* subtracting

by the mean and dividing by the standard deviation) and visualized using Complex Heatmap
version 2.6.2[92]. To assess how iTF-Microglia compare to other iPSC-Microglia and primary
microglia from a range of previous studies[17, 22, 24, 25, 93], raw fastqs obtained from the NCBI GEO
database) were subject to the same analysis pipeline stated above. Then, principal component
analysis was performed using microglia marker genes as input with DESeq2 version 1.30.1[94].
For differential gene expression analysis of LPS-treated vs. PBS-treated iTF-Microglia samples
and the *PFN1* overexpression vs. NTC iTF-Microglia DESeq2 version 1.30.1[94] was used to
calculate the log-fold-change and p-values and perform shrinkage of log-fold-change for
downstream visualization using ggplot2 version 3.3.3[88]. To compare both LPS-treatment up-
regulated and down-regulated differentially expressed genes (DEGs) in iTF-Microglia and
Brownjohn-iMG, DEGs that were significant ($p_{adj} < 0.05$) in at least one cell type were
visualized using VennDiagram (version 1.6.20).
1062
CROP-seq analysis. Alignment and gene expression quantification was performed on
scRNAseq libraries and sgRNA-enriched libraries using Cell Ranger (version 5.0.1, 10X
Genomics) with default parameters and reference genome GRCh38-3.0.0. Cellranger aggr was
used to aggregate counts belonging to the same sample across different GEM wells. The
resulting gene vs. cell barcode matrix contained 58,302 cells which had on average 41,827 reads
per cell, and a median of 3,346 genes per cell. sgRNA unique molecular identifier (UMI) counts
for each cell barcode were obtained using a previously described mapping workflow[86]. To
facilitate sgRNA identity assignment, a combination of demuxEM [95] and a z-score cut-off
method we previously described[15] were used such that only cells with a single sgRNA as
determined by both methods were carried forward in the analysis.
The raw gene vs. barcode matrix outputted by Cell Ranger was converted into a
SingleCellExperiment (SCE) object using the read10xCounts function from the DropletUtils
package version 1.10.3[96] in R (version 4.0.3). sgRNA assignments were appended to the SCE
metadata and filtered to only include cells with a single sgRNA, resulting in 28,905 cells. The
SCE was converted into a Seurat object using Seurat::as.Seurat version 4.0.1[97]. The data was
normalized and highly variable genes were identified using Seurat::SCTransform[98]. For initial
data exploration, principal-component analysis was performed using Seurat::RunPCA to
determine the number of principal components to retain. UMAP dimensional reduction using
Seurat::RunUMAP and clustering using Seurat::FindNeighbors and Seurat::FindClusters were
performed on the retained principal components with resolution = 0.7.
Initial data exploration revealed clusters that were not of interest due to a high proportion of
mitochondrial-encoding genes or disrupted microglia differentiation (Extended Data Fig.5A).
These clusters were removed from the downstream analysis and the remaining "microglia
cluster" population was normalized, clustered, and visualized using UMAP, as described above
with resolution = 0.25.
To determine the differentially expressed genes between UMAP clusters,
Seurat::FindAllMarkers was used. Single-cell heatmaps, ridge plots, rank plots, and UMAPs
were made using Seurat::DoHeatmap, Seurat::RidgePlot, Seurat::VizDimLoadings, and
Seurat::DimPlot or Seurat::FeaturePlot, respectively.
The relative proportion of cells with a given sgRNA in a given cluster compared to cells with
non-targeting control (NTC) sgRNAs in the given cluster was calculated and visualized using
Complex Heatmap version 2.6.2[92] (Supplemental Table 7).

For each CRISPRi target gene, the population of cells with the strongest knockdown (cells with expression of target gene less than the median expression of the target gene) was carried forward to perform differential gene expression analysis using Seurat::FindMarkers with parameters test.use = 't' (student's t-test), assay = "SCT", slot = "scale.data", to compare the Pearson residuals of cells[98] with knockdown sgRNAs versus non-targeting control cells. Genes with an adjusted p-value < 0.1 were deemed significant. The top 20 DEGs for each target gene which had >50 cells comprised the set of genes used to visualize convergent pathways using Complex Heatmap version 2.6.2[92].

The iTF-Microglia CROP-seq dataset was integrated with the previously published human scRNAseq dataset (Olah-hMG)[7] using Seurat[99]. Briefly, the Olah-hMG gene vs. cell barcode matrix and metadata were used to create a Seurat object and cells from surgery samples or with non-microglia identities as previously determined[7] were removed. Normalization and identification of highly variable genes was performed using Seurat::SCTransform with the same parameters as the iTF-Microglia. Next, integration features (3000 features) and integration anchors were identified for each Seurat object using Seurat::SelectIntegrationFeatures and Seurat::FindIntegrationAnchors and subsequent integration with identified anchors was performed using Seurat::IntegrateData. The integrated Seurat object was normalized, clustered, and visualized using UMAP, as described above with resolution = 0.25. Gene expression was visualized with UMAP using Seurat::FeaturePlot and the percentage of cells in the integrated SPP1-high cluster or SPP1-low clusters of either AD brain or control brain origin was calculated and significance was calculated on the cell counts using Fisher's exact test.

Image analysis with CellProfiler. Pipelines and example images are compiled in supplemental material and all analysis was performed using CellProfiler version 4.1.3. *Cell morphology metrics*: Nuclei were segmented as primary objects from Hoechst images. Cell segmentations were generated by propagating outward from nuclei objects until edges were identified in the phalloidin images. Area and shape metrics were calculated for each cell object. *Integrated F-actin intensity per cell*: for a given field of view, nuclei were segmented based on Hoechst images and total integrated SiR-actin intensity was summed. The resulting sum was divided and by the number of nuclei. *Longitudinal cell counts:* for a given field of view, nuclei were segmented based on Hoechst images acquired daily. *IBA1 intensity per cell:* This metric was determined similarly to the integrated F-actin intensity per nuclei; for a given field of view, the total integrated intensity of the IBA1 stain was divided by the number of segmented nuclei based on Hoechst.

Data Availability Statement
All screen datasets and RNA-transcriptomic datasets are publicly available in the CRISPRbrain data commons (http://crisprbrain.org/) (associated with Figures 1, 2, 4, 5, 6, 7 and Extended Data Figures 1, 4, 5, 6, 7). RNA sequencing datasets reported in this paper are in the process of being deposited on NCBI GEO. There are no restrictions on data availability.

Code Availability Statement
The MAGeCK-iNC bioinformatics pipeline for analysis of pooled screens available at https://kampmannlab.ucsf.edu/mageck-inc. The CellProfiler pipelines will be made available on request to the corresponding authors (MK), and will also be submitted to the CellProfiler

1140 depository of published pipelines (https://cellprofiler.org/examples/published_pipelines.html)
1141 upon publication.
1142
1143 **REFERENCES**
1144

1.   Butovsky, O. & Weiner, H.L. Microglial signatures and their role in health and disease.
Nat Rev Neurosci **19**, 622-635 (2018).

2.   Efthymiou, A.G. & Goate, A.M. Late onset Alzheimer's disease genetics implicates
microglial pathways in disease risk. *Mol Neurodegener* **12**, 43 (2017).

3.   Li, Q. & Barres, B.A. Microglia and macrophages in brain homeostasis and disease. *Nat*
Rev Immunol **18**, 225-242 (2018).

4.   Colonna, M. & Butovsky, O. Microglia Function in the Central Nervous System During
Health and Neurodegeneration. *Annu Rev Immunol* **35**, 441-468 (2017).

5.   Liddelow, S.A.*, et al.* Neurotoxic reactive astrocytes are induced by activated microglia.
Nature **541**, 481-487 (2017).

6.   Rodriguez-Gomez, J.A.*, et al.* Microglia: Agents of the CNS Pro-Inflammatory
Response. *Cells* **9** (2020).

7.   Olah, M.*, et al.* Single cell RNA sequencing of human microglia uncovers a subset
associated with Alzheimer's disease. *Nat Commun* **11**, 6129 (2020).

8.   Mathys, H.*, et al.* Temporal Tracking of Microglia Activation in Neurodegeneration at
Single-Cell Resolution. *Cell Rep* **21**, 366-380 (2017).

9.   Keren-Shaul, H.*, et al.* A Unique Microglia Type Associated with Restricting
Development of Alzheimer's Disease. *Cell* **169**, 1276-1290.e1217 (2017).

10.   Srinivasan, K.*, et al.* Untangling the brain's neuroinflammatory and neurodegenerative
transcriptional responses. *Nat Commun* **7**, 11295 (2016).

11.   Hammond, T.R.*, et al.* Single-Cell RNA Sequencing of Microglia throughout the Mouse
Lifespan and in the Injured Brain Reveals Complex Cell-State Changes. *Immunity* **50**, 253-
271.e256 (2019).

12.   Masuda, T.*, et al.* Spatial and temporal heterogeneity of mouse and human microglia at
single-cell resolution. *Nature* **566**, 388-392 (2019).

13.   Sankowski, R.*, et al.* Mapping microglia states in the human brain through the integration
of high-dimensional techniques. *Nat Neurosci* **22**, 2098-2110 (2019).

14.   Kampmann, M. CRISPR-based functional genomics for neurological disease. *Nat Rev*
Neurol **16**, 465-480 (2020).

15.   Tian, R*., et al.* CRISPR Interference-Based Platform for Multimodal Genetic Screens in
Human iPSC-Derived Neurons. *Neuron* **104**, 239-255 e212 (2019).

16.   Tian, R*., et al.* Genome-wide CRISPRi/a screens in human neurons link lysosomal failure
to ferroptosis. *Nat Neurosci* (2021).

17.   Abud, E.M*., et al.* iPSC-Derived Human Microglia-like Cells to Study Neurological
Diseases. *Neuron* **94**, 278-293 e279 (2017).

18.   Douvaras, P*., et al.* Directed Differentiation of Human Pluripotent Stem Cells to
Microglia. *Stem cell reports* **8**, 1516-1524 (2017).

19.   Muffat, J*., et al.* Efficient derivation of microglia-like cells from human pluripotent stem
cells. *Nature medicine* **22**, 1358-1367 (2016).

20.   Pandya, H*., et al.* Differentiation of human and murine induced pluripotent stem cells to
microglia-like cells. *Nature neuroscience* **20**, 753-759 (2017).

21.   Haenseler, W*., et al.* A Highly Efficient Human Pluripotent Stem Cell Microglia Model
Displays a Neuronal-Co-culture-Specific Expression Profile and Inflammatory Response. *Stem
cell reports* **8**, 1727-1742 (2017).

22.   Brownjohn, P.W*., et al.* Functional Studies of Missense TREM2 Mutations in Human
Stem Cell-Derived Microglia. *Stem cell reports* **10**, 1294-1307 (2018).

23.   McQuade, A*., et al.* Development and validation of a simplified method to generate
human microglia from pluripotent stem cells. *Mol Neurodegener* **13**, 67 (2018).

24.   Chen, S.W*., et al.* Efficient conversion of human induced pluripotent stem cells into
microglia by defined transcription factors. *Stem Cell Reports* **16**, 1363-1380 (2021).

25.   Zhang, Y*., et al.* An RNA-sequencing transcriptome and splicing database of glia,
neurons, and vascular cells of the cerebral cortex. *The Journal of neuroscience : the official
journal of the Society for Neuroscience* **34**, 11929-11947 (2014).

26.   Bennett, M.L*., et al.* New tools for studying microglia in the mouse and human CNS.
Proceedings of the National Academy of Sciences of the United States of America **113**, E1738-
1746 (2016).

27.   Hickman, S.E*., et al.* The microglial sensome revealed by direct RNA sequencing. *Nature
neuroscience* **16**, 1896-1905 (2013).

28.   Kierdorf, K*., et al.* Microglia emerge from erythromyeloid precursors via Pu.1- and Irf8-
dependent pathways. *Nat Neurosci* **16**, 273-280 (2013).

29.   Cuevas, V.D*., et al.* MAFB Determines Human Macrophage Anti-Inflammatory
Polarization: Relevance for the Pathogenic Mechanisms Operating in Multicentric Carpotarsal
Osteolysis. *J Immunol* **198**, 2070-2081 (2017).

30. Strohmeyer, R., Shelton, J., Lougheed, C. & Breitkopf, T. CCAAT-enhancer binding
protein-β expression and elevation in Alzheimer's disease and microglial cell cultures. *PLoS One*
9, e86617 (2014).

31. Masuda, T.*, et al.* Transcription factor IRF5 drives P2X4R+-reactive microglia gating
neuropathic pain. *Nat Commun* **5**, 3771 (2014).

32. Bohlen, C.J.*, et al.* Diverse Requirements for Microglial Survival, Specification, and
Function Revealed by Defined-Medium Cultures. *Neuron* **94**, 759-773.e758 (2017).

33. Horlbeck, M.A.*, et al.* Compact and highly active next-generation libraries for CRISPR-
mediated gene repression and activation. *Elife* **5** (2016).

34. Michaelson, M.D.*, et al.* CSF-1 deficiency in mice results in abnormal brain
development. *Development* **122**, 2661-2672 (1996).

35. Kondo, Y. & Duncan, I.D. Selective reduction in microglia density and function in the
white matter of colony-stimulating factor-1-deficient mice. *J Neurosci Res* **87**, 2686-2695
(2009).

36. Erblich, B., Zhu, L., Etgen, A.M., Dobrenis, K. & Pollard, J.W. Absence of colony
stimulation factor-1 receptor results in loss of microglia, disrupted brain development and
olfactory deficits. *PloS one* **6**, e26317 (2011).

37. Chitu, V.*, et al.* Microglial Homeostasis Requires Balanced CSF-1/CSF-2 Receptor
Signaling. *Cell Rep* **30**, 3004-3019 e3005 (2020).

38. Tan, F., Qian, C., Tang, K., Abd-Allah, S.M. & Jing, N. Inhibition of transforming
growth factor β (TGF-β) signaling can substitute for Oct4 protein in reprogramming and
maintain pluripotency. *J Biol Chem* **290**, 4500-4511 (2015).

39. Butovsky, O.*, et al.* Identification of a unique TGF-β-dependent molecular and functional
signature in microglia. *Nat Neurosci* **17**, 131-143 (2014).

40. Adler, A.S.*, et al.* CDK8 maintains tumor dedifferentiation and embryonic stem cell
pluripotency. *Cancer Res* **72**, 2129-2139 (2012).

41. Guerreiro, S., Privat, A.L., Bressac, L. & Toulorge, D. CD38 in Neurodegeneration and
Neuroinflammation. *Cells* **9** (2020).

42. Mayo, L.*, et al.* Dual role of CD38 in microglial activation and activation-induced cell
death. *J Immunol* **181**, 92-103 (2008).

43. Wang, Y.M.*, et al.* Blocking the CD38/cADPR pathway plays a double-edged role in
LPS stimulated microglia. *Neuroscience* **361**, 34-42 (2017).

44. Tanzer, M.C., Bludau, I., Stafford, C.A., Hornung, V. & Mann, M. Phosphoproteome
profiling uncovers a key role for CDKs in TNF signaling. *Nat Commun* **12**, 6053 (2021).

45.   Henry, K.L.*, et al.* CDK12-mediated transcriptional regulation of noncanonical NF-κB
components is essential for signaling. *Sci Signal* **11** (2018).

46.   Jin, Z., Wei, W., Yang, M., Du, Y. & Wan, Y. Mitochondrial complex I activity
suppresses inflammation and enhances bone resorption by shifting macrophage-osteoclast
polarization. *Cell Metab* **20**, 483-498 (2014).

47.   Salter, M.W. & Stevens, B. Microglia emerge as central players in brain disease. *Nat Med*
23, 1018-1027 (2017).

48.   Floden, A.M. & Combs, C.K. Microglia demonstrate age-dependent interaction with
amyloid-beta fibrils. *Journal of Alzheimer's disease : JAD* **25**, 279-293 (2011).

49.   Bliederhaeuser, C.*, et al.* Age-dependent defects of alpha-synuclein oligomer uptake in
microglia and monocytes. *Acta Neuropathol* **131**, 379-391 (2016).

50.   Hong, S.*, et al.* Complement and microglia mediate early synapse loss in Alzheimer
mouse models. *Science* **352**, 712-716 (2016).

51.   Lui, H.*, et al.* Progranulin Deficiency Promotes Circuit-Specific Synaptic Pruning by
Microglia via Complement Activation. *Cell* **165**, 921-935 (2016).

52.   Gilbert, L.A.*, et al.* Genome-Scale CRISPR-Mediated Control of Gene Repression and
Activation. *Cell* **159**, 647-661 (2014).

53.   Ramkumar, P.*, et al.* CRISPR-based screens uncover determinants of immunotherapy
response in multiple myeloma. *Blood Adv* **4**, 2899-2911 (2020).

54.   Wu, C.H.*, et al.* Mutations in the profilin 1 gene cause familial amyotrophic lateral
sclerosis. *Nature* **488**, 499-503 (2012).

55.   Schulz, D., Severin, Y., Zanotelli, V.R.T. & Bodenmiller, B. In-Depth Characterization
of Monocyte-Derived Macrophages using a Mass Cytometry-Based Phagocytosis Assay. *Sci Rep*
9, 1925 (2019).

56.   Zou, L.*, et al.* Profilin-1 is a negative regulator of mammary carcinoma aggressiveness.
Br J Cancer **97**, 1361-1371 (2007).

57.   Lu, E.*, et al.* Profilin 1 knockdown prevents ischemic brain damage by promoting M2
microglial polarization associated with the RhoA/ROCK pathway. *J Neurosci Res* **98**, 1198-1212
(2020).

58.   Masuda, T., Sankowski, R., Staszewski, O. & Prinz, M. Microglia Heterogeneity in the
Single-Cell Era. *Cell Rep* **30**, 1271-1281 (2020).

59.   Cappellano, G.*, et al.* The *Yin-Yang* of osteopontin in nervous system diseases:
damage *versus* repair. *Neural Regen Res* **16**, 1131-1137 (2021).

60.   Sala Frigerio, C*., et al.* The Major Risk Factors for Alzheimer's Disease: Age, Sex, and
Genes Modulate the Microglia Response to Aβ Plaques. *Cell Rep* **27**, 1293-1306.e1296 (2019).

61.   Szulzewsky, F*., et al.* Glioma-associated microglia/macrophages display an expression
profile different from M1 and M2 polarization and highly express Gpnmb and Spp1. *PLoS One*
10, e0116644 (2015).

62.   Geirsdottir, L*., et al.* Cross-Species Single-Cell Analysis Reveals Divergence of the
Primate Microglia Program. *Cell* **179**, 1609-1622.e1616 (2019).

63.   Zhan, L*., et al.* A MAC2-positive progenitor-like microglial population is resistant to
CSF1R inhibition in adult mouse brain. *Elife* **9** (2020).

64.   Zhan, L*., et al.* Proximal recolonization by self-renewing microglia re-establishes
microglial homeostasis in the adult mouse brain. *PLoS Biol* **17**, e3000134 (2019).

65.   Johnson, N.R*., et al.* Sex-specific life extension in tauopathy mice by CSF1R inhibition
causing selective microglial depletion and suppressed pathogenesis. *bioRxiv*,
2021.2003.2020.436288 (2021).

66.   Lambert, J.C*., et al.* Meta-analysis of 74,046 individuals identifies 11 new susceptibility
loci for Alzheimer's disease. *Nat Genet* **45**, 1452-1458 (2013).

67.   Yao, X*., et al.* Targeted genetic analysis of cerebral blood flow imaging phenotypes
implicates the INPP5D gene. *Neurobiol Aging* **81**, 213-221 (2019).

68.   Peng, Q*., et al.* TREM2- and DAP12-dependent activation of PI3K requires DAP10 and
is inhibited by SHIP1. *Sci Signal* **3**, ra38 (2010).

69.   Tsai, A.P*., et al.* INPP5D expression is associated with risk for Alzheimer's disease and
induced by plaque-associated microglia. *Neurobiol Dis* **153**, 105303 (2021).

70.   Pedicone, C*., et al.* Pan-SHIP1/2 inhibitors promote microglia effector functions essential
for CNS homeostasis. *J Cell Sci* **133** (2020).

71.   Huang, R.H*., et al.* Osteopontin Promotes Cell Migration and Invasion, and Inhibits
Apoptosis and Autophagy in Colorectal Cancer by activating the p38 MAPK Signaling Pathway.
Cell Physiol Biochem **41**, 1851-1864 (2017).

72.   Xu, J*., et al.* Multimodal single-cell/nucleus RNA sequencing data analysis uncovers
molecular networks between disease-associated microglia and astrocytes with implications for
drug repurposing in Alzheimer's disease. *Genome Res* (2021).

73.   Spangenberg, E*., et al.* Sustained microglial depletion with CSF1R inhibitor impairs
parenchymal plaque development in an Alzheimer's disease model. *Nat Commun* **10**, 3758
(2019).

74.   Sosna, J*., et al.* Early long-term administration of the CSF1R inhibitor PLX3397 ablates
microglia and reduces accumulation of intraneuronal amyloid, neuritic plaque deposition and
pre-fibrillar oligomers in 5XFAD mouse model of Alzheimer's disease. *Mol Neurodegener* **13**,
11 (2018).

75.   Hagan, N*., et al.* CSF1R signaling is a regulator of pathogenesis in progressive MS. *Cell*
Death Dis **11**, 904 (2020).

76.   Kana, V*., et al.* CSF-1 controls cerebellar microglia and is required for motor function
and social interaction. *J Exp Med* **216**, 2265-2281 (2019).

77.   Pons, V., Lévesque, P., Plante, M.M. & Rivest, S. Conditional genetic deletion of CSF1
receptor in microglia ameliorates the physiopathology of Alzheimer's disease. *Alzheimers Res*
Ther **13**, 8 (2021).

78.   Lin, Y.T*., et al.* APOE4 Causes Widespread Molecular and Cellular Alterations
Associated with Alzheimer's Disease Phenotypes in Human iPSC-Derived Brain Cell Types.
Neuron **98**, 1141-1154.e1147 (2018).

79.   TCW, J*., et al.* Cholesterol and matrisome pathways dysregulated in human
APOE ε4 glia. *bioRxiv*, 713362 (2019).

80.   Hasselmann, J*., et al.* Development of a Chimeric Model to Study and Manipulate
Human Microglia In Vivo. *Neuron* **103**, 1016-1033 e1010 (2019).

81.   Mancuso, R*., et al.* Stem-cell-derived human microglia transplanted in mouse brain to
study human disease. *Nature neuroscience* **22**, 2111-2116 (2019).

82.   Svoboda, D.S*., et al.* Human iPSC-derived microglia assume a primary microglia-like
state after transplantation into the neonatal mouse brain. *Proceedings of the National Academy of*
Sciences of the United States of America **116**, 25293-25303 (2019).

83.   Xu, R*., et al.* Human iPSC-derived mature microglia retain their identity and functionally
integrate in the chimeric mouse brain. *Nat Commun* **11**, 1577 (2020).

84.   Wang, C*., et al.* Scalable Production of iPSC-Derived Human Neurons to Identify Tau-
Lowering Compounds by High-Content Screening. *Stem cell reports* **9**, 1221-1233 (2017).

85.   Nagy, T. & Kampmann, M. CRISPulator: a discrete simulation tool for pooled genetic
screens. *BMC Bioinformatics* **18**, 347 (2017).

86.   Hill, A.J*., et al.* On the design of CRISPR-based single-cell molecular screens. *Nat*
Methods **15**, 271-274 (2018).

87.   Langmead, B., Trapnell, C., Pop, M. & Salzberg, S.L. Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. *Genome Biol* **10**, R25 (2009).

88.   Wickham, H. ggplot2 : Elegant Graphics for Data Analysis. in *Use R!,* 1 online resource
(XVI, 260 pages 232 illustrations, 140 illustrations in color (Springer International Publishing :
Imprint: Springer,, Cham, 2016).

89.   Patro, R., Duggal, G., Love, M.I., Irizarry, R.A. & Kingsford, C. Salmon provides fast
and bias-aware quantification of transcript expression. *Nat Methods* **14**, 417-419 (2017).

90.   Soneson, C., Love, M.I. & Robinson, M.D. Differential analyses for RNA-seq: transcript-
level estimates improve gene-level inferences. *F1000Res* **4**, 1521 (2015).

91.   Hasselmann, J*., et al.* Development of a Chimeric Model to Study and Manipulate
Human Microglia In Vivo. *Neuron* **103**, 1016-1033.e1010 (2019).

92.   Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinformatics* **32**, 2847-2849 (2016).

93.   Guttikonda, S.R*., et al.* Fully defined human pluripotent stem cell-derived microglia and
tri-culture system model C3 production in Alzheimer's disease. *Nat Neurosci* **24**, 343-354
(2021).

94.   Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

95.   Gaublomme, J.T*., et al.* Nuclei multiplexing with barcoded antibodies for single-nucleus
genomics. *Nat Commun* **10**, 2907 (2019).

96.   Lun, A.T.L*., et al.* EmptyDrops: distinguishing cells from empty droplets in droplet-
based single-cell RNA sequencing data. *Genome Biol* **20**, 63 (2019).

97.   Hao, Y*., et al.* Integrated analysis of multimodal single-cell data. *bioRxiv*,
2020.2010.2012.335331 (2020).

98.   Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-
seq data using regularized negative binomial regression. *Genome Biol* **20**, 296 (2019).

99.   Stuart, T*., et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-
1902.e1821 (2019).
1367
1368

1369
1370   **Figure 1 (legend overleaf)**

Figure 1: Rapid differentiation of iPSCs into microglia-like cells (iTF-Microglia) by transcription factor induction. a, Strategy for stable integration of six transcription factors integrated in AAVS1 and CLYBL loci by TALEN-mediated integration: The doxycycline-inducible reverse transcriptional activator (rtTA3G) is driven by the constitutive CAG promoter. Human MAFB, CEBPα and IRF8 are driven by the tet response element (TRE3G) in the AAVS1 locus. Human PU.1, CEBPβ and IRF5 are driven by TRE3G in the CLYBL locus. All transcription factors are separated from each other via T2A ribosome skipping sequences. **b,** Overview of the differentiation process for generating iTF-Microglia. *Top*: timeline with media and cytokines, *bottom*: representative phase-contrast images of cells on the indicated days. Scale bar: 100 μm. **c,** Expression of six inducible transcription factors during iTF-Microglia differentiation. Transcript abundance (TPM) of MAFB, CEBPα, IRF8 cassette and the PU.1, CEBPβ, IRF5 cassette at Day 0, Day 9 and Day 15 of differentiation. n = 3 biological replicates, p values from two-tailed Student's t-test. **d,** Representative immunofluorescence micrographs of iTF-Microglia on Day 8 of differentiation stained for microglia markers GPR34 and IBA1. Nuclei were labeled by Hoechst 33342. Scale bar: 100 μm. **e,** Expression of iPSC and microglia marker genes in iPSCs and derived iTF- Microglia on Day 9 and Day 15 of differentiation. The heatmap displays normalized and gene-centered transcripts per million (TPM) counts for selected genes (rows) for 3 biological replicates of timepoints (columns).iTF-Microglia express microglia homeostatic markers and activation markers, while losing their expression of iPSC markers. Asterisks highlight microglia-selective markers. **f,** Principal component analysis (PCA) on the expression of microglia marker genes of iTF-Microglia, human adult ex-vivo microglia[93], fetal and adult microglia[17], human myeloid cells[25], other iPSC-microglia[17, 22, 24] and iPSCs (this study and ref. [24]). Each dot reflects an independent biological sample. Colors represent the different cell types.

1396
1397
Figure 2: Functional characterization of iTF-Microglia. a, Phagocytosis of pHrodo-red-
labeled rat brain-derived synaptosomes by iTF-Microglia. Representative images at 0 h and 12 h
after synaptosome addition are shown. Treatment with 5 μM actin polymerization inhibitor
Cytochalasin D decreases phagocytosis. Scale bar: 100 μm. **b,** Phagocytosis of pHrodo-labeled
rat brain-derived synaptosomes with or without Cytochalasin D treatment was quantified by flow
cytometry at 0.5 h, 1.5 h and 2.5 h after synaptosome addition (mean +/- sd, n = three biological
replicates; p values from two-tailed Student's t-test). **c,** Morphological changes of iTF-Microglia
after LPS treatment are visualized by fluorescence microscopy. Samples were treated for 24 h
with 100 ng/ml LPS or buffer control and fixed samples were stained with AlexaFluor 488-

phalloidin for F-actin (green) and with Hoechst 33342 for nuclei (blue). Scale bar: 100 μm. **d,**
Transcriptomic changes caused by 50 ng/ml lipo-polysaccharide (LPS) treatment in Day 15 iTF-
Microglia (n = three biological replicates). Differentially expressed genes (p_{adj} < 0.05) are
labeled in black (increase). Other colors label genes associated with specific pathways that are
discussed in the main text. **e,** Cytokines secreted by iTF-Microglia. Analysis of cytokine array
signal (integrated density of dot blots) from supernatants of cultures treated with LPS or buffer
control (mean +/- sd, n = 6 biological replicates; p values from two-tailed Student's t-test).
Asterisk: GM-CSF is a component of the culture media. **f,** Co-culture with iPSC-derived
excitatory neurons promotes ramified morphology of iTF-Microglia. Representative fluorescence
micrographs at low and high magnification of Day 9 iTF- Microglia after 24 hours in co-culture.
iTF- Microglia express membrane-localized Lck-mNeonGreen (green). Neurons are stained for
the pre-synaptic marker synaptophysin (magenta). Nuclei are stained with Hoechst 33342 (blue).
Scale bars = 100 μm.
1420
1421

1422
1423
Figure 3: Gene knockdown and overexpression by CRISPRi and CRISPRa in iTF-
Microglia. a, Strategies for constitutive and inducible CRISPR interference (CRISPRi)/CRISPR
activation (CRISPRa) in iTF-Microglia. *Top*: For constitutive CRISPRi, a dCas9-BFP-KRAB
construct (catalytically dead Cas9 (dCas9) fused to BFP and the KRAB transcriptional repressor
domain) is expressed from the constitutive CAG promotor integrated into the CLYBL safe-
harbor locus. *Middle*: For inducible CRISPRi, dCas9-BFP-KRAB is tagged with ecDHFR
degrons. *Bottom*: For inducible CRISPRa, CAG promotor-driven ecDHFR-dCas9-VPH was
stably integrated into the CLYBL locus. VPH, activator domains containing 4X repeats of VP48,
P65 and HSF1. Addition of trimethoprim (TMP) stabilizes the inducible CRISPRi/a machineries.

b,c, Functional validation of (b) constitutive or (c) inducible CRISPRi activity via flow
cytometry of TFRC surface protein level stained iTF-Microglia expressing a TFRC-targeting
sgRNA or a non-targeting control (NTC) sgRNA at different days of differentiation (mean +/-
sd, n = 3 biological replicates; p values from two-tailed Student's t-test). (c) TMP was added to
induce CRISPRi activity where indicated. **d,** Functional validation of inducible CRISPRi activity
via TFRC immunofluorescence (IF) microscopy on Day 8. Top row, non-targeting (NTC)
sgRNA. Bottom row, sgRNA targeting *TFRC*. TFRC: red, F-actin: green, nuclei: blue. Scale bar
= 100 μm. **e,** Functional validation of inducible CRISPRa activity via flow cytometry of CXCR4
surface protein level staining in iTF-Microglia expressing *CXCR4* sgRNA or non-targeting
control (NTC) sgRNA (mean +/- sd, n = 3 biological replicates; p values from two-tailed
Student's t-test). TMP was added to induce CRISPRa activity where indicated.
1444

Figure 4: Identification of modifiers of survival and inflammation by CRISPRi screens.
a, Strategy for the CRISPRi screen to identify modifiers of survival/proliferation. iPSCs
expressing the inducible CRISPRi construct were transduced with an sgRNA library targeting
the "druggable genome". On Day 0, doxycyline and cytokines were added to induce microglial
differentiation and TMP was added to induce CRISPRi activity. Samples of cell populations
were taken at Day 0 and at Day 15 and frequencies of the cells expressing a given sgRNA were
determined by next-generation sequencing (NGS) to calculate Gene Scores quantifying the
survival/proliferation phenotype for each gene knockdown. **b,** Comparison of Gene Scores from
CRISPRi survival screens in iTF-Microglia (this study) vs. iPSC-derived neurons[15]. Each dot
represents a gene; genes are color-coded by pathways. **c,** Validation of the phenotype of CSF1R

knockdown. iTF-Microglia transduced with *CSF1R*-targeting or non-targeting control (NTC)
sgRNAs were imaged on different days after differentiation, and live cells were quantified based
on staining with Hoechst 33342. Data is shown as mean +/- sd, n = three wells per group, 7 fields
were imaged for each well. **d,** Strategy for a CRISPRi screen to identify modifiers of the
expression of CD38, a marker of reactive microglia. iPSCs expressing the inducible CRISPRi
construct were transduced with the druggable genome sgRNA library. On Day 0, doxycyline and
cytokines were added to induce microglial differentiation, and TMP was added to induce
CRISPRi activity. On Day 8, iTF-Microglia were stained for cell-surface levels of CD38 and
sorted by FACS into populations with low (bottom 30%) and high (top 30%) CD38 levels.
Frequencies of iTF-Microglia expressing a given sgRNA were determined in each population by
NGS. **e,** Volcano plot indicating knockdown phenotype and statistical significance (Mann-
Whitney U test) for genes targeted in the CD38 level screen. Dashed line indicates the cut-off for
hit genes (FDR = 0.1). Hit genes are shown in blue (knockdown decreases CD38 level) or red
(knockdown increases CD38 level), non-hit genes are shown in orange and "quasi-genes"
generated from random samples of non-targeting control sgRNAs are shown in grey. Hits of
interest are labeled. **f,** Validation of the phenotype of *MED1* and *CDK12* knockdown. CD38 cell
surface levels measured by flow cytometry of Day 8 iTF-Microglia targeting *MED1*, *CDK12*
compared to NTC sgRNA. n = 3 biological replicates; p values from two-tailed Student's t-test.
1475
1476
1477

1478

Figure 5: Identification of modifiers of phagocytosis by CRISPRi and CRISPRa screens.

a, Schematic of the screening strategy to identify modifiers of synaptosome phagocytosis. iPSCs

expressing inducible CRISPRi or CRISPRa constructs were transduced with an sgRNA library

targeting the druggable genome. On Day 0, doxycyline and cytokines were added to induce

microglial differentiation and TMP was added to induce CRISPRi activity. On Day 8, rat

synaptosomes labeled with pHrodo Red were added to the cells for 1.5 h and iTF-Microglia were
sorted based on fluorescence. Frequencies of cells expressing a given sgRNA in the low-
fluorescence and high-fluorescence populations were determined by next-generation sequencing
(NGS). **b**, Volcano blots summarizing knockdown and overexpression phenotypes and statistical
significance (Mann-Whitney U test) for genes targeted in the pooled phagocytosis screens. Left,
CRISPRi screen: right, CRISPRa screen. Dashed lines: Gene Score cutoff for hit genes (FDR =
0.1). Hit genes are shown in blue (knockdown decreases phagocytosis) or red (knockdown
increases phagocytosis), non-hit genes are shown in orange and "quasi-genes" generated from
random samples of non-targeting control sgRNAs are shown in grey. Hits of interest are labeled.
c, Competitive phagocytosis assay to test substrate specificity of *CD209* overexpression. Flow
cytometry measurement of phagocytosis of pHrodo-Red-labelled synaptosomes (*Left*, either
synaptosomes alone or together with beads) and green, fluorescent beads (*Right*, either beads
alone or together with synaptosomes) by iTF-Microglia expressing either non-targeting control
(NTC) sgRNAs or sgRNAs targeting *CD209*. Values represent mean +/- sd of n=3. Data was
analyzed using two-tailed Student's t-test. **d**, Representative fluorescent images demonstrating
higher F-actin staining in CRISPRa iTF-microglia at Day 8 with *PFN1* sgRNAs compared to
non-targeting control (NTC) sgRNAs (left). Right, integrated F-actin intensity per cell of
CRISPRa iTF-Microglia at Day 8 with *PFN1* sgRNAs or non-targeting control (NTC) sgRNAs.
(mean +/- sd, n = 5 fields of view from three different wells per sgRNA. P values from two-
tailed Student's t-test. **e**, Transcriptomic changes caused by *PFN1* overexpression in Day 8 iTF-
Microglia (n = 3 biological replicates). Differentially expressed genes ($p_{adj} < 0.05$) are labelled in
black. Other colors label genes associated with specific pathways that are discussed in the main
text. Alzheimer's disease (AD) risk genes are labelled with an asterisk.
1507
1508

1509
1510 **Figure 6 (legend overleaf)**

Figure 6: Single-cell RNA sequencing reveals distinct and disease-related microglia subclusters. a, Strategy for the CROP-seq screen. IPSCs expressing inducible CRISPRi machinery were transduced with a pooled library of 81 sgRNAs CROP-seq vector pMK1334. iPSCs are differentiated to iTF-Microglia and subjected to scRNAseq to obtain single-cell transcriptomes and identify expressed sgRNAs. **b**, UMAP projection of the 28,905 cells in the post-QC CROP-seq dataset. Cells are colored by sgRNA (*CDK8*-red, *TGFBR2*-orange) and cells with a high percentage of mitochondrial transcripts (blue). Microglia are labeled in green. Each dot represents a cell. **c**, UMAP projection depicting the 9 different clusters within the 19,834 microglia. Each dot represents a cell. The cells are color-coded based on their cluster membership. **d-e**, Ridge plots depicting iTF-Microglia clusters along PC1 (d) and PC2 (e). PC1 spans inflammation status (Interferon activated-homeostatic-chemokine activated) while PC2 spans proliferation status. **f**, Heatmap of iTF-Microglia clusters 1-9 and the relative expression of the top three differentially expressed genes of each cluster. **g**, UMAP projection of distinct marker expression of *CCL13* (*Left*) and *SPP1* (*Right*). *CCL13* is a marker for cluster 9 and *SPP1* is a marker for cluster 3. Cells are colored by the expression levels of the indicated gene. **h**, Phagocytic activity of iTF-Microglia in different states. Flow cytometry measurement of phagocytosis of pHrodo-Red-labelled synaptosomes (*Left*, Phagocytosis in CCL13^{high} and CCL13^{low} iTF-Microglia) (*Right*, Phagocytosis in SPP1^{high} and SPP1^{low} iTF-Microglia). Values represent mean +/- sd of n=3; p values from two-tailed Student's t-test. **i**, Integration of single-cell transcriptomes of iTF-Microglia and microglia from post-mortem human brains[7]. In the integrated UMAP, iTF-Microglia (*Left)* with high *SPP1* expression and human brain-derived microglia with high *SPP1* expression (*Right*) form a cluster (dashed outline). **j**, In brains from patients with Alzheimer's disease (AD), a higher fraction of microglia is in the SPP1^{high} cluster compared to control brains (data from Olah *et al*.[7]; P value from two-sided Fisher's exact test).

1537
Figure 7: CROP-seq reveals changes in cluster occupancy induced by gene knockdown.
a, sgRNA distribution across the iTF-Microglia clusters. UMAP projection depicts cells colored
by sgRNA. Cells with sgRNAs targeting *MAPK14* (blue), *CSF1R* (red), and *CDK12* (green), are
enriched in clusters 3, 6, and 9, respectively. Insert shows cluster 3. **b**, Changes in cluster
distribution after CRISPRi knockdown of targeted genes in iTF-Microglia. Heatmap with
hierarchical clustering of 37 target genes and non-targeting control (NTC) and their distribution
in clusters 1-9. **c,** Proportion of cells in cluster 9 (CCL13+) expressing either non-targeting
control (NTC) sgRNAs or sgRNAs targeting *CDK12*. **d**, Functional validation of increased

CCL13 level in iTF-Microglias expressing sgRNAs targeting *CDK12* compared to cells
expressing a non-targeting control (NTC) sgRNA. CCL13 levels were measured via flow
cytometry +/- 5h of GolgiPlug treatment. Values represent mean +/- sd of n = 3 biological
replicates; p values from two-tailed Student's t-test. **e,** Decreased synaptosome phagocytosis of
iTF-Microglia expressing sgRNAs targeting *CDK12* compared to cells expressing non-targeting
control (NTC) sgRNA. Phagocytosis is further reduced in the CCL13-high population of cells
expressing sgRNAs targeting *CDK12*. Phagocytosis was measured via flow cytometry with
additional staining for CCL13. Values represent mean +/- sd of n = 3 biological replicates; p
values from two-tailed Student's t-test. **f,** Proportion of cells in cluster 3 (SPP1+) expressing
either non-targeting control (NTC) sgRNAs or sgRNAs targeting *MAPK14* or *CSF1R*. **g-h,**
Functional validation of altered percentage of SPP1 positive cells in iTF-Microglias expressing
sgRNAs targeting *MAPK14* (g) or CSF1R (h) compared to cells expressing a non-targeting
control (*NTC*) sgRNA. SPP1 was measured via flow cytometry after treating cells for 5h with
GolgiPlug. Values represent mean +/- sd of n = 3 biological replicates; p values from two-tailed
Student's t-test. **i,** Survival of iTF-Microglia after treatment with various concentrations of
MAPK14 inhibitor Skepinone-L. Viable cells were quantified using the CellTiter-Glo assay 24h
after treatment with Skepinone-L. Values represent mean +/- sd of n = 12 biological replicates.
Data was analyzed by ANOVA. **j,** Percentage of SPP1-positive cells after 100 nM Skepinone-L
treatment. SPP1 was measured via flow cytometry after treating cells for 24 h or 36 h with
Skepinone-L and an additional 5 h with GolgiPlug. Values represent mean +/- sd of n = 3
biological replicates; p values from two-tailed Student's t-test. **k,** Survival of iTF-Microglia after
treatment with various concentrations of CSF1R inhibitor PLX3397. Viable cells were quantified
using the CellTiter-Glo assay, 24h after treatment with PLX3397. Values represent mean +/- sd
of n = 6 biological replicates; p values from two-tailed Student's t-test. **j,** Percentage of SPP1
positive cells after PLX3397 treatment. SPP1 was measured via flow cytometry after treating
cells for 24 h with PLX3397 and an additional 5 h with GolgiPlug. Values represent mean +/- sd
of n = 3 biological replicates; p values from two-tailed Student's t-test.
1573

1574

Extended Data Figure 1: Impact of Doxycycline removal on iTF-Microglia survival and sgRNA recovery in iPSC-derived microglia generated with different protocols. **a** Comparison of iTF-Microglia viability after Day 8 with different protocols. *Top*: timeline with different doxycycline supplementation paradigms, *bottom*: representative phase-contrast images at Day 15 with the indicated doxyccycline supplementation. Scale bar: 50 μm**. b**, Survival of iTF-Microglia at Day 15 after different doxycycline treatments indicated in a. Viable cells were quantified using the CellTiter-Glo assay. Values represent mean +/- sd of n = 12 biological replicates; p values from two-tailed Student's t-test. **c**, Principal component analysis (PCA) on the expression of microglia marker genes of iTF-Microglia, human adult ex-vivo microglia[93], fetal and adult microglia[17], human myeloid cells[25], other iPSC-microglia[17, 22, 24]. No iPSC samples were included. Each dot reflects an independent biological sample. Colors represent the different cell types. **d**, sgRNA recovery after transduction with a pooled sgRNA library in iPSCs and differentiation with two different iPSC-Microglia protocols. Strategy for the infection of iPSCs with an sgRNA library with 13,025 elements and timepoint of sgRNA recovery in iPSC-Microglia with the actual recovered counts of sgRNAs after next-generation-sequencing (NGS) from the protocol from Brownjohn *et al.*[22] (*Top*) and iTF-Microglia (*Bottom*).

1592
1593
Extended Data Figure 2: Phagocytosis capacity of iTF-Microglia and morphological
changes after LPS treatment. a-b, Phagocytosis of yellow-green (YG) beads (a) or pHRodo-
Red labelled synaptosomes (b) measured by flow cytometry. Histograms of YG-beads-FITC (a)
and Synaptosome-PE (b) after 1.5h of substrate exposure +/- 5μM Cytochalasin D (CytoD)
treatment. Controls are iTG-Microglia without substrate exposure. **c,** Phagocytosis of yellow-
green (YG) beads at different timepoints. Flow cytometric quantification of the percentage of
YG bead-positive cells at after 0.5 h, 1.5 h and 2.5 h of incubation with beads. Addition of 5 μM
CytoD decreases the percentage of YG bead-positive cells. N = three individual biological
replicates; p values from two-tailed Student's t-test. **d**, Morphological changes of iTF-Microglia

after LPS treatment. Swarm plots showing the automated quantification of microglia F-actin
staining in area, shape factor and perimeter with explanation of the three parameters. N = 16
wells from 3 individual differentiations; p values from two-tailed Mann-Whitney test. **e,**
Comparison of differentially expressed genes in response to LPS treatment in iTF-Microglia
versus iPSC-derived microglia (iMG) differentiated following a previously published protocol by
Brownjohn *et al.*[22].
1609

1610
1611
1612   **Extended Data Figure 3: Karyotyping of the monoclonal iTF-iPSC lines.** A normal karyotype
1613   was confirmed for monoclonal **a,** iTF-iPSCs, **b,** constitutive CRISPRi iTF-iPSC, **c,** inducible
1614   CRISPRi iTF-iPSC, **d,** inducible CRISPRa iTF-iPSC lines.

Extended Data Figure 4 (legend overleaf)

Extended Data Figure 4: Functional validation of CRISPRi/a activity in iPSCs and iTF-Microglia. a-b, Functional validation of constitutive (a) or inducible (b) CRISPRi activity via flow cytometry of TFRC surface protein level stained iPSCs expressing a TFRC-targeting sgRNA or a non-targeting control (NTC) sgRNA (mean +/- sd, n = 3 biological replicates; p values from two-tailed Student's t-test). TMP was added to induce CRISPRi activity where indicated. **c-d, Knockdown of TFRC in iPSCs with (a) the constitutive and (b) the inducible CRISPRi system.** qPCR quantification of the relative fold change of TFRC mRNA levels in CRISPRi-iPSCs expressing a TFRC sgRNA as compared to a non-targeting control sgRNA in the presence or absence of trimethoprim (TMP). (mean +/- sd, n = 3 biological replicates; p values from two-tailed Student's t-test). TFRC levels were normalized to the housekeeping gene GAPDH. **e-j Knockdown of three different genes in iTF-Microglia with (e,g,i) constitutive CRISPRi and (f,h,j) inducible CRISPRi.** qPCR quantification of the relative fold change of *TFRC* mRNA levels (e,f), *INPP5D* mRNA levels (g,h) or *PICALM* mRNA levels (I,j) in CRISPRi-iTF-Microglia expressing a *TFRC* sgRNA (e,f), *INPP5D* sgRNA (g,h) or *PICALM* sgRNA (I,j) compared to a non- targeting control sgRNA at different days of differentiation in the presence of TMP (mean +/- sd, n = 3 biological replicates). **k,** Functional validation of inducible CRISPRa activity via flow cytometry of CXCR4 surface protein level stained iPSCs expressing a CXCR4-targeting sgRNA or a non-targeting control (NTC) sgRNA (mean +/- sd, n = 3 biological replicates; p values from two-tailed Student's t-test). TMP was added to induce CRISPRi activity where indicated. **l-m,** qPCR quantification of the relative fold change of CXCR4 mRNA levels in inducible CRISPRa-iPSCs expressing a CXCR4 sgRNA as compared to a non-targeting control sgRNA in the presence or absence of trimethoprim (TMP), which stabilizes the DHFR degron. (mean +/- sd, n = 3 biological replicates; p values from two-tailed Student's t-test). CXCR4 levels were normalized to the housekeeping gene GAPDH.

Extended Data Figure 5: Knockdown of *CDK8* and *TGFBR2* induces proliferation and
decreases microglia markers in iPSC-derived microglia generated with different protocols
a, Comparison of Gene Scores from CRISPRi survival/proliferation screens in iTF-Microglia
(this study) vs. iPSCs[15]. Each dot represents a gene. **b-c**, IBA1 staining in Day 8 CRISPRi iTF-
Microglia containing sgRNAs targeting *CDK8* or *TGFBR2* compared to non-targeting control
(NTC) sgRNAs. **b**, Representative images. Scale bar = 50 μm. **c**, Quantification. Mean +/-sd, n =
6 fields of view from 2 different wells per sgRNA; p values from two-tailed Student's t-test. **d-e**,

IBA1 staining in Day 8 iMGs generated by the protocol from Brownjohn *et al.*, 2018 expressing sgRNAs targeting *CDK8* compared to non-targeting control (NTC) sgRNAs. **d**, Representative images. Scale bar = 50 μm. **e**, Quantification. Mean +/-sd, n =9 fields of view from 3 different wells per sgRNA; p values from two-tailed Student's t-test. **f**, Relative change in live cells of iTF-Microglia at Day 8 (*left*) and Day 15 (*right*) containing sgRNAs targeting *CDK8*, *CSF1R* or *TGFBR2* compared to non-targeting control sgRNAs. The inducible CRISPRi system was stabilized with TMP from Day 0 – Day 8 (*left*) or Day 8 – Day 15 (*right*). (mean +/-sd, n = 3 biological replicates; p values from two-tailed Student's t-test. **g,** CD38 cell surface levels measured by flow cytometry in iTF-Microglia 24 h treatement with 100 ng/mL LPS or PBS control.

1663
1664
Extended Data Figure 6: Validation of phagocytosis hits and overview of genes selected for
the CROP-seq screen based on primary screens. a, Comparing Gene Scores for hits from
phagocytosis CRISPRi and CRISPRa screens. Each dot represents a gene. **b-d**, Validation of
(b,c) CRISPRi hits and (d) CRISPRa hits in (b,d) iTF-Microglia or (c) iPSC-derived microglia
differentiated using an alternative protocol by Brownjohn *et al*.[22] Phagocytosis of pHrodo-
labelled synaptosomes by cells expressing either non-targeting control (NTC) sgRNAs or
sgRNAs targeting *CSF1R*, *INPP5D*, *PFN1* and *PFN2* was quantified by flow cytometry. Values
represent mean +/- sd of n = 3 biological replicates; p values from two-tailed Student's t-test. **e**,

Overexpression of CD209 (*left*) and PFN1 (*right*) with the inducible CRISPRa system in iTF-

Microglia. QPCR quantification of the relative fold change of CD209 and PFN1 mRNA levels in

iTF-Microglia expressing CD209 and PFN1 sgRNA as compared to a non-targeting control

sgRNA in the presence of TMP (mean +/- sd, n = 3 biological replicates; p values from two-

tailed Student's t-test). CD209 and PFN1 levels were normalized to the housekeeping gene

GAPDH. **f**, Binary heatmap of genes selected for the CROP-seq screen and their knockdown

phenotype in the CRISPRi survival, phagocytosis and inflammation screens. Red: KD increases

phenotype (positive hit). Blue: KD decreases phenotype (negative hit). Grey: not a significant

hit, p > 0.1.

1682

1683

Extended Data Figure 7: Characterization of microglia cluster signatures. a-c, UMAP projection representing single-cell transcriptomes, with cells colored based on (a) the percentage of mitochondrial transcripts, (b) the expressed sgRNAs, with sgRNAs targeting *CDK8* in orange, sgRNAs targeting *TGFBR2* in blue, non-targeting control sgRNAs (NTC) in green, and all other sgRNAs in grey, or (c) expression levels of *SOX2* (*Left*) or *CSF1R* (*Right*). **d**, Distribution of iTF-Microglia expressing non-targeting (NTC) sgRNAs across the 9 clusters described in Figure 6. **e-f**,the top 40 genes with the highest embedding values for (e) the first principal component (PC-1) and (f) the second principal component (PC-2), displayed in ranking order.

Top DEGs
(KD vs NTC)

Mitochondrial electron
transport chain

TGFβ regulation

Response to cytokine stimulus

Response to type 1 interferon

Ribosomal proteins

Avg diff
KD vs. NTC

−2

Cytoskeleton

Major histocompatibility process

Interferon signalling

CRISPRi
targeted
gene

1694
1695 **Extended Data Figure 8 (legend overleaf).**

Extended Data Figure 8: CROP-seq reveals transcriptomic changes in iTF-Microglia induced by gene knockdown. Changes in gene expression in response to CRISPRi knockdown of genes of interest in iTF-Microglia. Each column represents one CRISPRi-targeted gene. For each CRISPRi-targeted gene, cells with the strongest knockdown were selected and the top 20 differentially expressed genes in comparison to non-targeting control (NTC) sgRNA containing cells were selected. The merged set of these genes is represented by the rows. Rows and columns were clustered hierarchically based on Pearson correlation. Functionally related clusters of differentially expressed genes are labeled.

a

b

Extended Data Figure 9: Transcriptomic changes in iTF-Microglia induced by *CDK12*
knockdown in cluster 9 and in all other clusters. a, Changes in cluster distribution after
CRISPRi knockdown of targeted genes in iTF-Microglia. Distribution of cells according to the
37 targeted genes and non-targeting control (NTC) in clusters 1-9. **b**, Average differences of
gene expression induced by *CDK12* knockdown in cluster 9 compared to those in all other
clusters. Genes encoding phosphatidylserine (PS) recognition receptors are labeled in magenta
and Genes encoding MHC complex components are labeled in green.

1715

SUPPLEMENTARY TABLE LEGENDS

Supplementary Table 1. RNA-Seq Normalized Counts, Related to Figure 1. Gene-level counts per sample normalized to library size (transcript per million). Samples, in triplicate, include Day 0 iTF-iPSCs, Day 9 iTF-Microglia (PBS-treated and LPS-treated), Day 15 iTF-Microglia, and Day 9 Brownjohn-iMG (PBS-treated and LPS-treated). Columns are: Ensemble gene ID (ensembl_id), gene, all samples.

Supplementary Table 2. RNA-Seq LPS Differentially Expressed Genes, Related to Figure 2. Differentially expressed genes from comparing expression levels of LPS-treated cells to PBS-treated cells in Day 15 iTF-Microglia (first tab) and Brownjohn-iMG (second tab). Columns are: Ensembl gene ID (ensembl id), differentially expressed gene (gene), average expression over all samples (base mean), effect size estimate PBS vs. LPS ($\log_2$ fold change), $\log_2$ fold change standard error, p value, and adjusted p value. Tab 1 is iTF-Microglia, tab 2 is Brownjohn-iMG.

Supplementary Table 3. Primary Screen Phenotypes, Related to Figures 4, 5, and Extended Data Figure 6. Phenotypes from survival and FACS-based screens (survival, activation, and phagocytosis) are listed for all genes targeted in the H1 library. Columns are: targeted transcription start site (index), targeted gene (gene), knockdown phenotype, p value, and the gene score (product of phenotype $-\log_{10}$(p value)).

Supplementary Table 4. RNA-Seq, Differentially Expressed Genes as a result of *PFN1* overexpression in iTF-Microglia, Related to Figure 5. Differentially expressed genes of Day 8 iTF-Microglia overexpressing two different *PFN1* sgRNAs compared to non-targeting control sgRNA (NTC).

Supplementary Table 5. CROP-seq Pooled sgRNA Library, Related to Figures 6 and 7. Sequences for sgRNAs in CROP-seq pooled sgRNA library. Columns are: gene targeted for CRISPRi knockdown (target.gene), sgRNA short name as used in the paper (sgRNA.name), and sgRNA protospacer sequence (sgRNA.sequence).

Supplementary Table 6. Overview of CROP-seq results, Related to Figures 6 and 7.

Supplementary Table 7. CROP-seq Cluster Differentially Expressed Genes, Related to Figure 6. Differentially expressed genes (DEGs) for each UMAP cluster (1-9) compared to all other cluster, only positive values included. Columns are: gene, cluster, average log2 fold change, p value, adjusted p value

Supplementary Table 8. CROP-seq Target Gene Cluster Proportions, Related to Figure 7. Relative proportions of cells in clusters (rows) with a sgRNA targeting a given gene (columns) normalized to non-targeting control proportions (NTC).

Supplementary Table 9. CROP-seq Knockdown versus Control Differentially Expressed Genes, Related to Extended Data Figure 9. Differentially expressed genes, calculated using Student's T-test between cell with CRISPRi knockdown and non-targeting control (NTC) sgRNAs. Columns are: gene targeted for CRISPRi knockdown (TargetGene), differentially

1761 expressed gene (Gene), $\log_2$ counts per million, $\log_2$-fold change, p value, false discovery rate
1762 (FDR).
1763
1764 **Supplementary Table 10. Individual sgRNA sequences and primers.** qPCR and CROP-seq
1765 sgRNA enrichment primers, as well as individually cloned sgRNAs are listed. Columns are:
1766 name of sgRNA or primer (Name), section of sequence (sequence), sequence 5' to 3', use in this
1767 study (use).
1768
1769