1   Effects of tDCS on the attentional blink revisited: A statistical evaluation of a replication

2   attempt

3   Leon C. Reteig[1,2], Lionel A. Newman[1,3], K. Richard Ridderinkhof[1,2], & Heleen A. Slagter[1,4]

4   [1] Department of Psychology, University of Amsterdam

5   [2] Amsterdam Brain and Cognition, University of Amsterdam

6   [3] Department of Artificial Intelligence and Cognitive Engineering, University of Groningen

7   [4] Department of Applied and Experimental Psychology, Vrije Universiteit Amsterdam

8   Preprint submitted for peer-review.

9   Author Note

15   Abstract

The attentional blink (AB) phenomenon reveals a bottleneck of human information

processing: the second of two targets is often missed when they are presented in rapid

succession among distractors. A recent study by London & Slagter (*Journal of Cognitive

Neuroscience*, *33*, 756–68, 2021) showed that the size of the AB can be changed by applying

transcranial direct current stimulation (tDCS) over the left dorsolateral prefrontal cortex

(lDLPFC). Although AB size at the group level remained unchanged, the effects of anodal

and cathodal tDCS were negatively correlated: if a given individual's AB size decreased from

baseline during anodal tDCS, their AB size would increase during cathodal tDCS, and vice

versa. Here, we attempted to replicate this finding. Like London & Slagter, we found no

group effects of tDCS, but also no longer found a significant negative correlation. We present

a series of statistical measures of replication success, all of which confirm that both studies

are not in agreement. First, the correlation here is significantly smaller than a conservative

estimate of the original correlation. Second, the difference between the correlations is greater

than expected due to sampling error, and our data are more consistent with a zero-effect

than with the original estimate. Finally, the overall effect when combining both studies is

small and not significant. Our findings thus indicate that the effects of lDPLFC-tDCS on the

AB are less substantial than observed by London & Slagter (2021). Although this should be

quite a common scenario, null findings can be difficult to interpret and are still

under-represented in the brain stimulation and cognitive neuroscience literatures. An

important auxiliary goal of this paper is therefore to provide a tutorial for other researchers,

to maximize the evidential value from null findings.

Keywords: attentional blink, tDCS, dlPFC, individual differences, replication

Word count: 8076

Effects of tDCS on the attentional blink revisited: A statistical evaluation of a replication attempt

Introduction

The *attentional blink* (AB) phenomenon clearly demonstrates that our capacity to process incoming information is easily overwhelmed. The AB occurs when two targets are embedded in a rapidly presented stream of distractors (Raymond, Shapiro, & Arnell, 1992; for reviews, see Dux & Marois, 2009; Martens & Wyble, 2010). The first target (T1) is usually reported with little difficulty. When there is a longer lag between the two targets (> 500 ms; MacLean & Arnell, 2012), accuracy for the second target (T2) can be on par with the first. However, for shorter lags, T2 is most often missed—as if the attentional system momentarily faltered ("blinked").

While the AB might seem to be a fundamental bottleneck, it can under some circumstances be overcome. For example, the size of the AB can be reduced by distracting activities (Olivers, Nieuwenhuis, Olivers, & Nieuwenhuis, 2005; Olivers & Nieuwenhuis, 2006; Thomson, Ralph, Besner, & Smilek, 2015), or after following an intensive mental training program (Slagter et al., 2007). Others have tried to use non-invasive brain stimulation (Dayan, Censor, Buch, Sandrini, & Cohen, 2013) as a means to influence the AB. Several studies have shown that repetitive transcranial magnetic stimulation (TMS) can improve target perception in AB tasks (Arasanz, Staines, & Schweizer, 2012; Cooper, Humphreys, Hulleman, Praamstra, & Georgeson, 2004; Esterman et al., 2017). Yet, as TMS did not show a differential effect for targets presented at shorter or longer lags, it did not affect the AB itself.

Transcranial direct current stimulation (tDCS) is another brain stimulation technique that has gained traction in the past two decades. In tDCS, an electrical current flows between an anodal and cathodal electrode, which can affect the excitability of the underlying cortex (Gebodh et al., 2019). Anodal stimulation generally enhances cortical excitability,

while cathodal stimulation may have an inhibitory effect (Nitsche & Paulus, 2000, 2001)

(though note that this does not hold in all cases (Parkin, Bhandari, Glen, & Walsh, 2018),

and the underlying physiology is complex (Bikson, Paulus, Esmaeilpour, Kronberg, &

Nitsche, 2019; Liu et al., 2018; Stagg, Antal, & Nitsche, 2018)).

London and Slagter (2021) recently examined the effects of tDCS on the AB. They

applied anodal and cathodal tDCS over the left dorsolateral prefontal cortex (lDPLFC, with

the other electrode on the right forehead)—one of the core brain areas implicated in the AB

(Hommel et al., 2006; Slagter, Johnstone, Beets, & Davidson, 2010). At the group level,

tDCS did not appear to have any effects on AB size. However, anodal and cathodal tDCS

did appear to systematically affect the AB within individuals, as their effects were negatively

correlated. For a given individual, this negative correlation implies that when AB size

decreased (compared to a baseline measurement) during anodal tDCS, AB size would

increase during cathodal tDCS (or vice versa).

London and Slagter's (2021) findings complement earlier literature showing large

individual differences in both the AB (Willems & Martens, 2016) and effects of tDCS

(Krause & Cohen Kadosh, 2014). However, only one other tDCS study of the AB exists to

date, which in fact did find a group-level effect of stimulation, in contrast to London and

Slagter (2021) (Sdoia, Conversi, Pecchinenda, and Ferlazzo (2019)). Also, the negative

correlation between the effects of anodal and cathodal tDCS has so far only been reported in

London and Slagter (2021), and was based on an exploratory analysis. We thus decided to

conduct another study aiming to replicate this finding.

To foreshadow our results, like London and Slagter (2021), we did not find a group

effect of tDCS on the AB. However, the correlation between the effects of anodal and

cathodal tDCS was also not significant. Although this indicates that the two studies may

differ, a failure to reject the null hypothesis by itself does not tell us much (Harms & Lakens,

2018): it is crucial to also take measures of the uncertainty and effect size in both studies

into account (Simonsohn, 2015). That said, there is no consensus on when a replication can be considered successful or not (Brandt et al., 2014), let alone a single statistical test to definitely answer this question conclusively (Anderson & Maxwell, 2016).

Therefore, we employ a number of statistical methods to maximize the evidential value in these two studies. We ask three questions (cf. Zondervan-Zwijnenburg and Rijshouwer (2020)) that all aim to evaluate to what extent the present study is a successful replication of London and Slagter (2021; Zwaan, Etz, Lucas, and Donnellan, 2018; cf. Camerer et al., 2018; Open Science Collaboration, 2015). First, while the effect in our study was not significant, there might still be a meaningful effect that is simply smaller than anticipated. Therefore, we used equivalence testing (Lakens, Scheel, & Isager, 2018) to answer the question "*is the correlation in study 2 significantly small?*". Second, although our result differs from London and Slagter (2021), it could still be more consistent with their findings than with alternative explanations. So in addition, we asked "*is the correlation in study 2 different from study 1?*", and aimed to answer this question using prediction intervals (Patil, Peng, & Leek, 2016; Spence & Stanley, 2016) and replication Bayes factors (Verhagen & Wagenmakers, 2014; Wagenmakers, Verhagen, & Ly, 2016). Finally, it could be that the effect in our study alone is not sufficiently large, but the overall effect based on both studies is. This raises the question "*is the effect significant when combining study 1 and 2?*", which we addressed through meta-analysis (Goh, Hall, & Rosenthal, 2016; Quintana, 2015) and by pooling the data.

These questions address issues of reproducibility that are currently faced by many in the brain stimulation field (Héroux, Loo, Taylor, & Gandevia, 2017), and in the cognitive neuroscience community at large (Huber, Potter, & Huszar, 2019; Munafò et al., 2017). Therefore, aside from our focus on tDCS and the AB, an important auxiliary goal of this article is to provide a tutorial on the statistical evaluation of replication studies (also see Lakens, McLatchie, Isager, Scheel, and Dienes (2020), Zondervan-Zwijnenburg and Rijshouwer (2020)). We hope this may prove useful to other researchers who find themselves

in similar situations.[1]

Methods

Participants

Fourty-eight participants took part in total, eight of whom were excluded after the first session. One participant was excluded as a precaution because they developed an atypical headache after the first session, and we could not rule out this was related to the tDCS. Another stopped responding to our requests to schedule the second session. Another six participants were excluded because their mean T1 accuracy in the first session was too low, which implies they were not engaged in the task in the manner we expected, and would leave fewer trials to analyze, because our T2 accuracy measure included only trials in which T1 was seen. We used a cut-off of 63% T1 accuracy as an exclusion criterium, which was two standard deviations below the mean of a separate pilot study (n = 10).

This left a final sample of 40 participants (29 female, mean age = 20.94, $SD = 2.45$, range = 18–28). This sample size was determined a priori to slightly exceed London and Slagter (2021) (n = 34).

The experiment and recruitment took place at the University of Amsterdam. All procedures for this study were approved by the ethics review board of the Faculty for Social and Behavioral Sciences, and complied with relevant laws and institutional guidelines. All participants provided their written informed consent and were compensated with course credit or €10 per hour (typically €65 for completing two full sessions).

---

[1] We also aimed to examine the neurophysiological mechanisms underlying the individual differences in response to tDCS using EEG (Harty, Sella, & Cohen Kadosh, 2017; Krause & Cohen Kadosh, 2014; Li, Uehara, & Hanakawa, 2015). However, since we did not replicate the behavioral effect of tDCS on the AB of London and Slagter (2021), we were unable to fulfill this aim.

Procedure

The study procedures were identical to London and Slagter (2021): participants received anodal and cathodal tDCS in separate sessions (Figure 1), which typically took place exactly one week apart (cf. minimum of 48 hours in London and Slagter (2021)). The time in between served to keep the sessions as similar as possible, and to minimize the risk of tDCS carry-over effects. 18 participants received anodal tDCS in the first session and cathodal tDCS in the second, and vice versa for the remaining 22 participants.

Figure 1. **Experimental design**. After a baseline block without stimulation, participants performed the attentional blink task during 20 minutes of anodal (red) or cathodal (blue) tDCS, followed by a post-test block (also without stimulation). The second session (typically 7 days later) was identical, except that the tDCS polarity was reversed.

First, participants experienced the sensations induced by tDCS in a brief trial stimulation (see the tDCS section). Next, participants completed 20 practice trials of the task (see the Task section). For the main portion of the experiment, participants performed three blocks of the task (Figure 1): before tDCS (*baseline*), during anodal/cathodal tDCS (*tDCS*), and after tDCS (*post*). Finally, after completing the three blocks, participants filled in a questionnaire on tDCS-related adverse effects (see Table S1 and Figure S1 in the Supplementary Material).

151   Within each block of the task, participants took a self-timed break every 50 trials (~5

152 minutes); between the blocks, the experimenter walked in. Participants performed the task

153 for exactly 20 minutes during the *baseline* and *post* blocks. During the *tDCS* block, the task

154 started after the 1-minute ramp-up of the current was complete, and continued for 21

155 minutes (constant current, plus 1-minute of ramp-down).

Task

The attentional blink task (Figure 2) was almost identical to the one used in London

and Slagter (2021) and Slagter and Georgopoulou (2013), which in turn was based on a task

designed by Dux and Marois (2008). A rapid serial visual presentation stream of 15 letters

(cf. 17 letters in London and Slagter (2021)) was shown on each trial, using Presentation

software (Neurobehavioral Systems, Inc.). Each letter was displayed for 91.7 ms (11 frames

at 120 Hz) on a dark gray background. The letters were presented in font size 40 (font:

Courier New) at a viewing distance of 90 cm. On each trial, the letters were randomly

sampled without replacement from the alphabet, excluding the letters I, L, O, Q, U and V,

as they were too similar to each other. All distractor letters were mid-gray, whereas T1 and

T2 were colored. T1 was red and always appeared at position 5 in the stream. T2 was green

and followed T1 after either 2 distractors (*lag 3*) or 7 distractors (*lag 8*) (cf. lags 2, 4 and 10

in London and Slagter (2021)).

The letter stream was preceded by a fixation cross (same color as the letters) presented

for 1750 ms (cf. 480 ms in London and Slagter (2021)) and followed by another fixation cross

(cf. none in London and Slagter (2021)). Finally, the participant was prompted to type in

(using a standard keyboard) the letter they thought was presented as T1 ("Which letter was

red?"), followed by T2 ("Which letter was green?").

Trial duration varied slightly because both the T1 and T2 response questions were

self-paced, so some participants completed more trials than others depending on their

response times. On average, participants completed 130 short lag trials ($SD = 17$; range = 78–163) and 65 long lag trials ($SD = 9$; range = 39–87) per 20-minute block.

Figure 2. **Attentional blink task**. Participants viewed rapid serial visual presentation streams of 15 letters, all of which were distractors (gray letters) except for T1 and T2. T1 was presented in red at position 5; T2 was presented in green and followed T1 after 2 distractors (*lag 3*, inside the AB window) or 7 distractors (*lag 8*, outside the AB window). At the end of the trial, participants reported the identity of T1 and then T2 (self-paced).

tDCS

Transcranial direct current stimulation was delivered online (i.e. during performance of 180 the attentional blink task) using a DC-STIMULATOR PLUS (NeuroCare Group GmbH).

The current was ramped up to 1 mA in 1 minute, stayed at 1 mA for 20 minutes, and was

ramped down again in 1 minute.

One electrode was placed at F3 (international 10-20 system) to target the lDLPFC; the

other was placed over the right forehead, centered above the eye (approximately

corresponding to position Fp2). Both electrodes were 5 x 7 cm in size (35 cm^2), leading to a

current density of 0.029 mA/cm^2. The montage and tDCS parameters are identical to

London and Slagter (2021), the only exception being the conductive medium. We used

Ten20 conductive paste (Weaver and Company), because it was easier to apply concurrently

with the EEG equipment (see the EEG section); London and Slagter (2021) used saline

solution as a conductive medium, together with rubber straps to keep the electrodes in place.

Participants received either anodal tDCS (anode on F3, cathode on right forehead) or

cathodal tDCS (cathode on F3, anode on right forehead) in separate sessions. The procedure

was double-blinded: both the participant and the experimenters were unaware which polarity

was applied in a given session. The experimenter loaded a stimulation setting on the tDCS

device (programmed by someone not involved in data collection), without knowing whether

it was mapped to anodal or cathodal tDCS. In the second session, the experimenter loaded a

second setting mapped to the opposite polarity (half the dataset), or simply connected the

terminals of the device to the electrodes in the opposite way.

At the start of the experiment, participants received a brief trial stimulation, based on

which they decided whether or not they wanted to continue with the rest of the session. The

experimenter offered to terminate the experiment in case tDCS was experienced as too

uncomfortable, but none of the participants opted to do so. For the trial stimulation, the

current was ramped up to 1 mA in 45 seconds, stayed at 1 mA for 15 seconds, and was

ramped down again in 45 seconds.

EEG

We also recorded EEG during all three task blocks. As noted in the Introduction, we refrained from further analysis of the EEG data, given the absence of a behavioral effect. Instead, we are making the EEG data publicly available at this time.[2] The dataset is formatted according to the Brain Imaging Data Structure (BIDS) standard (Gorgolewski et al., 2016) for EEG (Pernet et al., 2019), to facilitate re-use. We include the raw data, as well as the fully preprocessed data and the MATLAB code that generated it.

Data analysis

Data were analyzed using R (Version 3.6.2; R Core Team, 2019)[3] from within RStudio (Version 1.1.463; RStudio Team, 2016).

Group-level analysis.   Repeated measures ANOVAs were conducted on T1 accuracy (percentage of trials where T1 was reported correctly) and T2|T1 accuracy (percentage of trials where T2 was reported correctly, out of the subset of T1-correct trials) using the *afex* package (Version 0.28-1; Singmann, Bolker, Westfall, Aust, & Ben-Shachar, 2021). The same factors were included for both repeated measures ANOVAs, following London and Slagter (2021): Lag (3, 8), Block (baseline, tDCS, post), Stimulation (anodal, cathodal), and the between-subject factor Session order (anodal tDCS in the first session vs. cathodal tDCS in the first session). Effect sizes are reported as generalized eta-squared ($\hat{\eta}^2_G$) (Bakeman, 2005). Greenhouse-Geisser-adjusted degrees of freedom (df^{GG}) and p-values are reported as a correction for sphericity violations.

---

[2] https://doi.org/10.18112/openneuro.ds001810.v1.1.0

[3] We, furthermore, used the R-packages *BayesFactor* (Version 0.9.12.4.2; Morey & Rouder, 2018), *broom* (Version 0.7.4.9000; Robinson, Hayes, & Couch, 2021), *cowplot* (Version 1.1.1; Wilke, 2020), *emmeans* (Version 1.5.4; Lenth, 2021), *here* (Version 1.0.1; Müller, 2020), *kableExtra* (Version 1.3.2; Zhu, 2021), *knitr* (Version 1.31; Xie, 2015), *papaja* (Version 0.1.0.9997; Aust & Barth, 2020), *psychometric* (Version 2.2; Fletcher, 2010), *pwr* (Version 1.3.0; Champely, 2020), and *tidyverse* (Version 1.3.0; Wickham et al., 2019).

Individual differences analysis.   We reproduced the analysis behind Figure 4 of

London and Slagter (2021), which showed a differential effect of anodal vs. cathodal tDCS at

the individual participant level. First, we calculated AB magnitude by subtracting T2|T1

accuracy at lag 3 from T2|T1 accuracy at lag 8. Next, change scores were created by

subtracting AB magnitude in the *baseline* block from the *tDCS* and the *post* blocks,

respectively. The change scores in the anodal and cathodal session were then correlated to

each other. Again following London and Slagter (2021), we computed a partial correlation

(using the *ggm* package (Version 2.5; Marchetti, Drton, & Sadeghi, 2020)), attempting to

adjust for variance due to Session order.

Replication analyses.   In contrast to London and Slagter (2021), the analysis

described in the previous section did not produce a significant correlation in our dataset.

Therefore, we conducted five follow-up analyses that aim to quantify to what extent our

results (do not) replicate London and Slagter (2021). These all provide a complementary

perspective on this question. First, we performed an equivalence test (1) to assess whether

the effect in the present study was significantly smaller than in London and Slagter (2021).

While this procedure is more focused on hypothesis testing, we also constructed prediction

intervals (2) to capture the range of effect sizes we can expect in a replication of London and

Slagter (2021). Both of these procedures are based on frequentist statistics, which cannot

directly quantify the amount of evidence for a (null) hypothesis. Therefore, we also

computed a replication Bayes factor that expresses whether the data in the present study are

more likely under the null hypothesis that the effect is absent, vs. the alternative hypothesis

that the effect is comparable to London and Slagter (2021). Finally, we directly combined

both studies and estimated the size of the overall effect, through meta-analysis (4) of both

correlations, and by computing a new correlation on the pooled dataset (5). More details on

each analysis can be found in the following sections, and the provided online resources (see

the Data, materials, and code availability section).

Equivalence tests. Equivalence tests can be used to test for the *absence* of an effect
of a specific size (see Lakens et al., 2018 for a tutorial). Usually, the effect size used for the
test is the smallest effect size of interest (the SESOI). Typically, equivalence tests are two
one-sided tests: one test of the null hypothesis that the effect exceeds the upper equivalence
bound (positive SESOI), and one that the effect exceeds the lower equivalence bound
(negative SESOI). However, a one-sided test is more appropriate here: London and Slagter
(2021) found that the effects of anodal and cathodal tDCS were anticorrelated, so we are only
interested in negative effect sizes. This is known as an inferiority test (Lakens et al., 2018).

We follow the "small telescopes" (Simonsohn, 2015) approach to set the SESOI to $r_{33\%}$:
the correlation that London and Slagter (2021) had 33% power to detect. The reasoning
behind this approach is that it is difficult to prove that an effect does not exist at all, but
easier to show that it is surprisingly small. An equivalence test can suggest that the effect is
unlikely to exceed $r_{33\%}$, such that the odds to detect it were stacked at least 2:1 against
London and Slagter (2021). That would not mean the effect does not exist at all, but it
would mean the original evidence for the effect is not very convincing, as "too small a
telescope" (in this case, an inadequate sample size) was used to reliably detect it.

There are many possible specifications of the SESOI, none of which are necessarily
wrong or right (Lakens et al., 2018). We favored the "small telescopes" approach because it
constitutes a relatively strict test—$r_{33\%}$ is much smaller than the original correlation in
London and Slagter (2021). Because the observed correlation in London and Slagter (2021)
could have overestimated the true correlation, it is prudent to set the SESOI to be smaller.
Furthermore, the approach was specifically designed to evaluate replication results
(Simonsohn, 2015), and has been used previously in large-scale replication studies (e.g.
Camerer et al., 2018). However, recent simulation studies have also demonstrated that
equivalence tests for replications can be either too liberal (especially in the case of
publication bias) (Muradchanian, Hoekstra, Kiers, & Ravenzwaaij, 2020), or too conservative
(especially in case of low sample sizes) (Linde, Tendeiro, Selker, Wagenmakers, &

Ravenzwaaij, 2020).

We conducted an inferiority test using the *TOSTER* package (Version 0.3.4; Lakens,

2017) against the null hypothesis that the correlation coefficient in the present study is at

least as negative as $-r_{33\%}$. At a standard alpha level of 0.05, the test is significant if the 90%

confidence interval around the observed correlation does not contain $r_{33\%}$. This would mean

that the observed correlation should be considered "statistically inferior", as it is then

significantly smaller (i.e. less negative) than $-r_{33\%}$.

Prediction interval. Prediction intervals contain a range of values we can expect a

new observation to fall within. In our case, the observation of interest is the correlation

between the effects of anodal and cathodal tDCS. This correlation is estimated based on a

sample, and is thus subject to sampling error: any two estimates of the correlation will

almost never be exactly the same. Prediction intervals aim to quantify how dissimilar two

estimates can be before we should be surprised.

Here, we construct a prediction interval around the original estimate of the correlation

in London and Slagter (2021). This prediction interval contains the range of correlation

coefficients we can expect to see in the present study, given the results of London and Slagter

(2021). The width of the interval depends on the sample sizes of both studies, as larger

samples will reduce variation in the estimates, leading to smaller prediction intervals (Patil

et al., 2016).

If the original study were replicated 100 times, 95 of the observed correlation

coefficients would fall within the 95% prediction interval (Patil et al., 2016). Note that this

definition is related to, but different from, a *confidence interval*, which quantifies uncertainty

about the (unknown) true correlation in the population (95 out of every hundred constructed

95% confidence intervals contain the true population parameter). Because prediction

intervals concern the next single observation, they make a more specific claim, and will be

wider than confidence intervals.

We calculated a 95% prediction interval for correlations, following Spence and Stanley

(2016), using the *predictionInterval* package (Version 1.0.0; Stanley, 2016).

Replication Bayes factor.   Bayes factors can be used to express the relative

evidence for the null (H_0) or alternative hypothesis (H_1) (Wagenmakers, Marsman, et al.,

2018). In a default Bayesian hypothesis test, H_0 states the effect size is absent (i.e. exactly

zero); H_1 states that the effect is present (specified further by a prior distribution of effect

sizes).

In a replication context, we want to decide between two different scenarios (Verhagen

& Wagenmakers, 2014). H_0 is the hypothesis of an idealized skeptic, who disregards the

information from the original study and believes the effect is absent. The alternative

hypothesis H_r belongs to an idealized proponent, who believes that the effect is exactly as in

the original study, i.e. their prior distribution is simply the posterior distribution of the

original study.

We used the replication Bayes factor test for correlations developed by Wagenmakers et

al. (2016). The replication Bayes factor BF_{0r} expresses evidence for H_0 : "the correlation is

0" relative to H_r : "the correlation is as in the original study". We use the interpretation

scheme from Wagenmakers, Love, et al. (2018), where $1 < BF_{0r} < 3$ consitutes "anecdotal

evidence" for H_0, $3 < BF_{0r} < 10$ ~ "moderate evidence", and $10 < BF_{0r} < 30$ ~ "strong

evidence".

Meta-analysis.   The outcomes from multiple studies on the same phenomenon can

be combined through meta-analysis. Here we compute a meta-analytic estimate of the

correlation based on the correlations observed here and as reported by London and Slagter

(2021), using the *metafor* package (Version 2.4-0; Viechtbauer, 2010). We weighed the

estimate by sample size, so the present study will have a slightly higher influence on the

meta-analytic effect size (because its sample size exceeds London and Slagter (2021)). We

specified the meta-analysis as a fixed-effects model, because both studies are highly similar

and from the same population (e.g., the experiments were conducted in the same location,

and the sample was from the same university student population). With a fixed-effects

analysis, we estimate the size of the effect *in the set of available studies*, meaning our

inferences cannot generalize beyond. A random-effects meta-analysis would be appropriate in

case the studies were more dissimilar, and if we sought an estimate of *the true effect in the*

population, but we would need more than just two studies for this approach. Note that while

meta-analyses are a powerful way to assess the overall effect in a series of studies, they are

particularly vulnerable to false positives when there is publication bias (Muradchanian et al.,

2020), or if the selection of studies (or any single study) is biased (Ueno, Fastrich, &

Murayama, 2016).

Pooling the data with London and Slagter (2021).   Another approach is to

pool the single-subject data from both studies, and to re-calculate the partial correlation on

the combined sample (n = 74). The main difference between the two studies is that London

and Slagter (2021) presented T2 at lags 2, 4 and 10; here we used lags 3 and 8. The long

lags (lag 8 vs. lag 10) should be fairly comparable, as they are both well outside the

attentional blink window (> 500 ms following T1; MacLean & Arnell, 2012). However, there

should be a sizeable performance difference at the short lags (lag 2 vs. lag 3), as the

attentional blink is larger at lag 2 than lag 3. Therefore, we opted to also create a "lag 3"

condition in the data from London and Slagter (2021), by averaging T2|T1 accuracy at lag 2

and lag 4. The difference from lag 2 to 4 (and 4 to 10) in London and Slagter (2021) looks

fairly linear (see their Figure 3), so this seems a fair approximation of "true" lag 3

performance. Afterwards we recomputed the partial correlation between AB magnitude

change scores as described previously (see the Individual differences analysis section).

Note that this analysis is tailored to this series of studies, and not generally advisable.

To get a more accurate estimate of the effect at lag 3, it is necessary to redo the analysis on

the larger, combined sample. But repeating a statistical test after collecting more data

("optional stopping") invalidates the interpretation of the p-value and can drastically

increase the false positive rate (Simmons, Nelson, & Simonsohn, 2011). This would only be acceptable when the analysis plan has been preregistered, and the false positive rate of sequential analyses is controlled (for potential solutions, see Lakens, 2014). We therefore do not report a p-value for this test, but only the effect size and its confidence interval.

Data, materials, and code availability

All of the data and materials from this study and the data from London and Slagter (2021) are available on the Open Science Framework[4]. The analysis code is available on GitHub[5] (and also from our OSF page), in the form of an R notebook detailing all the analyses that we ran for this project, along with the results. We also include an Rmarkdown (Xie, Allaire, & Grolemund, 2018) source file for this paper that can be run to reproduce the pdf version of the text, along with all the figures and statistics.

Results

Group-level analysis

Figure 3 shows the attentional blink (T2|T1 accuracy per lag) for each of the three blocks (pre, tDCS, post) and stimulation conditions separately. The summary statistics and ANOVA results for T2|T1 accuracy can be found in Tables 1 and 2. There was a clear attentional blink effect on average (main effect of Lag, $F[1, 38] = 432.11$, $p < .001$), as T2|T1 accuracy for lag 8 was higher than lag 3.

---

[4] https://doi.org/10.17605/OSF.IO/Y6HSF

[5] https://doi.org/10.5281/zenodo.4782446

Figure 3. **No effects of tDCS on the attentional blink at the group level**. There was a clear attentional blink effect: a lower % T2 accuracy (given T1 correct: *T2/T1*; dashed lines) for *lag 3* (T2 presented inside the attentional blink window) than *lag 8* (T2 presented outside the attentional blink window, on par with *T1 accuracy*). However, the attentional blink did not change systematically over stimulation conditions (*anodal*, *cathodal*) and blocks (*pre*, *tDCS*, *post*). T1 accuracy (solid lines) was also not affected by tDCS.

³⁷⁵ Effects of tDCS at the group-level should manifest as a three-way interaction between
³⁷⁶ Block, Stimulation, and Lag. As in London and Slagter (2021), this interaction was not
³⁷⁷ significant ($F[1.77, 67.17] = 2.77$, $p = .076$). However, the higher-order interaction with
³⁷⁸ Session order did reach significance ($F[1.77, 67.17] = 7.25$, $p = .002$). This interaction
³⁷⁹ appears to be mostly driven by a learning effect across sessions and blocks (the Session order
³⁸⁰ by Stimulation by Lag interaction). From the first to the second session, lag 3 performance
³⁸¹ improved, while lag 8 performance declined somewhat (leading to a smaller attentional
³⁸² blink). This change was stronger in participants that received anodal tDCS in the first
³⁸³ session, and less pronounced in the cathodal-first group. We do not consider this a genuine

Table 1

Descriptive statistics for T2|T1 accuracy

	First session: anodal (n = 18)		First session: cathodal (n = 22)	
	anodal	cathodal	anodal	cathodal
baseline				
lag 3	0.395 (0.149)	0.486 (0.167)	0.469 (0.211)	0.423 (0.168)
lag 8	0.826 (0.119)	0.819 (0.095)	0.854 (0.125)	0.855 (0.087)
tDCS				
lag 3	0.415 (0.168)	0.450 (0.165)	0.460 (0.201)	0.444 (0.188)
lag 8	0.787 (0.131)	0.819 (0.098)	0.830 (0.115)	0.849 (0.096)
post				
lag 3	0.437 (0.171)	0.451 (0.160)	0.453 (0.190)	0.447 (0.164)
lag 8	0.783 (0.118)	0.825 (0.098)	0.840 (0.145)	0.846 (0.120)

Note:

Values are "Mean (SD)"

effect of tDCS on the attentional blink, because there is no clear reason why these randomized groups should differ, and because the largest difference between the anodal and cathodal session occurred in the baseline block already (see Table 1).

We ran the same repeated measures ANOVA for T1 accuracy (Table 3). On average, participants correctly identified T1 in 82% of trials (Figure 3), which is comparable to previous studies using this task (86% in London & Slagter, 2021; 82% in Slagter & Georgopoulou, 2013). T1 accuracy was also slightly lower for lag 3 than lag 8 (main effect of Lag, $F[1, 38] = 29.23$, $p < .001$). There was a main effect of Block, reflecting that T1 accuracy decreased *within* a session ($F[1.89, 71.87] = 6.64$, $p = .003$). Finally, T1 performance also decreased *across* the sessions (interaction between Session order and

Table 2

Repeated measures ANOVA on T2/T1 accuracy

Effect	F	df_1^{GG}	df_2^{GG}	p	$\hat{\eta}_G^2$
Session order	0.33	1	38	.568	.006
Block	1.13	1.91	72.71	.328	.001
Stimulation	2.47	1	38	.125	.002
Lag	432.11	1	38	< .001	.634
Session order × Block	0.29	1.91	72.71	.741	.000
Session order × Stimulation	5.55	1	38	.024	.005
Session order × Lag	0.48	1	38	.494	.002
Block × Stimulation	0.28	1.91	72.51	.747	.000
Block × Lag	1.67	1.70	64.73	.199	.001
Stimulation × Lag	0.10	1	38	.751	.000
Session order × Block × Stimulation	1.93	1.91	72.51	.154	.001
Session order × Block × Lag	0.24	1.70	64.73	.752	.000
Session order × Stimulation × Lag	5.84	1	38	.021	.002
Block × Stimulation × Lag	2.77	1.77	67.17	.076	.001
Session order × Block × Stimulation × Lag	7.25	1.77	67.17	.002	.003

394 Stimulation: $F[1, 38] = 11.24$, $p = .002$).

395   In sum, we conclude that there is no significant effect of tDCS on the attentional blink

396 or T1 accuracy at the group level, in agreement with London and Slagter (2021).

Individual differences

Our main aim was to replicate London and Slagter (2021), who found a negative

correlation between AB magnitude change scores (comparing the *tDCS* and *baseline* blocks)

in the anodal and cathodal sessions ($r(31)$ = -.45, 95%CI[-0.68, -0.12], $p = .009$). Their

interpretation was that the effects of anodal and cathodal tDCS were anti-correlated: those

participants that improve their performance (i.e., show a smaller AB) due to anodal tDCS

tend to worsen due to cathodal tDCS, and vice versa.

We ran the same partial correlation (attempting to adjust for Session order) between

the anodal and cathodal AB magnitude change scores (*tDCS* - *baseline*) on our data (Figure

4). However, here the resulting correlation did not go in the same direction ($r(37)$ = .02,

95%CI[-0.30, 0.33]), and was not significant ($p = .899$). Or, adopting the terminology

suggested by LeBel, Vanpaemel, Cheung, and Campbell (2019): we obtained *no signal* (the

95% confidence interval includes 0, and therefore the correlation is not significant), and our

result is *inconsistent* (the 95% confidence interval excludes the point estimate of the

correlation in London and Slagter (2021)). In the next sections, we present a number of

follow-up analyses that further explore this difference in results between both studies (see

the Replication analyses subsection in the Methods section).

Is the correlation in study 2 significantly small?   London and Slagter (2021)

found a medium- to large correlation ($r = -.45$), but the correlation we find here is much

smaller ($r = .02$). We use equivalence testing (Lakens et al. (2018); see the Equivalence tests

subsection in the Methods section) to assess whether this correlation is significantly smaller

than a smallest effect size of interest (SESOI). Following the "small telescopes" approach

Table 3

Repeated measures ANOVA on T1 accuracy

Effect	F	df_1^{GG}	df_2^{GG}	p	$\hat{\eta}_G^2$
Session order	0.96	1	38	.332	.018
Block	6.64	1.89	71.87	.003	.010
Stimulation	0.00	1	38	.996	.000
Lag	29.23	1	38	< .001	.013
Session order × Block	0.60	1.89	71.87	.540	.001
Session order × Stimulation	11.24	1	38	.002	.030
Session order × Lag	0.04	1	38	.844	.000
Block × Stimulation	0.24	1.92	73.08	.777	.000
Block × Lag	1.91	1.93	73.20	.158	.001
Stimulation × Lag	0.31	1	38	.584	.000
Session order × Block × Stimulation	9.94	1.92	73.08	< .001	.007
Session order × Block × Lag	0.19	1.93	73.20	.821	.000
Session order × Stimulation × Lag	0.96	1	38	.333	.000
Block × Stimulation × Lag	0.31	1.86	70.75	.718	.000
Session order × Block × Stimulation × Lag	0.53	1.86	70.75	.580	.000

Figure 4. **The effects of anodal and cathodal tDCS are not correlated in the present study.** Data points show AB magnitude change scores (*tDCS - baseline*, *post - baseline*) for each participant in the study, in the *anodal* session (x-axis) and the *cathodal* session (y-axis). While London and Slagter (2021) found a negative partial correlation (for *tDCS - baseline*), suggesting opposite effects of anodal and cathodal tDCS, this effect appears to be absent here. The partial correlation coefficients (attempting to adjust for Session order) and p-values are printed in the upper left.

(Simonsohn, 2015), we set the SESOI to $r_{33\%}$, the correlation London and Slagter (2021) had 33% power to detect. Given their sample size of 34 participants, $r_{33\%} = .27$.

An inferiority test shows that the correlation here is significantly less negative than $-r_{33\%}$ ($p = .038$) (Figure 6), although only by a small margin. The effect is therefore "statistically inferior": the correlation does not exceed the lower equivalence bound ($-r_{33\%} = -.27$). By this definition, the correlation is too small (i.e. not negative enough) to be considered meaningful, indicating that we did not successfully replicate London and Slagter (2021).

Is the correlation in this study (study 2) different from the previous study

(study 1)?   To evaluate whether the correlation in the present study (study 2) was to be

expected based on London and Slagter (2021) (study 1), we constructed a 95% *prediction*

interval (PI) (Spence & Stanley, 2016), using the correlation in London and Slagter (2021)

and the sample size of both studies (see the Prediction interval subsection of the Methods

section).

The 95% PI[-0.82, -0.05] around the original correlation ($r = $ -.45) is very wide, so

almost any negative correlation would fall within it. However, it does not include the

replication result ($r = $ .02) (Figure 6). Assuming the results of both studies differ only due

to sampling error, the correlation observed here has a 95% chance to fall within the interval.

This means the correlation in our replication study is inconsistent with the correlation in

London and Slagter (2021), and thus that the replication should not be considered successful.

Another approach to quantify the difference between study 1 and 2 is to construct a

replication Bayes Factor (Wagenmakers et al., 2016). We use this Bayes factor to assess

evidence for H_0 : "the effects of anodal and cathodal tDCS are uncorrelated" relative to H_r :

"the effects of anodal and cathodal tDCS are anticorrelated as in London and Slagter (2021)"

(see the Replication Bayes factor subsection of the Methods section).

This replication Bayes factor expresses that the data are $BF_{0r} = 9.66$ times more likely

under H_0 than under H_r. This constitutes moderate to strong evidence that the effect is

absent vs. consistent with London and Slagter (2021), and thus that the effect did not

replicate.

Is the effect significant when combining study 1 and 2?   The sample size in

both study 1 (n = 34) and 2 (n = 40) is on the lower end. Therefore, we might get a more

accurate estimate of the size of the effect when combining both studies.

The meta-analytic estimate of the correlation is $r = $ -0.20, 95%CI[-0.42, 0.04], $p = $ .094

(Figure 6). So, when combining the correlation from London and Slagter (2021) and the

correlation observed here, the overall effect is no longer significant.

In addition, we pooled the data from both studies at the single-subject level and re-calculated the partial correlation on the combined sample (n = 74). To make the datasets as comparable as possible, we first averaged over T2|T1 accuracy at lags 2 and 4 in London and Slagter (2021), to mimic a "lag 3" condition cf. the present study (Figure 5).

Figure 5. **The effects of anodal and cathodal tDCS are also not correlated when pooling data from both studies.** As in Figure 4, data points show AB magnitude change scores (*tDCS - baseline*) in the *anodal* session (x-axis) and the *cathodal* session (y-axis), but now for each participant in London and Slagter (2021) (*study 1*) and the present study (*study 2*). While London and Slagter (2021) found a negative partial correlation, suggesting opposite effects of anodal and cathodal tDCS, this effect appears to be absent when based on the combined data of both studies. The partial correlation coefficient (attempting to adjust for Session order) is printed in the upper left.

The partial correlation based on the pooled data is $r(71) = -.18$. Thus, the correlation across a combination of both samples is a lot smaller than in London and Slagter (2021), and

460 slightly smaller still than the meta-analytic estimate that included the original correlation

461 from London and Slagter (2021) (Figure 6).

Figure 6. **Summary of all the replication analyses** (with exception of the replication Bayes Factor). The first two rows show the partial correlation (attemtping to adjust for Session order) between the AB magnitude change scores (tDCS - baseline) in the anodal and cathodal sessions, for London and Slagter (2021) and the present study. The first is significantly negative, the second is slightly positive and not significant, because its 95% confidence interval (CI) overlaps with zero. The third row shows the 90% CI around the correlation in the present study. Because this interval does not overlap with the "small telescopes" effect size, (indicated by the x: $-r_{33\%} = -.27$), this correlation is significantly smaller. The fourth row shows the 95% prediction interval (PI) around the correlation in London and Slagter (2021). Because this interval does not overlap with the correlation in the present study, both correlations are not consistent. The final two rows show the overall effect when the two correlations are meta-analyzed, and when one correlation is computed over the pooled data from both studies. Neither are significant (95% CI overlaps with zero). Thus, our replication analyses all suggest that we failed to replicate London & Slagter (2021), and when the results are examined in combination, no evidence in support of a negative relationship between the AB magnitude change scores (*tDCS - baseline*) in the anodal and cathodal sessions is obtained.

462 In sum, all the analyses we conducted seem to support the same conclusion: that the
463 current study is not a successful replication of London and Slagter (2021), and that therefore
464 the purported opposing effect of anodal and cathodal tDCS on the attentional blink within
465 individuals is called into question.

Discussion

London and Slagter (2021) were the first to uncover potential individual differences in
the effect of tDCS on the AB. They found a negative correlation between the effects of
anodal and cathodal tDCS on T2|T1 accuracy across individuals individuals (in the absence
of a group-level effect of tDCS). This analysis suggested an interesting pattern of individual
differences in response to tDCS: those that tended to benefit from anodal tDCS (i.e., whose
AB became smaller compared to baseline) would tend to worsen during cathodal tDCS, and
vice versa. This finding seemed plausible given the substantial and well-documented
individual differences in both the AB (Willems & Martens, 2016) and the effects of tDCS
(Krause & Cohen Kadosh, 2014).

We conducted a replication study, and again found no effect of tDCS at the group level.
However, in contrast to London and Slagter (2021), the correlation between the AB
magnitude change scores (tDCS - baseline) in the anodal and cathodal sessions here is not
significant, and not in the same direction. We therefore also computed several statistical
measures of replication success focused on the negative correlation between anodal and
cathodal tDCS. As discussed in more detail below, the field of electrical brain stimulation is
characterized by inconsistent findings across studies. It is hence important that researchers
in this field also start to incorporate methods that permit quantification of replication
success. It is our hope that the overview of main, complementary analyses that researchers
can use to maximize the evidential value from negative findings, provided here, will serve as
a guideline for future studies.

The measures of replication success showed that in our study the observed correlation

is smaller than in London and Slagter (2021), and than the smallest correlation they could

have reasonably detected (i.e., an equivalence test for the lower bound of $r_{33\%}$ (Simonsohn,

2015) was significant). The difference between the two studies is greater than expected based

on sampling error alone (i.e., the correlation in the present study falls outside of the 95%

prediction interval). Further, the data provide moderate to strong evidence for the null

hypothesis of zero correlation compared to the alternative that the correlation is as in

London and Slagter (2021) (i.e., we found a replication Bayes factor in favor of the null

hypothesis of ~10). Finally, combining both studies yields a smaller and non-significant

correlation (both in a meta-analysis and by pooling the data).

The overall picture is consistent: all measures indicate that the present study is not a

successful replication of London and Slagter (2021). But because both studies were similar in

design and sample size, it is not warranted to dismiss the findings of London and Slagter

(2021). Our analysis revealed substantial variation between both estimates of the effect, and

so the conclusion whether tDCS is effective should not be based on any single study, nor on

only a single replication study (Hedges & Schauer, 2019). In any series of studies, one should

expect to sometimes not find a significant result, purely due to random variation—even if

there is a true effect (Amrhein, Trafimow, & Greenland, 2019).

Still, the marked difference in results of both studies is surprising: they are similar

enough that the present study could be considered a direct replication of London and Slagter

(2021; Zwaan et al., 2018). We used the exact same electrode montage and tDCS parameters,

followed the same experimental design (Figure 1), ran the experiment in the same location

with the same participant population, and used a virtually identical task. Nevertheless, there

are some discrepancies between the studies that could have contributed to the different

outcome.

The most important difference is that London and Slagter (2021) presented T2 at lags

2, 4, and 10, whereas we used lags 3 and 8. This means that AB magnitude (the difference in T2|T1 accuracy between the shortest and longest lag) was on average smaller in the present study. We introduced the change for precisely this reason: to have about as many trials for the EEG analyses in which T2 was seen vs. missed, we needed the AB to be smaller. But this increase in average T2|T1 accuracy may have also reduced the between-subject variability that is essential for analyses that capitalize on individual differences. Indeed, from Figure 5 it seems that the sample in London and Slagter (2021) had a larger spread, at least in the change scores for AB magnitude.

Though the change in lags is probably the most important, the concurrent EEG measurement did introduce other differences as well. Each session took longer to complete, because the EEG setup required extra time, and the pace of the task was slowed down with longer inter-stimulus intervals. Finally, the current flow could have changed due to the presence of the EEG cap and electrodes, as well as the use of conductive paste instead of saline solution.

At the group level, our results are in agreement with London and Slagter (2021): we find no differential effects of anodal and cathodal tDCS on the AB. However, another study (Sdoia et al., 2019) did report a group-level effect of tDCS on the AB, but this study used different stimulation parameters and was fairly small (see London and Slagter (2021) for a more extensive discussion of the differences). At least with our AB task and fairly standard electrode montage and stimulation parameters (Santarnecchi et al., 2015), the AB does not appear to be very amenable to tDCS over the lDLPFC. Our null results are consistent with recent reviews and meta-analyses highlighting there is little evidence that prefrontal tDCS can be used to enhance cognition (Medina & Cason, 2017); or if so, that its effects on attention (Reteig, Talsma, Schouwenburg, & Slagter, 2017) and cognition more generally (Tremblay et al., 2014) are difficult to predict, rather small (Dedoncker, Brunoni, Baeken, & Vanderhasselt, 2016), and restricted to a limited subset of outcome measures and stimulation parameters (Boer et al., 2021; Imburgio & Orr, 2018).

Interpreting null results is always difficult, especially in brain stimulation studies

(Graaf & Sack, 2018). Ultimately, a myriad of possible underlying explanations may apply,

most of which we have no direct access to. For one, we cannot be sure that the current

density in the lDLPFC reached sufficient levels (Kim et al., 2014; Opitz, Paulus, Will,

Antunes, & Thielscher, 2015) based on our montage alone. More precise targeting of the

stimulation towards the lDPLFC (Datta et al., 2009) and modeling of the current flow

(Berker, Bikson, & Bestmann, 2013; Karabanov, Saturnino, Thielscher, & Siebner, 2019)

could provide some more confidence. Still, it would be difficult to verify that anodal tDCS

and cathodal tDCS indeed had an opposite effect on cortical excitability (Bestmann, de

Berker, & Bonaiuto, 2015; Bestmann & Walsh, 2017). This assumption holds in many cases,

and would provide a plausible explanation for the negative correlation in London and Slagter

(2021; Krause and Cohen Kadosh, 2014). But ideally, it should be validated with direct

measures of cortical excitability. Such measures are hard to obtain, although some recent

studies suggest that combining tDCS with magnetic resonance spectroscopy can be used for

this purpose (Antonenko et al., 2019; H. L. Filmer et al., 2019; Talsma, Broekhuizen,

Huisman, & Slagter, 2018). Combining tDCS with MRI may also help predict individual

differences in stimulation efficacy. One recent study found that individual variability in the

cortical thickness of left prefrontal cortex accounted for almost 35% of the variance in

stimulation efficacy among subjects (Hannah L. Filmer et al., 2019).

It is also possible that we do not find an effect because the studies did not have

sufficient statistical power (Minarik et al., 2016). Especially for the individual differences

analyses, the sample size in both studies is on the lower end. For example, to detect a

medium-sized correlation ($r = 0.3$) with 80% power, the required sample size is n = 84. We

do approach this sample size when combining both studies (n = 74). Also, the correlation in

London and Slagter (2021) was larger ($r = $ -0.45), but we cannot have much confidence in

the precision of this estimate: : the 95% confidence interval is so wide (-0.68 to -0.12) that it

is consistent with a large range of effect sizes, and therefore almost unfalsifiable (Morey &

Lakens, 2016). To estimate the size of a medium correlation ($r = 0.3$) within a narrower confidence band ($\pm$ 0.15 with 90% confidence), a sample size of n = 134 would be required (Schönbrodt & Perugini, 2013). Although our analyses suggest that the correlation is small if anything, this means we cannot accurately estimate how small—even with the combined sample.

These are all decisions to be made at the design stage, which can increase the value of a null result (Graaf & Sack, 2018). However, after the data are in, additional tools are available to increase the value of a null result (Harms & Lakens, 2018), especially in the case of a replication study (Simonsohn, 2015), which we have demonstrated here.

We hope our paper provides inspiration for others in the fields of brain stimulation and cognitive neuroscience. Many speak of a crisis of confidence (Héroux et al., 2017) and fear that the current literature is lacking in evidential value (Medina & Cason, 2017). This is certainly not a unique development, as these sentiments (Baker, 2015) and low rates of replication abound in many research areas (Camerer et al., 2018; Klein et al., 2018; Open Science Collaboration, 2015). But it is perhaps aggravated by the fact that the brain stimulation field has not matured yet (Parkin, Ekhtiari, & Walsh, 2015). At the same time, this also leaves room for the field to improve, and more work is necessary to determine the potential of electrical brain stimulation (Filmer, Mattingley, & Dux, 2020).

To make sure the literature accurately reflects the efficacy of the technique, it is crucial to combat publication bias. Positive results are heavily over-represented in most of the scientific literature, (Fanelli, 2012; Ferguson & Heene, 2012; Franco, Malhotra, & Simonovits, 2014), which has recently prompted a call to brain stimulation researchers to also publish their null results.[6] Preregistration and registered reports are other important methods to combat publication bias that the field of electrical brain stimulation should embrace (see e.g.,

---

[6] Research Topic in Frontiers, "Non-Invasive Brain Stimulation Effects on Cognition and Brain Activity: Positive Lessons from Negative Findings": https://www.frontiersin.org/research-topics/5535/non-invasive-brain-stimulation-effects-on-cognition-and-brain-activity-positive-lessons-from-negativ

Horne et al. (2021)). Publishing null results and replication studies—and making the most of their interpretation—is crucial to better this situation.

Acknowledgements

This work was supported by a Research Talent Grant from the Netherlands Organization for Scientific Research (NWO) (452-10-018, to HAS and KRR). We thank Raquel London for sharing all her experience and materials, as well as Daphne Box and Esther van der Giessen for their assistance in data collection.

References

Amrhein, V., Trafimow, D., & Greenland, S. (2019). Inferential Statistics as Descriptive Statistics: There Is No Replication Crisis if We Don't Expect Replication. *The American Statistician*, *73*(sup1), 262–270. https://doi.org/10.1080/00031305.2018.1543137

Anderson, S. F., & Maxwell, S. E. (2016). There's more than one way to conduct a replication study: Beyond statistical significance. *Psychological Methods*, *21*(1), 1–12. https://doi.org/10.1037/met0000051

Antonenko, D., Thielscher, A., Saturnino, G., Aydin, S., Ittermann, B., Grittner, U., & Flöel, A. (2019). Towards precise brain stimulation: Is electric field simulation related to neuromodulation? *Brain Stimulation.* https://doi.org/10.1016/j.brs.2019.03.072

Arasanz, C. P., Staines, W. R., & Schweizer, T. A. (2012). Isolating a cerebellar contribution to rapid visual attention using transcranial magnetic stimulation. *Frontiers in Behavioral Neuroscience*, *6*, 55. https://doi.org/10.3389/fnbeh.2012.00055

Aust, F., & Barth, M. (2020). *papaja: Create APA manuscripts with R Markdown.* Retrieved from https://github.com/crsh/papaja

Bakeman, R. (2005). Recommended effect size statistics for repeated measures designs. *Behavior Research Methods*, *37*(3), 379–384. https://doi.org/10.3758/BF03192707

Baker, M. (2015). 1,500 scientists lift the lid on reproducibility. *Nature*, *533*, 52–454.

Berker, A. O. de, Bikson, M., & Bestmann, S. (2013). Predicting the behavioral impact of transcranial direct current stimulation: issues and limitations. *Frontiers in Human Neuroscience*, *7*(October), 613. https://doi.org/10.3389/fnhum.2013.00613

Bestmann, S., de Berker, A. O., & Bonaiuto, J. (2015). Understanding the behavioural consequences of noninvasive brain stimulation. *Trends in Cognitive Sciences*, *19*(1), 13–20. https://doi.org/10.1016/j.tics.2014.10.003

Bestmann, S., & Walsh, V. (2017). Transcranial electrical stimulation. *Current Biology*, *27*(23), R1258–R1262. https://doi.org/10.1016/j.cub.2017.11.001

Bikson, M., Paulus, W., Esmaeilpour, Z., Kronberg, G., & Nitsche, M. A. (2019). Mechanisms of acute and after effects of transcranial direct current stimulation. In H. Knotkova, M. A. Nitsche, M. Bikson, & A. J. Woods (Eds.), *Practical guide to transcranial direct current stimulation* (pp. 81–113). Cham (Switzerland): Springer. https://doi.org/10.1007/978-3-319-95948-1

Boer, N. S. de, Schluter, R. S., Daams, J. G., Werf, Y. D. van der, Goudriaan, A. E., & Holst, R. J. van. (2021). The effect of non-invasive brain stimulation on executive functioning in healthy controls: A systematic review and meta-analysis. *Neuroscience & Biobehavioral Reviews*, *125*, 122–147. https://doi.org/10.1016/j.neubiorev.2021.01.013

Brandt, M. J., IJzerman, H., Dijksterhuis, A., Farach, F. J., Geller, J., Giner-Sorolla, R., . . . Veer, A. van 't. (2014). The Replication Recipe: What makes for a convincing replication? *Journal of Experimental Social Psychology*, *50*, 217–224. https://doi.org/10.1016/j.jesp.2013.10.005

Camerer, C. F., Dreber, A., Holzmeister, F., Ho, T. H., Huber, J., Johannesson, M., . . . Wu, H. (2018). Evaluating the replicability of social science experiments in Nature and Science between 2010 and 2015. *Nature Human Behaviour*, *2*(9), 637–644. https://doi.org/10.1038/s41562-018-0399-z

Champely, S. (2020). *Pwr: Basic functions for power analysis.* Retrieved from https://CRAN.R-project.org/package=pwr

Cooper, A. C., Humphreys, G. W., Hulleman, J., Praamstra, P., & Georgeson, M. (2004). Transcranial magnetic stimulation to right parietal cortex modifies the attentional blink. *Experimental Brain Research*, *155*(1), 24–29. https://doi.org/10.1007/s00221-003-1697-9

Datta, A., Bansal, V., Diaz, J., Patel, J., Reato, D., & Bikson, M. (2009). Gyri-precise head model of transcranial direct current stimulation: Improved spatial focality using a ring electrode versus conventional rectangular pad. *Brain Stimulation*, *2*(4), 201–207.e1. https://doi.org/10.1016/j.brs.2009.03.005

Dayan, E., Censor, N., Buch, E. R., Sandrini, M., & Cohen, L. G. (2013). Noninvasive brain stimulation: from physiology to network dynamics and back. *Nature Neuroscience*, *16*(7), 838–844. https://doi.org/10.1038/nn.3422

Dedoncker, J., Brunoni, A. R., Baeken, C., & Vanderhasselt, M. A. (2016). A systematic review and meta-analysis of the effects of transcranial direct current stimulation (tDCS) over the dorsolateral prefrontal cortex in healthy and neuropsychiatric samples: influence of stimulation parameters. *Brain Stimulation*, *9*(4), 501–517. https://doi.org/10.1016/j.brs.2016.04.006

Dux, P. E., & Marois, R. (2008). Distractor inhibition predicts individual differences in the attentional blink. *PloS One*, *3*(10), e3330. https://doi.org/10.1371/journal.pone.0003330

Dux, P. E., & Marois, R. (2009). The attentional blink: A review of data and theory.

Attention, Perception, and Psychophysics, 71(8), 1683–1700.

https://doi.org/10.3758/APP.71.8.1683

Esterman, M., Thai, M., Okabe, H., DeGutis, J., Saad, E., Laganiere, S. E., & Halko, M. A.

(2017). Network-targeted cerebellar transcranial magnetic stimulation improves

attentional control. *NeuroImage, 156*(May), 190–198.

https://doi.org/10.1016/j.neuroimage.2017.05.011

Fanelli, D. (2012). Negative results are disappearing from most disciplines and countries.

Scientometrics, 90(3), 891–904. https://doi.org/10.1007/s11192-011-0494-7

Ferguson, C. J., & Heene, M. (2012). A vast graveyard of undead theories: Publication bias

and psychological science's aversion to the null. *Perspectives on Psychological Science,*

7(6), 555–561. https://doi.org/10.1177/1745691612459059

Filmer, H. L., Ehrhardt, S., Bollmann, S., Mattingley, J. B., & Dux, P. E. (2019).

Accounting for individual differences in the response to tDCS with baseline levels of

neurochemical excitability. *Cortex, 115*, 324–334.

https://doi.org/10.1016/j.cortex.2019.02.012

Filmer, H. L., Ehrhardt, S. E., Shaw, T. B., Mattingley, J. B., & Dux, P. E. (2019). The

efficacy of transcranial direct current stimulation to prefrontal areas is related to

underlying cortical morphology. *NeuroImage, 196*, 41–48.

https://doi.org/10.1016/j.neuroimage.2019.04.026

Filmer, H. L., Mattingley, J. B., & Dux, P. E. (2020). Modulating brain activity and

behaviour with tDCS: Rumours of its death have been greatly exaggerated. *Cortex,*

123, 141–151. https://doi.org/10.1016/j.cortex.2019.10.006

Fletcher, T. D. (2010). *Psychometric: Applied psychometric theory.* Retrieved from

https://CRAN.R-project.org/package=psychometric

Franco, A., Malhotra, N., & Simonovits, G. (2014). Publication bias in the social sciences:

Unlocking the file drawer. *Science*, *345*(6203), 1502–1505.

https://doi.org/10.1126/science.1255484

Gebodh, N., Esmaeilpour, Z., Bikson, M., Adair, D., Fregni, F., & Schestattsky, P. (2019).

Transcranial direct current stimulation among technologies for low-intensity

transcranial electrical stimulation: classification, history, and terminology. In H.

Knotkova, M. A. Nitsche, M. Bikson, & A. J. Woods (Eds.), *Practical guide to*

transcranial direct current stimulation (pp. 3–43). Cham (Switzerland): Springer.

https://doi.org/10.1007/978-3-319-95948-1_1

Goh, J. X., Hall, J. A., & Rosenthal, R. (2016). Mini Meta-Analysis of Your Own Studies:

Some Arguments on Why and a Primer on How. *Social and Personality Psychology*

Compass, *10*(10), 535–549. https://doi.org/10.1111/spc3.12267

Gorgolewski, K. J., Auer, T., Calhoun, V. D., Craddock, R. C., Das, S., Duff, E. P., . . .

Poldrack, R. A. (2016). The brain imaging data structure, a format for organizing

and describing outputs of neuroimaging experiments. *Scientific Data*, *3*, 160044.

https://doi.org/10.1038/sdata.2016.44

Graaf, T. A. de, & Sack, A. T. (2018). When and how to interpret null Results in NIBS: A

taxonomy based on prior expectations and experimental design. *Frontiers in*

Neuroscience, *12*, 915. https://doi.org/10.3389/fnins.2018.00915

Harms, C., & Lakens, D. (2018). Making 'null effects' informative: statistical techniques and

inferential frameworks. *Journal of Clinical and Translational Research*, *3*, 382–393.

https://doi.org/10.18053/jctres.03.2017s2.007

Harty, S., Sella, F., & Cohen Kadosh, R. (2017). Transcranial Electrical Stimulation and

Behavioral Change: The Intermediary Influence of the Brain. *Frontiers in Human*

Neuroscience, *11*(March), 1–5. https://doi.org/10.3389/fnhum.2017.00112

Hedges, L. V., & Schauer, J. M. (2019). More Than One Replication Study Is Needed for

Unambiguous Tests of Replication. *Journal of Educational and Behavioral Statistics*,

44(5), 543–570. https://doi.org/10.3102/1076998619852953

Héroux, M. E., Loo, C. K., Taylor, J. L., & Gandevia, S. C. (2017). Questionable science

and reproducibility in electrical brain stimulation research. *PLoS ONE*, *12*(4),

e0175635. https://doi.org/10.1371/journal.pone.0175635

Hommel, B., Kessler, K., Schmitz, F., Gross, J., Akyürek, E., Shapiro, K., & Schnitzler, A.

(2006). How the brain blinks: towards a neurocognitive model of the attentional blink.

Psychological Research, *70*(6), 425–435. https://doi.org/10.1007/s00426-005-0009-3

Horne, K. S., Filmer, H. L., Nott, Z. E., Hawi, Z., Pugsley, K., Mattingley, J. B., & Dux, P.

E. (2021). Evidence against benefits from cognitive training and transcranial direct

current stimulation in healthy older adults. *Nature Human Behaviour*, *5*(1), 146–158.

https://doi.org/10.1038/s41562-020-00979-5

Huber, D. E., Potter, K. W., & Huszar, L. D. (2019). Less "story" and more "reliability" in

cognitive neuroscience. *Cortex*. https://doi.org/10.1016/j.cortex.2018.10.030

Imburgio, M. J., & Orr, J. M. (2018). Effects of prefrontal tDCS on executive function:

Methodological considerations revealed by meta-analysis. *Neuropsychologia*, *117*,

156–166. https://doi.org/10.1016/j.neuropsychologia.2018.04.022

Karabanov, A. N., Saturnino, G. B., Thielscher, A., & Siebner, H. R. (2019). Can

transcranial electrical stimulation localize brain function? *Frontiers in Psychology*, *10*,

213. https://doi.org/10.3389/fpsyg.2019.00213

Kim, J.-H., Kim, D.-W., Chang, W. H., Kim, Y.-H., Kim, K., & Im, C.-H. (2014).

Inconsistent outcomes of transcranial direct current stimulation may originate from

anatomical differences among individuals: electric field simulation using individual

MRI data. *Neuroscience Letters*, *564*, 6–10.

https://doi.org/10.1016/j.neulet.2014.01.054

Klein, R. A., Vianello, M., Hasselman, F., Adams, B. G., Adams, R. B., Alper, S., . . .

Nosek, B. A. (2018). Many Labs 2: Investigating variation in replicability across samples and settings. *Advances in Methods and Practices in Psychological Science*, *1*(4), 443–490. https://doi.org/10.1177/2515245918810225

Krause, B., & Cohen Kadosh, R. (2014). Not all brains are created equal: the relevance of individual differences in responsiveness to transcranial electrical stimulation. *Frontiers in Systems Neuroscience*, *8*(February), 25. https://doi.org/10.3389/fnsys.2014.00025

Lakens, D. (2014). Performing high-powered studies efficiently with sequential analyses. *European Journal of Social Psychology*, *44*(7), 701–710. https://doi.org/10.1002/ejsp.2023

Lakens, D. (2017). Equivalence tests: A practical primer for t-tests, correlations, and meta-analyses. *Social Psychological and Personality Science*, *1*, 1–8. https://doi.org/10.1177/1948550617697177

Lakens, D., McLatchie, N., Isager, P. M., Scheel, A. M., & Dienes, Z. (2020). Improving Inferences About Null Effects With Bayes Factors and Equivalence Tests. *The Journals of Gerontology: Series B*, *75*(1), 45–57. https://doi.org/10.1093/geronb/gby065

Lakens, D., Scheel, A. M., & Isager, P. M. (2018). Equivalence Testing for Psychological Research: A Tutorial. *Advances in Methods and Practices in Psychological Science*, *1*(2), 259–269. https://doi.org/10.1177/2515245918770963

LeBel, E. P., Vanpaemel, W., Cheung, I., & Campbell, L. (2019). A Brief Guide to Evaluate Replications. *Meta-Psychology*, *3*. https://doi.org/10.15626/MP.2018.843

Lenth, R. V. (2021). *Emmeans: Estimated marginal means, aka least-squares means.* Retrieved from https://CRAN.R-project.org/package=emmeans

Li, L. M., Uehara, K., & Hanakawa, T. (2015). The contribution of interindividual factors to variability of response in transcranial direct current stimulation studies. *Frontiers in*

Cellular Neuroscience, *9*(May), 181. https://doi.org/10.3389/fncel.2015.00181

Linde, M., Tendeiro, J., Selker, R., Wagenmakers, E.-J., & Ravenzwaaij, D. van. (2020). Decisions About Equivalence: A Comparison of TOST, HDI-ROPE, and the Bayes Factor. *PsyArXiv.* https://doi.org/10.31234/osf.io/bh8vu

Liu, A., Vöröslakos, M., Kronberg, G., Henin, S., Krause, M. R., Huang, Y., . . . Buzsáki, G. (2018). Immediate neurophysiological effects of transcranial electrical stimulation. *Nature Communications*, *9*(1), 5092. https://doi.org/10.1038/s41467-018-07233-7

London, R. E., & Slagter, H. A. (2021). No Effect of Transcranial Direct Current Stimulation over Left Dorsolateral Prefrontal Cortex on Temporal Attention. *Journal of Cognitive Neuroscience*, *33*(4), 756–768. https://doi.org/10.1162/jocn_a_01679

MacLean, M. H., & Arnell, K. M. (2012). A conceptual and methodological framework for measuring and modulating the attentional blink. *Attention, Perception, and Psychophysics*, *74*(6), 1080–1097. https://doi.org/10.3758/s13414-012-0338-4

Marchetti, G. M., Drton, M., & Sadeghi, K. (2020). *Ggm: Graphical markov models with mixed graphs.* Retrieved from https://CRAN.R-project.org/package=ggm

Martens, S., & Wyble, B. (2010). The attentional blink: Past, present, and future of a blind spot in perceptual awareness. *Neuroscience and Biobehavioral Reviews*, *34*(6), 947–957. https://doi.org/10.1016/j.neubiorev.2009.12.005

Medina, J., & Cason, S. (2017). No evidential value in samples of transcranial direct current stimulation (tDCS) studies of cognition and working memory in healthy populations. *Cortex*, *94*, 131–141. https://doi.org/10.1016/j.cortex.2017.06.021

Minarik, T., Berger, B., Althaus, L., Bader, V., Biebl, B., Brotzeller, F., . . . Sauseng, P. (2016). The importance of sample size for reproducibility of tDCS effects. *Frontiers in Human Neuroscience*, *10*(September), 453. https://doi.org/10.3389/FNHUM.2016.00453

Morey, R. D., & Rouder, J. N. (2018). *BayesFactor: Computation of bayes factors for common designs.* Retrieved from https://CRAN.R-project.org/package=BayesFactor

Morey, R., & Lakens, D. (2016). Why most of psychology is statistically unfalsifiable. https://doi.org/10.5281/zenodo.838685

Munafò, M. R., Nosek, B. A., Bishop, D. V. M., Button, K. S., Chambers, C. D., Percie du Sert, N., ... Ioannidis, J. P. A. (2017). A manifesto for reproducible science. *Nature Human Behaviour*, *1*(1), 0021. https://doi.org/10.1038/s41562-016-0021

Muradchanian, J., Hoekstra, R., Kiers, H., & Ravenzwaaij, D. van. (2020). *How Best to Quantify Replication Success? A Simulation Study on the Comparison of Replication Success Metrics.* MetaArXiv. https://doi.org/10.31222/osf.io/wvdjf

Müller, K. (2020). *Here: A simpler way to find your files.* Retrieved from https://CRAN.R-project.org/package=here

Nitsche, M. A., & Paulus, W. (2000). Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation. *The Journal of Physiology*, *527 Pt 3*(2000), 633–639. https://doi.org/10.1111/j.1469-7793.2000.t01-1-00633.x

Nitsche, M. A., & Paulus, W. (2001). Sustained excitability elevations induced by transcranial DC motor cortex stimulation in humans. *Neurology*, *57*(10), 1899–1901. https://doi.org/10.1212/WNL.57.10.1899

Olivers, C. N. L., Nieuwenhuis, S., Olivers, C. N. L., & Nieuwenhuis, S. (2005). The beneficial effect of concurrent task-irrelevant mental activity on temporal attention. *Psychological Science*, *16*(4), 265–269. https://doi.org/10.1111/j.0956-7976.2005.01526.x

Olivers, C. N., & Nieuwenhuis, S. (2006). The beneficial effects of additional task load, positive affect, and instruction on the attentional blink. *Journal of Experimental Psychology: Human Perception and Performance*, *32*(2), 364–379.

https://doi.org/10.1037/0096-1523.32.2.364

Open Science Collaboration. (2015). Estimating the reproducibility of psychological science. *Science*, *349*(6251), aac4716–1–aac4716–8. https://doi.org/10.1126/science.aac4716

Opitz, A., Paulus, W., Will, S., Antunes, A., & Thielscher, A. (2015). Determinants of the electric field during transcranial direct current stimulation. *NeuroImage*, *109*, 140–150. https://doi.org/10.1016/j.neuroimage.2015.01.033

Parkin, B. L., Bhandari, M., Glen, J. C., & Walsh, V. (2018). The physiological effects of transcranial electrical stimulation do not apply to parameters commonly used in studies of cognitive neuromodulation. *Neuropsychologia*, (September 2017), 1–8. https://doi.org/10.1016/j.neuropsychologia.2018.03.030

Parkin, B. L., Ekhtiari, H., & Walsh, V. F. (2015). Non-invasive human brain stimulation in cognitive neuroscience: a primer. *Neuron*, *87*(5), 932–945. https://doi.org/10.1016/j.neuron.2015.07.032

Patil, P., Peng, R. D., & Leek, J. T. (2016). What should researchers expect when they replicate studies? A statistical view of replicability in psychological science. *Perspectives on Psychological Science*, *11*(4), 539–544. https://doi.org/10.1177/1745691616646366

Pernet, C. R., Appelhoff, S., Gorgolewski, K. J., Flandin, G., Phillips, C., Delorme, A., & Oostenveld, R. (2019). EEG-BIDS, an extension to the brain imaging data structure for electroencephalography. *Scientific Data*, *6*, 103. https://doi.org/10.1038/s41597-019-0104-8

Quintana, D. S. (2015). From pre-registration to publication: A non-technical primer for conducting a meta-analysis to synthesize correlational data. *Frontiers in Psychology*, *6*, 1549. https://doi.org/10.3389/fpsyg.2015.01549

Raymond, J. E., Shapiro, K. L., & Arnell, K. M. (1992). Temporary suppression of visual

processing in an RSVP task: An attentional blink? *Journal of Experimental Psychology: Human Perception and Performance, 18*(3), 849–860.

R Core Team. (2019). *R: A language and environment for statistical computing.* Vienna, Austria: R Foundation for Statistical Computing. Retrieved from https://www.R-project.org/

Reteig, L. C., Talsma, L. J., Schouwenburg, M. R. van, & Slagter, H. A. (2017). Transcranial Electrical Stimulation as a Tool to Enhance Attention. *Journal of Cognitive Enhancement, 1*(1), 10–25. https://doi.org/10.1007/s41465-017-0010-y

Robinson, D., Hayes, A., & Couch, S. (2021). *Broom: Convert statistical objects into tidy tibbles.*

RStudio Team. (2016). RStudio: Integrated Development Environment for R. Boston, MA: RStudio, Inc. Retrieved from http://www.rstudio.com/

Santarnecchi, E., Brem, A.-K., Levenbaum, E., Thompson, T., Kadosh, R. C., & Pascual-Leone, A. (2015). Enhancing cognition using transcranial electrical stimulation. *Current Opinion in Behavioral Sciences, 4*, 171–178. https://doi.org/10.1016/j.cobeha.2015.06.003

Schönbrodt, F. D., & Perugini, M. (2013). At what sample size do correlations stabilize ? *Journal of Research in Personality, 47*, 609–612. https://doi.org/10.1016/j.jrp.2013.05.009

Sdoia, S., Conversi, D., Pecchinenda, A., & Ferlazzo, F. (2019). Access to consciousness of briefly presented visual events is modulated by transcranial direct current stimulation of left dorsolateral prefrontal cortex. *Scientific Reports, 9*(1), 10950. https://doi.org/10.1038/s41598-019-47527-4

Simmons, J. P., Nelson, L. D., & Simonsohn, U. (2011). False-positive psychology: Undisclosed flexibility in data collection and analysis allows presenting anything as

significant. *Psychological Science*, *22*(11), 1359–1366.

https://doi.org/10.1177/0956797611417632

Simonsohn, U. (2015). Small telescopes: Detectability and the evaluation of replication

results. *Social Science Research Network*, (January 2015), 1–11.

https://doi.org/http://dx.doi.org/10.2139/ssrn.2259879

Singmann, H., Bolker, B., Westfall, J., Aust, F., & Ben-Shachar, M. S. (2021). *Afex:*

Analysis of factorial experiments. Retrieved from

https://CRAN.R-project.org/package=afex

Slagter, H. A., & Georgopoulou, K. (2013). Distractor inhibition predicts individual

differences in recovery from the attentional blink. *PloS One*, *8*(5), e64681.

https://doi.org/10.1371/journal.pone.0064681

Slagter, H. A., Johnstone, T., Beets, I. A., & Davidson, R. J. (2010). Neural competition for

conscious representation across time: An fMRI study. *PLoS ONE*, *5*(5), e10556.

https://doi.org/10.1371/journal.pone.0010556

Slagter, H. A., Lutz, A., Greischar, L. L., Francis, A. D., Nieuwenhuis, S., Davis, J. M., &

Davidson, R. J. (2007). Mental training affects distribution of limited brain resources.

PLoS Biology, *5*(6), e138. https://doi.org/10.1371/journal.pbio.0050138

Spence, J. R., & Stanley, D. J. (2016). Prediction interval: What to expect when you're

expecting . . . a replication. *PLOS ONE*, *11*(9), e0162874.

https://doi.org/10.1371/journal.pone.0162874

Stagg, C. J., Antal, A., & Nitsche, M. A. (2018). Physiology of transcranial direct current

stimulation. *Journal of ECT*, *34*(3), 144–152.

https://doi.org/10.1097/YCT.0000000000000510

Stanley, D. (2016). *PredictionInterval: Prediction interval functions for assessing replication*

study results. Retrieved from

https://CRAN.R-project.org/package=predictionInterval

Talsma, L. J., Broekhuizen, J. A., Huisman, J., & Slagter, H. A. (2018). No evidence that baseline prefrontal cortical excitability (3T-MRS) predicts the effects of prefrontal tDCS on WM performance. *Frontiers in Neuroscience*, *12*, 481. https://doi.org/10.3389/fnins.2018.00481

Thomson, D. R., Ralph, B. C. W., Besner, D., & Smilek, D. (2015). The more your mind wanders, the smaller your attentional blink: An individual differences study. *Quarterly Journal of Experimental Psychology*, *68*(1), 181–191. https://doi.org/10.1080/17470218.2014.940985

Tremblay, S., Lepage, J.-F., Latulipe-Loiselle, A., Fregni, F., Pascual-Leone, A., & Théoret, H. (2014). The uncertain outcome of prefrontal tDCS. *Brain Stimulation*, *7*(6), 773–783. https://doi.org/10.1016/j.brs.2014.10.003

Ueno, T., Fastrich, G. M., & Murayama, K. (2016). Meta-analysis to integrate effect sizes within an article: Possible misuse and Type I error inflation. *Journal of Experimental Psychology: General*, *145*(5), 643–654. https://doi.org/10.1037/xge0000159

Verhagen, J., & Wagenmakers, E. J. (2014). Bayesian tests to quantify the result of a replication attempt. *Journal of Experimental Psychology: General*, *143*(4), 1457–1475. https://doi.org/10.1037/a0036731

Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software*, *36*(3), 1–48. Retrieved from https://www.jstatsoft.org/v36/i03/

Wagenmakers, E. J., Love, J., Marsman, M., Jamil, T., Ly, A., Verhagen, J., . . . Morey, R. D. (2018). Bayesian inference for psychology. Part II: Example applications with JASP. *Psychonomic Bulletin and Review*, *25*(1), 58–76. https://doi.org/10.3758/s13423-017-1323-7

Wagenmakers, E. J., Marsman, M., Jamil, T., Ly, A., Verhagen, J., Love, J., . . . Morey, R. D. (2018). Bayesian inference for psychology. Part I: Theoretical advantages and practical ramifications. *Psychonomic Bulletin and Review, 25*(1), 35–57. https://doi.org/10.3758/s13423-017-1343-3

Wagenmakers, E.-J., Verhagen, J., & Ly, A. (2016). How to quantify the evidence for the absence of a correlation. *Behavior Research Methods, 48*, 413–426. https://doi.org/10.3758/s13428-015-0593-0

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., . . . Yutani, H. (2019). Welcome to the tidyverse. *Journal of Open Source Software, 4*(43), 1686. https://doi.org/10.21105/joss.01686

Wilke, C. O. (2020). *Cowplot: Streamlined plot theme and plot annotations for 'ggplot2'*. Retrieved from https://CRAN.R-project.org/package=cowplot

Willems, C., & Martens, S. (2016). Time to see the bigger picture: Individual differences in the attentional blink. *Psychonomic Bulletin and Review, 23*(5), 1289–1299. https://doi.org/10.3758/s13423-015-0977-2

Xie, Y. (2015). *Dynamic documents with R and knitr* (2nd ed.). Boca Raton, Florida: Chapman; Hall/CRC. Retrieved from https://yihui.org/knitr/

Xie, Y., Allaire, J. J., & Grolemund, G. (2018). *R Markdown: The Definitive Guide.* Boca Raton, FL: CRC Press. https://doi.org/10.1016/B978-0-12-814447-3.00041-0

Zhu, H. (2021). *KableExtra: Construct complex table with 'kable' and pipe syntax.* Retrieved from https://CRAN.R-project.org/package=kableExtra

Zondervan-Zwijnenburg, M., & Rijshouwer, D. (2020). Testing Replication with Small Samples. In R. van de Schoot & M. Miočević (Eds.), *Small sample size solutions: A guide for applied researchers and practitioners.* Abingdon, Oxon ; New York, NY: Routledge.

941   Zwaan, R. A., Etz, A., Lucas, R. E., & Donnellan, M. B. (2018). Making replication

942   mainstream. *Behavioral and Brain Sciences, e120.*

943   https://doi.org/10.1017/S0140525X17001972

944   **Supplementary material**

945 **tDCS adverse events**

Figure S1. **tDCS adverse events**. Number of reports out of 89 sessions (either anodal or cathodal tDCS). Top row shows intensity ratings [*little, moderate, strong, very strong*]; bottom row shows participant's confidence that event was related to tDCS [*unlikely, possibly, likely, very likely*]. Adverse events are sorted in descending order of number of reports (for very rare events (five reports or fewer for a given polarity), some text counts have been removed to prevent overlap).

Table S1

Number of reports of tDCS side effects

	Intensity rating					Confidence rating				
	none	a little	moderate	strong	very strong	n/a	unlikely	possibly	likely	very likely
anodal session										
burning	23	11	8	4	0	23	0	2	7	14
dizziness	43	3	0	0	0	43	1	2	0	0
fatigue	22	9	12	2	1	24	13	8	1	0
headache	32	9	3	1	1	34	1	10	1	0
itching	18	17	7	4	0	18	1	4	8	15
nausea	43	2	1	0	0	43	1	1	1	0
pain	41	3	2	0	0	40	1	0	4	1
tingling	11	21	9	5	0	11	0	4	12	19
cathodal session										
burning	26	9	5	2	1	25	0	1	7	10
dizziness	40	2	0	1	0	39	0	2	1	1
fatigue	15	13	6	8	1	18	12	9	4	0
headache	30	7	4	2	0	31	0	8	2	2
itching	18	14	9	2	0	18	0	0	15	10
nausea	41	2	0	0	0	40	0	2	0	1
pain	37	4	0	2	0	36	0	1	3	3
tingling	5	25	11	2	0	5	0	1	15	22

[1] Intensity rating: To which degree were the following sensations present during stimulation?

[2] Confidence rating: To which degree do you believe this was caused by the stimulation?