1

Soil protists can actively redistribute beneficial bacteria along *Medicago*

***truncatula* roots**

Christopher J. Hawxhurst[Δa], Jamie L. Micciulla[Δb], Charles M. Bridges[b], Mikhael Shor[c] , Daniel J.

Gage[b] and Leslie M. Shor[a,d,*]

[Δ]Co-first Authors

[a]Department of Chemical & Biomolecular Engineering, University of Connecticut, Storrs, CT,

USA

[b]Department of Molecular and Cellular Biology, University of Connecticut, Storrs, CT, USA

[c]Department of Economics, University of Connecticut, Storrs, CT, USA

[d]Center for Environmental Sciences & Engineering, University of Connecticut, Storrs, CT, USA

*Corresponding Author: leslieshor@gmail.com

19 Christopher J. Hawxhurst and Jamie L. Micciulla contributed equally to this work. Author order

20 was determined alphabetically.

22 Keywords: Protist, Soil, Plant Beneficial Bacteria, Transport

Abstract

The rhizosphere is the region of soil directly influenced by plant roots. The microbial community

in the rhizosphere includes fungi, protists, and bacteria, all of which play a significant role in

plant health. The beneficial bacterium *Sinorhizobium meliloti* infects growing root hairs on

nitrogen-starved leguminous plants. Infection leads to the formation of a root nodule, where *S.*

meliloti converts atmospheric nitrogen to ammonia, a usable form of nitrogen for plants. *S.*

meliloti is often found in biofilms and travels slowly along the roots, leaving developing root

hairs at the growing root tips uninfected. Soil protists are an important component of the

rhizosphere system who prey on soil bacteria and have been known to egest undigested

phagosomes. We show that the soil protist, *Colpoda sp.*, can transport *S. meliloti* down *Medicago*

truncatula roots. By using pseudo-3D soil microcosms, we directly observed the presence of

fluorescently labelled *S. meliloti* along *M. truncatula* roots and tracked the displacement of the

fluorescence signal over time. Two weeks after co-inoculation, this signal was detected 52 mm,

on average, farther down the roots when *Colpoda sp.* was also present compared with the

experimental treatment that contained bacteria but not protists. Direct counts also showed that

protists are required for viable bacteria to reach the deeper sections of root systems in our

microcosms. Facilitating bacterial transport may be an important mechanism whereby soil

protists promote plant health. As a sustainable agriculture biotechnology, protist-facilitated

transport has the potential to boost efficacy of bacterial inoculants, thereby helping growers

avoid overuse of nitrogen fertilizers and enhance performance of climate-smart, no-till farming

practices.

Importance

Soil protists are an important part of the microbial community in the rhizosphere. Plants

grown with protists fare better than plants grown without protists. Mechanisms through which

protists support plant health include nutrient cycling, alteration of the bacterial community

through selective feeding, and consumption of plant pathogens. Here we provide data in support

of an additional mechanism: protists act as transport vehicles for bacteria in soil. We show that

protist-facilitated transport can deliver plant-beneficial bacteria to the growing tips of roots that

may otherwise be sparsely inhabited with bacteria originating from a seed-associated inoculum.

By co-inoculating *Medicago truncatula* roots with both *S. meliloti*, a nitrogen fixing legume

symbiont, and *Colpoda sp.*, a ciliated protist, we show substantial and statistically significant

transport with depth and breadth of bacteria-associated fluorescence as well as transport of viable

bacteria. Co-inoculation with shelf-stable encysted soil protists may be employed as a

sustainable agriculture biotechnology to better distribute beneficial bacteria and enhance the

performance of inoculants.

Introduction

The rhizosphere is the zone of soil surrounding plant roots that is under the influence of the root. This soil fraction contains 5 to 100 times more organisms per unit volume than adjacent bulk soil (1). Up to 40% of carbon assimilated by plants is contributed back to the soil as rhizo-deposits (2). In return for these carbon-rich root exudates, plants often benefit from microbial activity through a variety of mechanisms such as moisture retention, pathogen suppression, plant-hormone synthesis and the release of recalcitrant nutrients (3, 4). Furthermore, some soil dwelling bacteria, such as those from the family Rhizobiaceae, beneficially infect leguminous roots and directly provide fixed nitrogen in exchange for TCA cycle intermediates.

The functioning of soil and rhizosphere systems emerges, in part, from biological communities interacting in a spatially structured environment (5–12). In trying to better understand the importance of biological factors on the properties of soil, research has been dedicated to the study and interaction of different bacterial species and plants in spatially-constrained systems (6, 9, 10). Protists form an additional, important, and often overlooked component of the soil system. The presence of protists in addition to bacteria can promote and enhance size and weight of plant roots and shoots versus the presence of bacteria alone (13). Protist-associated enhancement to plant health can arise through several different mechanisms. In the "microbial loop" hypothesis, protists recycle limiting nutrients by grazing on the microbes fed by root exudates (14–18). Protists may also alter the microbial composition of the rhizosphere through selective grazing (18, 19). A third potential mechanism, investigated here, is that protists may improve plant health by acting as a transport facilitator: directly relocating bacteria or other "cargo" along downward-growing roots (20–22). Similar examples include

transport of fungal spores hitchhiking on bacterial flagella (23). Other examples are given in

recent reviews (24, 25).

Plant growth-promoting rhizobacteria (PGPR) are beneficial soil bacteria used in

agriculture to improve crop yield and disease resistance. PGPR-mediated nutritional effects

include symbiotic nitrogen fixation, non-symbiotic nitrogen fixation, release of recalcitrant

nitrogen from soils, phosphate solubilization, and iron delivery (26–29). Commercially-used

PGPR include many species of nitrogen-fixing bacteria: *Azospirillum* spp., *Azotobacter* spp.,

Pseudomonas spp., *Bacillus subtilis* strains, *Bacillus megaterium* strains and *Trichoderma* spp. (a

fungal genus) (29). Many additional species are under investigation in laboratory, greenhouse,

and field trials, but have not yet been commercialized.

PGPR are typically inoculated onto seeds before planting or they are directly applied to

the soil during planting. Soil bacteria usually grow as biofilms and spread very slowly. Thus,

bacteria inoculated onto, or near, seeds often do not leave the area of inoculation and do not

establish a population on the root large enough to provide beneficial services (30–33). Further,

symbiotic infection of root hairs with nitrogen-fixing rhizobia, such as *S. meliloti*, often occurs at

root hairs located near root tips. Because roots grow only near their tips, suitable locations for

infection in a seedling are located further and further from the soil surface as a seedling

develops. A lack of bacteria at the root tip will most often result in decreased rate of symbiotic

infection and nodulation (34).

Soil protists have the potential to act as transport vehicles for bacteria. Soybean roots can

grow at ~3000 µm per hour (35). For comparison, the spreading rate of biocontrol *P. fluorescens*

biofilms is less than 10 µm per hour (36), while ciliated protists can swim at speeds up to 400

µm per second for short periods (37). We have previously shown that protists move

fluorescently-labeled microspheres and bacteria in emulated soil micromodels (20). In situ,

protists are known to carry bacteria, most commonly as intracellular parasites, for example when

Legionella spp. infect aquatic amoeba or *Campylobacter* spp. infect ciliates (38, 39). Also,

bacterial transport by protists has been demonstrated in the soil amoeba *Dictyostelium*

discoideum, which can farm bacteria and carry them to new habitats (40, 41). Although protists

graze on bacteria, ingestion is not always fatal (38). Further, bacteria have been observed

attached to the outer surfaces of protists, thus providing a means for transport that does not

require ingestion (20). We propose protists contribute to the rhizosphere community, in part,

through their capacity to move bacteria. The goal of this study was to test the hypothesis that soil

protists, despite grazing, can enhance dispersal of bacteria along root systems. We measured the

spatial distribution of fluorescently labelled *S. meliloti* through soil-containing mesocosms over

time in the presence and absence of a soil-isolated, ciliated, protist from the genus *Colpoda*. Our

methods included both nondestructive direct imaging repeated over time and direct counts of

viable bacteria and protists collected from root systems at the conclusion of each experiment.

Experimental Materials and Methods

Model soil. Sterile model soil was comprised of play sand (Quikrete) and fine vermiculite

combined 1:1 in a soil mixer. The mix was fractionated by sieving and the 125-250 µm fraction

and 63-125 µm fraction were combined 1:2 to emulate the particle sized distribution of a model

sandy loam (11).

Plants. *Medicago truncatula* is a model legume that offers uniform growth and natural symbiosis

with *S. meliloti*. To synchronize the plants, *M. truncatula* cv. Jemalong seeds were sterilized by

successive washing with 95% ethanol for 30 min, 10% bleach for 15 min, and sterile diH$_2$O for

15 min on a shaking platform at 80 rpm. Next, seeds were imbibed in sterile diH$_2$O in the dark

for 24 h. Finally, seeds were rinsed in sterile diH$_2$O, transferred to an empty, sterile plastic Petri

dish, and held inverted in darkness overnight for germination.

Bacterial strain construction. The BioFAB plasmid pFAB2298 was kindly provided by Vivek

Mutalik (42). We amplified the expression cassette containing Ptrc-BCD2-mRFP1 with Phusion

polymerase (Fisher) as a single unit from pFAB2298 using primers 5'-

ATCTGCAGGCTTCCCAACCTTACCAG-3' and 5'-

CAGTATCGATAGTCAGTGAGCGAGGAAGC-3', which contain 5' *Pst*I and *Cla*I restriction

sites, respectively. We digested the resulting 1.244 kb amplicon with *Pst*I and *Cla*I (New

England Biolabs) and cloned it into pBlueScript II SK(+) to create pCMB21. We digested the

expression cassette from pCMB21 using AvrII and EcoRI-HF (New England Biolabs) and

cloned into pSEVA551 (*tetA*, RSF1010 *oriV*) to create pCMB35 (43). The plasmid pCMB35

contains the broad host range RSF1010 origin of replication, the *tetA* gene encoding a

tetracycline efflux transporter which confers tetracycline resistance, and a protein expression

cassette which employs the strong, constitutive Ptrc promoter and a bicistronic 5' UTR region

with two ribosome binding sites (RBS) which allows even expression levels across expressed

genes, in this case the gene *mRFP1*. Constitutive mRFP1 expression by Rm1021 allows the

location of bacteria within the device to be identified through fluorescence microscopy. We

introduced plasmid pCMB35 into *S. meliloti* strain Rm1021 by electroporation in a 1 mm gapped

cuvette at 2200 V and plated on TY medium (6 g/L tryptone (BD), 3 g/L yeast extract (BD), 0.38

g/L CaCl$_2$ (Acros), 15 g/L agar (BD)), amended with 500 μg/mL streptomycin sulfate (Acros)

and 5 μg/mL tetracycline hydrochloride (Sigma).

μ-rhizoslide construction. We constructed small-scale, soil-containing, mesocosms adapted

from the rhizoslide designed by Le Marié, *et al* (44). Mesocosms were comprised of a 3D printed

spacer with a hollow opening which was filled with model soil. We refer to our version as a "μ-

rhizoslide." After several design iterations to balance space for root growth, plant health, and

imaging consistency, the final spacer design was 0.5 cm thick, featuring a 1.5 cm wide, 7.5 cm

long main channel with a 1.5 mm connector at the top of the channel to stiffen the spacer and

limit deformation (Figure 1).

Our μ-rhizoslides facilitate direct optical imaging of the growing plant root system while

minimizing water-based transport effects. The μ-rhizoslide assembly consists of a 3D printed

spacer sandwiched between two 50×75 mm glass slides, with a 50×85 mm filter paper wick

located between the back of the 3D printed spacer and the rear microscope slide to allow for

plant hydration via capillary action without gravity driven, bulk water flow. Spacers were

designed using Solidworks 2018, imported into PreForm software and printed on a Formlabs

Form 2 stereolithography 3D printer using Clear Resin GPCL04 (Formlabs, Somerville, MA).

To improve biocompatibility, printed spacers were treated to leach labile resin or crosslinker

using a protocol adapted from Kadilak *et al*. (45). Treatment is necessary because un-crosslinked

monomer can have antimicrobial properties (46). Briefly, we soaked 3D printed spacers in an

isopropanol bath for 30 min, then rinsed with additional isopropanol from a squeeze bottle. This

procedure was repeated twice more to remove un-crosslinked resin before resting in a 60 °C

oven for 60 min.

Assembly of μ-rhizoslides proceeded as follows. The 3D printed spacer resting on a glass

slide was filled with sterile diH$_2$O-saturated sterile model soil, then covered with a filter paper

wick, and finally the μ-rhizoslide assembly was closed by affixing a second glass slide to the

back. Germinated seeds were placed in a small depression at the top of the soil column. We kept

the μ-rhizoslides separated by treatment in transparent plastic boxes with each μ-rhizoslide

positioned 45° off vertical with the filter paper wick above the soil to encourage root growth

directly against the glass on the front of the assembly. This arrangement also helped protect the

roots from light exposure, as we kept the μ-rhizoslide boxes in a growth chamber under a 16:8 h

light:dark regime at 23 °C. We allowed the plants to grow in the μ-rhizoslides for 2 d to ensure

plants became established prior to microbial inoculation and imaging.

Growth of protists and bacteria. The protists used here were identified by 18s rRNA

sequencing as a belonging to ciliate genus, *Colpoda*, which we refer to as UC1 (47). We isolated

UC1 from a bean rhizosphere. UC1 was maintained in 25 cm^3 tissue-culture flasks (Thermo

Scientific #130192) with only 5 mL of Page's saline solution (48) to promote oxygenation. For

experimental runs, protists were fed heat-killed *E. coli* DH5a, inoculated at $OD_{595} = 0.005$ as

measured in 100 μL of a suspension in a 96-well microtiter dish. Prior to plant inoculation, UC1

cultures were stored in darkness, at room temperature, for 3 weeks. Cysts were removed from the

cell culture flask using a sterile cell scraper, washed twice with Page's saline solution, then

resuspended in 100 μL of the same solution. Protist cysts were washed, concentrated, and

counted the day of inoculation. We estimated the concentration of the cysts by averaging the

number of cysts counted in three 1 μL spots under a microscope. For experimental runs using

trophozoites (protists in their active growing stage) encysted protist cultures were washed two

days before inoculation to allow for excystment. Once most protists had emerged from the cysts

and were active, they were concentrated and counted to enable delivery of a consistent number of

protists per replicate.

Bacteria inoculants were prepared as follows. A single colony of *S. meliloti* strain

Rm1021/pCMB35 was picked from a TY + tetracycline (5 μg/ml) plate, grown in a 125 mL flask

containing 25 mL of liquid TY + tetracycline (5 μg/ml) at 30 °C with shaking at 160 rpm for 3 d.

Cells were washed twice with Page's saline solution then resuspended in Page's to OD_{595} =

0.100. OD_{595} was determined in a 96-well microtiter dish using 100 µL of cell suspension.

Experimental design and setup. For each of the three experimental runs, 2-day seedlings were

inoculated directly at the shoot-root interface with 10 µL of one of three different solutions:

1. Page's saline solution (Control);

2. Page's saline solution containing 10^7 CFU Rm1021/pCMB35 (Bacteria Only);

3. Page's saline solution containing 10^7 CFU Rm1021/pCMB35 + 1.5×10^3 *Colpoda* UC1

202   (Bacteria + Protists).

Each treatment of each experimental run was replicated between 3 and 5 times for a total of 41

different µ-rhizoslides prepared and analyzed (see Table 1). To inoculate µ-rhizoslides, we

positioned chambers at an angle either wick side up, the same way they were incubated,

(Experimental Run 1) or wick side down (Experimental Runs 2 & 3) and held inverted for 1 hr.

Inoculating wick down prevented the formation of a large fluorescence bloom directly against

the forward-facing glass at the initial time point. Eliminating this large source of "background"

fluorescence at the initial time point made it easier to measure subsequent changes in the

fluorescence signal over time. Protists were either encysted (Experimental Run 1) or trophozoites

(Experimental Runs 2 & 3); the use of trophozoites prompted quicker onset of protist-facilitated

transport activity. The watering regime was either to water µ-rhizoslides weekly with complete

fertilizer (Experimental Run 1) or every 4[th] day, alternating between sterile diH_2O and a

nitrogen-free fertilizer (Experimental Runs 2 & 3). More frequent watering and reduced nutrient

amendments helped plants avoid water starvation and fungal overgrowth. All replicates for all

treatments were handled identically for a given experimental run and the effect of methods on

results are discussed. In all cases, µ-rhizoslides were positioned during plant growth at a 45°

angle with the filter paper above the soil (except when briefly repositioned for imaging, when the

front face was positioned down on the stage of an inverted microscope). Experimental Run 1

lasted two weeks post inoculation, then was halted because several of the plants had died,

prompting the modifications to the methodology described here. Experimental Runs 2 and 3

lasted three weeks post inoculation then were stopped because the roots in several replicates had

reached the end of the soil-filled channel.

Imaging and Image Analysis. On the day of inoculation and at intervals of 1 week subsequently

we imaged the entire soil area of each μ-rhizoslide using a Zeiss AXIO-Observer Z1 inverted

microscope (Carl Zeiss Inc., Germany). Collected images included tiled brightfield and

fluorescence images using a 5× Zeiss EC Plan-Neofluar 5×/0.16 M27 lens and an AxioCam

MRm Rev.3 camera (1348×1040 pixels, 6.45×6.45 μm/pixel) set to collect 585-binned photos

(2×2 binning) when imaging. The transmission-based brightfield light equipped on the

microscope could not illuminate through the 0.5 cm of soil, so we moved the brightfield lamp to

an external stand and illuminated directly from the front of the μ-rhizoslide. For fluorescence

imaging, we found 570 nm excitation and 620 nm emission filters gave the strongest

fluorescence signal for *S. meliloti* Rm1021/pCMB35. We used the Zeiss Zen Pro 2.3 software to

stitch a single 207-megapixel image ($29,341 \times 7,072$ pixels) from each set of 585 photos then

exported the tiled image as a 12-bit grayscale TIFF at full resolution.

We created composite images for each replicate of each experimental run to illustrate the

movement of bacteria with and without protists (Figure 2, Figure S1). Here, brightfield images

showing the fully-developed root were overlain with false-colored fluorescence images at each

timepoint to show the progression of the fluorescence signal over time. These composite images

are for illustration purposes only and were not used directly for quantitative image analysis.

For quantitative image analysis, stitched fluorescence images were imported into

MATLAB and converted into $29{,}341 \times 7{,}072$ matrices. The initial (i.e., Week 0) tiled

fluorescence image of each μ-rhizoslide was subtracted from the corresponding tiled

fluorescence images from subsequent time points to remove background fluorescence, such as

from mica particles in the soil. Any resulting negative pixel value was set equal to 0. Tiled

images were cropped to remove the non-soil portion, eliminating fluorescence from the 3D

printed spacers. As *S. meliloti* bacteria was predominately found along roots and towards the

center of the soil channel, cropping out the region adjacent to the spacer had little effect on the

bacterial fluorescence signal.

To quantify displacement of the fluorescence signal by treatment and time, we partitioned

the photo matrixes into sections each approximately 1 mm high in the longitudinal direction (i.e.,

along the root axis) and extending across the full breadth of the soil-filled channel. The

dimensions of each of these sections is $386 \times 7{,}072$ pixels. Next, we determined the proportion

of each section with a non-zero fluorescence signal. We define "fluorescent pixel fraction" as the

ratio of nonzero pixels to total pixels in a section.

We defined "longitudinal extent" as the depth of the 1-mm section furthest from the soil

surface where fluorescent pixel fraction > 0.1 (i.e., where more than 10% of pixels in the section

are non-zero, after correcting for background as described above). We defined "lateral extent" as

the greatest fluorescent pixel fraction observed for a given replicate at a given point in time,

regardless of depth, then multiplied this ratio by 1.5 cm to represent the width of a continuous

fluorescent region.

The statistical significance of adding protists on both longitudinal and lateral extent was

determined by nine binary comparisons for each (one for each pair of treatments in each week).

Here, data from the three experimental runs are pooled together and p-values are computed using

unpaired t-tests, two-sided, with unequal variance, and applying a Bonferroni correction for

multiple hypothesis tests. The effect of experimental run on longitudinal and lateral extent was

also determined using linear regression. This is important because of the small changes to the

experimental procedures employed. For each direction and week, we regressed longitudinal

extent or lateral extent on treatment and experimental run, with the control treatment and

Experimental Run 1 as the omitted reference category. All statistical analyses were done using

Stata SE 17 (StataCorp 2021, College Station, TX).

Enumeration of *Colpoda sp.* and *S. meliloti* in μ-rhizoslide. After final imaging, μ-rhizoslides

were disassembled and roots were separated from the bulk soil. Root systems (plant tissue plus

adhering soil) were bisected into upper ("shallow roots") and lower ("deeper roots") segments

beginning 2.5 cm below the root/shoot interface. Root weight was recorded, then each root

segment was suspended in $10 \times$ Page's saline solution by weight in 50 mL conical tubes then

vortexed for 1 min at medium speed on a tabletop vortex mixer.

For bacterial abundance, 10-fold serial dilutions of each vortexed root extract were

prepared in Page's saline solution then 10 μL spots of each dilution were placed onto TY +

tetracycline (5 μg/mL) agar plates in square grid-marked Petri dishes and incubated at 30 °C for

3 d. Colonies were counted, and bacterial abundance of *S. meliloti* strain Rm1021/pCMB35 on

each root system segments was estimated by multiplying colony counts by the corresponding

dilution factor.

To estimate protist abundance, 3-fold serial dilutions were carried out with octuple

replication in 96-well microtiter dishes using Page's saline solution + heat-killed *E. coli* DH5a

($OD_{595} = 0.050$). Microtiter dishes were incubated at room temperature in darkness for 1 week.

The presence or absence of protists in each well was recorded. For analysis, we assumed "at least

one" protist made it into any well that contained a population of protists after one week. The

Minimum Recovered Number per ml (MRN/ml) of protists was calculated as follows:

MRN/ml $= 10 * 3^{(\text{farthest positive dilution factor} - 1)}$. Significance was determined as described previously

with paired or unpaired t-tests as noted in the data tables.

Results and Discussion

Performance of µ-rhizoslides. The µ-rhizoslides described were constructed as a smaller

version of the rhizoslides designed by C. L. Marie (44) but optimized for microscopy and

increased throughput. We report root growth and plant-microbe interactions were readily

monitored on a standard inverted microscope using these smaller µ-rhizoslides. *M. truncatula*

roots occupied a region 15×75 mm and were imaged as a series of overlapping tiled images.

Brightfield microscopy was used to localize root tissue including visualization of primary,

secondary and tertiary roots, and fluorescence microscopy was effective in monitoring

displacement of a bacteria-associated fluorescence signal over time. Images were collected once

per week for three weeks after planting, and in our case, experimental duration was limited only

by growth of the plant. The µ-rhizoslide system described here is simple and inexpensive to

produce, can support seeds as large as 4 mm diameter, and can support experiments until root

growth extends beyond the chamber size.

Displacement of *S. meliloti* is enhanced by protists. In order to test the hypothesis that addition

of protists can enhance dispersal of bacteria along root systems, the nitrogen fixing symbiont *S.*

meliloti strain Rm1021 was transformed with plasmid pCMB35 which constitutively expressed

mRFP. The movement and growth of Rm1021/pCMB35 was nondestructively assayed by

fluorescence microscopy, and the role of protists in facilitating its transport was determined by

comparing displacement of the fluorescence signal over time across treatments. Typical of our

results (Figure 2) we observed fluorescence gradually spreading with depth and breadth along

and across the growing roots. Strikingly and repeatedly, we observed that the fluorescence signal

reached further down the primary axis of the growing roots (i.e., in the longitudinal direction)

and spread more side-to-side when in the presence of soil protists. Tiled images for all forty-one

μ-rhizoslides at all points in time are available in the Supplemental Materials, Figure S1.

To analyze these images quantitatively, fluorescent pixel fraction (ratio of non-zero to

total pixels) was quantified for thin vertical strips of the soil channel for each composite image.

A plot of fluorescent pixel fraction versus depth in the chamber illustrates the displacement of

the fluorescence signal in the longitudinal direction over time (Figure 3 and Figure S2). The

results show small values of fluorescent pixel fraction scattered fairly uniformly with depth for

Control and Bacteria Only treatments, with much larger values for fluorescent pixel fraction in

all replicates of the Bacteria + Protists treatments, but in particular in the shallow portion of the

soil channel, and at later points in time. For example, for Experimental Run 2 at Week 2, most

replicates show a wide band spanning 20-30 mm of depth where fluorescent pixel fraction > 0.5.

At Week 2 and Week 3 some replicates show appreciable fluorescent pixel fraction values

extending all the way to the end of the soil-filled channel. Meanwhile, for the other two

treatments (Control, where no microbes are added, and Bacteria Only), no replicate showed such

marked fluorescence.

The pattern of fluorescent pixel fraction versus depth (i.e., the longitudinal direction) in

Experimental Run 3 is quite similar to Experimental Run 2, with the greatest fluorescent pixel

fraction near the surface. For Experimental Run 1 the fluorescence pattern is somewhat different,

with much lower fluorescent pixel fraction values overall, yet still substantially higher values of

fluorescent pixel fraction in the Bacteria + Protists treatment than in the Control or the Bacteria

Only treatments. Referencing back to the images Experimental Run 1, Bacteria + Protists

Treatment (Figure S1.1C), shows in all five replicates marked fluorescence directly adjacent to

root tissue and extending along the full depth of the channel: this would result in low fluorescent

pixel fraction values even through the root may be fully colonized.

Because even a small number of bacteria that reach a favorable habitat can create robust

populations in time, the spatial extent of even a relatively small fluorescence signal is perhaps

more important to our analysis than the total amount of fluorescence. We defined "longitudinal

extent" as the greatest depth where fluorescent pixel fraction > 10%. Practically, exceeding this

threshold of 10% fluorescent pixels reduces the probability of spurious results and conceptually

may indicate some potential for bacterial colonization. A plot of longitudinal extent by replicate

and time (Figure 4) show the effect of protists on mobilizing bacteria across the entire root

system. For example, by Week 2 the median longitudinal extent in the Bacteria + Protists

treatment was 52 mm further down the root as compared with the Bacteria Only treatment. By

Week 3, while replicates are quite variable, median longitudinal extent reached the end of the

soil-filled channel (-75 mm) for the Bacteria + Protists treatment, but extended just 10 mm down

the root for the Bacteria Only treatment. The high variability in longitudinal extent at Week 3

may be due in part to differing morphology of the root systems that naturally give rise to varying

opportunities for bacterial colonization.

Roots branch as they grow and infectible root hairs can be found near the tips of all

growing roots; therefore, bacterial displacement *across* the major axis of root growth is also

important to promote performance of bacterial soil amendments. A plot of lateral extent (the

greatest value of fluorescent pixel fraction at any depth, converted to an equivalent width) by

replicate and time shows the effect of protists on spreading bacteria across the root system

(Figure 5). For the Bacteria + Protists treatment, the median lateral extent was about 8 mm by

Week 2, and about 12 mm by Week 3. These lateral extents are substantially greater than those

observed in the Bacteria Only treatment, where the median value of lateral extent was about 1

mm, and did not change much over time.

Even greater longitudinal and lateral extents may have been observed in a different plant

growth chamber, but as the restrictions of our setup constrain all treatments equally, the platform

clearly shows the effect of protists on fluorescence signal displacement. Statistical analysis of

these results shows that the differences in longitudinal extent and lateral extent among the

different treatments are highly statistically significant. Binary unpaired t-tests were done for all

combinations of treatment (Bacteria Only, Bacteria + Protists, and Control) for all points in time.

Both vertical and horizontal distances are significantly higher in the Bacteria + Protists treatment

than in the Bacteria Only treatment or in the Control in every week (Bonferroni corrected $p <$

0.001) (Table 2). Meanwhile, no significant differences are found between the BO treatment and

Control in any week (Bonferroni corrected $p > 0.1$).

These comparisons above were performed by grouping all experiments together.

However, we detailed important differences in how the bacteria were added, whether the protists

were "awake" when added, and in the watering condition—these methodological variations

certainly impacted bacterial transport. To test for the effect of experimental conditions (i.e.,

"Experimental Run") on our results, we performed a linear regression and found that even when

controlling for experimental run, the treatment effect of protists on longitudinal and lateral extent

is still highly significant (Table S1).

Although despite methodological differences protists always promote bacterial dispersal,

we observed interesting differences in the displacement patterns with experimental run that are

worth considering. In Experimental Run 1, where soil conditions were dry, the fluorescence

signal was found primarily along the roots. In Experimental Runs 2 and 3 we observed a more

prominent, wider, and somewhat shallower pattern of fluorescence. In examining the results

quantitatively, the most significant difference is the lateral spread of the fluorescence signal. For

Experimental Run 1, median lateral extents were 1.1 mm (Week 1) and 1.2 mm (Week 2). For

Runs 2 and 3, where plants were watered nearly twice as often, median lateral extent were 5.8

mm (Week 1) and 12.2 mm (Week 2) and 7.4 mm (Week 1) and 9.0 mm (Week 2), respectively.

These differences are statistically significant (p=0.02 or less). In terms of longitudinal extent,

differences apparent at Week 1 were not significant by Week 2 (Table S2). In Experimental Run

1, the reduced water application rate may have been associated with a less-branched root

morphology, as the plant sent roots downward in search of water, as well as fewer contiguous

water-saturated pathways through the soil pore volume other than directly along the root itself.

Both of these factors would tend to focus microbial activity immediately around the roots and its

associated mucilage and promote longitudinal transport at the expense of lateral.

Protists enhance dispersal of viable bacteria along root systems. After imaging was complete,

μ-rhizoslides were disassembled and the root systems (root tissue plus adhering soil) were

divided into shallow and deep segments. The bacterial abundance in the shallow segment, as

determined by plating for CFUs, averaged 1.02×10^7 in the Bacteria Only treatment and $1.44 \times$

10^7 in the Bacteria + Protists treatment (Table 3, Figure 6). These abundances are not statistically

significant (p = 0.315). In other words, despite predation by protists, there is a similar total

bacterial population in the shallow root system of the Bacteria Only and Bacteria + Protists

treatments, implying that access to habitat niches and/or substrates may limit bacterial population

size in the shallow portion of our setup and not the effects of predation.

In the deeper root system, we found 0.458×10^7 CFU bacteria in the Bacteria + Protists

treatment (Table 3, Figure 6). versus 0 CFU recovered in the Bacteria Only treatment. We only

recovered bacteria from deeper roots in replicates where protists were present. We found here

that bacteria were unable to reach the deeper roots in our setup without the assistance of protists.

These results support the fluorescence data in showing that protists not only enhance

displacement of a bacteria-associated fluorescence signal but also seem to directly enhance the

spatial distribution of recoverable, viable bacteria in growing root microcosms.

For completeness, we also enumerated protists in root system segments. We only

recovered protists in the Bacteria + Protists treatment, and report comparable numbers from both

shallow and deep root system segments (Table 4 and Figure 6). This result implies that protists

fully disperse within the soil system by the end of the experiment.

Conclusions, Limitations, and Applications. While previous work has shown the potential for

using protists as vehicles for transporting particles (20), here we demonstrate the potential for

protists to transport viable beneficial bacteria in a system containing a living plant and real soil.

In this report, we show protist-facilitated transport increases abundance of viable bacteria in the

deeper root system segment despite losses due to grazing. The imaging experiments showed

distinct patterns of fluorescence displacement both longitudinally and laterally over time – but

only when protists were present. In the absence of protists, there was no statistically significant

change in the fluorescence signal with depth or breadth versus the control, and the imaging

showed that the fluorescence signal remained within 5 mm of the inoculation site for the duration

of the experiment.

Because fluorescent protein levels will likely vary between bacterial cells due to

differences in plasmid number, growth rate, and cell size, the fluorescence signal is most

conservatively interpreted as an indicator of bacteria position, and not a direct measurement of

bacterial abundance or biomass. In addition, our fluorescence results may have been limited

because the fluorescent plasmid pCMB35, which is usually maintained with 5 μg/mL

tetracycline, may have not been retained because tetracycline was not added to the soil in the μ-

rhizoslides. We opted not to add tetracycline because of the possibility that it could adversely

impact plant health and development. Initially, we attempted to use a different fluorescent

protein-expressing *S. meliloti* strain with a plasmid that is maintained without selection, L5-

30/pDG77, but the resulting DsRed fluorescence signal was not compatible with our imaging

setup.

For all these reasons, our quantitative approach focused on the extent of the spatial

domain that contained an appreciable fluorescent signal (i.e., fluorescent pixel fraction > 0.1 by

section). We did not attempt to aggregate total fluorescence as a proxy for abundance or

biomass. The choice to emphasize spatial extent of fluorescence and not amount of fluorescence

helps to explain why longitudinal and lateral extent are dramatically different between treatments

yet we recover similar numbers of bacteria in the top (shallow root system) segment of the

Bacteria Only and Bacteria + Protists treatments. In the former, the entire population of bacteria

is limited to the initial inoculation site, whereas bacteria in the Bacteria + Protists treatment are

spread out over a larger area. Importantly, *S. meliloti* strain Rm1021/pCMB35 was not recovered

in the deeper system segment at all unless protists were present. The bacteria appear not to be

able to reach that area of the soil channel after three weeks of plant development and growth

without the assistance of protists.

The quantities of bacteria and protists recovered in our study represent minimum values

for several reasons. First, only bacteria and protists directly associated with roots and root-

adhering soil were quantified. The bulk soil, which was not evaluated by these methods, likely

contained additional organisms. Also, our methods may tend to under-represent the true

microbial abundance, especially for protists. For our MRN assay the true value may be

approximately three times higher because of the assumption that the last well of a dilution series

initially contained only one protist when in fact it may have contained two or more protists.

However, despite these important methodological details, the trend in our data strongly support

the hypothesis that protists enhance dispersal of bacteria along root systems, including bacteria

that are recoverable and viable. The effect of these bacteria on the roots and plant health is the

subject of a further study.

The protist used in this study, *Colpoda sp.*, a ciliate isolated from a legume rhizosphere,

is particularly successful in transporting bacteria. Ciliates most often filter feed, a mechanism of

promiscuous consumption of food particles and prey. Bacteria consumed by the ciliates are

packaged into a phagosome which is processed and eventually egested from the protist; a process

that is not well characterized. It is thought that egested phagosomes break down in the soil and

these may release viable bacteria which could go on to colonize the area where they were

released. How often bacteria can survive intake and egestion, and whether some species are

adept at this in not clearly understood. Some bacterial species, for example *Campylobacter*

jejuni, Listeria spp. and *Mycobacterium* spp. are known to escape digestion and colonize the

cytoplasm of protists that consumed them (38, 49). Ciliates have also been shown to emerge

from feeding on bacterial biofilms with a matrix of bacteria and EPS stuck to their cell bodies

(20). When the protists travel along plant roots consuming bacteria, both egestion of viable

bacteria and physical translocation of bacteria may contribute to protist-assisted transport.

The technology described here provides a simple, reproducible experimental system

which has the potential to aid in the understanding of rhizosphere biology. Of particular interest

is bacterial movement in the context of screening a variety of different bacteria/protist/plant/soil

combinations for their effects on such movement. For example, protist transportation of

Sinorhizobium sp. and its effect on nodulation can be studied using this system and parameters

that affect transport such as water saturation, soil composition and soil porosity can be readily

altered then tested and quantitated. Protist-facilitated transport may also be adapted for targeted

delivery of specialized microbes or encapsulated agrochemicals. With increased understanding

of protist-facilitated transport mechanisms and limitations under reproducible yet realistic

conditions we can begin developing new methods to support sustainable agriculture and have

increasing confidence of their utility for use in the field.

Acknowledgements. This work was supported by DOE award DE-SC0014522 (to LMS and

DJG), NSF award 1605816 (to LMS and DJG), and USDA National Institute of Food and

Agriculture award 2016-67013-24412 (to DJG and LMS).

References

1.   Berendsen RL, Pieterse CMJ, Bakker PAHM. 2012. The rhizosphere microbiome and plant health.
Trends Plant Sci.

2.   van Veen JA, Liljeroth E, Lekkerkerk LJA, van de Geijn SC. 1991. Carbon Fluxes in Plant-Soil
Systems at Elevated Atmospheric CO2 Levels. Ecol Appl 1:175–181.

3.   Guo Y-S, Furrer JM, Kadilak AL, Hinestroza HF, Gage DJ, Cho YK, Shor LM. 2018. Bacterial
Extracellular Polymeric Substances Amplify Water Content Variability at the Pore Scale. Front
Environ Sci 6:93.

4.   Haas D, Défago G. 2005. Biological control of soil-borne pathogens by fluorescent
pseudomonads. Nat Rev Microbiol 3:307–319.

5.   Baveye PC, Otten W, Kravchenko A, Balseiro-Romero M, Beckers É, Chalhoub M, Darnault C,
Eickhorst T, Garnier P, Hapca S, Kiranyaz S, Monga O, Mueller CW, Nunan N, Pot V, Schlüter S,
Schmidt H, Vogel H-J. 2018. Emergent Properties of Microbial Activity in Heterogeneous Soil
Microenvironments: Different Research Approaches Are Slowly Converging, Yet Major
Challenges Remain. Front Microbiol 9:1929.

6.   Jiang X, Zerfaß C, Feng S, Eichmann R, Asally M, Schäfer P, Soyer OS. 2018. Impact of spatial
organization on a novel auxotrophic interaction among soil microbes. ISME J 12:1443–1456.

7.   Ben-Jacob E, Finkelshtein A, Ariel G, Ingham C. 2016. Multispecies Swarms of Social
Microorganisms as Moving Ecosystems. Trends Microbiol 24:257–269.

8.   Young IM, Crawford JW. 2004. Interactions and self-organization in the soil-microbe complex.
Science 304:1634–7.

9.   Cho H, Jönsson H, Campbell K, Melke P, Williams JW, Jedynak B, Stevens AM, Groisman A,
Levchenko A. 2007. Self-Organization in High-Density Bacterial Colonies: Efficient Crowd
Control. PLoS Biol 5:e302.

10.   Kim HJ, Boedicker JQ, Choi JW, Ismagilov RF. 2008. Defined spatial structure stabilizes a
synthetic multispecies bacterial community. Proc Natl Acad Sci 105:18188–18193.

11.   Deng J, Orner EP, Chau JF, Anderson EM, Kadilak AL, Rubinstein RL, Bouchillon GM, Goodwin
RA, Gage DJ, Shor LM. 2015. Synergistic effects of soil microstructure and bacterial EPS on
drying rate in emulated soil micromodels. Soil Biol Biochem 83:116–124.

12.   Soufan R, Delaunay Y, Gonod LV, Shor LM, Garnier P, Otten W, Baveye PC. 2018. Pore-scale
monitoring of the effect of microarchitecture on fungal growth in a two-dimensional soil-like
micromodel. Front Environ Sci 6.

13.   Krome K, Rosenberg K, Bonkowski M, Scheu S. 2009. Grazing of protozoa on rhizosphere
bacteria alters growth and reproduction of Arabidopsis thaliana. Soil Biol Biochem 41:1866–1873.

14.   Bonkowski M, Cheng W, Griffiths BS, Alphei J, Scheu S. 2000. Microbial-faunal interactions in
the rhizosphere and effects on plant growth. Eur J Soil Biol 36:135–147.

15.   Bonkowski M, Clarholm M. 2012. Stimulation of plant growth through interactions of bacteria
522   and protozoa: Testing the auxiliary microbial loop hypothesis. Acta Protozool. Portal Czasopism
523   Naukowych Ejournals.eu.

524   16.   Henkes GJ, Kandeler E, Marhan S, Scheu S, Bonkowski M. 2018. Interactions of mycorrhiza and
525   protists in the rhizosphere systemically alter microbial community composition, plant shoot-to-
526   root ratio and within-root system nitrogen allocation. Front Environ Sci 6:117.

527   17.   Bouwman LA, Bloem J, van den Boogert PHJF, Bremer F, Hoenderboom GHJ, de Ruiter PC.
528   1994. Short-term and long-term effects of bacterivorous nematodes and nematophagous fungi on
529   carbon and nitrogen mineralization in microcosms. Biol Fertil Soils 17:249–256.

530   18.   Bonkowski M. 2004. Protozoa and plant growth: The microbial loop in soil revisited. New Phytol.
531   John Wiley & Sons, Ltd (10.1111).

532   19.   Rønn R, McCaig AE, Griffiths BS, Prosser JI. 2002. Impact of protozoan grazing on bacterial
533   community structure in soil microcosms. Appl Environ Microbiol 68:6094–105.

534   20.   Rubinstein RL, Kadilak AL, Cousens VC, Gage DJ, Shor LM. 2015. Protist-facilitated particle
535   transport using emulated soil micromodels. Environ Sci Technol 49:1384–1391.

536   21.   Shor LM, Chau JF, Gage DJ. October 2018. Methods and systems to replicate soil properties.
537   10101312. United States.

538   22.   Gage DJ, Shor LM, Gage JL, Micciulla JL, Rubinstein RL. 2017. Microbial carriers for targeted
539   delivery of agricultural payloads. US9603368B2. United States.

540   23.   Muok AR, Claessen D, Briegel A. 2021. Microbial hitchhiking: how Streptomyces spores are
541   transported by motile soil bacteria. ISME J 2021 1–10.

542   24.   Muok AR, Briegel A. 2021. Intermicrobial Hitchhiking: How Nonmotile Microbes Leverage
543   Communal Motility. Trends Microbiol 29:542–550.

544   25.   King W, Bell T. 2021. Can dispersal be leveraged to improve microbial inoculant success? Trends
545   Biotechnol https://doi.org/10.1016/J.TIBTECH.2021.04.008.

546   26.   Kuo AP, Bhattacharjee N, Lee Y-S, Castro K, Kim YT, Folch A. 2019. High-Precision
547   Stereolithography of Biomicrofluidic Devices. Adv Mater Technol 1800395.

548   27.   Rodríguez H, Fraga R. 1999. Phosphate solubilizing bacteria and their role in plant growth
549   promotion. Biotechnol Adv 17:319–339.

550   28.   Glick B. 2012. Plant growth-promoting bacteria: mechanisms and applications. Scientifica (Cairo)
551   2012:1–15.

552   29.   Berg G. 2009. Plant-microbe interactions promoting plant growth and health: perspectives for
553   controlled use of microorganisms in agriculture. Appl Microbiol Biotechnol 84:11–18.

554   30.   Wadisirisuk P, Danso SKA, Hardarson G, Bowen GD. 1989. Influence of Bradyrhizobium
555   japonicum Location and Movement on Nodulation and Nitrogen Fixation in Soybeans. Appl
556   Environ Microbiol 55:1711.

557   31.   Madsen EL, Alexander M. 1982. Transport of Rhizobium and Pseudomonas through Soil. Soil Sci
558   Soc Am J 46:557–560.

559   32.   López-García S, Vázquez T, Favelukes G, Lodeiro A. 2002. Rhizobial position as a main
560   determinant in the problem of competition for nodulation in soybean. Environ Microbiol 4:216–
561   224.

562   33.   Humphris SN, Bengough AG, Griffiths BS, Kilham K, Rodger S, Stubbs V, Valentine TA, Young
563   IM, SN H, AG B, BS G, K K, S R, V S, TA V, IM Y. 2005. Root cap influences root colonisation
564   by Pseudomonas fluorescens SBW25 on maize 54:123–130.

565   34.   Stougaard J. 2000. Regulators and Regulation of Legume Root Nodule Development. Plant
566   Physiol 124:531 LP – 540.

567   35.   Lund ZF. 1970. The Effect of Calcium and Its Relation to Several Cations in Soybean Root
568   Growth. Soil Sci Soc Am J 34:456–459.

569   36.   Korber DR, Lawrence JR, Sutton B, Caldwell DE. 1989. Effect of laminar flow velocity on the
570   kinetics of surface recolonization by Mot+ and Mot− Pseudomonas fluorescens. Microb Ecol
571   18:1–19.

572   37.   Wang W, Shor LM, LeBoeuf EJ, Wikswo JP, Taghon GL, Kosson DS. 2008. Protozoan migration
573   in bent microfluidic channels. Appl Environ Microbiol 74:1945–9.

574   38.   First MR, Park NY, Berrang ME, Meinersmann RJ, Bernhard JM, Gast RJ, Hollibaugh JT. 2012.
575   Ciliate Ingestion and Digestion: Flow Cytometric Measurements and Regrowth of a Digestion-
576   Resistant Campylobacter jejuni. J Eukaryot Microbiol 59:12–19.

577   39.   Best A, Kwaik YA. 2019. Evasion of phagotrophic predation by protist hosts and innate immunity
578   of metazoan hosts by Legionella pneumophila. Cell Microbiol 21.

579   40.   Brock DA, Douglas TE, Queller DC, Strassmann JE. 2011. Primitive agriculture in a social
580   amoeba. Nature 469:393–396.

581   41.   Disalvo S, Haselkorn TS, Bashir U, Jimenez D, Brock DA, Queller DC, Strassmann JE. 2015.
582   Burkholderia bacteria infectiously induce the protofarming symbiosis of Dictyostelium amoebae
583   and food bacteria. Proc Natl Acad Sci U S A 112:E5029–E5037.

584   42.   Mutalik VK, Guimaraes JC, Cambray G, Lam C, Christoffersen MJ, Mai QA, Tran AB, Paull M,
585   Keasling JD, Arkin AP, Endy D. 2013. Precise and reliable gene expression via standard
586   transcription and translation initiation elements. Nat Methods 10:354–360.

587   43.   Silva-Rocha R, Martínez-García E, Calles B, Chavarría M, Arce-Rodríguez A, De Las Heras A,
588   Páez-Espino AD, Durante-Rodríguez G, Kim J, Nikel PI, Platero R, De Lorenzo V. 2013. The
589   Standard European Vector Architecture (SEVA): A coherent platform for the analysis and
590   deployment of complex prokaryotic phenotypes. Nucleic Acids Res 41.

591   44.   Le Marié C, Kirchgessner N, Marschall D, Walter A, Hund A. 2014. Rhizoslides: Paper-based
592   growth system for non-destructive, high throughput phenotyping of root development by means of
593   image analysis. Plant Methods 10.

594   45.   Kadilak AL, Rehaag JC, Harrington CA, Shor LM. 2017. A 3D-printed microbial cell culture
595   platform with *in situ* PEGDA hydrogel barriers for differential substrate delivery. Biomicrofluidics
596   11:054109.

597   46.   Macdonald NP, Zhu F, Hall CJ, Reboud J, Crosier PS, Patton EE, Wlodkowic D, Cooper JM.
598   2016. Assessment of biocompatibility of 3D printed photopolymers using zebrafish embryo
599   toxicity assays. Lab Chip 16:291–7.

600   47.   Taerum SJ, Steven B, Gage DJ, Triplett LR. 2020. Validation of a PNA clamping method for
601   reducing host DNA amplification and increasing eukaryotic diversity in rhizosphere microbiome
602   studies. bioRxiv 2020.05.07.082966.

603   48.   Caspers H. 1978. F. C. Page: An Illustrated Key to Freshwater and Soil Amoebae with notes on
604   Cultivation and Ecology. – With 64 fig., 155 pp. The Ferry House, Far Sawrey, Ambleside,
605   Cumbria: Freshwater Biological Association, Scientific Publication No. 34. 1976. SBN 900 386
606   26 6, ISSN 0367-1857. £ 2.50. Int Rev der gesamten Hydrobiol und Hydrogr 63:289–289.

607   49.   Adékambi T, Salah S Ben, Khlif M, Raoult D, Drancourt M. 2006. Survival of environmental
608   mycobacteria in Acanthamoeba polyphaga. Appl Environ Microbiol 72:5974–5981.

610   Supplemental Material linked to the online version of the paper at Wiley-Blackwell:
611   •   Tables S1, S2
612   •   Figures S1, S2.

Figures and Tables

Table 1. Experimental design showing three independent experimental runs each with multiple replicates of each treatment. A grand total of 41 separate µRhizoslides were imaged on successive weeks, then sacrificed to enumerate bacteria and protists. Treatment abbreviations are Control (C); Bacteria Only (BO); and Bacteria + Protists (BP). Plants in Experimental Run 1 died during the third week leading to modified conditions for Experimental Runs 2 and 3.

	Experimental Run #1			Experimental Run #2			Experimental Run #3		
	C	BO	BP	C	BO	BP	C	BO	BP
Week 1 Week 2 Week 3	n = 5	n = 5	n = 5	n = 5	n = 5	n = 5	n = 3	n = 4	n = 4
	n/a: plants died								
Tiled Images				Fig. 2 (selected reps.)					
	Fig. S1.1A	Fig. S1.1B	Fig. S1.1C	Fig. S1.2A	Fig. S1.2B	Fig. S1.2C	Fig. S1.3A	Fig. S1.3B	Fig. S1.3C
Results	Fig. S2A			Fig. 3			Fig. S2B		
	Fig. 4, Fig. 5, Fig. 6								

Table 2: Pairwise significance (p-values[*]) comparing longitudinal extent (i.e., with plant depth, see Figure 4) and lateral extent (see Figure 5) by treatment and by week. Treatment abbreviations are Control (C), where no microbes are added; Bacteria Only (BO); and Bacteria + Protists (BP).

Treatment Pairs	Week 1	Week 2	Week 3
Longitudinal Extent			
BO v. C	p=0.120	p=0.869	p=0.351
BP v. C	p<0.001***	p<0.001***	p<0.001***
BP v. BO	p=0.001***	P<0.001***	p<0.001***
Lateral Extent			
BO v. C	p=0.138	p=0.590	p=0.349
BP v. C	p<0.001***	p<0.001***	p<0.001***
BP v. BO	p=0.001***	p<0.001***	p<0.001***

p-values of unpaired t-tests, two-sided, unequal variance with Bonferroni correction for multiple hypothesis tests
*** p<0.01, ** p<0.05, * p<0.10

Figures and Tables

Table 3: Bacterial abundance ($\times 10^7$ as colony forming units, or CFUs) averaged over all runs and replicates by root system segment (i.e., shallow v. deep) and treatment with statistical significance (p-value). Treatment abbreviations are Control (C), where no microbes are added; Bacteria Only (BO); and Bacteria + Protists (BP).

Treatment	Shallow	Deep	Total	Shallow v. Deep[+]
C	0	0	0	---
BO	1.02	0	1.02	p=0.001***
BP	1.44	0.458	1.90	p=0.008**
BP v. BO[*]	p=0.315	p=0.013**	p=0.080*	

[*]*BP v. BO: p-values of unpaired t-tests, two-sided, unequal variance*
[+]*Shallow v. Deep: p-values of paired t-tests, two-sided*
*** p<0.01, ** p<0.05, * p<0.10

Table 4: Protist abundance averaged across all runs and replicates. Abundance is estimated as minimum recovered number (MRN) of propagated extract dilutions by root system segment (i.e., shallow v. deep) and treatment with statistical significance (p-value).

Treatment	Shallow	Deep	Total	Shallow v. Deep[+]
C	0	0	0	---
BO	0	0	0	---
BP	233	185	419	p=0.026**

[+]*Shallow v. Deep: p-values of paired t-tests, two-sided*
*** p<0.01, ** p<0.05, * p<0.10

Figure Legends

Figure 1. μ-rhizoslide construction and use. A) Computer-aided design (CAD) drawing of the
rhizoslide assembly showing layers including 3D printed rhizoslide spacer and filter paper wick
between two glass slides. B) Schematic of assembled μ-rhizoslide. C) Photograph of μ-
rhizoslide in use.

Figure 2. Selected composite brightfield (at last imaging date) and fluorescence (false-
colored by week as indicated) images of soil channels for: A) Control: soil and plant
inoculated with 10 μL buffer: no microbes added; B) Bacteria Only: soil and plant inoculated
with 10 μL buffer containing 10^7 CFU bacteria; C) Bacteria + Protists: soil and plant inoculated
with 10 μL buffer containing 10^7 CFU bacteria plus 1.5 x 10^3 *Colpoda sp*. protists. Images for all
experimental runs and all replicates are available in Figure S1.

Figure 3. Spatial distribution of fluorescence signal over time. For each treatment (arranged
above in columns), the entire soil channel was imaged at different points in time (Week 0, Week
1, Week 2, Week 3). Here, "Fluorescent Pixel Fraction" (the ratio of non-zero pixels to total
pixels) was computed for binned strips (each measuring 1 mm high and 15 mm wide) of the soil
channel after background was subtracted (Week 0) and plotted with distance in the longitudinal
direction. Data from Experimental Run 2 with replicates distinguished by color. Full data from
the other two experimental runs can be found in Supplemental Material, Figure S2.

Figure 4. Longitudinal extent (the greatest depth where fluorescent pixel fraction > 10%)
by treatment and time. The longitudinal direction is along the major axis of root growth.
Experimental runs are grouped by color (Run 1 = blue circles: Run 2 = red squares; Run 3 =
yellow triangles). Top and bottom edges of each box represent upper and lower quartiles, and the
inside line is the sample median. Whiskers represent the non-outlier minimum and maximum
values. Outliers were determined as values more than 1.5 times the inner quartile range beyond
the box limits. Treatments are: Control (C) n=13; Bacteria Only (BO) n=14; Bacteria + Protists
(BP) n=14.

Figure 5. Lateral extent (the largest fluorescent pixel fraction at any depth converted to an
equivalent width) of the fluorescence signal by treatment and time. The lateral direction is
across the major axis of root growth. Experimental runs are grouped by color (Run 1 = blue
circles: Run 2 = red squares; Run 3 = yellow triangles). Top and bottom edges of each box
represent upper and lower quartiles, and the inside line is the sample median. Whiskers represent
the non-outlier minimum and maximum values. Outliers were determined as values more than
1.5 times the inner quartile range beyond the box limits. Treatments are: Control (C) n=13;
Bacteria Only (BO) n=14; Bacteria + Protists (BP) n=14.

Figures and Tables

Figure 6. Bacteria and protist abundance in the top ("shallow roots") and bottom ("deeper roots") segments of *M. truncatula* root systems. Bacteria abundance is represented by average colony forming units (CFUs) in plated serial dilutions of the root system extract. Protist abundance is represented by average minimum recovered number (MRN) of protists in propagated serial dilutions of the root system extract. Experimental runs are grouped by color (Run 1 = blue circles: Run 2 = red squares; Run 3 = yellow triangles). Top and bottom edges of each box represent upper and lower quartiles, and the inside line is the sample median. Whiskers represent the non-outlier minimum and maximum values. Outliers were determined as values more than 1.5 times the inner quartile range beyond the box limits. Treatments are: Control (C) n=13; Bacteria Only (BO) n=14; Bacteria + Protists (BP) n=14.

A   B   C

1 cm

Week 0
Week 1
Week 2
Week 3

1 cm