1

Soil protists can actively redistribute beneficial bacteria along *Medicago truncatula* roots

Christopher J. Hawxhurst[Δa], Jamie L. Micciulla[Δb], Charles M. Bridges[b], Leslie M. Shor[a,c], Daniel J. Gage[b#]

[Δ]Co-first Authors

[a]Department of Chemical & Biomolecular Engineering, University of Connecticut, Storrs, CT, USA

[b]Department of Molecular and Cellular Biology, University of Connecticut, Storrs, CT, USA

[c]Center for Environmental Sciences & Engineering, University of Connecticut, Storrs, CT, USA

#Address correspondence to Daniel Gage, Daniel.Gage@UConn.edu

Christopher J. Hawxhurst and Jamie L. Micciulla contributed equally to this work. Author order was determined alphabetically.

Keywords: Protist, Soil, Plant Beneficial Bacteria, Transport

Abstract

The rhizosphere is the region of soil directly influenced by plant roots. The microbial community

in the rhizosphere includes fungi, protists, and bacteria, all of which play a significant role in plant

health. The beneficial bacterium Sinorhizobium meliloti infects growing root hairs on nitrogen

starved leguminous plants. Infection leads to the formation of a root nodule, where S. meliloti

converts atmospheric nitrogen to ammonia, the usable form of nitrogen for plants. However, S.

meliloti, often found in biofilms, travels slowly; whereas infectible root hairs are found at the

growing root tip, potentially causing many root hairs to remain uninfected by S. meliloti when it

is delivered as a seed inoculant. Soil protists are an important component of the rhizosphere system

who prey on soil bacteria and have been known to egest undigested phagosomes. We show that

the soil protist, Colpoda sp., plays an important role in transporting S. meliloti down Medicago

truncatula roots. By using pseudo-3D soil microcosms we directly observed the presence of

fluorescently labelled S. meliloti along M. truncatula roots and track the displacement of bacteria

over time. In the presence of Colpoda sp., S. meliloti was detected 44 mm, on average, farther

down the roots, compared with the Bacteria Only Treatment. Facilitating bacterial transport may

be an important mechanism whereby soil protists promote plant health. Protist facilitated transport

as a sustainable agriculture biotechnology has the potential to boost efficacy of bacterial

inoculants, avoid overuse of nitrogen fertilizers, and enhance performance of no-till farming

practices.

Importance

Soil protists are an important part of the microbial community in the rhizosphere. Plants

grown with protists fare better than plants grown without protists. Mechanisms through which

protists support plant health include nutrient cycling, alteration of the bacterial community through

selective feeding and consumption of plant pathogens. Here we provide data in support of an

additional mechanism: protists act as a transport vectors for bacteria. We show that protist-

facilitated transport can deliver plant-beneficial bacteria to the growing tips of roots that may

otherwise be sparsely inhabited with bacteria from a seed-associated inoculum. By co-inoculating

Medicago truncatula roots with both *Sinorhizobium meliloti,* a nitrogen fixing legume symbiont,

and the ciliated protist, *Colpoda sp.,* we show bacteria are distributed an average of 44 mm farther

down plant roots than when the plants are inoculated by bacteria alone. Co-inoculation with shelf-

stable encysted soil protists may be employed as a sustainable agriculture biotechnology to better

distribute beneficial bacteria and enhance the performance of inoculants.

Introduction. The rhizosphere is the zone of soil surrounding plant roots that is under the influence of the root. This soil fraction contains 5 to 100 times more organisms per unit volume than adjacent bulk soil (1). Up to 40% of carbon assimilated by plants is contributed back to the soil as rhizo-deposits (2). In return for these carbon-rich root exudates, plants often benefit from microbial activity through a variety of mechanisms such as moisture retention, pathogen suppression, plant-hormone synthesis and the release of recalcitrant nutrients (3, 4). Furthermore, some soil dwelling bacteria, such as those from the family Rhizobiaceae, beneficially infect leguminous roots and directly provide fixed nitrogen in exchange for TCA cycle intermediates.

The functioning of soil and rhizosphere systems emerges, in part, from biological communities interacting in a spatially structured environment (5–11). In trying to better understand the importance of biological factors on the properties of soil, research has been dedicated to the study and interaction of different bacterial species and plants in spatially-constrained systems (6, 9, 10). Protists form an additional, important, and sometimes overlooked component of the soil system. The presence of protists in addition to bacteria can promote and enhance size and weight of plant roots and shoots over bacteria alone (12). Protist-associated enhancement to plant health can arise through several different mechanisms. In the "microbial loop" hypothesis, protists recycle limiting nutrients by grazing on the microbes fed by root exudates (13–17). Protists may also alter the microbial composition of the rhizosphere through selective grazing (17, 18). A third potential mechanism is that protists may improve plant health by acting as a transport vector: moving bacteria or other "cargo" along downward-growing roots (19–21). Similar examples include transport of fungal spores hitchhiking on bacterial flagella (37). Other examples are given in recent reviews (38, 39).

Plant growth promoting rhizobacteria (PGPR) are beneficial soil bacteria used in

agriculture to improve yields and disease resistance. PGPR-mediated nutritional effects include

symbiotic nitrogen fixation, non-symbiotic nitrogen fixation, release of recalcitrant nitrogen from

soils, phosphate solubilization, and iron delivery (2–5). Commercially-used PGPR include many

species of nitrogen-fixing bacteria: *Azospirillum* spp.*, Azotobacter* spp., *Pseudomonas* spp.,

Bacillus subtilis strains, *Bacillus megaterium* strains and *Trichoderma* spp. (a fungal genus) (2).

Many additional species are under investigation in laboratory, greenhouse, and field trials, but

have not yet been commercialized.

PGPR are typically inoculated onto seeds before planting or they are directly applied to the

soil during planting. Soil bacteria usually grow as biofilms and spread very slowly. Thus, bacteria

inoculated onto, or near, seeds often do not leave the area of inoculation and do not establish a

population on the root large enough to provide beneficial services (6–9, 22). Further, symbiotic

infection of root hairs with nitrogen-fixing rhizobia, such as *Sinorhizobium meliloti*, often occurs

at growing root hairs which are located growing at growing root tips. Since roots grow at their tips,

these are actively moving away from the soil surface where the seed was planted. The relative lack

of bacteria at the root tip can cause a decreased rate of symbiotic infection and nodulation (23).

Soil protists have the potential to act as transport vehicles for bacteria. Soybeans roots can

grow at ~3000 μm per hour (24). For comparison, the spreading rate of biocontrol *P. fluorescens*

biofilms is less than 10 μm per hour (25), while ciliated protists can swim at speeds up to 400 μm

per second for short periods (26). We have shown in vitro that protists can transport beads and

bacteria much faster than they could travel without protists (19). In situ, protists are known to carry

bacteria, most commonly as intracellular parasites, for example when *Legionella* spp. infect

aquatic amoeba or *Campylobacter* spp infect ciliates [28, 40]. Also, bacterial transport by protists

has been demonstrated in the soil amoeba *Dictyostelium discoideum*, which can carry bacteria to

new habitats (26, 27). Although protists graze on bacteria, ingestion is not always fatal (28).

Further, bacteria have been observed attached to the outer surfaces of protists, thus providing a

means for transport that does not require ingestion (19). We propose protists contribute to the

rhizosphere community, in part, through their capacity to spatially redistribute bacteria. The goal

of this study was to test the hypothesis that a protist species, despite grazing, could enhance

bacterial dispersal along root systems. We measured the spatial distribution of fluorescently

labelled *S. meliloti* through soil-containing mesocosms over time in the presence and absence of a

soil-isolated, ciliated, protist from the genus *Colpoda*.

Experimental Materials and Methods

Model soil. Sterile model soil was comprised of play sand (Quikrete) and fine vermiculite

combined 1:1 in a soil mixer. The mix was fractionated by sieving and the 125-250 μm fraction

and 63–125 μm fraction were combined 1:2 to emulate the particle sized distribution of a model

sandy loam (11).

Plants. *Medicago truncatula* is a model legume that offers uniform growth and natural symbiosis

with *S. meliloti.* To synchronize the plants, we sterilized the seeds by successive washing with

95% ethanol for 30 min, 10% bleach for 15 min, and sterile diH$_2$O for 15 min on a shaking platform

at 80 rpm. Next, we imbibed seeds in sterile diH$_2$O in the dark for 24 h. Then, we rinsed the seeds

in sterile diH$_2$O, transferred them to a sterile plastic petri dish, and stored them in an inverted dish

in darkness overnight for germination.

Bacterial strain construction. The BioFAB plasmid pFAB2298 was kindly provided by Vivek

Mutalik (29). We amplified the expression cassette containing Ptrc-BCD2-mRFP1 with Phusion

polymerase   (Fisher)   as   a   single   unit   from   pFAB2298   using   primers   5'-

ATCTGCAGGCTTCCCAACCTTACCAG-3' and 5'-

CAGTATCGATAGTCAGTGAGCGAGGAAGC-3', which contain 5' *Pst*I and *Cla*I restriction

sites, respectively. We digested the resulting 1.244 kb amplicon with *Pst*I and *Cla*I (New England

Biolabs) and cloned it into pBlueScript II SK(+) to create pCMB21. We digested the expression

cassette from pCMB21 using AvrII and EcoRI-HF (New England Biolabs) and cloned into

pSEVA551 (*tetA*, RSF1010 *oriV*) to create pCMB35 (30). We introduced plasmid pCMB35 into

S. meliloti strain Rm1021 by electroporation in a 1 mm gapped cuvette at 2200 V and plated on

TY medium (6 g/L tryptone (BD), 3 g/L yeast extract (BD), 0.38 g/L CaCl$_2$ (Acros), 15 g/L agar

(BD)), amended with 500 μg/mL streptomycin sulfate (Acros) and 5 μg/mL tetracycline

hydrochloride (Sigma).

μ-Rhizoslide construction. We constructed small-scale, soil-containing, mesocosms from the

rhizoslide designed by Le Marié, *et al* (31). Mesocosms were comprised of a 3D printed spacer

with a hollow opening which was filled with model soil. This was sandwiched between two 50 ×

75 mm glass slides. We refer to our version as a "μ-rhizoslide". After several design iterations to

balance space for root growth, plant health, and imaging consistency, the final spacer design

features a 1.5 cm wide, 7.5 cm long main channel with a 1.5 mm connector at the top of the channel

to stiffen the spacer and limit deformation (Fig. 1).

Our μ-rhizoslides allow for direct optical imaging of the growing plant root and minimize

any water-based transport effects. The μ-rhizoslide assembly (Fig 1A) consists of a 3D printed

spacer sandwiched between two 50 × 75 mm glass slides, with a 50 × 85 mm filter paper wick

added between the back of the 3D printed spacer and the rear microscope slide, to allow for plant

hydration without gravity driven, bulk, water flow. Spacers were designed using Solidworks 2018,

imported into PreForm software and printed on a Formlabs Form 2 stereolithography 3D printer

using Clear Resin GPCL04 (Formlabs, Somerville, MA). To improve biocompatibility, printed

spacers were treated to leach labile resin or crosslinker using a protocol adapted from Kadilak et

al. (32). Treatment is necessary because un-crosslinked monomer can have antimicrobial

properties (33). Briefly, we soaked 3D printed spacers in an isopropanol bath for 30 min, then

rinsed with additional isopropanol from a squeeze bottle. This treatment was repeated twice more

to ensure there was no remaining un-crosslinked resin.

Assembly of μ-rhizoslides proceeded as follows. The 3D printed spacer resting on a glass

slide was filled with sterile diH_2O-saturated sterile model soil, then covered with a filter paper

wick, and closed by affixing another glass slide on the front of the μ-rhizoslide assembly.

Germinated seeds were placed in a small depression at the top of the soil column. We kept the μ-

rhizoslides separated by treatment in transparent plastic boxes with each μ-rhizoslide positioned

45° off vertical with the filter paper wick on the top face to encourage root growth directly against

the glass slide on the bottom of the assembly. This arrangement also helped protect the roots from

light exposure, as we kept the μ-rhizoslide boxes in a growth chamber under a 16:8 h light:dark

cycle at 23 °C to maintain the plants. We allowed the plants to grow in the μ-rhizoslides for 2 d to

insure root establishment prior to inoculation and imaging.

Growth of protists and bacteria. The protists used in this experiment were identified by 18s

rRNA sequencing as a belonging to ciliate genus *Colpoda*, which we refer to as UC1 (34). UC1

was isolated from a bean rhizosphere and cultured in the lab in Page's Saline Solution (35). UC1

was maintained in 25 cm3 tissue-culture flasks (Thermo Scientific #130192) with 5 mL of Page's

Saline to allow for full oxygenation for optimal protist growth. Protists were fed heat-killed *E. coli*

DH5a, inoculated at OD595 = 0.005 as measured in 100 μL of a suspension in a 96-well microtiter

dish. Prior to plant inoculation, UC1 cultures were stored in darkness, at room temperature, for 3

weeks. Cysts were removed from the cell culture flask using a sterile cell scraper, washed twice

with Page's Saline Solution, then resuspended in 100 µL of the same medium. Protist cysts were

washed, concentrated, and counted the day of inoculation. We estimated the concentration of the

cysts by averaging the number of cysts counted in three 1 µL spots under a microscope. For

experiments using trophozoites (protists in their active growing stage) encysted protist cultures

were washed two days before inoculation to allow for excystment. Once most protists had emerged

from the cysts and were active they were concentrated, counted, and inoculated them onto plants

that had been previously established in µ-rhizoslides, as outlined above.

Bacteria inoculants were prepared as follows. A single colony of *S. meliloti* strain

Rm1021/pCMB35 was picked from a TY + tetracycline (5 µg/ml) plate, grown in a 125 ml flask

containing 25 ml of liquid TY + tetracycline (5 µg/ml) at 30 °C with shaking at 160 rpm for 3 d.

Cells were washed twice with Page's saline solution then resuspended in Page's to OD_{595} = 0.1.

OD_{595} was determined in a 96-well microtiter dish using 100 µL of cell suspension.

Experimental design. We prepared 10 µL inocula for five replicate µ-rhizoslides for each of the

following three treatments: 1) Page's medium alone (Control); 2) 10^7 CFU Rm1021/pCMB35

in Page's medium (Bacteria Only); 3) 10^7 CFU Rm1021/pCMB35 + 1.5 × 10^3 *Colpoda* UC1 in

Page's medium (Bacteria + Protist). To prevent a large fluorescent bloom at the inoculation site,

we delivered inocula directly onto the root/shoot interface with the µ-rhizoslide positioned wick-

side down, opposite of the orientation used for growth (described above), to avoid gravity-driven

flow of the inoculum along the front glass slide. Inoculated µ-rhizoslides were stored wick-side

down for one (1) hour post inoculation, then returned to their normal orientation so roots were

facing down for the remainder of the experiment. µ-rhizoslides were grouped by treatment and

positioned in the growth chamber as described above. Every 4th day, μ-rhizoslides were alternately

watered with either sterile diH$_2$O or a modified Hoagland's solution (19).

Imaging and Image Analysis. Once per week, we imaged the entire soil area of each μ-rhizoslide

using tiled brightfield and fluorescent images using a Zeiss AXIO-Observer Z1 inverted

microscope (Carl Zeiss Inc., Germany) with an AxioCam MRm Rev.3 camera (1348 × 1040 pixels,

6.45 × 6.45 μm/pixel). We used a 5× Zeiss EC Plan-Neofluar 5×/0.16 M27 lens and set the camera

to collect 585-binned photos (2 × 2 binning). The transmission-based brightfield light equipped on

the microscope could not illuminate through the 0.5 cm of soil, so we moved the brightfield lamp

to an external stand and focused it on the soil surface at the front of the μ-rhizoslide. For fluorescent

imaging, we found 570 nm excitation and 620 nm emission filters gave the strongest fluorescent

signal for *S. meliloti* Rm1021/pCMB35. We used the Zeiss Zen Pro 2.3 software to stitch a single

207-megapixel image (29341 × 7072 pixels) from each set of 585 photos. We then exported each

image as 12-bit grayscale TIFF at full resolution.

For image analysis, the stitched fluorescent images were imported into MATLAB and

converted into 29341 × 7072 matrices. Since we took steps to limit bacterial fluorescent

bloom on images taken the same day as the inoculation (Week 0), we were able to use the

fluorescent images from week 0 as a measure of background fluorescence from sources like the

3D printed spacer, and mica particles in the soil. We subtracted the Week 0 tiled fluorescent image

of each μ-rhizoslide from the tiled fluorescent images of all subsequent weeks to remove the

aforementioned background fluorescence. Any subsequent negative pixel value was set to 0. In

addition, tiled images were cropped to remove the non-soil portion, eliminating fluorescence from

the 3D printed spacers. As *S. meliloti* bacteria was predominately found towards the center of the

soil channel, along the roots, removing the background from the 3D spacer this way had a minimal

effect on the bacterial fluorescence signal.

We partitioned the photo matrixes into sections approximately 1 mm in length along the

root axis (386×7072 pixels), encompassing the full width of the soil region. Next, we determined

the fraction of the area of each section with a non-zero fluorescence signal. We define "normalized

fluorescent area" as the number of nonzero pixels divided by the total number of pixels in the

section. We then plotted the normalized fluorescent area versus distance of the section from the

soil-air interface (Fig 3). We define "furthest displacement" as the 1-mm partition with at least

10% normalized fluorescent area. We then plotted the mean furthest vertical and horizontal

displacement values of the five replicates for each treatment versus distance from the soil-air

interface with their standard deviation (Fig 4 & Fig 5). We used one-way ANOVA using the values

for furthest progress of the fluorescent signal for each treatment to determine the statistical

significance of the different treatments.

We created composite images for ease of visualization of the progress of the fluorescent

signal (Fig 2). We used the Week 3 brightfield images for each rhizoslide trial, and overlayed these

images with weekly greyscale fluorescent images colored coordinated to the week. These

composite images were not used for quantitative data analysis.

Post-imaging quantification of *Colpoda sp.* and *S. meliloti* in u-rhizoslides. After the final

imaging, μ-rhizoslides were disassembled and roots were separated from the bulk soil. Roots were

cut into "upper" and "lower" halves beginning 2.5 cm from the root/shoot interface. Root weight

was recorded, then the root sections were suspended in 10 times their weight in Page's medium in

50 ml conical tubes. Roots were vortexed for 1 minute at medium speed on a tabletop vortex mixer.

For bacteria abundance, 10-fold dilutions were made in Page's medium and 10 µL spots

were placed onto TY + tetracycline (5 µg/ml) agar plates and incubated at 30 °C for 3 days.

Colonies were counted, and the total number of *S. meliloti* strain Rm1021/pCMB35 was estimated

for each root section.

To estimate protist abundance, 3-fold dilutions were carried out in a 96 well microtiter dish

using Page's medium+ heat-killed *E. coli* DH5a (OD_{595} = 0.05). Eight replicate dilutions were

done for each root section. Microtiter dishes were incubated at room temperature in darkness for

1 week. Presence of protists in each well was recorded and assumed that "at least one" protist made

it into wells that contained protists. The Minimum Recovered Number per ml (MRN/ml) of protists

was back calculated as follows:

MRN/ml = $10 * 3^{(\text{farthest positive dilution factor} - 1)}$

Results

µ-rhizoslides allow convenient, long-term tracking of plant root development. µ-rhizoslides

were constructed as a smaller version of the rhizoslides designed by C. L. Marie (31), and

optimized for microscopy and increased through-put. Root growth and behavior is easily

monitored on a standard inverted microscopy with the smaller µ-rhizoslides. Plant roots occupy a

region 15×75 mm and can be readily imaged as a series of overlapping tiled images (Fig 2). When

not being imaged the µ-rhizoslides were kept at a 45-degree angle to promote root growth directly

against the glass slide. This successfully facilitated direct microscopy of the roots without

removing the surrounding soil. Brightfield microscopy was used to image root tissue of which

allowed monitoring of primary, secondary and tertiary roots. Images were collected once a week

for 4 weeks after planting. This demonstrated the ability of the μ-rhizoslide to support plant growth

for extended periods: a typical requirement of many root-growth experiments.

Distribution of *S. meliloti* on plant roots increases in the presence of protists. Inoculation of

seeds with beneficial bacteria, while effective under some circumstances, often fails to provide

maximum benefits to the grown plant because the bacteria are not effectively distributed along the

growing root system. Soil protists, which often prey on bacteria, may enhance bacterial movement

by physically moving them as a result of the feeding process, or by excreting still living bacteria

after they have been ingested. In order to test this hypothesis, the nitrogen fixing symbiont

Sinorhizobium meliloti strain Rm1021 was transformed with plasmid pCMB35 which directs the

constitutive expression of mRFP. The movement and growth of Rm1021/pCMB35 was tracked

by fluorescent microscopy in the μ-rhizoslides over the course of four weeks under three

conditions: Control (plant only); Bacteria (plant + Rm1021/pCMB35 only) and Bacteria + Protist

(plant + protist and Rm1021/pCMB35) (Fig 4). When comparing the progress of the fluorescent

signal, indicative of Rm1021/pCMB35, a significant difference between treatments was

demonstrated by ANOVA for all time points after Week 0 ($p < 0.005$). Rm1021/pCMB35 in the

Bacteria + Protist Treatment colonized an average of 44 mm more root when compared with the

other treatments by Week 2. Data from later time points showed increased variation in the

movement of Rm1021/pCMB35 along roots which was perhaps due to differing morphology of

the root systems between replicates that gave rise to variable opportunities for root colonization.

The lack of meaningful progress of the fluorescent signal in the Bacteria + Protist Treatment

between Weeks 2 and 3 may be due, in part, to loss of the fluorescent plasmid, which is usually

maintained with 5 μg/mL tetracycline selection. Tetracycline was not added to the μ-rhizoslides

to limit interference with plant development, or bacteria may be succumbing to increased predation

pressure due to increased numbers of protists.

We inoculated roots from the back of the μ-rhizoslides, thus the absence of fluorescent

signal in the Bacteria Only Treatment indicates that Rm1021/pCMB35 travelled less than the 5

mm distance between the back of the slide and the front when protists were absent. The higher

fluorescent signal for the Control Treatment compared with the Bacteria Only Treatment,

especially in Weeks 2 and 3, was likely due to root autofluorescence that was more pronounced

under this condition.

Quantification of protists and bacteria from plant roots corroborate imaging data and

indicate protist-assisted transport of inoculated bacteria. After imaging was complete, μ-

rhizoslides were disassembled and the plant roots were divided into upper and lower halves. Roots

and the attached soil were transferred to a volume of Page's Saline Solution that was 10x the mass

of the root sample and vortexed to disassociate microbial biomass from the roots. Dilutions were

spotted on a TY+Tc5 plates and incubated for 3 days to determine bacterial CFUs associate with

the root sections. Serial three-fold dilutions were also transferred to a 96-well plate containing

Page's Saline Solution and a heat-killed *E. coli* to promote protist growth and reproduction. After

one week, the <u>m</u>inimum <u>r</u>ecovered <u>n</u>umber (MRN) of protists was calculated from the highest

dilution that contained reproducing protists.

The overall bacterial abundance in the upper sections was equivalent in the Bacteria Only

and the Bacteria + Protist treatments despite predation by protists in the latter (Fig 6A). In the

lower section of the Bacteria Only treatment strain Rm1021/pCMB35 was not detected, while an

average of 1.14×10^5 CFUs of this strain, per 1.0 mg of root tissue, was detected in the lower

section of the Bacteria + Protist treatments (Fig 6A). It is important to note that while no

fluorescence was detected in the upper half of the Bacteria Only treated roots, Rm1021/pCMB35

was still recovered from the roots on which they were inoculated. Additionally, protists were only

observed in the Bacteria + Protist Treatment, with comparable numbers in both the upper and

lower sections (Fig 6B). More protists may have been present in these sections, as these values are

considered the minimum number of recovered protists, true values may be approximately 3x

higher because calculation of MRN assumes that the last well of a dilution series contained only

one protist at the start of the one-week growth period. These measurements also represent a

minimum because they represent bacteria and protists associated with roots; bulk soil which was

not evaluated by these methods likely contained additional organisms. Overall, these data indicate

that protists not only transported the bacteria along the roots, but that the bacteria were viable after

delivery to lower sections of the roots, something that could not be determined by imaging alone.

Discussion. While previous work has shown the potential of using protists as vehicles for

transporting particles (19), here we demonstrate their ability to transport beneficial bacteria in a

realistic soil system. This transport occurs despite bacterial population loss due to protist grazing.

Direct counts of bacteria and protists showed the same order-of-magnitude concentrations

of each in the upper sections of the Bacteria Only and the Bacteria + Protists treated μ-rhizoslides.

Contrary to this, the imaging experiments showed distinct movement of bacteria only in the

Bacteria + Protists treatment. This difference was likely seen because imaging allowed us to

discern only populations of bacteria that were growing at high densities, and direct counts allowed

us to quantify bacteria not directly assessable to the camera and those that were present in

concentrations too low to image. Importantly, *S. meliloti* strain Rm1021/pCMB35 was only

recovered from the lower halves of roots treated with Bacteria + Protists, supporting the hypothesis

that protists can assist in delivering beneficial bacteria throughout the rhizosphere allowing roots

to receive bacterial benefits much quicker and perhaps more uniformly than they would otherwise.

Additionally, visualization of bacterial distributions by fluorescent microscopy and image

analysis in the Bacteria + Protists treatment showed bacterial proliferation that extended both along

the root and horizontally from the root. For the Bacteria Only treatment, distribution was less than

5 millimeters from the initial inoculation site in any direction. On their own, bacteria did not spread

far enough from the initial inoculation site to reach the front of the μ-rhizoslide where the

fluorescence would be visible to the imaging equipment.

The protist used in this study, *Colpoda sp.*, a ciliate isolated from a legume rhizosphere, is

particularly successful in transporting bacteria. Ciliates commonly eat by means of filter feeding,

a mechanism of promiscuous consumption of food particles and prey. Bacteria consumed by the

ciliates are packaged into a phagosome which is processed and eventually egested from the protist;

a process that is not well characterized. It is thought that egested phagosomes break down in the

soil and may release viable bacteria which could go on to colonize the area where they were

released. How often bacteria can survive intake and egestion, and whether some species are adept

at this in not clearly understood. Some bacterial species, for example *Campylobacter jejuni* and

Mycobacterium spp., are known to escape digestion and colonize the cytoplasm of protists that

consumed them (28, 36). Ciliates have also been shown to emerge from feeding on bacterial

biofilms with a matrix of bacteria stuck to their cell bodies (19). When the protists travel along

plant roots consuming bacteria, both egestion of viable bacteria and physical translocation of

bacteria may contribute to protist-assisted transport.

The technology described here provides a simple, reproducible experimental system which

has the potential to aid in the understanding of rhizosphere dynamics. Of particular interest is

bacterial movement in the context of screening a variety of different bacteria/protist/plant/soil

combinations for their effects on such movement. For example, protist transportation of

Sinorhizobium sp. and its effect on nodulation can be studied using this system and parameters that

affect transport such as water saturation, soil composition and soil porosity can be readily and

tested and quantitated. Protist transport may also be adapted for targeted delivery of other microbes

or nano-coated agrochemicals. With increased understanding of such processes, and protist-

assisted transport mechanisms, we can begin developing new methods to support sustainable

agriculture that is more likely to function in the field.

Acknowledgements. This work was supported by DOE award DE-SC0014522 (to LMS and

DJG), NSF award 1605816 (to LMS and DJG), and USDA National Institute of Food and

Agriculture award 2016-67013-24412 (to DJG and LMS).

References

1. Berendsen RL, Pieterse CMJ, Bakker PAHM. 2012. The rhizosphere microbiome and plant health. Trends Plant Sci 7: 478-486.

2. van Veen JA, Liljeroth E, Lekkerkerk LJA, van de Geijn SC. 1991. Carbon fluxes in plant-soil systems at elevated atmospheric CO_2 levels. Ecol Appl 1:175–181.

3. Guo Y-S, Furrer JM, Kadilak AL, Hinestroza HF, Gage DJ, Cho YK, Shor LM. 2018. Bacterial extracellular polymeric substances amplify water content variability at the pore scale. Front Environ Sci 6:93.

4. Haas D, Défago G. 2005. Biological Control of Soil-borne Pathogens by Fluorescent Pseudomonads. Nat Rev Microbiol 3:307–319.

5. Baveye PC, Otten W, Kravchenko A, Balseiro-Romero M, Beckers É, Chalhoub M, Darnault C, Eickhorst T, Garnier P, Hapca S, Kiranyaz S, Monga O, Mueller CW, Nunan N, Pot V, Schlüter S, Schmidt H, Vogel H-J. 2018. Emergent properties of microbial activity in heterogeneous soil microenvironments: different research approaches are slowly converging, yet major challenges remain. Front Microbiol 9:1929.

6. Jiang X, Zerfaß C, Feng S, Eichmann R, Asally M, Schäfer P, Soyer OS. 2018. Impact of spatial organization on a novel auxotrophic interaction among soil microbes. ISME J 12:1443–1456.

7. Ben-Jacob E, Finkelshtein A, Ariel G, Ingham C. 2016. multispecies swarms of social microorganisms as moving ecosystems. Trends Microbiol 24:257–269.

8. Young IM, Crawford JW. 2004. Interactions and self-organization in the soil-microbe complex. Science 304:1634–7.

9. Cho H, Jönsson H, Campbell K, Melke P, Williams JW, Jedynak B, Stevens AM, Groisman A, Levchenko A. 2007. Self-organization in high-density bacterial colonies: efficient crowd control. PLoS Biol 5:e302.

10. Kim HJ, Boedicker JQ, Choi JW, Ismagilov RF. 2008. defined spatial structure stabilizes a synthetic multispecies bacterial community. Proc Natl Acad Sci 105:18188–18193.

11. Deng J, Orner EP, Chau JF, Anderson EM, Kadilak AL, Rubinstein RL, Bouchillon GM, Goodwin RA, Gage DJ, Shor LM. 2015. Synergistic effects of soil microstructure and bacterial EPS on drying rate in emulated soil micromodels. Soil Biol Biochem 83:116–124.

12. Krome K, Rosenberg K, Bonkowski M, Scheu S. 2009. grazing of protozoa on rhizosphere bacteria alters growth and reproduction of *Arabidopsis thaliana*. Soil Biol Biochem 41:1866–1873.

13. Bonkowski M, Cheng W, Griffiths BS, Alphei J, Scheu S. 2000. microbial-faunal interactions in the rhizosphere and effects on plant growth. Eur J Soil Biol 36:135–147.

14. Bonkowski M, Clarholm M. 2012. Stimulation of Plant Growth Through Interactions of
bacteria and protozoa: testing the auxiliary microbial loop hypothesis. Acta Protozool.
51(3):237-247. DOI: 10.4467/16890027AP.12.019.0765

15. Henkes GJ, Kandeler E, Marhan S, Scheu S, Bonkowski M. 2018. Interactions of mycorrhiza
and protists in the rhizosphere systemically alter microbial community composition, plant shoot-
to-root ratio and within-root system nitrogen allocation. Front Environ Sci 6:117.

16. Bouwman LA, Bloem J, van den Boogert PHJF, Bremer F, Hoenderboom GHJ, de Ruiter
PC. 1994. short-term and long-term effects of bacterivorous nematodes and nematophagous
fungi on carbon and nitrogen mineralization in microcosms. Biol Fertil Soils 17:249–256.

17. Bonkowski M. 2004. Protozoa and plant growth: the microbial loop in soil revisited. New
Phytol. 162: 617–631

18. Rønn R, McCaig AE, Griffiths BS, Prosser JI. 2002. Impact of protozoan grazing on
bacterial community structure in soil microcosms. Appl Environ Microbiol 68:6094–105.

19. Rubinstein RL, Kadilak AL, Cousens VC, Gage DJ, Shor LM. 2015. Protist-facilitated
particle transport using emulated soil micromodels. Environ Sci Technol 49:1384–1391.

20. Shor LM, Chau JF, Gage DJ. 2018. Methods and systems to replicate soil properties.

21. Gage DJ, Shor LM, Gage JL, Micciulla JL, Rubinstein RL. 2017. Microbial carriers for
targeted delivery of agricultural payloads. US9603368B2. US.

22. Humphris SN, Bengough AG, Griffiths BS, Kilham K, Rodger S, Stubbs V, Valentine TA,
Young IM. 2005. Root cap influences root colonization by *Pseudomonas fluorescens* SBW25 on
maize. FEMS Microbiol Ecol 54:123–130.

23. Stougaard J. 2000. regulators and regulation of legume root nodule development. Plant
Physiol 124:531 LP – 540.

24. Lund ZF. 1970. the effect of calcium and its relation to several cations in soybean root
growth. Soil Sci Soc Am J 34:456–459.

25. Korber DR, Lawrence JR, Sutton B, Caldwell DE. 1989. Effect of laminar flow velocity on
the kinetics of surface recolonization by Mot^+ and Mot^- *Pseudomonas fluorescens*. Microb Ecol
18:1–19.

26. Wang W, Shor LM, LeBoeuf EJ, Wikswo JP, Taghon GL, Kosson DS. 2008. Protozoan
migration in bent microfluidic channels. Appl Environ Microbiol 74:1945–9.

27. Wang W, Shor LM, LeBoeuf EJ, Wikswo JP, Kosson DS. 2005. Mobility of protozoa
through narrow channels. Appl Environ Microbiol 71:4628–4637.

28. First MR, Park NY, Berrang ME, Meinersmann RJ, Bernhard JM, Gast RJ, Hollibaugh JT.
2012. Ciliate Ingestion and Digestion: flow cytometric measurements and regrowth of a
digestion-resistant *Campylobacter jejuni*. J Eukaryot Microbiol 59:12–19.

29. Mutalik VK, Guimaraes JC, Cambray G, Lam C, Christoffersen MJ, Mai QA, Tran AB,
Paull M, Keasling JD, Arkin AP, Endy D. 2013. Precise and reliable gene expression via
standard transcription and translation initiation elements. Nat Methods 10:354–360.

30. Silva-Rocha R, Martínez-García E, Calles B, Chavarría M, Arce-Rodríguez A, De Las Heras
A, Páez-Espino AD, Durante-Rodríguez G, Kim J, Nikel PI, Platero R, De Lorenzo V. 2013. the
standard European vector architecture (SEVA): A coherent platform for the analysis and
deployment of complex prokaryotic phenotypes. Nucleic Acids Res 41: D666.

31. Le Marié C, Kirchgessner N, Marschall D, Walter A, Hund A. 2014. Rhizoslides: Paper-
based growth system for non-destructive, high throughput phenotyping of root development by
means of image analysis. Plant Methods 10: 13 (2014). https://doi.org/10.1186/1746-4811 10-13

32. Kadilak AL, Rehaag JC, Harrington CA, Shor LM. 2017. A 3D-printed microbial cell culture
platform with in situ PEGDA hydrogel barriers for differential substrate delivery.
Biomicrofluidics 11:054109.

33. Macdonald NP, Zhu F, Hall CJ, Reboud J, Crosier PS, Patton EE, Wlodkowic D, Cooper JM.
2016. Assessment of biocompatibility of 3D printed photopolymers using zebrafish embryo
toxicity assays. Lab Chip 16:291–7.

34. Taerum SJ, Steven B, Gage DJ, Triplett LR. 2020. Validation of a PNA clamping method for
reducing host DNA amplification and increasing eukaryotic diversity in rhizosphere microbiome
studies. Phytobiomes J 4:291-302. https://doi.org/10.1094/PBIOMES-05-20-0040-TA

35. Caspers H. 1978. F. C. Page: An illustrated key to freshwater and soil amoebae with notes on
cultivation and ecology. The Ferry House, Far Sawrey, Ambleside, Cumbria: Freshwater
Biological Association, Scientific Publication No. 34. 1976. ISSN 0367-1857. 63:289–289.

36. Adékambi T, Salah S Ben, Khlif M, Raoult D, Drancourt M. 2006. Survival of environmental
mycobacteria in *Acanthamoeba polyphaga*. Appl Environ Microbiol 72:5974–5981.

37. Muok, A.R., Claessen, D. & Briegel, A. (2021). Microbial hitchhiking: how *Streptomyces* spores are
transported by motile soil bacteria. ISME J: https://doi.org/10.1038/s41396-021-00952-8

38. A.R. Muok, A. Briegel. (2021) Intermicrobial Hitchhiking: How Nonmotile Microbes Leverage
Communal Motility. Trends in Microbiol 29: 542-550. https://doi.org/10.1016/j.tim.2020.10.005

39. William L. King, Terrence H. Bell. (2021). Can dispersal be leveraged to improve microbial inoculant
success? Trends in Biotech. https://doi.org/10.1016/j.tibtech.2021.04.008

40. Best, AM, Abu Kwaik, Y. Evasion of phagotrophic predation by protist hosts and innate immunity of
metazoan hosts by *Legionella pneumophila*. Cellular Microbiol. 2019; 21:e12971.
https://doi.org/10.1111/cmi.12971

Figure 1. µRhizoslide construction and use. A) CAD drawing of the rhizoslide assembly
showing layers including 3D printed rhizoslide spacer and filter paper wick between two glass
slides. B) Fully assembled rhizoslide mockup. C) Photograph of rhizoslide in use.

Figure 2 Fluorescent images of plants two weeks post-inoculation. A) Control treatment:
inoculated with 10 µL buffer only. B) Bacteria Only treatment: inoculated with 10 µL bacteria
suspension. C) Bacteria + Protist treatment: inoculated with 10 µL bacteria suspension with
Colpoda sp. added.

Figure 3 Imaging bacterial movement and growth following plant inoculation with bacteria or bacteria plus protists. For each replicate, the fraction of pixels registering fluorescence in each 1mm image slice is plotted. This normalized fluorescent area is plotted against distance along the root to show the spread of fluorescently labelled bacteria through the rhizosphere. The graphs for each treatment (Control, Bacteria Only, Bacteria + Protists) are plotted in rows, while weeks are shown in columns. Within each graph, replicates are shown by different color bars.

Figure 4 Position of furthest progress of the bacterial fluorescent signal following plant inoculation with bacteria or bacteria plus protists. Marked locations indicate the furthest location along the plant root axis where the fluorescent signal was at least 10% of the total area. A) Control treatment shows no significant shift in position with time ($p > 0.05$). Slight increase in fluorescent signal on some replicates is likely because of root autofluorescence due to nitrogen starvation. B) Bacteria Only treatment shows no shift in position of fluorescent signal with time ($p > 0.05$). C) Bacteria + Protist treatment shows approximately 12 mm of progress in fluorescent signal by Week 1, increasing to 44 mm on average by Week 2. Large deviations between replicates are likely due to root morphology. (n=9, error bars=1 STD).

Figure 5 Horizontal spread of the bacterial fluorescent signal following plant inoculation with bacteria or bacteria plus protists. The widest fluorescent area for each replicate was calculated for each week. Marked locations indicate the horizontal width of the fluorescent signal averaged across the replicates. Control and Bacteria Only treatments show widths below 2 mm, which is likely root autofluorescence. Bacteria + Protist treatment shows increasing fluorescence signal width between Weeks 0 and 2, while Weeks 2 and 3 are comparable. (n=9, error bars=1 STD).

498
Figure 6 Bacterial and protists counts performed three weeks following plant inoculation
with bacteria or bacteria plus protists. The upper and lower sections of *M. truncatula* root
systems were separated into top (TF) and bottom (BF) fractions and analyzed for: A) Average
bacterial CFUs per 1.0 mg root tissue and B) Average minimum recovered number (MRN) of
protists per 1.0 mg root tissue. Control (C) n=8, Bacteria Only (BO) n=9, Bacteria + Protists (BP)
n=9