SARS-CoV-2 spike P681R mutation, a hallmark of the Delta variant, enhances viral fusogenicity and pathogenicity

Akatsuki Saito[1,2,3,#], Takashi Irie[4,#], Rigel Suzuki[5,#], Tadashi Maemura[6,7#], Hesham Nasser[8,9,#], Keiya Uriu[10,#], Yusuke Kosugi[10,#], Kotaro Shirakawa[11], Kenji Sadamasu[12], Izumi Kimura[10], Jumpei Ito[10], Jiaqi Wu[13,14], Kiyoko Iwatsuki-Horimoto[6], Mutsumi Ito[6], Seiya Yamayoshi[6,15], Seiya Ozono[16], Erika P Butlertanaka[1], Yuri L Tanaka[1], Ryo Shimizu[8,17], Kenta Shimizu[5], Kumiko Yoshimatsu[18], Ryoko Kawabata[4], Takemasa Sakaguchi[4], Kenzo Tokunaga[16], Isao Yoshida[12], Hiroyuki Asakura[12], Mami Nagashima[12], Yasuhiro Kazuma[11], Ryosuke Nomura[11], Yoshihito Horisawa[11], Kazuhisa Yoshimura[12], Akifumi Takaori-Kondo[11], Masaki Imai[6,15], The Genotype to Phenotype Japan (G2P-Japan) Consortium, So Nakagawa[13,14]*, Terumasa Ikeda[8]*, Takasuke Fukuhara[5]*, Yoshihiro Kawaoka[6,7,15]*, Kei Sato[8,12]*

[1] Department of Veterinary Science, Faculty of Agriculture, University of Miyazaki, Miyazaki 8892192, Japan
[2] Center for Animal Disease Control, University of Miyazaki, Miyazaki 8892192, Japan
[3] Graduate School of Medicine and Veterinary Medicine, University of Miyazaki, Miyazaki 8892192, Japan
[4] Institute of Biomedical and Health Sciences, Hiroshima University, Hiroshima 7348551, Japan
[5] Department of Microbiology and Immunology, Graduate School of Medicine, Hokkaido University, Hokkaido 0608638, Japan
[6] Division of Virology, Institute of Medical Science, University of Tokyo, Tokyo 1088639, Japan
[7] Influenza Research Institute, Department of Pathobiological Sciences, School of Veterinary Medicine, University of Wisconsin-Madison, WI 53711, USA
[8] Division of Molecular Virology and Genetics, Joint Research Center for Human Retrovirus infection, Kumamoto University, Kumamoto 8600811, Japan
[9] Department of Clinical Pathology, Faculty of Medicine, Suez Canal University, Ismailia 41511, Egypt
[10] Division of Systems Virology, Department of Infectious Disease Control, International Research Center for Infectious Diseases, The Institute of Medical Science, The University of Tokyo, Tokyo 1088639, Japan
[11] Department of Hematology and Oncology, Graduate School of Medicine, Kyoto University, Kyoto 6068507, Japan
[12] Tokyo Metropolitan Institute of Public Health, Tokyo 1690073, Japan
[13] Department of Molecular Life Science, Tokai University School of Medicine, Kanagawa 2591193, Japan

[14] CREST, Japan Science and Technology Agency, Saitama 3220012, Japan
[15] The Research Center for Global Viral Diseases, National Center for Global Health
and Medicine Research Institute, Tokyo 1628655, Japan
[16] Department of Pathology, National Institute of Infectious Diseases, Tokyo
1628640, Japan
[17] Graduate School of Medical Sciences, Kumamoto University, Kumamoto 8600811,
Japan
[18] Institute for Genetic Medicine, Hokkaido University, Hokkaido 0600815, Japan
49
[#] These authors contributed equally
*Correspondences:
so@tokai.ac.jp (S.N.)
ikedat@kumamoto-u.ac.jp (T.Ikeda)
fukut@pop.med.hokudai.ac.jp (T.F)
yoshihiro.kawaoka@wisc.edu (Y.Kawaoka)
KeiSato@g.ecc.u-tokyo.ac.jp (K.Sato)

57
58   **Conflict of interest**: The authors declare that no competing interests exist.
59   **Keywords**: SARS-CoV-2; COVID-19; spike protein; B.1.617.2; Delta variant;
60   P681R; fusion

Abstract

During the current SARS-CoV-2 pandemic, a variety of mutations have been accumulated in the viral genome, and currently, four variants of concerns (VOCs) are considered as the hazardous SARS-CoV-2 variants to the human society[1]. The newly emerging VOC, the B.1.617.2/Delta variant, closely associates with a huge COVID-19 surge in India in Spring 2021[2]. However, its virological property remains unclear. Here, we show that the B.1.617.2/Delta variant is highly fusogenic, and notably, more pathogenic than prototypic SARS-CoV-2 in infected hamsters. The P681R mutation in the spike protein, which is highly conserved in this lineage, facilitates the spike protein cleavage and enhances viral fusogenicity. Moreover, we demonstrate that the P681R-bearing virus exhibits higher pathogenicity than the parental virus. Our data suggest that the P681R mutation is a hallmark that characterizes the virological phenotype of the B.1.617.2/Delta variant and is closely associated with enhanced pathogenicity.

Main

In December 2019, an unusual infectious disease, now called COVID-19, emerged in Wuhan, Hubei province, China[3,4]. SARS-CoV-2, the causative agent of COVID-19, has rapidly spread all over the world, and as of July 2021, SARS-CoV-2 is an ongoing pandemic: more than 180 million cases of infections have been reported worldwide, and more than 4 million people died of COVID-19[1].

During the current pandemic, SARS-CoV-2 has acquired a variety of mutations[5]. First, in the spring of 2020, a SARS-CoV-2 derivative harbouring the D614G mutation in its spike (S) protein has emerged and quickly become predominant[6]. Because the D614G mutation increases viral infectivity, fitness, and inter-individual transmissibility[7-12], the D614G-bearing variant has quickly swept out the original strain. Since the fall of 2020, some SARS-CoV-2 variants bearing multiple mutations have emerged and rapidly spread worldwide. As of June 2021, there have been at least five variants of concern (VOC): B.1.1.7 (Alpha), B.1.351 (Beta), P.1 (Gamma), B.1.427/429 (Epsilon; note that this variant is currently out of concerns/interest) and B.1.617.2 (Delta), and these lineages respectively emerged in the UK, South Africa, Brazil, the USA and India[13,14].

At the end of 2020, the B.1.617 lineage has emerged in India, and this variant is thought to be a main driver of a massive COVID-19 surge in India, which has peaked 400,000 infection cases per day[2]. The B.1.617 lineage includes three sublineages, B.1.617.1, B.1.617.2 and B.1.617.3, and a sublineage, B.1.617.2, is the latest VOC, the Delta variant[13,14]. Importantly, early evidence have suggested that the B.1.617.2/Delta may have an increased risk of hospitalization compared to the B.1.1.7 cases[15,16]. However, the virological features of this newly emerging VOC, particularly its infectivity and pathogenicity, remain unclear. Additionally, although recent studies have shown that the B.1.617.2/Delta variant is relatively resistant to the neutralising antibodies (NAbs) elicited by vaccination[17,18], the mutation(s) that are responsible for the virological features of this VOC are unaddressed. In this study, we demonstrate that the B.1.617.2/Delta is more pathogenic than the prototypic SARS-CoV-2 in a Syrian hamster model. We also show that the P681R mutation in the S protein is a hallmark mutation of this lineage. The P681R mutation enhances the cleavage of SARS-CoV-2 S protein and enhances viral fusogenicity. Moreover, we demonstrate that the P681R mutation is responsible for the higher pathogenicity of the B.1.617.2/Delta variant *in vivo*.

Phylogenetic and epidemic dynamics of the B.1.617 lineage

We set out to investigate the phylogenetic relationship of the three subvariants belonging to the B.1.617 lineage. We downloaded 1,761,037 SARS-CoV-2 genomes

and information data from the Global Initiative on Sharing All Influenza Data
(GISAID) database (https://www.gisaid.org; as of May 31, 2021). As expected, each
of three sublineages, B.1.617.1, B.1.617.2 and B.1.617.3, formed a monophyletic
cluster, respectively (**Fig. 1a, Extended Data Fig. 1**). We then analyzed the
epidemic of each of three B.1.617 sublineages. The B.1.617 variant, particularly
B.1.617.1, was first detected in India on December 1, 2020 (GISAID ID:
EPI_ISL_1372093) (**Fig. 1b-d**). Note that a SARS-CoV-2 variant (GISAID ID:
EPI_ISL_2220643) isolated in Texas, the USA, on August 10, 2020, was also
recorded to belong to the B.1.617.1. However, the S protein of this viral sequence
(GISAID ID: EPI_ISL_2220643) possesses neither L452R nor P681R mutations,
both of which are the features of the B.1.617 lineage. Therefore, the
EPI_ISL_2220643 sequence isolated in the USA may not be the ancestor of the
current B.1.617.1 lineage, and the EPI_ISL_1372093 sequence obtained in India
would be the oldest B.1.617 lineage.

The B.1.617.2 (GISAID ID: EPI_ISL_2131509) and B.1.617.3 (GISAID IDs:
EPI_ISL_1703672, EPI_ISL_1703659, EPI_ISL_1704392) were detected in India on
December 10, 2020 and February 13, 2021, respectively (**Fig. 1e, f**). The B.1.617.1
sublineage has peaked during February to April, 2021, in India, and then decreased
(**Fig. 1d**). Although the B.1.617.3 variant has sporadically detected in India (**Fig. 1f**),
the B.1.617.2/Delta lineage has become dominant in India since March 2021 and
spread all over the world (**Fig. 1e**). At the end of May 2021, 100%, 70% and 43.3%
of the deposited sequences in GISAID per day from India (May 7), the UK (May 21)
and the whole world (May 19) have been occupied by the B.1.617.2 sublineage (**Fig.
1e and Extended Data Table 1**).

We next investigated the proportion of amino acid replacements in the S
protein of each B.1.617 sublineage comparing with the reference strain (Wuhan-Hu-
1; GenBank accession no. NC_045512.2). As shown in **Fig. 1g**, the L452R and
P681R mutations were highly conserved in the B.1.617 lineage, and notably, the
P681R mutation (16,650/16,759 sequences, 99.3%) was the most representative
mutation in this lineage. These data suggest that the P681R mutation is a hallmark
of the B.1.617 lineage.

Prominent syncytia formation by the B.1.617.2/Delta variant

To investigate the virological characteristics of the B.1.617.2/Delta variant, we
conducted virological experiments using a viral isolate of B.1.617.2 (GISAID ID:
EPI_ISL_2378732) as well as a D614G-bearing B.1.1 isolate (GISAID ID:
EPI_ISL_479681) in Japan. In Vero cells, the growth of the B.1.617.2/Delta variant
was significantly lower than that of the B.1.1 isolate (**Fig. 2a**). Particularly, the levels

of viral RNA of the B.1.617.2/Delta variant at 48 hours postinfection (hpi) was more
than 150-fold lower than that of the B.1.1 isolate (**Fig. 2a**). On the other hand,
although the growth kinetics of these viruses was relatively comparable in
VeroE6/TMPRSS2 cells and Calu-3 cells (**Fig. 2a**), microscopic observations
showed that the B.1.617.2/Delta formed larger syncytia compared to the B.1.1 virus
(**Fig. 2b**). By measuring the size of the floating syncytia in the infected
VeroE6/TMPRSS2 culture, the syncytia formed by the B.1.617.2/Delta infection
were significantly (2.7-fold) larger than that by the B.1.1 infection (**Fig. 2b**).
Immunofluorescence assay further showed that the B.1.617.2/Delta-infected
VeroE6/TMPRSS2 cells exhibit larger multinuclear syncytia compared to the B.1.1
isolate (**Extended Data Fig. 2**). These results suggest that the B.1.617.2/Delta
variant is feasible for forming syncytia compared to the D614G-bearing B.1.1 virus.
163

Higher pathogenicity of the B.1.617.2/Delta variant in Syrian hamsters
To investigate the pathogenicity of the B.1.617.2/Delta variant, we conducted
hamster infection experiments using the B.1.617.2/Delta isolate and the B.1.1 isolate.
Although the viral RNA loads in the oral swab of the B.1.617.2/Delta-infected
hamsters were significantly lower than those of the B.1.1-infected hamsters at 1 and
4 days postinfection (dpi), these values were comparable any other dpi (**Fig. 2c**).
After infection with these viruses, infected hamsters significantly lost their body
weights from 2 dpi. The peak weight loss was 16% for the B.1.617.2/Delta and 13%
for the B.1.1, with the B.1.617.2/Delta isolate having a significantly greater weight
loss than the B.1.1 at 3 and 4 dpi (**Fig. 2d**). These results suggest that the
B.1.617.2/Delta has a higher pathogenicity than the B.1.1 isolate, despite relatively
comparable proliferative potential.
176

P681R mutation as the determinant of enhanced and accelerated fusogenicity
The P681R mutation in the S protein is a unique feature of the B.1.617 lineage
including the B.1.617.2/Delta variant (**Fig. 1g**). Because the P681R mutation is
located in the proximity of the furin cleavage site (FCS; residues RRAR positioned
between 682-5) of the SARS-CoV-2 S protein[19], we hypothesized that the P681R
mutation is responsible for the preference of cell-cell fusion, which leads to larger
syncytia formation. To address this possibility, we generated the P681R-bearing
artificial virus by reverse genetics (**Extended Data Fig. 3**) and preformed virological
experiments. Although the amounts of viral RNA in the culture supernatants of the
D614G/P681R-infected Vero and VeroE6/TMPRSS2 cells were significantly lower
than those of the D614G-infected cells in some time points, the growth of these two
viruses was relatively comparable (**Fig. 3a**). However, the size of floating syncytia in

the D614G/P681R-infected VeroE6/TMPRSS2 cells at 72 hpi was significantly larger
than that in the D614G mutant-infected cells (**Fig. 3b**). This observation well
corresponds to that in the culture infected with the B.1.617.2/Delta variant (**Fig. 2b**).
To clearly observe the syncytia formation, we further generated the GFP-
expressing replication-competent D614G and D614G/P681R viruses. The levels of
viral RNA in the supernatant and GFP-positive cells were similar in Vero,
VeroE6/TMPRSS2 and Calu-3 cells (**Extended Data Fig. 4**). However, at 24 hpi,
significantly larger GFP-positive adherent syncytia were observed in the
VeroE6/TMPRSS2 cells infected with the GFP-expressing D614G/P681R virus (**Fig.
3c**). Additionally, the size of GFP-positive floating syncytia at 72 hpi in the
VeroE6/TMPRSS2 cells infected with GFP-expressing D614G/P681R virus was
significantly bigger than that with GFP-expressing D614G virus (**Extended Data Fig.
5**). Moreover, GFP-positive syncytia were observed in the D614G/P681R-infected
but not in the D614G-infected Calu-3 cells at 72 hpi (**Extended Data Fig. 4c**). These
results suggest that the feature of the B.1.617.2/Delta virus observed in *in vitro* cell
culture experiments, particularly forming larger syncytia (**Fig. 2b** and **Extended Data
Fig. 2**), is well reproduced by the insertion of P681R mutation.
To directly investigate the effect of P681R mutation on the cleavage of
SARS-CoV-2 S protein, we prepared the HIV-1-based pseudoviruses carrying the
P681R mutation. Western blotting of the pseudoviruses prepared showed that the
level of cleaved S2 subunit was significantly increased by the P681R mutation
(**Extended Data Fig. 6a**), suggesting that the P681R mutation facilitates the furin-
mediated cleavage of SARS-CoV-2 S protein. We then performed the single-round
pseudovirus infection assay using the target HOS cells with or without TMPRSS2
expression. The infectivity of both the D614G and the D614G/P681R pseudoviruses
increased approximately 10-fold by the expression of TMPRSS2 in target cells
(**Extended Data Fig. 6b**). However, the infectivity of the D614G and the
D614G/P681R pseudoviruses were comparable regardless of TMPRSS2
expression (**Extended Data Fig. 6b**). These data suggest that the P681R mutation
does not affect the infectivity of viral particles.
We next addressed the effect of P681R mutation on viral fusogenicity by
cell-based fusion assay. In the effector cells (i.e., S-expressing cells), although the
expression level of the D614G/P681R S protein was comparable to that of the
D614G S, the level of the cleaved S2 subunit of the D614G/P681R mutant was
significantly higher than that of the D614G S (**Fig. 3d**). Consistent with the results in
the pseudovirus assay (**Extended Data Fig. 6**), these results suggest that P681R
mutation facilitates the S cleavage. Flow cytometry showed that the surface
expression level of the D614G/P681R S was significantly lower than the D614G S

(**Extended Data Fig. 7**). Nevertheless, the cell-based fusion assay using the target
cells without TMPRSS2 demonstrated that the D614G/P681R S is 2.1-fold more
fusogenic than the D614G S with a statistical significance (P = 0.0002 by Welch's t
test) (**Fig. 3e**). Moreover, a mathematical modeling analysis of the fusion assay data
showed that the initial fusion velocity of the D614G/P681R S (0.83 ± 0.03 per hour)
is significantly (2.8-fold) faster than that of the D614G S (0.30 ± 0.03 per hour; P =
4.0×10^{-6} by Welch's t test) (**Fig. 3f, g**). These data suggest that the P681R mutation
enhances and accelerates the SARS-CoV-2 S-mediated fusion. Furthermore, when
we use the target cells with TMPRSS2 expression, both the fusion efficacy (~1.2-
fold) and initial fusion velocity (~2.0-fold) were increased in both the D614G and
D614G/P681R S proteins (**Fig. 3f, g**). These results suggest that TMPRSS2
facilitates the fusion mediated by SARS-CoV-2 S and human ACE2, while the
TMPRSS2-dependent acceleration and promotion of viral fusion is not specific for
the P681R mutant.
241
Resistance to NAb-mediated antiviral immunity by the P681R mutation
The resistance to the NAb in the sera of COVID-19 convalescents and vaccinated
individuals is a hallmark characteristic of the VOCs (reviewed in [20,21]), and it has
been recently showed that the B.1.617.2/Delta variant is relatively resistant to the
vaccine-induced neutralisation[17,18]. To ask whether the P681R mutation contributes
to this virological phenotype, we performed the neutralisation assay. The
D614G/P681R pseudovirus was partially (1.2-1.5-fold) resistant to the three
monoclonal antibodies targeting the receptor binding domain of SARS-CoV-2 S
protein (**Extended Data Fig. 8a**). Additionally, the neutralisation experiments using
the 19 sera of second BNT162b2 vaccination showed that the D614G/P681R
pseudovirus is significantly resistant to the vaccine-induced NAbs compared to the
D614G pseudovirus (P < 0.0001 by Wilcoxon matched-pairs signed rank test;
Extended Data Fig. 8b and 9). These results suggest that the P681R-bearing
pseudovirus is relatively resistant to NAbs. Notably, in contrast to the neutralising
activity against cell-free viruses, the SARS-CoV-2 S-based fusion assay showed that
cell-cell infection is strongly resistant to the NAbs and the insensitivity to the NAbs
on cell-cell infection is not dependent on the P681R mutation (**Extended Data Fig.
8c**). Altogether, these findings suggest that the P681R mutation confers the NAbs
resistance upon cell-free viral particles and cell-cell infection is resistant to the NAb-
mediated antiviral action compared to cell-free infection.
262
P681R mutation as the determinant of higher pathogenicity of the Delta variant

To assess the impact of the P681R mutation on viral replication and the
pathogenicity of SARS-CoV-2, Syrian hamsters were intranasally infected with the
D614G and D614G/P681R viruses. The D614G-infected hamsters exhibited no
weight loss, although a slight decrease in body weight was observed for one of the
animals by 7 dpi (5.0%) (**Fig. 4a**). In contrast, all of hamsters infected with the
D614G/P681R virus experienced a gradual body weight loss and the animals
showed a significant weight loss of 4.7-6.9% at 7 dpi compared to the D614G virus
(P = 0.011, **Fig. 4a**). We also assessed pulmonary function in infected hamsters by
measuring enhanced pause (PenH), which is a surrogate marker for
bronchoconstriction or airway obstruction, by using a whole-body plethysmography
system. Syrian hamsters infected with the D614G and D614G/P681R viruses
showed the increases in the lung PenH value (**Fig. 4b**). At 7 dpi, the D614G/P681R-
infected animals had significantly higher PenH values compared with those of the
D614G-infected animals (P = 0.043). At 3 dpi, both viruses replicated efficiently in
the lungs and nasal turbinates of the infected animals and no significant difference
in viral replication was observed between the two groups (**Fig. 4c**). At 7 dpi, no
differences in viral titres in the nasal turbinates were found between the two groups;
however, the lung titres in the D614G/P681R-infected group were significantly higher
than those in the D614G-infected groups (P = 0.0013, **Fig. 4c**).
283
Discussion
Previous studies have demonstrated the close association of the FCS in the SARS-
CoV-2 S protein with viral replication mode and it is dependent on TMPRSS2.
Johnson et al. and Peacock et al. showed that the loss of FCS results in the increase
of viral replication efficacy in Vero cells while the attenuation of viral growth in the
Vero cells expressing TMPRSS2[22,23]. On the contrary, here we showed that the
replication efficacy of the B.1.617.2/Delta variant was severely decreased in Vero
cells compared to VeroE6/TMPRSS2 cells. More importantly, although the FCS-
deleted SARS-CoV-2 is less pathogenic than the parental virus[23], we revealed that
the B.1.617.2/Delta variant as well as the P681R-harbouring virus exhibit higher
pathogenicity. These findings suggest that the enhanced viral fusogenicity, which is
triggered by the P681R mutation, closely associates with viral pathogenicity.
It is evident that most VOCs considered so far have acquired mutations in
their S proteins, particularly in the RBD and N-terminal domain, to evade NAbs[20,21,24].
In sharp contrast, here we demonstrated that the B.1.617.2/Delta variant has
acquired a unique strategy to facilitate infection and evade antiviral immunity. The
P681R mutation that is highly conserved in this lineage enhances the efficacy of viral
fusion and further accelerates its speed of action. The P681R-mediated rapid

kinetics of viral fusion may attribute to not only immune evasion but also possibly
feasible the infection to exposed individuals.

304   Consistent with previous reports[25,26], here we showed that the cell-cell
infection mediated by the SARS-CoV-2 S protein is resistant to NAbs. The effect of
NAbs against cell-cell infection has been well studied in HIV-1 (*Retroviridae*)
infection, and it is well known that cell-cell infection is relatively more resistant to
NAbs compared to cell-free infection (reviewed in [27-29]). The resistance of cell-cell
spread against NAbs is not limited to HIV-1 but has been observed in the other
viruses such as vaccinia virus (*Poxviridae*)[30] and hepatitis C virus (*Flaviviridae*)[31],
suggesting that cell-cell infection is a common strategy for a variety of viruses to
evade antiviral humoral immunity. The fact that the B.1.617.2/Delta variant as well
as the P681R mutant efficiently form syncytia and the P681R mutant accelerates
and promotes cell-cell fusion suggests that switching the preference of viral
replication mode from cell-free infection to cell-cell infection may be a unique
strategy of the B.1.617.2/Delta variant to evade antiviral immunity.

317   Although the P681R mutant is highly fusogenic, the virus harbouring the
P681R mutation did not necessarily show higher growth compared to the parental
virus in *in vitro* cell cultures. Regarding this, the HIV-1 variants with higher
fusogenicity have been isolated from AIDS patients, but the enhanced fusogenicity
does not promote viral replication in *in vitro* cell cultures[32]. Similarly, the measles
virus (*Paramyxoviridae*) harbouring the deficient mutation in viral matrix protein[33]
and substitution mutations in viral fusion protein[34,35] are highly fusogenic and
efficiently expands via cell-cell fusion. However, the growth kinetics of these mutated
measles virus with higher fusogenicity in *in vitro* cell cultures is less efficient than the
parental virus[33]. Therefore, the discrepancy between the efficacy of viral growth in
in vitro cell cultures and viral fusogenicity is not specific for SARS-CoV-2. Rather,
the higher fusogenicity is associated with the severity of viral pathogenicity such as
HIV-1 encephalitis[36] and fatal subacute sclerosing panencephalitis, which is caused
by measles virus infection in brain[34,35]. Consistently, here we showed that the
B.1.617.2/Delta variant as well as the P681R mutant exhibited a higher fusogenicity
in vitro and enhanced pathogenicity *in vivo*. Our data suggest that the higher COVID-
19 severity and unusual symptoms caused by the B.1.617.2/Delta variant[15,16] are
partly due to the higher fusogenicity caused by the P681R mutation. Switching viral
infection mode by the P681R mutation may relate to the severity and/or unusual
outcome of viral infection, therefore, the epidemic of the SARS-CoV-2 variants
harbouring the P681R mutation should be surveyed in depth.

Author Contributions

A.S., T.Irie, R.Suzuki, H.N., K.U., Y.Kosugi, E.P.B., Y.L.T., R.Shimizu, K.Shimizu, R.K., T.Ikeda, T.F. and K.Sato performed the experiments.

R.Suzuki, T.M., K.I.-H., M.Ito, S.Y., M.Imai, K.Yoshimatsu, T.F. and Y.Kawaoka performed animal experiments.

K.Sadamasu, S.O., T.S., K.T., I.Y., H.A., M.N., and K.Yoshimura prepared experimental materials.

J.W. and S.N. performed molecular phylogenetic analysis.

K.Shirakawa, Y.Kazuma, R.N., Y.H., and A.T.-K. collected clinical samples.

A.S., T.Irie, M.Imai, S.N., T.Ikeda, T.F. Y.Kawaoka and K.S. designed the experiments and interpreted the results.

K.Sato wrote the original manuscript.

All authors reviewed and proofread the manuscript.

The Genotype to Phenotype Japan (G2P-Japan) Consortium contributed to the project administration.

Consortia

The Genotype to Phenotype Japan (G2P-Japan) Consortium: Mika Chiba, Hirotake Furihata, Haruyo Hasebe, Kazuko Kitazato, Haruko Kubo, Naoko Misawa, Nanami Morizako, Akiko Oide, Mai Suganami, Miyoko Takahashi, Kana Tsushima, Miyabishara Yokoyama, Yue Yuan

Acknowledgments

We would like to thank all members belonging to The Genotype to Phenotype Japan (G2P-Japan) Consortium. We thank Dr. Jin Gohda (The University of Tokyo, Japan) for providing reagents. An anti-HIV-1 p24 Monoclonal antibody (clone 183-H12-5C, Cat# ARP-3537) was obtained through the NIH HIV Reagent Program, NIAID, NIH (contributed by Drs. Bruce Chesebro and Kathy Wehrly). The super-computing resource was provided by Human Genome Center at The University of Tokyo and the NIG supercomputer at ROIS National Institute of Genetics.

This study was supported in part by AMED Research Program on Emerging and Re-emerging Infectious Diseases 20fk0108163 (to A.S.), 20fk0108401 (to T.F.), 21fk0108617 (to T.F.), 19fk0108113 (to Y.Kawaoka), JP20fk0108412 (to Y.Kawaoka), 20fk0108146 (to K.Sato), 20fk0108270 (to K.Sato) and 20fk0108413 (to T.Ikeda, S.N. and K.Sato); AMED Research Program on HIV/AIDS 21fk0410033 (to A.S.) and 21fk0410039 (to K.Sato); AMED Japan Program for Infectious Diseases Research and Infrastructure 20wm0325009 (to A.S.), 21wm0325009 (to A.S.) and 21wm0125002 (to Y.Kawaoka); JST A-STEP JPMJTM20SL (to T.Ikeda);

JST SICORP (e-ASIA) JPMJSC20U1 (to K.Sato); JST SICORP JPMJSC21U5 (to
K.Sato), JST CREST JPMJCR20H6 (to S.N.) and JPMJCR20H4 (to K.Sato); JSPS
KAKENHI Grant-in-Aid for Scientific Research C 19K06382 (to A.S.), 18K07156 (to
K.T.) and 21K07060 (to K.T.), Scientific Research B 18H02662 (to K.Sato) and
21H02737 (to K.Sato); JSPS Fund for the Promotion of Joint International Research
(Fostering Joint International Research) 18KK0447 (to K.Sato); JSPS Core-to-Core
Program JPJSCCA20190008 (A. Advanced Research Networks) (to K.Sato); JSPS
Research Fellow DC1 19J20488 (to I.K.); JSPS Leading Initiative for Excellent
Young Researchers (LEADER) (to T.Ikeda); ONO Medical Research Foundation (to
K.Sato); Ichiro Kanehara Foundation (to K.Sato); Lotte Foundation (to K.Sato);
Mochida Memorial Foundation for Medical and Pharmaceutical Research (to
K.Sato); Daiichi Sankyo Foundation of Life Science (to K.Sato); Sumitomo
Foundation (to K.Sato); Uehara Foundation (to K.Sato); Takeda Science Foundation
(to T.Ikeda and K.Sato); The Tokyo Biochemical Research Foundation (to K.Sato);
Mitsubishi Foundation (to T.Ikeda); Shin-Nihon Foundation of Advanced Medical
Research (to T.Ikeda); Tsuchiya Foundation (to T.Irie); a Grant for Joint Research
Projects of the Research Institute for Microbial Diseases, Osaka University (to A.S.);
an intramural grant from Kumamoto University COVID-19 Research Projects
(AMABIE) (to T.Ikeda); Intercontinental Research and Educational Platform Aiming
for Eradication of HIV/AIDS (to T.Ikeda); and Joint Usage/Research Center program
of Institute for Frontier Life and Medical Sciences, Kyoto University (to K.Sato).

References

WHO. "Coronavirus disease 2019". https://www.who.int/emergencies/diseases/novel-coronavirus-2019. (2021).

Singh, J., Rahman, S. A., Ehtesham, N. Z., Hira, S. & Hasnain, S. E. SARS-CoV-2 variants of concern are emerging in India. *Nat Med*, doi:10.1038/s41591-021-01397-4 (2021).

Wu, F. *et al.* A new coronavirus associated with human respiratory disease in China. *Nature* **579**, 265-269, doi:10.1038/s41586-020-2008-3 (2020).

Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* **579**, 270-273, doi:10.1038/s41586-020-2012-7 (2020).

Plante, J. A. *et al.* The variant gambit: COVID's next move. *Cell Host Microbe* **29**, 508-515, doi:10.1016/j.chom.2021.02.020 (2021).

Korber, B. *et al.* Tracking changes in SARS-CoV-2 spike: evidence that D614G increases infectivity of the COVID-19 virus. *Cell* **182**, 812-827, doi:10.1016/j.cell.2020.06.043 (2020).

Zhou, B. *et al.* SARS-CoV-2 spike D614G change enhances replication and transmission. *Nature* **592**, 122-127, doi:10.1038/s41586-021-03361-1 (2021).

Ozono, S. *et al.* SARS-CoV-2 D614G spike mutation increases entry efficiency with enhanced ACE2-binding affinity. *Nat Commun* **12**, 848, doi:10.1038/s41467-021-21118-2 (2021).

Yurkovetskiy, L. *et al.* Structural and functional analysis of the D614G SARS-CoV-2 spike protein variant. *Cell* **183**, 739-751 e738, doi:10.1016/j.cell.2020.09.032 (2020).

Hou, Y. J. *et al.* SARS-CoV-2 D614G variant exhibits efficient replication ex vivo and transmission in vivo. *Science* **370**, 1464-1468, doi:10.1126/science.abe8499 (2020).

Plante, J. A. *et al.* Spike mutation D614G alters SARS-CoV-2 fitness. *Nature*, doi:10.1038/s41586-020-2895-3 (2020).

Volz, E. *et al.* Evaluating the effects of SARS-CoV-2 spike mutation D614G on transmissibility and pathogenicity. *Cell* **184**, 64-75 e11, doi:10.1016/j.cell.2020.11.020 (2021).

CDC. "SARS-CoV-2 variant classifications and definitions (updated June 10, 2021)". https://www.cdc.gov/coronavirus/2019-ncov/variants/variant-info.html. (2021).

WHO. "Tracking SARS-CoV-2 variants". https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/. (2021).

PHE. Public Health England. "3 June 2021 Risk assessment for SARS-CoV-

2 variant: Delta (VOC-21APR-02, B.1.617.2)"
https://assets.publishing.service.gov.uk/government/uploads/system/upload
s/attachment_data/file/991135/3_June_2021_Risk_assessment_for_SARS-
CoV-2_variant_DELTA.pdf. (2021).

16   Sheikh, A. *et al.* SARS-CoV-2 Delta VOC in Scotland: demographics, risk of
hospital admission, and vaccine effectiveness. *Lancet* **397**, 2461-2462,
doi:10.1016/S0140-6736(21)01358-1 (2021).

17   Wall, E. C. *et al.* Neutralising antibody activity against SARS-CoV-2 VOCs
B.1.617.2 and B.1.351 by BNT162b2 vaccination. *Lancet*,
doi:10.1016/S0140-6736(21)01290-3 (2021).

18   Planas, D. *et al.* Reduced sensitivity of SARS-CoV-2 variant Delta to antibody
neutralization. *Nature*, doi:10.1038/s41586-021-03777-9 (2021).

19   Shang, J. *et al.* Cell entry mechanisms of SARS-CoV-2. *Proc Natl Acad Sci*
U S A **117**, 11727-11734, doi:10.1073/pnas.2003138117 (2020).

20   Harvey, W. T. *et al.* SARS-CoV-2 variants, spike mutations and immune
escape. *Nat Rev Microbiol*, doi:10.1038/s41579-021-00573-0 (2021).

21   Corti, D., Purcell, L. A., Snell, G. & Veesler, D. Tackling COVID-19 with
neutralizing monoclonal antibodies. *Cell*, doi:10.1016/j.cell.2021.05.005
(2021).

22   Peacock, T. P. *et al.* The furin cleavage site in the SARS-CoV-2 spike protein
is required for transmission in ferrets. *Nat Microbiol*, doi:10.1038/s41564-021-
00908-w (2021).

23   Johnson, B. A. *et al.* Loss of furin cleavage site attenuates SARS-CoV-2
pathogenesis. *Nature* **591**, 293-299, doi:10.1038/s41586-021-03237-4
(2021).

24   Piccoli, L. *et al.* Mapping Neutralizing and Immunodominant Sites on the
SARS-CoV-2 Spike Receptor-Binding Domain by Structure-Guided High-
Resolution Serology. *Cell* **183**, 1024-1042 e1021,
doi:10.1016/j.cell.2020.09.037 (2020).

25   Kruglova, N., Siniavin, A., Gushchin, V. & Mazurov, D. SARS-CoV-2 cell-to-
cell infection is resistant to neutralizing antibodies. *BioRxiv*, 442701 (2021).

26   Xia, S. *et al.* Inhibition of SARS-CoV-2 (previously 2019-nCoV) infection by a
highly potent pan-coronavirus fusion inhibitor targeting its spike protein that
harbors a high capacity to mediate membrane fusion. *Cell Res* **30**, 343-355,
doi:10.1038/s41422-020-0305-x (2020).

27   Dufloo, J., Bruel, T. & Schwartz, O. HIV-1 cell-to-cell transmission and broadly
neutralizing antibodies. *Retrovirology* **15**, 51, doi:10.1186/s12977-018-0434-
1 (2018).

28 Agosto, L. M., Uchil, P. D. & Mothes, W. HIV cell-to-cell transmission: effects
on pathogenesis and antiretroviral therapy. *Trends Microbiol* **23**, 289-295,
doi:10.1016/j.tim.2015.02.003 (2015).

29 Sattentau, Q. Avoiding the void: cell-to-cell spread of human viruses. *Nat Rev*
Microbiol **6**, 815-826, doi:10.1038/nrmicro1972 (2008).

30 Law, M., Hollinshead, R. & Smith, G. L. Antibody-sensitive and antibody-
resistant cell-to-cell spread by vaccinia virus: role of the A33R protein in
antibody-resistant spread. *J Gen Virol* **83**, 209-222, doi:10.1099/0022-1317-
83-1-209 (2002).

31 Timpe, J. M. *et al.* Hepatitis C virus cell-cell transmission in hepatoma cells in
the presence of neutralizing antibodies. *Hepatology* **47**, 17-24,
doi:10.1002/hep.21959 (2008).

32 Sterjovski, J. *et al.* Asn 362 in gp120 contributes to enhanced fusogenicity by
CCR5-restricted HIV-1 envelope glycoprotein variants from patients with
AIDS. *Retrovirology* **4**, 89, doi:10.1186/1742-4690-4-89 (2007).

33 Cathomen, T. *et al.* A matrix-less measles virus is infectious and elicits
extensive cell fusion: consequences for propagation in the brain. *EMBO J* **17**,
3899-3908, doi:10.1093/emboj/17.14.3899 (1998).

34 Ikegame, S. *et al.* Fitness selection of hyperfusogenic measles virus F
proteins associated with neuropathogenic phenotypes. *Proc Natl Acad Sci U*
S A **118**, doi:10.1073/pnas.2026027118 (2021).

35 Watanabe, S. *et al.* Mutant fusion proteins with enhanced fusion activity
promote measles virus spread in human neuronal cells and brains of suckling
hamsters. *J Virol* **87**, 2648-2659, doi:10.1128/JVI.02632-12 (2013).

36 Rossi, F. *et al.* The V1-V3 region of a brain-derived HIV-1 envelope
glycoprotein determines macrophage tropism, low CD4 dependence,
increased fusogenicity and altered sensitivity to entry inhibitors. *Retrovirology*
5, 89, doi:10.1186/1742-4690-5-89 (2008).

37 Matsuyama, S. *et al.* Enhanced isolation of SARS-CoV-2 by TMPRSS2-
expressing cells. *Proc Natl Acad Sci U S A* **117**, 7001-7003,
doi:10.1073/pnas.2002589117 (2020).

38 Ferreira, I. *et al.* SARS-CoV-2 B.1.617 emergence and sensitivity to vaccine-
elicited antibodies. *BioRxiv*, 443253 (2021).

39 Torii, S. *et al.* Establishment of a reverse genetics system for SARS-CoV-2
using circular polymerase extension reaction. *Cell Rep* **35**, 109014 (2021).

40 Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software
version 7: improvements in performance and usability. *Mol Biol Evol* **30**, 772-
780, doi:10.1093/molbev/mst010 (2013).

41 Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for
Phylogenetic Inference in the Genomic Era. *Mol Biol Evol* **37**, 1530-1534,
doi:10.1093/molbev/msaa015 (2020).
42 Reed, L. J. & Muench, H. A Simple Method of Estimating Fifty Percent
Endpoints. *Am J Hygiene* **27**, 493-497 (1938).
43 Motozono, C. *et al.* An emerging SARS-CoV-2 mutant evading cellular
immunity and increasing viral infectivity. *BioRxiv*, 438288 (2021).
44 Rausch, T., Fritz, M. H., Untergasser, A. & Benes, V. Tracy: basecalling,
alignment, assembly and deconvolution of sanger chromatogram trace files.
BMC Genomics **21**, 230, doi:10.1186/s12864-020-6635-8 (2020).
45 Shema Mugisha, C. *et al.* A simplified quantitative real-time PCR assay for
monitoring SARS-CoV-2 growth in cell culture. *mSphere* **5**,
doi:10.1128/mSphere.00658-20 (2020).
46 Niwa, H., Yamamura, K. & Miyazaki, J. Efficient selection for high-expression
transfectants with a novel eukaryotic vector. *Gene* **108**, 193-199,
doi:10.1016/0378-1119(91)90434-d (1991).
47 Ozono, S., Zhang, Y., Tobiume, M., Kishigami, S. & Tokunaga, K. Super-rapid
quantitation of the production of HIV-1 harboring a luminescent peptide tag. *J
Biol Chem* **295**, 13023-13030, doi:10.1074/jbc.RA120.013887 (2020).
48 Ikeda, T. *et al.* HIV-1 restriction by endogenous APOBEC3G in the myeloid
cell line THP-1. *J Gen Virol* **100**, 1140-1152, doi:10.1099/jgv.0.001276 (2019).
49 Ikeda, T. *et al.* HIV-1 adaptation studies reveal a novel Env-mediated
homeostasis mechanism for evading lethal hypermutation by APOBEC3G.
PLoS Pathog **14**, e1007010, doi:10.1371/journal.ppat.1007010 (2018).
50 Sultana, T. *et al.* Multiple Pathways To Avoid Beta Interferon Sensitivity of
HIV-1 by Mutations in Capsid. *J Virol* **93**, doi:10.1128/JVI.00986-19 (2019).
51 Kondo, N., Miyauchi, K. & Matsuda, Z. Monitoring viral-mediated membrane
fusion using fluorescent reporter methods. *Curr Protoc Cell Biol* **Chapter 26**,
Unit 26 29, doi:10.1002/0471143030.cb2609s50 (2011).
540

Figures & figure legends

Fig. 1. Molecular phylogenetic and epidemic dynamics of the B.1.617 lineage
during the pandemic.

a, A phylogenetic tree of the B.1.617 lineage. Bar, 0.0002 substitutions per site.
Bootstrap values, **, 100%; *, >70%. The uncollapsed tree is shown in **Extended**
Data Fig. 1.

b-f, Epidemic dynamics of the B.1.617 lineage. **b,** The numbers of sequences
deposited in GISAID per day for India (orange, left), UK (blue, middle), and the whole
world (gray, right). **c-f,** The percentages of each lineage deposited per day (**c**, all
B.1.617; **d**, B.1.617.1; **e**, B.1.617.2/Delta; **f**, B.1.617.3) from India (orange), the UK
(blue) and the whole world (gray) are shown. The date first identified is indicated.
The raw data are summarized in **Extended Data Table 1**.

g, Proportion of amino acid replacements in the B.1.617 lineage. The top 10
replacements conserved in the S protein of the B.1.617 and its sublineages are
summarized. The number in parenthesis indicates the number of sequences
included in each panel. The raw data are summarized in **Extended Data Table 2**.

Fig. 2. Virological features of the B.1.617.2/Delta variant *in vitro* and *in vivo*.
a, Growth kinetics of the B.1.617.2/Delta variant and a B.1.1 isolate. A viral isolate of B.1.617.2/Delta (GISAID ID: EPI_ISL_2378732) and a D614G-bearing B.1.1 isolate (GISAID ID: EPI_ISL_479681) [100 50% tissue culture infectious dose ($TCID_{50}$) for Vero cells and VeroE6/TMPRSS2 cells, 1,000 $TCID_{50}$ for Calu-3 cells] were inoculated and the copy number of viral RNA in the culture supernatant was quantified by real-time RT-PCR. The growth curves of the viruses inoculated are shown. Assays were performed in quadruplicate.
b, Syncytia formation. The syncytia in infected VeroE6/TMPRSS2 cells were observed at 72 hpi. (Top) Representative bright-field images of VeroE6/TMPRSS2 cells at 72 hpi are shown. Bars, 100 μm. (Bottom) The size of floating syncytia in B.1.1-infected (n = 217) and B.1.617.2/Delta-infected (n = 217) cultures are shown. The size of the floating single cells in uninfected culture (n = 177) was also shown as a negative control.
c, d, Syrian hamster infection with the B.1.617.2/Delta variant. Male hamsters were infected with 10^5 $TCID_{50}$ of the B.1.1 isolate (n = 6) and the B.1.617.2/Delta isolate (n = 12). Four hamsters at the same age were used for mock infection. The amount of viral RNA in the oral swab (**c**) and body weight (**d**) and were routinely measured.
In **a**, statistically significant differences (*, $P < 0.05$) versus the B.1.1 isolate were determined by Student's *t* test. NS, no statistical significance.
In **b**, statistically significant differences versus the B.1.1-infected culture (*, $P < 0.05$) and uninfected culture (#, $P < 0.05$) were determined by the Mann-Whitney U test.

In **c** and **d**, statistically significant differences were determined by the Mann-Whitney
U test, and those versus uninfected hamsters (*, $P < 0.05$) are indicated by asterisks.
The P value between the B.1.1 and the B.1.617.2/Delta at each dpi is indicated in
the figure.

Fig. 3. Virological features of the P681R-harbouring virus *in vitro.*

a, Growth kinetics of artificially generated viruses. The D614G and D614G/P681R
mutant viruses were generated by reverse genetics. These viruses (100 TCID$_{50}$)
were inoculated into Vero cells and VeroE6/TMPRSS2 cells, and the copy number
of viral RNA in the culture supernatant was quantified by real-time RT-PCR. The
growth curves of the viruses inoculated are shown. Assays were performed in
quadruplicate.

b, c, Syncytia formation. **b,** (Top) Floating syncytia in VeroE6/TMPRSS2 cells
infected with the D614G and D614G/P681R mutant viruses at 72 hpi are shown.
Bars, 200 µm. (Bottom) The size of floating syncytia in the D614G mutant-infected
(n = 63) and the D614G/P681R mutant-infected (n = 126) cultures are shown. **c,**
(Left) Adherent syncytia in VeroE6/TMPRSS2 cells infected with the GFP-
expressing D614G and D614G/P681R mutant viruses at 24 hpi are shown. Areas
enclosed with squares are enlarged in the bottom panels. Bars, 200 µm. (Right) The
size of adherent GFP$^+$ syncytia in the D614G mutant-infected (n = 111) and the
D614G/P681R mutant-infected (n = 126) cultures are shown.

d, Western blotting of the S-expressing cells. (Left) Representative blots of SARS-
CoV-2 full-length S and cleaved S2 proteins as well as ACTB as an internal control.
kDa, kilodalton. (Right) The ratio of S2 to the full-length S plus S2 proteins in the S-
expressing cells.

e-g, SARS-CoV-2 S-based fusion assay. Effector cells (S-expressing cells) and
target cells (ACE2-expressing cells or ACE2/TMPRSS2-expressing cells) were
prepared, and the fusion activity was measured as described in **Methods**. **e,** Kinetics

of fusion activity (experimental data). Assays were performed in quadruplicate, and

fusion activity (arbitrary unit) is shown. **f,** The kinetics of fusion velocity estimated by

a mathematical model based on the kinetics of fusion activity data (see **Methods**).

g, Initial velocity of the S-mediated fusion.

In **b,c**, statistically significant differences versus the D614G mutant-infected culture

(*, $P < 0.05$) and uninfected culture (#, $P < 0.05$) were determined by the Mann-

Whitney U test.

In **d**, a statistically significant difference (*, $P < 0.05$) versus the D614G S was

determined by Student's *t* test.

In **g,** statistically significant differences (*, $P < 0.05$) were determined by two-sided

Welch's *t* test.

Fig. 4. Enhanced pathogenicity by the P681R mutation in hamsters.

Syrian hamsters were intranasally inoculated with 10^4 TCID$_{50}$ (in 30 µl) of the D614G and D614G/P681R viruses.

a, Body weight changes in hamsters after viral infection. Body weights of virus-infected (n = 4 each) and uninfected hamsters (n = 3) were monitored daily for 7 days.

b, Pulmonary function analysis in infected hamsters. Enhanced pause (PenH), which is a surrogate marker for bronchoconstriction or airway obstruction, was measured by using whole-body plethysmography.

c, Virus replication in infected hamsters. Four hamsters per group were euthanized at 3 and 7 dpi for virus titration. Virus titres in the lungs (left) nasal turbinates (right) were determined by the plaque assay using VeroE6/TMPRSS2 cells. Points indicate data from individual Syrian hamsters.

Statistically significant differences were determined by the Mann-Whitney U test, and those versus uninfected hamsters (*, $P < 0.05$) are indicated by asterisks. The P value between the D614G and the D614G/P681R at each dpi is indicated in the figure.

Extended Data Fig. 1. A maximum-likelihood based phylogenetic tree of the
representative 334 SARS-CoV-2 sequences that belong to the B.1.617 lineage.
GISAID ID, exposure country, and sampling date were noted in each terminal node.
The country isolated (India, UK, or the other countries) and the PANGO sublineage
are labeled by colors as indicated in the figure. Red or blue circle on the branch was
shown in each internal node if the bootstrap value was ≥ 80 or ≥ 50 (n = 1,000).

Extended Data Fig. 2. Immunofluorescence staining of B.1.617.2/Delta-infected VeroE6/TMPRSS2 cells.

VeroE6/TMPRSS2 cells infected with the viruses indicated [multiplicity of infection (MOI) 0.01] were stained with anti-SARS-CoV-2 nucleocapsid (N) (green) and Hoechst (blue). (Top) Representative images at 48 hpi are shown. Areas enclosed with squares are enlarged in the bottom panels. DIC, differential interference contrast. Bars, 200 μm for low magnification panels; 50 μm for high magnification panels. (Bottom) The area of N-positive cells in B.1.1-infected (n = 50) and B.1.617.2/Delta-infected (n = 50) cultures are shown. A statistically significant difference versus the B.1.1-infected culture (*, $P < 0.05$) was determined by the Mann-Whitney U test.

Extended Data Fig. 3. Chromatograms of the mutated regions of SARS-CoV-2
viruses artificially generated by reverse genetics.
Chromatograms of nucleotide positions 23,399-23,407 (left) and 23,600-23,608
(right) of parental SARS-CoV-2 (strain WK-521, PANGO lineage A; GISAID ID:
EPI_ISL_408667) and the D614G (A23403G in nucleotide) and P681R (C23604G
in nucleotide) mutation are shown.

672
673
Extended Data Fig. 4. Growth kinetics of artificially generated GFP-expressing
viruses.
The GFP-expressing D614G and D614G/P681R mutant viruses were generated by
reverse genetics. These viruses (100 $TCID_{50}$ for Vero and VeroE6/TMPRSS2 cells,
100 or 1,000 $TCID_{50}$ for Calu-3 cells) were inoculated. The copy number of viral RNA
in the culture supernatant (**a**) and the level of GFP-positive cells (the percentage of
GFP-positive cells, for Vero and VeroE6/TMPRSS2 cells; the GFP intensity per well,
for Calu-3 cells) (**b**) are shown. (**c**) Representative images of the Calu-3 cells
infected with GFP-expressing viruses (100 $TCID_{50}$). Areas enclosed with circles are
enlarged in the right panels. Assays were performed in quadruplicate. Statistically
significant differences (*, $P < 0.05$) versus the D614G virus were determined by
Student's *t* test. NS, no statistical significance.

Extended Data Fig. 5. Syncytia formation in VeroE6/TMPRSS2 cells infected

with GFP-expressing viruses.

(Left) Floating syncytia in VeroE6/TMPRSS2 cells infected with the GFP-expressing

D614G and D614G/P681R mutant viruses (100 TCID$_{50}$) at 72 hpi are shown. Bars,

100 µm. (Right) The size of adherent GFP$^+$ syncytia in the D614G mutant-infected

(n = 147) and the D614G/P681R mutant-infected (n = 171) cultures are shown.

A statistically significant difference versus the D614G mutant-infected culture (*, $P <$

0.05) was determined by the Mann-Whitney U test.

Extended Data Fig. 6. Infectivity of the P681R-bearing pseudovirus.

a, Western blotting of pseudoviruses. (Left) Representative blots of SARS-CoV-2 full-length S and cleaved S2 proteins as well as HIV-1 p24 capsid as an internal control. kDa, kilodalton. (Right) The ratio of S2 to the full-length S plus S2 proteins on pseudovirus particles.

b, Pseudovirus assay. The HIV-1-based reporter virus pseudotyped with the SARS-CoV-2 S D614G or D614G/P681R was inoculated into HOS-ACE2 cells or HOS-ACE2/TMPRSS2 cells at 4 different doses (125, 250, 500 and 1,000 ng HIV-1 p24 antigen). Percentages of infectivity compared to the virus pseudotyped with parental S D614G (1,000 ng HIV-1 p24) in HOS-ACE2 cells are shown. The numbers on the bars of the HOS-ACE2/TMPRSS2 cell data indicate the fold change versus the HOS-ACE2 cell data. Assays were performed in quadruplicate. NS, no statistical significance.

Extended Data Fig. 7. Flow cytometry of the S-expressing cells.

(Left) Representative histogram of the S protein expression on the cell surface. The number in the histogram indicates the mean fluorescence intensity (MFI). (Right) The MFI of surface S on the S-expressing cells. A statistically significant difference (*, $P < 0.05$) versus the D614G S was determined by Student's *t* test.

Extended Data Fig. 8. Association of the P681R mutation on the sensitivity to NAbs.

Neutralisation assay was performed by using three RBD-targeting monoclonal antibodies (clones 8A5, 4A3 and CB6) (**a** and **c**) and 19 vaccinated sera (**b**). NAbs were used for the pseudovirus assay (**a** and **b**) and the S-based fusion assay (**c**). Pseudoviruses and effector cells (S-expressing cells) were treated with serially diluted NAbs or sera as described in **Methods**. The raw data of panel **b** is shown in **Extended Data Fig. 9**. NT_{50}, 50% neutralisation titre.

In **a**, the NT_{50} values of the D614G S (black) and D614G/P681R S (orange) are indicated.

In **b**, a statistically significant difference versus the D614G virus was determined by Wilcoxon matched-pairs signed rank test.

Extended Data Fig. 9. Neutralisation assay using 19 vaccinated sera.

Nineteen vaccinated sera were used for the neutralisation assay. The NT_{50} values of respective serum against the D614G S (black) and D614G/P681R S (orange) are indicated in each panel. The NT_{50} values are summarized in **Extended Data Fig. 8b**.

739   **Extended Data Table 1.** Number of daily deposited sequences in GISAID.

741   **Extended Data Table 2.** Percentage of the mutations detected in the S protein of

742   the B.1.617 lineage.

744   **Extended Data Table 3.** The SARS-CoV-2 genomic region encoded by each

745   template and the primers used for the preparation of each fragment for CPER.

Methods

Ethics Statement

For virus isolation, this study was approved by the Institutional Review Board of Tokyo Metropolitan Institute of Public Health, according to the Declaration of Helsinki 2013 (approval number 3KenKenKen-466). For the use of human specimen, all protocols involving human subjects recruited at Kyoto University were reviewed and approved by the Institutional Review Boards of Kyoto University (approval number G0697). All human subjects provided written informed consent. All experiments with hamsters were performed in accordance with the Science Council of Japan's Guidelines for Proper Conduct of Animal Experiments. The protocols were approved by the Institutional Animal Care and Use Committee of National University Corporation Hokkaido University (approval number 20-0123) and the Animal Experiment Committee of the Institute of Medical Science, the University of Tokyo (approval number PA19-75).

Collection of BNT162b2-Vaccinated Sera

Peripheral blood were collected four weeks after the second vaccination of BNT162b2 (Pfizer-BioNTech), and the sera of 19 vaccinees (average age: 38, range: 28-59, 26% male) were isolated from peripheral blood. Sera were inactivated at 56°C for 30 min and stored at –80°C until use.

Cell Culture

HEK293 cells (a human embryonic kidney cell line; ATCC CRL-1573), HEK293T cells (a human embryonic kidney cell line; ATCC CRL-3216), and HOS cells (a human osteosarcoma cell line; ATCC CRL-1543) were maintained in Dulbecco's modified Eagle's medium (high glucose) (Wako, Cat# 044-29765) containing 10% fetal bovine serum (FBS) and 1% PS. Vero cells [an African green monkey (*Chlorocebus sabaeus*) kidney cell line; JCRB0111] were maintained in Eagle's minimum essential medium (Wako, Cat# 051-07615) containing 10% FBS and 1% PS. VeroE6/TMPRSS2 cells [an African green monkey (*Chlorocebus sabaeus*) kidney cell line; JCRB1819][37] were maintained in Dulbecco's modified Eagle's medium (low glucose) (Wako, Cat# 041-29775) containing 10% FBS, G418 (1 mg/ml; Nacalai Tesque, Cat# G8168-10ML) and 1% PS. Calu-3 cells (a human lung epithelial cell line; ATCC HTB-55) were maintained in Minimum essential medium Eagle (Sigma-Aldrich, cat# M4655-500ML) containing 10% FCS and 1% PS. HOS-ACE2/TMPRSS2 cells, the HOS cells stably expressing human ACE2, was prepared as described previously[8,38]. HEK293-C34 cells, the *IFNAR1* KO HEK293 cells

784 expressing human ACE2 and TMPRSS2 by doxycycline treatment[39], were
785 maintained in Dulbecco's modified Eagle's medium (high glucose) (Sigma-Aldrich,
786 Cat# R8758-500ML) containing 10% FBS, 10 μg/ml blasticidin (InvivoGen, Cat# ant-
787 bl-1) and 1% PS.
788

Animal experiments
Syrian hamsters (Male, 4 weeks old) were purchased from Japan SLC Inc.
(Shizuoka, Japan). Baseline body weights were measured before infection. For the
virus infection in **Fig. 2c, d**, hamsters were euthanised by intramuscular injection of
a mixture of 0.15 mg/kg medetomidine hydrochloride (Domitor®, Nippon Zenyaku
Kogyo), 2.0 mg/kg midazolam (Dormicum®, Maruishi Pharmaceutical) and 2.5 mg/kg
butorphanol (Vetorphale®, Meiji Seika Pharma). The B.1.1 or B.1.167.2/Delta viruses
(10^5 $TCID_{50}$ in 100 μl) was intranasally infected under anesthesia. Body weights were
measured and oral swabs were collected under anesthesia with isoflurane
(Sumitomo Dainippon Pharma) daily. For the virus infection in **Fig. 4**, four hamsters
per group were intranasally inoculated with the D614G or the D614G/P681R viruses
(10^4 $TCID_{50}$ in 30 μl) under isoflurane anesthesia. Body weight was monitored daily
for 7 days. For virological examinations, four hamsters per group were intranasally
infected with the D614G or the D614G/P681R viruses (10^4 $TCID_{50}$ in 30 μl); at 3 and
7 dpi, the animals were euthanized and nasal turbinates and lungs were collected.
The virus titers in the nasal turbinates and lungs were determined by use of plaque
assays on VeroE6/TMPRSS2 cells.
806

Lung function
Respiratory parameters were measured by using a whole-body plethysmography
system (PrimeBioscience) according to the manufacturer's instructions. In brief,
hamsters were placed in the unrestrained plethysmography chambers and allowed
to acclimatize for 1 m before data were acquired over a 3-min period by using
FinePointe software.
813

Viral Genomes
All SARS-CoV-2 genome sequences and annotation information used in this study
were downloaded from GISAID (https://www.gisaid.org) as of May 31, 2021
(1,761,037 sequences). We first excluded the genomes with non-human hosts. We
obtained SARS-CoV-2 variants belonging to the B.1.617 lineage based on the
PANGO annotation (i.e. sublineages B.1.617.1, B.1.617.2/Delta, or B.1.617.3) for
each sequence in the GISAID metadata. Note that only one variant belonging to the
B.1.617 lineage (GISAID ID: EPI_ISL_1544002 isolated in India on February 25,

2021) was not used in the analysis because the variant is not assigned any three
sublineages possibly due to 212 undetermined nucleotides in the genome. To infer
epidemiology of the B.1.617 lineage (**Fig. 1b-1f**), we excluded genomes that
sampling date information are not available, and collected 2,855, 13,821, or 83
sequences belonging to the B.1.617.1, B.1.617.2/Delta, or B.1.617.3 sublineage,
respectively.
828
Phylogenetic Analyses
To infer the phylogeny of the B.1.617 sublineages, we screened SARS-CoV-2
genomes by removing genomes containing undetermined nucleotides at coding
regions. Since the number of genomes belonging to the sublineage B.1.617.1 or
B.1.617.2/Delta are large (i.e. 894 or 6152 sequences, respectively), we used 150
sequences randomly chosen for each sublineage. For the B.1.617.3 sublineage, 32
genomes were used. We used Wuhan-Hu-1 strain isolated in China on December
31, 2019 (GenBank ID: NC_045512.2 and GISAID ID: EPI_ISL_402125) and LOM-
ASST-CDG1 strain isolated Italy on February 20, 2020 (GISAID ID:
EPI_ISL_412973) as an outgroup. We then collected 334 representative SARS-
CoV-2 sequences, and aligned entire genome sequences by using the FFT-NS-1
program in MAFFT suite v7.407[40]. All sites with gaps in alignment are removed, and
the total length of alignment is 29,085 nucleotides. Maximum likelihood tree was
generated by IQ-TREE 2 v2.1.3 software with 1,000 bootstraps[41]. GTR+G
substitution model is utilized based on BIC criterion.
844
SARS-CoV-2 Preparation and Titration
A B.1.617.2/Delta isolate (GISAID ID: EPI_ISL_2378732) and a D614G-bearing
B.1.1 isolate (GISAID ID: EPI_ISL_479681) were isolated from SARS-CoV-2-
positive individuals in Japan. Briefly, 100 µl of the nasopharyngeal swab obtained
from SARS-CoV-2-positive individuals were inoculated into VeroE6/TMPRSS2 cells
in the biosafety level 3 laboratory. After the incubation at 37°C for 15 minutes, a
maintenance medium supplemented with Eagle's minimum essential medium
(FUJIFILM Wako Pure Chemical Corporation, Cat# 056-08385) including 2% FBS
and 1% PS was added, and the cells were cultured at 37°C under 5% CO_2. The
cytopathic effect (CPE) was confirmed under an inverted microscope (Nikon), and
the viral load of the culture supernatant in which CPE was observed was confirmed
by real-time RT-PCR. To determine viral genome sequences, RNA was extracted
from the culture supernatant using QIAamp viral RNA mini kit (Qiagen, Qiagen, Cat#
52906) . cDNA library was prepared by using NEB Next Ultra RNA Library Prep Kit

for Illumina (New England Biolab, Cat# E7530) and whole genome sequencing was
performed by Miseq (Illumina).

To prepare the working virus, 100 µl of the seed virus was inoculated into
VeroE6/TMPRSS2 cells (5,000,000 cells in a T-75 flask). At one hour after infection,
the culture medium was replaced with Dulbecco's modified Eagle's medium (low
glucose) (Wako, Cat# 041-29775) containing 2% FBS and 1% PS; at 2-3 days
postinfection, the culture medium was harvested and centrifuged, and the
supernatants were collected as the working virus.

The titre of the prepared working virus was measured as 50% tissue culture
infectious dose ($TCID_{50}$). Briefly, one day prior to infection, VeroE6/TMPRSS2 cells
(10,000 cells/well) were seeded into a 96-well plate. Serially diluted virus stocks were
inoculated to the cells and incubated at 37°C for 3 days. The cells were observed
under microscopy to judge the CPE appearance. The value of $TCID_{50}$/ml was
calculated with the Reed–Muench method[42].

SARS-CoV-2 Infection
One day prior to infection, Vero cells (10,000 cells), VeroE6/TMPRSS2 cells (10,000
cells) and Calu-3 cells (10,000 cells) were seeded into a 96-well plate. SARS-CoV-
2 was inoculated and incubated at 37°C for 1 h. The infected cells were washed, and
180 µl of culture medium was added. The culture supernatant (10 µl) was harvested
at indicated time points and used for real-time RT-PCR to quantify the viral RNA
copy number. To monitor the syncytia formed in infected cell culture, the bright-field
photos were obtained using ECLIPSE Ts2 (Nikon). The size of floating syncytia was
measured by "quick selection tool" in Photoshop CS5 (Adobe) as pixel, and the area
of floating syncytia was calculated from the pixel value. As for the GFP-expressing
recombinant viruses (**Extended Data Fig. 4b, c**), the bright-field and green
fluorescence photos were obtained using an All-in-One Fluorescence microscope
BZ-X800 (Keyence) at the indicated time points, and the GFP-fluorescence intensity
was analyzed by a BZ-X800 Analyzer (Keyence).

Immunofluorescence Staining
One day prior to infection, VeroE6/TMPRSS2 cells (200,000 cells) were seeded on
the coverslips put in 12-well plate and were infected with SARS-CoV-2 (2,000
$TCID_{50}$). At 48 hours postinfection, the cells were fixed with 4% paraformaldehyde
in phosphate buffer saline (PBS) (Nacalai Tesque, Cat# 09154-85) for 10 min at
room temperature. The fixed cells were permeabilized with 0.1% Triton X-100 in PBS
for 10 min, blocked with 10% FBS in PBS for overnight at 4°C, and then stained
using mouse anti-SARS-CoV-2 N monoclonal antibody (GeneTex, Cat#

GTX632269) for 1 h. After washing three times with PBS, cells were incubated with
an Alexa 488-conjugated anti-mouse IgG antibody (Jackson ImmunoResearch, Cat#
015-540-003) for 1 h. Nuclei were stained with Hoechst 33342 (Thermo Fisher
Scientific, Cat# H3570). The coverslips were mounted on glass slides using
Fluoromount-G (Southern Biotechnology, Cat# 0100-01) with Hoechst 33342.
Fluorescence microscopy was performed on a confocal laser microscope (A1RSi,
Nikon) and captured with NIS-Elements AR software (Nikon). The area of N-positive
cells was quantified using Fiji software implemented in Image J.
905

SARS-CoV-2 Reverse Genetics
Recombinant SARS-CoV-2 was generated by circular polymerase extension
reaction (CPER) as previously described[39,43]. In brief, 9 DNA fragments encoding
the partial genome of SARS-CoV-2 (strain WK-521, PANGO lineage A; GISAID ID:
EPI_ISL_408667)[37] were prepared by PCR using PrimeSTAR GXL DNA polymerase
(Takara, cat# R050A). A linker fragment encoding hepatitis delta virus ribozyme,
bovine growth hormone polyA signal and cytomegalovirus promoter was also
prepared by PCR. The corresponding SARS-CoV-2 genomic region and the
templates and primers of this PCR are summarized in **Extended Data Table 3**. The
10 obtained DNA fragments were mixed and used for CPER[39]. To prepare the GFP-
expressing replication-competent recombinant SARS-CoV-2, the fragment 9 in
which the *GFP* gene was inserted in the *ORF7a* frame was used instead of the
authentic F9 fragment (see **Extended Data Table 3**)[39].
To produce recombinant SARS-CoV-2, the CPER products were
transfected into HEK293-C34 cells using TransIT-LT1 (Takara, cat# MIR2300)
according to the manufacturer's protocol. At one day posttransfection, the culture
medium was replaced with Dulbecco's modified Eagle's medium (high glucose)
(Sigma-Aldrich, cat# R8758-500ML) containing 2% FCS, 1% PS and doxycycline (1
µg/ml; Takara, cat# 1311N). At six days posttransfection, the culture medium was
harvested and centrifuged, and the supernatants were collected as the seed virus.
To remove the CPER products (i.e., SARS-CoV-2-related DNA), 1 ml of the seed
virus was treated with 2 µl TURBO DNase (Thermo Fisher Scientific, cat# AM2238)
and incubated at 37°C for 1 h. Complete removal of the CPER products (i.e., SARS-
CoV-2-related DNA) from the seed virus was verified by PCR. The working virus was
prepared by using the seed virus as described above.
To generate recombinant SARS-CoV-2 mutants, mutations were inserted
in fragment 8 (**Extended Data Table 3**) using the GENEART site-directed
mutagenesis system (Thermo Fisher Scientific, cat# A13312) according to the
manufacturer's protocol with the following primers: Fragment 8_S D614G forward,

5'-CCA GGT TGC TGT TCT TTA TCA GGG TGT TAA CTG CAC AGA AGT CCC
TG-3'; Fragment 8_S D614G reverse, 5'- CAG GGA CTT CTG TGC AGT TAA CAC
CCT GAT AAA GAA CAG CAA CCT GG -3'; Fragment 8_S P681R forward, 5'-AGA
CTC AGA CTA ATT CTC GTC GGC GGG CAC GTA GTG TA-3'; and Fragment 8_S
P681R reverse, 5'-TAC ACT ACG TGC CCG CCG ACG AGA ATT AGT CTG AGT
CT-3', according to the manufacturer's protocol. Nucleotide sequences were
determined by a DNA sequencing service (Fasmac), and the sequence data were
analyzed by Sequencher version 5.1 software (Gene Codes Corporation). The
CPER for the preparation of SARS-CoV-2 mutants was performed using mutated
fragment 8 instead of parental fragment 8. Subsequent experimental procedures
correspond to the procedure for parental SARS-CoV-2 preparation (described
above). To verify insertion of the mutation in the working viruses, viral RNA was
extracted using a QIAamp viral RNA mini kit (Qiagen, cat# 52906) and reverse
transcribed using SuperScript III reverse transcriptase (Thermo Fisher Scientific,
cat# 18080085) according to the manufacturers' protocols. DNA fragments including
the mutations inserted were obtained by RT-PCR using PrimeSTAR GXL DNA
polymerase (Takara, cat# R050A) and the following primers: WK-521 23339-23364
forward, 5'-GGT GGT GTC AGT GTT ATA ACA CCA GG-3'; and WK-521 24089-
24114 reverse, 5'-CAA ATG AGG TCT CTA GCA GCA ATA TC-3'. Nucleotide
sequences were determined as described above, and sequence chromatograms
(**Extended Data Fig. 3**) were visualized using the web application Tracy
(https://www.gear-genomics.com/teal/)[44].

Real-Time RT-PCR
Real-time RT-PCR was performed as previously described[43,45]. Briefly, 5 µl of culture
supernatant was mixed with 5 µl of 2 × RNA lysis buffer [2% Triton X-100, 50 mM
KCl, 100 mM Tris-HCl (pH 7.4), 40% glycerol, 0.8 U/µl recombinant RNase inhibitor
(Takara, cat# 2313B)] and incubated at room temperature for 10 min. RNase-free
water (90 µl) was added, and the diluted sample (2.5 µl) was used as the template
for real-time RT-PCR performed according to the manufacturer's protocol using the
One Step TB Green PrimeScript PLUS RT-PCR kit (Takara, cat# RR096A) and the
following primers: Forward *N*, 5'-AGC CTC TTC TCG TTC CTC ATC AC-3'; and
Reverse *N*, 5'-CCG CCA TTG CCA GCC ATT C-3'. The copy number of viral RNA
was standardized with a SARS-CoV-2 direct detection RT-qPCR kit (Takara, cat#
RC300A). The fluorescent signal was acquired using a QuantStudio 3 Real-Time
PCR system (Thermo Fisher Scientific), a CFX Connect Real-Time PCR Detection
system (Bio-Rad), an Eco Real-Time PCR System (Illumina) or a 7500 Real Time
PCR System (Applied Biosystems).

Plasmid Construction

Plasmid Construction

A plasmid expressing the SARS-CoV-2 S D614G protein was prepared in our
previous study[8]. A plasmid expressing the SARS-CoV-2 S D614G/P681R mutant
was generated by site-directed mutagenesis PCR using pC-SARS2-S D614G[8] as
the template and the following primers: P681R Fw, 5'- CCA GAC CAA CAG CCG
GAG GAG GGC AAG GTC T-3' and P681R Rv, 5'-AGA CCT TGC CCT CCT CCG
GCT GTT GGT CTG G-3'. The resulting PCR fragment was digested with KpnI and
NotI and inserted into the KpnI-NotI site of the pCAGGS vector[46].

Pseudovirus Assay

Pseudovirus assay was performed as previously described[8,43]. Briefly, the
pseudoviruses, lentivirus (HIV-1)-based, luciferase-expressing reporter viruses
pseudotyped with the SARS-CoV-2 S protein and its derivatives, HEK293T cells (1
$\times 10^6$ cells) were cotransfected with 1 μg of psPAX2-IN/HiBiT[47], 1 μg of pWPI-Luc2[47],
and 500 ng of plasmids expressing parental S or its derivatives using Lipofectamine
3000 (Thermo Fisher Scientific, Cat# L3000015) or PEI Max (Polysciences, Cat#
24765-1) according to the manufacturer's protocol. At two days posttransfection, the
culture supernatants were harvested, centrifuged. The amount of pseudoviruses
prepared was quantified using the HiBiT assay as previously described[8,47]. The
pseudoviruses prepared were stored at –80°C until use. For the experiment, HOS-
ACE2 cells and HOS-ACE2/TMPRSS2 cells (10,000 cells/50 μl) were seeded in 96-
well plates and infected with 100 μl of the pseudoviruses prepared at 4 different
doses. At two days postinfection, the infected cells were lysed with a One-Glo
luciferase assay system (Promega, Cat# E6130), and the luminescent signal was
measured using a CentroXS3 plate reader (Berthhold Technologies) or GloMax
explorer multimode microplate reader 3500 (Promega).

1000

Western blotting

Western blotting was performed as previously described[48-50]. To quantify the level of
the cleaved S2 protein in the cells, the harvested cells were washed and lysed in
lysis buffer [25 mM HEPES (pH 7.2), 20% glycerol, 125 mM NaCl, 1% Nonidet P40
substitute (Nalacai Tesque, Cat# 18558-54), protease inhibitor cocktail (Nalacai
Tesque, Cat# 03969-21)]. After quantification of total protein by protein assay dye
(Bio-Rad, Cat# 5000006), lysates were diluted with 2 × sample buffer [100 mM Tris-
HCl (pH 6.8), 4% SDS, 12% β-mercaptoethanol, 20% glycerol, 0.05% bromophenol
blue] and boiled for 10 min. Ten microliter of the samples (50 μg of total protein)
were subjected to western blotting. To quantify the level of the cleaved S2 protein

on virions, 900 µl of the culture medium including the pseudoviruses were layered
onto 500 µl of 20% sucrose in PBS and centrifuged at 20,000 × g for 2 h at 4°C.
Pelleted virions were resuspended in 1× NuPAGE LDS sample buffer (Thermo
Fisher Scientific, Cat# NP0007) containing 2% β-mercaptoethanol, and the lysed
virions were subjected to western blotting. For the protein detection, following
antibodies were used: mouse anti-SARS-CoV-2 S monoclonal antibody (clone 1A9,
GeneTex, Cat# GTX632604), rabbit anti-ACTB monoclonal antibody (clone 13E5,
Cell Signaling, Cat# 4970), mouse anti-HIV-1 p24 monoclonal antibody (clone 183-
H12-5C, obtained from the HIV Reagent Program, NIH, Cat# ARP-3537),
horseradish peroxidase (HRP)-conjugated donkey anti-rabbit IgG polyclonal
antibody (Jackson ImmunoResearch, Cat# 711-035-152), and HRP-conjugated
donkey anti-mouse IgG polyclonal antibody (Jackson ImmunoResearch, Cat# 715-
035-150). Chemiluminescence was detected using SuperSignal West Femto
Maximum Sensitivity Substrate (Thermo Fisher Scientific, Cat# 34095) or Western
BLoT Ultra Sensitive HRP Substrate (Takara, Cat# T7104A) according to the
manufacturers' instruction. Bands were visualized using the image analyzer,
Amersham Imager 600 (GE Healthcare), and the band intensity was quantified using
Image Studio Lite (LI-COR Biosciences) or Image J.
1029
SARS-CoV-2 S-Based Fusion Assay
The SARS-CoV-2 S-based fusion assay was performed as previously described[43].
This assay utilizes a dual split protein (DSP) encoding *Renilla* luciferase (RL) and
GFP genes, and the respective split proteins, DSP_{1-7} and DSP_{8-11}, are expressed in
effector and target cells by transfection[49,51]. Briefly, on day 1, effector cells (i.e., S-
expressing cells) and target cells (i.e., ACE2-expressing cells) were prepared at a
density of 0.6 to 0.8 × 10^6 cells in a 6 well plate. To prepare effector cells, HEK293
cells were cotransfected with the expression plasmids for D614G S or D614G/P681R
(400 ng) with $pDSP_{1-7}$ (400 ng) using TransIT-LT1 (Takara, Cat# MIR2300). To
prepare target cells, HEK293 cells were cotransfected with pC-ACE2 (200 ng) and
$pDSP_{8-11}$ (400 ng). In addition to the plasmids above, selected wells of target cells
were also cotransfected with pC-TMPRSS2 (40 ng). On day 3 (24 h posttransfection),
16,000 effector cells were detached and reseeded into 96-well black plates
(PerkinElmer, Cat# 6005225), and target cells were reseeded at a density of
1,000,000 cells/2 ml/well in 6-well plates. On day 4 (48 h posttransfection), target
cells were incubated with EnduRen live cell substrate (Promega, Cat# E6481) for 3
h and then detached, and 32,000 target cells were applied to a 96-well plate with
effector cells. RL activity was measured at the indicated time points using a Centro
XS3 LB960 (Berthhold Technologies). The S proteins expressed on the surface of

effector cells were stained with rabbit anti-SARS-CoV-2 S monoclonal antibody
(GeneTex, Cat# GTX635654) and APC-conjugated goat anti-rabbit IgG polyclonal
antibody (Jackson ImmunoResearch, Cat# 111-136-144). Normal rabbit IgG
(SouthernBiotech, Cat# 0111-01) was used as a negative control. Expression levels
of surface S proteins were analyzed using a FACS Canto II (BD Biosciences). RL
activity was normalized to the mean fluorescence intensity (MFI) of surface S
proteins, and the normalized values are shown as fusion activity.
1056

Mathematical Modeling for Fusion Velocity Quantification
The following cubic polynomial regression model was employed to fit each of time-
series datasets (**Fig. 3e**):
$$y \sim b_0 + b_1 x + b_2 x^2 + b_3 x^3$$
The initial velocity of cell fusion was estimated from the derivative of the fitted cubic
curve.
1063

Neutralisation Assay
Virus neutralisation assay was performed on HOS-ACE2/TMPRSS2 cells using the
SARS-CoV-2 S pseudoviruses expressing luciferase (see "Pseudovirus Assay"
above). The viral particles pseudotyped with D614G S or D614G/P681R S were
incubated with serial dilution of heat-inactivated human serum samples or the RBD-
targeting NAbs (clones 8A5, 4A3 and CB6; Elabscience) at 37°C for 1 h. The
pseudoviruses without sera and NAbs were also included. Then, the 80 μl mixture
of pseudovirus and sera/NAbs was added into HOS-ACE2/TMPRSS2 cells (10,000
cells/50 μl) in a 96-well white plate and the luminescence was measured as
described above (see "Pseudovirus Assay" above). 50% neutralisation titre (NT$_{50}$)
was calculated using Prism 9 (GraphPad Software).
For the cell-cell fusion neutralisation assay, effector cells of the S-based
fusion assay (i.e., S-expressing cells) were incubated with the serially diluted
neutralizing antibodies targeting RBD (clones 8A5, 4A3 and CB6; Elabscience) at
37°C for 1 h. Then, target cells were applied and performed the S-based fusion
assay as described above (see "SARS-CoV-2 S-Based Fusion Assay" above).