Psychophysiological Evidence for Fear Extinction Learning via Mental Imagery

Xinrui Jiang[1], & Steven Greening[1,2,3]

Louisiana State University

Affiliations: [1]Louisiana State University, United States; [2]Pennington Biomedical Research

Center, United States; [3]University of Manitoba, Canada

Submitted to:

Corresponding author: Steven Greening

                          Department of Psychology

                          University of Manitoba

                          Email: steven.greening@umanitoba.ca

Abstract

Imagery-based extinction procedures have long been used in the treatments of fear-related conditions. The assumption here is that imagery can substitute for the perceptual stimuli in the extinction process. Yet, experimental validations of this assumption have been limited in number and some have relied exclusively on measures of autonomic reactivity without consideration of conscious feelings of fear. The current investigation sought to assess whether imagery-based exposure could lead to extinction of conditioned fear to the corresponding perceptual stimulus. Conditioned fear responses were measured by both a physiological (i.e., skin conductance response/SCR) and a subjective (i.e., self-reported fear) measure. Participants ($N = 56$) first underwent perceptual differential fear conditioning, then imagery extinction, then to perceptual extinction. SCR evidence was found for successful fear conditioning, generalization of fear from viewing to imagery, and most importantly, the absence of differential fear after imagery extinction upon re-exposure to the conditioned perceptual stimulus. Self-reported fear confirmed the acquisition and generalization of fear and provided evidence of a significant reduction in differential fear conditioning across extinction. Consistent with clinical evidence of the efficacy of imagery extinction and the existing limited experimental literature, the current study offers support for fear extinction to perceptual stimuli via imagery exposure.

Keywords: differential fear conditioning, extinction, mental imagery, SCR, self-reported fear

Psychophysiological Evidence for Fear Extinction Learning via Mental Imagery

1. Introduction

From as early as the practices of shamanism some 20,000 years ago (Achterberg, 2002) to today's imagery-based cognitive behavioral therapy techniques (Holmes, Arntz, & Smucker, 2007), mental imagery has been harnessed for its efficacy as a cognitive process that affects emotion perception and regulation. It is commonly referred to as a perceptual-like experience in the absence of external sensory input (Kosslyn, Thompson, & Ganis, 2006). Imagery is recognized as an important component of various fear-related affective conditions in terms of both their symptoms (e.g., intrusive images in depression, flashbacks in post-traumatic stress disorder or PTSD; Matthews, Collins, Thakkar, & Park, 2014; Weßlau & Steil, 2014), and clinical treatments (e.g., imaginal exposure, Abramowitz, 1996; Hecker, 1990; Powers, Halpern, Ferenschak, Gillihan, & Foa, 2010; Turner, Beidel, & Jacob, 1994; imagery rescripting, Arntz, 2012). The primary account for the imagery's impact in these examples is that it can substitute for the actual stimuli in affective processes to regulate fear responses. Yet, only a few experimental examinations have been done to assess this explanation (Mertens, Krypotos, & Engelhard, 2020).

While some have suggested that informational processing attributed to mental imagery is purely propositional, an abundance of behavioral and brain imaging research suggests that mental imagery is at least partially depictive or pictorial (Kosslyn, 1981, 2005; Kosslyn, Ganis, & Thompson, 2001). According the depictive theory, mental imagery generates a neural representation that functions like a weaker version of perception and is associated with the subjective experience of, for example, "seeing with the mind's eye" (Kosslyn et al., 2001). Thus, imagery of an object can "stand in" for the corresponding external stimulus in various cognitive

processes including associative learning and other affective processes (Lewis, O'Reilly, Khuu, & Pearson, 2013; Pearson & Kosslyn, 2015; Pearson, Naselaris, Holmes, & Kosslyn, 2015). On the other hand, the bio-informational theory noted that mental imagery contains not only the perceptual information of the imagined stimulus but also its conceptual information stored as propositions such as "relationships, descriptions, interpretations, labels, and tags" (Lang, 1977, 1979). The bio-informational theory posits that it is through the activation of this propositional network during imagery that emotional modifications occur (Lang, 1977, 1979). Regardless of the precise nature of mental imagery, it is acknowledged by both theoretical perspectives that mental imagery has the capacity to elicit emotional characteristics similar to its perceptual counterparts.

In addition to successful clinical applications and theoretical foundations, neuroimaging research has also provided evidence that mental imagery contributes to the elicitation or modification of emotional reactivity similarly to perception. For example, meta-analytic reviews of the brain regions associated with the emotional processing of perceptual stimuli report activations in regions such as medial prefrontal cortex (mPFC), anterior cingulate cortex (ACC), insula (INS), and amygdala (Kober et al., 2008; Murphy, Nimmo-Smith, & Lawrence, 2003; Phan, Wager, Taylor, & Liberzon, 2002). Importantly, activations of the same regions have been reported during elicitation of emotions via mental imagery, including mPFC (Schaefer et al., 2003), ACC, INS (Phan et al., 2002), and amygdala (Kreiman, Koch, & Fried, 2000). Taken together, the existing research appears to support the intuition that mental imagery of an emotional stimulus can substitute for its external presentation. In comparison to clinical evidence, theoretical discussions and neuroimaging findings, relatively less experimental research has investigated the emotion regulatory capacity of mental imagery in comparison with perception

despite it being a direct way to investigate the assumed correspondence between mental imagery and perception.

Among the existing studies that investigated the connection between mental imagery and emotion, many based their task paradigms on differential fear conditioning (Agren, Björkstrand, & Fredrikson, 2017; Greening et al., 2021; Grégoire & Greening, 2019; Koizumi et al., 2016; Reddan, Wager, & Schiller, 2018). Such paradigm typically involves one initially neutral conditioned stimulus (CS+) which is repeatedly paired with an aversive or threatening unconditioned stimulus (US; e.g., a loud noise or mild electric shock), and a second CS (CS-) which is never accompanied by the US (Carter, Hofstotter, Tsuchiya, & Koch, 2003). Upon such pairing, the CS+ but not the CS- exhibits a conditioned response (CR) that is similar to the type of response elicited by US (e.g., increased self-reported fear; increased skin conductance response, SCR; increased heart rate; or, escape behavior; Carter et al., 2003; Critchley, Mathias, & Dolan, 2002). The acquisition of differential fear conditioning is quantified as the difference in the CR between the CS+ versus the CS-. Greening et al. (2021) observed recently that visual imagery of conditioned stimuli elicited a significant differential fear conditioned response as measured by self-reported fear, SCR, and activation of the right INS. Following the acquisition of differential fear, fear extinction learning is commonly used for inhibiting the fear conditioning association (Milad & Quirk, 2012). It involves repeated presentations of the CS+ in the absence of the US such that a novel safety associated memory is formed, which leads to a reduction or suppression of the differential CR. Many imagery-based clinical interventions such as imaginal exposure (Abramowitz, 1996; Hecker, 1990; Powers et al., 2010; Turner et al., 1994) are premised on such extinction theorizing.

Though limited in number, existing research has provided supporting evidence for the efficacy of imagery exposure in the down-regulation of differential fear conditioning via fear extinction (e.g., Agren et al., 2017; Reddan et al., 2018). Agren et al. (2017) found that exposure to either an imagined or perceptual CS+ during extinction led to comparable degrees of reduction in differential SCR. Reddan et al. (2018) reported reductions in both threat-relevant neural predictive pattern expression and SCR upon re-exposure to conditioned auditory CS+ following imagined extinction during which participants were asked to play the conditioned CS+ tones in their head. Koizumi et al. (2016) also managed to reduce conditioned fear responses as measured by SCR and neural activations using counter conditioning by pairing implicit imagery of the CS+ with monetary reward using neurofeedback. Notably, the above cited studies all relied on physiological indices of differential conditioning without assessments of the subjective experiences of fear. Although there are important methodological reasons for avoiding the simultaneous acquisition of some measures of differential conditioning that may confound each other (e.g., SCR and fear potentiated startle; Sjouwerman, Niehaus, Kuhn, & Lonsdorf, 2016), from the perspective of the multi-system conceptualization of emotion it is important to collected multiple measures when possible. Specifically, emotion is expressed through both the conscious feeling of fear and also the corresponding behavioral and physiological expressions (LeDoux & Pine, 2016), and that over-reliance on any single response system will be limiting to our understanding of the emotion being studied (Frijda, 1986; Gross, 2013; Jiang, Burleigh, & Greening, 2021; Lang, 1993; Larsen & Prizmic-Larsen, 2006; LeDoux & Pine, 2016). For instance, Mauss and colleagues (2005) reported only moderate correlations between self-reported descriptions of the emotional experience and physiological responses. In addition, neuroimaging

evidence suggests that subjective and physiological expressions of fear may be reflective of different brain and cognitive mechanisms (Taschereau-Dumouchel, Kawato, & Lau, 2019).

The current study aimed to evaluate whether visual mental imagery could produce fear extinction following the acquisition of differential fear conditioning to visual objects, in terms of both self-reported fear and SCR. To do so, we used a within-subject design in which two CS+s and one CS- underwent acquisition of differential conditioning (Acquisition), followed by in vivo, perceptual extinction of one of the CS+s (CS+V) and imagery extinction of the other (CS+I) during Extinction Phase 1. Both of the CS+s were then viewed again in Extinction Phase 2. Considering the ample evidence supporting the effectiveness of imagery-based clinical interventions on the reduction of negative affect (Heyes, Lau, & Holmes, 2013; Holmes & Mathews, 2010; Weßlau & Steil, 2014) and the above research findings on imagery in fear down-regulation (Agren et al., 2017; Koizumi et al., 2016; Reddan et al., 2018), our overall prediction was that conditioned fear responses (as measured by both self-report and SCR) to perceived visual stimuli would reduce after repeated imaginary exposure to the same stimuli without the accompany of US. The accuracy of this overall prediction was evaluated on the basis of two specific experimental predictions. First, we hypothesized that imagining the CS+I during Extinction Phase 1 would elicit a significant CR compared to the CS-, as the fear association would generalize from viewing the CS+I during Acquisition to imagining the CS+I during Extinction Phase 1 (Greening et al., 2021). Second, if fear extinction learning via imagery exposure generalized back to viewing the CS+I, then we would observe significant and similar reductions in CRs associated with the CS+V and CS+I during early Extinction Phase 2 consistent with previous research (Agren et al., 2017; Reddan et al., 2018). Conversely, if imagery exposure

did not generalize to the visual CS+I, we would predict a persistent differential CR for the CS+I

in early Extinction Phase 2.

2. Method

2.1 Participants

A total of 61 Louisiana State University students taking undergraduate-level psychology

courses were recruited and completed this study. Five participants were removed from the final

analyses due to excessive noise in SCR throughout or a lack of SCR response to the US, i.e., the

shock, leaving a final sample of 56 individuals (32 females, mean age of 19.28, $SD = 1.18$;

demographic information is missing from 3 participants). This study was approved by the

Institutional Review Board of the Louisiana State University. Written informed consent was

given to each participant prior to the start of experiment sessions. All participants were

reimbursed by way of research credits.

2.2 Stimuli

All stimuli were created and presented in MATLAB R2018a with the Psychophysics

Toolbox extensions (Brainard, 1997; Kleiner et al., 2007; Pelli, 1997). Three capital letters were

selected as the CSs: letters "J" and "H" were paired with the US functioning as two distinct CS+s

and letter "F" was used as the CS- and it was never paired with the US. All stimuli were

presented at the center of the screen in 4-by-5 grids drawn on a grey background (figure 1A). A

single mild electric shock of 5-ms duration as the US was delivered to the distal phalanx of each

participant's ring finger and pinky on his/her non-dominant hand through attached electrodes

using the Biopac MP-150 system and AcqKnowledge software (for 5 participants, the shock

electrodes were placed on their index and middle fingers). Whenever the US was delivered it co-

terminated with the CS+. The intensity of the US was customized to each participant such that it

was rated as "uncomfortable but not painful." Shock intensity was checked and adjusted after the first half of Acquisition for each participant to avoid desensitization or sensitization to the US.

[Insert figure 1]

2.3 Procedures

Each participant first completed two training runs (12 trials/run) in which participants both viewed (figure 1C) or were instructed to imagine (figure 1D) the selected three letters. During each trial a probe appears, which was a single square in the grid filled-in in red (figure 1B) and participants were instructed identify via button press whether the probe was on or off the letter viewed/imagined. Specifically, they were instructed to press 1 if they thought the probe was on the letter and press 2 if they thought the probe was off the letter. This probe task was added to the training phrase to initially facilitate participants' attention to the task and ensure the instructions were understood regarding imagery. No shock was involved during this training phase of the experiment, which allows the trials in this phase to function as habituation trials for the CSs. However, no probes were present during the remaining phases of the experiment out of a concern that they could interfere with the differential conditioning (Carter et al., 2003).

During fear conditioning/Acquisition (12 trials/run x 4 runs; each CS letter, i.e., "J", "H", and "F", was presented 16 times total), all trials were view trials (i.e., there was no mental imagery). The two CS+ letters were paired with shock on 50% of the trials (i.e., there were 8 trials per CS+ that were reinforced with the US) and the CS- was never paired with shock. Participants were told that "for 2 out of the 3 letters, there is a chance you may get shocked" before the beginning of conditioning. In a given run, the order of stimulus presentation was pseudorandomized in a way that the first and last trials of each run were always CS- trials (i.e., letter "F"). Additionally, each run contained two reinforced trials per CS+ (i.e., CS+V and CS+I,

or the letters "J" and "H"). The second and third trials of each run were always reinforced shock

trials for each of the CS+V and the CS+I (the order is randomized). The second reinforced CS+

trial for both the CS+V and the CS+I always occurred in the second half of a given acquisition

run. In addition, no more than two consecutive trials were the same. Upon completion of

Acquisition, participants proceeded into the Extinction Phase 1 (12 trials/run x 2 runs; each CS

letter, i.e., "J", "H", and "F", was presented 8 times total) in which shock was removed from

CS+ presentations. One CS+ continued to be viewed (CS+V) and the other was imagined (CS+I)

for the entire phase. The assignment of which CS+ letter (i.e., letters "J" and "H") was the CS+V

versus the CS+I was counterbalanced across participants. The CS- (i.e., letter "F") was always

viewed during Extinction Phase 1. The last phase of the experiment was Extinction Phase 2 (12

trials/run x 2 runs; each CS letter, i.e., "J", "H", and "F", was presented 8 times total), in which

all stimuli were viewed without the presence of electric shocks. The order of stimulus

presentation for both Extinction Phase 1 and Extinction Phase 2 was pseudorandomized with no

more than two of the same CS trials in a row. Details of the trial schedule for each phase are

listed in table 1.

<center>[Insert table 1]</center>

2.4 Self-Report Measures

After Acquisition, Extinction Phase 1 and 2, participants were asked to complete a series

of self-report questionnaires on their level of fear towards receiving a shock, and on their

perceived likelihood of being shocked for each CS. They were instructed to report how much

they feared receiving a shock using a 7-point Likert scale (between 1 = "Not at All" and 7 =

"Very Much So"). A 10-point Likert scale (between 0% and 100% with intervals of 10%) was

provided for reporting estimations of shock likelihood.

2.5 Skin Conductance Response

During each trial, participants' SCRs were collected and sampled at 2000 Hz. SCR analysis was carried out in MATLAB R2018a (Version 9.4). A first-order Butterworth bandpass filter was applied during preprocessing with cut-off frequencies of .01 and 5 Hz (Bach, Flandin, Friston, & Dolan, 2010). The time series were then down-sampled to 100 Hz. Based on previous research (Grégoire & Greening, 2019; D. Schiller, Kanen, LeDoux, Monfils, & Phelps, 2013; Vervliet, Vansteenwegen, Baeyens, Hermans, & Eelen, 2005) SCRs to the CSs were calculated by subtracting baseline (1 second before stimulus onset) from peak amplitude (during 1-3.995 seconds after stimulus onset). Peaks were zeroed if their distance from the baseline was smaller than .02 µS. The difference scores were then square root transformed. Consistent with previous research; Grégoire & Greening, 2019; Reddan et al., 2018; Daniela Schiller et al., 2010), we only included unreinforced CS+ trials (i.e., trials without US) in subsequent analyses. Lastly, the first and last trials of the Acquisition run were always CS- trials, which were excluded from the analyses to avoid the potential confounding orienting (Daniela Schiller et al., 2010) or "novelty" effect (Lim & Pessoa, 2008) that can occur on the first trial of a run, and to ensure that there was an equal number of CS+ and CS- trials included in the analyses (Grégoire & Greening, 2019, 2020; Lonsdorf et al., 2017).

2.6 Data Analysis

Based on recommendations by Lonsdorf et al. (2017), in order to maintain the most generalizable sample possible we only excluded participants from the analysis who displayed no clear SCR response to the US or whose overall SCR data was of poor quality even after filtering and down-sampling. No outlier removal was undertaken. Mean values of square-rooted SCR were calculated for each CS types (i.e., CS+I, CS+V, and CS−) separated into Acquisition, early

Extinction Phase 1 (first half of trials), late Extinction Phase 1 (second half of trials), early

Extinction Phase 2 (first half of trials), and late Extinction Phase 2 (late half of trials). Self-

reported fear and shock estimations were collected at the end of each study phase and thus were

not broken down into early and late responses for each of the three CSs. Repeated

measures analyses of variance (ANOVAs) were conducted with Greenhouse–Geisser correction

for non-sphericity when needed. Post hoc paired *t*-tests without corrections were conducted when

significant interactions or main effects were found (Saville, 1990). Specific analyses are

described in the results section.

3. Results

3.1 Acquisition

SCR: Acquisition was analyzed using a repeated-measures ANOVA with CS type (i.e.,

CS+I, CS+V, and CS-) as the within-subject factor on mean SCR during Acquisition (figure 2).

There was a main effect of CS type, $F(2, 110) = 3.80$, $p = .025$, $\eta^2 = .06$. Post hoc *t*-tests revealed

that while there was a significant difference between the CS+I ($M = .23$, $SD = .21$) and the CS-

($M = .17$, $SD = .14$), $t(55) = 2.59$, $p = .012$, $d = .35$; the CS+V ($M = .20$, $SD = .19$) did not differ

from the CS- ($M = .17$, $SD = .14$), $t(55) = 1.40$, $p = .167$. Moreover, the two CS+s (i.e., CS+I and

CS+V) did not differ from each other, $t(55) = 1.46$, $p = .151$. To test the robustness of CS+I and

CS- difference, a Sensitivity Analysis from a Bayesian framework (Krypotos, Klugkist, &

Engelhard, 2017) was conducted contrasting the CS+I and CS- (two-sided) using prior

distributions based on the Cauchy distribution with the following scaling factors: 0.1, 0.3, 0.5,

0.7, 1, 1.5. The resulting Bayes Factors (BF_{10}) were: $BF_{10}(0.1) = 3.357$, $BF_{10}(0.3) = 4.178$,

$BF_{10}(0.5) = 3.658$, $BF_{10}(0.7) = 3.062$, $BF_{10}(1) = 2.381$, $BF_{10}(1.5) = 1.694$, respectively,

indicating moderate to anecdotal evidence for accepting the alternative hypothesis. Together,

based on the mean SCR, CS+I but not CS+V acquired differential fear at the group-level. Thus,

we did not include the CS+V in the subsequent analyses in the extinction phases with respect to

the SCR data.

[Insert figure 2]

Self-Report: The same repeated ANOVA was applied to self-reported fear (figure 3A)

and shock estimations (figure 3B). In contrast to SCR, these two self-report measures revealed

differential conditioning of both CS+s. Specifically, for self-reported fear there was a main effect

of CS-type, $F(1.38, 76) = 32.59$, $p < .0001$, $\eta^2 = .37$. Post hoc *t*-tests indicated that participants

reported higher fear ratings for both the CS+I ($M = 4.20$, $SD = 1.63$) and the CS+V ($M = 4.23$,

$SD = 1.67$) compared to the CS- ($M = 2.14$, $SD = 1.43$), with respective statistics of $t(55) = 6.61$,

$p < .0001$, $d = .88$, and $t(55) = 7.13$, $p < .0001$, $d = .95$. The two CS+s did not differ, $t(55) = -.12$,

$p = .901$. As for shock estimation, a main effect of CS type was also found, $F(1.71, 94.22) =$

59.95, $p < .0001$, $\eta^2 = .52$. Post hoc *t*-tests showed that participants reported higher likelihood of

shock for both the CS+I ($M = 49.82$, $SD = 17.11$) and the CS+V ($M = 49.64$, $SD = 18.39$)

compared to the CS- ($M = 17.86$, $SD = 20.95$): CS+I vs CS-, $t(55) = 8.45$, $p < .0001$, $d = 1.13$;

CS+V vs CS-, $t(55) = 8.07$, $p < .0001$, $d = 1.08$). The two CS+s, again, did not differ, $t(55) = .06$,

$p = .951$.

[Insert figure 3]

3.2 Extinction Phase 1: Differential Fear Generalization to CS+ Imagery

SCR: During Extinction Phase 1, participants imagined the CS+I and continued to view

the CS-, and the US was never delivered. A 2x2 ANOVA with CS-type (i.e., CS+I, CS-) and

timing (i.e., early trials, late trials) as the within subject factors were conducted on SCR data

during Extinction Phase 1 (figure 2B). This analysis revealed significant main effects for both

CS type, $F(1, 55) = 21.12$, $p < .0001$, $\eta_p^2 = .28$, and timing, $F(1, 55) = 41.45$, $p < .0001$, $\eta_p^2 = .43$.

There was no CS type x timing interaction, $F(1, 55) = 2.91$, $p = .094$. Post hoc t-tests indicated

that the CS+I was significantly different from the CS- during both early (CS+I: $M = .38$, SD

$= .32$; CS-: $M = .19$, $SD = .20$), $t(55) = 4.03$, $p < .001$, $d = .54$; and late (CS+I: $M = .22$, $SD = .24$;

CS-: $M = .11$, $SD = .12$), $t(55) = 3.55$, $p < .001$, $d = .47$ trials of Extinction Phase 1. Both the

CS+I and the CS- had greater SCR during early compared to their respective late trials of

Extinction Phase 1 (CS+I: $t(55) = 5.53$, $p < .0001$, $d = .74$; CS-: $t(55) = 3.10$, $p = .003$, $d = .41$).

As noted previously, the CS+V was not included in SCR analyses for Extinction Phase 1 and 2.

However, a graph containing all three CSs separated into early and late Extinction Phase 1 and 2

is provided in supplementary figure S1.

 Self-Report: Results of a one-way repeated measures ANOVA with CS-type (i.e., CS+I,

CS+V, and CS-) as the within-subject factor on self-reported fear (figure 3A) also supported

generalization of fear to imagery with a main effect of CS-type, $F(1.75, 96.49) = 15.88$, p

$< .0001$, $\eta_p^2 = .22$. Based on post hoc t-tests, CS+I ($M = 3.28$, $SD = 1.96$) and CS+V ($M = 3.16$,

$SD = 1.77$) were both rated significantly higher than CS- ($M = 1.92$, $SD = 1.45$), with $t(55) =$

4.36, $p < .0001$, $d = .58$, and $t(55) = 5.10$, $p < .0001$, $d = .68$, respectively. The two CS+s did not

differ from each other, $t(55) = .55$, $p = .588$. The same one-way repeated measures ANOVA

applied to shock estimations indicated that all CS's (i.e., CS+I, CS+V, and CS-) were reported to

be paired with shock close to 0% of the time during Extinction Phase 1 (CS+I: $M = 1.84$, $SD =$

7.16; CS+V: $M = 1.61$, $SD = 6.54$; CS-: $M = .91$, $SD = 3.94$) with no effect of CS type, $F(1.47,$

$81.02) = 1.15$, $p = .308$.

3.3 Extinction Phase 2: Imagery Exposure Led to Differential Fear Extinction Learning

SCR: During Extinction Phase 2, participants now viewed both the CS+I and the CS-, and as with Extinction Phase 1 the US was never delivered. We predicted that if mental imagery of CS+I was effective then we would observe no main effect of CS type. As predicted, a 2x2 ANOVA with CS-type (i.e., CS+I, CS-) and timing (i.e., early and late extinction) as the within subject factors (figure 2C) resulted in no interaction ($F(1, 55) = .52$, $p = .476$) nor main effects (CS type: $F(1, 55) = .02$, $p = .882$; timing: $F(1, 55) = .49$, $p = .485$). This suggests that CS+I and CS- did not differ during either early (CS+I: $M = .18$, $SD = .23$; CS-: $M = .16$, $SD = .19$) or late (CS+I: $M = .15$, $SD = .15$; CS-: $M = .16$, $SD = .17$) Extinction Phase 2.

Self-report: Regarding self-reported fear, a one-way repeated measures ANOVA with CS-type (i.e., CS+I, CS+V, and CS-) as the within-subject factor (figure 3A) revealed a main effect of CS type, $F(1.75, 96.24) = 14.25$, $p < .0001$, $\eta_p^2 = .21$. Importantly, a post hoc *t*-test revealed no significant difference in self-reported fear for the CS+I ($M = 1.70$, $SD = 1.06$) versus the CS+V ($M = 1.68$, $SD = 1.05$), $t(55) = .10$, $p = .924$. Additional post hoc *t*-tests revealed higher fear ratings for both the CS+I and the CS+V than the CS- ($M = 1.21$, $SD = .68$), with $t(55) = 2.89$, $p = .006$, $d = .39$, and $t(55) = 3.00$, $p = .004$, $d = .40$, respectively. Although this result may appear to suggest that self-reported fear was not extinguished, in the following section we carried out an additional analysis across the 3 main experimental phases to demonstrate that the magnitude of self-reported differential fear was reduced via both visual and imagery extinction. Regarding self-reported shock estimation, accurate estimations were reported (figure 3B) with respect to Extinction Phase 2 with close to 0% estimations for the CS+I ($M = .89$, $SD = 5.49$), the CS+V ($M = .38$, $SD = 2.67$), and the CS- ($M = .36$, $SD = 2.67$) with no main effect of CS type, $F(1, 55) = 1.83$, $p = .182$.

3.4 Self-Reported Fear Across Acquisition, Extinction Phase 1 and Extinction Phase 2

In order to evaluate the magnitude of differential fear conditioning as measured by self-report across the three main experimental phases we conducted a 3x2 ANOVA on the difference scores of self-reported fear with study phase (i.e., Acquisition, Extinction Phase 1, Extinction Phase 2) and CS contrast (i.e., CS+I minus CS-, CS+V minus CS-) as the within-subject factors. This revealed only a main effect of study phase, $F(1.71, 94.27) = 17.84$, $p < .0001$, $\eta_p^2 = .24$. Notably, there was neither a main effect of CS contrast, $F(1, 55) = .13$, $p = 721$, nor interaction, $F(1.45, 79.87) = .23$, $p = .724$. As there was no main effect of CS contrast and no interaction, the contrast scores were averaged together (i.e., producing one aggregate measure of the CS+ vs CS- difference) and compared between study phases. Post-hoc t-tests revealed significant reductions from Acquisition ($M = 2.07$, $SD = 2.48$) to Extinction Phase 1 ($M = 1.24$, $SD = 1.81$), $t(55) = 2.83$, $p = .007$, $d = .38$; and from Extinction Phase 1 to Extinction Phase 2 ($M = .47$, $SD = .77$), $t(55) = 3.73$, $p = .000$, $d = .50$. These results suggest that the magnitude of differential fear conditioning (CS+ minus CS-) as measured with self-reported fear significantly decreased throughout extinction (i.e., Extinction Phase 1 and 2). Moreover, the degree of reduction in differential self-reported fear was similar regardless of whether in vivo or imagery exposure was employed during Extinction Phase 1.

4. Discussion

Mental imagery has long played an important part in our understanding and treatments of various affective conditions. The assumption is that imagery performs like its perceptual counterpart and can exert regulatory effects on emotion in its place. While efforts have been made in the theorizing and assessment of imagery-based emotion regulatory strategies, experimental examination is lacking in comparison. The current investigation sought to determine if visual mental imagery could contribute to fear extinction as measured by both

physiology (i.e., SCR) and self-reported fear, as this has not been done in the current literature

(Agren et al., 2017; Reddan et al., 2018). First, we found that conditioned fear acquired by visual

stimuli generalized to visual imagery as indicated by elevated SCR and self-reported fear.

Second, following repeated visual imagery of the CS+I, there was no significant differential SCR

upon re-exposure to the CS+I compared to the CS-. In addition, based on self-reported fear,

imagery-based and perceptual extinction led to similar degrees of reductions in the differential

conditioned fear response, consistent with the SCR-based findings of Agren et al. (2017).

As the main research question of the current investigation is whether imagery exposure to

the CS+ leads to generalized fear reduction to the perceptual (i.e., visual) CS+, our observation

of significantly greater SCR for CS+I versus the CS- during acquisition allows us to make valid

inferences regarding mental imagery's impact on extinction despite the lack of clear differential

conditioning between the CS+V and the CS-. Our first observation supporting the utility of

imagery exposure during extinction learning comes from the observation of the generalization of

conditioned fear from perception to imagery during Extinction Phase 1 in terms of both SCR and

self-reported fear. An alternative possibility is that the elevation of SCR when imagining the

CS+I could be the result of the act of imagery, which may require more effort than simply

viewing in the case of CS-. The initial purpose of including the CS+V during Extinction Phase 1

was to help evaluate this possibility. On the other hand, previous research on the generalization

of differential fear conditioning from visual to imagined objects did not observe evidence of an

elevated SCR response to mental imagery compared to viewing, though they did observe a

significant SCR difference between the CS+ versus the CS- (Greening et al., 2021). Another

study by Agren et al. (2017) reported that while the conditioned response measured by SCR

difference between CS+ and CS- was initially larger in the imaginal extinction group, the

imagery and in vivo/perceptual extinction groups were comparable by late extinction. Based on these findings, it is unlikely that the SCR difference can be completely accounted for by the difference between viewing versus visual imagery per se, rather than being an expression of the generalized conditioned fear response.

The second piece of evidence supporting the use of mental imagery for fear extinction learning comes from Extinction Phase 2. Specifically, even during the early trials (i.e., the first four trials per condition) the CS+I no longer exhibited a greater SCR compared to the CS-. Had imagery fear extinction not affected the fear conditioned association between the perceptual in vivo CS+I and the CS- one would have predicted a persistent differential conditioning upon re-exposure to the in vivo CS+I during Extinction Phase 2, which was not observed. Unlike SCR, differential fear measured as self-reported fear did persist to some degree throughout extinction. However, this differential was significantly reduced from Acquisition to Extinction Phase 1 and from Extinction Phase 1 to Extinction Phase 2 irrespective of the type of CS+ (i.e., for both CS+I – CS- and CS+V – CS-). This is in line with previous research on self-reported fear (Lau et al., 2008; Shechner et al., 2015), and may reflect that individuals maintain stable declarative knowledge regarding the difference between the CS+s and the CS- even after extinction.

While the present findings provide evidence that mental imagery can facilitate extinction learning, there are at least two potential mechanistic explanations underpinning the observed effects. The first possibility is that mental imagery operates via depictive mechanisms (Lewis et al., 2013) such that during extinction, the imagery of the CS+ acts as a substitute for the in vivo (i.e., visual) stimulus (e.g., Greening et al., 2021). Alternatively, our results could be the result of propositional mechanisms, e.g., the controlled reasoning that viewing CS+ is no longer accompanied by the US since imagining CS+ was not (Mitchell, De Houwer, & Lovibond, 2009).

Future investigations are needed to further evaluate the underlying cognitive mechanisms of fear extinction during affective mental imagery, for example, by controlling for the potential confound of verbalization as has been done in previous research on mental imagery and emotion (Holmes & Mathews, 2005). See Mertens et al. (2020) for additional considerations when designing control conditions. Not only will these discoveries better our understanding of the operations of affective mental imagery, they would also be of great clinical significance (Andrade, May, & Kavanagh, 2012; Hirsch & Holmes, 2007).

Unexpectedly, only one of the two CS+s (i.e., CS+I) was found to exhibit threat related differential physiological reactivity (i.e., greater SCR for the CS+I vs. the CS-) following the Acquisition phase in the current study. Considering that the letter selections of the CS+I and the CS+V were counterbalanced across participants, this could not be attributed to differences between the letter stimuli of "J" and "H". One possibility is that the nature of the stimuli and the use of two CS+s in a within-subject designed made the stimulus discriminability for the purposes of differential conditioning difficult. Another potential contributor could be the length of the Acquisition phase. While a shock check was conducted during Acquisition, participants still exhibited habituation to the US towards the second half of Acquisition based on visual examination of trial-by-trial SCR (see supplementary figure S2). Additionally, compared to evolutionarily fear-relevant objects (e.g., images of snakes, human faces), stimuli lacking in evolutionary fear-relevance, such as the letter stimuli used in the current study, undergo fear conditioning less readily (Öhman & Mineka, 2001). Future research could consider using more fear-relevant objects (e.g., snakes or spiders), shorter acquisition procedures, or a between-group design to avoid require two CS+s (e.g., Agren et al., 2017; Reddan et al., 2018). The

inconsistency observed in the current study could also be a manifestation of the noise and measurement error in SCR.

Altogether, the present results from both SCR and self-reported fear suggest that repeated exposure to the mental image of an in vivo CS+ can facilitate extinction learning. Both the complementary findings and inferential similarities revealed between the subjective and physiological measures also demonstrate the importance of collecting multiple outcome measures based on a multi-component view of emotion (Frijda, 1986; Gross, 2013; Lang, 1993; Larsen & Prizmic-Larsen, 2006; LeDoux & Pine, 2016). The current study along with the existing experimental literature (Agren et al., 2017; Reddan et al., 2018) provide preliminary support for the potential clinical application of extinction-based imagery treatments in fear-related affective conditions.

References

Abramowitz, J. S. (1996). Variants of exposure and response prevention in the treatment of obsessive-compulsive disorder: A meta-analysis. *Behavior Therapy*, *27*(4), 583–600. https://doi.org/10.1016/S0005-7894(96)80045-1

Achterberg, J. (2002). *Imagery in healing: Shamanism and modern medicine*. Shambhala Publications.

Agren, T., Björkstrand, J., & Fredrikson, M. (2017). Disruption of human fear reconsolidation using imaginal and in vivo extinction. *Behavioural Brain Research*, *319*, 9–15. https://doi.org/10.1016/j.bbr.2016.11.014

Andrade, J., May, J., & Kavanagh, D. (2012). Sensory Imagery in Craving: From Cognitive Psychology to New Treatments for Addiction. *Journal of Experimental Psychopathology*, *3*(2), 127–145. https://doi.org/10.5127/jep.024611

Arntz, A. (2012). Imagery rescripting as a therapeutic technique: Review of clinical trials, basic studies, and research agenda. *Journal of Experimental Psychopathology*, *3*(2), 189–208. https://doi.org/10.5127/jep.024211

Bach, D. R., Flandin, G., Friston, K. J., & Dolan, R. J. (2010). Modelling event-related skin conductance responses. *International Journal of Psychophysiology*, *75*(3), 349–356. https://doi.org/10.1016/j.ijpsycho.2010.01.005

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436.

Carter, R. M., Hofstotter, C., Tsuchiya, N., & Koch, C. (2003). Working memory and fear conditioning. *Proceedings of the National Academy of Sciences*, *100*(3), 1399–1404. https://doi.org/10.1073/pnas.0334049100

Critchley, H., Mathias, C., & Dolan, R. J. (2002). Fear-conditioning in humans: the influence of

awareness and arousal on functional neuroanatomy. *Neuron*, *33*, 653–663.

Frijda, N. H. (1986). *The emotions*. Cambridge University Press.

Greening, S. G., Lee, T.-H., Gregoire, L., Burleigh, L., Robinson, T., Jiang, X., … Kaplan, J.

(2021). Fear in the mind's eye: Mental imagery can generate and regulate acquired

differential fear conditioned reactivity. *BioRxiv*, 2021.02.04.429795.

https://doi.org/https://doi.org/10.1101/2021.02.04.429795

Grégoire, L., & Greening, S. G. (2019). Opening the reconsolidation window using the mind's

eye: Extinction training during reconsolidation disrupts fear memory expression following

mental imagery reactivation. *Cognition*, *183*(November 2018), 277–281.

https://doi.org/10.1016/j.cognition.2018.12.001

Grégoire, L., & Greening, S. G. (2020). Fear of the known: semantic generalisation of fear

conditioning across languages in bilinguals. *Cognition and Emotion*, *34*(2), 352–358.

https://doi.org/10.1080/02699931.2019.1604319

Gross, J. J. (2013). *Handbook of emotion regulation*. Guilford publications.

Hecker, J. E. (1990). Emotional processing in the treatment of simple phobia: A comparison of

imaginal and in vivo exposure. *Behavioural Psychotherapy*, *18*(1), 21–34.

https://doi.org/10.1017/S0141347300017961

Hirsch, C. R., & Holmes, E. A. (2007). Mental imagery in anxiety disorders. *Psychiatry*, *6*(4),

161–165. https://doi.org/10.1016/j.mppsy.2007.01.005

Holmes, E. A., Arntz, A., & Smucker, M. R. (2007). Imagery rescripting in cognitive behaviour

therapy: Images, treatment techniques and outcomes. *Journal of Behavior Therapy and

Experimental Psychiatry*, *38*(4), 297–305. https://doi.org/10.1016/j.jbtep.2007.10.007

Holmes, E. A., & Mathews, A. (2005). Mental imagery and emotion: A special relationship?

Emotion. https://doi.org/10.1037/1528-3542.5.4.489

Jiang, X., Burleigh, L., & Greening, S. G. (2021). Complete the triangulation: Quantifying

differential fear conditioning with a noninterfering and sensitive behavioral measure along

with self☐report and physiological measures. *Psychophysiology*.

https://doi.org/10.1111/psyp.13831

Kleiner, M., Brainard, D. H., Pelli, D., Ingling, A., Murray, R., & Broussard, C. (2007). What's

new in Psychtoolbox-3. In *Perception* (Vol. 36). https://doi.org/10.1068/v070821

Kober, H., Barrett, L. F., Joseph, J., Bliss-moreau, E., Lindquist, K., & Wager, T. D. (2008).

Functional grouping and cortical - subcortical interactions in emotion. *NeuroImage*, *42*(2),

998–1031. https://doi.org/10.1016/j.neuroimage.2008.03.059

Koizumi, A., Amano, K., Cortese, A., Shibata, K., Yoshida, W., Seymour, B., … Lau, H. (2016).

Fear reduction without fear through reinforcement of neural activity that bypasses conscious

exposure. *Nature Human Behaviour*, *1*(1), 0006. https://doi.org/10.1038/s41562-016-0006

Kosslyn, S. M. (1981). The medium and the message in mental imagery: A theory.

Psychological Review, *88*(1), 46–66. https://doi.org/10.1037/0033-295X.88.1.46

Kosslyn, S. M. (2005). Mental images and the brain. *Cognitive Neuropsychology*, *22*(3–4), 333–

347. https://doi.org/10.1080/02643290442000130

Kosslyn, S. M., Ganis, G., & Thompson, W. L. (2001). Neural foundations of imagery. *Nature*

Reviews Neuroscience, *2*(9), 635–642. https://doi.org/10.1038/35090055

Kosslyn, S. M., Thompson, W. L., & Ganis, G. (2006). *The case for mental imagery*. Oxford

University Press.

Kreiman, G., Koch, C., & Fried, I. (2000). Imagery neurons in the human brain. *Nature*,

408(6810), 357–361. https://doi.org/10.1038/35042575

Krypotos, A.-M., Klugkist, I., & Engelhard, I. M. (2017). Bayesian hypothesis testing for human

threat conditioning research: an introduction and the condir R package. *European Journal

of Psychotraumatology*, *8*(sup1), 1314782. https://doi.org/10.1080/20008198.2017.1314782

Lang, P. J. (1977). Imagery and therapy: An information processing analysis of fear. *Behavior

Therapy*, *8*, 862–889. https://doi.org/10.1016/S0005-7894(77)80157-3

Lang, P. J. (1979). A Bio☐Informational Theory of Emotional Imagery. *Psychophysiology*, Vol.

16, pp. 495–512. https://doi.org/10.1111/j.1469-8986.1979.tb01511.x

Lang, P. J. (1993). The three-system approach to emotion. *The Structure of Emotion*, 18–30.

Larsen, R. J., & Prizmic-Larsen, Z. (2006). Measuring emotions: Implications of a multimethod

perspective. In *Handbook of multimethod measurement in psychology.* (pp. 337–351).

https://doi.org/10.1037/11383-023

Lau, J. Y. F., Lissek, S., Nelson, E. E., Lee, Y., Roberson-Nay, R., Poeth, K., … Pine, D. S.

(2008). Fear conditioning in adolescents with anxiety disorders: Results from a novel

experimental paradigm. *Journal of the American Academy of Child & Adolescent

Psychiatry*, *47*(1), 94–102. https://doi.org/10.1097/chi.0b01e31815a5f01

LeDoux, J. E., & Pine, D. S. (2016). Using neuroscience to help understand fear and anxiety: A

two-system framework. *American Journal of Psychiatry*, *173*(11), 1083–1093.

https://doi.org/10.1176/appi.ajp.2016.16030353

Lewis, D. E., O'Reilly, M. J., Khuu, S. K., & Pearson, J. (2013). Conditioning the mind's eye:

Associative learning with voluntary mental imagery. *Clinical Psychological Science*, *1*(4),

390–400. https://doi.org/10.1177/2167702613484716

Lim, S. L., & Pessoa, L. (2008). Affective Learning Increases Sensitivity to Graded Emotional

Faces. *Emotion*, *8*(1), 96–103. https://doi.org/10.1037/1528-3542.8.1.96

Lonsdorf, T. B., Menz, M. M., Andreatta, M., Fullana, M. A., Golkar, A., Haaker, J., … Merz, C.

   J. (2017). Don't fear "fear conditioning": Methodological considerations for the design and

   analysis of studies on human fear acquisition, extinction, and return of fear. *Neuroscience*

   and Biobehavioral Reviews, *77*, 247–285. https://doi.org/10.1016/j.neubiorev.2017.02.026

Matthews, N. L., Collins, K. P., Thakkar, K. N., & Park, S. (2014). Visuospatial imagery and

   working memory in schizophrenia. *Cognitive Neuropsychiatry*, *19*(1), 17–35.

   https://doi.org/10.1080/13546805.2013.779577

Mauss, I. B., Levenson, R. W., McCarter, L., Wilhelm, F. H., & Gross, J. J. (2005). The tie that

   binds? Coherence among emotion experience, behavior, and physiology. *Emotion*, *5*(2),

   175–190. https://doi.org/10.1037/1528-3542.5.2.175

Mertens, G., Krypotos, A.-M. M., & Engelhard, I. M. (2020). A review on mental imagery in

   fear conditioning research 100 years since the 'Little Albert' study. *Behaviour Research*

   and Therapy, *126*(August 2019), 103556. https://doi.org/10.1016/j.brat.2020.103556

Milad, M. R., & Quirk, G. J. (2012). Fear extinction as a model for translational neuroscience:

   Ten years of progress. *Annual Review of Psychology*, *63*, 129–151.

   https://doi.org/10.1146/annurev.psych.121208.131631

Mitchell, C. J., De Houwer, J., & Lovibond, P. F. (2009). The propositional nature of human

   associative learning. *Behavioral and Brain Sciences*, *32*(2), 183–198.

   https://doi.org/10.1017/S0140525X09000855

Murphy, F. C., Nimmo-Smith, I., & Lawrence, A. D. (2003). Functional neuroanatomy of

   emotions: A meta-analysis. *Cognitive, Affective, & Behavioral Neuroscience*, *3*(3), 207–233.

   https://doi.org/10.3758/CABN.3.3.207

Öhman, A., & Mineka, S. (2001). Fears, phobias, and preparedness: Toward an evolved module

of fear and fear learning. *Psychological Review*, *108*(3), 483–522.

https://doi.org/10.1037/0033-295X.108.3.483

Pearson, J., & Kosslyn, S. M. (2015). The heterogeneity of mental representation: Ending the

imagery debate. *Proceedings of the National Academy of Sciences of the United States of

America*, *112*(33), 10089–10092. https://doi.org/10.1073/pnas.1504933112

Pearson, J., Naselaris, T., Holmes, E. A., & Kosslyn, S. M. (2015). Mental Imagery: Functional

Mechanisms and Clinical Applications. *Trends in Cognitive Sciences*, *19*(10), 590–602.

https://doi.org/10.1016/j.tics.2015.08.003

Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics: transforming numbers

into movies. *Spatial Vision*, *10*(4), 437–442.

Phan, K. L., Wager, T., Taylor, S. F., & Liberzon, I. (2002). Functional neuroanatomy of

emotion: A meta-analysis of emotion activation studies in PET and fMRI. *NeuroImage*,

16(2), 331–348. https://doi.org/10.1006/nimg.2002.1087

Powers, M. B., Halpern, J. M., Ferenschak, M. P., Gillihan, S. J., & Foa, E. B. (2010). A meta-

analytic review of prolonged exposure for posttraumatic stress disorder. *Clinical

Psychology Review*, *30*(6), 635–641. https://doi.org/10.1016/j.cpr.2010.04.007

Reddan, M. C., Wager, T. D., & Schiller, D. (2018). Attenuating neural threat expression with

imagination. *Neuron*, *100*(4), 994-1005.e4. https://doi.org/10.1016/j.neuron.2018.10.047

Saville, D. J. (1990). Multiple Comparison Procedures: The Practical Solution. *The American

Statistician*, *44*(2), 174–180. https://doi.org/10.1080/00031305.1990.10475712

Schaefer, A., Collette, F., Philippot, P., Linden, M. Van der, Laureys, S., Delfiore, G., … Salmon,

E. (2003). Neural correlates of "hot" and "cold" emotional processing: A multilevel

approach to the functional anatomy of emotion. *NeuroImage*, *18*(4), 938–949.

https://doi.org/10.1016/S1053-8119(03)00009-0

Schiller, D., Kanen, J. W., LeDoux, J. E., Monfils, M.-H., & Phelps, E. A. (2013). Extinction during reconsolidation of threat memory diminishes prefrontal cortex involvement. *Proceedings of the National Academy of Sciences*, *110*(50), 20040–20045. https://doi.org/10.1073/pnas.1320322110

Schiller, Daniela, Monfils, M.-H. M. H., Raio, C. M., Johnson, D. C., Ledoux, J. E., & Phelps, E. A. (2010). Preventing the return of fear in humans using reconsolidation update mechanisms. *Nature*, *463*(7277), 49–53. https://doi.org/10.1038/nature08637

Shechner, T., Britton, J. C., Ronkin, E. G., Jarcho, J. M., Mash, J. A., Michalska, K. J., … Pine, D. S. (2015). Fear conditioning and extinction in anxious and nonanxious youth and adults: Examining a novel developmentally appropriate fear-conditioning task. *Depression and Anxiety*, *32*(4), 277–288. https://doi.org/10.1002/da.22318

Sjouwerman, R., Niehaus, J., Kuhn, M., & Lonsdorf, T. B. (2016). Don't startle me— Interference of startle probe presentations and intermittent ratings with fear acquisition. *Psychophysiology*, *53*(12), 1889–1899. https://doi.org/10.1111/psyp.12761

Taschereau-Dumouchel, V., Kawato, M., & Lau, H. (2019). Multivoxel pattern analysis reveals dissociations between subjective fear and its physiological correlates. *Molecular Psychiatry*. https://doi.org/10.1038/s41380-019-0520-3

Turner, S. M., Beidel, D. C., & Jacob, R. G. (1994). Social phobia: A comparison of behavior therapy and atenolol. *Journal of Consulting and Clinical Psychology*, *62*(2), 350–358. https://doi.org/10.1037/0022-006X.62.2.350

Vervliet, B., Vansteenwegen, D., Baeyens, F., Hermans, D., & Eelen, P. (2005). Return of fear in a human differential conditioning paradigm caused by a return to the original acquistion

context. *Behaviour Research and Therapy*, *43*(3), 323–336.

https://doi.org/10.1016/j.brat.2004.01.001

Weßlau, C., & Steil, R. (2014). Visual mental imagery in psychopathology - Implications for the

maintenance and treatment of depression. *Clinical Psychology Review*, *34*(4), 273–281.

https://doi.org/10.1016/j.cpr.2014.03.001

Author Notes

Funding: This research was supported a Louisiana Board of Regents – Research

Competitiveness Subprogram grant to S.G.G.

Conflict of Interest: The authors declare that they have no conflict of interest.

Figure 1. Letter stimuli (A), probes, used in the two training runs only (B), and trial designs for view (C) and imagine (D) trials in

Acquisition, Extinction Phase 1 and Extinction Phase 2.

Table 1. Trial schedule for each phase of the study.

	Training/Habituation	Acquisition	Extinction Phase 1	Extinction Phase 2
CS+I with US		8 view trials		
CS+V with US		8 view trials		
CS+I without US	4 view, 4 imagine trials	8 view trials	8 imagine trials	8 view trials
CS+V without US	4 view, 4 imagine trials	8 view trials	8 view trials	8 view trials
CS-	4 view, 4 imagine trials	16 view trials (8 excluded in analyses)	8 view trials	8 view trials

Figure 2. Mean square-rooted SCR for each CS type during Acquisition (A), and early and late Extinction Phase 1 (B) and Extinction Phase 2 (C). Due to insignificant difference between CS+V and CS- during Acquisition, CS+V was not included in Extinction Phase 1 and Extinction Phase 2. Error bars show ± 1 standard error. Each dot represents one subject.

$* p < .05.$ $** p < .01.$ $*** p < .001.$ $**** p < .0001.$

Figure 3. Mean self-reported fear ratings (A) and shock estimations (B) for each CS type. Error bars show ± 1 standard error. Each dot represents one subject.

p < .05. ** *p* < .01. *** *p* < .001. **** *p* < .0001.

Figure S1. Mean square-rooted SCR for each CS type during early and late Extinction Phase 1 (A) and Extinction Phase 2 (B). Error bars show ± 1 standard error. Each dot represents one subject.

Figure S2. Trial-by-trial square-rooted SCR for all three CS type during Acquisition, Extinction Phase 1, and Extinction Phase 2. Error bars show ± 1 standard error.