Exosomal TAR DNA binding protein 43 profile in canine model of
amyotrophic lateral sclerosis: A preliminary study in developing blood-
based biomarker for neurodegenerative diseases

Penelope Pfeiffer [1†], Joan R. Coates [2], Yajaira M. Esqueda[1] Andrew Kennedy[1],
Kyleigh Getchell[1], Myra McLenon[1], Edina Kosa[1], Abdulbaki Agbas[1, 3]*

[1]*Kansas City University, Kansas City MO;* [2]*University of Missouri-Columbia, MO;*
[3]*Heartland Center for Mitochondrial Medicine, Kansas City KS;* [†]*Mount Sinai Hospital,*
Chicago IL;
9
10 *Correspondence: Abdulbaki Agbas, aagbas@kansascity.edu; Tel.: +1 816-654-7614

11 †Current affiliation

12 **Biographical notes**

13
14 **Penelope Pfeiffer, DO:** Dr. Pfeiffer graduated from Kansas City University in 2016 and currently she
15 is working as staff physician in Mount Sinai Hospital, Chicago, IL

16
17 **Joan R. Coates**, DVM, MS, DACVIM–Neurology
18 Professor of Neurology and Neurosurgery
19 Joan Coates specializes in neurology and neurosurgery and is Section Head of the Neurology &
20 Neurosurgery Section/Service. Her research focus involves the study of canine degenerative
21 myelopathy as a disease model for amyotrophic lateral sclerosis (ALS, Lou Gehrig's disease).

22
23 **Yajaira M. Esqueda:** Yajaira M. Esqueda worked with Dr. Agbas as graduate students and currently,
24 she is searching other career opportunities.

25
26 **Andrew Kennedy:** Andrew Kennedy was a research student in Dr. Agbas' Lab and he is a medical
27 student at Kansas City University at present time.

28
29 **Kyleigh Getchell**: Kyleigh Getchell was a research student in Dr. Agbas' Lab and she is a medical
30 student at Kansas City University at present time.

31
32 **Myra McLenon:** Myra McLenon is currently working in Dr. Agbas' lab and applying for medical
33 school.

34
35 **Edina Kosa, MSc:** Edina Kosa is a Research Associate to Dr. Agbas. She functions as a laboratory
36 manager. She is involved in training of student researchers and research fellows in Dr, Agbas' lab.
37 She also contributes to the many aspects of the research projects.

38
39 **Abdulbaki Agbas**, MSc, PhD
40 Professor of Biochemistry in Basic Science Department of the Kansas City University. His lab does
41 research in Neuroscience and Biochemistry. Their current projects are (1) Blood-based early
42 biomarker development for neurodegenerative disease, (2) Bioenergetics in neurodegenerative
43 diseases, (3) platelet mitochondria respiration profile in patients with Alzheimer's and ALS diseases.

Exosomal TAR DNA binding protein 43 profile in canine model of amyotrophic lateral sclerosis: A preliminary study in developing blood-based biomarker for neurodegenerative diseases.

Abstract:

Objective: Blood-based biomarkers provide a crucial information in progress of neurodegenerative diseases with minimally invasive sampling method. Validated blood-based biomarker application in people with amyotrophic lateral sclerosis would derive numerous benefits. Canine degenerative myelopathy is a naturally occurring animal disease model to study the biology of human amyotrophic lateral sclerosis. Serum derived exosomes are potential carriers for cell-specific cargoes making them ideal venue to study biomarkers for a variety of diseases and biological processes. This study assessed the exosomal proteins that may be assigned as surrogate biomarker that may reflect biochemical changes in central nervous system.

Methods: Exosomes were isolated from canine serum using commercial exosome isolation reagents. Exosomes target proteins contents were analysed by Western blotting method.

Results: The profiles of potential biomarker candidates in spinal cord homogenate and that of serum-derived exosomes were found elevated in dogs with degenerative myelopathy as compare to control subjects.

Conclusions: Serum-derived exosomal biomolecules can serve as surrogate biomarkers in neuro degenerative diseases.

Keywords: TDP-43, amyotrophic lateral sclerosis, degenerative myelopathy, exosome, biomarker

Key Messages

• A canine with degenerative myelopathy can serve as a model animal to study human amyotrophic

lateral sclerosis.

• Serum-derived exosomes contains Transactive Response DNA Binding Protein 43 (TDP-43),

potential biomarker candidate.

• The levels of spinal cord TDP-43 proteins and that of serum-derived exosomes exhibited a similar

profiling. Therefore, serum derived exosomes may be used as a venue for establishing blood-

based biomarkers for neurodegenerative diseases.

1. Introduction

The aberrant protein aggregation in motor neurons is the hallmark of amyotrophic lateral

sclerosis (ALS). ALS is a progressive disease that directly affects motor neurons, leading to

loss of muscle function. About 5-10% of the inherited forms of ALS are linked to a mutation

in the SOD1 gene *(SOD1).* The misfolded mutant [Cu/Zn] superoxide dismutase (SOD1)

protein is believed to contribute to development of ALS, yet the role of misfolded non-mutant

SOD1 in the disease progress is unclear. Recently, chemically modified aberrant Transactive

Response DNA Binding Protein 43 (TDP-43) species were found to represent a major

accumulating protein in neuronal cytoplasmic inclusions (1, 2) and in exosomes (3, 4) in

Frontotemporal Lobar Degeneration (FTLD), and in ALS patients (5). Exosomes are nano-

size membranous vesicles that contains several macromolecules including aberrant

pathological proteins (6). The size and membranous structure of exosomes allow them to pass

through the blood brain barrier (7-9). These features make exosomes a potential platform in

which targeted biomolecules can be analysed.

The biomolecular link and clinical similarities between human ALS and canine degenerative

myelopathy (DM) suggested that human ALS hallmark signature proteins (i.e., SOD1 and TDP-43) have the same or similar profile in canine DM. These similarities may be employed to further biomarker validity studies such as reproducibility, specificity, sensitivity, and robustness. As a naturally occur disease in the pet population, canine DM has many similarities to human ALS and is thought to be a result, in part, to mutated SOD1 protein in dogs.   Thus, dogs affected with DM can serve as an animal disease model for ALS based on CNS size and complexity, and disease characteristics.

ALS is an incurable and fatal disease with a patients survival rate of 3-5 years once diagnosed (10). In most patients, the symptoms begin in the lower limbs (11, 12). Patients often complain of tripping, stumbling, foot drop, or a "slapping" gait. Upper limb involvement includes decreased dexterity in the fingers, cramping, stiffness, weakness and wrist drop (13). Early diagnosis of ALS is very difficult, which can delay time to treatment (14). Establishing a blood-based biomarker(s) will help clinicians with early diagnosis and initiating timely treatment. The pharmaceutical intervention with well-timing may improve the quality of life and lifespan for ALS patients.

ALS researchers have used rodents as an animal model with mutated SOD1 to study the pathology of the disease (15). The rodent model is of limited use as disease progression is very short (3-6 months) and is based on overexpression of the human gene for mutant SOD1. In addition, peripheral biomarker studies are difficult to perform in a longitudinal course due to limited access to serial samples of serum/exosomes in mice due to the small volume of blood. Canine DM is a late adult neurodegenerative disease accompanied by (*SOD1)* mutations (*SOD1:c.118A, SOD1:c52T*), and protein aggregation (16, 17). Early clinical signs include general proprioceptive ataxia, and spastic upper motor neuron paresis in the pelvic limbs with progression to flaccid tetraplegia and dysphagia (18-20). DM and ALS are

characterized by progressive multisystem neurodegeneration involving upper and lower

motor neurons. Canine DM phenotypically resembles upper motor onset ALS in reference to

clinical and histopathologic features. The distribution of lesions and clinical disease

progression in DM are similar to that reported for the UMN-onset ALS (21) with UMN signs

in DM-affected dogs progressing later to LMN signs (16, 20).

An abnormal accumulation of TDP-43 has been demonstrated in the cytoplasm and platelets

from ALS patients (22). The post translationally modified derivatives of TDP-43 (i.e., hyper-

phosphorylation and ubiquitination) have been studied in ALS (23, 24). Prior research has

established that the accumulation of TDP-43 in the cytosol was causal in the loss of function

in affected neurons (25, 26). However, the percent of TDP-43 in peripheral blood, which

represents the central nervous system (CNS) origin of TDP-43, is unknown. Therefore,

assessment of the confined TDP-43 in exosomes may reflect the changes in TDP-43 protein

better than that of large volume of serum or plasma where the target proteins are diluted. This

study explored the feasibility of using exosomes to assess potential biomarkers for DM such

as TDP-43, pTDP-43, and SOD1.

2. Materials and Methods
2.1. Animals

Previously -80^{0}C frozen spinal cord tissues from thoracic region and serum samples were

obtained from Dr. Joan R. Coates at the College of Veterinary Medicine, University of

Missouri-Columbia under the approval of institutional animal care and use committee

(IACUC# 10077). Pet owners signed an informed consent for samples being used for research

purposes.

Whole blood was collected from the jugular or cephalic veins. Blood was centrifuged and serum was placed in 500 microliter aliquot tubes. Aliquots were stored at -80^0C. The dog was necropsied for CNS tissue collections. The muscle was removed from the dorsal column of the spine. The dorsal lamina just medial to the articular processes was removed to expose the spinal cord. Specifically, the thoracic spinal cord was removed and sectioned accordingly. The tissues were immediately frozen on dry ice and stored at -80^0C.

Tissues were harvested from various breeds with diagnosed DM and age-matched dogs not affected with DM (Table 1 in Supplemental Data). All DM affected dogs were homozygous for the SOD1 E40K mutation (20).

2.2. Isolation of exosomes from canine serum

The isolation of exosomes from serum of DM and control dogs was performed using a commercially available particle precipitation method following manufacturer's protocol (miRCURY Exosome Serum/Plasma Kit, Qiagen #76603). The exosome isolation procedure was based on the capture of water molecules, which otherwise would form the hydrate envelope of particles in suspension. Mixing the starting sample with a proprietary precipitation buffer diminished the hydration of the subcellular particles and allowed the precipitation of even particles smaller than 100 nanometer with a low-speed centrifugation step (27). Before proceeding to the isolation of exosomes, the serum samples were first pelleted by centrifugation, and the supernatant was filtered through 0.20-micrometer filtration disk to remove residual cells, debris, platelets, large micro vesicles etc. The samples were incubated with a proprietary precipitation buffer for 1 hour at 4°C. After a 30-minute low speed centrifugation at 500 x g, the exosomes pellets were re-suspended for further analysis of their content. The isolated exosomes was verified by immunoprobing assay using an

antibody that recognized exosome surface protein, tumor susceptibility 101 protein (TSG101)

(Supplemental Figure-3). Exosome suspension was aliquoted and kept in -20^0C until use.

2.3. Spinal cord homogenate preparation

50-60 milligram thoracic spinal cord tissue (T3 section) was homogenized in an ice-cold buffer

(0.32 M sucrose, 0.5 mM $MgSO_4$, 10 mM ε-amino-n-caproic acid, protease inhibitor cocktail

(0.05% v/v) (Calbiochem #539134), phosphatase inhibitor cocktails set II (0.1% v/v)

(Calbiochem # 524625), 10 mM HEPES, pH 7.4) in a glass pestle-glass homogenizer (8-10

strokes). The tissue/homogenization buffer ratio was one (mg tissue) / 15 (µL buffer). The

homogenate was resuspended 8-10 times using a 1 mL syringe with a 26^{-gauge} needle to shear

DNA and liberate nuclear proteins. The homogenate was incubated on ice for 20 minutes and

subjected to centrifugation at 16,000 X g for 20 minutes at 4^0C. The supernatant containing all

soluble proteins, including nuclear proteins was collected into a clean tube. For purpose of the

protein assay, 5-10 µL aliquot was saved. Total protein estimation was analyzed by BCA assay

(ThermoFisher Scientific #23225).The sample was aliquoted into 100 µL volumes into

microfuge tubes and stored at -80°C until use.

2.4.. Immunoblotting analysis

10 µg of exosomal proteins were loaded onto a commercial 4-20% SDS/PAG gradient

gel. The electrophoresis was run at 100V for 75-80 min until the dye-front migrated to

0.5 cm from the bottom of the gel. Resolved proteins on the gel were transferred onto a

PVDF membrane by electro transfer unit at 60 V for 90 min. The membrane was

stained/destained for total protein staining according to manufacturer's protocol

(REVERT Total Protein Stain kit, LI-COR; Cat# 926-11016). The membrane was

blocked with a blocking agent (SEA BLOCK Blocking, Thermo Scientific, Cat #

UH2788881) for one hour at room temperature, followed by overnight incubation with

anti- TDP-43 Ab (1:1000 dilution; Proteintech Cat# 10782-2-AP), anti-P(S409/410)

TDP-43 Ab (1:1000 dilution; Proteintech Cat# 66318-1-Ig), and anti-SOD1 Ab (1:750

dilution; Proteintech Cat#10269-1-AP) on an orbital shaker at 4^0C. The next day the

membrane was incubated with Infrared (IR)-tagged antibody (1:10,000 dilution; LI-

COR Cat# 926-3211, 926-68071, 926-32210, 926-68070) for 1-2 hours at room

temperature. The protein bands were visualized in an image analyser (LI-COR Odyssey

Infrared Imager (Model No. 9120). The band intensity was normalized based on total

protein staining signals, and analysed by Image Studio image analysing program

(V.3.1, Li-COR Biosciences).

3. Results

3.1. Dogs with DM spinal cord homogenate show elevated TDP-43 and SOD1 in the thoracic

region of the spinal cord

The thoracic region of the spinal cord from companion dogs with DM (n=4) and from age

matched non-DM affected control dogs (n=4) were obtained from a tissue archive at the

University of Missouri College of Veterinary Medicine. The signalment of dogs used in this

study listed in **Table-1** in Supplemental Data. **Figure-1** shows that TDP-43 protein levels are

elevated in the DM spinal cord region as compared to that of the control dogs. A t-test analysis

indicated that differences between two groups were significant (P< 0.01). This observation

agrees of other published data that TDP-43 accumulation has been observed in human spinal

cord (23, 28). SOD1-immunoreactive aggregates were observed in ventral horn motor

neurons (29). This study confirmed that the spinal cord SOD1 aggregation profile occurred

in dogs with DM (Fig.2). In this study, the SOD1 aggregation profile was considered as

positive control.

3.2. Exosomal TDP-43 , phosphorylated TDP-43, and SOD1 are elevated in canine with DM

Frozen serum samples from dogs were obtained from a tissue archive at the University Of

Missouri College Of Veterinary Medicine. The signalments of dogs used in this study were

listed in **Table-2 and Table-3** in Supplemental Data. **Figure-3 (A, B)** showed that both total

TDP-43 and its phosphorylated derivative (pTDP-43) were elevated in exosomes of the dogs

affected with DM as compared to that of the control dogs. A significant difference ($P \leq 0.001$)

was found between DM affect and control dogs. These preliminary results suggest that

exosomes can be used to assess biomarker proteins such as TDP-43 and its phosphorylated

derivatives.

Serum-derived exosomal SOD1 protein levels were elevated in dogs affected with DM

as compare to that of control dogs. The difference between the two groups was

significant based on an unpaired t-test analysis ($P \leq 0.05$). We have observed that SOD1

proteins in canine serum- derived exosomes exhibited a heterogenic profile (i.e.,

monomer, trimer and tetramer) (Fig.4). Soluble monomers (~25 kDa) were considered

in the data analyses and graphing.

4. Discussion

Complexity of a biosample matrix and the invasive nature of tissue sampling led investigators

to develop peripheral tissue-based biomarkers for ALS. Blood-based biomarker initiatives

have gained popularity due to easy access for sampling with minimal invasion (i.e., venous

puncture), access to several blood cells, and subcellular structures. Once developed and

validated, blood-based biomarkers will be utilized for diagnosis, monitoring the prognosis,

and monitoring the treatment of ALS (30-32).

Canine degenerative myelopathy (DM) is an adult-onset, progressive neurodegenerative

disease that shares important similarities to ALS including clinical presentation and

progression, pathological features, and *SOD1* mutations (33). This naturally occurring disease

in the companion dog recapitulates some forms of human mutant SOD1 ALS. The E40K

SOD1 mutation is widespread among companion dogs (34) and leads to SOD1 aggregate

accumulations within cells, the putative mechanism of disease in DM and in both familial and

sporadic forms of human ALS (16, 35-37).

In this preliminary study, we demonstrated that serum-derived exosomes may be utilized for

assessing ALS hallmark proteins (i.e., TDP-43, pTDP-43 and SOD1), and compared to that

of thoracic spinal cord homogenate in DM affected dogs. We chose the thoracic region

because the disease pathology of canine DM is most severe in this region (33). This region

also reflects more TDP-43 accumulation relevant to pathology of ALS (38). Serum exosomes

were isolated from the dog cohort consisting of animals with DM and age matched non-DM

affected. Due to some technical obstructions and limited tissue amount, we were unable to

isolate sufficient exosomes from the spinal cord tissue from this cohort. We are currently

refining our techniques to increase the exosome yield.

This study demonstrated the feasibility of the followings: (i) effectively isolating exosomes

from serum, (ii) analyse exosomal TDP-43 and pTDP-43 protein profiles, and (iii) comparing

those profiles to spinal cord tissue (thoracic region) homogenate from the different cohort of

dogs with DM as part of proof- of- concept studies. TDP-43 and SOD1 are known as the

hallmark proteins for ALS (23, 24, 39). This study demonstrated that total homogenates of

spinal cord TDP-43 levels were elevated in dogs with DM (Fig.1). The SOD1 profile served

as a positive control and established that both proteins were elevated (Fig.2). The next logical

step would be to demonstrate sensitivity and specificity to establish biomarker reliability

within the study limitations due to lack of the biosample archive did not follow the specified

minimal information for studies of extracellular vesicles, MISEV2018 (40).

Exosomes are small stable single-membrane organelles that carry selected proteins, lipids,

nucleic acids, and glycoconjugates (6). Exosome biogenesis is a valuable mechanism to

assess the protein quality control (6). They are emerging as key mediators of communication

and waste management among neurons, glial cells and connective tissues during normal and

diseases conditions (41). Although exosomes are being developed as therapeutic agents in

multiple disease models, they can also serve as surveillance agents by transporting aberrant

proteins like hyper-phosphorylated TDP-43 or cytosolic TDP-43 that accumulates in cell

plasma. Using dog DM model may establish the link between spinal cord and serum/plasma

derived exosomes in terms of TDP-43 biochemistry. We have observed increased TDP-43

and SOD1 levels in exosomes derived from DM affected dog. SOD1 analysis was performed

for further confirming the spinal cord and serum exosome crosstalk in terms of the content

TDP-43 and its derivative.

The SOD1 protein profile revealed three SOD1 species (monomer, trimer, and tetramer),

although SOD1 resolved under the reducing conditions. Recent observations demonstrated

that trimer-SOD1 aggressively promoted cell death (42, 43). Because of their small size,

exosomes can easily pass the blood brain barrier to serve as accessible biomarkers of neuronal

dysfunction (7, 9, 44, 45). This pilot study provided some preliminary data support the notion

that spinal cord TDP-43 may be transported via exosomes, and that TDP-43 loaded exosomes

appear in blood. However, we did not perform spinal cord-derived exosomal TDP-43 and

SOD1 assessment. Hence, we do not claim yet serum-derived exosomal TDP-43 and SOD1

directly reflecting that of spinal cord. Studies are undergoing to further isolate spinal cord

derived exosomes appeared in serum and analyze their cargo content in terms of TDP-43 and

SOD1.

The concept of serum-derived exosome analysis for potential biomarker candidates will

eliminate very invasive tissue and cerebrospinal fluid sampling and allows investigators to

obtain repeated blood samples with a minimal invasive sampling method for exosome

isolation. Exosomes derived from serum represent global exosomes which include neuronal

derived exosomes (NDE) (46). A recent review has discussed on astrocyte-derived extracellular vesicles in the blood may be a useful tool for detecting astrocyte-specific biomarkers in neurological conditions (47, 48) Our current research is focusing on the development of a process, to selectively isolate the NDE population from global serum and spinal cord tissue derived exosomes.

The availability of a disease model that more closely reflects the human condition would provide an opportunity to test if the pathologic processes shown to contribute to the motor neuron disease in rodent models are likely to contribute to ALS. Such a model could also be used in second-generation preclinical trials to shield ALS patients from participation in clinical trials with little probability of success. In addition, studies with the proposed disease model may reveal disease processes that are shared with the ALS cases but do not occur, or are difficult to detect, in the rodents due to small size, short lifespan, or other issues. Ideally, the desired disease model would develop both upper and lower motor neuron signs due to a single spontaneous mutation in an orthologue of a human gene known to harbor ALS-causing mutations. This model would involve an animal species that is intermediate between rodents and humans in life expectancy and in the size and complexity of the CNS. It would be readily available on a wide variety of genetic backgrounds negating the need to maintain research colonies. It would display a disease phenotype that, similar to ALS, first manifests in mature individuals. In addition, the initial clinical signs would consistently appear at the same anatomic site and spread throughout the body in a uniform in pattern and rate. Canine DM has all these features and is therefore another disease model for understanding ALS. Companion dogs with DM represent a clinical population confounded by complexities in diagnosis, comorbidity and environmental and genetic diversity similar to those encountered in a human clinical trial setting. Thus, incorporation of veterinary clinical trials into the ALS

treatment development paradigm will enhance translational efficiency by identifying and

optimizing those therapies most likely to generate clinical benefit.

Acknowledgments: Authors acknowledge the Office for Research and Support Program at Kansas City University for supporting student research and providing logistic supports, Robert Liu, DO and Qwynton Johnson for providing technical help, Mrs. Jessica Sage for training summer research fellows, and Dr. Jan Talley and Ms. Tuba Agbas for reading and editing the manuscript.

Funding: Summer Research Fellowship Program (PP) provided by Kansas City University funded this research. Intramural grant to support (AA) for student researchers (AK ,MM, YME, and KG), and by Kansas City Consortium of Musculoskeletal Disorders (KCMD) (AA and JRC) under a pilot project grant (KCMD_FY21).

Disclosure of interest: The authors declare no conflict of interest.

ORCID

Abdulbaki Agbas orcid logo 0000-0003-3854-778

Author Contributions: Conceptualization, AA and JRC; methodology, AA, PP, AK,KG, YME, MM and EK.; formal analysis, PP,AK,KG,EK, and AA; writing, original draft preparation, AA; writing, review and editing, AA,MM, and JRC; supervision, AA and EK; All authors have read and agreed to the published version of the manuscript.

Data availability statement: The authors confirm that the data supporting the findings of this study are available within the article and its supplementary materials. The data are available on reasonable request from corresponding author (aagbas@kansascity.edu)

References

1.	Beyer L, Gunther R, Koch JC, Klebe S, Hagenacker T, Lingor P, et al. TDP-43 as structure-based biomarker in amyotrophic lateral sclerosis. Ann Clin Transl Neurol. 2021;8(1):271-7.
2.	Majumder V, Gregory JM, Barria MA, Green A, Pal S. TDP-43 as a potential biomarker for amyotrophic lateral sclerosis: a systematic review and meta-analysis. BMC Neurol. 2018;18(1):90.
3.	Iguchi Y, Eid L, Parent M, Soucy G, Bareil C, Riku Y, et al. Exosome secretion is a key pathway for clearance of pathological TDP-43. Brain : a journal of neurology. 2016;139(Pt 12):3187-201.

4.   Hosaka T, Yamashita T, Tamaoka A, Kwak S. Extracellular RNAs as Biomarkers of Sporadic Amyotrophic Lateral Sclerosis and Other Neurodegenerative Diseases. Int J Mol Sci. 2019;20(13).

5.   Suk TR, Rousseaux MWC. The role of TDP-43 mislocalization in amyotrophic lateral sclerosis. Molecular neurodegeneration. 2020;15(1):45.

6.   Pegtel DM, Gould SJ. Exosomes. Annu Rev Biochem. 2019;88:487-514.

7.   Saint-Pol J, Gosselet F, Duban-Deweer S, Pottiez G, Karamanos Y. Targeting and Crossing the Blood-Brain Barrier with Extracellular Vesicles. Cells. 2020;9(4).

8.   Morad G, Carman CV, Hagedorn EJ, Perlin JR, Zon LI, Mustafaoglu N, et al. Tumor-Derived Extracellular Vesicles Breach the Intact Blood-Brain Barrier via Transcytosis. ACS Nano. 2019;13(12):13853-65.

9.   Qu M, Lin Q, Huang L, Fu Y, Wang L, He S, et al. Dopamine-loaded blood exosomes targeted to brain for better treatment of Parkinson's disease. J Control Release. 2018;287:156-66.

10.   What is ALS? The ALS Association2014 [updated 02/16/2014. Available from: www.alsa.org.

11.   Wijesekera LC, Leigh PN. Amyotrophic lateral sclerosis. Orphanet J Rare Dis. 2009;4:3.

12.   Turner MR, Brockington A, Scaber J, Hollinger H, Marsden R, Shaw PJ, et al. Pattern of spread and prognosis in lower limb-onset ALS. Amyotrophic lateral sclerosis : official publication of the World Federation of Neurology Research Group on Motor Neuron Diseases. 2010;11(4):369-73.

13.   Masrori P, Van Damme P. Amyotrophic lateral sclerosis: a clinical review. Eur J Neurol. 2020;27(10):1918-29.

14.   Armon C. Amyotrophic lateral sclerosis Medscape2014 [Available from: <http://emedicine.medscape.com/article/1170097-overview>.

15.   Wegorzewska I, Baloh RH. TDP-43-based animal models of neurodegeneration: new insights into ALS pathology and pathophysiology. Neuro-degenerative diseases. 2011;8(4):262-74.

16.   Awano T, Johnson GS, Wade CM, Katz ML, Johnson GC, Taylor JF, et al. Genome-wide association analysis reveals a SOD1 mutation in canine degenerative myelopathy that resembles amyotrophic lateral sclerosis. Proc Natl Acad Sci U S A. 2009;106(8):2794-9.

17.   Holder AL, Price JA, Adams JP, Volk HA, Catchpole B. A retrospective study of the prevalence of the canine degenerative myelopathy associated superoxide dismutase 1 mutation (SOD1:c.118G > A) in a referral population of German Shepherd dogs from the UK. Canine genetics and epidemiology. 2014;1:10.

18.   Morgan BR, Coates JR, Johnson GC, Bujnak AC, Katz ML. Characterization of intercostal muscle pathology in canine degenerative myelopathy: a disease model for amyotrophic lateral sclerosis. Journal of neuroscience research. 2013;91(12):1639-50.

19.   Morgan BR, Coates JR, Johnson GC, Shelton GD, Katz ML. Characterization of thoracic motor and sensory neurons and spinal nerve roots in canine degenerative myelopathy, a potential disease model of amyotrophic lateral sclerosis. Journal of neuroscience research. 2014;92(4):531-41.

20.   Coates JR, Wininger FA. Canine degenerative myelopathy. Vet Clin North Am Small Anim Pract. 2010;40(5):929-50.

21.   Gordon PH, Cheng B, Katz IB, Mitsumoto H, Rowland LP. Clinical features that distinguish PLS, upper motor neuron-dominant ALS, and typical ALS. Neurology. 2009;72(22):1948-52.

22.   Sage J, Hall L, McVey A, Barohn RJ, Statland JM, Jawdat O, et al. Use of Capillary Electrophoresis Immunoassay to Serach for Potential Biomarkers of Amyotrophic Lateral Sclerosis in Human Platelets. JVisExp. 2020;156(e60638):1-9.

23.   Neumann M, Kwong LK, Lee EB, Kremmer E, Flatley A, Xu Y, et al. Phosphorylation of S409/410 of TDP-43 is a consistent feature in all sporadic and familial forms of TDP-43 proteinopathies. Acta neuropathologica. 2009;117(2):137-49.

24.   Neumann M, Sampathu DM, Kwong LK, Truax AC, Micsenyi MC, Chou TT, et al. Ubiquitinated TDP-43 in frontotemporal lobar degeneration and amyotrophic lateral sclerosis. Science. 2006;314(5796):130-3.

25.   Baloh RH. TDP-43: the relationship between protein aggregation and neurodegeneration in amyotrophic lateral sclerosis and frontotemporal lobar degeneration. The FEBS journal. 2011;278(19):3539-49.

26.   Jo M, Lee S, Jeon YM, Kim S, Kwon Y, Kim HJ. The role of TDP-43 propagation in neurodegenerative diseases: integrating insights from clinical and experimental studies. Exp Mol Med. 2020;52(10):1652-62.

27.   miRCURY Exosome kits handbook. In: Qiagen, editor. 2017. p. 7-8.

28.   Feneberg E, Charles PD, Finelli MJ, Scott C, Kessler BM, Fischer R, et al. Detection and quantification of novel C-terminal TDP-43 fragments in ALS-TDP. Brain Pathol. 2020:e12923.

29.   Katz ML, Jensen CA, Student JT, Johnson GC, Coates JR. Cervical spinal cord and motor unit pathology in a canine model of SOD1-associated amyotrophic lateral sclerosis. J Neurol Sci. 2017;378:193-203.

30.   Sun J, Carrero JJ, Zagai U, Evans M, Ingre C, Pawitan Y, et al. Blood biomarkers and prognosis of amyotrophic lateral sclerosis. Eur J Neurol. 2020;27(11):2125-33.

31.   Aydemir D, Ulusu NN. Importance of the serum biochemical parameters as potential biomarkers for rapid diagnosis and evaluating preclinical stage of ALS. Med Hypotheses. 2020;141:109736.

32.   Brodovitch A, Boucraut J, Delmont E, Parlanti A, Grapperon AM, Attarian S, et al. Combination of serum and CSF neurofilament-light and neuroinflammatory biomarkers to evaluate ALS. Sci Rep. 2021;11(1):703.

33.   Averill DR, Jr. Degenerative myelopathy in the aging German Shepherd dog: clinical and pathologic findings. J Am Vet Med Assoc. 1973;162(12):1045-51.

34.   Zeng R, Coates JR, Johnson GC, Hansen L, Awano T, Kolicheski A, et al. Breed distribution of SOD1 alleles previously associated with canine degenerative myelopathy. J Vet Intern Med. 2014;28(2):515-21.

35.   Rakhit R, Chakrabartty A. Structure, folding, and misfolding of Cu,Zn superoxide dismutase in amyotrophic lateral sclerosis. Biochim Biophys Acta. 2006;1762(11-12):1025-37.

36.   Forsberg K, Jonsson PA, Andersen PM, Bergemalm D, Graffmo KS, Hultdin M, et al. Novel antibodies reveal inclusions containing non-native SOD1 in sporadic ALS patients. PloS one. 2010;5(7):e11552.

37.   Bosco DA, Morfini G, Karabacak NM, Song Y, Gros-Louis F, Pasinelli P, et al. Wild-type and mutant SOD1 share an aberrant conformation and a common pathogenic pathway in ALS. Nat Neurosci. 2010;13(11):1396-403.

38.   Noto Y, Shibuya K, Sato Y, Kanai K, Misawa S, Sawai S, et al. Elevated CSF TDP-43 levels in amyotrophic lateral sclerosis: specificity, sensitivity, and a possible prognostic value. Amyotrophic lateral sclerosis : official publication of the World Federation of Neurology Research Group on Motor Neuron Diseases. 2011;12(2):140-3.

39.   Neumann M, Kwong LK, Truax AC, Vanmassenhove B, Kretzschmar HA, Van Deerlin VM, et al. TDP-43-positive white matter pathology in frontotemporal lobar

degeneration with ubiquitin-positive inclusions. Journal of neuropathology and experimental neurology. 2007;66(3):177-83.

40. Thery C, Witwer KW, Aikawa E, Alcaraz MJ, Anderson JD, Andriantsitohaina R, et al. Minimal information for studies of extracellular vesicles 2018 (MISEV2018): a position statement of the International Society for Extracellular Vesicles and update of the MISEV2014 guidelines. J Extracell Vesicles. 2018;7(1):1535750.

41. Song Z, Xu Y, Deng W, Zhang L, Zhu H, Yu P, et al. Brain Derived Exosomes Are a Double-Edged Sword in Alzheimer's Disease. Front Mol Neurosci. 2020;13:79.

42. Zhu C, Beck MV, Griffith JD, Deshmukh M, Dokholyan NV. Large SOD1 aggregates, unlike trimeric SOD1, do not impact cell viability in a model of amyotrophic lateral sclerosis. Proceedings of the National Academy of Sciences of the United States of America. 2018;115(18):4661-5.

43. Proctor EA, Fee L, Tao Y, Redler RL, Fay JM, Zhang Y, et al. Nonnative SOD1 trimer is toxic to motor neurons in a model of amyotrophic lateral sclerosis. Proceedings of the National Academy of Sciences of the United States of America. 2016;113(3):614-9.

44. Saeedi S, Israel S, Nagy C, Turecki G. The emerging role of exosomes in mental disorders. Transl Psychiatry. 2019;9(1):122.

45. Banks WA, Sharma P, Bullock KM, Hansen KM, Ludwig N, Whiteside TL. Transport of Extracellular Vesicles across the Blood-Brain Barrier: Brain Pharmacokinetics and Effects of Inflammation. Int J Mol Sci. 2020;21(12).

46. Malm T, Loppi S, Kanninen KM. Exosomes in Alzheimer's disease. Neurochem Int. 2016;97:193-9.

47. Upadhya R, Zingg W, Shetty S, Shetty AK. Astrocyte-derived extracellular vesicles: Neuroreparative properties and role in the pathogenesis of neurodegenerative disorders. J Control Release. 2020;323:225-39.

48. Long X, Yao X, Jiang Q, Yang Y, He X, Tian W, et al. Astrocyte-derived exosomes enriched with miR-873a-5p inhibit neuroinflammation via microglia phenotype modulation after traumatic brain injury. J Neuroinflammation. 2020;17(1):89.

2 **FIGURES**

13 **Figure-1**

23 **Figure-2**

11 **Figure-3**

22 **Figure-4**

Figure-1. Total TDP-43 levels of thoracic region of spinal cord homogenate. TDP-43 levels were elevated in the spinal cord homogenate from DM affected dogs (n=4, Supplemental Data / Table-1: dog # 1,2,3,4) as compared to that of the control dogs (n=4, Supplemental Data / Table-1: dog # 5,6,7,8) [One- sample t-test ($P<0.01$)]. The protein band intensity were normalized to total protein staining.

Figure-2. SOD1 levels of thoracic region of spinal cord homogenate. SOD1 levels were elevated in the spinal cord homogenate from DM affected dogs (n=4, Supplemental Data / Table-1: dog # 1,2,3,4) as compare to that of the control dogs (n=4, Supplemental Data / Table-1: dog # 5,6,7,8) [One-sample t-test ($P< 0.01$)]. The protein band intensities were normalized to total protein staining.

Figure 3. Serum derived exosomal total TDP-43 and phosphorylated TDP-43 profile in DM affected dogs. Both total TDP-43 (A) and phosphorylated TDP-43 (pTDP-43) (B) levels were elevated in serum-derived exosomes from DM affected dogs (n=12; Table-2 Supplemental Data: dog # 1,2,3,4,5,6,7,8,9,10,11,12) compared to control dogs (n=10; Table-3 Supplemental Data: dog# 1,2,3,4,5,6,7,8,9,10). One-sample t-test analysis revealed statistical significance for total TDP-43 ($P<0.0001$) and for pTDP-43 ($P<0.001$) Original immunoblot analysis pertinent to this figure provided in Supplemental Fig.1 and Fig.2. The protein band intensities were normalized to total protein staining.

Figure 4. Serum derived exosomal SOD1 profile in dogs with DM; (A) Exosomal SOD1 levels appear to be elevated in dogs with DM. The difference between control (n= 3, Table-3 Supplemental Data ; dog #1,2,4,5) and DM (n=4; Table-2 Supplemental Data: dog #1,2,3,4,6) groups did achieve statistically significance ($p <0.05$). (B) Monomer SOD1 proteins bands were analysed and graphed. The protein band intensities were normalized to total protein staining.