Genomic islands targeting *dusA* in *Vibrio* species are distantly

related to *Salmonella* Genomic Island 1 and mobilizable by IncC

conjugative plasmids

Romain Durand[1,#a], Florence Deschênes[1] and Vincent Burrus[1]*

[1] Département de biologie, Université de Sherbrooke, Sherbrooke, Québec,

Canada

[#a]Current Address: Institut de Biologie Intégrative et des Systèmes, Université

Laval, Québec, Canada

* Corresponding author

E-mail: vincent.burrus@usherbrooke.ca

Short title: *dusA*-specific genomic islands mobilizable by IncC conjugative

plasmids

Abstract

Salmonella Genomic Island 1 (SGI1) and its variants are significant contributors

to the spread of antibiotic resistance among *Gammaproteobacteria*. All known

SGI1 variants integrate at the 3' end of *trmE*, a gene coding for a tRNA

modification enzyme. SGI1 variants are mobilized specifically by conjugative

plasmids of the incompatibility groups A and C (IncA and IncC). Using a

comparative genomics approach based on genes conserved among members of

the SGI1 group, we identified diverse integrative elements distantly related to

SGI1 in several species of *Vibrio*, *Aeromonas*, *Salmonella*, *Pokkaliibacter*, and

Escherichia. Unlike SGI1, these elements target two alternative chromosomal

loci, the 5' end of *dusA* and the 3' end of *yicC*. Although they share many

features with SGI1, they lack antibiotic resistance genes and carry alternative

integration/excision modules. Functional characterization of IME*Vch*USA3, a

dusA-specific integrative element, revealed promoters that respond to AcaCD,

the master activator of IncC plasmid transfer genes. Quantitative PCR and

mating assays confirmed that IME*Vch*USA3 excises from the chromosome and is

mobilized by an IncC helper plasmid from *Vibrio cholerae* to *Escherichia coli*.

IME*Vch*USA3 encodes the AcaC homolog SgaC that associates with AcaD to

form a hybrid activator complex AcaD/SgaC essential for its excision and

mobilization. We identified the *dusA*-specific recombination directionality factor

RdfN required for the integrase-mediated excision of *dusA*-specific elements

from the chromosome. Like *xis* in SGI1, *rdfN* is under the control of an AcaCD-

responsive promoter. Although the integration of IME*Vch*USA3 disrupts *dusA*, it

provides a new promoter sequence and restores the reading frame of *dusA* for

proper expression of the tRNA-dihydrouridine synthase A. Phylogenetic analysis

of the conserved proteins encoded by SGI1-like elements targeting *dusA*, *yicC*,

and *trmE* gives a fresh perspective on the possible origin of SGI1 and its

variants.

Author summary

We identified integrative elements distantly related to the *Salmonella* Genomic

Island 1 (SGI1), a key vector of antibiotic resistance genes in

Gammaproteobacteria. SGI1 and its variants reside at the 3' end of *trmE*, share a

large, highly conserved core of genes, and carry a complex integron that confers

multidrug resistance phenotypes to their hosts. Unlike members of the SGI1

group, these novel genomic islands target the 5' end *dusA* or the 3' end of *yicC*,

lack multidrug resistance genes, and seem much more diverse. We showed here

that, like SGI1, these elements are mobilized by conjugative plasmids of the IncC

group. Based on comparative genomics and functional analyses, we propose a

hypothetical model of the evolution of SGI1 and its siblings from the progenitor of

IncA and IncC conjugative plasmids via an intermediate *dusA*-specific integrative

element through gene losses and gain of alternative integration/excision

modules.

Introduction

Integrative and mobilizable elements (IMEs) are discrete, mobile chromosomal regions that can excise from the chromosome and borrow the mating apparatus of helper conjugative elements to transfer to a new bacterial host [1,2]. IMEs are usually composed of two main functional modules. The site-specific recombination module contains genes and *cis*-acting sequences that mediate the integration of the IMEs into and their excision from the chromosome. The mobilization module includes the *cis*-acting origin of transfer (*oriT*) and usually encodes mobilization proteins required to initiate the conjugative transfer at *oriT* [1]. In its simplest form, the mobilization module only consists of an *oriT* locus mimicking the *oriT* of the helper element [3–5]. The excision of IMEs is elicited by conjugative plasmids or integrative and conjugative elements (ICEs). These helper elements encode the type IV secretion system (T4SS) that translocates the IME DNA into the recipient cell [1].

Several distinct families of IMEs have been described to date. Most encode beneficial traits for their host, such as resistance to antibiotics and heavy metals or bacteriocin synthesis [1,6]. *Salmonella* Genomic Island 1 (SGI1) is certainly one of the most studied IMEs. Though first described 20 years ago, SGI1 and its siblings have only recently gained a lot of attention due to their prevalence and prominent role in the spread of multidrug resistance [7,8]. The canonical 43-kb SGI1 resides at the 3' end of *trmE* (also known as *mnmE* or *thdF*) in *Salmonella enterica* serovar Typhimurium DT104 [9]. *trmE* encodes the 5-carboxymethylaminomethyluridine-tRNA synthase GTPase subunit. SGI1

variants have been reported in a wide array of *Gammaproteobacteria*, including

Proteus mirabilis (PGI1), *Acinetobacter baumannii* (AGI1), *Morganella*,

Providencia, *Enterobacter*, *Escherichia coli*, *Vibrio cholerae* (GI-15), and

Klebsiella pneumoniae [7,10,11]. Most variants carry a class I integron

structurally similar to the In104 integron of SGI1. In104 confers resistance to

ampicillin, chloramphenicol/florfenicol, streptomycin/spectinomycin,

sulfamethoxazole, and tetracycline [8,12]. SGI1 and its variants are an

epidemiological threat exacerbated by their specific mobilization by conjugative

plasmids of the incompatibility groups A (IncA) and C (IncC) [13,14]. IncC

plasmids contribute to the global circulation of multidrug resistance genes,

including NDM metallo-β-lactamase and carbapenemase genes, among a broad

range of *Gammaproteobacteria* [15,16]. The transcriptional activator AcaCD

encoded by IncC plasmids triggers the excision and mobilization of SGI1 [17,18].

SGI1 and most variants share a conserved core set of 28 genes, representing

27.4 kb, disrupted by insertion sequences and the class 1 integron inserted at

diverse positions (Fig 1, top) [7,9,12]. Thus far, the function of a few conserved

genes has been characterized. Together with the *cis*-acting recombination

site *attP,* the genes *int* and *xis* form the recombination module of SGI1

[13]. *int* encodes the site-specific tyrosine recombinase (integrase) that targets

the 3' end of *trmE. xis* encodes the recombination directionality factor (RDF or

excisionase) that enhances the excision reaction catalyzed by Int. The

mobilization module includes the mobilization genes *mpsAB* and the *oriT* located

upstream of *mpsA* [19]. *mpsA* encodes an atypical relaxase distantly related to

tyrosine recombinases. Unlike most characterized IMEs, SGI1 carries a replicon

composed of an iteron-based origin of replication (*oriV*) and the replication

initiator gene *rep* [20,21]. SgaCD, a transcriptional activator complex expressed

by SGI1 in response to a coresident IncC plasmid, controls *rep* expression

[21,22]. The excised replicative form of SGI1 destabilizes the helper plasmid by

an unknown process, and is further stabilized by its *sgiAT* addiction module

[20,22–24]. Finally, SGI1 encodes three mating pore subunits, TraN$_S$, TraH$_S$, and

TraG$_S$, that actively replace their counterparts in the T4SS encoded by the IncC

plasmid [25]. The substitution of TraG allows SGI1 to bypass the IncC-encoded

entry exclusion mechanism and transfer between cells carrying conjugative

plasmids belonging to the same entry exclusion group [26].

Given the high similarity between SGI1 variants integrated at *trmE*, we undertook

a search for distant SGI1-like IMEs in bacterial genomes using MpsA, TraG$_S$,

SgaC, and TraN$_S$ as baits. Here, we report the existence of distantly related IMEs

integrated at the 5' end of *dusA* in several species of *Vibrionaceae* and the 3' end

of *yicC* in several species of *Gammaproteobacteria*. We have examined the

interactions between an IncC plasmid and IME*Vch*USA3, a *dusA*-specific

representative IME from an environmental *V. cholerae* strain. The genetic

determinants required for the excision of IME*Vch*USA3 and its mobilization by

IncC plasmids were characterized. Finally, we took a fresh look at the emergence

and evolution of SGI1 and its siblings by conducting phylogenetic analyses and

proposed a hypothetical evolutionary pathway of putative IMEs resembling SGI1.

Results

Novel integrative elements (IEs) distantly related to SGI1 are inserted in

dusA and _yicC_ in various _Gammaproteobacteria_

To find novel SGI1-like elements, we searched the Refseq database using blastp

and the primary sequences of MpsA, TraG$_S$, SgaC, and TraN$_S$. Considering the

substitution of integration modules can change the integration site [27–29], the

integrase Int$_{trmE}$ was excluded from the analysis. We identified 24 distinct

integrative elements encoding homologs of the four bait proteins in 36 different

bacterial strains (Fig 1, Table 1 and S1 Table). 21 of these IEs are integrated into

the 5' end of _dusA_ in diverse _Vibrio_ species from various origins. The remaining

three are located at the 3' end of _yicC_ in _E. coli_, _Aeromonas veronii_, _P. mirabilis_,

S. enterica serovar Kentucky, and _Pokkaliibacter plantistimulans_. The size of the

IEs varies from 22.8 kb to 37.1 kb. The conserved genes _mpsA_ (together with

mpsB), _traG_, _traN_, and _sgaC_ remain in a syntenic order, though sporadically

separated by variable DNA (Fig 1).

Figure 1. Schematic representations of SGI1-related IEs. The position and
orientation of open reading frames (ORFs) are indicated by arrowed boxes.
Colors depict the function deduced from functional analyses and BLAST
comparisons. Potential AcaCD binding sites are represented by green angled
arrows. Each island is flanked by the _attL_ and _attR_ (vertical grey lines)
attachment sites when integrated into the 3' end of _trmE_ (light blue), the 5' end of
dusA (light green), or the 3' end of _yicC_ (pink). The annotation of _attL_ and _attR_
relative to _int_ is based on SGI1 (_trmE_) [9], IE_Aba_D1279779 of _Acinetobacter_

baumannii D1279779 (*dusA*) [30] and MGI*Vfl*Ind1 (*yicC*) [3]. Details regarding

ORFs are shown in S1 Dataset.

Consistent with the change of integration site, the respective *int* genes of SGI1

and the *dusA*- and *yicC*-specific IEs do not share any sequence similarity.

Furthermore, unlike SGI1, these novel IEs lack *xis* downstream of *int* (Fig 1).

Instead, *yicC*-specific IEs carry two small open reading frames (ORF) upstream

of the *attR* site. The putative translation product of the second one shares 35%

identity over 65% coverage with the excisionase RdfM of MGI*Vfl*Ind1 [31].

Although *dusA*-specific IEs lack *xis* and *rdfM*, all carry an ORF predicted to

encode a 76-aminoacyl residue protein containing the pyocin activator protein

PrtN domain (Pfam PF11112). Based on its size, position, predicted DNA-binding

function, conservation, and evidence presented below, we named this ORF *rdfN*.

Phylogenetic analysis of Int$_{yicC}$ proteins of *yicC*-specific SGI1-like IEs form a

cluster distinct from the integrases of IMEs mobilizable by IncC plasmids through

a MobI protein (Pfam PF19456), such as MGI*Vmi*1, and IMEs that mimic the *oriT*

of SXT/R391 ICEs, such as MGI*Vfl*Ind1 [3,17,32] (Fig 2A).

Figure 2. Integrases encoded by the *yicC*- and *dusA*-specific IEs. Maximum

likelihood phylogenetic analyses of Int$_{yicC}$ (A) and Int$_{dusA}$ (B). The trees are drawn

to scale, with branch lengths measured in the number of substitutions per site

over 400 and 359 amino acid positions for Int$_{yicC}$, and Int$_{dusA}$, respectively. The

helper elements and mechanism of mobilization are indicated for each lineage

according to the keys shown in the legend box of panel A. The inset of panel B

shows logo sequences of the repeats in *attL* and *attR* attachment sites. The

arrows indicate the island termini experimentally determined for IE*Aba*D1279779

by Farrugia *et al.* [30]. (C) Heatmap showing blastp identity percentages of

pairwise comparison of Int$_{dusA}$ representative proteins. Proteins accession

numbers are provided in S2 Dataset, except for IE*Aba*D1279779

(WP_000534871.1), IE*Ppr*Pf-5 of *Pseudomonas protegens* Pf-5

(WP_011060295.1), and IEs of *Burkholderia gladioli* BSR3 (WP_013697845.1),

Bradyrhizobium sp. BTAi1 (WP_012043559.1), *Agrobacterium* sp. H13-3

(WP_013636109.1), and *Neisseria gonorrhoeae* FA 1090 (EFF39980.1).

Phylogenetic analysis of Int$_{dusA}$ proteins confirmed that the integrases of these

IEs form a monophyletic group exclusive to the *Vibrionaceae* and distinct from

those encoded by other *dusA*-specific IEs found in other taxa, including

GI*Aca*Bra1 from *Aeromonas caviae* that is likely mobilizable by IncC plasmids via

a MobI protein [32] (Fig 2B). Int$_{dusA}$ proteins of the IEs identified here share at

least 75% identity, while identities drop below 60% with the non-*Vibrio* Int$_{dusA}$

proteins (Fig 2C). Sequence logos built using alignments of the *attL* and *attR*

chromosomal junctions revealed a 21-bp imperfect repeat at the extremities of

each IE (Fig 2B). This repeat is similar to the one reported for *dusA*-specific IEs

found in a broader range of species [30].

Three types of *dusA*-integrated SGI1-related elements

Blastn and blastp analyses using SGI1ΔIn104 as the reference confirmed that

the identified *dusA*-specific IEs share limited sequence similarities with SGI1

(S1A Fig). Besides the genes encoding MpsA, TraG, SgaC, and TraN, all carry

the auxiliary mobilization factor gene *mpsB* and the *oriT* sequence (Fig 1).

Secondary structure prediction of the aligned *oriT* sequences located upstream

of *mpsA* using RNAalifold revealed that despite the sequence divergence, the

structure of *oriT* with three stem-loops was strictly conserved (S2B Fig). In

contrast, *sgaD* is not strictly conserved and highly divergent from *sgaD* of SGI1

when present (Fig 1 and S1A Fig).

Comparison using IE*Vch*USA2 as the reference suggests that *dusA*-specific IEs

cluster into three distinct types as confirmed by the phylogenetic analysis of

concatenated MpsA-TraG-SgaC-TraN (Fig 3A, S1B and S3 Fig). Type 1 *dusA*-

specific IEs such as IME*Vch*USA3 are mainly found in *V. cholerae* and lack both

traH and *sgaD*. Type 2 IEs such as IE*Vch*USA2 lack *sgaD* but carry *traH* (Fig 1

and 3A). This lineage only includes two *dusA*-specific IEs of *V. cholerae* but also

closely related *yicC*-specific IEs such as IE*Eco*MOD1 and the *trmE*-specific

GI*Vch*O27-1. Finally, type 3 IEs such as GI*Vch*USA5 are the most distant from

the two other types and SGI1 (Fig 3A). Type 3 IEs carry both *traH* and *sgaD* and

reside in diverse *Vibrio* species. With the exception of a few outliers encoded by

IEs such as IE*Vch*N2817, IE*Vch*N2708 or IE*Ppl*Ind1, the proteins MpsA, TraG,

SgaC and TraN encoded by members of the same type typically share more than

95% identity (Fig 3B and S3). MpsA remains the least divergent protein between

the three types, sharing at least 65% identity between type 1 and type 3, and

from 64% to 93% with SGI1. In contrast, TraG and TraN are the most divergent

between types, ranging from 46% to 59% for TraG and from 46% to 76% for

TraN.

Worthy of note, these three distinct lineages of *dusA*-specific IEs are supported

by the phylogeny of the *oriT* sequences (S2A Fig). Again, *oriT* loci of type 3 IEs

strongly diverge from those of types 1 and 2, as well as from the *oriT* loci of the

highly homogenous SGI1 group.

Figure 3. Conserved genes support three main lineages of *dusA*-specific

SGI1-like IEs. (A) Maximum likelihood phylogenetic analysis of concatenated

MpsA-TraG-SgaC-TraN. The tree is drawn to scale, with branch lengths

measured in the number of substitutions per site over 2,637 amino acid positions.

Taxa corresponding to IEs targeting *trmE* and *yicC* are indicated by a light blue

circle and a red circle, respectively. All other taxa correspond to *dusA*-specific

IEs. Phylogenetic relationships of MpsA, TraG, SgaC and TraN proteins are

shown separately in S2 Fig. (B) Heatmaps showing blastp identity percentages of

pairwise protein comparisons for representatives of MpsA, TraG, SgaC, and

TraN. Proteins accession numbers and clusters are provided in S1 Table and S2

Dataset.

Variable features found in the *dusA*- and *yicC*-specific IEs

Most variable genes in the identified IEs encode proteins of unknown function. A

search for antibiotic resistance determinants using the Resistance Gene Identifier

server failed to reveal any known resistance gene. Several IEs encode putative

functions altering host processes and virulence, including the transport of ions

and small molecules (*ktrAB*, *trkH*, and *kdpD* for potassium uptake and *rcnAR* for

nickel/cobalt efflux in IE*Vch*Hai10, sulfite export in IE*Vch*N2817 and IE*Vch*Swe1),

c-di-GMP degradation (IE*Vch*Ban1), and fimbriae (IE*Vch*Ban1) (S1 Dataset).

None of the reported IEs carries the same replication module (*S004-rep-oriV*) as

canonical SGI1. Instead, five *dusA*-specific IEs belonging to the type 3 lineage

(IE*Vch*USA5, IE*Vch*Bra2, IE*Vpa*Chn1, IE*Vpa*Chn2, and IE*Vpa*Ban1a) encode a

putative replication initiator protein with the IncFII_repA domain (Pfam PF02387)

(Fig 1, S1 Dataset). IE*Vvu*USA1 encodes a putative helicase with an UvrD_C_2

domain (Pfam PF13538), whereas IE*Vch*Hai10, IE*Vch*USA2 and IE*Vch*Swe1

encode a putative ATP-dependent helicase (PcrA) and a putative ATP-

dependent endonuclease (YbjD). In addition, IE*Vch*N2786, IE*Sen*USA1 and

IE*Eco*MOD1 encode a predicted DEAD/DEAH box helicase (Pfam PF00270 and

PF00271). The three *yicC*-specific IEs encode a homolog of TrfA (Pfam

PF07042), the replication initiator protein of broad-host-range IncP plasmids [33].

No replicative functions could be ascribed with confidence to any gene carried by

the other *dusA*-specific IEs. Several IEs also encode toxin-antitoxin systems,

such as *sgiAT* and *higAB*, which likely enhance their stability (Fig 1). In the type 3

IEs IE*Vch*Bra2, IE*Vch*N2708, IE*Vpa*Chn1 and IE*Vpa*Ban1a, *sgiAT* is associated

with a gene coding for a putative abortive infection bacteriophage resistance

factor (Abi_2, Pfam PF07751). Likewise, IE*Vvu*USA1 carries a gene coding for a

different putative abortive infection bacteriophage resistance factor (AbiEii toxin,

Pfam PF13304).

Finally, IE*Vpa*Ban1a is integrated at *dusA* adjacent to a distinct IE, IE*Vpa*Ban1b,

in a tandem fashion. GI*Vpa*Ban1b codes for two predicted integrases sharing

44% and 27% identity with Int$_{dusA}$ of IE*Vpa*Ban1a. GI*Vpa*Ban1b encodes a

putative type I restriction-modification system, a MobA-like relaxase (MOB_{P1}), the

mobilization auxiliary factor MobC, and an RdfN homolog (Fig 1).

Non-canonical SGI1-like IEs carry AcaCD-responsive genes

Considering the divergence of the 24 new IEs from prototypical SGI1, we

wondered whether an IncC plasmid could mobilize them like SGI1. The hallmark

of IncC-dependent mobilization is the presence of AcaCD-responsive promoters

in IncC-mobilizable IEs. Hence, we searched for putative AcaCD-binding sites in

the sequences of *trmE*-specific IEs (prototypical SGI1 was used as the positive

control) and the *yicC-* and *dusA*-specific IEs. In these IEs, an AcaCD-binding

motif was predicted upstream of *traN*, *traHG* (or *traG*), *S018*, and *xis* (or *rdfM* or

rdfN) (Fig 1 and S4 Fig). Moreover, an AcaCD-binding motif was also predicted

upstream of *trfA* in the *yicC*-specific IEs.

We cloned the promoter sequences of *int*, *traN*, *traG*, *S018*, and *rdfN* of

IME*Vch*USA3 upstream of a promoterless *lacZ* reporter gene and monitored the

β-galactosidase activity with or without AcaCD. The promoter P_{int} was active

regardless of the presence of AcaCD (Fig 4A). In contrast, the four other

promoters exhibited weak activity in the absence of AcaCD. Upon induction of

acaDC expression, P_{traN} and P_{S018} remained unresponsive, while the activities of

P_{traG} and P_{rdfN} increased 40 and 400 times, respectively (Fig 4B). The inertia of

P_{traN} and P_{S018} toward AcaCD could result from single nucleotide substitutions in

the AcaCD binding site previously shown to be essential for recruiting the

activator [22]: CCS**A**AAWW instead of CCS**C**AAWW in P_{traN} and CCC**C**AAAA instead of

CCC**A**AAAA in P_{S018} (S4 Fig).

Figure 4. β-galactosidase activities of the promoters P_{int}, P_{traN}, P_{traG}, P_{S018}

and P_{rdfN} of IMEVchUSA3 transcriptionally fused to $lacZ$. (A) Colonies were

grown on LB agar with or without arabinose to induce *acaDC* expression from

pBAD-*acaDC*. (B) Induction levels of the same promoters in response to AcaCD.

β-galactosidase assays were carried out using the strains of panel A. Ratios

between the enzymatic activities in Miller units for the arabinose-induced versus

non-induced strains containing pBAD-*acaDC* are shown. The bars represent the

mean and standard error of the mean of three independent experiments.

Hence, despite their divergence and different integration sites, these IEs share

with SGI1 a common activation mechanism elicited by the presence of an IncC

plasmid.

IncC plasmids induce the excision and mobilization of IMEVchUSA3

Next, we tested whether a coresident IncC plasmid could trigger the excision of

IMEVchUSA3 from *dusA* in its original host, *V. cholerae* OY6PG08. The

derepressed IncC plasmid pVCR94^{Kn} Δ*acr2* [34] was introduced into OY6PG08

by conjugation from *E. coli* KH40. The Δ*acr2* mutation improves the efficiency of

interspecific transfer of the plasmid [35]. OY6PG08 Kn^R transconjugants were

tested by PCR to amplify the *attL* and *attR* chromosomal junctions, as well as the

attB and *attP* sites resulting from the excision of IMEVchUSA3 (S5A Fig).

IMEVchUSA3 was rarely retained in the transconjugants compared to the control

IncC-free OY6PG08 clones, suggesting it was unstable and rapidly lost in IncC$^+$

cells (S5B and S5C Figs).

To test the interspecific mobilization of IME*Vch*USA3 from *V. cholerae*

OY6PG08, we inserted a selection marker upstream of *traG* and used pVCR94^{Kn}

Δ*acr2* as the helper plasmid. IME*Vch*USA3^{Cm} transferred to *E. coli* CAG18439 at

a frequency of 7.01×10^{-5} transconjugant/donor CFUs. Amplification of *attL* and

attR using *E. coli*-specific primers confirmed that IME*Vch*USA3 integrates at

dusA in *E. coli* (S5D Fig).

Excision of *dusA*-specific IEs depends on *rdfN*

To further characterize the biology of IME*Vch*USA3, we measured its excision

rate and copy number by qPCR, with and without coresident pVCR94^{Sp}. We also

monitored its intraspecific transfer (*E. coli* to *E. coli*) in the same context.

Spontaneous excision of the island rarely occurred (<0.001% of the cells) (Fig

5A). In contrast, in the presence of the helper plasmid, the free *attB* site was

detected in more than 67% of the cells confirming that the IncC plasmid elicits

the excision of IME*Vch*USA3^{Kn}. Likewise, the presence of the plasmid resulted in

a ~3-fold increase of the copy number of IME*Vch*USA3^{Kn} (Fig 5B), suggesting

that the excised form of the island undergoes replication. The frequency of

transfer of IME*Vch*USA3^{Kn} was comparable to that of the helper plasmid

(~3.5×10^{-2} transconjugants/donor), while the frequency of cotransfer was more

than two logs lower (Fig 5C).

Figure 5. Effect of *acaDC* and *rdfN* on the IncC-dependent excision and

mobilization of IME*Vch*USA3. (A) IME*Vch*USA3^{Kn} excision rate corresponds to

the *attB*/chromosome ratio. (B) IME*Vch*USA3^{Kn} copy number corresponds to the

higA/chromosome ratio. For panels A and B, all ratios were normalized using the

control set to 1 and displayed in white. (C) Impact of *acaC*, *acaDC*, *sgaC* and

rdfN deletions on the mobilization of IME*Vch*USA3. Conjugation assays were

performed with CAG18439 (Tc) containing the specified elements as donor

strains and VB112 (Rf) as the recipient strain. The bars represent the mean and

standard error of the mean obtained from a biological triplicate. ¤ indicates that

the excision rate or transfer frequency was below the detection limit. Statistical

analyses were performed (on the logarithm of the values for panels A and C)

using a one-way ANOVA with Dunnett's multiple comparison test. For panels A

and B, statistical significance indicates comparisons to the normalization control.

Statistical significance is indicated as follows: ****, $P < 0.0001$; ***, $P < 0.001$; **,

$P < 0.01$; *, $P < 0.05$; ns, not significant. (D) Schematic representation of mini-IE

inserted at the 5' end of *dusA*. (E) RdfN acts as a recombination directionality

factor. Detection of *attB*, *attP*, *attL* and *attR* sites by PCR in colonies of *E. coli*

EC100 *dusA*::mini-IE in the presence or absence of *rdfN*. L, 1Kb Plus DNA

ladder (Transgen Biotech).

Thus far, the factors required to catalyze the excision of *dusA*-specific IEs have

not been examined [30]. Whereas all *dusA*-specific IEs lack *xis* downstream of

int, they carry a small ORF, here named *rdfN*, coding for a putative PrtN homolog

(Fig 1) [30]. The deletion of *rdfN* abolished the excision and replication of

IME*Vch*USA3[Kn]. Complementation by ectopic expression of *rdfN* from the

arabinose-inducible promoter P_{BAD} restored the wild-type excision level but not

the replication (Fig 5A and 5B). Likewise, deletion of *rdfN* abolished the

mobilization of IMEVchUSA3^{Kn} but had no impact on the transfer of the helper

plasmid (Fig 5C), confirming the specific role of $rdfN$ in the IE's mobility.

To confirm that $rdfN$ encodes the sole and only RDF of IMEVchUSA3, we

constructed mini-IE, a minimal version of IMEVchUSA3 that only contains int and

a spectinomycin-resistance marker. mini-IE is flanked by $attL$ and $attR$ and is

integrated at $dusA$ in $E.\ coli$ EC100 (Fig 5D). Using mini-IE, $attB$ and $attP$ were

detected only upon ectopic expression of $rdfN$ from pBAD-$rdfN$, confirming that

no other IMEVchUSA3-encoded protein besides Int and RdfN is required for the

excision of the element (Fig 5E). $rdfN$ is the essential RDF gene that favors the

excision of IMEVchUSA3 and, most likely, all $dusA$-specific IEs.

A SgaC/AcaD hybrid complex activates the excision and mobilization of

IMEVchUSA3

Next, we investigated the role of the transcriptional activator genes $acaC$ and

$sgaC$ in the mobilization of IMEVchUSA3. Deletion of $acaDC$ abolished the

excision and replication of IMEVchUSA3^{Kn}, confirming that its excision relies on

$rdfN$, whose expression is activated by AcaCD (Figs 4, 5A and 5B). The mutation

also confirmed that SgaC provided by IMEVchUSA3^{Kn} is insufficient by itself to

elicit $rdfN$ expression. The excision rate remained extremely low in cells that lack

the helper plasmid or cells that carry pVCR94^{Sp} $\Delta acaDC$. However,

IMEVchUSA3^{Kn} allowed the low-frequency transfer of pVCR94^{Sp} $\Delta acaDC$ [17]

(Fig 5C). Hence SgaC alone can activate to some degree the expression of the

transfer genes of the helper plasmid. In contrast, deletion of $acaC$ had no

significant impact on the excision, replication, and mobilization of

IME*Vch*USA3^{Kn}, or on the transfer of the helper plasmid (Figs 5A, 5B and 5C).

The primary sequences of AcaC and SgaC from IME*Vch*USA3 share 85%

identity over 94% coverage, whereas AcaC and SgaC from SGI1 share only 75%

identity over 92% coverage. Hence AcaD produced by the plasmid and SgaC

produced by the IME likely generate a functional chimeric transcriptional complex

that acts as a potent activator of *rdfN* and the transfer genes.

The transfer of IME*Vch*USA3^{Kn} Δ*sgaC* decreased nearly 3 logs compared to the

wild-type IE, despite the presence of *acaDC* on the helper plasmid (Fig 5C).

Moreover, deletion of both *acaC* and *sgaC* nearly abolished all transfer. Ectopic

expression of *sgaC* alone from pBAD-*sgaC* complemented these deletions to

wild-type levels (Fig 5C). These observations confirm that *sgaC*, not *acaC*,

combined with *acaD* produces a hybrid activator complex that is essential for the

excision and mobilization of the element.

IME*Vch*USA3 provides a new promoter and N-terminus for *dusA*

expression

Since *dusA*-specific IEs insert within the 5' end of *dusA*, we wondered whether

the gene remains expressed after the integration event. Sequence analysis of the

attR junction of *E. coli* K12 transconjugants revealed that IME*Vch*USA3 provides

a new 5' coding sequence that diverges significantly from the native *E. coli dusA*

gene (Fig 6A). This alteration of the 5' end of *dusA* results in a novel N-terminus

of identical length sharing 61% identity over the 35 initial amino acid residues

with native DusA. To test the expression of *dusA*, we constructed a translational

lacZ fusion to its fortieth codon downstream of the *attR* junction in *E. coli*

CAG18439 and BW25113 (Fig 6B). β-galactosidase assays revealed that *dusA*

remains expressed after integration in both strains, confirming that IME*Vch*USA3

provides a new promoter (Fig 6C). However, we observed a statistically

significant reduction of *dusA* expression resulting from the integration of the IE in

both strains, suggesting that the transcription or translation signals brought by

the IE are weaker than the original ones upstream of *E. coli dusA*.

Figure 6. IME*Vch*USA3 drives the expression of *dusA*. (A) Comparison of the

coding sequences of the 5' end of *dusA* in *E. coli* K12 MG1655 before (*attB* site)

and after (*attL* junction) the integration of IME*Vch*USA3. The core sequence of

the *attB* and *attL* recombination sites is indicated with red shading. The ATG start

codon of *dusA* is shown in bold. The sequence shown in blue is internal to

IME*Vch*USA3. Amino acid residues shown in red differ from the native N-

terminus of DusA. This sequence was obtained by sequencing the *attL* junction

of an *E. coli* CAG18439 *dusA*::IME*Vch*USA3^{Kn} transconjugant colony. (B)

Schematic representation of the *dusA*'-*lacZ* translational fusion for the detection

of *dusA* expression. (C) β-galactosidase activity of the *dusA*'-'*lacZ* fusion before

(-) and after (+) insertion of IME*Vch*USA3^{Kn} in *E. coli* CAG18439 (FD034) and

BW25113 (FD036). The bars represent the mean and standard error of the mean

of three independent experiments. Statistical analyses were performed using an

unpaired *t* test to compare the expression before and after integration of

IME*Vch*USA3^{Kn} for each strain. Statistical significance is indicated as follows: **,

P < 0.01; *, *P* < 0.05.

Discussion

SGI1-like elements integrated at the 3' end of *trmE* are widespread in a broad

range of *Enterobacteriaceae* and sporadically found in a few *Vibrio* species [7].

The integrase of SGI1 and its variants occasionally targets the intergenic region

between *sodB* and *purR* genes, a secondary attachment site [36]. Here, we

report the identification of distant SGI1-like elements that specifically target the 5'

end of *dusA* in multiple *Vibrio* species and the 3' end of *yicC* in

Enterobacteriaceae and *Balneatrichaceae*. Farrugia *et al.* [30] already described

IEs integrated at the 5' end of *dusA*, mostly prophages or phage remnants found

exclusively in *Alpha-*, *Beta-* and *Gammaproteobacteria*. These authors identified

IE*Vch*Ban1a and IE*Vch*Bra2 in *V. cholerae*, and several other IEs predicted to

encode conjugative functions in *Bradyrhizobium*, *Caulobacter*, *Mesorhizobium*,

Paracoccus, *Pseudomonas*, and *Rhodomicrobium* [30]. Our group recently

reported a *dusA*-specific IE in *Aeromonas caviae* 8LM potentially mobilizable by

IncC plasmids [32]. GI*Aca*8LM lacks *tra* genes but encodes a mobilization protein

(MobI) under the control of an AcaCD-responsive promoter. Together, these

reports confirm that *dusA* is an insertion hotspot for distinct families of mobile

elements across at least three *Proteobacteria* phyla.

Thus far, only the *dusA*-specific IEs in *A. baumannii* D1279779 and *P. protegens*

Pf-5 were shown to excise from the chromosome, albeit at a low level [30].

Neither IE has been tested for intercellular mobility. Here, we characterized

IME*Vch*USA3, a representative member of a subgroup of *dusA*-specific IEs

circulating in *Vibrio* species and distantly related to SGI1. We demonstrated that

IME*Vch*USA3 is mobilizable by IncC conjugative plasmids to *E. coli*. In the

presence of an IncC plasmid, this IME excises in practically all cells of the

population and becomes highly unstable (Figs 5B and 6A). We showed that its

excision was under the control of AcaCD provided by the IncC plasmid and

required *rdfN*, a gene whose expression is driven by an AcaCD-responsive

promoter (Fig 4). *rdfN* encodes a novel RDF distantly related to the pyocin

activator protein PrtN of *Pseudomonas*. *rdfN* seems to be ubiquitous, yet highly

divergent, in *dusA*-specific IEs reported by Farrugia *et al.* [30]. For instance,

RdfN (PrtN) encoded by the IE of *P. protegens* Pf-5 shares only 29% identity with

RdfN of IME*Vch*USA3, and their promoters are unrelated. Hence, the expression

of *rdfN* homologs encoded by different families of *dusA*-specific IEs is likely

controlled by different factors. Only the IEs that have evolved AcaCD-responsive

promoters for their RDF gene are expected to be mobilizable by IncC or related

plasmids.

Excision and mobilization of IME*Vch*USA3 occurred in the presence of a Δ*acaC*

but not a Δ*acaDC* mutant of the helper plasmid (Fig 5), confirming that *sgaC* of

the IME produces a functional activator subunit that can interact with AcaD

provided by the plasmid. Furthermore, we showed here that, unlike *acaC*, *sgaC*

plays a central role in the biology of IME*Vch*USA3 as the absence of *acaC* had

no effect on the excision or transfer of the IME, while the absence of *sgaC* in

spite of the presence of *acaC*, compromised its mobilization (Figs 5A, 5B and

5C). We recently showed that AcaD most likely stabilizes the binding of AcaC to

the DNA [22]. Therefore, AcaD and SgaC from IME*Vch*USA3 likely interact to

form a chimerical activator complex. This interaction could compensate for the

loss of *sgaD* in *yicC-* and type 1 and 2 *dusA*-specific IEs (Fig 1). The primary

sequences of AcaC and SgaC of IME*Vch*USA3 (type 1) share 85% identity. In

contrast, AcaC only shares 75% identity with SgaC of SGI1 and 64% identity with

SgaC of GI*Vch*USA5 (type 3), suggesting that retention of *sgaD* allowed faster

divergence of SgaC from AcaC. Retention of *sgaC* in the IEs could result from its

essential role as the elicitor of excision and replication reported for SGI1. Indeed,

although AcaCD binds to the promoters P_{xis} and P_{rep} of SGI1, it fails to initiate

transcription at these two promoters, unlike SgaCD [22]. Nonetheless, P_{xis} and

P_{rep} are not conserved in the IEs described here. *S004-rep* is missing, whereas

rdfN or *rdfM* replaced *xis* with novel AcaCD-responsive promoters (Fig 4 and S3

Fig). This observation raises intriguing questions regarding the recruitment of

functional gene cassettes and their assimilation in a regulatory pathway. How did

xis, *rdfN*, and *rdfM* acquire their AcaCD-responsive promoters? Is it by

convergent evolution? What are the signals driving *rdfN* expression and IE

excision in *dusA*-specific IEs resembling prophages?

Approximately 3 copies per cell of IME*Vch*USA3 were detected in the presence

of the helper IncC plasmid (Fig 5B), lower than the copy number reported for

SGI1 (~8 copies/cell) [20,22]. IME*Vch*USA3 lacks SGI1's replication module

(*S004-rep-oriV*); however, one of the multiple genes of unknown function could

encode an unidentified replication initiator protein. Notably, GI*Vch*O27-1 encodes

a putative replication protein with an N-terminal replicase domain (PF03090) and

a C-terminal primase domain (PriCT-1, PF08708) [20]. Multiple IEs also carry

putative replicons based on *repA* and *trfA* (Fig 1), suggesting that independent

replication is crucial in their lifecycle, perhaps to improve their stability in the

presence of a helper plasmid [20–24].

Farrugia *et al.* [30] hypothesized that *dusA*-specific IEs could restore the

functioning of DusA. We demonstrated here that IME*Vch*USA3 provides a new

promoter allowing expression of *dusA*, though at a lower level than in IME-free

cells, and restores the open reading frame with an altered N-terminus (Fig 6).

Similarly, the ICE SXT that targets the 5' end of the peptide chain release factor

3 (RF3) gene *prfC* provides a new promoter and N-terminus in both *V. cholerae*,

its original host, and *E. coli* [37]. In both cases, the consequences of the

alteration of the N-terminus on the activity of the protein remain unknown.

The relative positions of *int* and *rdfN*/*rdfM* across the *attP* site suggest that, to

remain functional, the recombination modules must be acquired or exchanged

when the IEs are in their excised circular form. The promiscuity of different

families of IEs targeting *yicC*, *dusA*, and *trmE* and mobilizable by IncC plasmids

could act as the catalyst for these recombination events. During entry into a new

host cell by conjugation, IncC plasmids elicit the excision of such IEs and

promote homologous recombination between short repeated sequences in

response to double-stranded break induced by host defense systems (CRISPR-

Cas3) [34]. Hence the diversification of IncC plasmid-mobilizable IEs could be a

side effect of the DNA repair mechanism used by these plasmids.

Unlike SGI1 and its siblings, all *dusA*-specific SGI1-like IEs reported here lack

antibiotic resistance genes. Furthermore, SGI1 variants are prevalent in several

pathogenic species and relatively well-conserved, whereas their *dusA*-specific

relatives are scarce and highly divergent. These observations suggest that

despite the considerable functional resemblances between *trmE-* and *dusA*-

specific SGI1-like IEs, the epidemiological success of the SGI1 lineage has

directly stemmed from the acquisition of class I integrons conferring multidrug

resistance by forerunner elements such as SGI0 [38]. Based on the phylogenetic

relationships between the core proteins MpsA, TraG, SgaC and TraN, *oriT* loci,

and integrase proteins (Figs 2 and 3, and S2A and S3 Figs), we propose a

hypothetical evolutionary pathway leading to the emergence of the different types

of IEs described here (Fig 7). The diversity of *dusA*-specific IEs and relative

homogeneity of the SGI1 group suggest that the latter originated from the

progenitor of IncA and IncC plasmids via a *dusA*-specific IE intermediate.

Figure 7. Proposed hypothetical evolutionary pathway of SGI1-like IEs. The

sequence of events was inferred from the phylogenetic trees presented in this

study, site of integration and conservation of *traH* and *sgaD* in the IEs. The

proposed pathway ignores the gene cargo and presumes that the IE lineages

evolved from the progenitor of IncA and IncC plasmids. The *dusA*-specific

recombination module was chosen as the progenitor to minimize gain/loss and

recombination events. Green and red arrows indicate gene gains and losses,

respectively. The orange dashed line indicates a probable recombination event

from which stemmed GI*Vch*O27-1.

Materials and Methods

Bacterial strains and media

Bacterial strains and plasmids used in this study are described in Table 2. Strains

were routinely grown in lysogeny broth at 37°C in an orbital shaker/incubator and

were preserved at -75°C in LB broth containing 20% (vol/vol) glycerol. Antibiotics

were used at the following concentrations: ampicillin (Ap), 100 µg/ml;

chloramphenicol (Cm), 20 µg/ml; erythromycin (Em), 200 µg/ml; kanamycin (Kn),

10 µg/ml for single-copy integrants of pOP*lacZ*-derived constructs, 50 µg/ml

otherwise; nalidixic acid (Nx), 40 µg/ml; rifampicin (Rf), 50 µg/ml; spectinomycin

(Sp), 50 µg/ml; tetracycline (Tc), 12 µg/ml. Diaminopimelate (DAP) was

supplemented to a final concentration of 0.3 mM when necessary.

Mating assays

Conjugation assays were performed as previously described [25]. However,

mixtures of donor and recipient cells were incubated on LB agar plates at 37°C

for 4 hours. Donors and recipients were selected according to their sole

chromosomal markers. When required, mating experiments were performed

using LB agar plates supplemented with 0.02% arabinose to induce expression

of pBAD30-derived complementation vectors. Frequencies of transconjugant

formation were calculated as ratios of transconjugant per donor CFUs from three

independent mating experiments.

Molecular biology

Plasmid DNA was prepared using the QIAprep Spin Miniprep Kit (Qiagen),

according to manufacturer's instructions. Restriction enzymes used in this study

were purchased from New England Biolabs. Q5 DNA polymerase (New England

Biolabs) and EasyTaq DNA Polymerase (Civic Bioscience) were used for

amplifying cloning inserts and verification, respectively. PCR products were

purified using the QIAquick PCR Purification Kit (Qiagen), according to

manufacturer's instructions. *E. coli* was transformed by electroporation as

described by Dower *et al.* [39] in a Bio-Rad GenePulser Xcell apparatus set at 25

µF, 200 Ω and 1.8 kV using 1-mm gap electroporation cuvettes. Sanger

sequencing reactions were performed by the Plateforme de Séquençage et de

Génotypage du Centre de Recherche du CHUL (Québec, QC, Canada).

Plasmids and strains constructions

Plasmids and oligonucleotides used in this study are listed in Table 2 and S2

Table, respectively. IME*Vch*USA3^{Cm} was constructed by inserting the *pir*-

dependent replication RP4-mobilizable plasmid pSW23T [40] at locus

CGT85_RS05425 of *V. cholerae* OYP6G08 (Genbank NZ_NMSY01000009) by

homologous recombination. Briefly, CGT85_RS05425 was amplified using primer

pair dusAigEcoRIF/dusAigEcoRIR. The amplicon was digested with EcoRI and

cloned into EcoRI-digested pSW23T using T4 DNA ligase. The resulting plasmid

was confirmed by restriction profiling and DNA sequencing, then introduced into

the DAP-auxotrophic *E. coli* β2163 [40] by transformation. This strain was used

as a donor in a mating assay to transfer the plasmid into *V. cholerae* OYP6G08,

generating IME*Vch*USA3^{Cm}. Single-copy integration of the pSW23T derivative

was confirmed by PCR and antibiotic resistance profiling.

IME*Vch*USA3^{Kn} was constructed from IME*Vch*USA3^{Cm}. Briefly, pVCR94^{Kn} Δ*acr2*

was transferred from the DAP-auxotrophic *E. coli* KH40 into OYP6G08 bearing

IME*Vch*USA3^{Cm}. After selection on LB agar medium supplemented with

chloramphenicol and kanamycin, Cm^R Kn^R *V. cholerae* OYP6G08

transconjugants were confirmed by growth on thiosulfate-citrate-bile salts-

sucrose (TCBS) agar medium (Difco). In *V. cholerae*, the integration and excision

of the IME were confirmed by amplification of the *attL*, *attR*, *attB*, and *attP* sites

with primer pairs oRD4/ORD6, oRD1/oRD3, oRD1/oRD6, and oRD4/oRD3,

respectively. IME*Vch*USA3^{Cm} was then mobilized from OYP6G08 to *E. coli*

CAG18439. In *E. coli*, the integration and excision of the IME were confirmed by

amplification of the *attL*, *attR*, *attB* and *attP* sites with primer pairs oRD4/ORD5,

oRD2/oRD3, oRD2/oRD5 and oRD4/oRD3, respectively. IMEVchUSA3^{Kn} was

constructed by replacing pSW23T with a single kanamycin resistance marker

using the one-step chromosomal gene inactivation technique with primer pair

dusAscarNoFRTf/dusAscarNoFRTr and pKD13 as the template. The deletions

Δ*sgaC* and Δ*prtN* in IME*Vch*USA3^{Kn} were obtained using the primer pairs

oFD26r/oFD26f and DelprtNr/DelprtNf, and pKD3 and pVI36 as the templates,

respectively. The Δ*dapA* deletion mutant of *E. coli* MG1655 was constructed

using primer pair FwDeltaDapA-MG1655/ RvDeltaDapA-MG1655 and pKD3 as

the template. The Δ*lacZ* mutation was introduced in *E. coli* CAG18439 using

primer pair lacZW-B/lacZW-F and plasmid pKD4 as the template. The *dusA'*-

'lacZ fusion was introduced in *E. coli* BW25113 and CAG18439 using primer pair

oDF15/oDF16 and pVI42B as the template. The fortieth codon of *dusA* was fused

to the eighth codon of *lacZ* downstream of the *attB* site. The λRed recombination

system was expressed using either pSIM6, pSIM9 or pKD46 [41,42]. When

appropriate, resistance cassettes were excised from the resulting constructions

using the Flp-encoding plasmid pCP20 [43]. All deletions were validated by

antibiotic profiling and PCR.

Fragments encompassing promoter regions upstream of *int, traN, traG, s018* and

rdfN were amplified using primer pairs oFD6.f/oFD6.r, oFD1.f/oFD1.r,

oFD3.f/oFD3.r, oFD5.f/oFD5.r and oFD4.f/oFD4.r, respectively, and genomic

DNA from *E. coli* CAG18439 *dusA*::IME*Vch*USA3^{Kn} as the template. The

amplicons were digested with PstI/XhoI and cloned into PstI/XhoI-digested

pOP*lacZ* [17]. The resulting constructs were single-copy integrated into the *attB*$_\lambda$

chromosomal site of *E. coli* BW25113 using pINT-ts [44]. To construct the

expression vectors pBAD-*rdfN* and pBAD-*sgaC*, PCR fragments containing *rdfN*

or *sgaC* were amplified from genomic DNA of *E. coli* CAG18439 bearing

IME*Vch*USA3 as the template and primer pairs prtNEcoRIf/prtNHindIIIrev and

oFD38r/oFD44f, respectively. The PCR fragments were digested by either EcoRI

or SacI, and HindIII and cloned into pBAD30 cut with the same enzymes.

mini-IE was constructed as follows. The 1,591-bp fragment of excised circular

IME*Vch*USA3^{Kn} that contains *attP-int* was amplified using primer pair

oVB12/oVB10 and genomic DNA from *E. coli* CAG18439 *dusA*::IME*Vch*USA3^{Kn}

as the template. The 1,421-bp fragment of pVI36 that contains *aadA7* was

amplified using primer pair oVB11/oVB13. Both fragments were joined using the

PCR-based overlap extension method [45]. After the final PCR amplification

using oVB12/oVB13, the amplicon was purified, digested with SacI, and ligated.

The ligation mixture was then transformed into *E. coli* EC100. Transformant

colonies were selected on LB agar supplemented with spectinomycin. The

constitutive expression of *int* and the absence of replicon prompted the

spontaneous integration of mini-IE at the 5' end of *dusA* in EC100.

All final constructs were verified by PCR and DNA sequencing by the Plateforme

de Séquençage et de Génotypage du Centre de Recherche du CHUL (Québec,

QC, Canada).

qPCR assays

qPCR assays for quantification of excision and copy number of IME*Vch*USA3^{Kn}

were carried out as described previously [22] with the following modification.

attB$_{dusA}$ (241 bp) and *higA* (229 bp) of IME*Vch*USA3^{Kn} were quantified using

primer pairs attBdusAqPCRfwd/ attBdusAqPCRrev and higAqPCRfwd/

higAqPCRrev, respectively (S2 Table). The excision rate and copy number of

IME*Vch*USA3^{Kn} were calculated as the ratio of free *attB$_{dusA}$* site per chromosome

and as the ratio of *higA* per chromosome, respectively. The data were analyzed

and normalized using all three chromosomal genes *dnaB*, *hicB* and *trmE* as

references and the qBase framework as described previously [22,46].

β-galactosidase assays

The assays were carried out on LB agar plates supplemented with 5-bromo-4-

chloro-3-indolyl-β-D-galactopyranoside (X-gal) or in LB broth using *o*-nitrophenyl-

β-D-galactopyranoside (ONPG) as the substrate as described previously [32].

acaDC expression from pBAD-*acaDC* was induced by adding 0.2% arabinose to

a refreshed culture grown to an OD_{600} of 0.2, followed by a 2-h incubation at

37°C with shaking prior to cell sampling.

Comparative analyses

Sequences were obtained using blastp against the Genbank Refseq database

with the primary sequences of key proteins MpsA, $TraG_S$, SgaC, $TraN_S$ of SGI1

(Genbank AAK02039.1, AAK02037.1, AAK02036.1, AAK02035.1, respectively),

and Int_{dusA} of IE*Vch*Bra2 (Genbank EEO15317.1) and Int_{yicC} of IE*Eco*MOD1

(Genbank WP_069140142.1). Hits were exported, then sorted by accession

number to identify gene clusters that likely belong to complete IEs. Sequences of

IEs were manually extracted and the extremities were identified by searching for

the direct repeats contained in *attL* and *attR* sites. When an IE sequence

spanned across two contigs (e.g., IE*Vch*Hai10 and IE*Ppl*Ind1), the sequence was

manually assembled. IE sequences were clustered using cd-hit-est with a 0.95

nucleotide sequence identity cut-off [47]. Some of the annotated sequences were

manually curated to correct missing small open reading frames such as *mpsB*,

and inconsistent start codons. Pairwise comparisons of Int, MpsA, TraG, SgaC

and TraN proteins were generated with blastp using sets of representative

proteins selected after clustering using cd-hit with a 0.95 sequence identity cut-

off (Int, MpsA, TraG, SgaC) or a 0.90 sequence identity cut-off (TraN) [47].

Heatmaps showing the blastp identity scores were drawn using the Python library

seaborn v0.11.1 [48]. Circular blast representations (blast atlases) were

generated with the Blast Ring Image Generator (BRIG) 0.95 [49], with blastn or

blastp, against SGI1ΔIn104 and IE*Vch*USA2, with an upper identity threshold of

80% and a lower identity threshold of 60%. Antibiotic resistance gene prediction

was conducted using the Resistance Gene Identifier (RGI) software and CARD

3.1.3 database [50]. AcaCD binding motifs were identified using FIMO and MAST

[51] with the AcaCD motif matrix (S1 Matrix) described previously [17]. Logos for

attL and *attR* repeats were generated with MAST [51] using alignments of

sequences flanking the IE_{dusA} elements identified in this work.

Phylogenetic analyses

Evolutionary analyses were conducted in MEGA X [52] and inferred by using the

maximum likelihood method based on the JTT (MpsA or SgaC proteins), LG

(Int_{dusA}, Int_{yicC}, TraG or $RepA_{IncFII}$ proteins) or WAG (TraN) matrix-based models

[53–55]. Protein sequences were aligned with Muscle [56]. Aligned sequences

were trimmed using trimal v1.2 using the automated heuristic approach [57].

Initial tree(s) for the heuristic search were obtained automatically by applying

Neighbor-Join and BioNJ algorithms to a matrix of pairwise distances estimated

using a JTT model, and then selecting the topology with the superior log

likelihood value. A discrete Gamma distribution was used to model evolutionary

rate differences among sites (5 categories) for Int_{dusA} (parameter = 3.5633), Int_{yicC}

(parameter = 2.6652), SgaC (parameter = 1.4064), TraG (parameter = 1.9005)

and TraN (parameter = 1.6476) proteins. For Int_{dusA}, MpsA and TraG, the rate

variation model allowed for some sites to be evolutionarily invariable ([+I], 7.81%

sites for Int_{dusA}, 44.62% sites for MpsA and 5.22% sites for $TraG_S$). The trees are

all drawn to scale, with branch lengths measured in the number of substitutions

per site. In all trees, bootstrap supports are shown as percentages at the

branching points only when > 80%.

oriT sequences were obtained manually using the previously identified *oriT* of

SGI1 as the reference [19], then clustered using cd-hit-est with a 1.0 nucleotide

sequence identity cut-off. Sequences were then aligned using Muscle and a

NeighborNet phylogenetic network was built using SplitsTree4 [58] with default

parameters (Uncorrected_P method for distances and EqualAngle drawing

method). The secondary structures of the aligned *oriT* sequences were predicted

using RNAalifold v2.4.17 from the ViennaRNA package [59]. Default options

were used (including no RIBOSUM scoring), except for the following: no

substituting "T" for "U" (--noconv), no lonely pairs (--noLP), no GU pairs (--noGU)

and DNA parameters (-P DNA). The predicted Vienna output and the annotated

alignment were merged into a predicted secondary structure of SGI1 *oriT* color-

coded to display the inter-island diversity.

Statistical analyses and figures preparation

Numerical data presented in graphs are available in S3 Dataset. Prism 8

(GraphPad Software) was used to plot graphics and to carry out statistical

analyses. All figures were prepared using Inkscape 1.0 (https://inkscape.org/).

Acknowledgments

We are grateful to Yann Boucher for the kind gift of *Vibrio cholerae* OYP6G08

and Kévin T. Huguet for technical assistance. We thank Nicolas Rivard and

David Roy for their insightful comments on the manuscript.

References

References

1.   Bellanger X, Payot S, Leblond-Bourget N, Guédon G. Conjugative and
mobilizable genomic islands in bacteria: evolution and diversity. FEMS
Microbiol Rev. 2014;38: 720–760. doi:10.1111/1574-6976.12058

2.   Guédon G, Libante V, Coluzzi C, Payot S, Leblond-Bourget N. The Obscure
World of Integrative and Mobilizable Elements, Highly Widespread Elements
that Pirate Bacterial Conjugative Systems. Genes. 2017;8: 337.
doi:10.3390/genes8110337

3.   Daccord A, Ceccarelli D, Burrus V. Integrating conjugative elements of the
SXT/R391 family trigger the excision and drive the mobilization of a new
class of *Vibrio* genomic islands. Mol Microbiol. 2010;78: 576–588.
doi:10.1111/j.1365-2958.2010.07364.x

4.   Daccord A, Ceccarelli D, Rodrigue S, Burrus V. Comparative analysis of
mobilizable genomic islands. J Bacteriol. 2013;195: 606–614.
doi:10.1128/JB.01985-12

5.   Waldor MK. Mobilizable genomic islands: going mobile with *oriT* mimicry.
Mol Microbiol. 2010;78: 537–540. doi:10.1111/j.1365-2958.2010.07365.x

6.   Carraro N, Rivard N, Ceccarelli D, Colwell RR, Burrus V. IncA/C Conjugative
Plasmids Mobilize a New Family of Multidrug Resistance Islands in Clinical
Vibrio cholerae Non-O1/Non-O139 Isolates from Haiti. mBio. 2016;7:
e00509-16. doi:10.1128/mBio.00509-16

7.   de Curraize C, Siebor E, Neuwirth C. Genomic islands related to *Salmonella*
genomic island 1; integrative mobilisable elements in *trmE* mobilised in trans
by A/C plasmids. Plasmid. 2021;114: 102565.
doi:10.1016/j.plasmid.2021.102565

8.   Mulvey MR, Boyd DA, Olson AB, Doublet B, Cloeckaert A. The genetics of
Salmonella genomic island 1. Microbes and Infection. 2006;8: 1915–1922.
doi:10.1016/j.micinf.2005.12.028

9.   Boyd D, Peters GA, Cloeckaert A, Boumedine KS, Chaslus-Dancla E,
Imberechts H, et al. Complete nucleotide sequence of a 43-kilobase
genomic island associated with the multidrug resistance region of
Salmonella enterica serovar Typhimurium DT104 and its identification in
phage type DT120 and serovar Agona. J Bacteriol. 2001;183: 5725–5732.
doi:10.1128/JB.183.19.5725-5732.2001

10. Grim CJ, Hasan NA, Taviani E, Haley B, Chun J, Brettin TS, et al. Genome
sequence of hybrid *Vibrio cholerae* O1 MJ-1236, B-33, and CIRS101 and

comparative genomics with *V. cholerae*. J Bacteriol. 2010;192: 3524–3533.
doi:10.1128/JB.00040-10

11. Cummins ML, Hamidian M, Djordjevic SP. *Salmonella* Genomic Island 1 is
Broadly Disseminated within Gammaproteobacteriaceae. Microorganisms.
2020;8: 161. doi:10.3390/microorganisms8020161

12. Hall RM. *Salmonella* genomic islands and antibiotic resistance in *Salmonella*
enterica. Future Microbiol. 2010;5: 1525–1538. doi:10.2217/fmb.10.122

13. Doublet B, Boyd D, Mulvey MR, Cloeckaert A. The *Salmonella* genomic
island 1 is an integrative mobilizable element. Mol Microbiol. 2005;55: 1911–
1924. doi:10.1111/j.1365-2958.2005.04520.x

14. Douard G, Praud K, Cloeckaert A, Doublet B. The *Salmonella* genomic
island 1 is specifically mobilized in trans by the IncA/C multidrug resistance
plasmid family. PLoS ONE. 2010;5: e15302.
doi:10.1371/journal.pone.0015302

15. Harmer CJ, Hall RM. The A to Z of A/C plasmids. Plasmid. 2015;80: 63–82.
doi:10.1016/j.plasmid.2015.04.003

16. Wu W, Feng Y, Tang G, Qiao F, McNally A, Zong Z. NDM Metallo-β-
Lactamases and Their Bacterial Producers in Health Care Settings. Clin
Microbiol Rev. 2019;32: e00115-18. doi:10.1128/CMR.00115-18

17. Carraro N, Matteau D, Luo P, Rodrigue S, Burrus V. The master activator of
IncA/C conjugative plasmids stimulates genomic islands and multidrug
resistance dissemination. PLoS Genet. 2014;10: e1004714.
doi:10.1371/journal.pgen.1004714

18. Kiss J, Papp PP, Szabó M, Farkas T, Murányi G, Szakállas E, et al. The
master regulator of IncA/C plasmids is recognized by the *Salmonella*
Genomic island SGI1 as a signal for excision and conjugal transfer. Nucleic
Acids Res. 2015;43: 8735–8745. doi:10.1093/nar/gkv758

19. Kiss J, Szabó M, Hegyi A, Douard G, Praud K, Nagy I, et al. Identification
and Characterization of *oriT* and Two Mobilization Genes Required for
Conjugative Transfer of *Salmonella* Genomic Island 1. Front Microbiol.
2019;10: 457. doi:10.3389/fmicb.2019.00457

20. Huguet KT, Rivard N, Garneau D, Palanee J, Burrus V. Replication of the
Salmonella Genomic Island 1 (SGI1) triggered by helper IncC conjugative
plasmids promotes incompatibility and plasmid loss. PLoS Genet. 2020;16:
e1008965. doi:10.1371/journal.pgen.1008965

21. Szabó M, Murányi G, Kiss J. IncC helper dependent plasmid-like replication
of Salmonella Genomic Island 1. Nucleic Acids Research. 2021;49: 832–
846. doi:10.1093/nar/gkaa1257

22. Durand R, Huguet KT, Rivard N, Carraro N, Rodrigue S, Burrus V. Crucial
role of *Salmonella* genomic island 1 master activator in the parasitism of
IncC plasmids. Nucleic Acids Res. 2021; in press. doi:10.1093/nar/gkab204

23. Harmer CJ, Hamidian M, Ambrose SJ, Hall RM. Destabilization of IncA and
IncC plasmids by SGI1 and SGI2 type *Salmonella* genomic islands.
Plasmid. 2016;87–88: 51–57. doi:10.1016/j.plasmid.2016.09.003

24. Huguet KT, Gonnet M, Doublet B, Cloeckaert A. A toxin antitoxin system
promotes the maintenance of the IncA/C-mobilizable *Salmonella* Genomic
Island 1. Sci Rep. 2016;6: 32285. doi:10.1038/srep32285

25. Carraro N, Durand R, Rivard N, Anquetil C, Barrette C, Humbert M, et al.
Salmonella genomic island 1 (SGI1) reshapes the mating apparatus of IncC
conjugative plasmids to promote self-propagation. PLOS Genetics. 2017;13:
e1006705. doi:10.1371/journal.pgen.1006705

26. Humbert M, Huguet KT, Coulombe F, Burrus V. Entry Exclusion of
Conjugative Plasmids of the IncA, IncC, and Related Untyped Incompatibility
Groups. J Bacteriol. 2019;201: e00731-18. doi:10.1128/JB.00731-18

27. Roberts AP, Johanesen PA, Lyras D, Mullany P, Rood JI. Comparison of
Tn*5397* from Clostridium difficile, Tn*916* from *Enterococcus faecalis* and the
CW459*tet*(M) element from *Clostridium perfringens* shows that they have
similar conjugation regions but different insertion and excision modules.
Microbiology. 2001;147: 1243–1251. doi:10.1099/00221287-147-5-1243

28. Burrus V, Pavlovic G, Decaris B, Guédon G. The ICE*St1* element of
Streptococcus thermophilus belongs to a large family of integrative and
conjugative elements that exchange modules and change their specificity of
integration. Plasmid. 2002;48: 77–97. doi:10.1016/s0147-619x(02)00102-6

29. Bioteau A, Durand R, Burrus V. Redefinition and unification of the SXT/R391
Family of integrative and conjugative elements. Appl Environ Microbiol.
2018;84: pii: e00485-18. doi:10.1128/AEM.00485-18

30. Farrugia DN, Elbourne LDH, Mabbutt BC, Paulsen IT. A novel family of
integrases associated with prophages and genomic islands integrated within
the tRNA-dihydrouridine synthase A (*dusA*) gene. Nucleic Acids Res.
2015;43: 4547–4557. doi:10.1093/nar/gkv337

31. Daccord A, Mursell M, Poulin-Laprade D, Burrus V. Dynamics of the SetCD-
regulated integration and excision of genomic islands mobilized by

integrating conjugative elements of the SXT/R391 family. J Bacteriol.
2012;194: 5794–5802. doi:10.1128/JB.01093-12

32. Rivard N, Colwell RR, Burrus V. Antibiotic Resistance in *Vibrio cholerae*:
Mechanistic Insights from IncC Plasmid-Mediated Dissemination of a Novel
Family of Genomic Islands Inserted at *trmE*. mSphere. 2020;5: e00748-20.
doi:10.1128/mSphere.00748-20

33. Thorsted PB, Shah DS, Macartney D, Kostelidou K, Thomas CM.
Conservation of the genetic switch between replication and transfer genes of
IncP plasmids but divergence of the replication functions which are major
host-range determinants. Plasmid. 1996;36: 95–111.
doi:10.1006/plas.1996.0037

34. Roy D, Huguet KT, Grenier F, Burrus V. IncC conjugative plasmids and
SXT/R391 elements repair double-strand breaks caused by CRISPR-Cas
during conjugation. Nucleic Acids Res. 2020;48: 8815–8827.
doi:10.1093/nar/gkaa518

35. Carraro N, Sauvé M, Matteau D, Lauzon G, Rodrigue S, Burrus V.
Development of pVCR94ΔX from *Vibrio cholerae*, a prototype for studying
multidrug resistant IncA/C conjugative plasmids. Front Microbiol. 2014;5: 44.
doi:10.3389/fmicb.2014.00044

36. Doublet B, Golding GR, Mulvey MR, Cloeckaert A. Secondary Chromosomal
Attachment Site and Tandem Integration of the Mobilizable *Salmonella*
Genomic Island 1. Fairhead C, editor. PLoS ONE. 2008;3: e2060.
doi:10.1371/journal.pone.0002060

37. Hochhut B, Waldor MK. Site-specific integration of the conjugal *Vibrio*
cholerae SXT element into *prfC*. Mol Microbiol. 1999;32: 99–110.

38. de Curraize C, Siebor E, Neuwirth C, Hall RM. SGI0, a relative of
Salmonella genomic islands SGI1 and SGI2, lacking a class 1 integron,
found in *Proteus mirabilis*. Plasmid. 2019; 102453.
doi:10.1016/j.plasmid.2019.102453

39. Dower WJ, Miller JF, Ragsdale CW. High efficiency transformation of *E. coli*
by high voltage electroporation. Nucleic Acids Res. 1988;16: 6127–6145.

40. Demarre G, Guérout A-M, Matsumoto-Mashimo C, Rowe-Magnus DA,
Marlière P, Mazel D. A new family of mobilizable suicide plasmids based on
broad host range R388 plasmid (IncW) and RP4 plasmid (IncPalpha)
conjugative machineries and their cognate *Escherichia coli* host strains. Res
Microbiol. 2005;156: 245–255. doi:10.1016/j.resmic.2004.09.007

41. Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products. Proc Natl Acad Sci U S A.
2000;97: 6640–6645.

42. Datta S, Costantino N, Court DL. A set of recombineering plasmids for
gram-negative bacteria. Gene. 2006;379: 109–115.
doi:10.1016/j.gene.2006.04.018

43. Cherepanov PP, Wackernagel W. Gene disruption in *Escherichia coli*: TcR
and KmR cassettes with the option of Flp-catalyzed excision of the
antibiotic-resistance determinant. Gene. 1995;158: 9–14. doi:10.1016/0378-
1119(95)00193-a

44. Haldimann A, Wanner BL. Conditional-replication, integration, excision, and
retrieval plasmid-host systems for gene structure-function studies of
bacteria. J Bacteriol. 2001;183: 6384–6393. doi:10.1128/JB.183.21.6384-
6393.2001

45. Senanayake SD, Brian DA. Precise large deletions by the PCR-based
overlap extension method. Mol Biotechnol. 1995;4: 13.
doi:10.1007/BF02907467

46. Hellemans J, Mortier G, De Paepe A, Speleman F, Vandesompele J. qBase
relative quantification framework and software for management and
automated analysis of real-time quantitative PCR data. Genome Biology.
2007;8: R19. doi:10.1186/gb-2007-8-2-r19

47. Li W, Godzik A. Cd-hit: a fast program for clustering and comparing large
sets of protein or nucleotide sequences. Bioinformatics. 2006;22: 1658–
1659. doi:10.1093/bioinformatics/btl158

48. Waskom M. seaborn: statistical data visualization. JOSS. 2021;6: 3021.
doi:10.21105/joss.03021

49. Alikhan N-F, Petty NK, Ben Zakour NL, Beatson SA. BLAST Ring Image
Generator (BRIG): simple prokaryote genome comparisons. BMC
Genomics. 2011;12: 402. doi:10.1186/1471-2164-12-402

50. Alcock BP, Raphenya AR, Lau TTY, Tsang KK, Bouchard M, Edalatmand A,
et al. CARD 2020: antibiotic resistome surveillance with the comprehensive
antibiotic resistance database. Nucleic Acids Res. 2020;48: D517–D525.
doi:10.1093/nar/gkz935

51. Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, et al. MEME
SUITE: tools for motif discovery and searching. Nucleic Acids Res. 2009;37:
W202-208. doi:10.1093/nar/gkp335

52.   Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular
Evolutionary Genetics Analysis across Computing Platforms. Mol Biol Evol.
2018;35: 1547–1549. doi:10.1093/molbev/msy096

53.   Jones DT, Taylor WR, Thornton JM. The rapid generation of mutation data
matrices from protein sequences. Bioinformatics. 1992;8: 275–282.
doi:10.1093/bioinformatics/8.3.275

54.   Whelan S, Goldman N. A General Empirical Model of Protein Evolution
Derived from Multiple Protein Families Using a Maximum-Likelihood
Approach. Mol Biol Evol. 2001;18: 691–699.
doi:10.1093/oxfordjournals.molbev.a003851

55.   Le SQ, Gascuel O. An Improved General Amino Acid Replacement Matrix.
Mol Biol Evol. 2008;25: 1307–1320. doi:10.1093/molbev/msn067

56.   Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and
high throughput. Nucleic Acids Res. 2004;32: 1792–1797.
doi:10.1093/nar/gkh340

57.   Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for
automated alignment trimming in large-scale phylogenetic analyses.
Bioinformatics. 2009;25: 1972–1973. doi:10.1093/bioinformatics/btp348

58.   Huson DH, Bryant D. Application of Phylogenetic Networks in Evolutionary
Studies. Molecular Biology and Evolution. 2006;23: 254–267.
doi:10.1093/molbev/msj030

59.   Lorenz R, Bernhart SH, Höner zu Siederdissen C, Tafer H, Flamm C,
Stadler PF, et al. ViennaRNA Package 2.0. Algorithms Mol Biol. 2011;6: 26.
doi:10.1186/1748-7188-6-26

60.   Kirchberger PC, Orata FD, Nasreen T, Kauffman KM, Tarr CL, Case RJ, et
al. Culture-independent tracking of *Vibrio cholerae* lineages reveals complex
spatiotemporal dynamics in a natural population. Environ Microbiol. 2020;22:
4244–4256. doi:10.1111/1462-2920.14921

61.   Heidelberg JF, Eisen JA, Nelson WC, Clayton RA, Gwinn ML, Dodson RJ,
et al. DNA sequence of both chromosomes of the cholera pathogen *Vibrio
cholerae.* Nature. 2000;406: 477–483. doi:10.1038/35020000

62.   Singer M, Baker TA, Schnitzler G, Deischel SM, Goel M, Dove W, et al. A
collection of strains containing genetically linked alternating antibiotic
resistance elements for genetic mapping of *Escherichia coli.* Microbiol Rev.
1989;53: 1–24.

63.   Ceccarelli D, Daccord A, René M, Burrus V. Identification of the Origin of
Transfer (*oriT*) and a New Gene Required for Mobilization of the SXT/R391

Family of Integrating Conjugative Elements. J Bacteriol. 2008;190: 5328–
5338. doi:10.1128/JB.00150-08

64. Grenier F, Matteau D, Baby V, Rodrigue S. Complete genome sequence of
Escherichia coli BW25113. Genome Announc. 2014;2: pii: e01038-14.
doi:10.1128/genomeA.01038-14

65. Garriss G, Waldor MK, Burrus V. Mobile antibiotic resistance encoding
elements promote their own diversity. PLoS Genet. 2009;5: e1000775.
doi:10.1371/journal.pgen.1000775

66. Guzman LM, Belin D, Carson MJ, Beckwith J. Tight regulation, modulation,
and high-level expression by vectors containing the arabinose P_{BAD}
promoter. J Bacteriol. 1995;177: 4121–4130.

Table 1. Main features of the IEs described in this study.

IE Name	Organism[1]	Size (bp)	Target site	Genbank accession number
GI*Vch*Rus1	*V. cholerae* 1	30,204	*dusA*	NZ_SMZE01000022
IE*Vch*Aus1	*V. cholerae* A12JL36W30	27,410	*dusA*	NZ_VIOZ01000074
IE*Vch*USA5	*V. cholerae* OYP2C05	28,706	*dusA*	NZ_NMTM01000021
IME*Vch*USA3	*V. cholerae* OYP6G08	30,910	*dusA*	NZ_NMSY01000009
IE*Vch*A215[2]	*V. cholerae* A215 sv Inaba	29,933	*dusA*	CWPR01000020.1
IE*Vch*USA2	*V. cholerae* 692-79	29,931	*dusA*	MIPA01000024
IE*Vch*N2751[2]	*V. cholerae* N2751	30,018	*dusA*	NZ_VSGL01000012
IE*Vch*N2744[2]	*V. cholerae* N2744	30,134	*dusA*	NZ_VSGF01000021.1
IE*Vch*N2708	*V. cholerae* N2708	27,248	*dusA*	NZ_VSFQ01000013
IE*Vch*N2786	*V. cholerae* N2786	24,658	*dusA*	NZ_VSHP01000008
IE*Vch*N2817	*V. cholerae* N2817	28,717	*dusA*	NZ_VSIM01000004
IE*Vch*Chn1	*V. cholerae* N2787	27,195	*dusA*	NZ_VSHQ01000015
IE*Vch*Ban1	*V. cholerae* EM-1676-A	36,519	*dusA*	NZ_KB662834
IE*Vch*Hai10	*V. cholerae* 2012Env-2	37,162	*dusA*	NZ_JSTD01000059/60
IE*Vch*Swe1	*V. cholerae* 11116	28,100	*dusA*	NZ_MDYK01000006
IE*Vch*Bra2	*V. cholerae* TMA-21	28,230	*dusA*	ACHY01000008
IE*Vme*USA1	*V. metoecus* 07-2435	23,518	*dusA*	NZ_LCUE01000016
IE*Vpa*Chn1	*V. parahaemolyticus* GIMxtf41-2013.07	28,589	*dusA*	NZ_MRWJ01000014
IE*Vpa*Chn2	*V. parahaemolyticus* C2_8	25,944	*dusA*	NZ_NNLT01000047
IE*Vpa*Ban1a	*V. parahaemolyticus* NIHCB0757	29,418	*dusA*	AVPX01000004
IE*Vvu*USA1	*V. vulnificus* VA-WGS-18041	30,228	*dusA*	NZ_RBZL01000019
IE*Eco*MOD1	*E. coli* MOD1-EC5437	25,611	*yicC*	NZ_NLPO01000006
IE*Sen*USA1	*S. enterica* Kentucky ARS-CC8289	25,970	*yicC*	NZ_MCPS01000044
IE*Ppl*Ind1	*P. plantistimulans* L1E11	22,777	*yicC*	NZ_LAPT01000132/095
GI*Vch*O27-1	*V. cholerae* 10432-62	26,646	*trmE*	CP010812

946 [1] *V.*, *Vibrio*; *E.*, *Escherichia*; *S.*, *Salmonella*; *P.*, *Pokkaliibacter*

947 [2] Not represented in Fig 1 as nucleotide sequences contain gaps

948 **Table 2. Strains and elements used in this study.**

Strains or elements	Relevant genotype or phenotype[a]	Source or reference
V. cholerae		
OY6PG08	Environmental, Oyster Pond, MA, USA, August 2009	[60]
N16961	O1 El Tor	[61]
E. coli		
β2163	(F–) RP4-2-Tc::Mu Δ*dapA*::(*erm-pir*) (Kn Em)	[40]
CAG18439	MG1655 *lacZU118 lacI42*::Tn*10* (Tc)	[62]
BW25113	F⁻ Δ(*araD-araB*)567, Δ*lacZ4787*(::*rrnB-3*), λ⁻, *rph-1*, Δ(*rhaD-rhaB*)568, *hsdR514*	[41]
EC100	F⁻ *mcrA Δ(mrr-hsdRMS-mcrBC) Φ80dlacZΔM15 ΔlacX74 recA1 endA1 araD139 Δ(ara, leu)7697 galU galK λ⁻ rpsL (Sm^R) nupG*	Epicentre, Madison Wis.
KH40	MG1655 Δ*dapA*::*cat* (Cm)	This study
VB112	Rf-derivative of MG1655	[63]
GG56	Nx-derivative of BW25113	[35,64]
FD034	CAG18439 Δ*lacZ dusA'-'lacZ-aad7* (Tc Sp)	This study
FD036	GG56 *dusA'-'lacZ-aad7* (Nx Sp)	This study
Plasmids		
pKD3	Cm^R PCR template for one-step chromosomal gene inactivation (Cm)	[41]
pKD4	Kn^R PCR template for one-step chromosomal gene inactivation (Kn)	[41]
pKD13	Kn^R PCR template for one-step chromosomal gene inactivation (Kn)	[41]
pVI36	Sp^R PCR template for one-step chromosomal gene inactivation (Sp)	[63]
pVI42B	pVI36 BamHI::P_{lac}-*lacZ* (Sp)	[65]
pSW23T	pSW23::*oriT*$_{RP4}$; *oriV*$_{R6Kγ}$ (Cm)	[40]
pOP*lacZ*	pAH56 *lacZ* (Kn)	[17]
pBAD30	*ori*$_{p15A}$ *bla araC* P_{BAD} (Ap)	[66]
pBAD-*acaDC*	pBAD30::*acaDC* (Ap)	[17]
pBAD-*rdfN*	pBAD30::*rdfN* (Ap)	This study
pBAD-*sgaC*	pBAD30::*sgaC* (Ap)	This study
pVCR94^Kn Δ*acr2*	Δ*acr2* mutant of pVCR94^Kn (Su Kn)	[34]
pVCR94^Sp	Sp^R derivative of pVCR94 (Su Sp)	[17]
pVCR94^Sp Δ*acaC*	Δ*acaC* mutant of pVCR94^Sp (Su Sp)	[17]
pVCR94^Sp Δ*acaDC*	Δ*acaDC* mutant of pVCR94^Sp (Su Sp)	[17]
Integrative elements		
IME*Vch*USA3		This study
IME*Vch*USA3^Cm	IME*Vch*USA3 CGT85_RS05425ΩpSW23T (Cm)	This study
IME*Vch*USA3^Kn	Kn^R derivative of IME*Vch*USA3 (Kn)	This study
IME*Vch*USA3^Kn Δ*sgaC*	Δ*sgaC* mutant of IME*Vch*USA3 (Kn)	This study
IME*Vch*USA3^Kn Δ*rdfN*	Δ*rdfN* mutant of IME*Vch*USA3 (Kn)	This study
mini-IE	*attP-int-aad7* derived from IME*Vch*USA3 (Sp)	This study

949 [a]Ap, ampicillin; Cm, chloramphenicol; Em, erythromycin, Kn, kanamycin; Nx, Nalidixic

950 acid; Rf, rifampin; Sm, streptomycin; Sp, spectinomycin; Su, sulfamethoxazole; Tc,

951 tetracycline; Tm, trimethoprim; ts, thermosensitive.

Supporting information

S1 Fig. Comparative sequence analysis of SGI1-like *dusA*-specific IEs. Blastn and blastp atlases using either SGI1ΔIn104 (A) or IE*Vch*USA2 (B) as the reference. Coding sequences appear on the outermost circle in blue for the positive strand and red for the negative strand, with the *oriT* depicted as a grey arc. All other sequences are represented only according to their homology with the reference, with full opacity corresponding to 100% identity and gaps indicating identity below 60%. The order of the IEs in the atlases is indicated according to the color keys shown in the inset of panel B.

S2 Fig. NeighborNet phylogenetic network and predicted secondary structure of 39 *oriT* loci of SGI1-like IEs. Each IE's integration site and type are annotated. The sequence of canonical SGI1 (Genbank AF261825.2) was used as a reference to show the predicted secondary structure of all *oriT* sequences. Pairs can be perfectly conserved, imperfectly conserved (1/39 not conserved), not conserved (> 1/39), or an A-T or G-C pair only. In the latter case, the sequence is not conserved, but the predicted local secondary structure is.

S3 Figure. Maximum likelihood phylogenetic analysis of key proteins of SGI1-related IEs. The trees for MpsA (A), TraG (B), SgaC (C) and TraN (D) proteins are drawn to scale, with branch lengths measured in the number of substitutions per site over 321, 1,145, 188, and 968 amino acid positions, respectively. For clarity, the lengths of the branches linking the two groups in panels A and C were artificially divided by 8 and 4, respectively. Taxa corresponding to IEs targeting *trmE* and *yicC* are shown by a light blue circle and a red circle, respectively. All other taxa correspond to *dusA*-specific IEs. Proteins accession numbers are provided in S1 Table and S2 Dataset.

S4 Figure. Alignment of AcaCD-responsive promoters predicted in IEs targeting

***dusA*, *yicC* and *trmE*.** Promoter sequences are grouped based on the function of the

expressed genes as follows: (A) RDFs; (B) mating pair stabilization; (C) mating pair

formation and stabilization; (D) unknown. AcaCD binding sites are shown in green. Logo

sequences and *p*-values were generated by MAST [51]. Known transcription start sites

are shown in blue [17,22]. Predicted Shine-Dalgarno sequences are shown in pink. The

initiation start codon is shown in bold letters.

S5 Figure. Excision of IME*Vch*USA3 is enhanced in IncC⁺ cells. (A) Model of

excision of IME*Vch*USA3. (B and C) Detection of *attB*, *attP*, *attL* and *attR* sites by PCR

in colonies of *V. cholerae* OYO6G08 bearing (lanes 9 to 16) or lacking (lanes 1 to 8)

pVCR94^{Kn} Δ*acr2.* Control lanes: L, 1Kb Plus DNA ladder (Transgen Biotech); +,

V. cholerae N16961 genomic DNA. (D) Detection of *attB*, *attP*, *attL* and *attR* sites by

PCR in transconjugant colonies of *E. coli* CAG18439 (lanes 1 to 4). L, 100bp Plus II

DNA Ladder (Transgen Biotech)

S1 Table. Features of the identified IEs and associated strains.

S2 Table. Oligonucleotides used in this study.

S1 Dataset. Features of ORFs in the identified IEs.

S2 Dataset. Clusters generated by cd-hit for Int, MpsA, TraG, SgaC and TraN.

S3 Dataset. Numerical data presented in Figs 4, 5 and 6.

S1 Matrix. AcaCD motif matrix to identify AcaCD binding sites.

A IME*Vch*USA3 excision rate

B IME*Vch*USA3 copy number

C

D

E

A

E. coli K12 *dusA attB* site

```
TAATCTCTACATTTGAAA ATG CAC GGT AAT TCT GAA ATG CAA AAA ATC AAC CAA ACC AGC GCA ATG CCT GAA
                    M   H   G   N   S   E   M   Q   K   I   N   Q   T   S   A   M   P   E

AAA ACT GAC GTT CAC TGG AGT GGT CGG TTT AGC G TT GCA CCA ATG CTC GAC TGG A CG GAC AGA CAT
K   T   D   V   H   W   S   G   R   F   S   V   A   P   M   L   D   W   T   D   R   H
```

E. coli K12 *dusA*::IME*Vch*USA3^{Kn} *attR* junction

```
AGTTGAGATTAGTTAAGA ATG AGC CAG AAA CAG AAT ACA GAT AAA TCT TTG CAA GAC AGT GAG TTA TCT AGT
                    M   S   Q   K   Q   N   T   D   K   S   L   Q   D   S   E   L   S   S

AAT ATT GAT GTG TCA TTA AGT CGT AGA TTC AGC TG GCC CCC ATG CTT GAC TGG A CG GAC AGA CAT
N   I   D   V   S   L   S   R   R   F   S   V   A   P   M   L   D   W   T   D   R   H
```

B

C