Main Manuscript for

MRP5 and MRP9 Play a Concerted Role in Male Reproduction and Mitochondrial Function

Ian Chambers[1], Praveen Kumar[2], Jens Lichtenberg[3], Pengcheng Wang[2], Jianshi Yu[2], John Phillips[4], Maureen Kane[2], David Bodine[3], Iqbal Hamza[1]

[1]Department of Animal & Avian Sciences and Department of Cell Biology & Molecular Genetics, University of Maryland, College Park, MD, USA

[2]Department of Pharmaceutical Sciences, University of Maryland School of Pharmacy, Baltimore, MD, USA

[3]Genetics and Molecular Biology Branch, National Human Genome Research Institute, National Institutes of Health, Bethesda, MD, USA

[4]Department of Medicine, University of Utah School of Medicine, Salt Lake City, UT, USA

Corresponding Author: Iqbal Hamza

Email: hamza@umd.edu

Author Contributions:

IGC: Conceptualization, Methodology, Investigation, Formal analysis, Validation, Visualization, Writing—original draft, Writing—review and editing

PK: Conceptualization, Methodology, Investigation, Formal analysis, Validation

JL: Methodology, Investigation, Formal analysis, Validation, Visualization

PW: Methodology, Investigation, Formal analysis

JY: Methodology, Investigation, Formal analysis

JP: Conceptualization, Methodology, Investigation, Formal analysis, Validation, Resources, Funding acquisition, Writing—review and editing

MAK: Conceptualization, Methodology, Investigation, Formal analysis, Validation, Resources, Funding acquisition, Writing—review and editing

DB: Conceptualization, Methodology, Investigation, Formal analysis, Validation, Resources, Funding acquisition, Writing—review and editing

IH: Conceptualization, Methodology, Investigation, Formal analysis, Validation, Visualization, Resources, Software, Supervision, Funding acquisition, Writing—original draft, Writing—review and editing

35 **Competing Interest Statement:**

36 IH is the President and Founder of Rakta Therapeutics Inc. (College Park, MD), a company
37 involved in the development of heme transporter-related diagnostics. He declares no other
38 competing financial interests.

39
Classification: Biological Sciences, Genetics

Keywords: Multidrug Resistance Proteins (MRPs), Reproduction, Heme, Mitochondria
Associated Membranes (MAMs), Fertility

This PDF file includes:

Main Text
Figures 1 *to* 6
47

Abstract
Multidrug Resistance Proteins (MRPs) are transporters that play critical roles in cancer even
though the physiological substrates of these enigmatic transporters are poorly elucidated. In
Caenorhabditis elegans, MRP5/ABCC5 is an essential heme exporter as *mrp-5* mutants are
unviable due to their inability to export heme from the intestine to extra-intestinal tissues. Heme
supplementation restores viability of these mutants but fails to restore male reproductive deficits.
Correspondingly, cell biological studies show that MRP5 regulates heme levels in the mammalian
secretory pathway even though MRP5 knockout (KO) mice do not show reproductive phenotypes.
The closest homolog of MRP5 is MRP9/ABCC12, which is absent in *C. elegans* raising the
possibility that MRP9 may genetically compensate for MRP5. Here, we show that MRP5 and
MRP9 double KO mice are viable but reveal significant male reproductive deficits. Although
MRP9 is highly expressed in sperm, MRP9 KO mice show reproductive phenotypes only when
MRP5 is absent. Both ABCC transporters localize to mitochondrial-associated membranes
(MAMs), dynamic scaffolds that associate the mitochondria and endoplasmic reticulum.
Consequently, DKO mice reveal abnormal sperm mitochondria with reduced mitochondrial
membrane potential and fertilization rates. Metabolomics show striking differences in metabolite
profiles in the DKO testes and RNA-seq show significant alterations in genes related to
mitochondrial function and retinoic acid metabolism. Targeted functional metabolomics reveal
lower retinoic acid levels in the DKO testes and higher levels of triglycerides in the mitochondria.
These findings establish a model in which MRP5 and MRP9 play a concerted role in regulating
male reproductive functions and mitochondrial sufficiency.
69

Significance Statement
MRPs are typically implicated in cancer biology. Here, we show that MRP9 and MRP5 localize to
mitochondrial-associated membranes and play a concerted role in maintaining mitochondrial
homeostasis and male reproductive fitness. Our work fills in significant gaps in our understanding
of MRP9 and MRR5 with wider implications in male fertility. It is plausible that variants in these
transporters are associated with male reproductive dysfunction.
76
77

Main Text

Introduction

Multidrug Resistance-associated Proteins (MRPs) belong to the type C subclass of the
ATP Binding Cassette (ABCC) transporters, a highly conserved family of proteins that have been
studied for decades for their role in cancers (1-7). MRPs are notoriously upregulated in a number
of malignancies because of their ability to efflux a wide range of synthetic and chemotherapeutic
agents, conferring cancer cells resistance to drug treatments (3, 8-11). Due to their critical
importance in human health and clinical outcomes, myriads of MRP substrates have now been
identified, particularly from *in vitro* studies of tumor cell lines (3, 4, 6, 12, 13). However, the
promiscuity of drugs effluxed in this context is likely a functional consequence of cancer cells
hijacking these active transporters rather than that being their true evolutionary role. Therefore, a
major outstanding item remaining in the MRP field is the determination of true physiological
functions and substrates for a number of these transporters.
Previous work from our group shed some light on the function of one such ABCC
transporter, MRP5/ABCC5, where studies in worms, yeast, zebrafish and mammalian cells
demonstrated it to be a heme exporter (14). Prior to this discovery, the role of MRP5 appeared to
be that of cyclic nucleotide export from cells as demonstrated by transport of cGMP in membrane
vesicles expressing human MRP5 *in vitro* (12, 15). However, the generation of MRP5 null mice
demonstrated little to no contribution in cGMP efflux and no observable overt phenotypes, leaving
the physiological relevance of this transporter unclear outside its *in vitro* or cancer cell context
(16, 17). This is not the case in *C. elegans*, a heme auxotroph, where MRP-5 is absolutely
essential for intercellular heme trafficking and knockout of *mrp*-5 results in heme deficiency and
embryonic lethality (14).
While cell biological studies in mouse embryonic fibroblasts from MRP5 KO mice
confirmed that MRP5 indeed plays a conserved role in heme export i.e. transporting heme into
the secretory pathway for incorporation into hemoproteins or exporting it from the cell across the
plasma membrane (14, 18-20), MRP5 KO mice do not show any heme-related phenotypes *in*
vivo. A postulated hypothesis to explain these differences could be genetic compensation by
other closely related transporters, which are phylogenetically conserved in higher order
eukaryotes but absent in the *C. elegans* genome (14, 21). A candidate for this proposed
compensation and further functional delineation is MRP9/ABCC12, the closest relative of MRP5
and one of the last of the ABCC proteins to have been discovered (6, 22, 23). MRP9 is highly
expressed in male reproductive tissues but fails to effectively efflux drugs *in vitro*, and its
endogenous function remains to be elucidated (6, 24, 25).
In this work, we generate a MRP9 knockout and MRP9 / MRP5 double knockout (DKO)
mouse models to fully characterize these transporters' physiological functions. Loss of MRP9 and
MRP5 disrupts mitochondrial homeostasis and regulatory pathways in the testes, with RNA-seq
and metabolomic analyses revealing large-scale perturbation in genes and metabolism related to
mitochondrial function. Together, these results establish a concerted and critical role for MRP9
and MRP5 in male reproduction and mitochondrial metabolism, implying that the ancestral role of
MRP5 in *C. elegans* reproduction was co-opted in the vertebrate ancestor.

Results

Loss of MRP9 in mice induces the expression of MRP5 in the testes
C. elegans mrp-5(ok2067) deletion mutants are lethal unless rescued with dietary heme
supplementation (14). Serendipitously, we found that even in the presence of heme *mrp-*
5(ok2067) male worms are defective in mating **(Figures S1A and S1B)**. These worms exhibit
defective mating apparatus with stunted tail fan formation, reduced numbers of rays, and spicules
which fail to retract despite normal mating behaviors. Given that hermaphrodite *mrp-5* knockout
worms mate normally with wildtype males (not shown), it would appear the defect is paternal. For
these reasons, we hypothesized that MRP-5 may have additional functions related to male

reproduction outside of its traditional role in heme transport and homeostasis. MRP5 knockout
mice however show no reproductive deficiencies, prompting inquiry into other ABCC transporters
that may genetically compensate for the loss of MRP5 in higher order vertebrates.
Phylogenetic analysis of ABCC transporters across metazoans confirmed that members
of the short MRP clade (ABCCs - 4, 5, 7, 11, 12) cluster tightly together across species indicating
their evolutionary relation (6, 14). Within this subclade, *C. elegans* has only MRP-5 whereas most
vertebrates have an additional paralog, MRP9 **(Figure 1A)**, which is conserved in all higher-order
vertebrates but remains to be fully characterized *in vivo.* We therefore generated MRP9 mutant
mice via CRISPR/Cas9 to explore MRP9's endogenous functions. Utilizing guide RNAs targeted
to the second exon of *Abcc12* we were able to generate 11 founder animals with various indels at
the target site and a 17 bp insertion line was selected for further characterization **(Figures 1B**
and S1C). RT-PCR confirmed that the frameshift-inducing mutation was transcribed with no
alternate start sites or splice variants identified **(Figure S1D)**.
Abcc12-/- animals are viable and reach adulthood with no overt phenotypes.
Immunoblots of lysates from testes of adult mice confirmed no detectable MRP9 protein in
knockout animals **(Figure 1C)**. Immunohistochemistry of tissue sections showed MRP9 in both
the maturing and developed spermatozoa of seminiferous tubules and caudal epidymis,
respectively, consistent with the findings reported by Ono et al **(Figure 1D)** (24). Recent single
cell RNAseq datasets of the testis niche further confirmed these findings as the gene expression
profile of *Abcc12* is distinctly germ cell specific **(Figure S1E)** (26). This is in contrast to *Abcc5*,
which appears to be normally expressed in early spermatogonia precursors as well as Sertoli and
Leydig cells **(Figure S1E)**. Immunoblotting of lysates from additional tissues of wildtype and KO
mice revealed the testes as the only organ with any appreciable amounts of both, MRP9 and
MRP5 **(Figure 1C)**. Importantly, MRP9 KO testes showed elevated levels of MRP5 raising the
notion that MRP5 could be compensating for MRP9 deficiency in the testes **(Figures 1E and 1F)**.

MRP9 and MRP5 are required for male reproductive fitness

MRP9 KO mice do not show any statistically significant defects in reproduction, with
expected litter sizes and normal Mendelian F1 segregation when intercrossed [n = 166, Chi-
squared = 0.976, *P* value = 0.614] **(Figure 2A)**. MRP9 KO crossed to MRP5 KO mice
successfully generated DKO mice which reached adulthood with no overt phenotypes. Although
DKO mice were viable, they were deficient in their reproductive capacity; the average number of
pups from intercrosses of DKO mice were significantly reduced compared to WT [n > 33 litters, *P*
value < 0.0001] **(Figure 2B)**. Interestingly, fecundity was restored back to single KO levels when
DKO females were crossed with WT or double heterozygous males alone, suggestive that the
failure may be paternal [n > 21 litters, *P* value < 0.0001] **(Figure 2B)**. Furthermore, DKO male
mice setup to breed had high incidences of penile prolapse and clogging of their urogenital tracks
with seminal coagulum (five DKOs compared to zero WT male mice), regardless of the female
genotype **(Figure 2C)**. These findings, taken together with our protein expression data, prompted
a closer investigation into male fitness as the major defect behind the DKO reproductive failure.
Indeed, sperm from DKO mice had significantly lower *in vitro* fertilization (IVF) rates compared to
wildtype **(Figure 2D)**. Interestingly this may be attributed to sperm fitness / quality rather than
quantity as DKO sperm showed trends of decreased motility with no differences in total sperm
production **(Figures S2A, S2B and 2E)**.

MRP9 is localized to mitochondrial associated membranes in the testes

To understand why DKO male mice have reduced fecundity and sperm fitness we further
characterized the function of these proteins in this context. Given their distinct spatiotemporal
gene expressions **(Figure S1E)**, we hypothesized that these MRPs may play overlapping
subcellular roles during spermatid maturation. The Borst group had shown that MRP9 is localized
to the sperm midpiece (24, 27, 28), and when expressed in HEK293 closely resembled an ER
localization (24). In polarized MDCKII cells, transfection of *ABCC12* showed a distinct intracellular
distribution of MRP9 in stark contrast to MRP5 that localized primarily to the basolateral surface

of the plasma membrane **(Figure 3A)**. To determine if this was also the case *in vivo* or merely a
consequence of ectopic expression*,* we performed subcellular fractionation of lysates from mouse
testes. Immunoblotting showed that MRP9 protein was actually significantly enriched in 9,000 *x g*
fractions which pellet mitochondria **(Figure 3B)**. This result was surprising given previously
published findings (24, 29, 30), and that MRP9 lacks any obvious mitochondrial targeting
sequences. However, an explanation for this unexpected result could be that mitochondrial-
associated membranes (MAMs), the contact sites of the ER which are intimately associated with
the mitochondria, are enriched in our crude mitochondrial fractions (31-34).
Next, we performed Airyscan super resolution microscopy on HeLa cells transiently
transfected with human *ABCC12* **(Figure 3C)**. 3D rendering revealed that MRP9 was in close
proximity or at the interface of the mitochondria (Mitotracker) and ER (Calnexin) **(Figure 3C and
Movie S1)**. To validate this localization, we performed Percoll-based differential centrifugations to
further separate crude mitochondrial fractions into purified mitochondria and MAMs from testes
tissue homogenates (32). Immunoblots of distinct testes fractions revealed that MRP9 was in fact
not present in purified mitochondria but significantly enriched in MAMs, characterized by the
markers Mitofusin 2 and Calnexin **(Figure 3D)**. Interestingly, probing for MRP5 from these same
fractions indicated that it too was enriched in the MAMs as well as total membranes **(Figure 3E)**.
These results indicate that subcellular expression patterns of MRPs may vary dependent on
tissue contexts.

Multi-omics of DKO testes reveals extensive perturbation and mitochondrial dysfunction
In order to systematically dissect why the combined loss of MRP9 and MRP5 manifests
in male reproductive dysfunction, we sought to understand what substrates may be transported
by these proteins and how their absence may impact homeostatic pathways in the testes. To
achieve this, we performed untargeted metabolomics and RNAseq from paired testes of the same
mice to integrate any differential metabolites and gene expression changes **(Figure S3A)**. Global
untargeted metabolomics of testes indicated significant perturbation, with 24 of the top 25
differential species measured from negative ion mode acquisition of aqueous phase metabolites
showing accumulation in the DKO **(Figure 4A)**. Dramatic differences in metabolite profiles were
evident however regardless of the extraction method (aqueous vs organic) or the ion mode of
acquisition (negative vs positive) **(Figures S3B and S3C)**. Analysis of these data allowed for the
identification of mass-to-charge ratios (m/z) of hyper-accumulated candidate metabolites.
However, given our sampling was untargeted, a particular *m/z* could represent multiple
possibilities of actual biological compounds even given the high mass accuracy and resolution of
the Q-TOF platform. As a means to associate these putative species with specific metabolites, all
significant m/z's identified from our six untargeted datasets were pooled and processed using
MetaboAnalyst Mummichog analyses (35). From the species of interest analyzed, 65 metabolites
were assigned possible KEGG IDs **(Table S1)**. Amongst these, succinyl-CoA, which is also an
essential TCA cycle energy source and a heme precursor, had the largest fold-change increase
in DKO testes (36, 37).
RNAseq analysis from the same tissues found 3,736 differentially expressed genes
between WT and DKO, with a number of top genes differentially expressed including
uncharacterized and testis-specific predicted genes (*Gm10354*, *Gm9758*, *5430401F13Rik*) as
well as mitochondrial respiratory genes (*Atp5k*, *Ndufa1*, *Atp5e*) **(Figure 4B)**. To unbiasedly
determine the pathways and Gene Ontology (GO) terms most significantly affected in these mice
we performed Ingenuity Pathway Analysis (IPA) **(Figures 4C and S3D)**. The majority of the top
GO canonical pathways identified by IPA were intimately interconnected and overlapped with
mitochondrial function (**Figures 4C and 4D**). Of note, the top two pathways identified, EIF2
Signaling and Protein Ubiquitination, are critical for mitigating unfolded protein response and
mitochondrial dysfunction associated with iron and heme deficiencies (38).
To generate metabolite-gene relationships, we integrated our metabolomics and RNAseq
data using MetaboAnalyst Joint Pathway Analysis. The merged data analysis revealed significant
mitochondrial pathway dysfunction with energy production components of the cell (TCA Cycle,

Pyruvate Metabolism, and Glycolysis) showing the highest impact **(Figure 4E)**. Taken together,
these findings reinforce a role for MRP9 and MRP5 in mitochondrial metabolism in the testes.

EIF2a phosphorylation and associated signaling pathways mitigate mitochondrial
dysfunction

235   Having identified mitochondrial dysfunction as the overarching factor from our integrative
analysis, we sought to determine the specific mechanisms that underlie this dysfunction in the
testes. Mining our RNAseq dataset, we built an analysis pipeline to determine and curate fold
changes of all genes associated with a given pathway from published GO databases for
downstream investigation and mapping. In querying all genes associated with the mitochondria
and mitochondrial dysfunction, we identified over 200 dysregulated genes (**Figure 5A**). We then
asked what individual genes or pathways may be involved in responding to such a significant
level of disruption. Given the top-related pathway identified from our IPA studies was EIF2
signaling, we investigated all associated genes and found 95 that were perturbed in DKO testes
(**Figure 5B**). EIF2 signaling and EIF2a phosphorylation by the heme responsive kinase HRI
(*Eif2ak1*) are established mechanisms for regulating translation to reduce mitochondrial stress
and unfolded protein response (UPR) in situations of intracellular heme deficiency (38-41). In
particular this model is well documented in the erythron, where heme is critical as it becomes the
limiting factor for incorporation into hemoglobin for red blood cell differentiation **(Figure S4A)**.
Limited studies have characterized a similar role in mitigating mitochondrial dysfunction in
neurons, but very little is known about HRI and EIF2a signaling in other tissues such as the testes
(39, 42-45). Indeed, analysis of the testes single cell RNAseq dataset determined that both
factors are expressed during spermatid maturation **(Figure S4B)**, and immunoblotting of total
testes homogenates confirmed significant EIF2a protein expression **(Figure 5C)** (26). Importantly,
EIF2a phosphorylation was significantly increased in the DKO mice, substantiating the
involvement of the EIF2a pathway in mitigating UPR and mitochondrial dysfunction in the testes
(Figure 5C and 5D). We next asked if the observed EIF2 signaling in the testes of DKO mice was
in response to heme deficiency, as is the hallmark in the erythron. If so, one would expect that
iron/heme homeostasis should also be perturbed in the testes of these mice. Subsequent probing
with our pipeline pathway revealed heme and iron homeostasis related genes were in fact
significantly altered **(Figure S4C)**. To determine if heme levels themselves were also impacted,
we measured total heme in male reproductive tissues. Heme levels in the caudal epididymides,
the storage sites for mature sperm, showed a significant decrease **(Figure 5E)**, while whole
testes and seminal vesicles showed no differences **(Figure S4D and S4E)**. These results
suggest that there is insufficient heme specifically in spermatozoa of DKO mice.

265   A critical component of the canonical EIF2 signaling cascade for maintaining
mitochondrial function and redox homeostasis is the alteration of mTOR signaling to inhibit
proliferation and enable differentiation (38, 46). Indeed, the mTOR signaling pathway was one of
the most significant GO terms identified by IPA **(Figure 4C and S3D)**, which we also validated
with our analysis pipeline **(Figure S5A)**. Upregulation of EIF-4E, a highly abundant translation
regulation factor in the testes, has been shown to play an important role in spermatogenesis
through regulation of capping stage-specific mRNAs during germ cell development (47). A
modulator of these related pathways is retinoic acid, which has been implicated in mTOR/EIF-4E
cascade and plays a key role in a variety of biological contexts including regulation of genes
impacting mitochondrial function (48-52). Principal targets of retinoic acid include key OxPhos
and respiratory genes such as NADH dehydrogenase subunits, cytochrome c oxidases and the
ATPases, some of the top differentially expressed genes from our RNAseq **(Figure 4B)** (51, 53).
Investigation of retinoic acid related genes revealed significant perturbation including decreases
in the cytosolic chaperone *Crabp1*, responsible for delivering retinoic acid for catabolism, as well
as binding receptors (RAR, RXRs), suggestive of dysregulated retinoic acid homeostasis **(Figure**
S5B) (54-56). Importantly, retinoic acid signaling is essential for male germ cell differentiation,
exerting key control over meiotic initiation and spermatogenesis, with defects in retinoid
homeostasis a well-established cause of male infertility (57-60). Taken together, we hypothesized
that retinoic acid homeostasis/signaling may play an important role in the phenotypes observed in

our DKO mice. Quantification of the major components of the retinoid pathway revealed that total
levels of vitamin A (retinol) and storage esters were not perturbed in the testes **(Figure S5C and**
S5D), but all-*trans* retinoic acid levels were significantly reduced DKOs compared to wildtype
(Figure S5E). Other retinoic acid isomers were unchanged (13-*cis* retinoic acid, 9,13-di-*cis*
retinoic acid) or below the limit of detection (9-*cis* retinoic acid) (data not shown).
To further validate if retinoic acid binding transcription factors and cascade regulators
were indeed impacting mitochondrial genes of interest, we performed unbiased motif discovery
on regulatory sequences of all up-regulated and down-regulated genes from our pathway pipeline
(Figure 5A and S5F). The top five statistically significant DNA sequences identified were further
analyzed and queried against known binding motifs. With two independent analysis platforms
(MEME suite and HOMER) we identified significant enrichment of conserved retinoic acid related
binding motifs including putative RXRs and RORA binding sequences **(Figure S5F)** (61). These
findings suggest a mechanistic link between transcripts related to mitochondrial dysfunction,
retinoid signaling genes, and decreased retinoic acid levels, corresponding with a specific
enrichment of retinoic acid binding motifs in the DKO mice.

Ablation of MRP9 and MRP5 significantly damages sperm mitochondria

To determine whether mitochondrial disruption could be inducing the gene expression
and metabolomic changes that in-turn impact sperm fertilization in the DKO testes, we analyzed
mitochondria oxidative phosphorylation complexes in total testes lysates. However, all five
complexes appeared to be normal and unchanged in DKOs **(Figure S6A)**. We next fractionated
the testes and performed high-throughput targeted metabolomics on each isolated fraction,
allowing for the identification and quantification of specific metabolites within each subcellular
niche **(Figure S6B)**. We found that the mitochondria and MAM fractions of the DKO testes hyper-
accumulate triglycerides, indicative of mitochondrial metabolism dysfunction **(Figure S6C and**
S6D) (62, 63).
Mature caudal epididymal sperm were then collected from WT and DKO adult mice by
swim-out for mitochondrial membrane potential ($\Delta\Psi_M$) analysis and transmission electron
microscopy (TEM). Staining of sperm with JC-1 revealed that DKO mice have reduced J
aggregates and $\Delta\Psi_M$ compared to WT **(Figure 7A, 7B and S7A)**. Strikingly, TEM cross sections
of sperm midpieces revealed DKO mice also have highly vacuolated and aberrant mitochondria,
in contrast to the ubiquitously electron dense cristae in the WT **(Figure 7C)**. These defects were
not observed in the single KOs or double heterozygous animals **(Figure S7B)**.

Discussion

ABCC proteins are one of the most well-studied class of transporters due to their critical
roles in cancer biology, multidrug resistance and associated human diseases (3, 64-67). Taken
together, our studies fill significant gaps in our understanding of not only of MRP9 but also MRP5,
uncovering novel endogenous functions related to reproduction. Given MRP5 is genetically
dispensable in the mouse model with low gene expression in male reproductive tissues, no
studies have examined its protein levels or function in the testes (14, 68-70). Klein et al showed a
weak MRP5 signal in Leydig cells of rats but nothing in macaque or human testis (71). We show
that MRP5 is detectable in the testes, but that it also appears to be enriched in mitochondrial
associated membranes along with MRP9. Though MRP5 still localizes to basolateral plasma
membranes in polarized MDCKII cells, our tissue fractionation studies bring to light the possibility
of additional novel tissue dependent expression patterns and subcellular localizations for this well
studied transporter. We envision that these results may have significant implications when it
comes to cancer and the distinct roles MRPs may play dependent upon tissue / cell type. We
propose the following model for the concerted function of MRP9 and MRP5 in the testes **(Figure**
7D): In the absence of MRP9 and MRP5, metabolites such as heme and/or other substrates, are
inappropriately trafficked or distributed causing mitochondrial damage. This dysfunction induces
signaling through the EIF2 and mTOR pathways to regulate protein translation and UPR with
corresponding changes in retinoic acid levels and signaling. Germ cells differentiate into mature

spermatozoa, but these are functionally defective as they fail to fertilize normal oocytes due to
mitochondrial insufficiencies.
MRP9 is an enigmatic ABCC transporter. It has a unique expression pattern, no known
drug substrates, and the last in the ABCC class still to be characterized (6, 24, 25). In the current
study, we fill in some of the gaps in our knowledge by generating an MRP9 knockout mouse
model to explore its endogenous functions. We find that MRP9 is specifically expressed in
developing spermatids and mature spermatozoa in mice and unlike most other MRPs, MRP9
does not localize to the plasma membrane irrespective of cell polarization, consistent with the
published findings of Ono et al (24). Contrary to other reported studies however, MRP9 is not
endogenously expressed in any cell lines tested in our hands, including primary mouse
embryonic fibroblasts, and MRP9 protein cannot be induced by sodium butyrate treatment
(Figure S8) as was reported recently (72).
Transient transfections and tissue fractionations provide strong evidence that MRP9
partitions specifically to mitochondrial-associated membranes of the testes, a distinct
compartment characterized by mitochondrial outer membrane and endoplasmic reticulum
markers. This is physiologically reasonable as MRP9 is localized to developing and mature
spermatozoa which are known to be devoid of traditional ER structures (73, 74). Though the ER
is depleted in the process of differentiation, vestiges of ER membranes likely remain, particularly
those in tight contact with mitochondria, and have been postulated to concentrate in the neck of
the sperm midpiece to serve as a store for Ca^{2+} necessary for sperm activation (74, 75). Indeed,
ER resident proteins such as ERp29 and Calreticulin have also been demonstrated to be present
in sperm and play essential roles in acrosome reaction, fertilization and stress responses (74,
75). We envision MRP9 may reside at these interfaces adjacent to mitochondria of the midpiece,
effluxing substrates essential for regulating mitochondrial function and/or trafficking metabolites
between these compartments as spermatogonia develop.
Loss of MRP9 alone has no discernable impact on male reproductive function and
knockout mice are viable. Additionally, histological inspection of the testis and epididymis of these
mice reveal effective spermatogenesis, with normal spermatid progression in seminiferous
tubules and mature sperm in their convoluted ducts. Unsurprisingly, we also see no
morphological defects evident by electron microscopy of sperm from MRP9 single knockouts, as
these male mice are perfectly capable of siring progeny even in a MRP5 heterozygous
background. Any discernable impact on reproduction is only found when both MRP9 and MRP5
are ablated. Indeed, DKO mice show significant reproductive failures due to decreased male
fecundity. Defects in sperm fertilization rates and high incidences of penile prolapse and seminal
coagulum clogging contribute to their decrease in reproductive output as measured by litter size.
This is likely due to the DKO sperm's inability to effectively fertilize, as supported by the
drastically reduced IVF rates.
In addition to decreased motility trends, we show substantial changes in the composition
of metabolites retained in the testes as well as significant perturbations in gene expression as a
consequence of the loss of MRP9 and MRP5. Importantly, our integrative analysis of these
approaches demonstrate that energy production and mitochondrial function were the most
impacted pathways in the DKO testes, consistent with our localization studies. Targeted high-
throughput metabolomics of subcellular fractions also revealed substantial changes at the
subcellular level. While others have shown that triglyceride species can accumulate at the whole
tissue level during mitochondrial dysfunction, we show that these lipid changes are occurring at
the mitochondria and MAMs in the DKO testes (62, 63, 76, 77).
The top pathway identified by our gene expression analysis was EIF2 signaling, which we
validated *in vivo* as a response to mitochondria dysfunction and/or induced by alterations in heme
levels in maturing sperm. Indeed, heme and iron related genes were differentially expressed in
the testes, and quantification of heme in the caudal epididymis of DKO mice showed significantly
decreased levels. In a similar vein to the well-established models of unfolded protein response in
the erythron, we envisage a necessity for heme sharing and partitioning as mitochondria are
rapidly being recruited and cells are undergoing proliferation for spermatogenesis and
differentiation. Though this translational regulation via EIF2 signaling has been well-studied in

other contexts, virtually nothing is known about its role in the testes. The only published study to
our knowledge relates to EIF2s3y, which was shown to be essential for regulating protein
proliferation in spermatogenesis (78). It is yet to be determined whether or not this subunit plays
any direct role to our findings here, although it is also significantly upregulated in the DKO testes
(P = 0.022).
Beyond EIF2 signaling, IPA identified several additional pathways: Protein Ubiquitination
(essential for unfolded protein response), mTOR Signaling, Regulation of EIF4, Oxidative
Phosphorylation, Germ Cell-Sertoli Cell Junction Signaling (an essential component of sperm
differentiation), and Mitochondrial Dysfunction. It is plausible that all-*trans* retinoic acid may be a
regulator in all these pathways for the following reasons: 1) it is essential in male fertility as one of
the primary signaling molecules from Sertoli cells for progression of spermatogenesis; 2) it is
critical for regulating gene expression of essential oxidative phosphorylation and mitochondrial
genes; and 3) it has been shown to modulate mTOR and EIF4 regulation (51, 52, 57). Consistent
with this notion, we find genes associated with retinoids and retinoic acid levels significantly
decreased in DKO testes. Furthermore, targets of gene expression perturbation have significant
enrichment of retinoic acid-related binding motifs for transcriptional regulation, albeit modulation
via these motifs may be indirect. RORA has been reported to act as a constitutive activator of
transcription in the absence of ligand and all-*trans* retinoic acid does not bind to RXR with high
affinity as compared to 9-*cis* retinoic acid (79, 80).
These systemic changes in the testes of DKO mice are in-line with the striking
morphological abnormality of their mitochondria at the ultrastructural level. The impact of such
evident dysfunction at the sperm midpiece has significant effects on male fecundity. The
mitochondrial sheath is essential for reproductive fitness as these tightly wrapped mitochondria
are critical for ATP production and maximize efficiency of energy delivery, directly correlated with
sperm swimming velocities across species (27, 28, 81-84). However, this immense output comes
with a price - because spermatozoa are so densely packed with mitochondria, they are also
epicenters for the production of detrimental free radicals/reactive oxygen species (27, 81, 83, 85).
If not balanced and mitigated, this can cause myriads of problems including lipid peroxidation,
which has been shown to damage the sperm and decrease their mitochondrial membrane
potential (27, 81, 83, 86). Loss of MRP5 and MRP9 alters mitochondrial homeostasis not only
inducing vacuolated and aberrant mitochondria, but also resulting in a significant reduction in
sperm mitochondrial membrane potential as observed by JC-1 staining. This relationship between
mitochondrial membrane potential and sperm function directly correlates with mitochondrial
health, sperm motility, and modulation of respiratory chain activity and oxidative phosphorylation
pathways (27, 82, 87-89). Concordantly, male fecundity relies heavily on this organelle and its
membranes being fully intact and functional for the overall health of the spermatozoa and their
fertilization capacity, cementing why DKO males show such significant loss of biological function
(27, 81-83). Male infertility is a significant problem globally, affecting over 70 million people and
accounting for 50% of documented issues in infertile couples (82, 90, 91). To our knowledge,
mutations or SNP variants of MRPs have not yet been investigated for any correlation to cases of
male reproductive dysfunction. Our findings here would suggest that human patients with
infertility issues, typically assayed for and characterized by mitochondrial dysfunctions, may have
hypomorphic mutations in MRP9 and MRP5.
432
433

Materials and Methods

General methods

Animals
All mice used were housed in Animal Sciences Departmental facilities at the University of
Maryland and kept in standard 12-hour light-dark cycles. Both sexes were used for all
experiments unless stipulated otherwise, such as the analysis of male sex organs and requisite

male reproduction phenotypes. ABCC12/MRP9 mutant mice were generated via CRISPR/Cas9
RNA injections at the National Human Genome Research Institute (NHGRI). Guide RNAs **(Table**
S3) were purchased from Sage Laboratories, St Louis, MO and Cas9 RNA was purchased from
Trilink Biotechnologies, San Diego, CA. The guide RNA and Cas9 RNA were combined at a
concentration of 5 ng/μl each in 10 mM Tris, 0.25 mM EDTA pH 7.5 for pronuclear injection.
Performed using standard procedures, fertilized eggs were collected from super-ovulated
C57BL/6J females approximately nine hours after mating with C57BL/6J male mice. Pronuclei
were injected with a capillary needle with a 1-2 μm opening pulled with a Sutter P-1000
micropipette puller. The RNAs were injected using a FemtoJet 4i (Eppendorf) with continuous
flow estimated to deposit approximately 2 pl of solution. Injected eggs were surgically transferred
to pseudo-pregnant CB6 F1 recipient females. DNA was then obtained from founder (F0) animals
by tail biopsy and sequenced to determine mutations. F0 animals carrying mutations were
crossed to C57BL/6 animals and the resulting heterozygous F1 animals were either intercrossed
to generate homozygous mutant animals or back crossed to C57BL/6 mice for propagation.
MRP5 mutant mice, a gift from Dr. Piet Borst (Netherlands Cancer Institute), had previously been
outcrossed into full C57BL/6 background and were bred to MRP9 mutant mice to generate double
knockouts (DKOs). MRP5 and MRP9 mice were genotyped from tail genomic DNA via PCR and
RFLP methods (see **Table S4** for complete primer sequences). In brief, tail snips were taken from
weaned 21-day old (P21) pups and digested. The MRP5 allele was determined as described
previously via amplification of specific MRP5 and hygromycin cassette primers (14). MRP9
mutants having a +17bp insertion were PCR amplified and subsequently restriction enzyme
digested with AvaII (specific to the mutation). All subsequent protocols described below involving
these animals were approved by the Institutional Animal Care and Use Committee (IACUC) at the
University of Maryland, College Park.

Standard Immunoblotting Methods
For specific detection of MRP5 and MRP9 protein lysates, as well all other general proteins
unless otherwise specified, the following protocol was used: Lysis in lysis buffer (0.75% NP40,
0.5% Triton X-100, 0.1% SDS, 1.75 mM EDTA, 1.25 mM EGTA, 0.25% DOC, 150 mM NaCl, 50
mM HEPES, 25 mM Tris-Cl pH 7.5) with 3x protease inhibitors (cOmplete Protease Inhibitor
Cocktail EDTA-free - 1 mM phenylmethylsulfonyl fluoride, 4 mM benzamidine, 2 μg/ml leupeptin,
and 1 μg/ml pepstatin, Roche Diagnostics, cat. number 1187358001) and 3x phosphatase
inhibitors (Millipore Inhibitor Cocktail Set, cat. number 524627) and sonicated twice. Protein
homogenates were then clarified via centrifugation at 15,000 $x\ g$ for 15 min at 4°C and insoluble
pellets discarded. Total protein concentration of the supernatant was measured using the
Bradford assay (Bradford Reagent Protein Assay Dye Reagent, Bio-Rad, cat. number 5000006).
Unboiled protein samples were then mixed with 4x Laemmli sample buffer and 100 mM
dithiothreitol (DTT), and 100 μg of protein/lane was loaded and separated in an SDS-PAGE gel
and transferred to a nitrocellulose membrane. Proteins were then cross-linked to membranes by
UV treatment and stained with ponceau S solution for 3 minutes for protein loading control before
incubation in blocking buffer (5% nonfat dry milk in 0.05% PBS-Tween 20 [PBS-T]) for 1-hour
rocking at room temperature. Blots were then incubated overnight at 4°C in blocking buffer
containing primary antibody of interest; anti-MRP5 and anti-MRP9 rat monoclonal antibodies are
always used at a concentration of 1:200 unless specified otherwise. The following day blots are
rinsed once quickly (2 min) and then washed three times (15 min) in PBS-T and subsequently
incubated for one hour with horseradish peroxidase (HRP)-conjugated goat anti-rat IgG
secondary antibody at 1:20,000 (Invitrogen cat. number 31470). Blots were then rinsed again
once quickly, followed by three washes (20 min) with PBS-T and then developed using enhanced
chemiluminescence (SuperSignal West Pico PLUS Reagent, Pierce, cat. number 34580) and
detected using ChemiDocTM Imaging Systems (Bio-Rad). If subsequent re-probing was
necessary, blots were then stripped with 500 mM NaOH for 10min and rinsed quickly with DI H_2O
twice, and then PBS-T three times (5min) before blocking again for one hour to repeat the
process with subsequent antibodies.

Subcellular fractionation and MAM isolation
Mitochondrial Associated Membranes (MAMs) isolation and subcellular fractionizations were
based off of Wieckowski et al 2009 Nature Methods Protocol with minor modifications for
optimization of testes tissue. Age matched wildtype and double knock out male mice were
anesthetized with intraperitoneal injection of ketamine/xylene and then systemic perfusion was
performed via standard cardiac puncture method. Perfusion was performed with IB-1 buffer (225
mM mannitol, 75 mM sucrose, 0.5% BSA, 0.5 mM EGTA and 30 mM Tris–HCl pH 7.4) using 25
ml total pushed through a 30 ml syringe with a 26G needle (BD Precision Glide, cat. number
305111) into the left ventricle of the beating heart, which upon completion resulted in completely
cleared liver and kidneys. Livers and testes were then immediately harvested and placed in IB-3
buffer (225 mM mannitol, 75 mM sucrose and 30 mM Tris–HCl pH 7.4) on ice at 4°C and the
remaining isolation was performed entirely at 4°C in a cold room. A single liver or four pooled
testes were necessary to achieve sufficient yield of purified MAMs, which were minced and
combined and then lysed in IB-1 buffer via 10 strokes in a 22 gauge 5ml glass duall dounce
homogenizer. The homogenate was then spun down twice at 740 x g to remove nuclei and
cellular debris. The remaining supernatant was spun at 9,000 x g for the isolation of crude
mitochondria, which was subsequently washed and spun down twice at 10,000 x g. The first
10,000 x g spin was in IB-2 buffer (225 mM mannitol, 75 mM sucrose, 0.16% BSA and 30 mM
Tris–HCl pH 7.4) and then the second in IB-3 for further purification prior to a 100,000 x g spin
through a Percoll (Sigma, cat. number P1644) gradient to separate MAMs from mitochondria. The
original supernatant from the 9,000 x g spin was also spun down at 100,000 x g to concentrate
the remaining smaller organelles, plasma membranes and cytosolic proteins labeled "Membrane
Prep." Each of the isolated fractions "740 x g", "Membrane Prep", "Mitochondria" and "MAMs" as
well as total initial homogenate were subsequently lysed in lysis buffer and following SIM, ran
side by side in a Criterion Precast Gradient 4-15% SDS-PAGE gel for western blotting probing of
key intracellular organellar markers and proteins of interest: HSP60 (Abcam, cat. number
ab46798), Calnexin (Abcam, cat. number ab22595), Mitofusin 2 (Invitrogen 7H42L13, cat.
number 702768), MRP5 and MRP9.

Histology and immunohistochemistry
Paraffin-embedded tissues were sectioned at 4 μm and processed for immunohistochemistry
(IHC) as previously described in Pek et al with minor modifications (92). Slides were first
rehydrated in descending gradient from pure xylene through ethanol series (100%, 95%, 80%,
70%) and followed by endogenous HRP quenching before heat-induced epitope retrieval in
citrate buffer pH 6 (Target Retrieval Solution, DAKO, cat. number S1700). After epitope retrieval,
sections were then rinsed and blocked with avidin, biotin and protein blocking following Vector
labs ABC elite staining kit (Vector labs, cat. number PK-6104) prior to incubation with monoclonal
rat anti-MRP9 antibodies at 1:50 overnight at 4°C. The next day sections were then incubated
with secondary biotinylated anti-rat antibody for 30 minutes at room temperature. ABC tertiary
reagent was added and then signals were detected by DAB substrate incubation per
manufacturer recommendations (DAB/Metal Concentrate Peroxidase Kit. Thermo Fischer, cat.
number 1859346 and 1856090). Finally, slides were lightly counterstained with hematoxylin and
then dehydrated in opposite ascending series and mounted with DPX Mountant (Sigma, cat.
number 06522-100).

Heme quantification
To extract heme from perfused mouse tissues, a minimum of 30 mg of caudal epididymides
(either three or four pooled together), an entire testis or seminal vesicle were dounce
homogenized fresh in a minimum of five volumes of lysis solution containing 50 mM Tris/HCl pH
8.0, 5 mM $CaCl_2$, 50 mM NaCl, 1% Triton X-100 and 0.5% Proteinase K. Proteinase K was added
fresh to the buffer and the homogenate was incubated overnight at 37°C with moderate shaking.
The next day the Proteinase K digest is sonicated for 1 min (10 W, pulse 0.5 sec), centrifuged at
11,000 x g for 45 minutes and then the supernatant was collected for heme quantification. 20 µl
of the tissue lysate sample is mixed with 180 µl of 2 M oxalic acid and sample tubes are then

heated in a water bath set at 100°C for 30 minutes while a duplicate set of samples remain at
room temperature as blank controls. Fluorescence of porphyrins were then read using
Ex400/10nm, Em645/40nm on a BioTek microplate reader (BioTek, Synergy HT) and heme
content calculated based off of a heme standard curve and subtraction of blank values (parallel
unheated samples in oxalic acid) from samples.

EIF2α phosphorylation quantification
Eight age matched wildtype and double knock out male mice were sacrificed by cervical
dislocation, and whole testes were dissected out and rinsed in ice cold PBS. A single testis was
then homogenized in 500 µl standard lysis buffer with 5x protease inhibitors and 3x phosphatase
inhibitors, via 8 strokes in a 20 gauge 2 ml glass duall dounce homogenizer, transferred to 1.5 ml
Eppendorf tube and placed on ice for 15 minutes, vortexing for 15 seconds every 5 minutes.
Afterward, the homogenate is then sonicated for 10 seconds, twice, prior to proceeding with
standard clarifying spin for standard immunoblotting. For all samples, 150 µg of protein was then
separated on Criterion Precast Gradient 4-15% SDS-PAGE gels (Bio-Rad, cat. number 5671083)
and probed for phosphorylated EIF2α protein levels with rabbit anti-Eif2a S51 phosphorylation
polyclonal antibody (1:500 dilution) (Cell Signaling Technologies, cat. number D9G8) and Total
EIF2α protein levels with rabbit anti-EIF2a polyclonal antibody (1:10,000 dilution) (Cell Signaling
Technologies, cat. number D7D3). Additional blots were probed with rat anti-MRP5 NC3
monoclonal antibody (1:200) and rat anti-MRP9 M9I-27 monoclonal antibody (1:200).
Quantification of EIF2α S51 phosphorylation was performed by Image Lab 6.1 software (BioRad)
and normalized to total EIF2α protein.
565

Mouse reproduction studies

Breeding schema for general fecundity and fertility
All breeding studies were performed in harem setups with one stud male mouse and two female
mice per cage. These harems were monitored over a period of at least 6 months to track
breeding of ABCC5/ABCC12 double knockout mice (-/-, -/-) as well as the requisite control
intercrosses of heterozygous-homozygous (+/-, -/-) and homozygous-heterozygous (-/-, +/-) mice.
Once successful pregnancies were determined, females were separated from males and
individually housed for monitoring fecundity. New female mice were then rotated in the male mice
cages to maintain setups of two female mice per stud male. Males unable to produce any
pregnancies after two months were removed and replaced. Pregnant dams were monitored at
least twice daily for birth of pups and overall health, litter size was recorded at birth and any
subsequent mortality logged through weaning was reported. Fecundity was calculated based off
the number of breeding setups required to achieve same number of pregnancies as well as the
number of offspring per litter and average productivity of litters per harem breeding setup. All
genetic segregation and statistics for MRP5 and MRP9 alleles were calculated on weaned P21
mice.

***In vitro* fertilization (IVF) and general sperm characterization**
All IVF was conducted in HTF media (EmbryoMax, Sigma, cat. number MR-070), supplemented
with 3 mg/ml of BSA. Males were mated with females and assessment of mating was determined
by the presence of a vaginal plug. Males that were listed as plug positive were allowed to rest a
total of three days prior to IVF. Spermatozoa was collected from individual males and placed in
90 µl of TYH media supplemented with beta-methyl-cyclodextrin to induce capacitation (93).
Sperm samples were allowed to capacitate for a total of 45-60 minutes. A 10 µl sample was
diluted in 1400 µl of IVF media and assessed on an IVOS system (Hamilton Thorne CASA -
Computer Assisted Sperm Analyzer). Total sperm concentration, percent progressive motility and
percent rapid motile cells was calculated. Visual assessment of morphology was also assessed,
and IVF was conducted and standardized to an insemination dose of 1.0×10^6 sperm/ml for each
male processed. Oocytes and sperm were incubated for a total of 5 hours. Presumptive zygotes

were rinsed through HTF media to remove accessory sperm and cultured overnight in KSOM
media plus amino acids (94). The next morning, total number of two cell embryos were
determined and a fertilization rate calculated based on the number of two cell embryos per total
number of oocytes inseminated.

Sperm "swim-out" collection
Age matched male mice were sacrificed by cervical dislocation, and entire reproductive tracts
were dissected out intact to isolate the caudal epididymis. Caudal epididymides (CE) were
trimmed away from fat pads and the testes, rinsed once quickly (dipping) in DPBS and then
dabbed dried (single touch) on a kimwipe before being placed in a 35 mm dish with mineral oil.
Prewarmed DPBS with added 100 mg/L $MgCl_2$ and $CaCl_2$ (swim-out solution) to 37°C was then
added to the mineral oil in close proximity to the CEs for sperm swim-out. Swim-out was achieved
by snipping the CEs with dissecting micro-scissors and dragging the exposed globule with a 22G
needle from the mineral oil into the swim-out solution. The entire petri dish was then covered and
then placed in the 37°C incubator for 30min to allow for purified sperm to fully swim-out into the
solution. Then sperm were collected with a 1ml pipettor and spun down gently in a 1.5 ml
Eppendorf tube at 333 $x\,g$ for 3 minutes, followed by resuspension in fresh swim-out buffer and
rocked gently for 2 minutes. This rinsing was repeated the same way again twice, and after the
third rinse and spin down the sperm are ready for downstream analysis.

Sperm transmission electron microscopy (TEM) imaging
Swim-out collected sperm were resuspended in a final 2% PFA and 1% Glutaraldehyde solution
for fixation, and then rinsed in 2 changes of 0.1 M Sodium Cacodylate buffer with 2.4% sucrose
and 8 mM $CaCl_2$. Samples were then postfixed in 2% Osmium Tetroxide in 0.1 M Sodium
Cacodylate buffer with 2.4% sucrose and 8 mM $CaCl_2$. Samples were rinsed thoroughly with
distilled water, and then en bloc stained in saturated aqueous Uranyl Acetate. Samples were then
dehydrated through a graded ethanol series (50%,70%, 95%, 95%, 100%, 100%, 100%, 100%).
Samples were transitioned into absolute Acetone for 3 changes. Then infiltrated in a 1:1 (plastic:
acetone) and then 3:1 (plastic: acetone) dilutions of Embed 812 and followed with 3 changes of
100% plastic prior to embedding. Samples were cure 24 hours in a 60°C oven. Once removed
from the oven samples were thick sectioned with glass knives to determine best area to thin
section, on a Leica EM UC6 Ultramicrotome. A diamond knife (Diatome) was used to thin section
samples and sections were placed on 150 hex copper grids. Grids were stained sequentially with
Aqueous Saturated Uranyl Acetate followed by Reynold's Lead Citrate. Sections were examined
at 120 KV on a JEOL 1400 plus Transmission Electron Microscope at 6000x magnification and
digital images were taken on a GATAN Orius camera. Representative cross-sectional planes of
the sperm midpieces were taken for an n > 15 FOVs for each animal sample.

Sperm JC1 staining and flow cytometry
JC-1, a fluorescent, cationic, lipophilic dye, was used to demonstrate differences in mitochondrial
membrane potential (MitoProbeTM JC-1 Assay Kit for Flow Cytometry, Invitrogen cat. number
M34152). In brief, sperm collected fresh from swim-out were resuspended in 500 µM swim-out
solution with final concentration of 3 µM JC-1 dye and gently rocked at 37°C for 20 minutes in the
dark prior to two rounds of rinses (333 $x\,g$ for 3 minutes). Per manufacturer instructions, CCCP
(carbonyl cyanide m-chlorophenyl hydrazine) 25 µM final concentration was used as positive
control to disrupt mitochondrial potential and to confirm that the JC-1 response was indeed
sensitive. JC-1 exhibits potential-dependent accumulation of red fluorescent J-aggregates in
mitochondria, indicated by a fluorescence emission shift from green (~529 nm) to red (~590 nm).
In this way, mitochondrial depolarization can be inferred by a decrease in the red/green
fluorescence intensity ratio, so sperm were analyzed using a BD Biosciences FACSCanto II and
gated for green (FITC) and red (PE) fluorescence from a 488 laser. Percentage of sperm with
defective mitochondrial membrane potential FITC-pos PE-neg were gated from a minimum of

200,000 events per replicate. Representative sperm from each gating strategy were also sorted
directly onto microscope slides using BD Biosciences FACSAria II and imaged.
645

Mammalian cell culture methods
All mammalian cells were cultured continuously at 37°C in a humidified incubator with 5% CO_2.
Immortalized fibroblasts were generated from mouse embryos isolated at E12.5 by sacrificing
pregnant females and culturing the mouse embryonic fibroblasts (MEFs) as described previously
(14, 95). Once established, these primary cells from wildtype, MRP9 single knockout and double
knockout mice were immortalized by retroviral infection of MEFs with conditioned media from Ψ2-
U195 cells producing the SV40 large T antigen. For overexpression in cell culture, plasmids for
either *ABCC5* or *ABCC12* constructs were transfected into cell lines using PolyJet transfection
reagent (SignaGen Laboratories, cat. number SL100688). Previously, human *ABCC5* ORF was
cloned into pcDNA3.1(+)zeo and pEGFP-N1 plasmids for mammalian expression (14). Likewise,
human *ABCC12* ORF was cloned into pcDNA3.1(+)zeo via EcorI and XbaI in a similar manner
and subsequently epitope tagged with FLAG. Cells that were transfected with either vector or
MRP expressing constructs could then be harvested up to 48 hours later for western blotting
(protein expression), microscopy (localization) or other downstream assays. For attempting
induction of endogenous expression of MRP5 or MRP9 in cell lines possibly capable of
production (i.e. T-47D, A549, HCT116 and GC-2spd), cells were treated for 24 hours with 2 mM
Sodium Butyrate (NaB) following methods from Bin Shi et al 2020 and then analyzed for
expression and subcellular localization (72). For western blotting, cells were harvested in
standard lysis buffer and followed standard immunoblotting procedure as described above. For
polarization of MDCKII cells, trypsinized cells were seeded onto transwell cell culture inserts
(Corning, COSTAR Polystyrene 6 well plates 24mm inserts, cat. number 3450) and allowed to
grow to a monolayer in standard growth media (DMEM supplemented with 10% FBS) changed
daily. Prior to reaching confluence, MDCKII cell's monolayer integrity was determined daily by
measuring the transepithelial electrical resistance (TEER) using an EVOM^2 Epithelial
Voltohmmeter device (World Precision Instruments, cat. number EVOM2). MDCKII cells were
considered polarized and used for imaging when TEER values exceeded 200 Ω/cm^2 resistance
and were validated by checking for markers of polarization. Immunofluorescence was performed
as follows: cells seeded onto coverslips (or transwells) were rinsed with ice cold DPBS (and
subsequently in between each additional step) and immediately fixed in freshly prepared 4% PFA
for greater than 30 minutes; cells were then quenched with 0.1 M ethanolamine for 5 minutes,
twice; permeabilized with 0.1% Triton-X 100 in DPBS solution for 10 minutes; and blocked in
blocking solution of 3% BSA:Superblock (SuperBlock Buffer in PBS, Thermo Scientific, cat.
number 37515) for greater than one hour at room temperature or overnight at 4°C; incubated with
primary antibody (rat monoclonal anti-MRP9 M9I-38, 1:10 in blocking solution) overnight at 4°C;
secondary antibody (Alexa fluorophore-conjugated goat anti-rat, 1:2000 in blocking solution) for 1
hour at RT; and subsequently counterstained with DAPI (1:30,000 of 5 mg/ml) for 3 minutes and
mounted using Antifade reagent (ProLong Gold Antifade Reagent, Invitrogen, cat. number
P36934). Images were taken and processed using an Airyscan980 SR confocal microscope
(Zeiss).
685

Metabolomics methods

Global untargeted metabolomics of mouse testes
Tissue was first extracted either in aqueous or organic methods. For aqueous extractions, 10
volumes of methanol:H_2O (3:1) was added to the tissue sample and homogenized with ceramic
beads (Precellys®24-Dual, PeqLab). Then 200 μl of the homogenate was transferred to a new
tube where it was then vortexed and centrifuged at 15,000 RPM for 10 min at 4°C. The
supernatant was transferred to sample vial for ultra high performance liquid chromatography
coupled with mass spectrometry (UHPLC-MS) analysis. Analytes were separated with either a

CSH C18 column or a BEH amide column for more nonpolar and polar species, respectively. For
the first aqueous method, separation was achieved using an ACQUITY UPLC CSH C18 Column
(1.7 µm, 2.1 mm X 100 mm). Mobile phase A was water (contained 0.1% formic acid) and mobile
phase B was acetonitrile (contained 0.1% formic acid). The gradient was: 0 to 1 min, 5% B; 1.1 to
10 min, 5% to 95 % B; 10.1 to 13 min, 95 % B; 13.1 to 13.5 min, 95 to 5% B; 13.6 to 15 min, hold
at 5% B. The flow rate was 0.5 mL/min. The column was maintained at 50°C and the auto-
sampler was kept at 10°C. A 5 µL injection was used for all samples. For the second method of
aqueous analysis, the separation was achieved using an ACQUITY UPLC BEH amide Column
(1.7 µm, 2.1 mm X 100 mm). Mobile phase A was water (contained 0.1% formic acid) and mobile
phase B was acetonitrile (contained 0.1% formic acid). The gradient was: 0 to 1 min, 99% B; 1 to
10 min, 99% to 30 B; 10.1 to 12 min, hold at 99% B. The flow rate was 0.5 mL/min. The column
was maintained at 45°C and the auto-sampler was kept at 10°C. A 5 µL injection was used for all
samples. For both aqueous sample preps, data was acquired in both positive (HDMSe) and
negative (HDMSe) mode. The capillary voltages were separated for positive (0.8 kV) and
negative (0.8 kV) and sampling cone voltage was 40 V. Nitrogen at a flow of 800 L/h was used as
the desolvation gas with a constant desolvation temperature of 500°C. The source temperature
was set at 150°C. Data were acquired over the m/z range of 50-1000. For organic extraction
(lipid) methods, likewise 10 volumes of methanol:H_2O (3:1) was added to the tissue sample and
homogenized with ceramic beads (Precellys®24-Dual, PeqLab). However, now after
homogenization 400 ul was transferred to a new tube and 500 µL of ice-cold methyl-tert-butyl
ether (MTBE) is added, prior to incubation at 650 RPM for 1 hour. Additionally, 500 µL of ice-cold
water was then added and incubated at 650 RPM for 15 min before phase separation was
completed by centrifugation for 8 min at 8,000 RPM at 4°C. The top (organic) layer was removed
and dried at room temperature under nitrogen. The recovered lipids were reconstituted in 200 µL
of isopropanol:acetonitrile:water (2:1:1, v/v/v) and were transferred to sample vial for UPLC-MS
analysis. The separation was achieved using a CORTECS HILIC Column (2.7 µm, 2.1 mm X 100
mm). Mobile phase A was water/acetonitrile (5:95, v/v) with 10 mM ammonium acetate and
mobile phase B water/acetonitrile (50:50, v/v) with 10 mM ammonium acetate. The gradient was
ramped from 0.1% to 20% B in 10 min, ramped to 80% B in 3 min, ramped back down to 0.1% B
and held for 3 min. The flow rate was 0.5 mL/min. The column was maintained at 30°C and the
auto-sampler was kept at 10°C. A 5 µL injection was used for all samples. Like the aqueous
phase, data was acquired in both positive (HDMSe) and negative (HDMSe) mode. The capillary
voltages were separated for positive (2.8 kV) and negative (1.9 kV) and sampling cone voltage
was 30 V. Nitrogen at a flow of 900 L/h was used as the desolvation gas with a constant
desolvation temperature of 550°C. The source temperature was set at 120°C. Data were
acquired over the m/z range of 100-1500. Positive and negative mode data were basally
analyzed by Progenesis QI and putative metabolite identification was searched by using HMDB
and LIPID MAPS databases (delta < 5 ppm). The output marker lists were then manually sorted
and annotated for more sophisticated downstream analysis with MetaboAnalyst (96-98).
Multivariate data analysis, volcano plots, Variable Importance in Projection (VIP) heat maps and
mummichog algorithm analysis to putatively annotate metabolites were performed using
MetaboAnalyst 4.0 (35).

Measurement of retinoids from reproductive tissues
Two separate analyses were performed to determine the retinoid profile in wild type and double
knockout mice. Testes and seminal vesicles were assayed for retinoic acid isomers and retinol
and total retinyl esters. Tissue samples were stored in -80°C until processed. Only glass
containers, pipettes, and syringes were used to handle retinoids. Extraction of retinoids was
performed under yellow lights using a two-step liquid-liquid extraction that has been described in
detail previously using 4,4-dimethyl-RA as an internal standard for RA and retinyl acetate as an
internal standard for retinol and total retinyl ester (99-102). Briefly, for the extraction of retinoids,
tissue was homogenized in 1 mL of 0.9% NaCl (normal saline) and to each homogenate aliquot,
3 mL of 0.025 M KOH in ethanol was added to the homogenate followed by addition of 10 mL
hexane to the aqueous ethanol phase. The samples were vortexed and centrifuged for 1 to 3 min

at 1,000 rpm in a Dynac centrifuge (Becton Dickinson) to facilitate phase separation and pellet
precipitated protein. The hexane (top) phase containing nonpolar retinoids [retinol and total retinyl
esters (RE)] was removed. Then 4 M HCl (200 µL) was added to the remaining aqueous ethanol
phase, samples were vortexed, and then polar retinoids (RA) were removed by extraction with a
second 10 mL aliquot of hexane as described above. Organic hexane phases were evaporated
under nitrogen while heating at approximately 30°C in a water bath (model N-EVAP 112,
Organomation Associates, Berlin, MA, USA). All samples were resuspended in 60 µL acetonitrile.
Levels of RA were determined by liquid chromatography-multistage tandem mass spectrometry
(LC-MRM[3]) which is an LC-MS/MS method utilizing two distinct fragmentation events for
enhanced selectivity (99). RA was measured using a Shimadzu Prominence UFLC XR liquid
chromatography system (Shimadzu, Columbia, MD) coupled to an AB Sciex 6500+ QTRAP
hybrid triple quadrupole mass spectrometer (AB Sciex, Framingham, MA) using atmospheric
pressure chemical ionization (APCI) operated in positive ion mode as previously described (99).
For the LC separation, the column temperature was controlled at 25°C, the autosampler was
maintained at 10°C and the injection volume was typically 20 µL. All separations were performed
using an Ascentis Express RP-Amide guard cartridge column (Supelco, 50 × 2.1 mm, 2.7 µm)
coupled to an Ascentis Express RP-Amide analytical column (Supelco, 100 × 2.1 mm, 2.7 µm).
Mobile phase A consisted of 0.1% formic acid in water, and mobile phase B consisted of 0.1%
formic acid in acetonitrile. Endogenously occurring retinoid isomers including all-trans-retinoic
acid (RA), 9-*cis* retinoic acid, 13-*cis* retinoic acid, and 9,13-di-*cis* retinoic acid are resolved using
a gradient separation at a flow rate of 0.4 mL min^{-1} with gradient conditions described previously
(99). The APCI source conditions and MRM[3] detection parameters were as previously described
where the MRM[3] transition for RA was m/z 301.1 → m/z 205.1 → m/z 159.1 and for 4,4-
dimethyl RA was m/z 329.2 → m/z 151.2 → m/z 100.0 (99).
Levels of retinol and total retinyl esters (RE) were quantified via HPLC-UV according to previously
published methodology (102, 103). Note: total retinyl esters is 90+% retinyl palmitate. Retinol and
RE were resolved by reverse-phase chromatography (Zorbax SB-C18, 4.6 × 100 mm, 3.5 µm) on
a Waters Acquity UPLC H-class system and were quantified by UV absorbance at 325 nm.
Analytes were separated at 1 mL min^{-1} with a gradient separation as described previously with a
typical injection volume of 30 µL. The amount of retinoic acid (RA), retinol (ROL) and total retinyl
ester (RE) was normalized per gram of tissue.

Targeted metabolomics of subcellular fractionation of testes
High-Throughput Targeted Metabolomics were performed on subcellular fractions isolated from
WT and DKO testes. This subcellular fractionation was performed as described above with minor
modifications, generating "Total Homogenate", "740 *x g*", "Membrane Prep", "Mitochondria" and
"MAMs" fractions. In order to maintain compatibility with downstream mass spectrometry analysis,
each fraction was collected in ice cold 0.9% saline for its final centrifugation and subsequently
was rinsed and diluted in ice cold 0.9% saline instead of lysis buffer prior to freezing. The
concentration of metabolites and lipids were then determined with mass spectrometry using
MxP® Quant 500 kit (Biocrates Life Sciences AG, Innsbruck, Austria). The kit enables
quantification of approximately 630 endogenous metabolites belonging to 26 different biochemical
classes: alkaloids (1), amine oxides (1), amino acids (20), amino acids related compounds (30),
bile acids (14), biogenic amines (9), carbohydrates (1), carboxylic acids (7), cresols (1), fatty
acids (12), hormones (4), indoles and derivatives (4), nucleobases and related compounds (2),
vitamins (1), acylcarnitines (40), lysophosphatidylcholines (14), phosphatidylcholines (76),
sphingomyelins (15), ceramides (28), dihydroceramides (8), hexosylceramides (19),
dihexosylceramides (9), trihexosylceramides (6), cholesteryl esters (22), diglycerides (44),
triglycerides (242). The method combines direct infusion coupled with tandem mass spectrometry
(DI – MS/MS) and reverse phase liquid chromatography coupled with tandem mass spectrometry
workflows (LC – MS/MS). Extraction was carried out per manufacturer's instructions. Tissue
samples were homogenized with ceramic beads (Precellys®24-Dual, Bertin Technologies SAS,
France) prior to analysis. Briefly, mitochondrial fractions and tissue homogenates were thawed
over ice and 30 µl of the sample was loaded onto the filter plate in 96 well format, impregnated

with deuterated internal standards and dried under a stream of nitrogen in a positive pressure
manifold. A 5% solution of phenylisothiocyanate in ethanol:water:pyridine (1:1:1, v:v:v) was added
for derivatization of metabolites. Extraction was done with 5 mM ammonium acetate in methanol.
The extracts were eluted with positive pressure manifold, followed by dilution steps for LC-MS/MS
and DI-MS/MS acquisitions. The tandem mass spectrometry platform consisted of a Waters I
Class Ultra performance liquid chromatography (UPLC) coupled to a Waters TQ-XS tandem
quadrupole mass spectrometer (Waters Corporation, Milford, MA). MetIDQ software (Biocrates
Life Sciences AG, Innsbruck, Austria) was used to register the samples, analyze the data and
validate the assay. All reagents used were of analytical grade and mobile phases used were of
LC-MS grade.

Transcriptomics and associated methods

RNAseq of mouse testes
RNA was extracted from frozen tissues following standard Trizol chloroform protocol (Invitrogen,
cat. number 15596026) and aqueous fractions were then taken directly into Qiagen RNA plus
extraction kit (Qiagen, cat. number 74136) for maximum purity. Total RNA quality and quantity
were tested on an Agilent Bioanalyzer 2100 System (Agilent cat. number G2939BA) using RNA
Nano Chip On-Chip Electrophoresis trays for samples with RIN scores greater than 8 prior to
processing for library prep/construction and sequencing. RNA-seq was performed at NISC using
Illumina NovaSeq6000. FastQC, version 0.11.7, was used as an additional bioinformatics quality
control on output reads from sequencing. The reads were then aligned to the mouse mm10
GRCh38 reference genome for gene annotation using STAR (Spliced Transcripts Alignment to a
Reference), version 2.7.0f, and numbers of reads in transcripts per million (TPM) mapped to
genes were counted generating matrixes for all samples. Differentially expressed genes were
then identified via both manual export, sorting and fold change annotation, as well as processed
via DESeq2, version 1.12.3, for more complex analysis. DESeq2 was run using R (version 3.6.1)
and Bioconductor (version 3.4) with BioInstaller (version 1.24.0) for volcano plotting and statistical
analysis of differential changes utilizing adjusted *P* value < 0.01 and False Discovery Rate (FDR)
cutoffs of 0.05. The complete RNA sequencing results have been depositing to the NCBI Gene
Expression Omnibus under accession under GSE176740.

Pathway analysis
Gene Ontology (GO) enrichment and network analysis of RNAseq results were performed using
both Ingenuity Pathway Analysis (IPA) software package (Qiagen version 1-16) and manually
developed excel pipeline workflows. For IPA analysis, statistically significant gene lists by tissue
were manually filtered for minimum threshold of 10 TPM, sorted by *P* value and uploaded along
with gene expression fold change for analysis. For pipeline pathway analysis, gene lists for GO
terms and associated pathways of interest were downloaded from MGI (104-106), RGD (107) as
well as manually added from literature review of relevant publications. These all-encompassing
lists were then used as target gene IDs for filtered call look up functions from RNAseq gene
expression RSEM matrixes. In brief, genes with matching IDs which passed threshold of
expression in at least one sample condition and were statistically significant, were output in a
heatmap with *P* value and relative fold change displayed.

Motif analysis
The MEME Suite and HOMER motif platforms were used to unbiasedly identify conserved motifs
via discovery analysis of enriched sequences from differentially expressed genes. More
specifically, we generated gene lists (either globally or mitochondria targeted) and divided them
into upregulated and downregulated genes and extracted their associated sequence FASTA files.
Then 1000bp upstream of ORF and 50bp downstream were used to run MEME or HOMER via
Perl scripts on NIH High Performance Cluster (HPC) to discover the top 5 ungapped enriched
putative motif sequences. These sequences were then further analyzed with the web-based suite

tools TomTom and GoMo for MEME motif findings and manually curated for motif verification,
which were further validated by checking and annotating output from the HOMER suite.

Additional Bioinformatics and statistical analyses
All alignment and evolutionary analyses were conducted using MEGA X software version 10.2.4
(108-111). All data throughout are expressed as means ± SEM unless otherwise stated. Means of
groups were assessed by using two-sided, unpaired student's t-test or the equivalent analyses
where applicable unless otherwise stated. A *P* value of < 0.05 when compared to baseline values
or genotypes (WT) was considered statistically significant. All analyses were performed using
Prism 9 software (GraphPad, version 9.1.0).
858

859 **Acknowledgments**

860
We would like to acknowledge Ms. Lisa Garrett for conducting the pronuclear injections at
NHGRI, NIH to generate the *ABCC12* mouse. We also thank Mr. Simon Beardsley for sharing the
C. elegans mrp-5 data. We acknowledge the Imaging Core Facility in the Department of Cell
Biology and Molecular Genetics at the University of Maryland, College Park supported by Award
Number 1S10OD025223-01A1 from the National Institutes of Health. This work was also done in
part by utilizing the computational resources of the National Institutes of Health HPC Biowulf
cluster (http://hpc.nih.gov). The MRP5 (M5I-10, M5-NC3) and MRP9 (M9I-27, M9I-38, M9II-3)
antibodies used throughout were a generous gift from Piet Borst / Koen van de Wetering
(Netherlands Cancer Institute). This work was supported by funding from the National Institutes of
Health DK85035 and DK125740 (IH); HD077260 (MAK); the Utah Center for Iron and Heme
Disorders was supported by funding from DK110858 (JP); and the Intramural Program of the
National Human Genome Research Institute (DB). The funders had no role in study design, data
collection and analysis, decision to publish, or preparation of the manuscript.
874
875

References

1. M. Kool *et al.*, Analysis of expression of cMOAT (MRP2), MRP3, MRP4, and MRP5,
homologues of the multidrug resistance-associated protein gene (MRP1), in human
cancer cell lines. *Cancer Res* **57**, 3537-3547 (1997).
2. M. A. McAleer, M. A. Breen, N. L. White, N. Matthews, pABC11 (Also Known as MOAT-C
and MRP5), a Member of the ABC Family of Proteins, Has Anion Transporter Activity but
Does Not Confer Multidrug Resistance When Overexpressed in Human Embryonic
Kidney 293 Cells. *Journal of Biological Chemistry* **274**, 23541-23548 (1999).
3. P. Borst, R. Evers, M. Kool, J. Wijnholds, A family of drug transporters: the multidrug
resistance-associated proteins. *J Natl Cancer Inst* **92**, 1295-1302 (2000).
4. G. D. Kruh, M. G. Belinsky, The MRP family of drug efflux pumps. *Oncogene* **22**, 7537-
7552 (2003).
5. P. M. Jones, A. M. George, The ABC transporter structure and mechanism: perspectives
on recent research. *Cellular and Molecular Life Sciences CMLS* **61**, 682-699 (2004).
6. A. J. Slot, S. V. Molinski, S. P. Cole, Mammalian multidrug-resistance proteins (MRPs).
Essays Biochem **50**, 179-207 (2011).
7. J.-Y. Lee, D. M. Rosenbaum, Transporters Revealed. *Cell* **168**, 951-953 (2017).
8. B. Sarkadi, M. Müller, Z. Holló, The multidrug transporters—proteins of an ancient
immune system. *Immunology Letters* **54**, 215-219 (1996).
9. I. B. Holland, M. A. Blight, ABC-ATPases, adaptable energy generators fuelling
transmembrane movement of a variety of molecules in organisms from bacteria to
humans. *Journal of Molecular Biology* **293**, 381-399 (1999).
10. S. Cole *et al.*, Overexpression of a transporter gene in a multidrug-resistant human lung
cancer cell line. *Science* **258**, 1650-1654 (1992).
11. S. P. C. Cole, Targeting Multidrug Resistance Protein 1 (MRP1, ABCC1): Past, Present,
and Future. *Annual Review of Pharmacology and Toxicology* **54**, 95-117 (2014).
12. C. A. Ritter *et al.*, Cellular export of drugs and signaling molecules by the ATP-binding
cassette transporters MRP4 (ABCC4) and MRP5 (ABCC5). *Drug Metab Rev* **37**, 253-278
(2005).
13. R. G. Deeley, C. Westlake, S. P. Cole, Transmembrane transport of endo- and xenobiotics
by mammalian ATP-binding cassette multidrug resistance proteins. *Physiol Rev* **86**, 849-
899 (2006).
14. T. Korolnek, J. Zhang, S. Beardsley, G. L. Scheffer, I. Hamza, Control of metazoan heme
homeostasis by a conserved multidrug resistance protein. *Cell Metab* **19**, 1008-1019
(2014).
15. G. Jedlitschky, B. Burchell, D. Keppler, The multidrug resistance protein 5 functions as an
ATP-dependent export pump for cyclic nucleotides. *J Biol Chem* **275**, 30069-30074
(2000).
16. G. Reid *et al.*, Characterization of the Transport of Nucleoside Analog Drugs by the
Human Multidrug Resistance Proteins MRP4 and MRP5. *Molecular Pharmacology* **63**,
1094-1103 (2003).
17. R. S. Jansen, S. Mahakena, M. de Haas, P. Borst, K. van de Wetering, ATP-binding
Cassette Subfamily C Member 5 (ABCC5) Functions as an Efflux Transporter of
Glutamate Conjugates and Analogs. *J Biol Chem* **290**, 30429-30440 (2015).
18. X. Yuan *et al.*, Regulation of intracellular heme trafficking revealed by subcellular
reporters. *Proc Natl Acad Sci U S A* **113**, E5144-5152 (2016).

19. R. K. Donegan, C. M. Moore, D. A. Hanna, A. R. Reddi, Handling heme: The mechanisms
underlying the movement of heme within and between cells. *Free Radical Biology and*
Medicine **133**, 88-100 (2019).
20. I. G. Chambers, M. M. Willoughby, I. Hamza, A. R. Reddi, One ring to bring them all and
in the darkness bind them: The trafficking of heme without deliverers. *Biochim Biophys*
Acta Mol Cell Res **1868**, 118881 (2021).
21. A. R. Reddi, I. Hamza, Heme Mobilization in Animals: A Metallolipid's Journey. *Accounts*
of Chemical Research **49**, 1104-1110 (2016).
22. J. Tammur *et al.*, Two new genes from the human ATP-binding cassette transporter
superfamily, ABCC11 and ABCC12, tandemly duplicated on chromosome 16q12. *Gene*
273, 89-96 (2001).
23. T. K. Bera *et al.*, MRP9, an unusual truncated member of the ABC transporter
superfamily, is highly expressed in breast cancer. *Proc Natl Acad Sci U S A* **99**, 6997-7002
(2002).
24. N. Ono *et al.*, Multidrug resistance-associated protein 9 (ABCC12) is present in mouse
and boar sperm. *Biochem J* **406**, 31-40 (2007).
25. H. Shimizu *et al.*, Characterization of the mouse Abcc12 gene and its transcript encoding
an ATP-binding cassette transporter, an orthologue of human ABCC12. *Gene* **310**, 17-28
(2003).
26. M. Jung *et al.*, Unified single-cell analysis of testis gene regulation and pathology in five
mouse strains. *Elife* **8** (2019).
27. A. Amaral, B. Lourenço, M. Marques, J. Ramalho-Santos, Mitochondria functionality and
sperm quality. *REPRODUCTION* **146**, R163-R174 (2013).
28. N.-H. Gu, W.-L. Zhao, G.-S. Wang, F. Sun, Comparative analysis of mammalian sperm
ultrastructure reveals relationships between sperm morphology, mitochondrial
functions and motility. *Reproductive Biology and Endocrinology* **17**, 66 (2019).
29. D. Stojanovski, M. Bohnert, N. Pfanner, M. van der Laan, Mechanisms of Protein Sorting
in Mitochondria. *Cold Spring Harb Perspect Biol* **4** (2012).
30. J. Lee, D. H. Kim, I. Hwang, Specific targeting of proteins to outer envelope membranes
of endosymbiotic organelles, chloroplasts, and mitochondria. *Frontiers in Plant Science* **5**
(2014).
31. J. E. Vance, MAM (mitochondria-associated membranes) in mammalian cells: lipids and
beyond. *Biochim Biophys Acta* **1841**, 595-609 (2014).
32. M. R. Wieckowski, C. Giorgi, M. Lebiedzinska, J. Duszynski, P. Pinton, Isolation of
mitochondria-associated membranes and mitochondria from animal tissues and cells.
Nature Protocols **4**, 1582-1590 (2009).
33. P. Pinton, Mitochondria-associated membranes (MAMs) and pathologies. *Cell Death Dis*
9, 413 (2018).
34. M. Yang *et al.*, Mitochondria-Associated ER Membranes – The Origin Site of Autophagy.
Frontiers in Cell and Developmental Biology **8** (2020).
35. S. Li *et al.*, Predicting Network Activity from High Throughput Metabolomics. *PLOS*
Computational Biology **9**, e1003123 (2013).
36. D. Phillips, A. M. Aponte, S. A. French, D. J. Chess, R. S. Balaban, Succinyl-CoA synthetase
is a phosphate target for the activation of mitochondrial metabolism. *Biochemistry* **48**,
7140-7149 (2009).

37.   I. Martínez-Reyes, N. S. Chandel, Mitochondrial TCA cycle metabolites control
968   physiology and disease. *Nature Communications* **11**, 102 (2020).
969   38.   J. J. Chen, S. Zhang, Heme-regulated eIF2alpha kinase in erythropoiesis and
970   hemoglobinopathies. *Blood* **134**, 1697-1707 (2019).
971   39.   A. P. Han *et al.*, Heme-regulated eIF2alpha kinase (HRI) is required for translational
972   regulation and survival of erythroid precursors in iron deficiency. *EMBO J* **20**, 6909-6918
973   (2001).
974   40.   J. Igarashi *et al.*, Elucidation of the Heme Binding Site of Heme-regulated Eukaryotic
975   Initiation Factor 2α Kinase and the Role of the Regulatory Motif in Heme Sensing by
976   Spectroscopic and Catalytic Studies of Mutant Proteins. *Journal of Biological Chemistry*
977   **283**, 18782-18791 (2008).
978   41.   R. N. V. S. Suragani *et al.*, Heme-regulated eIF2α kinase activated Atf4 signaling pathway
979   in oxidative stress and erythropoiesis. *Blood* **119**, 5276-5284 (2012).
980   42.   H. Mellor, K. M. Flowers, S. R. Kimball, L. S. Jefferson, Cloning and characterization of
981   cDNA encoding rat hemin-sensitive initiation factor-2 alpha (eIF-2 alpha) kinase.
982   Evidence for multitissue expression. *Journal of Biological Chemistry* **269**, 10201-10204
983   (1994).
984   43.   J. J. Berlanga, S. Herrero, C. de Haro, Characterization of the Hemin-sensitive Eukaryotic
985   Initiation Factor 2α Kinase from Mouse Nonerythroid Cells*. *Journal of Biological
986   Chemistry* **273**, 32340-32346 (1998).
987   44.   A. B. Page *et al.*, Persistent eIF2alpha(P) is colocalized with cytoplasmic cytochrome c in
988   vulnerable hippocampal neurons after 4 hours of reperfusion following 10-minute
989   complete brain ischemia. *Acta Neuropathol* **106**, 8-16 (2003).
990   45.   T. Tsuyama, A. Tsubouchi, T. Usui, H. Imamura, T. Uemura, Mitochondrial dysfunction
991   induces dendritic loss via eIF2α phosphorylation. *Journal of Cell Biology* **216**, 815-834
992   (2017).
993   46.   S. Zhang *et al.*, HRI coordinates translation necessary for protein homeostasis and
994   mitochondrial function in erythropoiesis. *eLife* **8**, e46976 (2019).
995   47.   Y. Miyagi *et al.*, Abundant expression of translation initiation factor EIF-4E in post-
996   meiotic germ cells of the rat testis. *Laboratory investigation; a journal of technical
997   methods and pathology* **73**, 890-898 (1995).
998   48.   J. Marill, N. Idres, C. C. Capron, E. Nguyen, G. G. Chabot, Retinoic acid metabolism and
999   mechanism of action: a review. *Curr Drug Metab* **4**, 1-10 (2003).
49.   N. Y. Kedishvili, Enzymology of retinoic acid biosynthesis and degradation: Thematic
Review Series: Fat-Soluble Vitamins: Vitamin A. *Journal of Lipid Research* **54**, 1744-1760
(2013).
50.   T. J. Cunningham, G. Duester, Mechanisms of retinoic acid signalling and its roles in
organ and limb development. *Nature Reviews Molecular Cell Biology* **16**, 110-123 (2015).
51.   H. B. Everts, C. D. Berdanier, Regulation of Mitochondrial Gene Expression by Retinoids.
IUBMB Life (International Union of Biochemistry and Molecular Biology: Life) **54**, 45-49
(2002).
52.   R. Puttagunta, S. Di Giovanni, Retinoic acid signaling in axonal regeneration. *Front Mol
Neurosci* **4**, 59 (2011).
53.   C. D. Berdanier, H. B. Everts, C. Hermoyian, C. E. Mathews, Role of vitamin A in
mitochondrial gene expression. *Diabetes Res Clin Pract* **54 Suppl 2**, S11-27 (2001).

54.   N. NOY, Retinoid-binding proteins: mediators of retinoid action. *Biochemical Journal*
348, 481-495 (2000).

55.   J. L. Napoli, Cellular retinoid binding-proteins, CRBP, CRABP, FABP5: Effects on retinoid
metabolism, function and related diseases. *Pharmacology & therapeutics* **173**, 19-33
(2017).

56.   R.-Z. Liu *et al.*, CRABP1 is associated with a poor prognosis in breast cancer: adding to
the complexity of breast cancer cell response to retinoic acid. *Molecular Cancer* **14**, 129
(2015).

57.   C. A. Hogarth, M. D. Griswold, The key role of vitamin A in spermatogenesis. *J Clin Invest*
120, 956-962 (2010).

58.   J. K. Amory *et al.*, Isotretinoin administration improves sperm production in men with
infertility from oligoasthenozoospermia: a pilot study. *Andrology* **5**, 1115-1123 (2017).

59.   J. T. Busada *et al.*, Retinoic acid regulates Kit translation during spermatogonial
differentiation in the mouse. **397**, 140-149 (2015).

60.   M. D. Griswold, Spermatogenesis: The Commitment to Meiosis. *Physiol Rev* **96**, 1-17
(2016).

61.   C. Peluso-Iltis, J. Osz, N. Rochel, "DNA recognition by retinoic acid nuclear receptors".
(Elsevier, 2020), 10.1016/bs.mie.2020.03.001, pp. 235-260.

62.   S. Vankoningsloo *et al.*, Mitochondrial dysfunction induces triglyceride accumulation in
3T3-L1 cells: role of fatty acid beta-oxidation and glucose. *J Lipid Res* **46**, 1133-1149
(2005).

63.   Y. Kawano, D. E. Cohen, Mechanisms of hepatic triglyceride accumulation in non-
alcoholic fatty liver disease. *Journal of Gastroenterology* **48**, 434-441 (2013).

64.   Y. Toyoda *et al.*, MRP class of human ATP binding cassette (ABC) transporters: historical
background and new research directions. *Xenobiotica* **38**, 833-862 (2008).

65.   Y. K. Zhang, Y. J. Wang, P. Gupta, Z. S. Chen, Multidrug Resistance Proteins (MRPs) and
Cancer Therapy. *AAPS J* **17**, 802-812 (2015).

66.   A. Lorico *et al.*, Disruption of the murine MRP (multidrug resistance protein) gene leads
to increased sensitivity to etoposide (VP-16) and increased levels of glutathione. *Cancer*
Res **57**, 5238-5242 (1997).

67.   J. Wijnholds *et al.*, Increased sensitivity to anticancer drugs and decreased inflammatory
response in mice lacking the multidrug resistance-associated protein. *Nature medicine*
3, 1275-1279 (1997).

68.   C. J. F. De Wolf *et al.*, cGMP transport by vesicles from human and mouse erythrocytes.
FEBS Journal **274**, 439-450 (2007).

69.   G. Jedlitschky, B. Burchell, D. Keppler, The Multidrug Resistance Protein 5 Functions as
an ATP-dependent Export Pump for Cyclic Nucleotides. *Journal of Biological Chemistry*
275, 30069-30074 (2000).

70.   P. Borst, C. de Wolf, K. van de Wetering, Multidrug resistance-associated proteins 3, 4,
and 5. *Pflugers Arch* **453**, 661-673 (2007).

71.   D. M. Klein, S. H. Wright, N. J. Cherrington, Localization of Multidrug Resistance-
Associated Proteins along the Blood-Testis Barrier in Rat, Macaque, and Human Testis.
Drug Metabolism and Disposition **42**, 89-93 (2014).

72.   B. Shi *et al.*, Effect of sodium butyrate on ABC transporters in lung cancer A549 and
colorectal cancer HCT116 cells. *Oncol Lett* **20**, 148-148 (2020).

73. S. Costello *et al.*, Ca2+-stores in sperm: their identities and functions. *Reproduction*
(Cambridge, England) **138**, 425-437 (2009).

74. K. K. Karna, Y. S. Shin, B. R. Choi, H. K. Kim, J. K. Park, The Role of Endoplasmic Reticulum
Stress Response in Male Reproductive Physiology and Pathology: A Review. *The World*
Journal of Men's Health **37** (2019).

75. X. Ying *et al.*, Endoplasmic reticulum protein 29 (ERp29), a protein related to sperm
maturation is involved in sperm-oocyte fusion in mouse. *Reprod Biol Endocrinol* **8**, 10-10
(2010).

76. T. Klopstock *et al.*, Mitochondrial DNA mutations in multiple symmetric lipomatosis.
Molecular and Cellular Biochemistry **174**, 271-275 (1997).

77. A. Muñoz-Málaga *et al.*, Lipomatosis, proximal myopathy, and the mitochondrial 8344
mutation. A lipid storage myopathy? *Muscle & Nerve* **23**, 538-542 (2000).

78. S. Mazeyrat *et al.*, A Y-encoded subunit of the translation initiation factor Eif2 is
essential for mouse spermatogenesis. *Nature Genetics* **29**, 49-53 (2001).

79. J. M. Harris, P. Lau, S. L. Chen, G. E. Muscat, Characterization of the retinoid orphan-
related receptor-alpha coactivator binding interface: a structural basis for ligand-
independent transcription. *Mol Endocrinol* **16**, 998-1012 (2002).

80. G. Allenby *et al.*, Retinoic acid receptors and retinoid X receptors: interactions with
endogenous retinoic acids. *Proc Natl Acad Sci U S A* **90**, 30-34 (1993).

81. N. Srivastava, M. Pande, Mitochondrion: Features, functions and comparative analysis of
specific probes in detecting sperm cell damages. *Asian Pacific Journal of Reproduction* **5**,
445-452 (2016).

82. K. Nowicka-Bauer *et al.*, Sperm mitochondrial dysfunction and oxidative stress as
possible reasons for isolated asthenozoospermia. *J Physiol Pharmacol* **69** (2018).

83. J. D. D. A. Losano *et al.*, Spermatic mitochondria: role in oxidative homeostasis, sperm
function and possible tools for their assessment. *Zygote* **26**, 251-260 (2018).

84. G. E. Olson, V. P. Winfrey, Structural organization of surface domains of sperm
mitochondria. *Molecular Reproduction and Development* **33**, 89-98 (1992).

85. M. Maneesh, H. Jayalekshmi, Role of reactive oxygen species and antioxidants on
pathophysiology of male reproduction. *Indian J Clin Biochem* **21**, 80-89 (2006).

86. Adam J. Koppers, Lisa A. Mitchell, P. Wang, M. Lin, R. J. Aitken, Phosphoinositide 3-
kinase signalling pathway involvement in a truncated apoptotic cascade associated with
motility loss and oxidative DNA damage in human spermatozoa. *Biochemical Journal*
436, 687-698 (2011).

87. A. Amaral, J. Ramalho-Santos, Assessment of mitochondrial potential: implications for
the correct monitoring of human sperm function. *Int J Androl* **33**, e180-186 (2010).

88. A. Amaral *et al.*, Identification of proteins involved in human sperm motility using high-
throughput differential proteomics. *J Proteome Res* **13**, 5670-5684 (2014).

89. A. Alamo *et al.*, Mitochondrial Membrane Potential Predicts 4-Hour Sperm Motility.
Biomedicines **8**, 196 (2020).

90. K. Hwang, R. C. Walters, L. I. Lipshultz, Contemporary concepts in the evaluation and
management of male infertility. *Nat Rev Urol* **8**, 86-94 (2011).

91. J. Fainberg, J. A. Kashanian, Recent advances in understanding and managing male
infertility. *F1000Res* **8** (2019).

92. R. H. Pek *et al.*, Hemozoin produced by mammals confers heme tolerance. *eLife* **8**
(2019).

93.   N. Nakagata, "Cryopreservation of Mouse Spermatozoa and In Vitro Fertilization" in
Transgenic Mouse Methods and Protocols*, M. H. Hofker, J. van Deursen, Eds. (Humana
Press, Totowa, NJ, 2011), 10.1007/978-1-60761-974-1_4, pp. 57-73.
94.   Y. Ho, K. Wigglesworth, J. J. Eppig, R. M. Schultz, Preimplantation development of mouse
embryos in KSOM: Augmentation by amino acids and analysis of gene expression.
Molecular Reproduction and Development **41**, 232-238 (1995).
95.   D. A. Williams, M. F. Rosenblatt, D. R. Beier, R. D. Cone, Generation of murine stromal
cell lines supporting hematopoietic stem cell proliferation by use of recombinant
retrovirus vectors encoding simian virus 40 large T antigen. *Molecular and Cellular*
Biology **8**, 3864-3871 (1988).
96.   J. Chong *et al.*, MetaboAnalyst 4.0: towards more transparent and integrative
metabolomics analysis. *Nucleic Acids Res* **46**, W486-W494 (2018).
97.   J. Chong, M. Yamamoto, J. Xia, MetaboAnalystR 2.0: From Raw Spectra to Biological
Insights. *Metabolites* **9**, 57 (2019).
98.   J. Chong, D. S. Wishart, J. Xia, Using MetaboAnalyst 4.0 for Comprehensive and
Integrative Metabolomics Data Analysis. *Current Protocols in Bioinformatics* **68**, e86
(2019).
99.   J. W. Jones, K. Pierzchalski, J. Yu, M. A. Kane, Use of fast HPLC multiple reaction
monitoring cubed for endogenous retinoic acid quantification in complex matrices. *Anal*
Chem **87**, 3222-3230 (2015).
100.   M. A. Kane, N. Chen, S. Sparks, J. L. Napoli, Quantification of endogenous retinoic acid in
limited biological samples by LC/MS/MS. *Biochem J* **388**, 363-369 (2005).
101.   M. A. Kane, A. E. Folias, C. Wang, J. L. Napoli, Quantitative profiling of endogenous
retinoic acid in vivo and in vitro by tandem mass spectrometry. *Anal Chem* **80**, 1702-
1708 (2008).
102.   M. A. Kane, J. L. Napoli, Quantification of endogenous retinoids. *Methods Mol Biol* **652**,
1-54 (2010).
103.   M. A. Kane, A. E. Folias, J. L. Napoli, HPLC/UV quantitation of retinal, retinol, and retinyl
esters in serum and tissues. *Anal Biochem* **378**, 71-79 (2008).
104.   C. J. Bult, J. A. Blake, C. L. Smith, J. A. Kadin, J. E. Richardson, Mouse Genome Database
(MGD) 2019. *Nucleic Acids Res* **47**, D801-d806 (2019).
105.   T. F. Hayamizu, M. Mangan, J. P. Corradi, J. A. Kadin, M. Ringwald, The Adult Mouse
Anatomical Dictionary: a tool for annotating and integrating data. *Genome Biol* **6**, R29
(2005).
106.   C. L. Smith, J. T. Eppig, The mammalian phenotype ontology: enabling robust annotation
and comparative analysis. *Wiley Interdiscip Rev Syst Biol Med* **1**, 390-399 (2009).
107.   J. R. Smith *et al.*, The Year of the Rat: The Rat Genome Database at 20: a multi-species
knowledgebase and analysis platform. *Nucleic Acids Res* **48**, D731-d742 (2020).
108.   N. Saitou, M. Nei, The neighbor-joining method: a new method for reconstructing
phylogenetic trees. *Mol Biol Evol* **4**, 406-425 (1987).
109.   E. Zuckerkandl, L. Pauling, "Evolutionary Divergence and Convergence in Proteins" in
Evolving Genes and Proteins*, V. Bryson, H. J. Vogel, Eds. (Academic Press, 1965),
https://doi.org/10.1016/B978-1-4832-2734-4.50017-6, pp. 97-166.
110.   S. Kumar, G. Stecher, M. Li, C. Knyaz, K. Tamura, MEGA X: Molecular Evolutionary
Genetics Analysis across Computing Platforms. *Mol Biol Evol* **35**, 1547-1549 (2018).

111.   G. Stecher, K. Tamura, S. Kumar, Molecular Evolutionary Genetics Analysis (MEGA) for
1149   macOS. *Mol Biol Evol* **37**, 1237-1239 (2020).

1150
1151

1152   **Figures and Tables**

1153

1154
1155

Figure 1. Ablation of MRP9 in mice induces MRP5 expression in the testes
{A} Amino acid FASTA sequences for MRPs in humans, mice, zebrafish and worms were aligned
and used to generate the optimal phylogenetic tree shown. The evolutionary history was inferred
using the Neighbor-Joining method. The tree is drawn to scale, with branch lengths in the same
units as those of the evolutionary distances used to infer the phylogenetic tree. The evolutionary
distances were computed using the Poisson correction method and are in the units of the number
of amino acid substitutions per site. **{B}** Gene loci schematic of 5' UTR and the first four exons of
Abcc12 indicating the CRISPR target site in second exon for the generation of knockout MRP9
mice. **{C}** Immunoblot of protein lysates from transfected HeLa cells and Wildtype and MRP9 KO
tissues probed for MRP9 and MRP5. **{D}** MRP9 immunohistochemistry analysis of paraffin-
embedded testes tissue sections from WT and MRP9 KO mice. Tissue sections were probed for
MRP9 and lightly counterstained with hematoxylin. Images are representative of at least 3 mice;
scale bar equals 100 μm. **{E}** Immunoblot of protein lysates from wildtype and MRP9 KO testes
(n=6) probed for MRP5, MRP9 and Tubulin. **{F}** Quantification of MRP5 protein expression in
wildtype and MRP9 KO testes from subpanel E, normalized to Tubulin; *P* value = 0.0061.
1171

1172
1173

Figure 2. DKO mice are viable but have reduced male reproductive fecundity

{A} Genotyped progeny from MRP5-/- and MRR9+/- intercrosses (left), n = 190, Chi-squared = 0.568 and *P* value = 0.753; Genotyped progeny from MRP5+/- and MRR9-/- intercrosses (right), n = 166, Chi-squared = 0.976 and *P* value = 0.614. **{B}** Quantification of the number of pups weaned per litter of intercrossed WT, DKO, MRP5-/- MRR9+/-, MRP5+/- MRR9-/-, and DKO females crossed with WT or double het mice, **** *P* value < 0.0001. **{C}** Representative images of penile prolapse (left) and expressed seminal coagulum (right) in DKO male mice setup for breeding. **{D}** *In vitro* fertilization rate of WT and DKO sperm incubated with WT oocytes for 5 hours at a concentration of 1.0 x 10^6 sperm/ml. Next day total number of two cell embryos were determined and fertilization rate was calculated based on dividing by the total number of oocytes inseminated for each male, ** *P* value = 0.0019. **{E}** Total concentration of spermatozoa calculated by IVOS system computer assisted sperm analyzer.

1186

Figure 3. MRP9 is enriched in mitochondrial associated membranes (MAMs) of the testes

{A} MDCKII cells grown on transwell filters transfected with basolateral marker sodium taurocholate cotransporting polypeptide *NTCP-GFP*, apical marker syntaxin 3 *SYN3-GFP*, *ABCC12* or *ABCC5*. Polarization of the monolayers was additionally confirmed prior to fixation by measurement of trans-epithelial electrical resistance. A representative confocal section (XY) is depicted along with the composite stacks in the side panel views (XZ and YZ), scale bar equals 5 μm. **{B}** Immunoblot analysis of crude mitochondrial fractionation from testes of mice. Membrane was probed for MRP9 and HSP60, a mitochondrial matrix protein marker. **{C}** Airyscan super resolution microscopy and 3D rendering of HeLa cells transfected with human *ABCC12* for 48 hours and treated with Mitotracker Deep Red FM immediately prior to fixation and immunofluorescence. Antibody probing for MRP9 and Calnexin was followed by Alexa-488 and Alexa-568 secondary antibodies respectively followed by DAPI counter staining prior to mounting. **{D}** Immunoblot analysis of subcellular fractionation from testes of mice. Membranes were probed for MRP9 (M9II-3 antibody) and MFN2, CANX, HSP60 markers of outer mitochondrial membrane, ER membrane, and mitochondrial matrix respectively. **{E}** Immunoblot of subcellular fractionation from testes of wildtype, MRP9 heterozygous and double knockout mice. Membranes were probed for MRP9 (M9I-27 antibody) and MRP5.

1207
1208

Figure 4. Metabolomics and transcriptomics reveal pervasive perturbation in DKO testes

{A} Variable Importance in Projection (VIP) Heatmap of the top 25 differential metabolites in testes from aqueous phase extractions on C18 column run in negative mode. **{B}** Volcano plot of gene expression changes from DKO and WT testes; significance threshold set at -log(P value) > 5. **{C}** Top twelve Gene Ontology (GO) Pathways perturbed in DKO testes based on Qiagen IPA analysis. **{D}** Key overlapping canonical pathways in testes based on Qiagen IPA analysis. **{E}** MetaboAnalyst pathway impact analysis from combining RNAseq differential gene expression changes and metabolomics mummichog putative KEGG IDs of the testes; top five pathways by P value are annotated.

1219
1220

Figure 5. Loss of MRP5 and MRP9 induces EIF2 signaling to mitigate mitochondrial dysfunction

{A} Output of pathway pipeline analysis investigating all "mitochondria" and "mitochondrial disfunction" GO pathway gene lists. Gene lists were curated for integration with RNA expression data. Real expression was validated by thresholding a minimum 10 transcripts per million and statistically significant genes ($P < 0.05$) fold change were output in volcano plot and heatmap with 200 total genes identified, sorted by ascending P value. **{B}** Output of pathway pipeline analysis investigating all "EIF2 signaling" and "translational regulation" GO pathway gene lists. Gene lists were curated for integration with RNA expression data. Real expression was validated by thresholding a minimum 10 transcripts per million and statistically significant genes ($P < 0.05$) fold change were output in volcano plot and heatmap with 95 total genes identified, sorted by ascending P value. **{C}** Immunoblot analysis of testes lysates from DKO and WT mice probed for MRP5, MRP9, Serine 51 phosphorylated EIF2a and total EIF2a protein. Blot is representative of three separate experiments, each lane represents an individual mouse, n=8 mice per genotype. **{D}** Quantification of DKO and WT phosphorylated EIF2a normalized to total EIF2a protein in subpanel C, **** P value < 0.0001. **{E}** Heme content of pooled caudal epididymides from DKO and WT mice by oxalic acid quantification, n=15 animals per genotype (1.5 mice or 3 epididymides per replicate), * P value = 0.0275.

1239

1240
1241

Figure 6. Spermatozoa are severely compromised by the combined loss of MRP5 and MRP9

{A} Representative sorting analysis of WT and DKO sperm stained with JC-1. Q3 is gating for populations that are FITC-pos and PE-neg representing a reduction J-aggregates / red fluorescence and therefore decreased mitochondrial membrane potential. **{B}** Quantification of sperm stained with JC-1 and gated for populations that are FITC-pos and PE-neg (from subpanel 6a), n=5 animals, ** *P* value = 0.0055. **{C}** TEM of swum-out caudal epididymal spermatozoa from WT (top) and DKO (bottom). Cross sections of the sperm midpiece visualize cristae and mitochondrial morphology of the mitochondrial sheath. Yellow arrow heads highlight vacuolated and aberrant mitochondria. Representative images from at least 15 FOVs per sample, scalebar equal 1 μm. **{D}** Model figure – the combined ablation of MRP9 and MRP5 causes inappropriate trafficking or distribution of metabolites such as heme and/or other substrates inducing mitochondrial damage. This dysfunction induces signaling through the EIF2 and mTOR pathways to regulate protein translation and UPR with corresponding changes in retinoic acid levels and signaling. Germ cells differentiate into mature spermatozoa, but these are functionally defective as they fail to fertilize normal oocytes due to mitochondrial insufficiencies.

1258