Title Autogenous and anautogenous *Culex pipiens* bioforms exhibit insulin-like peptide signaling

pathway gene expression differences that are not dependent upon larval nutrition

Running Title Nutrition and ILP expression in *Cx. pipiens* autogeny

Alys M. Cheatle Jarvela[1], Katherine L. Bell[1], Anna Noreuil[1], and Megan L. Fritz[1,2]*

[1] Department of Entomology, University of Maryland College Park, 4112 Plant Sciences

Building, College Park, MD 20042

[2] Institute for Advanced Computer Studies, University of Maryland College Park

*corresponding author

Acknowledgements We thank the University of Maryland's Department of Cell Biology and

Molecular Genetics Imaging Core for the use of their Roche LightCycler qPCR machine and

Amy Beaven for expert training on the instrument. Thea Bliss conducted the wing

measurements. We thank Leslie Pick (LP; UMD Dept of Entomology) for thoughtful discussions

during the initial stages of this project. Kevin Vogel reviewed and provided constructive

feedback on an early draft of our manuscript. AMCJ received support from NIH R01GM113230

to LP. This work was funded by University of Maryland startup funds and NIH R01AI125622A to

MLF.

Article Type Original Article

Abstract

Culex pipiens form *pipiens* and *Cx. pipiens* form *molestus* differ in their ability to produce eggs

without a bloodmeal. Autogenous mosquitoes, such as the *molestus* bioform of *Cx. pipiens,*

depend on nutrition acquired as larvae instead of a bloodmeal to fuel the energy intensive

process of vitellogenesis, which requires abundant production of yolk proteins. In

anautogenous mosquito systems, ovary ecdysteroidogenic hormone (OEH) and insulin-like

peptides (ILPs) transduce nutritional signals and trigger egg maturation in response to a

bloodmeal. It is unclear to what extent the process is conserved in autogenous mosquitoes and

how the bloodmeal trigger has been replaced by teneral reserves. Here, we measured the

effects of a series of nutritional regimens on autogeny, time to pupation, and survival in *Cx.*

pipiens form *molestus* and form *pipiens*. We find that abundant nutrients never result in

autogenous form *pipiens* and extremely poor food availability rarely eliminates autogeny from

form *molestus*. However, the number of autogenous eggs generated increases with nutrient

availability. Similarly, using qPCR to quantify gene expression, we find several differences in the

expression levels of *ilps* between bioforms that are reduced and delayed by poor nutrition, but

not extinguished. Changes in *OEH* expression do not explain bioform-specific differences in

autogeny. Surprisingly, the source of most of the gene expression differences correlated with

autogeny is the abdomen, not the brain. Overall, our results suggest that autogeny is

modulated by nutritional availability, but the trait is encoded by genetic differences between

forms and these impact the expression of ILPs.

Keywords insulin-like peptides, *Culex pipiens*, form *molestus*, form *pipiens*, autogeny

Introduction

Blood is a requirement for egg production in most mosquito vectors of disease, and acquisition of blood drives pathogen transmission. This reproductive strategy is known as anautogeny. Yet egg production in the absence of a blood meal, or autogeny, has been described in dozens of mosquito species (Vinogradova, 2000). Autogeny can be obligatory, where females always lay their first (and, in some species, subsequent) batches of eggs without a blood meal, or facultative, where females may lay an autogenous egg raft dependent on their larval environmental conditions (Attardo et al., 2005; Provost-Javier et al., 2010; Tsuji et al., 1990). While it is well-established that autogenous reproduction is under genetic control (Aslamkhan and Laven, 1970; Krishnamurthy and Laven, 1961; Mori et al., 2008; O'Meara et al., 1969; Spielman, 1957; Trpis, 1978), the molecular basis for this unique reproductive strategy remains unknown. Comparing the molecular and physiological tradeoffs underlying autogenous and anautogenous reproduction has strong potential to elucidate the key genetic changes that led to multiple instances of reproductive divergence within and among mosquito species (O'Meara, 1985).

Due to its epidemiological relevance, the reproductive physiology of anautogenous mosquitoes has been investigated extensively. Egg development requires that yolk proteins produced by the fat body are deposited into developing oocytes, a process known as vitellogenesis. In anautogenous mosquitoes, vitellogenesis is tightly suppressed until a blood meal is obtained (reviewed in Attardo et al., 2005; Hansen et al., 2014). Four key signaling pathways, juvenile hormone III (JH), ecdysone, insulin-signaling, and target of rapamycin (TOR), interact to regulate the expression of yolk protein precursors (YPPs) in the fat body, vitellogenesis, and ovarian maturation (reviewed in Attardo et al., 2005; Hansen et al., 2014). JH acts on the fat body and ovaries during maturation, influencing the ability of a female to activate *YPP* genes (Noriega, 2004; Zou et al., 2013). Once a female obtains a blood meal, the steroid hormone ecdysone is the major hormone that regulates egg maturation via activation of

target genes, such as the predominant *YPP* gene, *vitellogenin* (*Vg*) (reviewed in Roy et al.,

2016). Ecdysone is regulated synergistically by ovary ecdysteroidogenic hormone (OEH) and

insulin-signaling. After feeding, stretch receptors in the gut trigger the release of OEH from the

brain into the hemolymph, which stimulates the production and release of ecdysone by follicle

cells of the ovary (Dhara et al., 2013; Hagedorn et al., 1979, 1975). Insulin-like peptides (ILPs),

a group of evolutionarily conserved peptide hormones, may also influence *YPP* gene expression

both directly and indirectly (Hansen et al., 2014).Together, insulin pathway and ecdysone

signaling activate *YPP* expression in the fat body (Roy et al., 2007). In *Aedes aegypti*, one

member of the insulin-like peptide family, ILP3, has been demonstrated to be essential for the

production of ecdysone, providing indirect reinforcement of *YPP* expression (Brown et al., 2008,

Dhara et al. 2013, Vogel et al. 2015). ILP3 also stimulates serine protease activity in the midgut

for blood meal digestion, contributing to the availability of amino acids required to produce YPPs

(Gulia-Nuss et al., 2012). The TOR signaling pathway detects these increases in amino acid

levels and contributes to the activation of *YPP* gene expression (Hansen et al., 2004).

Ultimately, vitellogenesis is not initiated in anautogenous females until these nutrient sensing

pathways signal the availability of yolk building blocks after a blood meal.

In contrast to anautogenous reproduction, the physiology of ovarian maturation in

autogenous mosquitoes has received less attention (Gulia-Nuss et al., 2015, 2012; Kassim et

al., 2012; Provost-Javier et al., 2010). One of the most well-studied autogenous species is the

rockpool mosquito, *Georgecraigius atropalpus* (formerly *Aedes* and *Ochlerotatus*)(Gulia-Nuss et

al., 2015, 2012; Telang et al., 2013, 2006). *G. atropalpus* females that are decapitated within 6

hours of pupal emergence fail to initiate vitellogenesis, demonstrating that hormones secreted

from the brain are critical to autogenous reproduction in this species. When injected, OEH fully

restores and ILP3 partially restores reproductive maturation in these decapitated females

(Gulia-Nuss et al., 2012), demonstrating that the roles of these hormones in *G. atropalpus* are

analogous to what has been found in the primarily anautogenous *Ae. aegypti* after they have

acquired a blood meal.

In *G. atropalpus* and other autogenous mosquitoes, nutritional status interacts with

genetic factors, including those underlying hormonal secretion from the brain, to influence

autogenous egg production. Females emerging from crowded, or nutrient poor larval habitats

may fail to reproduce autogenously or have low fecundity (Kassim et al., 2012; Krishnamurthy

and Laven, 1961; Lounibos et al., 1982; O'Meara and Krasnick, 1970; Trpis, 1978). Such

nutritional control of trait expression is observed in other insect species (Casasa and Moczek,

2018; Chandra et al., 2018; Wheeler and Frederik Nijhout, 1983), and in some cases, the

molecular mechanism underlying this nutritional regulation is known. For example, horn size in

taurus scarab beetles (*Onthophagus taurus*) is dependent on nutritional state, which is

transduced by insulin signaling (Casasa and Moczek, 2018). In ants, upregulation of *insulin-like*

peptide 2 (*ilp2*) occurs in reproductive as opposed to non-reproductive individuals and larval

nutrition can influence adult ILP2 expression levels (Chandra et al., 2018). In these and other

examples, an emerging theme is that insulin signaling is a critical component of the mechanism

that ties phenotype to nutritional status (reviewed in Nijhout and McKenna, 2018). In

autogenous mosquitoes, it is unclear how and whether the nutritional state of females impacts

the expression of ILPs and other genes underlying induction of vitellogenesis and ovarian

maturation.

Here, we examine the interaction between nutrition, gene expression, and ovarian

maturation in a species of mosquito which is polymorphic for autogeny, *Culex pipiens*. Within

the species, there are two interfertile and morphologically indistinguishable bioforms that exhibit

divergent reproductive strategies (*Culex pipiens* form *pipiens* and form *molestus*, hereafter

pipiens and *molestus*) (Harbach et al., 1984; Spielman, 2001; Spielman and Wong, 1973). The

form *molestus* is thought to be facultatively autogenous, while the form *pipiens* is anautogenous

(Roubaud, 1929). Although these populations are genetically distinct (Fonseca et al., 2004;

Kent et al., 2007; Yurchenko et al., 2020), the molecular mechanisms underlying their divergent

reproductive strategies have not been identified.

We begin to address this knowledge gap by quantifying changes in expression of key

nutrient-sensing pathway genes for the autogenous *molestus*, as compared to the

anautogenous *pipiens*. Multiple genes previously demonstrated to play a critical role in ovarian

maturation, including *oeh*, the full suite of *ilp*s, their downstream effector *foxo,* as well as *Vg1b*,

were examined for expression level differences in female heads and abdomens for four days

following adult emergence. We also measured gene expression in females that experienced

nutrient rich and poor larval conditions, which produced dramatic changes in time spent as

larvae and fecundity in autogenous *molestus*. Expression patterns for several of these genes

differed temporally, by form, and in a body segment specific way. Interestingly, insulin-signaling

pathway genes whose expression differed by form also show delayed or dampened expression

in poorly-fed *molestus* individuals, but never matched the expression levels observed in well-fed

pipiens. This demonstrates the relatively greater importance of genotype over nutrition in

autogenous egg production for *Cx. pipiens*. Furthermore, this work establishes a system that

will allow investigation of the key genetic variants which halt ovarian maturation in some females

yet facilitate it in others. Such variants would make attractive targets for novel genetic control

measures that limit ovarian maturation, egg production, and ultimately vector population growth.

Results and Discussion

Bioinformatic and phylogenetic analyses reveal six Culex ilp *orthologs*

At the start of this work, three *ilps* had been identified in *Cx. pipiens* (named *ilp1*, *ilp2*,

and *ilp5*)(Sim and Denlinger, 2009), but based on the numbers in *Aedes aegypti* (n = 8; (Riehle

et al., 2006)), *Anopheles gambiae* (n = 5; (Riehle et al., 2002)), *Anopheles stephensi* (n = 5;

(Marquez et al., 2011)), and *Drosophila melanogaster* (n = 8; Brogiolo et al., 2001; Colombani et

al., 2012; Garelli et al., 2012; Grönke et al., 2010), it was predicted that more remained to be

discovered (Sharma et al., 2019; Sim and Denlinger, 2009). We identified six *Culex ilp* genes

by performing a HMMR analysis in Vectorbase using an alignment of all eight *Drosophila* and

Aedes ilps against the CpipJ2.4 geneset. A phylogenetic analysis of these six *Culex* ILPs

determined their orthology to established sequences from *Drosophila*, *Aedes*, and *Anopheles*

(Fig. 1). We confirmed the identity of previously isolated *Cpip-ilp1* and *-ilp5* genes, as well as

those from the *Culex quinquefasciatus* Johannesburg genome that were included in other's

phylogenetic analyses (Marquez et al., 2011; Sim and Denlinger, 2009). One gene previously

named *Cpip-ilp2* (Genbank accession ACM66967.1) was orthologous to *ilp3* in other mosquito

species. Comparison of "*Cpip-ilp2*" to *Cqui-ilp2* and *Cqui-ilp3* further supports its orthology with

ilp3 (Fig. S1). Therefore, we refer to this sequence as *Cpip-ilp3* in this work.

One new *ilp* gene was identified and had likely been overlooked in previous work

because it is unique to *Culex* (XP_001868260.1). We named this gene *ilp4* because it is

syntenic with *ilp*s *1-3*, as is the case in other Dipteran genomes (Fig. S2). It is not a clear

ortholog of any other Dipteran *ilp4*, however, and likely arose by a lineage-specific gene

duplication event. All of the genes in this syntenic cluster have sequence features that define

them as "insulin-like peptides", such as a long, cleavable C peptide (Okamoto and Yamanaka,

2015). *Cpip-ilp5* and other mosquito *ilp5* orthologs are not syntenic in the genome with other *ilp*s

and contain distinct sequence features that define them as orthologs of *Dmel-ilp7* (Krieger et al.,

2004; Okamoto and Yamanaka, 2015; Riehle et al., 2006). Our phylogenetic analysis upholds

this grouping (Fig. 1). Similarly, *Cqui-ilp6* is in a clade with *Aaeg-ilp6* in our analysis and both

are located elsewhere in the genome, not in a syntenic cluster with other ILP genes. This is

also true for *Dmel-ilp6*, which is considered "IGF-like" based on sequence features such as its

short C peptide. Although our phylogeny does not group mosquito *ilp6* genes with *Dmel-ilp6*,

the short C peptides, extended C-terminus following the A domain, and genomic locations

suggest that they can also be considered "IGF-like" genes (Riehle et al., 2006). In sum, there

are six *ilp* genes in *Cx. pipiens* assemblage species: *ilps1-4* are true insulin-like peptides, *ilp5* is

similar to *Drosophila*'s atypical *ilp7*, and *ilp6* is best described as an IGF-like peptide.

Autogenic ovarian maturation is enhanced by a nutrient rich larval environment

Prior to quantifying gene expression, we confirmed that the timing and degree of *Culex*

ovarian maturation was consistent with previous observations of autogenous and anautogenous

mosquito populations. Dissections of adult *molestus* and *pipiens* females conducted over 96h

post emergence (PE) showed that ovarian maturation in *molestus* progressed beyond that of

pipiens by 48h PE and was complete by 96h PE, regardless of mating status (Fig. 2). Failure of

mating status to impact autogenous ovarian maturation agreed well with previous studies of

another *molestus* population and *G. atropalpus* (Gulia-Nuss et al., 2012; Kassim et al., 2012;

Spielman, 1957). Furthermore, our time course of ovarian maturation was consistent with

studies showing *molestus* follicle length reaches its maximum between 80-105h post-

emergence (Spielman, 1957), and egg deposition begins at 120h (Kassim et al., 2012).

Nutrient rich and poor aquatic environments are known to impact body size, larval

development, mortality, and egg production in autogenous mosquitoes (Kassim et al., 2012;

Lounibos et al., 1982; O'Meara and Krasnick, 1970; Telang and Wells, 2004). Here, we

examined both anautogenous *pipiens* and autogenous *molestus* for these traits in a

comparative framework under identical environmental conditions. Rather than rearing larvae

together in the same environment, as was previously done (Kassim et al., 2012; Spielman,

1957), we reared each larva in a single well of cell culture plate to avoid the confounding effects

of competition on trait expression. Each larva was consistently fed one of four diet treatments

ranging from 0.25mg desiccated liver powder (LP) + 0.14mg dry yeast (DY) per larva (extra-low)

to 1.07mg LP + 0.53mg DY (high) every other day. For both forms, larval survivorship was high

(> 85%) and did not differ by diet treatment (Fig. S3; Tables S1 & S2). Wing length, which

serves as a proxy measure of female body size and teneral reserves (Telang et al., 2006) was

positively correlated with larval nutrient availability, however (Table 1; Tables S3 & S4).

Together, this indicated that our diet treatments effectively manipulated larval nutrition without

inducing mortality.

As expected based on previous work (Attardo et al., 2005), larval development time was

negatively correlated with nutrient availability for both forms (Fig. S4; Tables S5 & S6). When

fed the highest diet treatment, form *pipiens* and *molestus* larvae developed at similar rates

(Tables S5 & S7), where mean days to pupation were 7.2 (s.d. = 0.8, n = 208) and 7.8 (s.d. 1.1,

n = 222), respectively. At the lowest diet treatments, larval development times lengthened to

10.2 (s.d. = 1.9, n = 202) and 13.1 (s.d. 2.6, n = 200) days, respectively, and there was higher

variation in development time for *molestus*. We reasoned that sex-specific differences in

nutrient acquisition, particularly for *molestus,* could account for this additional variation. In a

separate experiment, we quantified male and female development times for both *molestus* and

pipiens, and our results revealed a three-way interaction between sex, form, and diet treatment

(Fig. 3; Table S8 & S9). Female *molestus* larvae compensated for very low nutrient availability

by lengthening their development time beyond what we observed for *pipiens* females and males

of both forms (Fig. 3). This was consistent with previous studies of autogenous mosquitoes

(Lounibos et al. 1982, Kassim et al. 2012), but unlike our results, *molestus* larvae from previous

studies suffered increased mortality (up to 38.7%), likely due to the more severe starvation

conditions imposed for the lowest diet treatments (Kassim et al., 2012).

Ovarian dissections revealed that larval nutrition was positively correlated with the

numbers of elongated follicles produced by each *molestus* female (Tables S10 & S11), while

pipiens females never showed signs of ovarian maturation (Table S12). Form *molestus* females

raised at the highest diet treatment matured, on average, 47.6 (s.d. = 14.3, n = 62) follicles

whereas those from the lowest diet treatment matured only 14.3 (s.d. = 9.6, n = 51; Table S12,

Fig. S4). We also quantified the probability that *molestus* females failed to produce any mature

follicles according to diet treatment. From the highest to lowest diet treatment, 1.6 %, 3.2%,

2.8% and 19.6% of *molestus* females showed no signs of ovarian maturation. Our statistical

analysis indicated that the lowest diet treatment had greater potential to halt ovarian maturation

in *molestus* females than did the other diet treatments (Tables S13 & S14), confirming that

expression of autogeny is the result of a genotype by environment interaction. Finally, our

observation that poorly nourished *molestus* females lengthened larval development time led us

to quantify the relationship between time spent as larvae and the numbers of elongated follicles

produced. We tested for a positive correlation between development time and numbers of

elongated follicles produced by females reared under the low and extra low diet treatments but

found little evidence of this (Fig. S5; Tables S15 & S16).

Altogether, these results support and add to previous work, showing that autogenic

follicle development in *Culex* is a form-specific trait that is quantitatively impacted and

sometimes eliminated by larval diet (Fig. S4). Poor nutritional conditions lengthen larval

development in a pronounced, sex-specific way for autogenous female *molestus*, compared to

anautogenous *pipiens* reared under identical conditions (Fig. 3; Table S8). Yet increased larval

development time does not necessarily result in strong increases in the numbers of elongated

follicles produced by poorly nourished females. These results suggest there is a threshold level

of nutrient acquisition that must be met prior to pupation in autogenous *molestus* females. In

our hands, when nutritional requirements were met and female *molestus* pupated, most initiated

follicle maturation upon eclosion. Extended larval development times observed for *molestus*

females appeared to be more important for progression from larva to pupa than for increasing

reproductive output, however (Fig. S5).

Gene expression changes associated with autogeny are frequently observed in abdominal

tissue samples and are damped or delayed under poor nutritional conditions

After observing significant, yet non-lethal impacts of diet on larval development and

reproduction in our *Culex* forms, we compared the expression patterns of nine genes involved in

nutrient-sensing and ovarian maturation for females raised under nutrient rich and poor

conditions using our previous experimental design. These gene candidates included *ilps 1-6*,

foxo, *Vg1b,* and *oeh*. Autogenous *molestus* and anautogenous *pipiens* were reared individually

under the highest or lowest diet treatments as described above, and non-bloodfed females were

collected at 24-hour intervals PE in groups of ten, according to form and diet treatment. Heads

and abdomens were divided into pooled tissue samples for RNA extraction, cDNA synthesis,

and qPCR analysis. Relative fold changes in gene expression are reported with respect to 0-

24h PE well-fed form *pipiens* (Fig. 4-6)*.* Statistical significance of fold change differences

between forms for each time point were always examined by two-way ANOVA corrected for

multiple comparisons. This also allowed us to assess significance of fold change differences

between nutritional regimens within a bioform. With two discussed exceptions, we found no

significant differences associated with diet treatment.

Previous work suggested that insulin signaling is an attractive candidate pathway for

understanding the regulation of autogeny. For example, *ilp1* and *ilp5* are downregulated in

diapausing *Cx. pipiens* females, which shuts down their ovarian maturation (Sim and Denlinger,

2009). RNAi knock-down of *ilp1* also results in cessation of ovarian development in females not

programmed for diapause (Sim and Denlinger, 2009). Furthermore, ILP3 promotes ovarian

maturation in both the autogenous *G. atropalpus* (Gulia-Nuss et al., 2012), as well as

anautogenous *Ae. aegypti* following a blood meal (Brown et al., 2008). Brain medial

neurosecretory cells are the dominant source of ILP expression not only in mosquitoes, but in

insects in general (reviewed by Okamoto and Yamanaka, 2015). While ILP activity is controlled

primarily by the release of these neuropeptides into the hemolymph, we predicted that

genetically encoded regulatory changes between *Culex* bioforms could affect transcription

levels, and subsequently, ILP activity. Therefore, we expected to see upregulation of *ilps*,

especially *ilp3*, in the heads of *molestus* females, with the strongest upregulation occurring in

those that were well-fed.

Females of both bioforms expressed *ilp1*, *ilp2*, *ilp3*, *ilp4* and *ilp6* in the head tissues,

regardless of larval feeding conditions (Fig. 4A-E, Fig. 6). *ilp5* was the only insulin-like peptide

that was undetectable in the head (data not shown). Expression of *ilp1* and *ilp6* increased over

time but did so to the same extent in all groups, showing no correlation with autogenous egg

production (Fig. 4A and E). Expression levels of *ilp1*, *ilp3*, *ilp4,* and *ilp6* did not differ between

bioforms or feeding conditions in head tissues (Fig. 4A-E), even though ILP3 release from the

brain is associated with mosquito reproduction in other systems (Brown et al., 2008). One

exception was that *ilp1* expression was significantly different between well-fed and poorly-fed

molestus at 48-72h PE (log2 = -0.57 vs. 0.5, p = 0.018)(Fig. 4A). This likely reflects a slight

delay initiating upregulation of *ilp1* in poorly-fed *molestus*, which is ultimately upregulated in

both diet treatments at 72-96h PE (log2 = 1.67 vs. 1.94). Interestingly, *ilp2* experienced a ten-

fold downregulation in well-fed *molestus* at 24-48h PE vs. well-fed *pipiens* at the same time

point (log2 = -3.03 vs. log2 = 0.33, p = 0.014), and more modest downregulation in poorly-fed

molestus (log2 = -1.64)(Fig. 4B). The downstream effector of ILP signaling, *foxo*, was also

detected in all head samples (Fig. 4F). Well-fed *molestus* downregulated *foxo* by nine-fold at

24-48h PE with respect to equivalently staged *pipiens* (log2 = -2.7 vs. 0.47, p = 0.021) and at

48-72h PE (log2 = -1.81 vs. 1.31, p = 0.023). Weaker downregulation of *foxo* was observed in

poorly-fed *molestus* at those same time points (log2 = -1.57 and -0.52).

We also screened abdominal tissues as a potential alternative source of *ilp* expression.

While not as well-known, there is a precedent for *ilp* expression originating from ovaries, fat

body, and other abdominal tissues in many diverse insects (Okada et al., 2019; Okamoto et al.,

2009a, 2009b). Insects can express ILPs directly from their fat bodies to signal nutritional

conditions in a post-feeding stage. For example, in the beetle *Gnatocerus cornutus*, instead of

the fat body signaling to the brain to synthesize and release ILPs in response to nutritional input,

the fat body directly produces ILP to stimulate post-feeding growth (Okada et al., 2019). In

mosquitoes, ILP expression of abdominal origin has also been characterized. For example,

expression of ILP4 and ILP7 are detectable in ovary tissue in *Aedes* (Riehle et al., 2006).

Furthermore, in female *Anopheles gambiae*, an antibody against ILP1/3/4 detected expression

in abdominal ganglia that run along the body wall (Marquez et al., 2011). This same study

identified ILP1/3/4 immunoreactivity in neuronal axons along the midgut and observed changes

in the expression of several ILPs in the abdomen and midgut in response to sugar deprivation

and exposure to human insulin via a bloodmeal. We reasoned that *ilp*s of similar abdominal or

neural origin may be expressed in *molestus* prior to bloodfeeding and quantified gene

expression in whole abdomens (Fig. 5). *ilp1* was significantly upregulated at 72-96h PE in both

well and poorly-nourished *molestus* (log2 = 1.5 and 1.37 respectively) with respect to well-fed

pipiens of the same age (log2 = -1.78) (p = 0.0014 vs. p = 0.02) (Fig. 5A). *ilp3* expression

increased 53-fold in well-fed *molestus* compared to well-fed *pipiens* at 48-72h PE (log2 = 1.91

vs. -3.83, p = 0.003). The *ilp3* expression level differences between well-fed *molestus* and

pipiens gradually rose to 97-fold by 72-96h PE (log2 = 3.39 vs. -3.21, p = 0.0002) (Fig. 5C).

Poorly-nourished *molestus* significantly upregulated *ilp3* starting at 48-72h PE (16-fold, log2 =

0.24, p = 0.048), but expression levels did not reach those of well-fed *molestus* until 72h PE

(60-fold, log2 = 2.71, p = 0.002). Oddly, the expression of *ilp3* in *molestus* was not significantly

different when compared to poorly nourished *pipiens*. No differences in *ilp2, ilp4* or *ilp6*

expression were observed between bioforms or feeding conditions (Fig. 5B-F).   Finally, as was

the case for head tissues, *foxo* expression was also briefly downregulated in the abdomens of

well-fed *molestus.* We observed over three-fold lower expression in well-fed *molestus* vs. well-

fed *pipiens* at 0-24h PE (log2 = -1.86 vs. 0, p = 0.040) (Fig. 5G).

Interestingly, *ilp5* was only expressed in abdominal tissues, and primarily by the *pipiens*

bioform (Fig. 5E). Predominantly abdominal expression is conserved in *Aedes* and *Anopheles*

mosquitoes (Krieger et al., 2004; Riehle et al., 2006), so it was unsurprising that *ilp5* was not

detectable in head tissues. At any given time, expression was three to five times lower in

molestus than well-fed *pipiens* of the same stage (log2 on average was approximately -2 vs 0, p

< 0.05 in each case). While expression was equally low in *molestus* irrespective of nutritional

regimen, in *pipiens*, poor nutrition was associated with modest, but not significant, upregulation

of *ilp5*. These form and dietary differences in *ilp5* expression are especially intriguing. Previous

work found that the expression of *ilp5* was associated with a non-diapause state in *Culex*

pipiens (Sim and Denlinger, 2009). Indeed, one of the primary differences between the *pipiens*

and *molestus* bioforms is the ability to undergo diapause (Denlinger and Armbruster, 2014;

Vinogradova, 2000), although it is notable that knock-down of ILP5 had no effect on diapause

phenotypes in previous work (Sim and Denlinger, 2009). Currently, the functional role of *ilp5* in

pipiens is not well understood, but clues from *Aedes* suggest it is critical to acquisition of teneral

reserves. For example, CRISPR-Cas9 ablation of *ilp5* in *Aedes* resulted in bigger mosquitoes

with larger stores of lipids accumulated during larval stages (Ling and Raikhel, 2018). Because

increases in female size and teneral reserves are critical for autogeny (Chambers and Klowden,

1994; Gulia-Nuss et al., 2015; Telang et al., 2006), the significant reduction in *ilp5* expression

that we observed in the *molestus* bioform is potentially important for explaining a gain of

autogenic ability.

To summarize our major findings for insulin signaling pathway genes, we observed brief

downregulation of *ilp2* and *foxo* in the heads of females beginning at 24h PE. Upregulation of

ilp1 and *ilp3* occurred in the abdomens of *molestus* females after 48h PE, the latter of which

experienced more dramatic upregulation. Abdominal expression of *foxo* was mildly

downregulated in well-fed *molestus* females. Interestingly, *ilp5*, whose function is currently not

well understood in *Cx. pipiens*, was only detected in the abdominal tissue, and strongly

downregulated in *molestus* throughout the course of our experiment.

We then determined the time course of *Vg1b* expression to determine whether insulin-

signaling pathway genes could be regulators of its expression. In anautogenous systems, *Vg*

expression is repressed until a bloodmeal is acquired (Hansen et al., 2014). FOXO is the

downstream effector of insulin signaling. It is a forkhead box transcription factor that represses

the expression of *Vg* in many insects until the insulin signaling pathway is activated (Sheng et

al., 2011). Insulin signaling triggers phosphorylation of FOXO and deportation from the nucleus

to the cytoplasm, relieving the repression of *Vg* genes. In *molestus*, *Vg1b* was strongly

upregulated (p < 0.0001) in abdominal tissues immediately upon eclosion in well-nourished

samples and by 24-48h PE in poorly-nourished samples (Fig. 5H). Well-fed and poorly-fed

molestus were significantly different at 0-24h PE (p = 0.011), owing to the delayed onset of

upregulation that occurs in the poorly-fed sample. This agreed well with previous work in *Cx.*

tarsalis which demonstrated robust upregulation of this gene within a day of emergence in

autogenous females (Provost-Javier et al., 2010). Both well- and poorly-nourished *molestus*

samples ultimately attained changes in expression greater than 65,000 times that of newly

eclosed *pipiens*. It is notable that *Vg1b* expression peaked at 24-48h PE (log2 = 15.29) in well-

fed *molestus* but was delayed until 48-72h PE (log2 = 16) in poorly-fed *molestus*. No

upregulation occurs in *pipiens* in any feeding condition or at any time point (Fig. 5H).

Interestingly, *Vg1b* is detected before adulthood in *Cx. tarsalis,* and in *Ae. albopictus, Vg* has an

additional role in repressing blood-seeking behavior in young, sugar-fed females (Dittmer et al.,

2019; Provost-Javier et al., 2010). We can infer that peak *Vg* transcription must occur before

48h PE because ovary maturation is visible by this point (Fig. 2). It may also be that very early

upregulation of *Vg1b* serves to block blood-seeking behavior prior to or in addition to egg-

producing functions.

We next examined expression patterns of insulin-signaling pathway genes in light of

Vg1b's temporal dynamics. An early-acting difference of potential importance for rapid *Vg1b*

activation was the downregulation of *foxo* in 0-24h PE well-fed *molestus* abdomens.

Downregulation of this conserved repressor of vitellogenesis genes could make it easier to

activate *Vg1b* expression without bloodmeal-induced activation, even though it is typically

regulated at the post-transcriptional level. However, the difference was only seen in well-fed

molestus while autogeny occurred in both feeding regimens. Downregulation of *ilp5* also

occurred early in *molestus*, but was observed in both diets tested, making it a better candidate

regulator of *Vg1b*. While significantly upregulated in the abdomens of *molestus* females*, ilp1*

and *ilp3* are not likely to be involved in upregulation of *Vg1b*. We did not observe form-specific

expression differences at these genes until after autogenic egg production was in progress (Fig.

5A, 5C, Fig. 6). *ilp3* was not significantly upregulated until 48h PE in well-fed *molestus* and did

not achieve a similar fold increase until 72h PE in poorly-fed *molestus*. By this point, *Vg1b*

expression had passed its peak in both feeding conditions. Likewise, upregulation of *ilp1*

specific to *molestus* did not occur until 72h PE. It may still reinforce the process once it is in

progress since this *ilp* gene has already been shown to regulate ovarian development in *Culex*

mosquitoes. Knock-down of expression of ILP1 has been demonstrated to halt ovarian

development in non-diapausing *pipiens* and lower endogenous expression levels are observed

in diapausing females (Sim and Denlinger, 2009). It is possible that changes in *ilp1* and *ilp3*

expression have some important function in autogeny unrelated to *Vg1b* activation that will be

determined by additional functional studies.

In *Aedes*, OEH, a neuroparsin type neuropeptide, activates the insulin signaling pathway

in response to a bloodmeal, but bypasses the insulin receptor to do so (Dhara et al., 2013).

More recent work demonstrated ecdysone signaling influences *ilp* expression in response to a

bloodmeal, suggesting that it could also be important during autogeny (Ling and Raikhel, 2021).

Based on previous predictions, we expected that OEH stimulation of egg production would be a

conserved feature of mosquito reproduction (Vogel et al., 2015). We measured expression of

OEH in the heads of females from both forms for 0-96h PE, with the expectation that it would be

uniquely upregulated in *molestus* females. We did not detect expression in abdomens during

pilot experiments. Although it was implicated in autogenous reproduction in other mosquito

species (Brown et al., 2008; Dhara et al., 2013; Gulia-Nuss et al., 2015, 2012), *OEH* underwent

very modest upregulation in the heads of *molestus* females throughout our time course relative

to equivalently-staged *pipiens* (Fig. 4G, Fig. 6). OEH upregulation in the heads of *molestus*

females preparing for autogenous egg production occurred after *Vg1b* upregulation and was not

statistically significant compared to expression in equivalent-staged *pipiens* females (Fig. 4G,

Fig. 6). In *G. atropalpus,* ecdysteroid is secreted by the ovaries immediately upon eclosion and

peaks during the first day of emergence under high-food conditions, and experience a ~12 hour

delay of peak secretion under low-food conditions (Telang et al., 2006). This is controlled by

OEH from the brain released within 6 hours of eclosion and produced during pupation and the

first few hours post-eclosion (Fuchs et al., 1980; Gulia-Nuss et al., 2012). It is possible that our

wide 0-24 hour sampling window prevented detection of this early peak in OEH expression, or

that OEH peak expression occurs during pupation for *molestus.* It may also be that OEH

activity is predominantly regulated by release of peptide, and that transcriptional regulation is

not a good indicator of its potential role in *molestus* autogeny.

416   Here, we characterized the effects of nutrition on time to pupation, autogenous egg

production, and expression of genes connected to sensing of nutritional state in *Cx. pipiens.*

We confirmed that genetic factors interacted with nutritional status to regulate autogenous egg

production in *molestus*, while even well-fed *pipiens* never produced eggs autogenously. In

agreement with this trait expression data, we observed strong and significant bioform-specific

gene expression differences, which were only mildly influenced by nutrition. The comparison

between autogenous *molestus* and anautogenous *pipiens* revealed changes in gene expression

patterns in the abdomen (*ilp1*, *ilp3*, *ilp5*), head (*ilp2*), or both (*foxo*) (Fig. 6). Some of these

coincided with dramatic upregulation of *Vg1b* (ex. *foxo*, *ilp5*), while others occurred afterward

(ex. *ilp1*, *ilp3*). Gene expression differences in well- vs. poorly-nourished *molestus* could be

characterized as dampened or delayed rather than statistically significant changes in

expression, in agreement with incomplete loss of autogeny in our lowest diet treatment. Future

work on the functional consequences of these gene expression differences, especially that of

ilp5, will enhance our understanding of the evolution of autogeny and loss of diapause as

bioform-specific traits.

Data Availability

Scripts and data used to examine expression of autogeny under different dietary conditions can

be found at https://github.com/mcadamme/Culex_Nutrition_Exp. Data from gene expression

analysis can be found at: link to be added following acceptance.

Experimental Procedures

Gene identification and orthology

We identified six *Culex ilp* genes by performing a HMMR analysis in Vectorbase

(https://legacy.vectorbase.org/hmmer) using an alignment of all eight *Drosophila* and *Aedes ilps*

against the CpipJ2.4 geneset (Arensburger et al., 2010; Giraldo-Calderón et al., 2015). We next

constructed phylogenetic trees to establish gene orthology. Putative *Culex* ILPs plus previously

characterized insect ILP sequences were aligned in MUSCLE

(https://www.ebi.ac.uk/Tools/msa/muscle/)(Edgar, 2004). Tree topology was determined by

TOPALI v2.5 using both a MrBayes algorithm (Model: JTT + G, Runs: 2 Generations: 500000,

Sample Freq.: 10, Burnin: 25%) and maximum PhyML algorithm (Model: JTT + G, 100

bootstrap runs)(Anisimova and Gascuel, 2006; Milne et al., 2004; Ronquist and Huelsenbeck,

2003). Synteny of *ilp* genes was examined using Genomicus Metazoa (web-code version: 2014-

07-06, database version: 30.01, https://www.genomicus.biologie.ens.fr/genomicus-metazoa-

30.01/cgi-bin/search.pl)(Louis et al., 2013).

Mosquitoes

A below-ground population (form *molestus*) of *Cx. pipiens* was obtained from Calumet,

IL, and has been in colony since 2009 (Mutebi and Savage, 2009) and maintained following

Fritz et al. (Fritz et al., 2015, 2014). An above-ground population (form *pipiens*) of *Cx. pipiens*

was obtained from Evanston, IL, and has been in colony since 2016 (Noreuil and Fritz 2021).

Our laboratory-reared *molestus* had been cultured for 8 years without requiring a blood meal,

but our *pipiens* never produced egg rafts prior to blood feeding during its 2 years in culture. All

adults and larvae were maintained in an environmental chamber with a L:D photo period of

16h:8h, temperature of 25±1∘C, and humidity of 50±10%. Adults for each form were kept in

separate 60 x 60 x 60 cm white BugDorm-2 insect rearing cages (Megaview Science Education

Services Co., Taichung, Taiwan) and were provided with a 10% sucrose solution at all times.

Imaging of ovarian development

Pupae were picked from respective larval rearing pans, placed in individual tubes until

eclosion, then moved to cages containing a 10% sucrose resource where females were allowed

to age with or without males for one to four days. Ovaries were dissected under a

stereomicroscope (Olympus Corporation, model SZ61, Center Valley, PA, USA) at 3.0-4.5 X

magnification on a petri dish filled with 70% ethanol. Images were made using the Olympus

cellSens Entry microscope-mounted imaging system.

Impacts of genotype and nutrition on survivorship, larval development and ovarian maturation

To produce sufficient mosquitoes for experimentation, each colony was blood fed at 12-

14 days of age using an artificial membrane feeder. Goose blood (Na-heparinated, obtained

from Lampire Biological) sweetened with 50% sucrose solution (8:1 ratio) was provided through

a pork sausage casing for a minimum of 1 hour. Approximately 72 hours after blood feeding

females were provided with a cup of dechlorinated tap water and allowed to lay eggs for 48

hours. For each colony, the resulting egg rafts were placed in 27.0 x 19.4 x 9.5 cm plastic pans

with 800ml of dechlorinated tap water and were supplied with 0.16g of liver powder (BD

DifcoTM Desiccated, Powdered Beef Liver) and 0.08g of powdered inactive dry yeast (Genesee

Scientific) suspended in 8ml of RO water. After 48 hours, 60 individuals from each colony were

placed in one of four diet treatments using a 3mL plastic transfer pipette. Those treatments were

as follows: extra low (0.25mg desiccated liver powder (LP), 0.14mg dry yeast (DY), 4ml RO

water per larva), low (0.53mg LP, 0.27mg DY, 4ml RO water per larva), medium (0.80mg LP,

0.40mg DY, 4ml RO water per larva), and high (1.07mg LP, 0.53mg DY, 4ml RO water per

larva). To avoid competition between individuals, larvae were placed in a single well of a 12-well

culture plate. This resulted in 5 plates (12 x 5 = 60 individuals) per treatment, per form. Plates

were held in an environmental chamber with the same conditions as described above.

Individuals were fed every other day. Diet fed on days subsequent to the initial loading of the

plates contained the same mass of food as described above but was suspended in 100µl of RO

water instead of 4ml. Water was changed completely, and the diet was re-loaded as initially

described at day 8 or 10 of the experiment to avoid build-up of waste within each well. The

number of pupae present was recorded daily, and mortality was recorded every other day

during feeding. Pupae were removed from the well and placed in a small plastic cup within a 30

x 30 x 30 white Bugdorm-2 cage. Adults were provided with a 10% sucrose solution and held

under the previously described environmental conditions. Wing and ovarian dissections were

performed on adult females, beginning at 4 days PE. A single wing was removed from each of

3 females per diet treatment, per form and photographed using an Olympus cellSens Entry

microscope-mounted imaging system. Wing length measurements were made from the allular

notch to the tip of the wing using ImageJ software (Schneider et al., 2012). Ovaries were also

removed from at least 20 *molestus* females and 5 *pipiens* females per diet treatment, and the

numbers of follicles advanced beyond Christopher's stage IIb present were recorded (O'Meara

and Krasnick, 1970). This experiment was repeated four times.

In a separate experiment, we measured the impacts of diet, form, and sex on larval

development time. For each bioform and diet treatment (high or extra low), 60 larvae were

placed into wells of the five 12-well cell culture plates and allowed to develop as described

above. Upon pupation, individuals were transferred to a 2ml clear plastic lysis tube plugged

with cotton wool. We monitored emergence every 48h and scored both larval development time

and sex of the emerged adult. This experiment was repeated three times.

Statistical analysis of mortality, pupation, and ovarian maturation

Bayesian generalized linear models (GLMs) were always fit using the package BRMS (v.

2.12.0 (Bürkner, 2017)) in R (R Core Team, 2016). Posterior probability distributions of model

parameters were estimated using Markov Chain Monte Carlo (MCMC). For each model, we ran

4 chains with 6,000 steps, a burn-in of 2,000 steps, and saved every other step resulting in a

total of 8,000 samples drawn from the posterior probability distribution. To ensure a stable

sampling distribution had been reached, we visually examined trace plots, estimated effective

sample size, and calculated Gelman and Rubin's convergence diagnostic (Gelman and Rubin,

1992). Details of the models and their underlying distributions can be found in Tables S1-6, S8-

11, and S13-16.

RNA isolation and qPCR

To compare gene expression levels between species, developmental time points, and

nutritional states, we first extracted total RNA from pools of tissue derived from ten like

individuals using TRIzol Reagent (Invitrogen, Carlsbad, CA) according to the manufacturer's

instructions. Larvae from each bioform were reared individually in 12-well cell culture plates and

fed either high or extra-low diet treatments. Pupae were placed in individual 2mL lysis tubes,

allowed to emerge, and adult females were collected every 24 hours. From these females,

replicate pools of ten heads (bearing intact chemosensory appendages) and ten abdomens

were separately dissected on a small petri dish filled with dry ice for each diet treatment. Pooled

tissue was stored in sterile 1.5 mL microcentrifuge tubes containing TRIzol Reagent at -80°C

until RNA isolation was performed. One minor modification to this is that we back-extracted the

aqueous RNA-containing layer with chloroform an additional time to remove trace amounts of

TRIzol components that interfered with downstream steps. Once the resultant RNA pellet was

dissolved in 20 µL of nuclease-free water, the RNA was treated to remove residual genomic

DNA contamination with the Turbo DNA-free kit (Invitrogen, Carlsbad, CA) according to the

manufacturer's protocol. RNA yield was assessed following genomic DNA removal using a

Nanodrop Lite spectrophotometer (Thermo Fisher Scientific, Waltham, MA). 250ng of RNA was

used to synthesize cDNA with the iScript kit (BioRad, Hercules, CA). cDNA was diluted five-fold

in nuclease-free water for use as a qPCR template, and total RNA was diluted to an equivalent

dilution factor (2ng/µL) for –RT controls. qPCR was performed on a LightCycler 480 real-time

PCR cycler (Roche, Basel, CH) using Luna Universal qPCR master mix (NEB, Ipswich, MA).

Primer sequences can be found in Table S17. Results were analyzed by the ΔΔCt method

(Livak and Schmittgen, 2001). Reference gene (*EF1A*) was selected using the Normfinder

algorithm (Andersen et al., 2004) from a panel of six candidate reference genes selected from

relevant literature (Ling and Salvaterra, 2011; Sim and Denlinger, 2009; Van Hiel et al., 2009).

Statistical significance of qPCR data determined by 2-way ANOVA of ΔΔCts with Tukey's

multiple comparisons test using Graphpad software.

References

Andersen, C.L., Jensen, J.L., Ørntoft, T.F., 2004. Normalization of Real-Time Quantitative Reverse Transcription-PCR Data: A Model-Based Variance Estimation Approach to Identify Genes Suited for Normalization, Applied to Bladder and Colon Cancer Data Sets. Cancer Res. 64, 5245–5250. https://doi.org/10.1158/0008-5472.CAN-04-0496

Anisimova, M., Gascuel, O., 2006. Approximate likelihood-ratio test for branches: A fast, accurate, and powerful alternative. Syst. Biol. 55, 539–552. https://doi.org/10.1080/10635150600755453

Arensburger, P., Megy, K., Waterhouse, R.M., Abrudan, J., Amedeo, P., Antelo, B., Bartholomay, L., Bidwell, S., Caler, E., Camara, F., Campbell, C.L., Campbell, K.S., Casola, C., Castro, M.T., Chandramouliswaran, I., Chapman, S.B., Christley, S., Costas, J., Eisenstadt, E., Feschotte, C., Fraser-Liggett, C., Guigo, R., Haas, B., Hammond, M., Hansson, B.S., Hemingway, J., Hill, S.R., Howarth, C., Ignell, R., Kennedy, R.C., Kodira, C.D., Lobo, N.F., Mao, C., Mayhew, G., Michel, K., Mori, A., Liu, N., Naveira, H., Nene, V., Nguyen, N., Pearson, M.D., Pritham, E.J., Puiu, D., Qi, Y., Ranson, H., Ribeiro, J.M.C., Roberston, H.M., Severson, D.W., Shumway, M., Stanke, M., Strausberg, R.L., Sun, C., Sutton, G., Tu, Z.J., Tubio, J.M.C., Unger, M.F., Vanlandingham, D.L., Vilella, A.J., White, O., White, J.R., Wondji, C.S., Wortman, J., Zdobnov, E.M., Birren, B., Christensen, B.M., Collins, F.H., Cornel, A., Dimopoulos, G., Hannick, L.I., Higgs, S., Lanzaro, G.C., Lawson, D., Lee, N.H., Muskavitch, M.A.T., Raikhel, A.S., Atkinson, P.W., 2010. Sequencing of Culex quinquefasciatus establishes a platform for mosquito comparative genomics. Science 330, 86–88. https://doi.org/10.1126/science.1191864

Aslamkhan, M., Laven, H., 1970. Inheritance of autogeny in the Culex pipiens complex. Pak. J. Zool. 2, 121–147.

Attardo, G.M., Hansen, I.A., Raikhel, A.S., 2005. Nutritional regulation of vitellogenesis in mosquitoes: Implications for anautogeny. Insect Biochem. Mol. Biol., Genetic manipulation of insects 35, 661–675. https://doi.org/10.1016/j.ibmb.2005.02.013

Brogiolo, W., Stocker, H., Ikeya, T., Rintelen, F., Fernandez, R., Hafen, E., 2001. An evolutionarily conserved function of the Drosophila insulin receptor and insulin-like peptides in growth control. Curr. Biol. CB 11, 213–221. https://doi.org/10.1016/s0960-9822(01)00068-9

Brown, M.R., Clark, K.D., Gulia, M., Zhao, Z., Garczynski, S.F., Crim, J.W., Suderman, R.J., Strand, M.R., 2008. An insulin-like peptide regulates egg maturation and metabolism in the mosquito Aedes aegypti. Proc. Natl. Acad. Sci. U. S. A. 105, 5716–5721. https://doi.org/10.1073/pnas.0800478105

Bürkner, P.-C., 2017. brms: An R Package for Bayesian Multilevel Models Using Stan. J. Stat. Softw. 80, 1–28. https://doi.org/10.18637/jss.v080.i01

Casasa, S., Moczek, A.P., 2018. Insulin signalling's role in mediating tissue-specific nutritional plasticity and robustness in the horn-polyphenic beetle Onthophagus taurus. Proc. R. Soc. B Biol. Sci. 285, 20181631. https://doi.org/10.1098/rspb.2018.1631

Chambers, G.M., Klowden, M.J., 1994. Nutritional Reserves of Autogenous and Anautogenous Selected Strains of Aedes albopictus (Diptera: Culicidae). J. Med. Entomol. 31, 554–560. https://doi.org/10.1093/jmedent/31.4.554

Chandra, V., Fetter-Pruneda, I., Oxley, P.R., Ritger, A.L., McKenzie, S.K., Libbrecht, R., Kronauer, D.J.C., 2018. Social regulation of insulin signaling and the evolution of eusociality in ants. Science 361, 398–402. https://doi.org/10.1126/science.aar5723

Clifton, M.E., Noriega, F.G., 2012. The fate of follicles after a blood meal is dependent on previtellogenic nutrition and juvenile hormone in Aedes aegypti. J. Insect Physiol. 58, 1007–1019. https://doi.org/10.1016/j.jinsphys.2012.05.005

Colombani, J., Andersen, D.S., Léopold, P., 2012. Secreted peptide Dilp8 coordinates

Drosophila tissue growth with developmental timing. Science 336, 582–585.
https://doi.org/10.1126/science.1216689
Denlinger, D.L., Armbruster, P.A., 2014. Mosquito diapause. Annu. Rev. Entomol. 59, 73–93.
https://doi.org/10.1146/annurev-ento-011613-162023
Dhara, A., Eum, J.-H., Robertson, A., Gulia-Nuss, M., Vogel, K.J., Clark, K.D., Graf, R., Brown,
M.R., Strand, M.R., 2013. Ovary ecdysteroidogenic hormone functions independently of
the insulin receptor in the yellow fever mosquito, Aedes aegypti. Insect Biochem. Mol.
Biol. 43, 1100–1108. https://doi.org/10.1016/j.ibmb.2013.09.004
Dittmer, J., Alafndi, A., Gabrieli, P., 2019. Fat body–specific vitellogenin expression regulates
host-seeking behaviour in the mosquito Aedes albopictus. PLOS Biol. 17, e3000238.
https://doi.org/10.1371/journal.pbio.3000238
Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. Nucleic Acids Res. 32, 1792–1797. https://doi.org/10.1093/nar/gkh340
Feinsod, F.M., Spielman, A., 1980. Nutrient-mediated juvenile hormone secretion in
mosquitoes. J. Insect Physiol. 26, 113–117. https://doi.org/10.1016/0022-
1910(80)90050-5
Fonseca, D.M., Keyghobadi, N., Malcolm, C.A., Mehmet, C., Schaffner, F., Mogi, M., Fleischer,
R.C., Wilkerson, R.C., 2004. Emerging vectors in the Culex pipiens complex. Science
303, 1535–1538. https://doi.org/10.1126/science.1094247
Fritz, M.L., Walker, E.D., Miller, J.R., Severson, D.W., Dworkin, I., 2015. Divergent host
preferences of above- and below-ground Culex pipiens mosquitoes and their hybrid
offspring. Med. Vet. Entomol. 29, 115–123. https://doi.org/10.1111/mve.12096
Fritz, M.L., Walker, E.D., Yunker, A.J., Dworkin, I., 2014. Daily blood feeding rhythms of
laboratory-reared North American Culex pipiens. J. Circadian Rhythms 12, 1.
https://doi.org/10.1186/1740-3391-12-1
Fuchs, M.S., Sundland, B.R., Kang, S.-H., 1980. In vivo induction of ovarian development in
Aëdes atropalpus by a head extract from Aëdes aegypti. Int. J. Invertebr. Reprod. 2,
121–129. https://doi.org/10.1080/01651269.1980.10553347
Garelli, A., Gontijo, A.M., Miguela, V., Caparros, E., Dominguez, M., 2012. Imaginal discs
secrete insulin-like peptide 8 to mediate plasticity of growth and maturation. Science
336, 579–582. https://doi.org/10.1126/science.1216735
Gelman, A., Rubin, D.B., 1992. Inference from Iterative Simulation Using Multiple Sequences.
Stat. Sci. 7, 457–472. https://doi.org/10.1214/ss/1177011136
Giraldo-Calderón, G.I., Emrich, S.J., MacCallum, R.M., Maslen, G., Dialynas, E., Topalis, P.,
Ho, N., Gesing, S., VectorBase Consortium, Madey, G., Collins, F.H., Lawson, D., 2015.
VectorBase: an updated bioinformatics resource for invertebrate vectors and other
organisms related with human diseases. Nucleic Acids Res. 43, D707-713.
https://doi.org/10.1093/nar/gku1117
Grönke, S., Clarke, D.-F., Broughton, S., Andrews, T.D., Partridge, L., 2010. Molecular
Evolution and Functional Characterization of Drosophila Insulin-Like Peptides. PLOS
Genet. 6, e1000857. https://doi.org/10.1371/journal.pgen.1000857
Gulia-Nuss, M., Elliot, A., Brown, M.R., Strand, M.R., 2015. Multiple factors contribute to
anautogenous reproduction by the mosquito Aedes aegypti. J. Insect Physiol. 82, 8–16.
https://doi.org/10.1016/j.jinsphys.2015.08.001
Gulia-Nuss, M., Eum, J.-H., Strand, M.R., Brown, M.R., 2012. Ovary ecdysteroidogenic
hormone activates egg maturation in the mosquito Georgecraigius atropalpus after adult
eclosion or a blood meal. J. Exp. Biol. 215, 3758–3767.
https://doi.org/10.1242/jeb.074617
Hagedorn, H.H., O'Connor, J.D., Fuchs, M.S., Sage, B., Schlaeger, D.A., Bohm, M.K., 1975.
The ovary as a source of alpha-ecdysone in an adult mosquito. Proc. Natl. Acad. Sci. U.
S. A. 72, 3255–3259. https://doi.org/10.1073/pnas.72.8.3255

Hagedorn, H.H., Shapiro, J.P., Hanaoka, K., 1979. Ovarian ecdysone secretion is controlled by a brain hormone in an adult mosquito. Nature 282, 92–94. https://doi.org/10.1038/282092a0

Hansen, I.A., Attardo, G.M., Park, J.-H., Peng, Q., Raikhel, A.S., 2004. Target of rapamycin-mediated amino acid signaling in mosquito anautogeny. Proc. Natl. Acad. Sci. U. S. A. 101, 10626–10631. https://doi.org/10.1073/pnas.0403460101

Hansen, I.A., Attardo, G.M., Rodriguez, S.D., Drake, L.L., 2014. Four-way regulation of mosquito yolk protein precursor genes by juvenile hormone-, ecdysone-, nutrient-, and insulin-like peptide signaling pathways. Front. Physiol. 5. https://doi.org/10.3389/fphys.2014.00103

Harbach, R.E., Harbach, R.E., Harrison, B.A., Gad, A.M., 1984. Culex (Culex) Molestus Forskål (Diptera: Culicidae): neotype designation, description, variation, and taxonomic status. Proc. Entomol. Soc. Wash. 86, 521–542.

Kassim, N.F.A., Webb, C.E., Russell, R.C., 2012. Is the expression of autogeny by Culex molestus Forskal (Diptera: Culicidae) influenced by larval nutrition or by adult mating, sugar feeding, or blood feeding? J. Vector Ecol. 37, 162–171. https://doi.org/10.1111/j.1948-7134.2012.00213.x

Kent, R.J., Harrington, L.C., Norris, D.E., 2007. Genetic Differences Between Culex pipiens f. molestus and Culex pipiens pipiens (Diptera: Culicidae) in New York. J. Med. Entomol. 44, 50–59.

Krieger, M.J.B., Jahan, N., Riehle, M.A., Cao, C., Brown, M.R., 2004. Molecular characterization of insulin-like peptide genes and their expression in the African malaria mosquito, Anopheles gambiae. Insect Mol. Biol. 13, 305–315. https://doi.org/10.1111/j.0962-1075.2004.00489.x

Krishnamurthy, B.S., Laven, H., 1961. A Note on Inheritance of Autogeny in Culex Mosquitos. Bull. World Health Organ. 24, 675–677.

Ling, D., Salvaterra, P.M., 2011. Robust RT-qPCR Data Normalization: Validation and Selection of Internal Reference Genes during Post-Experimental Data Analysis. PLoS ONE 6. https://doi.org/10.1371/journal.pone.0017762

Ling, L., Raikhel, A.S., 2021. Cross-talk of insulin-like peptides, juvenile hormone, and 20-hydroxyecdysone in regulation of metabolism in the mosquito Aedes aegypti. Proc. Natl. Acad. Sci. 118. https://doi.org/10.1073/pnas.2023470118

Ling, L., Raikhel, A.S., 2018. Serotonin signaling regulates insulin-like peptides for growth, reproduction, and metabolism in the disease vector Aedes aegypti. Proc. Natl. Acad. Sci. 115, E9822–E9831. https://doi.org/10.1073/pnas.1808243115

Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods San Diego Calif 25, 402–408. https://doi.org/10.1006/meth.2001.1262

Louis, A., Muffato, M., Roest Crollius, H., 2013. Genomicus: five genome browsers for comparative genomics in eukaryota. Nucleic Acids Res. 41, D700-705. https://doi.org/10.1093/nar/gks1156

Lounibos, L.P., Van Dover, C., O'Meara, G.F., 1982. Fecundity, autogeny, and the larval environment of the pitcher-plant mosquito, Wyeomyia smithii. Oecologia 55, 160–164. https://doi.org/10.1007/BF00384482

Marquez, A.G., Pietri, J.E., Smithers, H.M., Nuss, A., Antonova, Y., Drexler, A.L., Riehle, M.A., Brown, M.R., Luckhart, S., 2011. Insulin-like peptides in the mosquito Anopheles stephensi: Identification and expression in response to diet and infection with Plasmodium falciparum. Gen. Comp. Endocrinol. 173, 303–312. https://doi.org/10.1016/j.ygcen.2011.06.005

Milne, I., Wright, F., Rowe, G., Marshall, D.F., Husmeier, D., McGuire, G., 2004. TOPALi: software for automatic identification of recombinant sequences within DNA multiple

alignments. Bioinforma. Oxf. Engl. 20, 1806–1807.
https://doi.org/10.1093/bioinformatics/bth155
Mori, A., Romero-Severson, J., Black, W.C., Severson, D.W., 2008. Quantitative trait loci
determining autogeny and body size in the Asian tiger mosquito (Aedes albopictus).
Heredity 101, 75–82. https://doi.org/10.1038/hdy.2008.32
Mostowy, W., Foster, W., 2004. Antagonistic effects of energy status on meal size and egg-
batch size of Aedes aegypti (Diptera: Culicidae). J. Vector Ecol. J. Soc. Vector Ecol. 29,
84–93.
Mutebi, J.-P., Savage, H.M., 2009. Discovery of Culex pipiens pipiens form molestus in
Chicago. J. Am. Mosq. Control Assoc. 25, 500–503. https://doi.org/10.2987/09-5910.1
Nijhout, H.F., McKenna, K.Z., 2018. The distinct roles of insulin signaling in polyphenic
development. Curr. Opin. Insect Sci. 25, 58–64.
https://doi.org/10.1016/j.cois.2017.11.011
Noriega, F.G., 2004. Nutritional regulation of JH synthesis: a mechanism to control reproductive
maturation in mosquitoes? Insect Biochem. Mol. Biol., Molecular and population biology
of mosquitoes 34, 687–693. https://doi.org/10.1016/j.ibmb.2004.03.021
Okada, Y., Katsuki, M., Okamoto, N., Fujioka, H., Okada, K., 2019. A specific type of insulin-like
peptide regulates the conditional growth of a beetle weapon. PLOS Biol. 17, e3000541.
https://doi.org/10.1371/journal.pbio.3000541
Okamoto, N., Yamanaka, N., 2015. Nutrition-dependent control of insect development by
insulin-like peptides. Curr. Opin. Insect Sci., Global change biology * Molecular
physiology 11, 21–30. https://doi.org/10.1016/j.cois.2015.08.001
Okamoto, N., Yamanaka, N., Satake, H., Saegusa, H., Kataoka, H., Mizoguchi, A., 2009a. An
ecdysteroid-inducible insulin-like growth factor-like peptide regulates adult development
of the silkmoth Bombyx mori. FEBS J. 276, 1221–1232. https://doi.org/10.1111/j.1742-
4658.2008.06859.x
Okamoto, N., Yamanaka, N., Yagi, Y., Nishida, Y., Kataoka, H., O'Connor, M.B., Mizoguchi, A.,
2009b. A fat body-derived IGF-like peptide regulates postfeeding growth in Drosophila.
Dev. Cell 17, 885–891. https://doi.org/10.1016/j.devcel.2009.10.008
O'Meara, G.F., 1985. Gonotrophic Interactions in Mosquitoes: Kicking the Blood-Feeding Habit.
Fla. Entomol. 68, 122–133. https://doi.org/10.2307/3494335
O'meara, G.F., Jr, C., B, G., 1969. Monofactorial inheritance of autogeny in Aedes atropalpus.
Mosq. News 29.
O'Meara, G.F., Krasnick, G.J., 1970. Dietary and Genetic Control of the Expression of
Autogenous Reproduction in Aedes Atropalpus (Coq.) (Diptera: Culicidae)1. J. Med.
Entomol. 7, 328–334. https://doi.org/10.1093/jmedent/7.3.328
Provost-Javier, K.N., Chen, S., Rasgon, J.L., 2010. Vitellogenin gene expression in autogenous
Culex tarsalis. Insect Mol. Biol. 19, 423–429. https://doi.org/10.1111/j.1365-
2583.2010.00999.x
R Core Team, 2016. R: A language and environment for statistical computing. R Foundation for
Statistical Computing.
Riehle, M.A., Fan, Y., Cao, C., Brown, M.R., 2006. Molecular characterization of insulin-like
peptides in the yellow fever mosquito, Aedes aegypti: expression, cellular localization,
and phylogeny. Peptides 27, 2547–2560. https://doi.org/10.1016/j.peptides.2006.07.016
Riehle, M.A., Garczynski, S.F., Crim, J.W., Hill, C.A., Brown, M.R., 2002. Neuropeptides and
Peptide Hormones in Anopheles gambiae. Science 298, 172–175.
https://doi.org/10.1126/science.1076827
Ronquist, F., Huelsenbeck, J.P., 2003. MrBayes 3: Bayesian phylogenetic inference under
mixed models. Bioinformatics 19, 1572–1574.
https://doi.org/10.1093/bioinformatics/btg180
Roubaud, E., 1929. Autogenous Cycle of Winter Generations of Culex pipiens L. C. r. Acad. Sci.

188.

Roy, S., Smykal, V., Johnson, L., Saha, T.T., Zou, Z., Raikhel, A.S., 2016. Chapter Five -
Regulation of Reproductive Processes in Female Mosquitoes, in: Raikhel, Alexander S.
(Ed.), Advances in Insect Physiology, Progress in Mosquito Research. Academic Press,
pp. 115–144. https://doi.org/10.1016/bs.aiip.2016.05.004

Roy, S.G., Hansen, I.A., Raikhel, A.S., 2007. Effect of insulin and 20-hydroxyecdysone in the fat
body of the yellow fever mosquito, Aedes aegypti. Insect Biochem. Mol. Biol. 37, 1317–
1326. https://doi.org/10.1016/j.ibmb.2007.08.004

Schneider, C. A., Rasband, W. S., Eliceiri, K. W., 2012. NIH Image to ImageJ: 25 years of
image analysis. Nat. Methods, 9(7), 671–675.

Sharma, A., Nuss, A.B., Gulia-Nuss, M., 2019. Insulin-Like Peptide Signaling in Mosquitoes:
The Road Behind and the Road Ahead. Front. Endocrinol. 10.
https://doi.org/10.3389/fendo.2019.00166

Sheng, Z., Xu, J., Bai, H., Zhu, F., Palli, S.R., 2011. Juvenile Hormone Regulates Vitellogenin
Gene Expression through Insulin-like Peptide Signaling Pathway in the Red Flour
Beetle, Tribolium castaneum. J. Biol. Chem. 286, 41924–41936.
https://doi.org/10.1074/jbc.M111.269845

Sim, C., Denlinger, D.L., 2009. A shut-down in expression of an insulin-like peptide, ILP-1, halts
ovarian maturation during the overwintering diapause of the mosquito Culex pipiens.
Insect Mol. Biol. 18, 325–332. https://doi.org/10.1111/j.1365-2583.2009.00872.x

Spielman, A., 2001. Structure and seasonality of nearctic Culex pipiens populations. Ann. N. Y.
Acad. Sci. 951, 220–234. https://doi.org/10.1111/j.1749-6632.2001.tb02699.x

Spielman, A., 1957. The inheritance of autogeny in the Culex pipiens complex of mosquitoes..
Am. J. Epidemiol. 65, 404–425. https://doi.org/10.1093/oxfordjournals.aje.a119878

Spielman, A., Wong, J., 1973. Environmental Control of Ovarian Diapause in Culex pipiens.
https://doi.org/10.1093/AESA/66.4.905

Telang, A., Li, Y., Noriega, F.G., Brown, M.R., 2006. Effects of larval nutrition on the
endocrinology of mosquito egg development. J. Exp. Biol. 209, 645–655.
https://doi.org/10.1242/jeb.02026

Telang, A., Rechel, J.A., Brandt, J.R., Donnell, D.M., 2013. Analysis of ovary-specific genes in
relation to egg maturation and female nutritional condition in the mosquitoes
Georgecraigius atropalpus and Aedes aegypti (Diptera: Culicidae). J. Insect Physiol. 59,
283–294. https://doi.org/10.1016/j.jinsphys.2012.11.006

Telang, A., Wells, M.A., 2004. The effect of larval and adult nutrition on successful autogenous
egg production by a mosquito. J. Insect Physiol. 50, 677–685.
https://doi.org/10.1016/j.jinsphys.2004.05.001

Trpis, M., 1978. Genetics of hematophagy and autogeny in the Aedes scutellaris complex
(Diptera: Culicidae). J. Med. Entomol. 15, 73–80.
https://doi.org/10.1093/jmedent/15.1.73

Tsuji, N., Okazawa, T., Yamamura, N., 1990. Autogenous and Anautogenous Mosquitoes: a
Mathematical Analysis of Reproductive Strategies. J. Med. Entomol. 27, 446–453.
https://doi.org/10.1093/jmedent/27.4.446

Van Hiel, M.B., Van Wielendaele, P., Temmerman, L., Van Soest, S., Vuerinckx, K.,
Huybrechts, R., Broeck, J.V., Simonet, G., 2009. Identification and validation of
housekeeping genes in brains of the desert locust Schistocerca gregaria under different
developmental conditions. BMC Mol. Biol. 10, 56. https://doi.org/10.1186/1471-2199-10-
56

Vinogradova, A.B., 2000. Culex Pipiens Pipiens Mosquitoes: Taxonomy, Distribution, Ecology,
Physiology, Genetics, Applied Importance and Control. Pensoft Publishers, Sofia.

Vogel, K.J., Brown, M.R., Strand, M.R., 2015. Ovary ecdysteroidogenic hormone requires a
receptor tyrosine kinase to activate egg formation in the mosquito Aedes aegypti. Proc.

Natl. Acad. Sci. 112, 5057–5062. https://doi.org/10.1073/pnas.1501814112
Wheeler, D.E., Frederik Nijhout, H., 1983. Soldier determination in Pheidole bicarinata: Effect of
methoprene on caste and size within castes. J. Insect Physiol. 29, 847–854.
https://doi.org/10.1016/0022-1910(83)90151-8
Yurchenko, A.A., Masri, R.A., Khrabrova, N.V., Sibataev, A.K., Fritz, M.L., Sharakhova, M.V.,
2020. Genomic differentiation and intercontinental population structure of mosquito
vectors Culex pipiens pipiens and Culex pipiens molestus. Sci. Rep. 10.
https://doi.org/10.1038/s41598-020-63305-z
Zou, Z., Saha, T.T., Roy, S., Shin, S.W., Backman, T.W.H., Girke, T., White, K.P., Raikhel, A.S.,
2013. Juvenile hormone and its receptor, methoprene-tolerant, control the dynamics of
mosquito gene expression. Proc. Natl. Acad. Sci.
https://doi.org/10.1073/pnas.1305293110
816
817
818

819 **Tables**

821 **Table 1: Mean wing length is positively correlated with nutrient availability in larval *Cx.***

822 ***pipiens.*** Means and standard deviations were calculated using one wing for 12 individuals per

823 form, per diet treatment. Correlations between wing length and larval nutrition were determined

824 using a Bayesian generalized linear model (Tables S3 & S4).

Form	Diet Treatment	Mean Wing Length (mm)	Standard Deviation
Cx. pipiens form *molestus*	High	3.43	0.21
	Medium	3.22	0.10
	Low	3.06	0.11
	Extra Low	2.84	0.09
Cx. pipiens form *pipiens*	High	3.66	0.15
	Medium	3.51	0.16
	Low	3.32	0.15
	Extra Low	3.16	0.14

Figures

Figure 1: Phylogenetic Tree of Dipteran ILPs establishes *Culex ilp* orthology. Six distinct

Culex ilp genes (bold) were assigned orthology based on clustering with their closest homologs

in other species, indicated by colored boxes around each clade. *Cqui-ilp4* is a lineage-specific

gene with no homology to other Dipteran *ilp* genes (orange box). Only *Cqui-ilp5* has well

supported orthology to a *Dmel-ilp* gene (*ilp7*) (green box). Tree topology determined by TOPALI

v2.5 using a MrBayes algorithm. Posterior probabilities listed at nodes to indicate statistical

support. When the node was also supported by a maximum likelihood tree (>50) (TOPALI,

PhyML algorithm, JTT + G, 100 bootstrap runs), the bootstrap support is listed as the second

value at the node. Genbank accession numbers listed next to gene names.

Figure 2: Ovarian development up to 96 hours post adult emergence in anautogenous
pipiens **and autogenous** *molestus* **females.** Panels A-D depict ovarian development in
anautogenous, form *pipiens* females in which follicles remain in resting stage (Christopher's
Stage - IIb) until a blood meal is taken. Panels E-H depict autogenous ovarian development for
form *molestus* females provided the opportunity to mate. Panels I-L show ovarian development
in autogenous *molestus* females denied a mating opportunity. A 0.1 mm scale is shown in
panels E and H for reference. ov, ovary; mg, midgut; fb, fatbody; mt, malpighian tubules; cr,
crop.

Figure 3: Days until pupation, per treatment, for *molestus* and *pipiens*. Males and females

are shown separately. Unfilled points represent observations for individual larvae, while filled

points represent outlier observations (1.5X greater or less than the interquartile range).

Figure 4: Downregulation of *ilp2* and *foxo* in head tissues are correlated with autogeny

phenotype. qPCR was used to determine whether expression of *ilp1* (A), *ilp2* (B), *ilp3* (C), *ilp4*

(D), *ilp6* (E), *foxo* (F), and *oeh* (G), differ between bioforms and feeding conditions over the first

four days of adult development, when autogenous egg production occurs in *molestus*. Only *ilp2*

(B) and *foxo* (F) are significantly differentially expressed, as determined by a two-way ANOVA

of log2 fold change values. p values are indicated by *s as presented in the key. Fold change

gene expression determined by ΔΔCt method, normalizing first to housekeeping gene *EF1a*,

then to *pipiens* high food 0-24 hr reference sample. Data are shown on a log2 scale such that

"0" means no change vs. the reference sample. "High" and "low" indicate diet types used in

larval rearing. Error bars indicate SEM. Each data point is the mean of 3-4 biological

replicates.

Figure 5: Multiple gene expression differences in the abdomen correlate with autogeny

phenotype when biotypes and feeding conditions are compared. qPCR was used to

determine whether expression of *ilp1* (A), *ilp2* (B), *ilp3* (C), *ilp4* (D), *ilp5* (E), *ilp6* (F), *foxo* (G),

and *Vg1b* (H) differ between bioforms and feeding conditions over the first four days of adult

development, when autogenous egg production occurs in *molestus*. *ilp1* is upregulated in

molestus at 72-96 hrs post-eclosion (A). *ilp2* expression is stable between bioforms, feeding

conditions and time points (B). *ilp3* is upregulated in molestus from 48-96 hrs post-eclosion (C).

ilp4 expression is stable between bioforms, feeding conditions and time points (D). *ilp5* is

significantly downregulated at all timepoints in *molestus*, irrespective of feeding condition (E).

ilp6 expression is stable between bioforms, feeding conditions and time points (F). *foxo* is

initially downregulated in well-fed *molestus* (G). *Vg1b* is enormously upregulated in molestus,

but with a slight delay in onset in low-food conditions (H). p values are indicated by *s as

presented in the key. Fold change gene expression determined by ΔΔCt method, normalizing

first to housekeeping gene *EF1a*, then to *pipiens* high food 0-24 hr reference sample. Data are

shown on a log2 scale such that "0" means no change vs. the reference sample. "High" and

"low" indicate diet types used in larval rearing. Error bars indicate SEM. Each data point is the

mean of 3-4 biological replicates.

Figure 6: Heat Map Summary of Fold Change Gene Expression. Data from Figures 4 and 5 shown as fold changes in expression (as opposed to log2 scale in original figures) and are grouped to show genes that are up and downregulated during autogenic egg production (listed first in each grouping) vs. those that remain stable across forms (listed last in each grouping). *Vg1b* expression is shown as 1/100th of actual fold change values to allow for visualization on the same scale as other measured genes.