_N_-glycolylneuraminic acid serum biomarker levels are elevated in

breast cancer patients at all stages of disease.

Lucy K. Shewell[1#], Christopher J. Day[1#], Jamie R. Kutasovic[2], Jodie L.

Abrahams[1^], Jing Wang[1], Jessica Poole[1], Colleen Niland[2], Kaltin Ferguson[2],

Jodi M. Saunus[2], Sunil R. Lakhani[2, 3], Mark von Itzstein[1], James C. Paton[4],

Adrienne W. Paton[4], and Michael P Jennings[1*].

[1]Institute for Glycomics, Griffith University, Gold Coast, Australia

[2]UQ Centre for Clinical Research, Faculty of Medicine, The University of

Queensland, Herston, QLD Australia

[3]Pathology Queensland, The Royal Brisbane and Women's Hospital, Herston,

QLD Australia

[4]Research Centre for Infectious Diseases, Department of Molecular and

Biomedical Science, University of Adelaide, Adelaide, Australia

^Current address: Glycosciences Laboratory, Department of Metabolism,

Digestion and Reproduction, Imperial College London, London, UK

#These authors contributed equally

*To whom correspondence may be addressed:

Prof Michael P. Jennings, Institute for Glycomics, Griffith University, Gold

Coast campus, QLD 4222, Australia, +61 7 5552 7050, Australia, +61 7 5552

7050, m.jennings@griffith.edu.au

Abstract

Background

Normal human tissues do not express glycans terminating with the sialic acid

N-glycolylneuraminic acid (Neu5Gc), yet Neu5Gc-containing glycans have

been consistently found in human tumor tissues, cells and secretions and

have been proposed as a cancer biomarker. We engineered a Neu5Gc-

specific lectin called SubB2M, and previously reported elevated Neu5Gc

biomarkers in serum from ovarian cancer patients using a Surface Plasmon

Resonance (SPR)-based assay. Here we report an optimized SubB2M SPR-

based assay and use this new assay to analyse sera from breast cancer

patients for Neu5Gc levels.

Methods

To enhance specificity of our SPR-based assay, we included a non-sialic acid

binding version of SubB, $SubB_{A12}$, to control for any non-specific binding to

SubB2M, which improved discrimination of cancer-free controls from early-

stage ovarian cancer. We analysed 96 serum samples from breast cancer

patients at all stages of disease compared to 22 cancer-free controls using

our optimized $SubB2M_{-A12}$-SPR assay. We also analysed a collection of

serum samples collected at 6 monthly intervals from breast cancer patients at

high risk for disease recurrence or spread.

Results

Analysis of sera from breast cancer cases revealed significantly elevated

levels of Neu5Gc biomarkers at all stages of breast cancer. We show that

Neu5Gc serum biomarker levels can discriminate breast cancer patients from

cancer-free individuals with 98.96% sensitivity and 100% specificity. Analysis

of serum collected prospectively, post-diagnosis, from breast cancer patients

at high risk for disease recurrence showed a trend for a decrease in Neu5Gc

levels immediately following treatment for those in remission.

Conclusions

Neu5Gc serum biomarkers are a promising new tool for early detection and

disease monitoring for breast cancer that may complement current imaging-

and biopsy-based approaches.

Keywords: *N*-glycolylneuraminic acid, Neu5Gc, biomarker, breast cancer,

diagnostic

Background

Aberrant glycosylation is one of the hallmarks of cancer cells. Normal human tissues do not express glycans terminating with the sialic acid *N*-glycolylneuraminic acid (Neu5Gc) as humans express an inactive cytidine monophosphate *N*-acetylneuraminic acid (Neu5Ac) hydroxylase (CMAH) enzyme (1, 2); the only enzyme known to convert Neu5Ac to Neu5Gc. Nevertheless, Neu5Gc-containing glycans have been consistently found in human tumor tissues, cells and secretions (3-10), and have been proposed as a tumor biomarker (6, 11, 12).

Little progress has been made towards the development of a Neu5Gc biomarker-based assay for cancer detection and patient monitoring due to the lack of sufficiently sensitive and specific tools to detect this potential glyco-marker in a clinically relevant biological sample. We have improved upon the current methods for the detection of Neu5Gc (13-15) by developing a lectin with enhanced sensitivity and specificity for this glycan in the context of complex biological samples. This new lectin is derived from the B-subunit of the Shiga toxigenic *Escherichia coli* (STEC) Subtilase cytotoxin (SubAB), which recognizes α2-3 linked Neu5Gc (16). We used structure aided design to engineer this lectin to ablate the recognition of Neu5Ac and to expand the recognition from only α2-3-linked Neu5Gc to include both α2-3 and α2-6 Neu5Gc linkages to substituent sugars (17). This improved Neu5Gc-specific lectin was called SubB2M (17, 18). In a SPR-based SubB2M assay, we previously reported that the serum of ovarian cancer patients at all stages of disease has elevated levels of Neu5Gc-containing biomarkers compared to cancer-free females (19). This demonstrated the potential utility of Neu5Gc-

containing biomarkers in the early detection of ovarian cancer, for which there

is currently no universally applicable blood-based biomarker. The best

currently available biomarker for ovarian cancer is the human cancer antigen

125 (CA125), also known as MUC16, a heavily glycosylated mucin (20).

Serum CA125 levels are elevated in approximately 80% of ovarian cancer

cases at the time of diagnosis (21). However, CA125 serum levels may also

be elevated in non-malignant conditions including pregnancy, endometriosis,

ovarian cysts, pelvic inflammatory disease and in the follicular phase of the

menstrual cycle (22). As a screening tool for ovarian cancer, longitudinal

measurement of CA125 levels has been shown to improve sensitivity and

specificity for early detection. However, outcomes from the largest ovarian

cancer screening trial to date, the UK Collaborative Trial of Ovarian Cancer

Screening (UKCTOCS) show that screening including CA125 did not

significantly reduce mortality (23, 24). Hence CA125 is currently the only

approved ovarian cancer serum biomarker, limited to monitoring response to

therapy and disease recurrence in post-menopausal women (23).

Breast cancer is the most frequently diagnosed cancer among women

worldwide and is the leading cause of cancer death in over 100 countries (25).

Detection of breast cancer at the early stages is associated with better patient

outcomes including lower morbidity and lower mortality rates (26).

Mammography is currently the main screening tool for the early detection of

breast cancer; however, this method has limitations. For example, the

sensitivity of mammography in women with dense breasts is reduced from 85

% to 47-64 % (27), yet these women have an increased risk of developing

breast cancer (28). Cancer antigen 15-3 (CA 15-3) is the most widely used

serum biomarker for breast cancer, and is approved for monitoring treatment

efficacy only, due to the low sensitivity in early detection (29, 30). CA 15-3 is a

secreted form of MUC1, a heavily glycosylated mucin (31, 32). High levels of

circulating CA 15-3 have been found in breast cancer patients (33); however,

like CA125 for ovarian cancer, levels of serum CA 15-3 are also elevated in

other physiological conditions, such as pregnancy (34) and coronary heart

disease (35). Despite decades of research, there is no single serum

biomarker that has proved useful for the early detection or monitoring of

recurrence in breast cancer (36, 37).

In this study we developed an improved SubB2M-based SPR assay

methodology and used this assay to analyze serum samples collected from

breast cancer patients to determine whether detection of Neu5Gc biomarkers

may be relevant in screening for and monitoring of breast cancer.

Methods

Expression and purification of SubB2M and SubB$_{A12}$

The recombinant SubB2M and SubB$_{A12}$ proteins were expressed and purified

as previously described (17, 38). Briefly, SubB2M and SubB$_{A12}$ were

expressed in *E. coli* BL21 (DE3) cells transformed with the SubB2M or

SubB$_{A12}$ expression constructs, respectively, as His$_6$-tagged fusion proteins,

which were then purified by Ni-NTA affinity chromatography.

Glycan array analysis of SubB2M and SubB$_{A12}$

Neu5Ac/Neu5Gc glycan array slides were purchased from Z-Biotech (Aurora,

Colorado, USA). A 16-subarray slide array was used and glycan array

analysis of SubB2M and SubB$_{A12}$ was performed as described previously (18)

and as described in Supplementary Table 1. The full list of glycans on the

array can be found at https://www.zbiotech.com/neu5gc-neu5ac-n-glycan-

array.html.

Development and use of the SubB2M-$_{A12}$-SPR assay for Neu5Gc serum

biomarkers

SPR was conducted using the Biacore S200 system (GE) with immobilization

of SubB2M and SubB$_{A12}$ performed essentially as described previously (19).

For glycan analysis, SubB2M was immobilized through flow cells 2 and 3 and

SubB$_{A12}$ was immobilized through flow cell 4 (capture levels: 5000-6000

Response Units (RU) onto a series S sensor chip CM5 (GE) using the NHS

capture kit. Glycans purchased from Chemily Glycoscience (Atlanta, GA)

were analysed across a five-fold dilution series in PBS at a maximum

concentration of 20µM. Analysis was run using single cycle analysis and

double reference subtraction on the Biacore S200 evaluation software.

For glycoprotein analysis, SubB2M was immobilized through flow cells 2 and

3 and SubB$_{A12}$ was immobilized through flow cell 4. Flow cell 1 was run as a

blank immobilization. After immobilization, a start up cycle of 0.5 % normal

human serum (Sigma-Aldrich, Cat No. H4522) was run over the immobilized

SubB proteins for 10 steps of 30 seconds at 30 µL/minute flow rate to

condition the chip. A final wash of 10 mM Tris/1 mM EDTA was run for 30

seconds at a 30 µL/minute flow rate prior to beginning the data collection.

SPR analysis was performed using multi-cycle analysis and double reference

(values from flow cell 1 and 0.5 % normal human serum only) subtraction

using the Biacore S200 evaluation software. At least two independent SPR

runs were performed for each sample set.

For analysis of human sera, samples were diluted 1:200 in PBS and analyzed

in duplicate in each SPR run as described above. RU values obtained for

each serum sample with SubB$_{A12}$ (flow cell 4) were subtracted from the RU

values obtained with SubB2M from flow cells 2 and 3 and averaged to obtain

the final RUs used for conversion to GPUs. Two independent SPR runs were

performed for each sample set.

Development of Glycoprotein Units (GPUs) standard curve for normalization

of data from the SubB2M-$_{A12}$-SPR assays

SPR was conducted as described above with SubB2M immobilized through

flow cells 2 and 3 and SubB$_{A12}$ immobilized through flow cell 4. To generate

an internal calibration curve, bAGP and CA125 were combined at starting

concentrations of 15 µg/ml and 15 units/ml, respectively, in 0.5 % normal

human serum. For further detail, see Supplementary Methods.

Mass spectrometry glycomic analysis of standard glycoproteins

To confirm the presence of Neu5Gc on the glycoprotein standards (bAGP and

human CA125) *N* and *O*-glycans were released and analysed by PGC-LC-

MS/MS as previously described (39, 40). For further details, see

Supplementary Methods.

Human serum samples

Victorian Cancer Biobank samples

Serum samples from cancer-free (normal) females and serum samples from

patients with Stage I (n=12), Stage II (n=11), Stage IIIC (n=10) and Stage IV

ovarian cancer (n=14) were obtained from the Victorian Cancer Biobank and

have been described previously (19). 24 serum samples from patients with

Stage I breast cancer, 24 with Stage II breast cancer, 24 with Stage III breast

cancer and 24 with Stage IV breast cancer were also obtained from the

Victorian Cancer Biobank under application 17020. As described in our

previous study (19), 'normal' controls are defined as patients with an apparent

non-malignancy diagnosis at the time the sample is taken. The serum

samples were collected immediately pre-operatively using Serum Separation

Tubes (BD) and were processed and stored at -80 °C within two hours of

collection. The patient data and serum samples used in this project were

provided by the Victorian Cancer Biobank with informed consent from all

donors and use of the samples was approved by the Griffith University HREC

(GU Ref No: 2017/732) in accordance with the National Statement on Ethical

Conduct in Human Research. The majority of the breast cancers in the cohort

were the most common form of breast cancer, invasive ductal carcinoma. The

remainder included 8 cases of invasive lobular carcinoma and 6 cases of

mucinous carcinoma. Information for each of the ovarian cancer serum

samples used in this study can be found in our previous publication (19) while

information regarding each of the breast cancer serum samples can be found

in Supplementary Table 2.

<u>Circulating Biomarkers of Relapse in Breast Cancer (Circ.BR) cohort</u>

Circ.BR was established in 2013 as part of the Brisbane Breast Bank (41).

Patients with breast cancer who are at high risk for disease recurrence or

spread (inclusion criteria below) are followed prospectively for 5 years, with

serial collection of blood samples taken at 6 monthly intervals and tumor

tissue collected at the time of surgery. Human research ethics committees of

The University of Queensland (ref. 2005000785) and The Royal Brisbane and

Women's Hospital (2005/022) approved the study with written informed

consent obtained from each subject. Serial blood samples from 9 patients

who experienced a relapse (median 4 samples per patient, range 3-9) and 6

patients who were free from recurrence (median 7 samples per patient, range

6-8) at the time of the study were analyzed, with a median follow-up of 19.2

months for the relapse group, and 43.9 months for the recurrence-free group.

Detailed information for each patient in the Circ.BR cohort are shown in

Supplementary Table 3.

Circ.BR inclusion criteria

Invasive breast cancer, grade 3 or grade 2 (score >6) invasive cancer and

axillary lymph node positive; OR grade 1 and grade 2 (score ≤6) invasive

cancer with adverse features such as tumor size >5cm; Family history

(NBOCC group 3)/gene carrier; or previous history of breast cancer.

Statistical analysis

All statistical analyses were performed using GraphPad Prism 8.0. The mean

GPUs between normal serum samples compared to cancer patient serum

samples were analyzed by two-tailed, unpaired *t*-tests, with a *P* value of <0.05

considered significant. Optimal cut-off values from Receiver operating

characteristics (ROC) analyses were determined by maximizing the sum of

specificity and sensitivity.

Results

Development of the optimized SubB2M-$_{A12}$-SPR assay for detection of

Neu5Gc serum biomarkers.

In a previous study, we analyzed serum samples from ovarian cancer patients

with our SubB2M lectin via the highly sensitive method of SPR (19). This

assay is based on label-free detection of Neu5Gc biomarkers that bind to the

SubB2M lectin and elicit a response in SPR (see Figure 1). In this study we

aimed to improve this SPR-based assay by including a parallel analysis of all

samples with a non-sialic acid binding version of SubB called SubB$_{A12}$.

SubB$_{A12}$ has the most critical amino acid residue of the B subunit required for

binding Neu5Gc mutated from a serine to an alanine (Ser12 > Ala12) (16).

Mutation of this Ser residue abolishes interactions with the C1 carboxylate

group of sialic acid and thus the SubB$_{A12}$ mutant cannot bind any sialylated

glycans. Any binding to SubB$_{A12}$ observed with serum samples must be due to

non-sialic acid-dependent interactions of serum components with the SubB

protein, for example the binding of antibodies that may recognize the SubB

portion of the SubAB toxin. The lack of binding to sialylated glycans by the

SubB$_{A12}$ mutant has been described previously (16) and was further

confirmed herein with an analysis of SubB$_{A12}$ specificity using a Neu5Ac/Gc

glycan microarray, where negligible binding was observed to either Neu5Ac or

Neu5Gc-containing glycans by this mutant protein compared to SubB2M

(Table S1 and Figure S1). The kinetics of the interaction of SubB2M and

SubB$_{A12}$ using SPR analysis with a range of paired synthetic oligosaccharides

presenting either a Neu5Ac or Neu5Gc are shown in **Figure 1A** and confirms

the loss of all sialic acid binding by SubB$_{A12}$. In the optimized SubB2M-$_{A12}$-

SPR assay for the analysis of serum samples, the SPR Response Units

(RUs) detected with SubB$_{A12}$ for each serum sample are subtracted from the

RUs detected with SubB2M to control for any non-Neu5Gc-dependent binding

of serum components to SubB2M **(Figure 1B)**.

To further refine our SubB2M-$_{A12}$-SPR assay we established a standard curve

using a mixture of two commercially available Neu5Gc-containing

glycoproteins at known concentrations to be included as an internal control in

each SPR analysis to normalize all data from all studies to common units. As

the identity and nature of the Neu5Gc glycoconjugates detected in the ovarian

cancer patient serum samples from our 2018 study (19) is currently unknown,

we selected a combination of glycoproteins representing a high molecular

weight glycoprotein with low Neu5Gc glycosylation and a lower molecular

weight glycoprotein with high Neu5Gc glycosylation. These glycoproteins

were the human tumor antigen CA125 and bovine AGP (bAGP), respectively.

We have previously confirmed the presence of Neu5Gc on bAGP (17) and

CA125 (18), and this was reconfirmed for both of the control glycoproteins by

mass spectrometry analysis **(Figure S2)**. The standard curve generated by

this mixture of glycoproteins diluted into 0.5 % normal human serum was used

to calibrate the SubB2M-$_{A12}$-SPR assay. The response units (RUs) obtained

for each serum sample was converted to Glycoprotein Units (GPUs)

(representative standard curve shown in **Figure S3**).

In our 2018 ovarian cancer study (19) we analyzed serum samples from

subjects in Stage I (n=12), Stage II (n=11), Stage IIIC (n=10) and Stage IV

ovarian cancer (n=14) as well as serum samples from 22 cancer-free females.

In the current study, we reanalyzed this sample set using our optimized

SubB2M-$_{A12}$-SPR assay **(Figure 2)**. Figure 2A shows the data before

subtraction of non-specific binding to SubB$_{A12}$ from each serum sample while

Figure 2B shows the data after subtraction of SubB$_{A12}$ responses. As we saw

with our original analysis of this sample set (19), significantly elevated serum

Neu5Gc biomarker levels were detected at all stages of ovarian cancer

compared to cancer-free female controls.

Receiver-Operating Characteristic (ROC) analyses were performed on the

serum Neu5Gc levels detected with the optimized SubB2M-$_{A12}$-SPR assay

before and after SubB$_{A12}$ subtraction **(Supplementary Table 4, Figure S4)**.

Subtraction of binding due to SubB$_{A12}$ improved the ability of our SPR-based

assay to distinguish cancer-free individuals from ovarian cancer patients at all

stages to 100 % specificity and 100 % sensitivity.

Serum Neu5Gc levels can discriminate breast cancer patients from

cancer-free individuals with high specificity and sensitivity

We then used the refined SubB2M-$_{A12}$-SPR assay to determine whether

elevated levels of Neu5Gc biomarkers could be detected in serum from

patients with breast cancer compared to cancer-free controls. We analyzed a

collection of breast cancer serum samples across all stages of disease (24

Stage I, 24 Stage II, 24 Stage III and 24 Stage IV) with the same set of

cancer-free females used for the ovarian cancer analyses. The SubB2M-$_{A12}$-

SPR analysis **(Figure 3)** shows that significantly elevated levels of Neu5Gc

biomarkers were detected in serum samples from all stages of disease.

Detailed clinical information for each of the serum samples can be found in

Supplementary Table 2.

ROC analyses were performed to assess the ability of serum Neu5Gc levels

detected with the optimized SubB2M-$_{A12}$-SPR assay to discriminate cancer-

free females from patients from each stage of breast cancer (**Supplementary**

Table 5, Figure S5). When all stages are considered as one group with an

optimal ROC cut-off value (>10.49 GPUs), as would be the case for a

diagnostic screen, the SubB2M-$_{A12}$-SPR assay has 98.96% sensitivity and

100% specificity to distinguish patients with breast cancer across all stages of

disease from cancer-free individuals (**Figure 4**). In summary, this test

achieves 100% specificity and 100% sensitivity for patients with Stage II-IV

disease, however, due to one individual data point in the Stage I group,

below the limit of detection of our assay, the overall sensitivity did not reach

100%. These data indicate that the detection of serum Neu5Gc-biomarkers

with SubB2M has potential to detect breast cancer at all stages of disease.

Analysis of serum Neu5Gc levels using SubB2M has potential utility for

treatment monitoring in breast cancer

The Circ.BR cohort is a collection of breast cancer patients with serum

samples collected at 6 monthly intervals, allowing us the opportunity to

analyze Neu5Gc biomarker levels over the course of disease in these patients

at high risk for disease recurrence or spread. Detailed clinical information for

each patient can be found in Supplementary Table 3. Analysis of the available

serum samples from 15 cases (6 cases in remission, 9 cases with relapse)

from this cohort showed a trend for a decrease in Neu5Gc levels immediately

following the first line of treatment in the cases who did not have a tumor

recurrence, but only in some of the recurrence cases. Figure 5 shows a

representative plot from one remission case and one relapse case, with the

remaining cases shown in Supplementary Figure S6. These data demonstrate

the potential utility of the assessment of serum Neu5Gc levels to monitor

treatment response during breast cancer and warrants further exploration and

validation.

Discussion

CA125 is currently the best performing serum biomarker for ovarian

cancer, but due to its limitations, it is not currently used as a screening tool

(23). Using our new, optimized SubB2M-$_{A12}$-SPR assay, we are able to

distinguish cancer-free females from ovarian cancer patients at all stages of

disease with 100 % specificity and 100 % specificity.

The SubB2M-$_{A12}$-SPR assay is an improved version of our previously

described SubB2M-SPR assay (19). Firstly, we included subtraction of non-

specific binding of serum components to SubB$_{A12}$. The inclusion of the

SubB$_{A12}$ mutant improved the specificity and sensitivity of our assay to

distinguish cancer-free females from early-stage ovarian cancer patients.

Secondly, we now report Neu5Gc biomarker levels as GPUs. Including a

standard curve in each SPR analysis allowed us to convert Neu5Gc

biomarker levels in patient serum samples to GPUs to control for assay-to-

assay variability and to accurately compare data from independent studies.

While screening mammography is used worldwide for early detection of

breast cancer, this method has limitations, including accessibility for women in

low-income countries (42), and there is currently no single serum biomarker

for the early detection of breast cancer. Our analysis of 96 breast cancer

patient serum samples from Stages I-IV, showing that serum Neu5Gc levels

can distinguish patients with early-stage breast cancer from cancer-free

individuals with high specificity and high sensitivity, and has perfect

discriminative ability to distinguish later stage breast cancer from cancer-free

individuals, indicates that SubB2M detection of serum Neu5Gc levels has the

potential to be developed into a tool for the early detection of breast cancer. It

is important to note that this cohort, whilst comprising mostly invasive ductal

carcinoma, also included the less common invasive lobular carcinoma, and

six cases of mucinous carcinoma. Mucinous carcinoma of the breast is a rare

form of invasive ductal carcinoma, making up 2-3% of breast cancer patients.

That all of these cancers could be differentiated from cancer-free controls by

our assay supports the broad applicability of the approach. Our analysis of

serially collected serum samples from the Circ.BR cohort showed that

SubB2M detection of Neu5Gc levels also has potential to be used to monitor

response to treatment, although further investigation of its utility, using a

larger cohort, is required.

Neu5Gc is well known to be found in human cancer (in cells, tissues and

secretions) (3, 5-9, 43), and may be found in trace amounts in some healthy

human tissues (44). The source of Neu5Gc in humans is assumed to be

dietary incorporation following consumption of animal-derived food. Dietary

acquisition of Neu5Gc and metabolic incorporation into human cells forms the

basis of the 'xenosialitis' hypothesis, which proposes that autoantibodies

against Neu5Gc at the cell surface promote inflammation and pathology (see

reference 45 for a recent review) (45). The direct biochemical evidence for

dietary uptake and expression in humans involved three human subjects

consuming Neu5Gc in the form of porcine submaxillary mucin (44).

Alternatively, It has been proposed that human cancer cells can produce

endogenous Neu5Gc by providing electron donors that circumvent the CMAH

inactivating truncation found in the human enzyme (46). The current study

and our previous data (19) report elevated levels of Neu5Gc in serum can

differentiate breast and ovarian cancer patients (n=143) from cancer-free

controls, and supports the hypothesis of endogenous production of this

biomarker by tumor cells in addition to potential dietary uptake.

Our development of the purpose-engineered SubB2M lectin implemented

in the SubB2M-$_{A12}$-SPR assay provides a sensitive, specific and practical

reagent for the measurement of Neu5Gc-containing glycoproteins in serum,

which are an under-explored class of circulating tumor biomarkers. The

detection of Neu5Gc glycosylation via SubB2M may also be used in

conjunction with existing biomarker assays to improve specificity and

sensitivity and represents a promising area for future studies.

Conclusions

Minimally invasive, rapid and reliable methods for the early detection of

ovarian and breast cancer will have a significant impact on patient survival

rates. Biomarkers that can accurately and specifically monitor response to

treatment and disease progression would also drastically improve patient

outcomes. Our SubB2M lectin assay offers the opportunity to exploit

measurement of Neu5Gc biomarkers as a path to achieve these outcomes.

Declarations

Ethics approval and consent to participate

The patient data and serum samples used in this project were provided by the

Victorian Cancer Biobank with informed consent from all donors and use of

the samples was approved by the Griffith University HREC (GU Ref No:

2017/732). Human research ethics committees of The University of

Queensland (ref. 2005000785) and The Royal Brisbane and Women's

Hospital (2005/022) approved the Circ.BR study with written informed consent

obtained from each subject.

Consent for publication

Not applicable

Availability of data and materials

All data generated or analysed during this study are included in this published

article and its supplementary information files.

Competing interests

The following authors (JCP, AWP, CJD and MPJ) declare that they are

named inventors on a patent on the SubB2M technology (WO2018085888A1)

and a second patent (JCP, AWP, CJD, LKS and MPJ) for improvements to

the SubB2M test (2021901444). Both of these patents were licensed to

BARD1 Life Sciences, VIC, Australia in 2020.

Funding

This work was supported by the National Health and Medical Research

Council (NHMRC) (Program Grant 1071659 to JCP and MPJ, Project Grant

1084050 to AWP, Principal Research Fellowship 1138466 to MPJ) and the

US Department of Defense (Breakthrough Award Level 1 W81XWH-20-1-

0527 to MPJ, CJD, LKS, JLA, JRK, SRL, JCP and AWP). The Brisbane

Breast Bank and Circ.BR study are supported by the NHMRC (Program grant

APP1113867 to SRL) and the Royal Brisbane and Women's Hospital

Foundation grant (to SRL).

Author's contributions

LKS curated all serum samples, analysed and interpreted SPR data,

produced figures, contributed to design of the study and was a major

contributor to writing of the manuscript, CJD acquired and analysed SPR and

glycan array data, produced figures, contributed to conceptualisation and

design of the study and contributed to writing of the manuscript, JRK

contributed to analysing and interpreting Circ.BR data and writing of the

manuscript, JLA acquired and interpreted mass spectrometry data and

contributed to review and editing of the manuscript, JW contributed to curation

of serum samples, SPR analysis and review and editing of the manuscript, JP

acquired glycan array data and contributed to review and editing of the

manuscript, CN contributed to the Circ.BR cohort and read and approved the

final manuscript, KF contributed to the Circ.BR cohort and read and approved

the final manuscript, JMS contributed to the Circ.BR cohort and read and

approved the final manuscript SRL contributed to the Circ.BR cohort and read

and approved the final manuscript, MvI provided resources and read and

approved the final manuscript, JCP contributed to conceptualisation and

design of the study, provided SubB2M and SubB$_{A12}$ proteins and reviewed

and edited the manuscript, AWP contributed to conceptualisation and design

of the study, provided SubB2M and SubB$_{A12}$ proteins and reviewed and edited

the manuscript, MPJ conceptualised and designed the study, contributed to writing of the manuscript and provided supervision of the study.

Acknowledgements

We thank the Victorian Cancer Biobank for providing all patient data and serum samples. The Victorian Cancer Biobank, through the Cancer Council Victoria as Lead Agency, is supported by the Victorian Government through the Victorian Cancer Agency, a business unit of the Department of Health and Human Services.

References

1   Varki NM, Varki A. Diversity in cell surface sialic acid presentations: implications for biology and disease. *Lab Invest*. 2007;87(9):851-857.

2   Chou HH, Hayakawa T, Diaz S, et al. Inactivation of CMP-N-acetylneuraminic acid hydroxylase occurred prior to brain expansion during human evolution. *Proceedings of the National Academy of Sciences of the United States of America*. 2002;99(18):11736-11741.

3   Inoue S, Sato C, Kitajima K. Extensive enrichment of N-glycolylneuraminic acid in extracellular sialoglycoproteins abundantly synthesized and secreted by human cancer cells. *Glycobiology*. 2010;20(6):752-762.

4   Samraj AN, Laubli H, Varki N, Varki A. Involvement of a non-human sialic Acid in human cancer. *Front Oncol*. 2014;4:33.

5   Higashi H, Hirabayashi Y, Fukui Y, et al. Characterization of N-glycolylneuraminic acid-containing gangliosides as tumor-associated Hanganutziu-Deicher antigen in human colon cancer. *Cancer research*. 1985;45(8):3796-3802.

6   Malykh YN, Schauer R, Shaw L. N-Glycolylneuraminic acid in human

tumours. *Biochimie*. 2001;83(7):623-634.

7   Furukawa J, Tsuda M, Okada K, et al. Comprehensive Glycomics of a

Multistep Human Brain Tumor Model Reveals Specific Glycosylation

Patterns Related to Malignancy. *PloS one*. 2015;10(7):e0128300.

8   Tzanakakis GN, Nikitovic D, Katonis P, Kanakis I, Karamanos NK.

Expression and distribution of N-acetyl and N-glycolylneuraminic acids

in secreted and cell-associated glycoconjugates by two human

osteosarcoma cell lines. *Biomedical chromatography : BMC*.

2007;21(4):406-409.

9   Tzanakakis GN, Syrokou A, Kanakis I, Karamanos NK. Determination

and distribution of N-acetyl- and N-glycolylneuraminic acids in culture

media and cell-associated glycoconjugates from human malignant

mesothelioma and adenocarcinoma cells. *Biomedical chromatography*

: BMC. 2006;20(5):434-439.

10   Labrada M, Dorvignit D, Hevia G, et al. GM3(Neu5Gc) ganglioside: an

evolution fixed neoantigen for cancer immunotherapy. *Seminars in*

Oncology. 2018;45(1):41-51.

11   Marquina G, Waki H, Fernandez LE, et al. Gangliosides expressed in

human breast cancer. *Cancer research*. 1996;56(22):5165-5171.

12   Higashi H, Sasabe T, Fukui Y, Maru M, Kato S. Detection of

gangliosides as N-glycolylneuraminic acid-specific tumor-associated

Hanganutziu-Deicher antigen in human retinoblastoma cells. *Japanese*

journal of cancer research : Gann. 1988;79(8):952-956.

13   Priego-Capote F, Orozco-Solano MI, Calderon-Santiago M, Luque de

Castro MD. Quantitative determination and confirmatory analysis of N-

acetylneuraminic and N-glycolylneuraminic acids in serum and urine by

solid-phase extraction on-line coupled to liquid chromatography-

tandem mass spectrometry. *Journal of chromatography A*.

2014;1346:88-96.

14   Kitajima K, Varki N, Sato C. Advanced Technologies in Sialic Acid and

Sialoglycoconjugate Analysis. *Topics in current chemistry*.

2015;367:75-103.

15   Diaz SL, Padler-Karavani V, Ghaderi D, et al. Sensitive and specific

detection of the non-human sialic Acid N-glycolylneuraminic acid in

human tissues and biotherapeutic products. *PloS one*.

2009;4(1):e4241.

16   Byres E, Paton AW, Paton JC, et al. Incorporation of a non-human

glycan mediates human susceptibility to a bacterial toxin. *Nature*.

2008;456(7222):648-652.

17   Day CJ, Paton AW, Higgins MA, et al. Structure aided design of a

Neu5Gc specific lectin. *Scientific reports*. 2017;7(1):1495.

18   Wang J, Shewell LK, Paton AW, Paton JC, Day CJ, Jennings MP.

Specificity and utility of SubB2M, a new N-glycolylneuraminic acid

lectin. *Biochemical and biophysical research communications*.

2018;500(3):765-771.

19   Shewell LK, Wang JJ, Paton JC, Paton AW, Day CJ, Jennings MP.

Detection of N-glycolylneuraminic acid biomarkers in sera from patients

with ovarian cancer using an engineered N-glycolylneuraminic acid-

544   specific lectin SubB2M. *Biochemical and biophysical research*

545   *communications*. 2018;507(1-4):173-177.

546   20   Yin BW, Lloyd KO. Molecular cloning of the CA125 ovarian cancer

547   antigen: identification as a new mucin, MUC16. *The Journal of*

548   *biological chemistry*. 2001;276(29):27371-27375.

549   21   Bast RC, Jr., Klug TL, St John E, et al. A radioimmunoassay using a

550   monoclonal antibody to monitor the course of epithelial ovarian cancer.

551   *The New England journal of medicine*. 1983;309(15):883-887.

552   22   Goonewardene T, Hall MR, Rustin GJS. Management of asymptomatic

553   patients on follow-up for

554 ovarian cancer with rising CA-125 concentrations. *Lancet Oncology*.

555   2007;8:813-821.

556   23   Jacobs I, Menon U, Ryan Aea. Ovarian cancer screening and mortality

557   in the UK Collaborative Trial of Ovarian Cancer Screening

558   (UKCTOCS): a randomised controlled trial. . *Lancet*. 2016;387:945-

559   956.

560   24   Menon U, Gentry-Maharaj A, Burnell M, et al. Ovarian cancer

561   population screening and mortality after long-term follow-up in the UK

562   Collaborative Trial of Ovarian Cancer Screening (UKCTOCS): a

563   randomised controlled trial. *Lancet*. 2021.

564   25   Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A.

565   Global cancer statistics 2018: GLOBOCAN estimates of incidence and

566   mortality worldwide for 36 cancers in 185 countries. *CA: a cancer*

567   *journal for clinicians*. 2018;68(6):394-424.

26   Groot MT, Baltussen R, Uyl-de Groot CA, Anderson BO, Hortobagyi

GN. Costs and health effects of breast cancer interventions in

epidemiologically different regions of Africa, North America, and Asia.

The breast journal. 2006;12 Suppl 1:S81-90.

27   Kolb TM, Lichy J, Newhouse JH. Comparison of the performance of

screening mammography, physical examination, and breast US and

evaluation of factors that influence them: an analysis of 27,825 patient

evaluations. *Radiology*. 2002;225(1):165-175.

28   Boyd NF, Guo H, Martin LJ, et al. Mammographic density and the risk

and detection of breast cancer. *The New England journal of medicine*.

2007;356(3):227-236.

29   Cheung KL, Graves CR, Robertson JF. Tumour marker measurements

in the diagnosis and monitoring of breast cancer. *Cancer treatment*

reviews. 2000;26(2):91-102.

30   Harris L, Fritsche H, Mennel R, et al. American Society of Clinical

Oncology 2007 update of recommendations for the use of tumor

markers in breast cancer. *Journal of clinical oncology : official journal of*

the American Society of Clinical Oncology. 2007;25(33):5287-5312.

31   Gendler SJ, Lancaster CA, Taylor-Papadimitriou J, et al. Molecular

cloning and expression of human tumor-associated polymorphic

epithelial mucin. *The Journal of biological chemistry*.

1990;265(25):15286-15293.

32   Nath S, Mukherjee P. MUC1: a multifaceted oncoprotein with a key role

in cancer progression. *Trends in molecular medicine*. 2014;20(6):332-

342.

33   Devine PL, McGuckin MA, Ramm LE, Ward BG, Pee D, Long S.

Serum mucin antigens CASA and MSA in tumors of the breast, ovary,

lung, pancreas, bladder, colon, and prostate. A blind trial with 420

patients. *Cancer*. 1993;72(6):2007-2015.

34   Han SN, Lotgerink A, Gziri MM, Van Calsteren K, Hanssens M, Amant

F. Physiologic variations of serum tumor markers in gynecological

malignancies during pregnancy: a systematic review. *BMC medicine*.

2012;10:86.

35   Li X, Xu Y, Zhang L. Serum CA153 as biomarker for cancer and

noncancer diseases. *Progress in molecular biology and translational*

science. 2019;162:265-276.

36   Tang SS, Gui GP. Biomarkers in the diagnosis of primary and recurrent

breast cancer. *Biomarkers in medicine*. 2012;6(5):567-585.

37   Loke SY, Lee ASG. The future of blood-based biomarkers for the early

detection of breast cancer. *European journal of cancer*. 2018;92:54-68.

38   Paton AW, Srimanote P, Talbot UM, Wang H, Paton JC. A new family

of potent AB(5) cytotoxins produced by Shiga toxigenic Escherichia

coli. *The Journal of experimental medicine*. 2004;200(1):35-46.

39   Jensen P, Karlsson N, Kolarich D, Packer N. Structural analysis of N-

and O-glycans released from glycoproteins. *Nat Protoc*.

2012;7(7):1299-1310.

40   Abrahams J, Campbell M, Packer N. Building a PGC-LC-MS N-glycan

retention library and elution mapping resource. *Glycoconjugate journal*.

2018;35(1):15-29.

41   McCart Reed AE, Saunus JM, Ferguson K, Niland C, Simpson PT,

Lakhani SR. The Brisbane Breast Bank. *Open Journal of Bioresources*.

2018;5:5.

42   Rivera-Franco MM, Leon-Rodriguez E. Delays in Breast Cancer

Detection and Treatment in Developing Countries. *Breast Cancer:*

Basic and Clinical Research. 2018;12:1-5.

43   Samraj AN, Pearce OM, Laubli H, et al. A red meat-derived glycan

promotes inflammation and cancer progression. *Proceedings of the*

National Academy of Sciences of the United States of America.

2015;112(2):542-547.

44   Tangvoranuntakul P, Gagneux P, Diaz S, et al. Human uptake and

incorporation of an immunogenic nonhuman dietary sialic acid.

Proceedings of the National Academy of Sciences of the United States

of America. 2003;100(21):12045-12050.

45   Dhar C, Sasmal A, Varki A. From "Serum Sickness" to "Xenosialitis":

Past, Present, and Future Significance of the Non-human Sialic Acid

Neu5Gc. *Frontiers in immunology*. 2019;10:807.

46   Bousquet PA, Sandvik JA, Jeppesen Edin NF, Krengel U. Hypothesis:

Hypoxia induces de novo synthesis of NeuGc gangliosides in humans

through CMAH domain substitute. *Biochemical and biophysical*

research communications. 2018;495(1):1562-1566.

Figures

Figure 1. Characterization of the SubB2M-$_{A12}$-SPR assay workflow used in this study. A) SPR analysis of Neu5Ac/Neu5Gc glycan pairs. NCDI: No concentration dependent interaction with glycan up to 20 μM.

B) Optimized SubB2M-$_{A12}$-SPR assay. Serum from cancer-free (normal) females and ovarian cancer patients were analyzed by SPR with SubB2M immobilized onto the surface of the sensor chip through flow cells 2 and 3 and SubB$_{A12}$ immobilized onto the sensor surface through flow cell 4.

Figure 2. Analysis of cancer-free and Stage I-IV ovarian cancer patient serum samples with the optimized SubB2M-$_{A12}$-SPR assay. 22 serum samples from cancer-free (normal) females, 12 patients with Stage I ovarian cancer, 11 with Stage II ovarian cancer, 10 with Stage IIIC ovarian cancer and 14 with Stage IV ovarian cancer were analyzed by the optimized SubB2M-SPR assay. The mean GPUs from duplicate analyses for each serum sample determined **A)** before and **B)** after subtraction of binding due to SubB$_{A12}$ are shown. Error bars = ± 1 SD from the mean for each group. Statistical analysis was performed using two-tailed unpaired *t*-tests. **** = *P*-value <0.0001 compared to Normal. Duplicate, independent assays were performed with both showing the same trends. One representative assay is shown. **C)** Descriptive statistics of data from ovarian cancer patient serum samples and cancer-free controls.

Figure 3. SubB2M-$_{A12}$-SPR analysis of cancer-free and Stage I – IV breast cancer serum samples. A) 22 serum samples from cancer-free females, 24

Stage I, 24 Stage II, 24 Stage III and 24 Stage IV serum samples were

analyzed by SubB2M-$_{A12}$-SPR assay. The mean GPUs from duplicate

analyses for each serum sample are shown. Error bars = ± 1 SD from the

mean for each group. Statistical analysis was performed using two-tailed

unpaired *t*-tests. **** = *P*-value <0.0001 compared to Normal. Duplicate,

independent assays were performed with both showing the same trends. One

representative assay is shown. Only values above 0 are shown. **B)** Cancer-

free controls compared to early stage breast cancer samples only (Stage I

and II samples only from Figure 3A). **C)** Descriptive statistics of data from

breast cancer patient serum samples and cancer-free controls.

Figure 4. ROC curve depicting the ability of serum Neu5Gc levels

determined by the optimized SubB2M-$_{A12}$-SPR assay to distinguish

breast cancer patients from cancer-free (normal) individuals. Sensitivity%

(true positive rate; ability to detect disease) is plotted against 100 %-

specificity% (false positive rate or 100 %-true negative rate; ability to detect

lack of disease). ROC analyses were performed with the data shown in Figure

3A using Graphpad Prism 8.0.

Figure 5. Representative plots of serum Neu5Gc levels determined by

SubB2M-$_{A12}$-SPR assay for A) one remission case and B) one relapse

case from the Circ.BR cohort. The mean GPUs from duplicate analyses for

each serum sample are shown. Error bars = ± 1 SD from the mean for each

group. Two independent assays were performed with both showing the same

trends with results from one assay presented. Plots for the remaining cases

are shown in Supplementary Figure S6. Clinical information for each patient is

shown in the top right of each plot with treatment history and metastases

overlaid. ALND: Axillary lymph node dissection, ILC: Invasive Lobular

Carcinoma, Mast: mastectomy, SNB: sentinel node biopsy, WLE: wide local

excision, XRT: radiation therapy.

A

Structure	Symbol	SubB2M	SubB$_{A12}$
Neu5Ac		NCDI	NCDI
Neu5Gc		$K_D = 18.4nM\pm2.9$	NCDI
lactose		NCDI	NCDI
lactosamine		NCDI	NCDI
Neu5Ac-α2-3-lac		NCDI	NCDI
Neu5Gc-α2-3-lac		$K_D = 16.8nM\pm2.1$	NCDI
Neu5Ac-α2-6-lac		NCDI	NCDI
Neu5Gc-α2-6-lac		$K_D = 8.6nM\pm0.22$	NCDI
sLeX-Ac		NCDI	NCDI
sLeX-Gc		$K_D = 92.4nM\pm4.2$	NCDI

B

	Normal		Stage I		Stage II		Stage III		Stage IV	
Number of samples	22		12		11		10		14	
SubB$_{A12}$ subtraction	Before	After	Before	After	Before	After	Before	After	Before	After
Minimum	5.38	0.668	7.67	8.553	8.10	8.509	19.99	27.55	11.95	18.49
Maximum	9.02	5.25	16.50	20.62	14.61	17.11	34.75	59.89	75.02	120.6
Mean	7.31	3.72	11.48	13.88	10.89	12.63	26.71	41.42	30.7	47.83
SD	0.933	1.04	2.58	3.434	2.20	2.758	5.4	11.00	17.37	30.14

	Normal	Stage I	Stage II	Stage III	Stage IV
Number of samples	22	24	24	24	24
Minimum	0.4233	-3.674	10.91	19.00	29.66
Maximum	10.08	37.86	30.74	73.66	203.4
Mean	5.430	22.25	18.18	39.96	85.60
SD	2.076	8.953	5.659	13.28	51.21

A   Remission

WLE+SNB

XRT

Docetaxel,
Cyclophosphamide

GPUs

ILC
ER+PR+HER2-
Grade 2
T1N1c
Age Dx: 53.2 yrs

Time (months)

B   Relapse

Mast+ALND

Docetaxel
Carboplatin
Herceptin

Lung   RIP

GPUs

Mixed ductal/micropapillary
ER-PR-HER2+
Grade 3
Stage T3N3a
Age Dx: 51.7 yrs
Age death: 53.9 yrs
Survival: 2.2 years

Time (months)