1 Luminosity thresholds of colored surfaces are determined by their upper-limit
2 luminances empirically internalized in the visual system
3

4 Takuma Morimoto[1]*, Ai Numata[2], Kazuho Fukuda[3] and Keiji Uchikawa[4]
5
6 1: Department of Experimental Psychology, University of Oxford, Oxford, UK
7 2: Department of Information Processing, Tokyo Institute of Technology, Yokohama, Japan
8 3: Department of Information Design, Kogakuin University, Tokyo, Japan
9 4: Human Media Research Center, Kanagawa Institute of Technology, Atsugi, Japan
10
11 *Corresponding author: takuma.morimoto@psy.ox.ac.uk
12 Address: New Radcliffe House, Radcliffe Observatory Quarter, Woodstock Road, Oxford OX2 6GG
13

Keywords: Luminosity threshold, Color vision, Optimal color

Abstract

We typically have a fairly good idea whether a given object is self-luminous or illuminated, but it

is not fully understood how we make this judgement. This study aimed to identify determinants of

the luminosity threshold, a luminance level at which a surface begins to appear self-luminous. We

specifically tested a hypothesis that our visual system knows the maximum luminance level that

a surface can reach under the physical constraint that a surface cannot reflect more light than

any incident light and applies this prior to determine the luminosity thresholds. Observers were

presented with a 2-degree circular test field surrounded by numerous overlapping colored circles

and luminosity thresholds were measured as a function of (i) the chromaticity of the test field, (ii)

the shape of surrounding color distribution and (iii) the color of the illuminant of the surrounding

colors. We found that the luminosity thresholds peaked around the chromaticity of test illuminants

and decreased as the purity of the test chromaticity increased. However, the loci of luminosity

thresholds across chromaticities were nearly invariant to the shape of the surrounding color

distribution and generally resembled the loci drawn from theoretical upper-limit luminances and

upper-limit luminance boundaries of real objects. These trends were particularly evident for illuminants on the black-body locus and did not hold well under atypical illuminants such as magenta or green. These results support the idea that our visual system empirically internalizes the gamut of surface colors under natural illuminants and a given object appears self-luminous when its luminance exceeds this internalized upper-limit luminance.

1. Introduction

Most objects in the real world are visible because they reflect light. Some objects however emit light themselves; such self-luminous objects typically have a distinct appearance (e.g., traffic lights visually stand out in a scene). However, any light reaching our retina is indiscriminately encoded by three classes of cone signals regardless of whether the light is reflected from a surface or directly emitted from a light source. Thus, judging whether a given object is self-luminous presents a mathematically underdetermined problem to the visual system. The goal of this study is to reveal how our visual system overcomes this computational challenge and generates a luminous percept.

Self-luminous objects normally have a glowing appearance distinct from the appearance of illuminated surfaces. This qualitative difference was formally introduced as a mode of color appearance (Katz, 1935). The original description finely discriminates various categories, but this study concerns two modes: surface-color mode and aperture-color mode, which respectively correspond to the qualities of color appearance for an illuminated surface and a self-luminous object. Color appearance has mostly been studied in the surface-color mode; only a limited number of studies have investigated the nature of the aperture-color mode (e.g., Uchikawa, Uchikawa & Boynton, 1989).

One common approach is to measure the transition luminance between surface-color mode and

aperture-color mode, which is known as the *luminosity threshold.* Past studies have investigated

what factors might govern this threshold. In an early study, Ullman (1976) extensively discussed

potential determinants of luminosity thresholds: highest intensity in a scene, absolute intensity of

the stimulus, local or global contrast, intensity comparison with the average intensity of the scene,

and lightness computation that emphasizes a transient intensity change over space while ignoring

a gradual intensity change. It was concluded that although each factor plays a role, none of these

factors are sufficient to predict luminosity thresholds. Bonato & Gilchrist (1994) reported

quantitative observations that an achromatic surface appears luminous when it has roughly 1.7

times the luminance of a surface that would be perceived as white. For chromatic stimuli, it was

repeatedly shown that luminosity thresholds were negatively correlated with stimulus purity in a

series of studies (Evans, 1959; Evans & Swenholt, 1967; Evans & Swenholt, 1968; Evans &

Swenholt, 1969). Speigle & Brainard (1996) measured luminosity thresholds using real colored

objects placed under illuminants of different color temperatures. They supported Evans's

consistent observation about the chromaticity-dependent nature of luminosity thresholds and

showed that the color of the illuminant also affects luminosity thresholds. More recently, Uchikawa

et al. (2001) pointed out that the brightness of colored surfaces rather than their physical

luminance is highly correlated with the luminosity thresholds of colored surfaces. These studies

well characterized the properties of a test stimulus and of surrounding contexts that have an

impact on luminosity thresholds.

One important open question in the field is whether our visual system bases self-luminous

judgements purely on heuristics that extract statistics from the external world. Such a strategy is

prevalent in many other visual judgements. For example, the famous anchoring theory determines a reference based on simple statistics in a given scene (i.e., the highest luminance in a scene is defined as white) which has been successful in explaining empirical results involving lightness judgements (Gilchrist & Bonato, 1995; Gilchrist et al. 1999). If our visual system takes a heuristic-based strategy, luminosity thresholds should be susceptible to scene content – for example a combination of surface reflectances that happen to be present in a scene. Alternatively, the visual system might additionally use an internal reference for luminosity judgements that is more robust to the variety of available scene contents. For instance, it was shown that our visual system might use statistical regularities about the possible range of surface colors and illuminant colors (Judd et al., 1964) as a prior to solve an ill-posed problem such as color constancy (Maloney & Wandell, 1986). Also, there are suggestions that color contrast and assimilation arise simply from learning statistical regularities in external environments (Lotto & Purves, 2000; Long & Purves, 2003). The success of these prior-based approaches implies a possibility that humans might take a similar strategy when making self-luminosity judgements.

Thus, one primary focus in this study is to reveal whether luminosity thresholds are determined purely based on rigid heuristics that rely on the statistics of external stimuli or whether the visual system additionally uses internal references to make a luminosity judgement. We specifically built a hypothesis based on the latter view: the visual system internalizes the physical gamut of surface colors under various illuminants and refers to this knowledge when judging whether a given surface is self-luminous. This physical gamut of surface colors is visualized by optimal colors (MacAdam, 1935a; MacAdam 1935b), which will be detailed in the General Method section. In short, optimal colors are the colors with the highest luminance that can be produced by reflected light under a given illuminant, for each possible chromaticity. It is assumed that the visual system

estimates the illuminant color and chooses the gamut under the estimated illuminant. In a more

general sense, this hypothesis could be treated as a Bayesian framework where the visual system

monitors the scene illuminant and selects which prior to use based on the estimated illuminant.

This hypothesis was specifically designed based on observations made in a series of color

constancy experiments (Uchikawa et al, 2012; Fukuda & Uchikawa 2014; Morimoto et al, 2016;

Morimoto et al, 2021). In these studies, we developed a model for illuminant estimation that

operated on the assumption that the visual system internalizes the gamut of surface colors under

various illuminants (i.e., distribution of optimal colors) and the model accounted for observers'

estimations of illuminants reasonably well in a variety of conditions. One interpretation of

luminosity thresholds is that the visual system takes the upper-limit boundary of surface colors as

the point beyond which objects are self-luminous. Thus, we speculated that the loci of luminosity

thresholds measured under different illuminants might resemble the locus of optimal colors.

We note that this study is also built on previous efforts made by Evans (1958) and Speigle and

Brainard (1996) for the following reasons. Evans (1958) in part of his analyses first made a

comparison between luminosity thresholds and the optimal color locus, which is the primary

purpose of this study. Though it was concluded that luminosity thresholds do not well align with

optimal color locus, the research used a simple stimulus configuration where a colored surface

was presented with a uniform background. Thus, we believe it is worth testing the accountability

of the optimal color model under a wider variety of conditions where richer cues to the illuminant

are provided. Speigle and Brainard (1996) was the first study to directly suggest that luminosity

thresholds are strongly influenced by illuminant color. They further modeled observers' luminosity

thresholds using the upper-limit luminance of physically plausible surfaces in the real world

estimated by a linear model that uses basis reflectance functions obtained via a principal

component analysis of Munsell papers. The locus obtained via their method corresponds to a

practical upper-limit luminance in the real world as opposed to a theoretical upper-limit luminance

defined by optimal colors. Nevertheless, we believe that their suggestion shows conceptual

similarity to our hypothesis.

In this study, we conducted three experiments to test our hypothesis. In each experiment, we

presented a 2-degree circular colored test field surrounded by many overlapping colored circles.

We measured luminosity thresholds as a function of test chromaticities. Experiment 1 was

designed to test the degree to which luminosity thresholds were influenced by the color statistics

of surrounding stimuli, in this case the geometry of the color distribution. In Experiment 2, we

tested the effect of the illuminant as well as the shape of the surrounding color distribution to

reveal whether the luminosity threshold loci agree with the optimal color locus under different

illuminants (3000K, 6500K and 20000K). In Experiment 3, we measured the loci of luminosity

thresholds under atypical illuminants (magenta and green) to investigate whether the loci of

luminosity thresholds over chromaticities might differ between chromatically typical and atypical

illuminants.

2. General Method

2.1. Computation of physical upper-limit luminance at a given chromaticity

We can compute the theoretical upper-limit luminance at each chromaticity by calculating the

chromaticity and the luminance of its optimal colors. Here we provide a basic idea of optimal color,

but a more detailed description is available elsewhere (e.g., Uchikawa et al., 2012, Morimoto, et

al., 2021). An optimal color is a hypothetical surface having a steep spectral reflectance function

as shown in Figures 1 (a) and (b). There are two types (band-pass and band-stop) and they can

have only 0% or 100% reflectances. Changing λ_1 and λ_2 generates numerous optimal colors ($\lambda_1 <$

λ_2). To give concrete examples we generated three illuminants of black body radiation: 3000K,

6500K and 20000K. Then 7,644 optimal colors were rendered under these illuminants as shown

by small dots in Figures 1 (c) and (d). Panel (c) shows L/(L+M) in MacLeod-Boynton (MB)

chromaticity diagram (MacLeod & Boynton, 1979) vs luminance distributions. Panel (d) shows

$\log_{10}$S/(L+M) vs. luminance distributions. To calculate cone excitations, we used the Stockman &

Sharpe cone fundamentals (Stockman & Sharpe, 2000).

In the real world surface reflectances must be less than 1.0 at any wavelength due to physical

constraints, and thus an optimal color has a higher luminance than any other surface that has the

same chromaticity. Thus, no real surface can exceed this optimal-color distribution. To show this

concretely, in Figures 1(c) and (d) we plotted 49,667 objects in the standard object color spectra

database for color reproduction evaluation (SOCS, ISO/TR 16066:2003).

From optimal color distributions we see that the physical upper-limit luminance is dependent on

the chromaticity. The peak of an optimal color distribution always corresponds to a full-white

surface (1.0 reflectance across all wavelengths), which thus corresponds to the chromaticity and

intensity of the illuminant itself (so-called white point of the illuminant). For this reason, when the

color temperature of the illuminant changes, the whole optimal color distribution shifts towards

the chromaticity of the illuminant without drastically changing its overall shape. Optimal colors

with a higher purity have lower luminance, as they have a narrower-band reflectance, and

consequently the distribution spreads out as the purity increases. Importantly, once all optimal

colors are calculated, we can look for the physical upper-limit luminance at any chromaticity by

looking for the luminance of the optimal color at that chromaticity. Interestingly it is notable that

the distribution of real objects (SOCS dataset) shows a somewhat similar shape to the optimal

color distribution.

Figure 1: (a), (b) Example optimal colors of band-pass and band-stop types, respectively. (c), (d) L/(L+M) vs.

luminance and $\log_{10}S/(L+M)$ vs. luminance distributions, respectively, for optimal colors and the SOCS

reflectance dataset rendered under 3000K, 6500K and 20000K. [one-column figure]

2.2. Estimation of the upper-limit luminance at a given chromaticity for real

surfaces

The theoretical upper-limit luminance can be computed through the calculation of optimal colors,

but the upper-limit luminance for real objects needs to be estimated. Thus, we analyzed 49,672

surface reflectances from the SOCS reflectance database. This dataset includes reflectances

from a wide range of categories of natural and man-made objects: "photo" (2304 samples),

"graphic" (30,624), "printer" (7856); "paints" (229); "flowers" (148); "leaves" (92); "faces" (8049);

and "Krinov datasets" (370) including natural objects which were measured in a separate study

(Krinov, 1947). We then excluded reflectances that contained a value higher than 1.0 at any

wavelength as they might include fluorescent substances. As a result, one reflectance from the

printer category and 4 reflectances from the paints category were excluded.

The remaining 49,667 surfaces were then rendered under 6500K and their chromaticity and

luminance were calculated. The luminance value was normalized by that of a full-white surface

(100% reflectance at any wavelength). As shown in Figure 2 (a), we plotted the chromaticity of all

surfaces on the MacLeod-Boynton chromaticity diagram, where L/(L+M) is the horizontal axis and

$\log_{10}$ S/(L+M) is the vertical axis. We defined a grid of 25×25 bins and classified 49,667 colors

into corresponding bins. Then, for each bin, the maximum luminance across all colors that belong

to the bin was defined as the upper-limit luminance of real objects. This procedure was repeated

for all 625 bins. The upper-left and lower-left subpanels in panel (b) show the upper-limit

luminance for optimal colors (for comparison purposes) and for real objects. As seen here the loci

of the upper-limit luminance for real objects were not smooth. We assumed that this is an artifact

due to the limited availability of reflectance samples in the database rather than the nature of

reflectances of real objects. Thus, we smoothed the upper-limit luminances by spatial filtering with

3×3 convolutional filters (each pixel has the value of 1/9). The lower-right subpanel depicts the

smoothed data. Note that this upper-limit luminance heatmap is dependent on the color of the

illuminant. Thus we repeated the same procedure for other black-body illuminants with color

temperatures from 3000K to 20000K with 500 K steps. Both the optimal color locus and real object

locus unsurprisingly peak at the chromaticity of the illuminant shown by the red cross symbol. The

upper-limit luminance of real objects decreases more sharply as the stimulus purity increases

than that of optimal colors. We can refer to these look-up-tables to find the upper-limit luminance

of real objects for an arbitrary chromaticity under illuminants of a range of color temperatures.

Note that this upper-limit luminance of real objects corresponds to the proposed model by Speigle

and Brainard (1996) at a conceptual level though they estimated the boundary using a linear

model rather than the 'big data' approach taken here.

Figure 2: How to estimate the upper-limit luminance for real objects using the SOCS spectral reflectance

dataset. A 25×25 grid was first drawn on the MacLeod-Boynton chromaticity diagram. For each grid bin, we

searched for the surface that has the highest luminance as shown in the right part of panel (a), which was

defined as the upper-limit luminance for that chromaticity bin. Panel (b) shows the locus of upper-limit

luminance for optimal color, real objects (raw), and real objects (smoothed) under 6500K illuminant. The

lightness indicates the upper-limit luminance value for the chromaticity bins. The pale green color indicates

that there is no data in that bin. [one-column figure]

2.3. Observers

Four observers (KK, MI, TM and YK) participated in Experiment 1. KK and YK were also recruited

for Experiment 2 as well as two new observers (KS and NT). KK, KS and YK participated in

Experiment 3. Observers except for KS were naïve to the purpose of all experiments. Observers'

ages ranged between 22 and 57 (*mean* 31.4, *s.d.* 13.2). Observers were all Japanese. All

observers had corrected visual acuity and normal color vision as assessed by Ishihara pseudo-

isochromatic plates. Before the experiments, informed consent was obtained from each observer.

Observers were offered to take several breaks during the experiments, and observers could stop

the participation at any point during the experiments.

2.4. Stimulus Configuration

The stimulus configuration is shown in Figure 3. The color distribution of the surrounding stimuli

and the chromaticities used for the test field are detailed in each experimental section. The spatial

pattern was shuffled for each trial.

Figure 3: Example of stimulus configuration. The center circle is the test stimulus, and its luminance was

adjusted by observers. Each circle had a diameter of 2 degrees in visual angle, and the whole image

subtended 15×15 degrees. The surrounding color distribution is detailed in each experimental section. [one-

column figure]

2.5. Apparatus

Data collection was computer-controlled and all experiments were conducted in a dark room. Stimuli were presented on a cathode ray tube (CRT) monitor (BARCO, Reference Calibrator V, 21 inches, 1844 × 1300 pixels, frame rate 95Hz) controlled with ViSaGe (Cambridge Research Systems), which allows a 14-bit intensity resolution for each of the RGB phosphors. We performed gamma correction using a ColorCAL (Cambridge Research Systems) and spectral calibration was performed with a PR650 spectroradiometer (Photo Research inc.). Observers were positioned 114 cm from the CRT monitor and the viewing distance was maintained with a chin rest. Observers were asked to view the stimuli binocularly.

2.6. General procedure

Observers first dark-adapted for 2 mins and then adapted to an adaptation field for 30 seconds. The adaptation field was the full uniform screen that had either a chromaticity of 6500K (Experiments 1 and 3) or the chromaticity of the test illuminant (Experiment 2), and in either case the luminance was equal to the mean luminance value across surrounding stimuli. Then, the first trial began. We drew surrounding stimulus circles so that they had a specific color distribution as detailed in each experimental section. The 2-degree circular test field was presented at the center of the screen. The test field was never occluded by surrounding stimuli. The observers' task was to adjust the luminance of the test field to the level at which the surface-color mode changed to the aperture-color mode using a keyboard with three possible luminance steps (± 0.5, ± 1.0, or ± 5.0 cd/m^2). The ambiguity regarding the criterion to judge the transition between surface-color mode and aperture-color mode was reported in a past study (Speigle & Brainard, 1996, Uchikawa et al., 2001). This is mainly because the transition is not sharp, and there is a range that a surface

can appear a mixture of surface-color mode and aperture color mode. Considering this reported

ambiguity, we instructed observers as follows: "Your task is to adjust the luminance of the center

test field so that the test field appears to be at the midpoint between the upper-limit of the surface

color mode and the lower-limit of the aperture color mode." The upper-limit of the surface color

mode and the lower-limit of the aperture color mode were described to observers as the limit at

which the test field completely appears as an illuminated surface and the limit at which the test

field completely appears as a light source, respectively. All observers agreed that this was a

reasonable judgment. Also, we note that our criterion is analogous to criteria used in past studies

(Bonato & Gilchrist,1994; Evans, 1959; Evans & Swenholt, 1967, 1968, 1969; Speigle & Brainard,

1996; Ullman, 1976). During the experiments, observers were instructed to view the whole

stimulus rather than fixate at a specific point to avoid local retinal adaptation. The initial luminance

value for the test field was randomly chosen from 2.0, 5.0, 8.0, 11.0, 14.0, 17.0, 20.0, 23.0, 26.0

and 29.0 cd/m^2. Specific experimental conditions are detailed in each experimental section.

3. Experiment 1

3.1 Surrounding color distribution, test illuminant and test chromaticity

In a natural scene, the colors of objects tend to cluster around the white point of the illuminant

and the density of colors decreases as purity increases. Consequently, the color distribution tends

to form a mountain-like shape as shown in Figure 1 (d). The aim of Experiment 1 was to

investigate how the loci of luminosity thresholds change when thresholds are measured in a scene

that has an atypical color distribution shape. In an extreme case where observers rely purely on

internal criteria to judge the self-luminosity of a surface, luminosity thresholds should not change

at all regardless of the surrounding color distribution. However, in contrast if observers make a

self-luminous judgement using surrounding colors, for example by estimating the upper

luminance boundary from the surrounding distribution, luminosity thresholds should largely

change depending on the shape of the surrounding color distribution.

Figure 4 (a) shows the five surrounding color distributions used in Experiment 1. The 6500K

illuminant on the black-body locus was chosen as the test illuminant in this experiment. We first

defined the *natural* color distribution in the upper-left subpanel and then transformed the

distribution to generate 4 atypical color distributions (*reverse*, *flat*, *slope+* and *slope-*) in the

following ways. First, to construct the *natural* color distribution, we started with a dataset of 574

spectral reflectances of natural objects (Brown, 2003). Out of the 574 reflectances, 516

reflectances were inside the chromaticity gamut of the experimental CRT monitor when rendered

under the 6500K test illuminant. All stimuli were presented via a ViSaGe, which had the technical

constraint that only 253 colors could be simultaneously presented. Thus, we selected 253

reflectance samples out of 516 reflectances. The reflectance spectra in the Brown dataset were

clustered around a white point in a chromaticity diagram; therefore, if we randomly sample from

those spectra, it generates a biased distribution with more data points around the white point.

Thus the 253 reflectances were selected such that, when rendered under 6500K, they were

approximately spatially and uniformly distributed across a chromaticity diagram: L/(L+M) and

S/(L+M).

To generate the other color distributions (*reverse*, *flat*, *slope+* and *slope-*), we independently

scaled each of the 253 reflectances by a scalar value to manipulate the luminance while keeping

the chromaticity constant. The inserted image in each subpanel shows an example of surrounding

stimuli that has the corresponding color distribution. Note that the spatial layout of the surrounding

stimuli was shuffled for each trial. For all distributions, the intensity of the test illuminant was

determined so that a full-white surface (i.e., 100% reflectance across all visible wavelengths) had

a luminance of 35.0 cd/m^2 under the test illuminant. We note that surrounding colors had relatively

low luminance values. For the test field to appear self-luminous, the test field needs to have a

substantially higher luminance than the surrounding colors. Thus, the choice of surround

luminances was unavoidable in order to ensure that observers could make a satisfactory

adjustment at any tested chromaticities within the luminance range allowed by our experimental

monitor.

For the center test field, we chose 9 reflectances out of the 253 so that they fell closely along the

black-body locus when placed under a 6500K illuminant. The panel (b) in Figure 4 shows these

9 test chromaticities at which luminosity thresholds were measured. The chromaticity of two

reflectances are slightly off from the black-body locus. This is because we could not find

reflectance samples that exactly fall on the locus.

Figure 4: (a) 5 color distribution sets for surrounding stimuli. An inserted image at the top shows an example

stimulus configuration for each color distribution. The vertical solid black line indicates the L/(L+M) value of

6500K test illuminant. Black cross symbols indicate mean cone response across all 253 surrounding stimuli.

(b) 9 test chromaticities at which the luminosity thresholds were measured. [1.5-column figure]

3.3 Procedure

One block consisted of 9 settings to measure thresholds at all 9 test chromaticities in random

order. There were 5 blocks in each session to test all 5 distribution shapes. The order of

distribution condition was randomized. All observers completed 20 sessions in total (i.e., 20

repetitions for each data point). They completed 10 sessions per day and thus the experiment

was conducted in two days.

3.3 Results

Figure 5 shows the results for Experiment 1. Colored symbols with error bars indicate each observer's setting. Each data point is the average across 20 repetitions. The average across 4 observers is shown as black circles. There was some variation across individuals. Furthermore, the experimental design was to try to collect reliable data from a small number of participants. Thus, we discuss results individually. The magenta circles and the line show luminances of optimal colors at test chromaticities when rendered under the test 6500K illuminant (the optimal color locus). In other words, if the visual system uses the optimal color to judge whether a surface emits a light, the observer's settings should match the magenta line. The blue circles and the line show a smoothed upper-limit luminance locus of real objects, estimated from the SOCS reflectance dataset as shown in Figure 2, which more rapidly decreases as it gets away from the white point than the optimal color locus does. For simplicity, we hereafter refer to the magenta and blue lines as predictions of the optimal color model and the real object model, respectively.

First, the loci of luminosity thresholds for all observers had a mountain-like shape regardless of surrounding color distribution. The loci generally peaked around the chromaticity of the test illuminant (the vertical black solid line) and the luminosity thresholds decreased as the test chromaticity moved away from the white point. Although there were some individual differences, especially in the overall setting level (e.g., KK generally had higher thresholds than others) and in the peak chromaticity, the luminosity thresholds generally seem to more resemble the prediction of the optimal color model than that of the real object model in this experiment. This is consistent with the hypothesis that the visual system knows the upper boundary of the optimal color distribution and judges that a given surface is self-luminous when its luminance exceeds the luminance of optimal colors.

Figure 5: Colored square symbols indicate averaged settings across 20 repetitions for each observer. The

error bars represent ±1 S.E. across 20 repetitions. The black circles represent settings averaged across

observers (n = 4). The magenta circles denote the luminance of the optimal color at the test chromaticities

and thus indicate the physical upper-limit luminance. The blue line shows the upper-limit luminance of real

objects estimated from the SOCS reflectance dataset. The vertical solid line shows the chromaticity of the

test illuminant (6500K). The black cross symbol indicates the mean LMS value across surrounding stimuli.

[1.5-column figure]

To quantify the similarity between observers and models, we calculated Pearson's correlation

coefficient between observer settings and model predictions over the 9 test chromaticities.

Figure 6 shows summary matrices of the correlation coefficients. We calculated correlation

coefficients for each observer and discuss them on an individual basis.

The magenta and blue symbols represent the optimal color model and the real object model,

respectively. In addition, we evaluated a model which judges the surface as self-luminous when

its luminance exceeds that of the surrounding color distribution. The luminosity thresholds

estimated from such a model should show much similarity to the shape of the surrounding color

distribution. For example, in the *reverse* condition, the luminosity threshold should be lowest at

the white point and increase as the saturation of the test stimulus increases. This model is

labelled as the "surrounding color" model in Figure 6. Note that this is a simplistic model and we

are not trying to claim that the visual system takes such a strategy. Instead, our goal here is to

build a framework in which we quantitatively predict an observer's behavior if she/he judges the

luminosity thresholds solely based on surrounding stimuli presented in each trial without using

any prior about the statistics of the real world.

The cyan star symbols in some cells indicate the highest correlation-coefficient value across the

3 tested models. The cyan arrows below each subpanel indicate the model that received the

highest number of cyan stars across the 4 observers.

Overall, since the observer settings are stable across all distribution conditions, the correlation

coefficient patterns are also similar between the optimal color and real object models whose

predictions are both not affected by surrounding colors. However, the correlation coefficients for

the surrounding color model strongly depend on distribution condition as predicted. Specific

trends are as follows. For observers KK and YK, the loci of the luminosity thresholds showed

the highest correlation with the optimal color model for all distributions. For TM, the real object

model was the best predictor in all distributions except for the *flat* condition. For MI, the optimal

color model showed the highest correlation for *reverse*, *flat* and *slope+* conditions while the real

object model showed the highest correlation for *natural* and *slope-* conditions. If we summarize

these trends based on the number of cyan arrows each model received, the optimal color model

is the best predictor in Experiment 1.

Figure 6: The matrices of Pearson's correlation coefficients calculated between observer settings and

model predictions over 9 test chromaticities. Each subpanel represents each distribution condition (*natural*,

reverse, *flat*, *slope+* and *slope-*). The color of each individual cell indicates the correlation coefficient as

denoted by the color bar. The cyan star symbol indicates the highest correlation coefficient across the 3

models. The cyan arrows at the bottom of each subpanel show the model that received the highest

number of cyan star marks, indicating a good candidate model of human observers' strategy to judge the

self-luminosity of a surface. [one-column figure]

The major finding in this experiment is that the loci of luminosity thresholds are nearly invariant

regardless of the shape of the surrounding color distribution. This result supports the idea that

observers use an optimal color distribution as an internal reference to determine the luminosity

thresholds. In the Appendix, we also provide two other alternative models that predicts luminosity

thresholds based on post-receptoral signals or cone signals of the test field alone, but it was

shown that these models did not predict the luminosity threshold well in Experiment 1. In

Experiment 2, we tested whether this observation holds under different illuminants which shift the

peak of the optimal color distribution as shown in Figure 1. If the visual system indeed uses

optimal colors, changes in luminosity thresholds should reflect changes in the optimal color

distribution.

4. Experiment 2

4.1 Surrounding color distribution, test illuminant and test chromaticity

We employed *natural*, *reverse* and *flat* distributions of surrounding colors. For test illuminants, we

used 3000K, 6500K and 20000K on the black-body locus. Out of the 253 reflectances we used in

Experiment 1, only 180 samples were inside the chromaticity gamut of the experimental CRT

monitor under all test illuminants and those 180 samples were used as surrounding stimuli in

Experiment 2. Panel (a) in Figure 7 shows all 9 test surrounding conditions (3 distributions × 3

test illuminants). Though we found that surrounding color distribution has no systematic effects

on luminosity thresholds in Experiment 1, we again manipulated the distribution shapes in

Experiment 2 to investigate if this finding held under different illuminants.

We then selected 15 surface reflectances from the 180 reflectances. Panel (b) shows the 15 test

chromaticities when rendered under each test illuminant at which the luminosity threshold was

measured.

Figure 7: (a) 9 color distributions for surrounding stimuli (3 test illuminants × 3 distributions). Inserted image shows an example stimulus configuration. The vertical solid black line indicates the L/(L+M) value of each test illuminant. Black cross symbols indicate mean cone response values across the 180 surrounding stimuli. (b) 15 test chromaticities at which the luminosity threshold was measured. [1.5-column figure]

4.2 Procedure

One block consisted of 15 consecutive settings to measure thresholds for all test chromaticities presented in random order. There were 9 blocks in one session to test all conditions (3 illuminants × 3 distributions). The order of conditions was randomized. All observers completed 10 sessions in total. The experiment was conducted in three days.

4.3 Results

The black line in Figure 8 shows the mean setting across 4 observers. The rest of the data

presentation follows the results in Experiment 1. For clarity, only the averaged setting is shown

here, but the individual observers' data is presented in Figure A1 in the Appendix.

First, the mean settings showed that the loci of luminosity thresholds were again mountain-like in

shape, and the influence of the shape of the surrounding color distribution was almost absent,

supporting the findings in Experiment 1. It is also noticeable that the peak chromaticity of the

mean setting in each panel shifted towards the illuminant chromaticity shown as vertical solid

lines.

Figure 8: The black circle symbols represent averaged observer settings (n = 4). The error bars represent

±1 S.E. across the 4 observers. The optimal color loci are plotted as magenta circles. The blue circles

show the upper-limit luminance of real objects. The red, black and blue vertical solid lines show the

chromaticities of the 3000K, 6500K and 20000K test illuminants, respectively. The black cross symbol

indicates the mean LMS value across surrounding stimuli. Individual observer data is shown in the

Appendix. The region surrounded by a rectangle in the 20000K condition is further discussed in Figure 9.

[1.5-column figure]

It should be noted that the peak chromaticity of the luminosity threshold loci for 20000K was

slightly shifted to the right along the L/(L+M) dimension from the chromaticity of the test illuminant.

This trend was generally consistent across observers as shown in Figure A1 (Appendix). One

potential reason could be that observers misestimated the illuminant color from the surrounding

colors. Human color constancy is often imperfect, and thus we speculated that observers'

luminance settings might better agree with the optimal color or real object model rendered under

an illuminant estimated by each observer instead of a ground-truth illuminant (20000K). In fact,

misestimates of illuminant color was also reported to be an important factor in predicting

luminosity thresholds by Speigle and Brainard (1996). The estimated illuminant is typically

measured using a technique such as achromatic adjustment (Brainard, 1998), but these data

were not collected in this study. Thus, we assumed that the peak chromaticity of observer settings

indicated the observer's estimated illuminant.

We first calculated the chromaticities of illuminants from 3000K to 20000K in 500K steps. Then,

for each observer and for each condition independently, we searched for the color temperature

that had the closest chromaticity to the peak chromaticity of the luminosity thresholds. Table 1

summarizes the color temperatures of the estimated illuminants in each condition. In the 3000K

condition, estimated illuminants matched the ground-truth color temperature for most observers.

For 6500K, there was a slight variation across observers. It is notable that in the 20000K condition

observers estimated color temperatures substantially lower than those of the ground-truth,

meaning illuminant color was estimated to be less blue. This could be because perceptual

differences between stimuli rendered under 20000K and under 6500K is smaller than the

difference between 3000K and 6500K, as perceptual sensitivities are reported to be worse for

bluish illuminants and surfaces (Pearce et al., 2014; Winkler et al., 2015).

Table1: Estimated illuminant by each observer judged from the chromaticity at which luminosity thresholds

peaked. The top row shows the color temperatures of ground-truth illuminants, and the other numbers

indicate the color temperature of estimated illuminants.

		3000K	6500K	20000K
Natural	KK	3000	5500	10500
	KS	3000	7000	8500
	NT	3000	5000	10500
	YK	3000	5500	12000
Reverse	KK	3000	6500	7500
	KS	4000	7000	7500
	NT	3000	5500	12000
	YK	3500	5000	10500
Flat	KK	3000	7000	8500
	KS	3000	5000	8500
	NT	3000	5500	12000
	YK	3000	6500	12000

509 Then, we drew optimal color loci under these estimated color temperatures. This concept is

510 depicted in Figure 9. Intuitively speaking this procedure allows us to estimate an optimal color

511 locus that the observer presumably used during the task, so that the peak of this new optimal

color locus coincides with the peak of the measured locus of luminosity thresholds. The cyan

curve shows the optimal colors under the estimated illuminant and seems to predict mean

observer settings better than the optimal color locus under the ground-truth illuminant (20000K).

Figure 9: Optimal color models based on the ground-truth illuminant (magenta) and based on estimated

illuminants for the averaged observer setting (cyan). It is shown that observers' settings are better

explained by the optimal color model that allows misestimation of illuminants by observers. [one-column

figure]

Figure 10 depicts the correlation coefficient matrices for all conditions. We compared

correlations from 5 models: (i) the optimal color model and (ii) the real object model under the

ground-truth illuminant, (iii) the optimal color model and (iv) the real object model under the

estimated illuminant, and (v) the surrounding color model. Again, the cyan star symbol in some

cells denotes the highest correlation across the 5 models for that participant. The cyan arrows

below each subpanel point to the model that has the highest number of cyan stars – the overall

best model for that condition.

Figure 10: The matrices of Pearson's correlation coefficient calculated between observer settings and

model predictions over 15 test chromaticities. The cyan star symbol indicates the highest correlation

coefficient across the 5 models. The cyan arrows at the bottom of each subpanel show the model that

received the highest number of cyan star symbols. [1.5-column figure]

Overall, the surrounding color model does not show high correlation with observer settings in

any condition, agreeing with the trends in Experiment 1. The optimal color model and the real

object model seem to show high correlation, and it depends on the condition which model

correlates better. For the *natural*-3000K condition, the highest correlation was found for the real

object models, consistently across all observers. For the *reverse*-3000K condition, all observers

except NT were best correlated with the optimal color model under the estimated illuminant

while for the *flat*-3000K condition the votes were split between the optimal color and real object

models. For *natural*-6500K, KK and NT were well predicted by the optimal color model under

the ground-truth illuminant, but the other two observers were better correlated with the real

object model. For the *reverse*-6500K condition, the optimal color and the real object model both

showed high correlations. The real object model, when used with the estimated illuminant,

predicted observer settings best for the *flat*-6500K condition. It is notable that for the 20000K

condition, the optimal color model under the estimated illuminant was consistently the best

predictor. The optimal color model under the ground-truth illuminant also shows much lower

correlations, suggesting that observers' misestimates of the illuminant play a role in predicting

luminosity thresholds. In summary both the optimal color model and the real object model

showed fairly good agreement with human observers' settings.

Experiments 1 and 2 collectively suggested that both the optimal color locus and the real object

locus seemed to be good candidate determinants of luminosity thresholds. We also note that

two other alternative simplistic models again did not predict luminosity thresholds well in

Experiment 2 (shown in Appendix). One noteworthy feature in Experiments 1 and 2 is that we

used illuminants on the blue-yellow axis that are typically found in natural environments. We

also used chromaticities on the black-body locus for the test field. If we assume that the visual

system learns the locus of optimal color distribution or real object distribution by observing

colors in natural environments, the luminosity thresholds under atypical illuminants may not

agree well with the prediction of the optimal color model or the real object model. We directly

tested this hypothesis in Experiment 3.

5. Experiment 3

Experiment 3 tested whether luminosity thresholds resembled the optimal color locus under

atypical illuminants. We also chose a wider range of test chromaticities from the black-body

locus and a locus that is orthogonal to the black-body locus.

5.1 Surrounding color distribution, test illuminant and test chromaticity

We employed *natural*, *reverse* and *flat* distributions for the surrounding stimuli. For test illuminants,

we used magenta and green illuminants. We chose two color filters (Rosco, R44 "Middle Rose"

and R4460 "Calcolor 60 Green") through which the 6500K illuminant was passed to obtain the

spectra shown in Figure 11 (a). The chromaticities of these illuminants largely deviate from black-

body locus as shown in panel (b). Out of the 574 spectral reflectances of natural objects collected

by Brown, 251 reflectances were inside the chromaticity gamut of the CRT monitor under both

illuminants. For the surrounding stimuli, we sampled 180 reflectances out of the 251 reflectances

and created each distribution following the manipulation used in Experiments 1 and 2.

Experiments 1 and 2 showed no effect of surrounding color distribution but Experiment 3 also

included this manipulation to confirm that the findings also held under atypical illuminants.

Figure 11: (a) Spectra for magenta and green illuminants used in Experiment 3. (b) Chromaticities of both

illuminants. The black-body locus and the chromaticity of 6500K illuminant are shown for comparison

purposes. [one-column figure]

Panel (a) in Figure 12 shows surrounding distributions for all 6 test conditions (3 distributions $\times$ 2

test illuminants). The intensities of the test illuminants were chosen so that the average luminance

across the 180 colors matched 2.5 cd/m^2.

In this experiment, the test chromaticities were chosen so that they varied along two directions:

(i) the black-body locus (shown as circles) and (ii) an axis approximately orthogonal to the black-

body locus (shown as triangles) depicted in panel (b). First, 8 reflectances were selected from the

180 and were used under both illuminant conditions. Then, we sampled 5 different reflectances

separately for each illuminant condition from the reflectances that can be presented only under

either magenta or green illuminant. Thus, these reflectance samples are not shared between

illuminant conditions. This choice was made to choose test chromaticities on the locus orthogonal

to black-body locus as widely as possible. In panel (b), the 5 data points surrounded by a red

edge represent the 5 reflectances that were not shared between illuminant conditions. There were

7 chromaticities for each axis, but one chromaticity was used for both axes (plotted as a black

square). The chromaticities of natural objects tend to spread along the black-body locus, and the

purpose of this design was to test whether luminosity thresholds measured at atypical

chromaticities would deviate from the prediction of the optimal color model or the real object model.

Figure 12: (a) 6 color distributions for surrounding stimuli (2 test illuminants × 3 distributions). Inserted

image shows an example stimulus configuration. The vertical solid black line indicates the L/(L+M) value of

test illuminant. Black cross symbols indicate mean cone response values across 180 surrounding stimuli.

(b) 13 test chromaticities at which the luminosity threshold was measured. Symbols with a red edge

indicates reflectances that were not shared between magenta and green illuminants. [one-column figure]

5.2 Procedure

One block consisted of 13 consecutive settings and thresholds were measured for all test

chromaticities in random order. Each session comprised 6 blocks to test all distribution × illuminant

conditions. The order of condition was randomized. All observers completed 10 sessions in total.

Observers conducted 5 sessions per day and thus the experiment were completed in two days.

5.3 Results

Figure 13 shows the results. The left 6 panels depict luminosity thresholds measured at

chromaticities along the black-body locus (black circles and square in panel (b), Figure 12) while

the right 6 panels indicate thresholds at chromaticities along the orthogonal locus (black triangles

and square in panel (b), Figure 12).

We first look at the left two columns. For the magenta illuminant condition, observers' settings

again show a mountain-like shape. Also, one can see that settings are not dependent on the

surrounding color distribution. However, in this condition the optimal color model and the real

object model show a relatively flat locus. For the green illuminant, observer settings appear flat.

However luminosity thresholds for subject KS show a fairly different trend from the other

observers, and the locus is not well predicted by the optimal color locus nor the real object locus,

which was not observed in Experiments 1 and 2.

Figure 13: Observer settings in Experiment 3. The left two columns plot luminosity thresholds measured at

test chromaticities on the black-body locus (circle and square symbols in panel (b), Figure 12). The right

two columns show results for test chromaticities on the locus orthogonal to black-body locus (triangle and

square symbols in panel (b), Figure 12). Colored square symbols indicate averaged settings across 10

repetitions for each observer. The error bar plots ±1 S.E across 10 repetitions. The black circle symbols

plot average observer settings (n = 3). The magenta circle symbols denote the optimal color locus and the

blue circles show the real object locus. The vertical solid line represents the chromaticity of the test

illuminant. The black cross symbol indicates the mean LMS value across surrounding stimuli. [two-column

figure]

When the test chromaticities are on the axis orthogonal to the black-body locus (right two

columns), for the magenta condition all observers' settings might appear to resemble the optimal

color locus. However, for the green illuminant condition, KS again shows a different trend from

the other observers and observers do not all agree with either model prediction.

Figure 14 allows us to compare the correlation coefficient across models and conditions. For

black-body reflectances shown under the magenta illuminant (the leftmost column), the optimal

color model overall showed good correlations for the *natural* condition, while the real object model

showed good correlations for the *reverse* and *flat* conditions. For the *natural* condition, one

observer (KS, not naïve) had the highest correlation with the surrounding color model, which was

not observed in Experiments 1 and 2 in which illuminants on the black-body locus were used as

test illuminants. For black-body reflectances shown under the green illuminant (the second

leftmost column), in most cells correlation coefficients appeared considerably low. Although the

optimal color model consistently had the highest correlation for all distribution conditions (average

coefficient across 9 cells is 0.578), the correlation coefficient is not so high if we consider that the

correlation for the optimal color model was 0.901 in Experiment 1 (averaged across 5 distributions

× 4 observers). Also, in Experiment 2, correlations were 0.746 for the optimal color model of the

ground-truth illuminant and 0.837 for the estimated illuminant (average across 9 conditions × 4

observers in both cases).

For the reflectances on the axis orthogonal to black-body locus under the magenta illuminant (the

second-from-the-right column), the trend seemed to be close to that of the test chromaticities on

the black-body locus (leftmost column), but the correlation coefficients overall seemed to be lower.

For the green-*natural* condition, the surrounding color model shows a high correlation with

observers KK and KS. It is notable that the optimal color model shows nearly zero or even

negative correlations. For the *reverse* condition, the real object model showed the best correlation,

but their values were not high (0.577, average across 3 observers). For the *flat* condition, we did

not find a consistently good model. It may be worth noting that for the green illuminant condition,

correlation coefficients for test chromaticities sampled from the locus orthogonal to black-body

locus are overall lower than those for test chromaticities sampled from black-body locus.

672
Figure 14: The matrices of Pearson's correlation coefficient calculated between observer settings and model

predictions over 7 test chromaticities in Experiment 3. The cyan star symbols indicate the highest correlation

coefficient across the 3 models. The cyan arrows at the bottom of each subpanel indicate the model that

received the highest number of cyan stars. [1.5-column figure]

In summary these results suggested that although the optimal color and real object models can

account for observer settings to some extent, overall coefficient values were substantially lower

than those observed in Experiments 1 and 2. Also the surrounding color model showed good

correlations in some cases. These results might imply that the visual system does not have a rigid

internal reference about upper-limit luminance under atypical illuminants and sometimes relies on

external cues such as the color of the surrounding stimuli. Also, for green illuminant condition, we

found a trend that the predictions of optimal color model were particularly worse when test

chromaticities were sampled from the axis orthogonal to the black body locus.

Finally, we summarize results from the three experiments to test whether correlation coefficients

of the optimal color model are higher for typical illuminants (Experiments 1 and 2) than atypical

illuminants (Experiment 3). For each observer, we averaged the correlation coefficient of the

optimal color model across all condition in Experiments 1 and 2 (14 conditions), which served as

a summary statistic for typical illuminants. For the 20000K condition in Experiment 2, we used the

correlation coefficient value of the optimal color model under the estimated illuminant as it

predicted observer settings substantially better than the model under the ground-truth illuminant.

We also calculated averaged correlation coefficients across all conditions in Experiment 3 (8

conditions) per observer. The averaged correlation coefficients across all observers were 0.879

± 0.0114 (average ±1 S.D.) for typical illuminants and 0.525 ± 0.155 for atypical illuminants.

Welch's t-test (one-tailed, no assumption about equal variance) showed that the optimal color

model has a significantly higher correlation for typical illuminants than atypical illuminants (t(2.01)

= 3.94, p = 0.0290). Also we performed the same analysis using correlation coefficients for the

real object model which showed the same trend (t(2.84) = 2.93, p = 0.0326).

These results are consistent with the idea that human observers empirically learn the upper-limit

luminance through observing colors in natural environments and use the criterion to judge

whether a given surface is self-luminous or not. Since magenta and green illuminants are

uncommon in natural environments, the visual system does not know the upper limit of surface

colors under those illuminants. Moreover, in the Appendix, we show that a simplistic model

predicts the luminosity threshold in Experiment 3 as well as the optimal color model. A potential

interpretation would be that when the scene illuminant has an atypical color, the visual system

makes a self-luminous judgement based on simple statistics.

5. General Discussion

This study investigated potential determinants of luminosity thresholds. Our three experiments

showed that the loci of luminosity thresholds have a mountain-like shape that peaks around the

illuminant color and decreases as stimulus purity increases, showing a striking similarity to optimal

color and real object loci. A simple alternative strategy which bases judgements on the

surrounding color distribution did not explain observers' settings well. Rather observers seem to

hold an internal representation about the luminance at which a surface should reach self-

luminosity. Moreover, such similarity between luminosity threshold and optimal color or real object

loci was higher when surfaces were placed under illuminants along the blue-yellow direction than

magenta and green illuminants that are atypical in natural environments. These support an idea

that the visual system empirically internalizes the gamut of surface colors through an observation

of colors in daily life. Going back to the original question of whether the visual system relies on a

heuristic or internal reference for luminosity judgements, the present study generally supports the

internal reference hypothesis.

We also note that some properties of the test field we did not consider in the present study are

known to affect luminosity judgement. For instance, it was reported that surfaces with smaller

areas appeared to emit light at lower luminance levels (Bonato & Gilchrist, 1999). It has also been

reported that surround stimuli are more likely to affect luminosity threshold if the surround stimuli

are presented at the same depth as the test field (e.g., Yamauchi & Uchikawa, 2005). Thus, in

future studies, it would be desired to expand our model to modify the prediction of luminosity

thresholds based on factors such as stimulus size and relative depth in a way that agrees with

human luminosity judgements.

One consistent trend across the three experiments was that observer settings were above the

physical limit (i.e., prediction of the optimal color model) in most cases. We suspect that there are

at least two reasons for this. First, observers' criterion to judge the luminosity threshold in this

study was the midpoint between the upper-limit of the surface color mode where a test field purely

appears as an illuminated surface and the lower-limit of the aperture color mode where a test field

purely appears as a light source. If we instead used a different criterion, for example to set the

luminance so that the test field appears simply as the upper-limit of the surface color mode, we

would have seen a smaller discrepancy. Second, it is possible that observers overestimated the

intensity of the test illuminant. Just as humans cannot directly access the chromaticity of test

illuminant, the intensity is also not directly known to observers. In fact, in other color constancy

studies (Morimoto, Fukuda & Uchikawa 2016; Morimoto et al., 2021), we have repeatedly found

that observers tend to overestimate illuminant intensities and the degree of overestimation

substantially varied across individuals. When the illuminant intensity is overestimated, the

observers' internal upper-limit luminance should accordingly increase, which could account for an

observed discrepancy between model predictions and observer settings. Moreover, we also note

that the limit drawn by optimal-color or real-object model corresponds to the upper-boundary as

a surface color, but the models are not designed to exactly predict how the test field should appear

when the luminance of the test field exceeds the predicted value by the models. Thus, it is

inherently difficult to directly compare the model prediction and observers' settings, and we think

that they rather need to be compared relatively (for example, using correlation coefficient). We

also found that individual differences in this study were mainly found in gain rather the shape

of the observer settings (though for some conditions in Experiment 3 shape differences were also

evident). This individual variation in gain could also be due to the variability in estimating illuminant

intensity.

Color constancy is often described as a visual ability to identify the same surface under different

illuminants. A surface reflects a light, and that reflected light enters our eyes. Because the

reflected light is a product of surface and illuminant components, color constancy is often framed

as a process in which our visual system estimates the influence of the illuminant. The "brightest

is white" heuristic, which assumes that a surface with the highest luminance provides the closest

information about the illuminant color, has been known as an influential approach in estimating

illuminant color (Land, 1977). However, self-luminous objects do not carry information about the

scene illuminant, which might cause a misestimation of the illuminant if included in a scene. In

general, when we receive an intense light from a surface, there are two ways to interpret this.

One is that the surface is placed under an intense illuminant and the other is that the surface is

self-luminous. This example highlights the need for luminous percepts to be incorporated into the

process of color constancy. In fact, Fukuda & Uchikawa (2014) showed that a surface appearing

in aperture-color mode does not have a strong influence on observers' estimates of the illuminant.

We chose a set of colored circles as experimental stimuli to directly test our hypothesis while

excluding any other cues. However, it is reported that changing a material property could affect

the mode of color appearance (Kuriki, 2015). Also, our experimental stimuli were simulated to be

uniformly illuminated by a single illuminant, but in natural environments the spectra hitting an

object surface changes from one direction to another (Morimoto et al., 2019). The presence of

multiple illuminants means that we need to consider multiple optimal color distributions, and thus

the loci of luminosity thresholds measured under such an environment might also change. Despite

a growing amount of research on material perception (Fleming, 2013), luminosity perception is

little studied in the field. While our choice of stimuli was necessary for experimental control, it will

be interesting whether our finding can be applied to a wider range of stimuli that have complex

material properties and are illuminated in non-uniform ways.

One closely related phenomenon to self-luminous perception would be brightness perception of

colored objects. The Helmholtz-Kohlrausch effect is that stimuli with high purity appear to have

high brightness even if luminance is kept the same. The effect was reported under a variety of

viewing conditions (Nayatani et al., 1991; Donoforio, 2011). However, it is unclear why a color

with high purity appears brighter. Curiously, as observed in the present study, the same trend

holds for luminosity thresholds: a surface with high purity reaches the limit of surface color mode

at a lower luminance level. Thus, if we take a strategy to determine the brightness of colored

stimuli in comparison to the theoretical upper-limit luminance at the chromaticity we could account

for the Helmholtz-Kohlrausch effect. Uchikawa et al. (2001) directly focused on this relationship

and argued that saturated colors appear brighter because the visual system knows that it has a

lower limit and brightness might be determined in proportion to the theoretical upper-limit

luminance.

Identifying the range of natural colors has been a major focus especially in the field of color

science (e.g. Pointer, 1980). While the limit of chromaticity has been well characterized, less is

known regarding the luminance limit. In this study, we used the SOCS reflectance dataset as a

reference to draw an upper-luminance boundary for real objects. The database covers a wide

range of color space as it includes man-made materials such as ink which can have narrow-band

reflectances. We do not intend to claim that the SOCS dataset in any sense represents all

plausible natural reflectance spectra. Yet, our separate analysis based on 16 hyperspectral

images (Nascimento, 2002; Foster, 2006) showed that colors in those images were mostly

covered in the gamut of the SOCS dataset. Also, to our knowledge we have not encountered

another dataset that has a larger color gamut than the SOCS dataset. We also found that if we

restrict samples to natural objects, the color gamut largely shrinks (see Figure 2 (b) in Morimoto

et al., 2016), and upper-limit luminance estimated only from natural samples would not predict

obtained luminosity thresholds in this study. Also, in this study, we used a smoothed upper-limit

luminance. We confirmed that if we instead used raw unsmoothed data, the correlation coefficient

was lower in almost all tested conditions. These results show that a precise evaluation of the

abundance of reflectance samples in real world seems to play a key role in understanding the

luminosity percept. When more reflectance datasets become available in the future, the gamut of

real objects may need to be re-evaluated.

In summary, our results showed a mysterious similarity between luminosity thresholds and optimal

colors. Yet it is difficult to make a conclusive statement as to whether the optimal color model is

better in accounting for luminosity thresholds than the real object model. This is partially because

the optimal color locus well resembles the locus of real objects, leading to high correlation

between predictions from two models. Furthermore, an intrinsically more challenging question

would be how our visual system learns the optimal color locus because optimal colors do not exist

in the real world. Considering this point, one plausible theory would be that our visual system

learns the plausible range of surface colors by seeing colors in daily life and empirically

internalizes the gamut of surface colors. Then, a given surface appears self-luminous when its

luminance exceeds the upper-limit luminance empirically internalized in the visual system. This

study presents a potential link between our perceptual judgment and statistical properties of the

real world.

Acknowledgement

All authors thank Laysa Hedjar for her careful edit of the language throughout the manuscript.

This work was supported by JSPS KAKENHI Grant Number JP19K22881, JP17K04503 and

26780413. TM is supported by a Sir Henry Wellcome Postdoctoral Fellowship awarded from the

Wellcome Trust (218657/Z/19/Z) and the Junior Research Fellowship from Pembroke College,

University of Oxford. This research was funded in whole, or in part, by the Wellcome Trust

(218657/Z/19/Z). For the purpose of open access, the author has applied a CC BY public

copyright licence to any Author Accepted Manuscript version arising from this submission.

Data access

The   raw   experimental   data   are   available   in   a   data   repository   at

https://doi.org/10.5281/zenodo.5590120. Codes to reproduce figures in the main manuscript are

available at https://github.com/takuma929.

Reference

Bonato F., & Gilchrist, A. L. (1994). The Perception of Luminosity on Different Backgrounds and

in Different Illuminations. Perception. 23, 9, 991-1006.

Bonato, F., & Gilchrist, A. L. (1999). Perceived area and the luminosity threshold. Perception &

Psychophysics, 61, 786–797.

Brainard. D. H. (1998). Color constancy in the nearly natural image 2 Achromatic loci. Journal of

the Optical Society of America A, 15, 2, 307–325.

Brown, R. Background and illuminants: the yin and yang of colour constancy. in Colour

Perception: Mind and the Physical World, R. Mausfeld and D. Heyer, eds. (Oxford University,

2003), 247–272.

Buchsbaum, G. (1980). A spatial processor model for object colour perception. Journal of the

Franklin Institute, 310, 1, 1–26.

Donofrio, R.L. (2011). Review Paper: The Helmholtz-Kohlrausch effect. Journal of the Society for

Information Display, 19, 658-664.

Evans, R. M. (1959). Fluorescence and Gray Content of Surface Colors. J. Opt. Soc. Am. 49,

1049-1059

Evans, R. M. & Swenholt, B. K. (1967). Chromatic strength of colors: dominant wavelength and

purity. J. Opt. Soc. Am. 58, 580-584. J. Opt. Soc. Am. 58, 580-584.

Evans, R. M. & Swenholt, B. K. (1968). Chromatic Strengths of Colors, Part II. The Munsell

System. J. Opt. Soc. Am. 58, 580-584.

Evans, R. M. & Swenholt, B. K. (1969). Chromatic Strength of Colors, III. Chromatic Surrounds

and Discussion. J. Opt. Soc. Am. 59, 628-634.

Fleming, R. W. (2013). Material Perception. Annual Review of Vision Science, 3, 365–388.

Fukuda, K. & Uchikawa, K. (2014). Color constancy in a scene with bright colors that do not have

a fully natural surface appearance. Journal of the Optical Society of America A, 31, 4, A239–

A246.

Gilchrist, A. & Bonato, F. (1995). Anchoring of lightness values in center-surround displays.

Journal of Experimental Psychology: Human Perception and Performance, 21, 6, 1427-1440.

Gilchrist, A., Kossyfidis, C., Bonato, F., Agostini, T., Cataliotti, J., Li, X., Spehar, B., Annan, V. &

Economou, E. (1999). An anchoring theory of lightness perception. Psychological Review, 106,

795–834.

Judd, D. B., MacAdam, D. L. Wyszecki, G., Budde, H. W., Condit, H. R., Henderson S. T., and

Simonds J. L. (1964) "Spectral distribution of typical daylight as a function of correlated color

temperature," J. Opt. Soc. Am. 54(8), 1031–1040

Kuriki, I. (2015) Effect of material perception on mode of color appearance. Journal of Vision,

15(8), 4.

Krinov, E. (1947) Spectral reflectance properties of natural formations. Tech. rep., National

Research Council of Canada. Technical Translation: TT-439.

Land, E. H. (1977). The Retinex Theory of Color Vision, Scientific American, 237, 6, 108 -128.

Long, F., & Purves, D. (2003) Natural scene statistics as the universal basis of color context effects.

Proceedings of the National Academy of Sciences, 100, 25, 15190-15193.

Lotto, R. B. & Purves, D. (2000) An empirical explanation of color contrast. Proceedings of the

National Academy of Sciences, 97, 23, 12834-12839.

MacAdam, D. (1935a). The theory of the maximum visual efficiency of colored materials. Journal

of the Optical Society of America A, 25, 249–252.

MacAdam, D. (1935b). Maximum visual efficiency of colored materials. Journal of the Optical

Society of America A, 25, 361–367.

MacLeod, D. I. A., & Boynton, R. M. (1979) Chromaticity diagram showing cone excitation by

stimuli of equal luminance. Journal of the Optical Society of America A, 69, 1183–1186.

Maloney, L. T. & Wandell, B. A. (1986) Color constancy: a method for recovering surface spectral

reflectance Journal of the Optical Society of America A, 3, 1, 29–33.

Morimoto, T., Fukuda, K., & Uchikawa, K. (2016) "Effects of surrounding stimulus properties on

color constancy based on luminance balance", Journal of the Optical Society of America A, 33,

3, A214–A227.

Morimoto, T., Kishigami, S., Linhares, J. M. M., Nascimento, S. M. C., & Smithson, H. E. (2019).

Hyperspectral environmental illumination maps: characterizing directional spectral variation in

natural environments. Optics Express, 27, 32277-32293.

Morimoto, T., Kusuyama T., Fukuda, K. & Uchikawa, K. (2021). Human color constancy based on

the geometry of color distributions. Journal of Vision, 27, 21(3):7. doi:

https://doi.org/10.1167/jov.21.3.7.

Nascimento, S.M.C., Ferreira, F.P., & Foster, D.H. (2002). Statistics of spatial cone-excitation

ratios in natural scenes. Journal of the Optical Society of America A, 19, 1484-1490.

Foster, D.H., Amano, K., Nascimento, S.M.C., & Foster, M.J. (2006). Frequency of metamerism

in natural scenes. Journal of the Optical Society of America A, 23, 2359-2372.

Nayatani Y., Umemura, Y., Sobagaki, H., Takahama, K. & Hashimoto, K. (1991). Lightness

perception of chromatic object colors. Color research and application, 16, 1.

Pearce B., Crichton S., Mackiewicz M., Finlayson G. D., & Hurlbert A. (2014). Chromatic

Illumination Discrimination Ability Reveals that Human Colour Constancy Is Optimised for Blue

Daylight Illuminations. PLoS ONE 9(2): e87989.

Pointer, M. R. (1980) The gamut of real surface colours. Col. Res. Appl. 5, 145–155.

Speigle, J. M. & Brainard, D. H. (1996). Luminosity thresholds: effects of test chromaticity and

ambient illumination. Journal of the Optical Society of America A, 13, 436-451

Stockman, A. and Sharpe, L. T. (2000) The spectral sensitivities of the middle- and long-

wavelength-sensitive cones derived from measurements in observers of known genotype.

Vision Research, 40, 1711–1737.

Uchikawa, K., Koida, K., Meguro, T., Yamauchi Y., & Kuriki, I. (2001). Brightness, not luminance,

determines transition from the surface-color to the aperture-color mode for colored lights.

Journal of the Optical Society of America A, 18, 737–746.

Uchikawa, H., Uchikawa, H. & Boynton, R. M. (1989). Influence of achromatic surrounds on

categorical perception of surface colors. Journal of the Optical Society of America A, 29, 7, 881–

890.

Uchikawa, K., Fukuda, K., Kitazawa, Y., & MacLeod, D. I. A. (2012). Estimating illuminant color

based on luminance balance of surfaces. Journal of the Optical Society of America A, 29, 2,

A133–A143.

Ullman, S. (1976) On visual detection of light sources. Biological Cybernetics, 21, 205–211.

Winkler A. D., Spillmann L., Werner J. S., & Webster M. A. (2015) Asymmetries in blue-yellow

color perception and in the color of 'the dress'. Current Biology, 29, 25(13), R547-R548.

Yamauchi, Y. & Uchikawa, K. (2005). Depth information affects judgment of the surface-color

mode appearance. Journal of Vision, 5, 515-524.

Appendix

Individual observer settings in Experiment 2

In the main text, we presented mean observer settings for Experiment 2. Figure A1 shows the individual observer settings. There is some individual variation, but overall the trend was similar across individuals.

Figure A1: Individual observer settings in Experiment 2. Colored square symbols indicate the averaged setting across 10 repetitions for each observer. The error bar indicates ±1 S.E across 10 repetitions. The magenta circles denote the optimal color locus and the blue circles show the real object locus. The red, black and blue vertical solid lines show the chromaticities of the 3000K, 6500K and 20000K test illuminants, respectively. The black cross symbol indicates mean LMS value across surrounding stimuli. Note that the horizontal range differs across panels. [two-column figure]

Implementation of other past models

Here, we compare two simplistic models that predict luminosity thresholds based on post-receptoral signals or cone signals of the test field. Note that these two models were also implemented as candidate models in Speigle & Brainard (1996).

The first model is a post-receptoral model which assumes that the visual system monitors the

weighted sum of the three types of cone signals of the test field. The test field appears self-

luminous when $w_L L_T + w_M M_T + w_S S_T >= 1$, where L_T, M_T and S_T denote cone signals of the test

field and w_L, w_M and w_S denote weightings for each class of cone signal ($w_L > 0$, $w_M > 0$, and w_S

> 0). Then, we optimized weightings w_L, w_M and w_S which produced the minimum root mean

square error between model prediction and mean observers' setting.

The second model is a generalized Evans's model (Evans, 1959) which assumes that a test field

appears self-luminous when any of L_T, M_T or S_T cone signals exceed a certain criterion level.

Thus, the model predicts that a test field is self-luminous when $L_T >= c_L$, $M_T >= c_M$ or $S_T >= c_S$.

The goal of the optimization here was to find criterion c_L, c_M and c_S that minimized the root mean

square error between model prediction and mean observers' setting.

For both models, the optimization procedure was performed separately for each experimental

condition. In other words, for example in Experiment 2, we performed optimization procedures 9

times in total (3 illuminants×3 distributions).

Figures A2, A3 and A4 depict the prediction of the post-receptoral model and the generalized

Evans's model as well as the optimal color model in Experiments 1, 2 and 3. To obtain the

prediction of the optimal color model, we used the ground-truth illuminant for Experiments 1 and

3, but for Experiment 2, we used the estimated illuminant so that the peak is matched between

the prediction of optimal color model and the mean observer settings (for more details, see the

Results section in Experiment 2 in the main text).

For Experiment 1, we see that the post-receptoral model predicts a linear luminosity threshold

locus over L/(L+M) which did not lead to a high correlation coefficient (shown at the right upper

corner in each panel). The generalized Evans's model came closer to observer settings, but in all

conditions the correlation coefficient was lower than that of the optimal color model. Welch's t-test

(one-tailed, no assumption about equal variance) on averaged correlation coefficients across the

5 conditions showed that the optimal color model has a significantly higher correlation than the

Evans' model ($t(4.64) = 4.57$, $p = 0.0072$).

A quite similar trend is shown in Experiment 2 though in one condition (*flat* and 20000K), the

Evans's model exceeded the optimal color model's correlation coefficient. However, again

Welch's t-test on averaged correlation coefficients across the 9 conditions showed a significantly

higher correlation coefficient for the optimal color model than the Evans's model ($t(8.95) = 3.72$,

p = 0.0048).

In contrast, for Experiment 3, we see that the Evans's model showed higher correlations than the

optimal color model especially in green illuminant conditions. Welch's t-test on averaged

correlation coefficients across the 12 conditions showed that there is no significant difference

between the Evans's model and the optimal color model ($t(12.2)$ = 1.80, p = 0.0962).

In summary, it is evident that for Experiments 1 and 2 the optimal color model predicts human

observer settings better than the two alternative models considered here. In Experiment 3, there

was no significant difference in correlation coefficients between the Evans's model and the optimal

color model. One interpretation would be that when the scene illuminant is atypical, human

observers rely on the simple statistics such as cone signals because the visual system does not

know the optimal color locus under the atypical illuminant.

1002

1003

1004

Figure A2: Predictions from the post-receptoral model, the generalized Evans's model and the optimal color

model in Experiment 1. Each model prediction was scaled to give the minimum root mean square error

between model prediction and mean observer setting to compare their shapes more easily. The correlation

coefficients between model prediction and mean observer settings are shown at the top right corner in each

panel (in the order of post-receptoral model, generalized Evans's model and optimal color model from top to

bottom). We used a ground-truth illuminant (i.e., 6500K) to obtain the prediction from the optimal color model.

[1.5-column figure]

1012

1013

1014

Figure A3: Predictions from the post-receptoral model, the generalized Evans's model, and the optimal color

model in Experiment 2. Each model prediction was scaled to give the minimum root mean square error

between model prediction and mean observer setting. The correlation coefficients between model prediction

and mean observer setting are shown at the right top corner in each panel. For the optimal color model, we

used an estimated illuminant whose peak matched that of the observer settings (see Results section in

Experiment 2 in the main text for more details). [1.5-column figure]

1021

1022

1023

1024 Figure A4: Predictions from the post-receptoral model, the generalized Evans's model, and the optimal color

1025 model in Experiment 3. Each model prediction was scaled to give the minimum root mean square error

1026 between model prediction and mean observer setting. The correlation coefficient between model prediction

1027 and mean observer setting is shown at the top right corner in each panel. We used a ground-truth illuminant

1028 (i.e. magenta or green) for the optimal color model. [two-column figure]

1029

1030

1031

1032

1033

49