Natural history of two carpenter ants, *Camponotus renggeri* and *C. rufipes* (Hymenoptera:

Formicidae), in an Atlantic Forest reserve

Names of authors:

Miguel Piovesana PEREIRA-ROMEIRO[1][$], Gabriel Tofanelo VANIN[1][$], Marianne AZEVEDO-

SILVA[2], Gustavo Maruyama MORI[1]*

Keywords: Neotropics, Atlantic forest, Hymenoptera, foraging area, nesting architecture, daily

activity, polydomy

Addresses of authors

1 – Instituto de Biociências, Universidade Estadual Paulista (UNESP), 11380–972 São Vicente

SP, Brazil

2 – Programa de Pós-Graduação em Ecologia, Universidade Estadual de Campinas, 13083–862

Campinas SP, Brazil

[$] Contributed equally

* gustavo.mori@unesp.br

Abstract

Widespread species face a wide variety of environmental challenges and their morphology,

behavior, and natural history may change across their range. However, not rarely, natural history

research is restricted to one or few locations. That is the case for *Camponotus renggeri* and *C.*

rufipes. Both species occur across South America in different ecosystems, but most research on

these species is restricted to the Brazilian savanna, known as Cerrado. Here, we describe the

home range, nesting habits, and activity schedule of *C. renggeri* and *C. rufipes* in an Atlantic

Forest reserve in SE Brazil. *C. renggeri* foraged exclusively during nighttime and *C. rufipes*

remained active throughout the nycthmeron, but with little intensity during daylight hours. Most

nests of both species were composed of dry straw, and home ranges varied from nearly 0.91 m²

(*C. renggeri*) to 1.79 m² (*C. rufipes*). Foraging areas overlapped, especially in *C. renggeri*. Our

field study reinforces the importance of natural history and what it adds to our knowledge on the

ecology and behavior of *C. renggeri* and *C. rufipes* in Atlantic Forest.

The closely related carpenter ants *Camponotus renggeri* (Emery, 1894) and *C. rufipes* (Fabricius, 1775) have been extensively studied in terms of physiology (TAKAHASHI-DEL-BIANCO & al. 1998, GALIZIA & al. 1999), natural history (RONQUE & al. 2016, RONQUE & al. 2018) and genetics (AZEVEDO-SILVA & al. 2015, RONQUE & al. 2016, DE AGUIAR & al. 2017, AZEVEDO-SILVA & al. 2020). These species are morphologically similar, being distinguished mainly by the brightness of their tegument and by the color of their legs. Both species are commonly found on vegetation, where they feed on liquid resources from extrafloral nectaries and trophobionts (OLIVEIRA & FREITAS 2004, RONQUE & al. 2018). Despite similarities in distribution and interactions with plants, these species differ with respect to nest architecture, home ranges, and breeding systems. *C. renggeri* occupies nests built in dead tree trunks or underground, whereas *C. rufipes* may also construct nests out of dry straw (RONQUE & al. 2016, RONQUE & al. 2018, AZEVEDO-SILVA & al. 2020). Both species occur across South America in different ecosystems (JANICKI & al. 2016), but most research on these species is restricted to the Brazilian savanna, known as Cerrado.

The main goal of this study was to investigate the natural history of *C. renggeri* and *C. rufipes* in the Atlantic Forest. Specifically, we aimed to describe their home range, nesting habits, and activity schedule. For both species, our findings unveiled consistent differences in home ranges and nest architecture when compared to previous findings, although we observed similarities in natural history between populations from the Cerrado and Atlantic Forest. Our study is the first step towards a broader understanding of how the ecology and behavior of tropical ant species may vary across different ecosystems.

Fieldwork was carried out in an Atlantic Forest fragment surrounded by the sea and urban environment, at Xixová-Japuí State Park, São Vicente, São Paulo in Southeast Brazil (23°59'33.3" S 46°23'21.7" W). The Park is dominated by secondary ombrophilous forests, and some areas are at initial successional stage, with abundant herbaceous plants and low canopy cover (Figure 1A). The region has a typically hot and very humid climate, with mean annual temperature of 23.6 °C and mean annual precipitation of 2500 mm, being classified as tropical rainforest climate (*Af*) (PEEL & al. 2007). Observations were performed from February to April 2019 and from May to July 2019. The study area was divided into 10 plots (400 m² each), where

ants were searched for 20 minutes in each plot. We used sardine baits to find *C. renggeri* and *C.*

rufipes nests. In each plot where we were able to find these species, from three to five nests of

each were tagged and characterized for nest architecture, home range, and activity schedule.

Daily activity schedule was described by the total number of workers entering and exiting nests

during sessions of 30 min, every 2 h, over a period of 24 h. For home range estimation, 4 *C.*

renggeri and 3 *C. rufipes* nests were observed for a total maximum of 16 h, encompassing their

peak activity period. Workers were followed from nest entrance to the furthest point before

returning to nest. These points were tagged and had their distance to the nest measured and

relative geographical location determined with a compass. Home range was estimated using

minimum convex polygons generated with the package adehabitatHR (CALENGE 2006) in R (R

CORE TEAM. 2020). Due to low nest persistence, all observations of *C. renggeri* were carried out

in the dry season, whereas observations of *C. rufipes* were performed during rainy season.

During fieldwork, additional natural history records were made for both species using video,

photographs, and notes (total of 32 h of observations for each species).

Nests were found in only 3 plots, and none of them had nests of both species. A total of 15 *C.*

renggeri (12 nests in the same plot, one in a second plot and two nests opportunistically found

outside of the plots) and 9 *C. rufipes* nests (all in the same plot) were identified and tagged. Of

the total, 8 *C. renggeri* and 6 *C. rufipes* nests were unoccupied by the end of this research (July

2019), indicating low nest persistence (Figure 1B, C). Both species built their nests using mainly

dry straw (*C. renggeri*: N = 15 nests; *C. rufipes*: N = 8 nests), but both could use additional

materials, such as freshly fallen and dry leaves (Table 1). Many of these nests were suspended

above ground, sustained by twigs (Figure 2). As for the activity schedule, the period of peak

activity for both species was nocturnal, increasing after sunset (18 h) and decreasing before

sunrise (05 h) (Figure 3). *C. rufipes* had a mean home range value of 1.79 m², and *C. renggeri*

had a mean value of 0.91 m² (Table 2). Major workers of *C. renggeri* and *C. rufipes* were found

engaging in aggressive behavior towards each other when crossing paths, resulting in the

decapitation of *C. rufipes* (see Supporting information S1). These observations suggest that

competition may occur between the two species.

Our findings in Atlantic Forest differ from those reported for other ecosystems, indicating
that environment may play an important role in shaping *C. renggeri* and *C. rufipes* behavior. All
but one *Camponotus rufipes* nests characterized in this study were made of dry straw, similarly
to the observed in the Cerrado and in the Argentinian "Chaco" (WEIDENMÜLLER & al. 2009,
RONQUE & al. 2018). Conversely, *C. renggeri* also presented this same nest architecture,
differing from the nests in dead trunks reported for this species in Cerrado (RONQUE & al. 2018),
whereas no nests of such type were found in the Atlantic Forest. It is known that vegetation may
play a role in regulating availability of resources for nesting in ants (RONQUE & al. 2018). The
Atlantic Forest fragment where this research was conducted is dominated by herbaceous plants,
thus, dead trunks is not as abundant as leaf litter. This may explain the prevalence of nests made
of dry straw over other nest architectures (SILVESTRE & al. 2003). The difference found for *C.*
renggeri nests between different ecosystems indicates that this species is able to modify its
nesting behavior in response to resource availability in different environments. Additionally, our
findings expand the repertoire of nesting strategies reported for the *Camponotus* genus (PFEIFFER
& LINSENMAIR 2000, SANTOS & al. 2005, TSCHINKEL 2005, YAMAMOTO & DEL-CLARO 2008,
SANTOS & DEL-CLARO 2009).

Camponotus rufipes and *C. renggeri* presented smaller home ranges compared to other carpenter
ants (PFEIFFER & LINSENMAIR 2000, YAMAMOTO & DEL-CLARO 2008, RONQUE & al. 2018).
However, despite the impracticability of a direct comparison due to slight methodological
differences, we observed much smaller values (Table 2) than previously findings reported for the
same species (RONQUE & al. 2018). In the Cerrado, *C. rufipes* foraging area doubled during rainy
season, whereas *C. renggeri* home range was not affected by different seasons (RONQUE & al.
2018). Given our observations occurred in different seasons for *C. rufipes* and *C. renggeri*, it is
unclear whether the home range estimated for *C. rufipes* is indeed larger than *C. renggeri* in the
Atlantic Forest or if it is an effect of seasonality. However, the differences reported between
biomes may be related to many factors, such as temperature, humidity, colony size, competition,
and food availability (HÖLLDOBLER 1987, BREED & al. 1990, GORDON 1995, MCGLYNN & al.
2003, RONQUE & al. 2018).

Beyond seasonal changes in foraging, daily cycles may also influence a colony´s activity. We

showed that *C. renggeri* foraged exclusively during night, whereas *C. rufipes* remained active throughout the nycthmeron, with little activity during the day. Similar patterns were reported for this species in Cerrado, with *C. renggeri* activity being negatively affected by temperature and positively by humidity, whereas *C. rufipes* was not affected by these environmental conditions (RONQUE & al. 2018). Indeed, nocturnal behavior is common among tropical ant species, presumably because it allows them to avoid high temperatures and low humidity during the day (PFEIFFER & LINSENMAIR 2000, RONQUE & al. 2018). Additionally, temporal segregation can be a result of competition and resource sharing, particularly in exudate-feeding *Camponotus* species (DEL-CLARO & OLIVEIRA 1999, OLIVEIRA & al. 1999, RONQUE & al. 2018), even though most species can combine exudate with animal prey in their diets (OLIVEIRA & FREITAS 2004, RONQUE & al. 2018). These mechanisms could explain not only the difference in daily activity, but also the observed spatial partition between *C. renggeri* and *C. rufipes*. Accordingly, we recorded an aggressive encounter of major workers of *C. renggeri* and *C. rufipes*, which further suggest an antagonistic relationship between these species.

This work focused on the description of natural history traits of *C. renggeri* and *C. rufipes* in the Atlantic Forest, an ecosystem where the ecology of these species is poorly explored. We consistently showed that the behavior of ants may vary across biomes, shedding new light on natural history of *Camponotus* species in tropical ecosystems. We explored which differences in the behavior of species may emerge in different environments and showed how ant behavior may vary across ecosystem. Identifying and understanding such variation is crucial to elaborate more complex and robust questions regarding ant ecology. Finally, we expect that our work encourages further investigation on ant natural history, which would be important to increase our knowledge on tropical systems.

Acknowledgments

We thank Instituto Florestal (COTEC 260108–010.270/2018), Instituto Chico Mendes de Conservação da Biodiversidade (SISBIO 66314-1), and Conselho de Gestão do Patrimônio Genético (A025CBF) for sampling and genetic resources access licenses. Also, we are grateful to Dr. Mariane Ueda Vaz Ronque and Prof. Paulo Sérgio Oliveira for their insightful comments and

suggestions on previous versions of this manuscript. Additional thanks to many colleagues for

field assistance. This research was funded by the Conselho Nacional de Desenvolvimento

Científico e Tecnológico (CNPq – PIBIC/Unesp #48498, #48010, and #54099) and Fundação de

Amparo à Pesquisa do Estado de São Paulo (FAPESP 2017/18291-2; 2019/12646-9; 20/15636-

1).

References

AGUIAR, H.J.A.C. DE, BARROS, L.A.C., ALVES, D.R., MARIANO, C.D.S.F., DELABIE, J.H.C. &
POMPOLO, S.D.G. 2017: Cytogenetic studies on populations of *Camponotus rufipes*
(Fabricius, 1775) and *Camponotus renggeri* Emery, 1894 (Formicidae: Formicinae). –
PLoS ONE 12: 1-18.

ANDERSON, J. 2017: Why Ecology Needs Natural History. – American Scientist 105: 290.

AZEVEDO-SILVA, M., MORI, G.M., CARVALHO, C.S., CÔRTES, M.C., SOUZA, A.P. & OLIVEIRA,
P.S. 2020: Breeding systems and genetic diversity in tropical carpenter ant colonies:
different strategies for similar outcomes in Brazilian Cerrado savanna. – Zoological Journal
of the Linnean Society 190: 1020-1035.

AZEVEDO-SILVA, M., MORI, G.M., SOUZA, A.P. & OLIVEIRA, P.S. 2015: Microsatellites for two
Neotropical dominant ant species, *Camponotus renggeri* and *C. rufipes* (Hymenoptera:
Formicidae). – Conservation Genetics Resources 7: 459-462.

BREED, M.D., ABEL, P., BLEUZE, T.J. & DENTON, S.E. 1990: Thievery, home ranges, and
nestmate recognition in *Ectatomma ruidum*. – Oecologia 84: 117-121.

BURY, B.R. 2006: Natural history, field ecology, conservation biology and wildlife management:
time to connect the dots. – Herpetological Conservation and Biology 1: 56-61.

CALENGE, C. 2006: The package "adehabitat" for the R software: A tool for the analysis of space
and habitat use by animals. – Ecological Modelling 197: 516-519.

DEBOUT, G., SCHATZ, B., ELIAS, M. & MCKEY, D. 2007: Polydomy in ants: What we know, what
we think we know, and what remains to be done. – Biological Journal of the Linnean
Society 90: 319-348.

DEL-CLARO, K. & OLIVEIRA, P.S. 1999: Ant-Homoptera Interactions in a Neotropical Savanna:
The Honeydew- Producing Treehopper, *Guayaquila xiphias* (Membracidae), and Its
Associated Ant Fauna on *Didymopanax vinosum* (Araliaceae). – Biotropica 31: 135.

ELLIS, S., PROCTER, D.S., BUCKHAM-BONNETT, P. & ROBINSON, E.J.H. 2017: Inferring
polydomy: a review of functional, spatial and genetic methods for identifying colony
boundaries. – Insectes Sociaux 64: 19-37.

GALIZIA, C.G., MENZEL, R. & HÖLLDOBLER, B. 1999: Optical imaging of odor-evoked
glomerular activity patterns in the antennal lobes of the ant *Camponotus rufipes*. –
Naturwissenschaften 86: 533-537.

GORDON, D.M. 1995: The development of an ant colony's foraging range. – Animal Behaviour

49: 649-659.

HÖLLDOBLER, B. 1987: Communication and competition in ant communities. In: Evolution and
coadaptation in biotic communities (Proceedings of the Second International Symposium
held in conjunction with the International Prize for Biology, Tokyo, 1986).

HÖLLDOBLER, B. & WILSON, E.O. 1990: The Ants.

HOLWAY, D.A. & CASE, T.J. 2000: Mechanisms of dispersed central-place foraging in
polydomous colonies of the Argentine ant. – Animal Behaviour 59: 433-441.

JANDT, J.M. & GORDON, D.M. 2016: The behavioral ecology of variation in social insects. –
Current Opinion in Insect Science 15: 40-44.

JANICKI, J., NARULA, N., ZIEGLER, M., GUÉNARD, B. & ECONOMO, E.P. 2016: Visualizing and
interacting with large-volume biodiversity data using client-server web-mapping
applications: The design and implementation of antmaps.org. – Ecological Informatics 32:
185-193.

JEANSON, R. & WEIDENMÜLLER, A. 2014: Interindividual variability in social insects - proximate
causes and ultimate consequences. – Biological Reviews 89: 671-687.

JOHNSON, K.G., BROOKS, S.J., FENBERG, P.B., GLOVER, A.G., JAMES, K.E., LISTER, A.M.,
MICHEL, E., SPENCER, M., TODD, J.A., VALSAMI-JONES, E., YOUNG, J.R. & STEWART, J.R.
2011: Climate change and biosphere response: Unlocking the collections vault. –
BioScience 61: 147-153.

KELLER, L. & ROSS, K.G. 1993: Phenotypic plasticity and "cultural transmission" of alternative
social organizations in the fire ant Solenopsis invicta. – Behavioral Ecology and
Sociobiology 33: 121-129.

LAVOIE, C. 2013: Biological collections in an ever changing world: Herbaria as tools for
biogeographical and environmental studies. – Perspectives in Plant Ecology, Evolution and
Systematics 15: 68-76.

MCGLYNN, T.P., SHOTELL, M.D. & KELLY, M.S. 2003: Responding to a Variable Environment:
Home Range, Foraging Behavior, and Nest Relocation in the Costa Rican Rainforest Ant
Aphaenogaster araneoides. – Journal of Insect Behavior 16: 687-701.

MCKEON, S., WEBER, L., ADAMS, A.J. & FLEISCHNER, T.L. 2020: Human Dimensions: Natural
History as the Innate Foundation of Ecology. – The Bulletin of the Ecological Society of
America 101: 1-7.

OLIVEIRA, P.S. & FREITAS, A.V.L. 2004: Ant-plant-herbivore interactions in the neotropical

cerrado savanna. – Naturwissenschaften 91: 557-570.

OLIVEIRA, P.S., RICO-GRAY, V., DÍAZ-CASTELAZO, C. & CASTILLO-GUEVARA, C. 1999:
Interaction between ants, extrafloral nectaries and insect herbivores in Neotropical coastal
sand dunes: Herbivore deterrence by visiting ants increases fruit set in *Opuntia stricta*
(Cactaceae). – Functional Ecology 13: 623-631.

PEEL, M.C., FINLAYSON, B.L. & MCMAHON, T.A. 2007: Updated world map of the Köppen-
Geiger climate classification. – Hydrology and Earth System Sciences 11: 1633-1644.

PFEIFFER, M. & LINSENMAIR, K.E. 2000: Contributions to the life history of the Malaysian giant
ant *Camponotus gigas* (Hymenoptera, Formicidae). – Insectes Sociaux 47: 123-132.

PURCELL, J., PELLISSIER, L. & CHAPUISAT, M. 2015: Social structure varies with elevation in an
Alpine ant. – Molecular Ecology 24: 498-507.

R CORE TEAM. 2020: R: A language and environment for statistical computing.

ROBINSON, E.J.H. 2014: Polydomy: The organisation and adaptive function of complex nest
systems in ants. – Current Opinion in Insect Science 5: 37-43.

RONQUE, M.U. V, AZEVEDO-SILVA, M., MORI, G.M., SOUZA, A.P. & OLIVEIRA, P.S. 2016: Three
ways to distinguish species: Using behavioural, ecological, and molecular data to tell apart
two closely related ants, *Camponotus renggeri* and *Camponotus rufipes* (Hymenoptera:
Formicidae). – Zoological Journal of the Linnean Society 176: 170-181.

RONQUE, M.U.V., FOURCASSIÉ, V. & OLIVEIRA, P.S. 2018: Ecology and field biology of two
dominant *Camponotus* ants (Hymenoptera: Formicidae) in the Brazilian savannah. – Journal
of Natural History 52: 237-252.

SANTOS, J.C. & DEL-CLARO, K. 2009: Ecology and behaviour of the weaver ant *Camponotus*
(Myrmobrachys) *senex*. – Journal of Natural History 43: 1423-1435.

SANTOS, J.C., YAMAMOTO, M., OLIVEIRA, F.R. & DEL-CLARO, K. 2005: Behavioral repertory of
the weaver ant *Camponotus* (Myrmobrachys) *senex* (Hymenoptera: Formicidae). –
Sociobiology 46: 27-37.

SILVESTRE, R., BRANDÃO, C.R.F., ROSA, R. & SILVA, D. 2003: Grupos funcionales de hormigas:
el caso de los gremios del Cerrado. In Introducción a las hormigas de la región Neotropical.

SNELL-ROOD, E.C. 2013: An overview of the evolutionary causes and consequences of
behavioural plasticity. – Animal Behaviour 85: 1004-1011.

TAKAHASHI-DEL-BIANCO, M., HEBLING, M.J.A. & BUENO, O.C. 1998: Respiratory metabolism of
Camponotus rufipes ants: Brood and adults. – Comparative Biochemistry and Physiology -

243   A Molecular and Integrative Physiology 119: 529-532.

244 TEWKSBURY, J.J., ANDERSON, J.G.T., BAKKER, J.D., BILLO, T.J., DUNWIDDIE, P.W., GROOM,
245   M.J., HAMPTON, S.E., HERMAN, S.G., LEVEY, D.J., MACHNICKI, N.J., RIO, C.M. DEL,
246   POWER, M.E., ROWELL, K., SALOMON, A.K., STACEY, L., TROMBULAK, S.C. & WHEELER,
247   T.A. 2014: Natural history's place in science and society. – BioScience 64: 300-310.

248 TSCHINKEL, W.R. 2005: The nest architecture of the ant, *Camponotus socius*. – Journal of Insect
249   Science 5.

250 WEIDENMÜLLER, A., MAYR, C., KLEINEIDAM, C.J. & ROCES, F. 2009: Preimaginal and Adult
251   Experience Modulates the Thermal Response Behavior of Ants. – Current Biology 19:
252   1897-1902.

253 WINGFIELD, J.C. 2013a: Ecological processes and the ecology of stress: The impacts of abiotic
254   environmental factors. – Functional Ecology 27: 37-44.

255 WINGFIELD, J.C. 2013b: The comparative biology of environmental stress: Behavioural
256   endocrinology and variation in ability to cope with novel, changing environments. – Animal
257   Behaviour 85: 1127-1133.

258 YAMAMOTO, M. & DEL-CLARO, K. 2008: Natural history and foraging behavior of the carpenter
259 ant *Camponotus sericeiventris* Guérin, 1838 (Formicinae, Campotonini) in the Brazilian tropical
260 savanna. – Acta Ethologica 11: 55-65.
261
262

263 **Figures and tables captions**

264 **Table 1:** Nest architecture of *Camponotus renggeri* and *Camponotus rufipes* in an Atlantic
265 Forest Reserve, southeastern Brazil.

Nest architecture	No. of nests	
	Camponotus renggeri (N = 15)	*Camponotus rufipes* (N = 9)
Dry straw	2	4
Dry straw and dry leaves	10	4
Dry straw and fresh leaves	3	0
Underground	0	1

267 **Table 2:** Home ranges of *Camponotus renggeri* and *Camponotus rufipes* in an Atlantic Forest
268 Reserve, southeastern Brazil.

Species	Nº of nests	Season	Mean n° ± SD of foragers sampled	Mean area ± SD of home ranges (m²)	Minimum and maximum sizes of home ranges (m²)
Camponotus renggeri	4	Dry/cold	12.5 ± 3.84	0.91 ± 1.10	0.13 – 2.81
Camponotus rufipes	3	Rainy/hot	27.67 ± 8.38	1.79 ± 0.79	1 – 2.87

● Unoccupied nest   ● Active nest   ★ Observed nests   --- PEXJ trail   ······ Transportation route

Figure 1: (A) General view of the Atlantic Forest reserve physiognomy in which the study was carried out. Distribution map of (B) *Camponotus renggeri* and (C) *C. rufipes* nests. Dots represent tagged nests, darker circles around them represent the mean foraging range for each species in the Atlantic Forest; lighter circles represent the standard deviation. Stars indicate nests observed in the research. Orange represents nests which were unoccupied by the end of the research, and green represents occupied nests by the end of the research. The larger dotted lines indicate trails of the at Xixová-Japuí State Park (PEXJ), in Southeast Brazil. The smaller dotted line represents a transportation route between nests.

280 **Figure 2:** *Camponotus renggeri* dry straw nest found in an Atlantic Forest Reserve, southeastern

281 Brazil. Note the presence of whole leaves and intertwining branches in the architecture.

Figure 3: Daily activity of *Camponotus renggeri* (N = 4) and *C. rufipes* (N = 3) colonies in an Atlantic Forest reserve, southeastern Brazil. Daily activity was calculated by the number of ants leaving/entering the nests (Mean ± SE). Air temperature and humidity for each day were collected in the Brazil's National Institute for Space Research (INPE) database (Mean), and

288 humidity data was missing from some hours of the day. The arrows indicate sunrise (white) and

289 sunset (black). N/A stands for not available.

290   **Supporting information S1**: Aggressive encounter between major workers of *C. renggeri* and

291   *C. rufipes*, resulting in the decapitation of *C. Rufipes*.