Complexity of Genomic Epidemiology of Carbapenem-Resistant *Klebsiella pneumoniae* Isolates in Colombia Urges the Reinforcement of Whole Genome Sequencing-Based Surveillance Programs

Sandra Yamile Saavedra[1], Johan Fabian Bernal[2], Efraín Montilla-Escudero[1], Stefany Alejandra Arévalo[2], Diego Andrés Prada[3], María Fernanda Valencia[2], Jaime Enrique Moreno[3], Andrea Melissa Hidalgo[1], Monica Abrudan[4], Silvia Argimón[4], Mihir Kekre[4], Anthony Underwood[4], David M Aanensen[4], Carolina Duarte[1] and Pilar Donado-Godoy[2**] and the NIHR Global Health Research Unit on Genomic Surveillance of Antimicrobial Resistance[a]

[1] Grupo de Microbiología, Instituto Nacional de Salud (INS), Bogotá, Colombia

[2] Colombian Integrated Program for Antimicrobial Resistance Surveillance – Coipars, CI Tibaitatá, Corporación Colombiana de Investigación Agropecuaria (AGROSAVIA), Tibaitatá – Mosquera, Cundinamarca, Colombia

[3] Grupo de Microbiología, Dirección de Investigación en Salud Pública, Instituto Nacional de Salud, Bogotá, Colombia.

[4] Center for Genomic Pathogen Surveillance – NIHR Global Health Research Unit (GHRU). Wellcome Sanger Institute. United Kingdom

[a] Members of the NIHR Global Health Research Unit on Genomic Surveillance of Antimicrobial Resistance are listed in the Acknowledgments.

Keywords: *Klebsiella pneumoniae*; Antimicrobial resistance; Carbapenemases; Whole genome sequence (WGS).

Running title: Genomic epidemiology of CRKP in Colombia

Corresponding author: Pilar Donado-Godoy, Colombian Integrated Program for Antimicrobial Resistance Surveillance – Coipars, CI Tibaitatá, Corporación Colombiana de Investigación

Agropecuaria (AGROSAVIA), Tibaitatá – Mosquera, Cundinamarca, Colombia. Phone: +57- 1- 422-

7300 Ext 1702. Email: pidonado@agrosavia.co.

Alternate corresponding author:

Carolina Duarte, Grupo de Microbiología, Instituto Nacional de Salud (INS), Avenida calle 26 No.

51-20 - Zona 6 CAN. Bogotá, D.C, Colombia. Phone (+571) 220 7700 Ext 1420. Email:

cduarte@ins.gov.co.

40-word summary: In Colombia, the dissemination of carbapenemases in carbapenem-resistant

Klebsiella pneumoniae is attributed to horizontal gene transfer and successful circulation of CG258,

and, to a lesser extent, other clones such as ST307, ST147, and ST152.

ABSTRACT

Background: Carbapenem-resistant *Klebsiella pneumoniae* (CRKP) is an emerging public health problem. This study explores the specifics of CRKP epidemiology in Colombia based on whole genome sequencing (WGS) of the National Reference Laboratory at Instituto Nacional de Salud (INS)'s 2013-2017 sample collection.

Methods: A total of 425 CRKP isolates from 21 departments were analyzed by HiSeq-X10®Illumina high-throughput sequencing. Bioinformatic analysis was performed, primarily using the pipelines developed collaboratively by the National Institute for Health Research Global Health Research Unit (GHRU) on Genomic Surveillance of AMR, and AGROSAVIA.

Results: Of the 425 CRKP isolates, 91.5% were carbapenemase-producing strains. The data support a recent expansion and the endemicity of CRKP in Colombia with the circulation of 7 high-risk clones, the most frequent being CG258 (48.39% of isolates). We identified genes encoding carbapenemases bla_{KPC-3}, bla_{KPC-2}, bla_{NDM-1}, bla_{NDM-9}, bla_{VIM-2}, bla_{VIM-4}, and bla_{VIM-24}, and various mobile genetic elements (MGE). The virulence of CRKP isolates was low, but colibactin (*clb3*) was present in 25.2% of isolates, and a hypervirulent CRKP clone (CG380) was reported for the first time in Colombia. ST258, ST512, and ST4851 were characterized by low levels of diversity in the core genome (ANI> 99.9%).

Conclusions: The study outlines complex CRKP epidemiology in Colombia. CG258 expanded clonally and carries specific carbapenemases in specific MGEs, while the other high-risk clones (CG147, CG307, and CG152) present a more diverse complement of carbapenemases. The specifics of the Colombian situation stress the importance of WGS-based surveillance to monitor evolutionary trends of STs, MGE, and resistance and virulence genes.

ABBREVIATIONS

CRKP – Carbapenem resistant *Klebsiella pneumoniae*

KPC – *Klebsiella pneumoniae* carbapenemase

FUNDING

This work was supported by Official Development Assistance (ODA) funding from the National Institute of Health Research [award ref: 16/136/111].

This research was commissioned by the National Institute of Health Research using Official Development Assistance (ODA) funding. The views expressed in this publication are those of the authors and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health.

CONFLICT OF INTEREST

The authors: No reported conflicts of interest. All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest.

Main manuscript

INTRODUCTION

Klebsiella pneumoniae (KP) is a pathogen that causes community and hospital-acquired infections worldwide; it is a threat to public health due to its high levels of antimicrobial resistance [1-3]. Carbapenem-resistant *K. pneumoniae* (CRKP) causes untreatable infections and high mortality [2]. The emergence and global spread of extremely resistant *K. pneumoniae* highlight the need for a greater understanding of resistance epidemiology, for which whole genome sequencing (WGS) methods provide a cost-effective alternative that complements routine methods [4]. Furthermore, the epidemic dynamics of CRKP differ across geographical regions, and it is important to understand local evolutionary and epidemiological variation to establish the most appropriate strategies of surveillance, infection control, and antibiotic stewardship [5, 6].

Colombia's record of carbapenem use is one of the world's highest. [7]. According to the last Instituto Nacional de Salud (INS) nationwide report, the use of these antimicrobials exceeds more than 20 times the defined daily dose (DDD) of several countries in Europe and North America [7, 8].

In Colombia, *K. pneumoniae* is the most frequent pathogen found in intensive care units, with resistance to carbapenems reported in up to 15,6 % of isolates [9-12]. The first reports of CRKP associated with KPC-2 were in 2005, and with KPC-3 in 2008 [13, 14]. Colombia is considered KPC endemic [15]. The CRKP clones associated with KPC-type carbapenemases commonly reported in Colombia are CG258, CG307, and CG14/15 [5, 16]. Other carbapenemases identified in CRKP are VIM and NDM-1 (reported since 2011 and 2012 respectively). [12, 17, 18]. The presence of different evolutionary mechanisms in the challenged health care system of Colombia, a medium-income country, may have created the conditions for a significant CRKP epidemic [5].

The present study aimed to extend the scope of previous CRKP reports in Colombia through WGS of CRKP isolates from the 21 most populated departments of Colombia, reported to the National Reference Laboratory (NRL), Instituto Nacional de Salud (INS) between 2013 and 2017. The

objective was to increase our understanding of CRKP epidemiology in Colombia to support prevention and control strategies.

MATERIALS AND METHODS

Bacterial Isolates

The NRL manages Colombia's AMR surveillance program and adjusts the criteria for isolate submission according to the epidemiological behavior observed in the country and epidemiological alerts from the Pan American Health Organization (PAHO) (Supplementary Methods).

Between 2013 and 2017, the NRL received 811 clinical isolates of *K. pneumoniae* non-susceptible to carbapenems. After sample selection (Supplementary Materials), 425 isolates remained that met the criteria: 1) from health care-associated infections (HAI) origin; 2) year of isolation between 2013 and 2017; 3) phenotypic confirmation *of K. pneumoniae* bacterial identification by Vitek-2 (bioMérieux); 4) resistance to carbapenems determined by AST disk diffusion using CLSI guidelines for the corresponding year; and 5) genotypic confirmation of carbapenemases in *K. pneumoniae*.

Whole Genome Sequencing

Genomic DNA was extracted from single-colony cultures using PureLink™ Genomic DNA Mini Kit (Invitrogen, USA). The DNA integrity was checked by electrophoresis in a 1% (w/v) agarose gel, the DNA purity was estimated using NanoDrop® spectrophotometer (Nanodrop Technologies Inc., USA), and the concentration was determined with a Qubit Fluorometer (Invitrogen, USA). Subsequently, 1000 ng of DNA was shipped to the Wellcome Sanger Institute (Hinxton, UK) to sequence using Illumina HiSeq-X10 (San Diego, USA). DNA concentrations were confirmed using AccuBlue Broad Range assay (Biotium, Inc., Fremont, USA), followed by normalization and DNA library construction. DNA was sheared into 450-550 bp fragments using Covaris LC 220 ultrasonicator (Brighton, UK), followed by PCR-based library preparation using Illumina adaptors and 384-indexed tags (NEBNext Ultra II FS DNA library kit). Afterward, size-selection, amplification, purification, and multiplexing were carried out, and libraries were pooled. Pool was quantified and normalized down to 4nM using Biomek NXP workstation for (automated liquid

handling; California, USA), Agilent Bioanalyzer 2100 (California, USA), and Roche Lightcycler 480)

(Utah, USA), before denaturation and loading on the Illumina platform.

Quality Control and Bioinformatic Analysis

All genome sequence data were processed using versioned Nextflow workflows and associated

Docker containers covering *de novo* assembly, mapping-based SNP phylogeny, MLST assignment,

and AMR determinant detection. The workflow structure is summarized in protocols [19]. Major

quality control parameters are available in Supplementary Table 1, including file size, number of

contigs, genome size, N50. AMR determinants were detected [19], and virulence factor prediction

was performed using ARIBA software (github.com/sanger-pathogens/ariba) in conjunction with the

VFDB database (github.com/haruosuz/vfdb). MSLT prediction was assigned with ARIBA software

using PubMLST databases downloaded on December 20, 2019. The isoforms of Tn*4401* were

determined with TETyper (github.com/aesheppard/TETyper) and annotated with Prokka v1.14.5.

Additional AMR determinants, virulence factors, and typing (K-locus, wzi, and O-locus) were

detected and species predicted using Kleborate, and Kaptive as implemented on the PathogenWatch

platform (https://pathogen.watch) [20, 21].

SNP variants were called relative to a reference genome (*K. pneumoniae* strain K2044 (NCBI RefSeq

NZ_CP026011.1)) and a maximum likelihood phylogeny inferred from pseudogenome alignment of

the variants using IQ_TREE with GTR+G model and 1000 bootstrap replicates [19, 22]. High-quality

views of trees and data were rendered using the ggtree R package (bioconductor.org/packages/ggtree).

For clustering, clonal groups were defined using core genome SNP analysis using Prokka v1.14.5

gene annotation and a 70-90% ANI between isolates.

Raw sequence data were deposited at the European Nucleotide Archive under accession PRJEB29742

and visualized with Microreact (https://microreact.org/project/vcsgT8Ic4/66d1f105) [23].

RESULTS AND DISCUSSION

Geographic Spread of High-Risk Clones in Colombia

The 425 CRKP isolates analyzed in this study originated from 21 departments of Colombia, representing 90% of the total population (Figure 1a). These isolates belonged to 129 institutions (114 hospitals and 15 microbiology laboratories). The level of medical services was determined for 102 institutions, of which 68, 19 and 15 provided high, medium-high, and medium levels respectively. Of all isolates, 93.9% (n = 399) were multidrug-resistant (MDR), and 40.5% (n=172) were resistant to at least one antibiotic from all classes tested (Supplementary Table 2).

A total of 80 sequence types (STs) were identified; 42 STs were reported for the first time in Colombia, including 10 novel STs: STs 4838-4841, 4851, 4852, 4984–4987 (Supplementary Table 2). The STs were classified into 42 clonal groups (CGs) and 9 singletons (Figure 1b). The most frequent CGs were: CG258, CG147, CG152, CG14/15, CG307, CG17/20, and CG13 (Figure 1b and Supplementary Table 3). These include known epidemic ("high-risk") STs that are over-represented in the global public genome collections. Of the 7 most frequent CGs of this study, 6 had been previously reported in Colombia [5, 15, 16, 21, 24-29]. To the best of our knowledge, this is the first report of ST152 in the Americas.

CG258, the main CG identified in this study (48.9%, n=208), presented the broadest geographic distribution (18 departments). CG17/20 (2.6%, n=11) was reported in 7 departments. CG147 (5.4%, n=23) and CG 14/15 (5.2%, n=22) were present in 6 departments each, and CG152 (5.4%, n=23) in 5 departments. Three geographically close departments, Antioquia, Santander, and Valle del Cauca (60%, n=255) reported the simultaneous presence of the 7 main CGs.

All but one of the 7 main CGs showed increased and recent dissemination compared to previous reports in Colombia. Of CG258, ST11 was reported in 10 departments in this study, contrasting with previous reports, in which it was identified in only 2 departments [16, 30]. ST512 which was previously only detected in Medellin, is now reported in 4 departments [5,16]. CG147 and CG307, which had been sporadically reported in Antioquia, were reported in 11 and 6 departments, respectively [16, 26, 29, 31]. CG152 is reported for the first time in Colombia in 5 departments (Supplementary Figure 1a). In this study, the prevalence of CG14/15 (5.2%) is lower compared to previous reports, of 30% between 2005 and 2007 and 8.3% between 2012 and 2014 [5]. This study

represents the broadest geographical analysis of CRKP in Colombia and confirms their endemic status.

Most of the STs found in Colombia in this study have been reported globally. ST258, the dominant high-risk ST worldwide, was first reported in Colombia in 2006 following global clonal propagation [15, 21, 24, 25]. CG258 (ST258, ST11, ST512) has been identified as responsible for 68% of 112 *K. pneumoniae* outbreaks worldwide, reported between 2005 and 2016 [32]. The percentage of CRKP that belongs to CG258 in this study is lower than those reported in the USA (70%) by the Centers for Disease Control and Prevention (CDC), or in Israel (90%) [33-35]. ST11 is a prevalent clone in Asia, while its spread in Latin America had been only reported in Brazil [36-39]. ST512 is endemic in Italy, Israel, and Greece, ST147 in India, Italy, Tunisia, and Greece, and CG307 in Italy and the United States [16, 29, 31, 40-42]. Since differences in fitness, epidemicity, and host-related factors between CGs have been reported, molecular identification of specific high-risk clones in a region is of the utmost importance to implement appropriate infection control programs [32].

Detection and Distribution of Antimicrobial Resistance Mechanisms and Mobile Genetic Elements

Carbapenemase genes were identified in 389 CRKP genomes; 36 isolates were non-carbapenemase producers.

bla$_{KPC}$

Among carbapenemase, *bla*$_{KPC}$ was identified in 294 isolates, and *bla*$_{NDM}$ and *bla*$_{VIM}$ in 63 and 20, respectively. Co-production of *bla*$_{KPC+NDM}$ was found in 11 isolates and co-production of *bla*$_{KPC+VIM}$ in 1 isolate (Figure 1a). These findings confirm that, unlike in other regions of the world, *bla*$_{KPC}$ continues to dominate in Colombia, and *bla*$_{OXA-48-like}$ does not play a role in carbapenem resistance. They also indicate a recent expansion of *bla*$_{NDM}$ in Colombia.

We found two variants of *bla*$_{KPC}$ – *bla*$_{KPC-3}$ (39%, n=168) and *bla*$_{KPC-2}$ (30%, n=126) – across 64 different STs in multiple departments (Figure 1a). Additionally, a wide variety of mobile genetic elements with *bla*$_{KPC}$ were identified. The most common plasmid replicons in *bla*$_{KPC}$-positive isolates were IncFIB(K) (86.7%), IncFII(K) (56.8%), ColRNAI (53.1%), IncR (39.1%), and IncFII (32.3%)

(Table 1, Figure 2c). From the KPC-carrying isolates, 68% showed the presence of the Tn*4401*

transposon. In the isolates carrying bla_{KPC-2}, the gene was located in three Tn*4401* isoforms and in

non-Tn*4401* elements (NTEKPC) (n=52) (Table1, Figure 2d). In contrast, bla_{KPC-3} was localized in

Tn*4401*a in ST512 and ST1519 isolates and in Tn*4401*b in 82.5% of ST258. The clonal lineage

ST258/512 accounts for 47% of all KPC genes reported in this study, while this percentage is >70%

in the EuSCAPE sample collection [43]. The variant bla_{KPC-2} is associated with a greater variety of

mobile genetic elements than in previous reports and is found in different CGs, including CG307,

CG14/15, and CG147 (Table1), suggesting that its propagation is associated with successful

Horizontal Gene Transfer (HGT) [5,17]. In contrast, bla_{KPC-3} is correlated with the presence of two

types of Tn*4401* elements in the clonal lineage ST258/512, suggesting dissemination by clonal

expansion. These observations confirm and extend the suggested compartmentalization of the genes

encoding for the two KPC variants in Colombia by Rojas *et al.* [5]. This horizontal transfer and

promiscuity of KPC-2-carrying mobile genetic elements might have appeared in Colombia by

selective pressure due to increased use of carbapenems [7].

*bla*NDM

Two variants of bla_{NDM} were found: bla_{NDM-1} (n=60) and bla_{NDM-9} (n=3). This is the first report of

bla_{NDM-9} in the Americas. Also, it is the first time that the dissemination of bla_{NDM-1} is reported in 12

departments in Colombia (Figure 1a). The two variants were associated mainly with ST152 (33%)

and ST11 (27%) (Supplementary Figure 1b). An IncC plasmid was carried by 98.4% of bla_{NDM}

isolates. Co-location of both determinants (IncC-bla_{NDM}) in the same contig could be confirmed in

15.9% of isolates, using Artemis (Supplementary Figure 2). The IncC replicon is common in bla_{NDM}

isolates worldwide and has been previously reported in Colombia [28, 44]. Other isolates showed

incomplete replicons due to short read sequencing limitations. The genetic context was identical in all

isolates, including variants bla_{NDM-1} and bla_{NDM-9} (Supplementary Figure 2). A Tn*Ec*-like transposon

from Tn*3* family was identified in all genomic contexts with NDM genes (within 3093 bp) [28]. We

conclude that the presence of NDM is associated with the expansion of the clones ST152 and ST11

and horizontal transfer of NDM-1 via the vehicle of an IncC plasmid.

*bla*VIM

Three bla_{VIM} variants – bla_{VIM-2} (n=13), bla_{VIM-4} (n=6) and bla_{VIM-24} (n=1) – were reported in 20 isolates from 7 departments, belonging to 8 STs, mainly ST147 (23.8%, n=5), ST11 (19%, n=4), and ST15 (19%, n=4). The 3 bla_{VIM} variants were found within 2 different class 1 integrons, known to be responsible for the mobilization of VIM genes (Supplementary Figure 2) [42,45]. Our results contrast with those reported globally, where bla_{VIM-1} is the most common variant in VIM-carrying CRKP [42]. The observation that bla_{VIM-2} is the most common allele of VIM in *P. aeruginosa* in Colombia suggests that HGT may have been responsible for the intra-species transfer of the gene [45]. Further work will be carried out to investigate this hypothesis and the potential directionality of the transfer.

Co-productions

The 11 isolates with co-production of $bla_{KPC+NDM}$ came from 3 departments and were associated with 8 STs. In the isolates co-harboring $bla_{KPC-3+NDM-1}$, it was observed that bla_{KPC-3} incorporated with Tn*4401*a exhibited a genetic environment, including [trpR (Tn3)-ISPsy42(Tn3)-ISKpn7(IS21)-blaKPC-3-ISKpn6(IS1182)-aacA4], and that the vehicle for bla_{NDM-1} was similar (TnEc-like). Co-producers of $bla_{KPC-2+NDM-1}$ contained the same bla_{NDM-1} platform (TnEc-like), but the bla_{KPC-2} genetic environment revealed a diversity of genomic contexts. In the isolate co-producers of $bla_{KPC-3+VIM-24}$, the bla_{VIM-24} gene was found associated with class-1 integron in IncC, and bla_{KPC-3} exhibited Tn*4401*b complemented with Tn*2* transposases (Supplementary Figure 2). CRKPs with co-production of at least 2 genes have been previously reported worldwide, including bla_{KPC} and bla_{VIM} in Italy, $bla_{KPC-2+NDM-1}$ in China and Brazil, and $bla_{NDM-1+OXA-181}$ in Singapore [46]. The 3 co-productions reported in this study have been previously sporadically reported in Colombia [11,12,28], but this is the first report of the full replicon. Monitoring the dynamics of CRKPs with co-productions in the surveillance program is essential because of the potential risks they present, and since the way they emerge and whether they are only transient or have transmission capacity remain uncertain [43, 46].

CRKP isolates not harboring carbapenemases

In 36 CRKP isolates not harboring a carbapenemase gene in their genomes, the resistance to carbapenems could be attributed to the presence of extended spectrum β-lactamase (ESBL) genes (86%, n = 31), especially $bla_{CTX-M-15}$ (22/31); in 24 of the 31 isolates, these ESBL genes were in combination with porin truncations (Supplementary Table 2).

Porin truncations or mutations were found in 52% of all isolates. These truncations result in a non-functional pore that acts in concert to lower carbapenem concentrations, limiting antibiotic influx [47]. In one isolate, no resistance mechanisms were identified (Supplementary Table 3).

The enzymatic hydrolysis of carbapenems by carbapenemases was identified as the primary mechanism of carbapenem resistance, and the vast majority of carbapenemase producer isolates (91%, n=385) were resistant simultaneously to all screening carbapenems (IPM, MEM, ETP) (Supplementary Figure 3). Possible explanations include the preferential use of carbapenems in hospitals for the treatment of serious infections caused by carbapenem resistance isolates, as well as the circulation of bla_{KPC-2} and bla_{KPC-3}-harboring isolates promoting the progression of carbapenem resistance in Colombia, which is a KPC endemic country.

Additional AMR determinants to aminoglycosides (n=390), fluoroquinolones (n=378), phenicols (n=317), ESBL and/or AmpC (n=248), and tetracyclines (n=219) were observed in the majority of isolates (Supplementary Table 2).

Our results outline the increasing complexity of the CRKP population, resistance mechanisms, and lineages in Colombia, which will require adaptations of the conventional infection control measures that were successful in areas where the objective was to curb the expansion of a single dominant clone.

Detection of Virulence Mechanisms and Serotypes

We observed a low prevalence of acquired virulence mechanisms among CRKP isolates in this study, which is in line with global trends [48]. However, we additionally observed an over-representation of colibactin genes among the isolates and the first observation in Colombia of hyper-virulent CRKP. The CG258 isolates exhibited different types of K-locus (KL), *wzi*, and O-locus (OL). The most common profile was KL107-*wzi*154:O2v2 (STs 258, 512, 1519, and 4851), present in 62.5% of isolates (Figure 2a, Figure 2b, Supplementary Table 3) [21, 40, 49, 50]. The profile KL105-*wzi*75:O2v2, detected in 24% of isolates, is unusual for ST11 [21]. In China, where ST11 is endemic, the dominant KLs are KL47 and KL64, but the occurrence of capsule switching has been suggested in the case of this particular ST [37, 51]. In CG258, 76.9% of isolates were carriers of yersiniabactin

(*ybt*), mainly of the *ybt*17 (105 isolates of ST258 and 2 of ST4851) and *ybt*10 (50 isolates of ST11 and 1 of ST258) lineages (Figure 2a, Figure 2b, Supplementary Table 3). Details for other CGs are shown in Figure 2a and Supplementary Table 3.

It is remarkable to note that isolates of ybt17 / colibactin (*clb3*) lineages represented 25.2% of all isolates, while the spread of *clb3* among CRKP is usually around 10% [52]. However, most isolates *ybt*17+*clb*3+ST258 have been reported in the literature as not being colibactin producers, due to a deletion in *clb*J/*clb*K [50,52].

Only one isolate (RB 877) from this study exhibited determinants associated with hypervirulence, such as *ybt*16, aerobactin (*iuc*2), salmochelin (*iro*2), and *rmpA_3* (KpVP2). This ST380 isolate was recovered in Antioquia in 2013 and is a producer of KPC-2, with the profile KL2-*wzi*203:O1v1 (Table 1, Supplementary Table 2). CG380 has been classified as a hypervirulent clone [48]. This is the first time a hypervirulent CRKP is reported in Colombia.

The two cases of virulence elements reported in this study illustrate the risk of convergence of resistance and virulence, either through the acquisition of virulence plasmid(s) by CRKP or through the acquisition of resistance mobile genetic element(s) by virulent clone. The latter appears to be more difficult [52]. This emphasizes the need for genomic surveillance programs to include both resistance and virulence locus information.

Genetic Diversity Among CRKP Isolates

The phylogenetic analysis by SNP reflected the MLST distribution into STs as described above. Additionally, it highlights the genomic diversity within each ST. Of interest, we observed ST11, ST147, ST152, and ST307 with low nucleotide divergence (0.05 - 0.47%), suggesting clonality (Figure 1b). However, a deeper analysis by ST showed distinct clades within each ST (Supplementary Figure 4).

Among the 208 isolates of CG258 (48.9% of the total isolates) grouped into 7 STs (Figure 1b and 2b), the ST258, ST512, and ST4851 were characterized by low levels of phylogenetic diversity (ANI> 99.9%). This indicates a recent and common evolutionary ancestor that supports their grouping. The remaining STs of CG258 showed greater diversity (ANI 99.7-99.8%) among pairs. ST11 isolates

carried a wide variety of carbapenemases (KPC, NDM, VIM). To our knowledge, this is the first report of $bla_{NDM+VIM}$-carrying ST11 isolates in Colombia, although they have been reported worldwide [42, 44, 53]. There were several distinct clades of ST11, with different serotypes, resistance genes, and plasmid replicons (Supplementary Figure 4). This opens the possibility of multiple introductions or emergence of ST11 into Colombia and should be further investigated by contextualizing the genomes with global collections.

The ST147 showed several clades with differences in carbapenemase genes, related to Inc-type elements, discarding the clonality of the group (Supplementary Figure 4). Similarly, ST307 showed different clades some with bla_{KPC}, bla_{NDM}, or bla_{VIM}, alongside IncFIB mobile elements and ESBL genes. The apparent clonality may come from the acquisition of carbapenemase genes in multiple events, related to mobile elements. This is in accordance with ST147 found related to Inc-type plasmids, and ST307 to $bla_{OXA48-like}$ plasmid genes, both related to carbapenemase genes HGT [43]. ST152 was found as an emerging clone with a genomic profile characterized by bla_{NDM}, several Inc-type mobile elements, and hypervirulence provided by the presence of yersiniabactin, similar to that reported in Saudi Arabia and other countries as a hypervirulent ST of importance in urinary tract infections [54]. All previous analyses enhances the importance of WGS for AMR surveillance, as differential findings between STs and genes reveal non-clonality that may be translated into differential treatment of CRKP infections.

Among this study's limitations were: having only voluntary surveillance isolates at the national level; and limited data of clinical and epidemiological information for a more exhaustive investigation. Nevertheless, the information obtained is essential for national surveillance and provides insights into pathogen genetics for planning effective surveillance.

CONCLUDING REMARKS

This study allowed a detailed description of CRKP epidemiology in Colombia that should be taken into account when establishing efficient WGS-based surveillance programs in the country.

There are 7 predominant high-risk STs in Colombia. In addition to the previously described CG258 with KPC, we showed a complex, recent and significant expansion of multiple clones of interest and carbapenemases, urging reinforcement of the CRKP genomic surveillance process.

The dissemination of carbapenemase genes is based on a complex combination of mechanisms. KPC-3 occurs by the expansion of CG258, mainly ST258 and ST512 carrying an IncFII replicon plasmid with the insertion of transposon Tn*4401*-like. The spread of KPC-2 and VIM genes was associated with the horizontal transfer of different plasmid replicons and mobile elements. NDM-1 dissemination was associated with the expansion of clones CG152 and ST11 (CG258) and possible HGT related to IncC. The high proportion of HTG mechanisms in disseminating carbapenem resistance genes could have resulted from higher selective pressure due to overuse of antibiotics in a challenged health care system. This result shows the importance of including mobile genetic elements in the CRKP surveillance program in Colombia.

The hypervirulent strain report and the over-representation of the colibactin virulence factor urge close monitoring of the potential convergence of virulence and resistance of CRKP in Colombia.

ACKNOWLEDGMENTS

Members of the NIHR Global Health Research Unit on Genomic Surveillance of Antimicrobial Resistance: Khalil Abudahab, Harry Harste, Dawn Muddyman, Ben Taylor, Nicole Wheeler, and Sophia David of the Centre for Genomic Pathogen Surveillance, Big Data Institute, University of Oxford, Old Road Campus, Oxford, United Kingdom and Wellcome Genome Campus, Hinxton, UK; Alejandra Arevalo, and Erik C. D. Osma Castro of the Colombian Integrated Program for Antimicrobial Resistance Surveillance – Coipars, CI Tibaitatá, Corporación Colombiana de Investigación Agropecuaria (AGROSAVIA), Tibaitatá – Mosquera, Cundinamarca, Colombia; K. L. Ravikumar, Geetha Nagaraj, Varun Shamanna, Vandana Govindan, Akshata Prabhu, D. Sravani, M. R. Shincy, Steffimole Rose, and Ravishankar K.N of the Central Research Laboratory, Kempegowda Institute of Medical Sciences, Bengaluru, India; Iruka N Okeke, Anderson O. Oaikhena, Ayorinde O. Afolayan, Jolaade J Ajiboye, and Erkison Ewomazino Odih of the Department of Pharmaceutical Microbiology, Faculty of Pharmacy, University of Ibadan, Oyo State, Nigeria; Celia Carlos, Marietta L. Lagrada, Polle Krystle V. Macaranas, Agnettah M. Olorosa, June M. Gayeta, and Elmer M. Herrera of the Antimicrobial Resistance Surveillance Reference Laboratory, Research Institute for Tropical Medicine, Muntinlupa, the Philippines; Ali Molloy, alimolloy.com; John Stelling, The Brigham and Women's Hospital; and Carolin Vegvari, Imperial College London.

The authors acknowledge all public health laboratories from the National Reference Laboratory (NRL) for providing the isolates CRKP obtained under the previous National Surveillance by the Laboratory of Antimicrobial Resistance Surveillance Network.
We acknowledge Angela Sofía García, MSc, part of the Colombian Genomic Unit of AGROSAVIA, for the valuable suggestions on the text to enrich the content of this article.

REFERENCES

1. Bengoechea JA, Sa Pessoa J. Klebsiella pneumoniae infection biology: living to counteract host defences. *FEMS Microbiol Rev* **2018**; 43: 123–144. Available at: https://doi.org/10.1093/femsre/fuy043.

2. Caneiras C, Lito L, Melo-Cristino J, Duarte A. Community- and Hospital-Acquired Klebsiella pneumoniae Urinary Tract Infections in Portugal: Virulence and Antibiotic Resistance. *Microorganisms* **2019**; 7. Available at: https://doi.org/10.3390/microorganisms7050138.

3. WHO. Global Priority List of Antibiotic-Resistant Bacteria to Guide Research, Discovery, and Development of New Antibiotics. 2017; Available at:

https://www.who.int/medicines/publications/WHO-PPL-Short_Summary_25Feb-ET_NM_WHO.pdf?ua=1.

4. Köser CU, Ellington MJ, Peacock SJ. Whole-genome sequencing to control antimicrobial resistance. *Trends Genet* **2014**; 30: 401–407. Available at: https://doi.org/10.1016/j.tig.2014.07.003.

5. Rojas LJ, Weinstock GM, De La Cadena E, et al. An Analysis of the Epidemic of Klebsiella pneumoniae Carbapenemase-Producing K. pneumoniae: Convergence of Two Evolutionary Mechanisms Creates the "Perfect Storm". *J Infect Dis* **2017**; 217:82–92. Available at: https://doi.org/10.1093/infdis/jix524.

6. Argimón S, Masim MAL, Gayeta JM, et al. Integrating whole-genome sequencing within the National Antimicrobial Resistance Surveillance Program in the Philippines. *Nat Commun* **2020**; 11: 1–15. Available at: https://doi.org/10.1038/s41467-020-16322-5. Accessed 7 January 2021.

7. Gomez Rubio A. Consumo de antibioticos en el ámbito hospitalario. Colombia 2017. 2017. Available at: https://www.ins.gov.co/buscador-eventos/Informesdeevento/CONSUMO DE ANTIBIÓTICOS 2017.pdf.

8. European Centre for Disease Prevention and Control. Antimicrobial consumption in the EU / EEA Annual Epidemiological Report for 2019 Key facts. Stockholm: 2020. Available at: https://www.ecdc.europa.eu/sites/default/files/documents/Antimicrobial-consumption-in-the-EU-Annual-Epidemiological-Report-2019.pdf .

9. INS. Resultados del Programa de Vigilancia por Laboratorio de Resistencia antimicrobiana en Infecciones Asociadas a la Atención en Salud (IAAS) 2015. 2016. Available at: https://www.ins.gov.co/BibliotecaDigital/informe-vigilancia-por-laboratorio-resistencia-antimicrobiana-y-whonet-iaas-2015.pdf

10. INS. Resultados del Programa de Vigilancia por Laboratorio de Resistencia antimicrobiana en Infecciones Asociadas a la Atención en Salud (IAAS) 2016. **2017**. Available at: http://www.ins.gov.co/buscador-eventos/Informacin de laboratorio/Informe Vigilancia por Laboratorio Resistencia Antimicrobiana y Whonet IAAS 2016.pdf

11. INS. Resultados del Programa de Informe de Resultados de la Vigilancia por Laboratorio de Resistencia antimicrobiana en Infecciones Asociadas a la Atención en Salud (IAAS) 2017. 2018. Available at: http://www.ins.gov.co/buscador-eventos/Informacin de laboratorio/Informe Vigilancia por Laboratorio Resistencia Antimicrobiana y Whonet IAAS 2017.pdf

12. Ovalle M V, Saavedra SY, González MN, Hidalgo AM, Duarte C, Beltrán M. Results of the national surveillance of antimicrobial resistance of Enterobacteriaceae and Gram negative bacilli in health care-associated infections in Colombia, 2012-2014. *Biomedica* **2017**; 37: 473–485. Available at: http://dx.doi.org/10.7705/biomedica.v34i2.3432.

13. Villegas M V, Lolans K, Correa A, et al. First detection of the plasmid-mediated class A carbapenemase KPC-2 in clinical isolates of Klebsiella pneumoniae from South America. *Antimicrob Agents Chemother* **2006**; 50: 2880–2882. Available at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1538657/pdf/0186-06.pdf.

14. Lopez JA, Correa A, Navon-Venezia S, et al. Intercontinental spread from Israel to Colombia of a KPC-3-producing Klebsiella pneumoniae strain. *Clin Microbiol Infect* **2011**; 17: 52–56. Available at: https://doi.org/10.1111/j.1469-0691.2010.03209.x.

15. Munoz-Price LS, Poirel L, Bonomo RA, et al. Clinical epidemiology of the global expansion of Klebsiella pneumoniae carbapenemases. *Lancet Infect Dis* **2013**; 13: 785–796. Available at: https://doi.org/10.1016/S1473-3099(13)70190-7.

16. Ocampo AM, Chen L, Cienfuegos A V, et al. A Two-Year Surveillance in Five Colombian Tertiary Care Hospitals Reveals High Frequency of Non-CG258 Clones of CarbapenemResistant

Klebsiella pneumoniae with Distinct Clinical Characteristics. *Antimicrob Agents Chemother* **2016**; 60:

332–342. Available at: https://doi.org/10.1128/AAC.01775-15.

17. Montealegre MC, Correa A, Briceño DF, et al. Novel VIM metallo-beta-lactamase variant, VIM-

24, from a Klebsiella pneumoniae isolate from Colombia. *Antimicrob Agents Chemother* **2011**; 55:

2428–2430. Available at: https://doi.org/10.1128/AAC.01208-10.

18. Escobar Pérez JA, Olarte Escobar NM, Castro-Cardozo B, et al. Outbreak of NDM-1-producing

Klebsiella pneumoniae in a neonatal unit in Colombia. *Antimicrob Agents Chemother* **2013**; 57:

1957–1960. Available at: https://doi.org/10.1128/AAC.01447-12.

19. Underwood A. GHRU (Genomic Surveillance of Antimicrobial Resistance) Retrospective 1

Bioinformatics Methods. protocols.io: 2020. Available at:

https://gitlab.com/cgps/ghru/pipelines/dsl2/pipelines/amr_prediction. Accessed 7 January 2021.

20. Lam MMC, Wick RR, Wyres KL, Holt KE. Genomic surveillance framework and global

population structure for Klebsiella pneumoniae. bioRxiv **2020**; :2020.12.14.422303. Available at:

http://biorxiv.org/content/early/2020/12/14/2020.12.14.422303.abstract

https://www.biorxiv.org/content/biorxiv/early/2020/12/14/2020.12.14.422303.full.pdf.

21. Wyres KL, Wick RR, Gorrie C, et al. Identification of Klebsiella capsule synthesis loci from

whole genome data. *Microb Genom* **2016**; 2: e000102. Available at:

https://doi.org/10.1099/mgen.0.000102.

22. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A Fast and Effective Stochastic

Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol Biol Evol* **2014**; 32:268–274.

Available at: https://doi.org/10.1093/molbev/msu300.

23. Argimón S, Abudahab K, Goater RJE, et al. Microreact: visualizing and sharing data for genomic

epidemiology and phylogeography. *Microb Genom* **2016**; 2: e000093. Available at:

https://doi.org/10.1099/mgen.0.000093.

24. Chen L, Mathema B, Chavda KD, DeLeo FR, Bonomo RA, Kreiswirth BN. Carbapenemase-

producing Klebsiella pneumoniae: molecular and genetic decoding. *Trends Microbiol* **2014**; 22: 686–

696. Available at: https://doi.org/10.1016/j.tim.2014.09.003.

25. Pitout JDD, Nordmann P, Poirel L. Carbapenemase-Producing Klebsiella pneumoniae, a Key Pathogen Set for Global Nosocomial Dominance. *Antimicrob Agents Chemother* **2015**; 59: 5873–5884. Available at: https://doi.org/10.1128/AAC.01019-15.

26. Cienfuegos-Gallet A V, Ocampo de Los Ríos AM, Sierra Viana P, et al. Risk factors and survival of patients infected with carbapenem-resistant Klebsiella pneumoniae in a KPC endemic setting: a case-control and cohort study. *BMC Infect Dis* **2019**; 19: 830. Available at: https://doi.org/10.1186/s12879-019-4461-x.

27. D'Apolito D, Arena F, Conte V, et al. Phenotypical and molecular assessment of the virulence potential of KPC-3-producing Klebsiella pneumoniae ST392 clinical isolates. *Microbiol Res* **2020**; 240: 126551. Available at: https://doi.org/10.1016/j.micres.2020.126551.

28. Rojas LJ, Wright MS, De La Cadena E, et al. Initial Assessment of the Molecular Epidemiology of blaNDM-1 in Colombia. *Antimicrob Agents Chemother* **2016**; 60: 4346–4350. Available at: https://doi.org/10.1128/AAC.03072-15.

29. Peirano G, Chen L, Kreiswirth BN, Pitout JDD. Emerging Antimicrobial Resistant High-Risk clones among Klebsiella pneumoniae: ST307 and ST147. *Antimicrob Agents Chemother* **2020**; 64: e01148-20. Available at: http://aac.asm.org/content/64/10/e01148-20.abstract https://aac.asm.org/content/64/10/e01148-20.long.

30. Garcia-Fulgueiras V, Zapata Y, Papa-Ezdra R, et al. First characterization of K. pneumoniae ST11 clinical isolates harboring blaKPC-3 in Latin America. *Rev Argent Microbiol* **2019**; Available at: https://doi.org/10.1016/j.ram.2019.10.003.

31. Villa L, Feudi C, Fortini D, et al. Diversity, virulence, and antimicrobial resistance of the KPC-producing Klebsiella pneumoniae ST307 clone. **2017**; 3. Available at: https://doi.org/10.1099/mgen.0.000110.

32. Navon-Venezia S, Kondratyeva K, Carattoli A. Klebsiella pneumoniae: A major worldwide source and shuttle for antibiotic resistance. *FEMS Microbiol Rev* **2017**; 41: 252–275.

33. Kitchel B, Rasheed JK, Patel JB, et al. Molecular epidemiology of KPC-producing Klebsiella pneumoniae isolates in the United States: Clonal expansion of multilocus sequence type 258.

Antimicrob Agents Chemother **2009**; 53: 3365–3370. Available at:

https://doi.org/10.1128/AAC.00126-09.

34. Schwaber MJ, Lev B, Israeli A, et al. Containment of a Country-wide Outbreak of Carbapenem-Resistant Klebsiella pneumoniae in Israeli Hospitals via a Nationally Implemented Intervention. *Clin Infect Dis* **2011**; 52: 848–855. Available at: https://doi.org/10.1093/cid/cir025. Accessed 7 January 2021.

35. Pathogenwatch. Genomes KPN ST258. Available at:

https://pathogen.watch/genomes/all/map?genusId=570&mlst=258&speciesId=573. Accessed 24 March 2021.

36. Qi Y, Wei Z, Ji S, Du X, Shen P, Yu Y. ST11, the dominant clone of KPC-producing Klebsiella pneumoniae in China. *J Antimicrob Chemother* **2011**; 66: 307–312. Available at:

https://doi.org/10.1093/jac/dkq431.

37. Zhou K, Xiao T, David S, et al. Novel Subclone of Carbapenem-Resistant Klebsiella pneumoniae Sequence Type 11 with Enhanced Virulence and Transmissibility, China. *Emerg Infect Dis* **2020**; 26: 289–297.

38. Pereira PS, de Araujo CF, Seki LM, Zahner V, Carvalho-Assef AP, Asensi MD. Update of the molecular epidemiology of KPC-2-producing Klebsiella pneumoniae in Brazil: spread of clonal complex 11 (ST11, ST437 and ST340). *J Antimicrob Chemother* **2013**; 68: 312–316. Available at: https://doi.org/10.1093/jac/dks396.

39. Pathogenwatch. Genomes KPN ST11. Available at:

https://pathogen.watch/genomes/all/map?genusId=570&mlst=11&speciesId=573. Accessed 24 March 2021.

40. David S, Reuter S, Harris SR, et al. Epidemic of carbapenem-resistant Klebsiella pneumoniae in Europe is driven by nosocomial spread. *Nat Microbiol* **2019**; 4: 1919–1929. Available at:

https://doi.org/10.1038/s41564-019-0492-8.

41. Pathogenwatch | Genomes KPN ST512. Available at:

https://pathogen.watch/genomes/all/map?genusId=570&mlst=512&speciesId=573. Accessed 24 March 2021.

42. Matsumura Y, Peirano G, Devinney R, et al. Genomic epidemiology of global VIM-producing Enterobacteriaceae. *J Antimicrob Chemother* **2017**; 72:2249–2258. Available at: https://doi.org/10.1093/jac/dkx148.

43. David S, Cohen V, Reuter S, et al. Integrated chromosomal and plasmid sequence analyses reveal diverse modes of carbapenemase gene spread among Klebsiella pneumoniae. *Proc Natl Acad Sci U S A* **2020**; 117: 25043–25054. Available at: https://doi.org/10.1073/pnas.2003407117.

44. Wu W, Feng Y, Tang G, Qiao F, McNally A, Zong Z. NDM Metallo-β-Lactamases and Their Bacterial Producers in Health Care Settings. *Clin Microbiol Rev* **2019**; 32. Available at: https://doi.org/10.1128/CMR.00115-18.

45. Correa A, Del Campo R, Perenguez M, et al. Dissemination of high-risk clones of extensively drug-resistant Pseudomonas aeruginosa in colombia. *Antimicrob Agents Chemother* **2015**; 59: 2421–2425. Available at: https://doi.org/10.1128/AAC.03926-14.

46. Gao H, Liu Y, Wang R, Wang Q, Jin L, Wang H. The transferability and evolution of NDM-1 and KPC-2 co-producing Klebsiella pneumoniae from clinical settings. *EBioMedicine* **2020**; 51. Available at: https://doi.org/10.1016/j.ebiom.2019.102599.

47. Wong JLC, Romano M, Kerry LE, et al. OmpK36-mediated Carbapenem resistance attenuates ST258 Klebsiella pneumoniae in vivo. *Nat Commun* **2019**; 10. Available at: http://dx.doi.org/10.1038/s41467-019-11756-y.

48. Wyres KL, Wick RR, Judd LM, et al. Distinct evolutionary dynamics of horizontal gene transfer in drug resistant and virulent clones of Klebsiella pneumoniae. *PLoS Genet* **2019**; 15: e1008114. Available at: https://doi.org/10.1371/journal.pgen.1008114.

49. Lam MMC, Wyres KL, Judd LM, et al. Tracking key virulence loci encoding aerobactin and salmochelin siderophore synthesis in Klebsiella pneumoniae. *Genome Med* **2018**; 10: 77. Available at: https://doi.org/10.1186/s13073-018-0587-5.

50. Wyres KL, Lam MMC, Holt KE. Population genomics of Klebsiella pneumoniae. *Nat Rev Microbiol* **2020**; 18: 344–359. Available at: https://doi.org/10.1038/s41579-019-0315-1.

51. Zhao Q, Guo L, Wang L-F, Zhao Q, Shen D-X. Prevalence and characteristics of surgical site hypervirulent Klebsiella pneumoniae isolates. *J Clin Lab Anal* **2020**; e23364. Available at: https://doi.org/10.1002/jcla.23364.

52. Lam MMC, Wick RR, Wyres KL, et al. Genetic diversity, mobilisation and spread of the yersiniabactin-encoding mobile element ICEKp in Klebsiella pneumoniae populations. *Microb Genom* **2018**; 4: e000196. Available at:

https://www.microbiologyresearch.org/content/journal/mgen/10.1099/mgen.0.000196. Accessed 7 January 2021.

53. Roach D, Waalkes A, Abanto J, et al. Whole Genome Sequencing of Peruvian Klebsiella pneumoniae Identifies Novel Plasmid Vectors Bearing Carbapenem Resistance Gene NDM-1. *Open Forum Infect Dis* **2020**; 7: ofaa266. Available at: https://doi.org/10.1093/ofid/ofaa266.

54. Al-Agamy MH, El-Mahdy TS, Radwan HH, Poirel L, Garcia-Perdomo HA. Cooccurrence of NDM-1, ESBL, RmtC, AAC(6′)-Ib, and QnrB in Clonally Related Klebsiella pneumoniae Isolates Together with Coexistence of CMY-4 and AAC(6′)-Ib in Enterobacter cloacae Isolates from Saudi Arabia. *Biomed Res Int* **2019**; 2019: 6736897. Available at:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6699326/pdf/BMRI2019-6736897.pdf.

Tables

Table 1. Characteristics of carbapenem-resistant *K. pneumoniae* isolates.

Resistance mechanisms to carbapenems	Number of isolates (% of total)	Number of STs	Number of departments	Carbapenemase gene variants	Kpc element (Tn*4401*)	Plasmid Replicon Type
Carbapenemase *bla*KPC-like	294 (69.1)	64 STs (ST258, n=106); (ST11, n=34); (ST 512, n=32); (other STs, n=122)	17 departments (Antioquia=77) (Atlántico=2) (Bogotá=5) (Bolivar=5) (Boyacá=11) (Caldas=8) (Caquetá=1) (Casanare=5) (Cauca=8) (Córdoba=3) (Cundinamarca=12) (Huila=7) (Meta=9) (Nariño=15) (N. Santander=17) (Quindio=3) (Risaralda=10) (Santander=50) (Valle del Cauca=46)	KPC-2 (n=126) KPC-3 (n=168)	Tn*4401*_truncC (n=7) Tn*4401*a (n=48) Tn*4401*b (n=145) No-Tn*4401* (n=94)	IncFIB_K = 255 IncFII_K = 167 ColRNAI = 156 IncR = 115 IncFII = 95 IncI2 = 70 IncN = 46 IncFIB_pQil = 35 IncX3 = 33 Others = 215 (25 replicons)
Carbapenemase *bla*NDM-like	63 (14.8)	8 STs (ST11, n=17); (ST152, n=21); (others STs, n=25)	12 departments (Antioquia=10) (Atlantico=1) (Bogota n=3) (Caldas=1) (Cauca=2)	NDM-1 (n=60) NDM-9 (n=3)	-	Inc._1 = 62 IncFIB_K = 51 IncFII_K = 35 ColRNAI = 22 IncR = 17 FIA_pBK30683 = 13

			(Cundinamarca=1)			Others= 38 (12 replicons)
			(Huila=2)			
			(Nariño=2)			
			(N.Santander=5)			
			(Risaralda= 9)			
			(Santander n=21)			
			(Valle del Cauca n=6)			
Carbapene mase *bla***VIM- like**	20 (4.7)	8 STs (ST11, n=4); (ST15, n=4); (ST70, n=4) (ST147, n=5); (ST273, n=1); (ST307, n=1); (ST340, n=2); (ST833, n=2);	7 departments (Antioquia=4) (Atlántico=1) (Bogotá=2) (Boyacá=1) (Cesar=1) (N. Santander=2) (Santander=9)	VIM-2 (n=13); VIM-4 (n=6); VIM-24 (n=1)	-	Inc._1 = 20 IncFIB_K = 15 IncR = 14 Col_pHAD28 = 7 Col440I = 6 IncFII_K = 6 FIA_pBK30683 = 4 ColRNAI = 3 Others = 15 (9 replicons)
Co- production of carbapene mases *bla***KPC- like &** *bla***NDM- like**	11 (2.6)	8 STs (ST11, n=2); (ST25, n=2); (ST76, n=1); (ST152, n=1); (ST219, n=1); (ST258, n=1); (ST340, n=1); (ST392, n=2);	3 departments (Antioquia, n=3); (Santander, n=2); (Huila, n=1)	KPC-2+NDM-1 (n=9); KPC-3+NDM-1 (n=2)	Tn*4401*a (n=1) Tn*4401*b (n=2) No-Tn*4401* (n=8)	IncFIB_K = 8 IncFII_K = 8 Inc._1 = 6 IncN = 5 IncR = 4 ColRNAI = 3 Col_pHAD28 = 2 FIA_pBK30683 = 1 IncFIB_pQil = 1 IncFII = 1 IncI2 = 1 IncM1 = 1
Co- production of carbapene mases *bla***KPC- like &**	1 (0.23)	1 ST (ST3660, n=1)	1 department (Bogota n=1)	KPC2+VIM-4 (n=1)	Tn*4401*b (n=1)	Inc._1 = 1 IncFIB_K = 1 IncFII_K = 1 IncR = 1

ESBL gene + porin defects, but no carbapene mase	22 (5.2)	18 STs (ST11 n=1); (ST15 n=1); (ST25 n=1) (ST29 n=1) (ST35 n=1) (ST37 n=1) (ST40 n=1) (ST70 n=2) (ST147 n=1) (ST258 n=1) (ST300 n=1) (ST307 n=1) (ST340 n=1) (ST405 n=1) (ST629 n=1) (ST4841 n=2) (ST4986 n=1) (ST4987 n=1)	9 departments (Antioquia=8) (Bolivar=1) (Caldas=1) (Huila=1) (Nariño=2) (N. Santander=1) (Santander=5) (Sucre=1) (Valle del Cauca = 2)	-	-	IncFIB_K = 15 IncFII_K = 15 IncR = 6 IncFIB_pKPHS1 = 4 IncFIB_pNDM_Mar = 4 IncHI1B_pNDM_MAR = 2 FIA_pBK30683 = 2 Col_pHAD28 = 2 Others = 17 (14 replicons)
ESBL gene, but no carbapene mase	7 (1.64)	6 STs (ST11 n=1); (ST39 n=1); (ST40 n=1) (ST147 n=2) (ST1373 n=1) (ST1758 n=1)	3 departments (Santandern=3); (Antioquia n=2) (Atlantico n=2)		-	IncFIB_K = 7 IncFII_K = 3 IncR = 3 Col_pHAD28 = 2 Col156 = 1 Col440I = 1 Col440II = 1 FIA_pBK30683 = 1 IncFIB_pKPHS1 = 1 IncN = 1
Porin defects, but no carbapene mases	6 (1.41)	3 STs (ST1198 n=1); (ST70 n=3); (ST14 n=2)	2 departments (Antioquia n=3) (Nariño,n=3);	-	-	IncFIB_K = 6 Inc._1 = 3 IncFIB_pKPHS1 = 3 IncFII_K = 3 IncR = 1 IncX4_1 = 1

ColRNAI = 1

No Carbapene mase, No Porin, No BLEE	1 (0.23)	1 STs (ST4984 n=1);	1 department (Antioquia n=1)	-	-	IncFIB_K = 1 IncR = 1

FIGURE LEGENDS

Figure 1. Geographical distribution of mechanisms of resistance to carbapenems in *Klebsiella pneumoniae* and core genome tree in Colombia, 2013-2017. a) Map depicting the 21 departments that submitted isolates for surveillance. The isolates were classified according to the mechanism of resistance to carbapenem by color (see key). Pie chart shows the proportions of different mechanisms of resistance to carbapenems in each department. b) Circular representation of the transformed phylogenetic tree showing the genetic relationships among 421 Colombian *K. pneumoniae* CRKP isolates. The phylogenetic tree was built using 421/425 isolates; 4 were excluded from the analysis because they presented a high number of N in the multiple alignments. The 421 isolates were classified into 80 sequence types (STs) grouped in 42 CG and 9 singletons. Due to the number of different CGs, for the graph, CGs with ≥6 isolates were selected for easy viewing on the tree: CG258, CG13, CG14/15, CG17/20, CG25, CG39, CG45, CG70, CG76, CG147, CG152, and CG307. The remaining CG and singletons were classified as other-CG.

Figure 2. Clonal group heat maps, virulence factors, resistance profile, carbapenemases, replicons, and alluvial of the CG258. The heatmap analyzes different clonal groups (each row of the heatmap represents a clonal group). a) Heatmap displaying the distribution of antimicrobials and the mechanism of resistance to carbapenems (y-axis) in the clonal group. b) Heatmap displaying the distribution of O locus, K locus (*wzi*), and virulence factors (y-axis) found in the clonal group. c) Heatmap displaying the distribution of the replicon type (y-axis) found in the clonal group. d) Alluvial diagram showing the "flow" of the presence of the most important transposable elements associated with carbapenemase and ST of CG258.

FIGURES

Figure 1.

Figure 2.

a) Distribution of resistance profile and carbapenemase by clonal group

d) Distribution of ST, carbapenemase, mobile elements, O locus, K locus, wzi in 258 clonal group

b) Distribution of O locus, K locus (*wzi*) and virulence factor by clonal group

c) Distribution of replicon markers by clonal group CG258 and non CG258